Title: Single Cell Transcriptional Profiling of Phox2b-Expressing Geniculate Ganglion Neurons

Authors: Catherine B. Anderson and Eric D. Larson

Affiliations: Department of Otolaryngology, University of Colorado Anschutz Medical Campus, Aurora CO 80045

Article Type: Original Report

Corresponding Author:

Eric D. Larson

University of Colorado, Department of Otolaryngology

12700 E. 19th Ave, MS 8606

Aurora, CO 80045

Key Words: Phox2b, geniculate ganglion, single cell sequencing, taste

Abstract

The sense of taste is fundamental for survival as harmful substances can be discriminated and prevented from

entering the body. Taste buds act as chemosensory sentinels and detect bitter, salty, sweet, sour, and umami

substances and transmit signals to afferent nerve fibers. Whether a single gustatory nerve fiber selectively is

responsive to a single taste modality (through taste receptor cell activation) is a point of contention in the field..

In the present study, we present a method for single cell RNA sequencing of gustatory geniculate ganglion

neurons and compare the results obtained to two prior published works. Additionally, independent reanalysis of

the raw data from these previous studies confirms molecular heterogeneity of ganglion neurons. Multiple

gustatory clusters are found, and we compare cluster markers identified by the original works and those

identified in the present study. Across all datasets and analyses, specific clusters show a high degree of

correlation including a somatosensory cluster (*Phox2b-*, *Piezo2+*, *Fxyd2+*), a potential sweet-best cluster

(*Phox2b+*, *Spon1+*, *Olfm3+*), and a potential sour-best cluster (*Phox2b+*, *Penk+*, *Htr3a+*). Additionally, a

putative mechanosensitive gustatory cluster with an unknown functional role is identified (*Phox2b+*, *Piezo2+*,

Calb1+). Other gustatory clusters (*Phox2b+*) are more varied across analyses, but are marked by *Olfm3*.

Which, if any, clusters comprise umami-best, bitter-best, or salty-best fibers will require further study.

Introduction:

Taste buds detect sapid molecules in the mouth and are the initiators of gustation. Each taste bud contains 50-100 mature taste cells, which are classified as three separate types. Type I cells are the least understood, but act as supporting cells and may transduce some portion of salty taste. Type II cells are well-understood and transduce signals in response to bitter, sweet, or umami stimuli. Type II cells communicate with afferent nerve fibers through release of ATP via non-traditional synaptic mechanisms (Finger et al., 2005; Huang et al., 2007; Romanov et al., 2007, 2018; Taruno et al., 2013). Type III cells form classical synapses with afferent nerve fibers and are responsible for sour and portions of salt taste detection. Due to the different nature of how Type II and Type III cells communicate with afferent fibers, it is likely that the innervating fibers have different molecular and physiological properties. Indeed, we have shown that afferent fibers that express the serotonin receptor 5-HT3A preferentially contact serotonergic Type III cells (Stratford et al., 2017). This subset of afferent fibers responds to exogenous serotonin as well as ATP while other gustatory fibers respond only to ATP (Larson et al., 2015). These data suggest at least 2 subpopulations of gustatory afferent nerve fibers.

We previously showed that the majority of geniculate ganglion neurons show excitatory responses to exogenously applied ATP and about 25% to serotonin via P2X2/P2X3 and 5HT3A receptors, respectively (Larson et al., 2015; Vandenbeuch et al., 2015). Older studies using patch clamp electrophysiology of rat gustatory geniculate ganglion neurons also showed excitatory responses to ACh, serotonin, substance P, and GABA in a small percentage of cells (King and Bradley, 2000; Koga and Bradley, 2000). These data suggest that multiple classes of neurons exist at the physiological level and that likely, these classes could be reflected at the molecular level.

Lingual taste fields are innervated by branches of the facial (cranial nerve VII) and glossopharyngeal nerve (cranial nerve IX). The anterior tongue receives input from the chorda tympani branch of the facial nerve. The cell bodies of the chorda tympani reside in the geniculate ganglion. In addition to the chorda tympani, this ganglion houses the cell bodies of the greater superficial petrosal and the nervus intermedius which receive gustatory input from the soft palate and somatosensory input from the external auditory meatus (Mtui et al., 2011).

Gustatory geniculate ganglion neurons are delineated by expression of the transcription factor, *Phox2b* (Dvoryanchikov et al., 2017; Ohman-Gault et al., 2017; Rios-Pilier and Krimm, 2019) which molecularly defines a neuron as gustatory or non-gustatory. The first study to characterize the transcriptome of single geniculate ganglion neurons utilized the Fluidigm C1 to profile 96 neurons from wildtype mice (Dvoryanchikov et al., 2017). They found about 2/3 of the cells express *Phox2b*, consistent with histological studies. Within the gustatory cells, three main clusters were identified, all of which expressed *Phox2b* and *P2rx3*. Other cluster markers included T1: *Olfm3*, *Itm2a*, *Hspb3*; T2: *Cd302*, *Kcnip1*, *Slc39a11*; T3: *Lypd1*, *Penk*, and *Trhr*. The authors then characterized the physiological response profiles of isolated ganglion neurons to exogenous ATP and 5-HT to match functional profiling with transcriptional profiling and demonstrated multiple physiological profiles that nicely matched expression of P2X and 5-HT receptor mRNA. This study was the first to characterize geniculate ganglion neurons at the molecular level using scRNA-seq.

More recently, Zhang et al. performed scRNA-sequence analysis on a larger cohort of geniculate

ganglion neurons (Zhang et al., 2019). They profiled over 400 neurons using the Cel-seq (Hashimshony et al.,

2012) method and identified 7 clusters. Based on cluster markers, they made multiple transgenic and knockout

mice to test the role of certain transcripts in taste detection. Molecular and physiological evidence is presented

for classes of sweet-best and sour-best neurons, marked predominantly by expression of *Spon1* and *Penk*

transcripts, respectively. Using a floxed Gcamp model, the authors recorded in vivo calcium signals in

geniculate ganglion neurons and showed that *Spon1* and *Penk* neurons selectively responded to sweet and

sour lingual stimuli, respectively. Knockout of *Cdh4* and *Cdh13* eliminated detection of umami and bitter

molecules, respectively, as assessed by two-bottle taste preference testing. Inhibition of *Egr2*-neurons using a

Egr2-cre/Tetanus toxin model eliminated detection of NaCl. However, it is unclear how protein products of

these transcripts are involved in taste signaling.

In the present study, we performed transcriptional profiling of geniculate ganglion neurons using an

alternative method. We show that the neurons we collected expressed many transcripts overlapping with

currently published datasets. Additionally, we performed a secondary analysis of published datasets from

Dvoryanchikov et al 2017 and Zhang et al 2019 to further explore geniculate ganglion neuron clusters in

attempt to understand more about the role of ganglion neurons in taste perception. Lastly, we show that

combining all three datasets preserves cell clustering using the new analysis methods in Seurat V3 (Butler et

al., 2018; Stuart et al., 2019).

Materials and Methods:

Animals

All animal procedures were conducted under a protocol approved by the University of Colorado Animal Care

and Use Committee. Phox2bcre mice were purchased from Jax (B6(Cg)-Tg(Phox2b-cre)3Jke/J; stock #

016223) and bred to Rosa26/tdTomato mice (B6.Cg-Gt(ROSA)26Sortm14(CAG-tdTomato)Hze/J; stock #

007914). Crossed mice of both gender were used between the ages of 8 and 24 weeks.

Immunofluorescence

Animals were anesthetized using urethane and transcardially perfused with 4% paraformaldehyde in

phosphate buffer (0.1M; PB) or euthanized with carbon dioxide. Geniculate ganglia and tongues were

extracted and fixed for 1 or 4 hours at room temperature before incubating in 20% sucrose in PB overnight at

4°C. Tissue was embedded and frozen in Optimal Cutting Temperature (OCT) and sectioned at a thickness of

12-16 μm using a cryostat. After > 24 hours at -20°C, slides were thawed, rehydrated with PBS, and blocked

using 2% normal donkey serum in blocking buffer (PBS plus 0.3% TritonX100, 1% BSA). Primary antibodies

were applied overnight at 4°C. After thorough washing fluorescent secondary antibodies were applied at room

temperature for 3 hours before counterstaining with DAPI and mounting with FluoroMount G (Southern

Biotech). Slides were imaged using a Leica SP8 using 20x (NA 0.75) and 63x (NA 1.4) oil-immersion

objectives.

Single cell preparation

Geniculate ganglia of three carbon dioxide-euthanized mice were rapidly extracted and placed in minimum essential medium with Earle's balanced salts (MEM/EBSS; Hyclone) supplemented with 1.25 mg/ml trypsin (Sigma-Aldrich) and 2.5 mg/ml collagenase A (Roche Diagnostics) for 30 minutes at 35°C. Digested ganglia were washed with MEM/EBSS, and gently triturated with a fire-polished glass pipette. The suspension was passed through a 50 μm filter and loaded as the top layer of a Percoll gradient (12.5 and 28%, Sigma-Aldrich; Malin et al., 2007). The gradient was spun at 1300x*g* for 10 minutes using a swinging bucket rotor. The cell pellet at the bottom of the tube was washed with MEM/EBSS before resuspending in 2 mL MEM/EBSS.

Single cell capture and RNA extraction

A QiaScout Microraft Array (Qiagen) was coated with poly-D-lysine (0.02 mg/ml) for 2 hours and washed with MEM/EBSS. The geniculate ganglion cell suspension was placed in the microraft array and the cells allowed to settle for 20 minutes. The microraft array was secured using a custom set up to the stage of an Olympus IX71 inverted microscope equipped with a 10X objective and the QiaScout instrument (Qiagen). Cells were visually inspected and microrafts containing a single tdTomato-positive (or negative in some cases) geniculate ganglion neuron were extracted and transferred to a PCR tube containing lysis buffer as part of the QIAseq FX Single Cell RNA Library Kit (Qiagen). Once in lysis buffer, cells were lysed by heating the tube to 95°C for 3 minutes followed by an infinite hold at 4°C until 24 individual samples were collected. The collection protocol was performed across two independent experiments, resulting in 48 individual sequencing libraries (42 tdTomato+, 6 tdTomato-).

Library preparation and sequencing

Sequencing libraries were prepared using the QIAseq FX Single Cell RNA Library Kit (Qiagen) according to manufacturer instructions. Cells were lysed, followed by gDNA removal, reverse transcription, ligation, and whole transcriptome amplification. After amplification, libraries were enzymatically fragmented (~300 bp fragments) followed by adapter ligation. Library size was evaluated using a DNA tapestation and concentration was determined using qPCR (QIAseq Library Quant Array). Libraries were pooled at equimolar concentration and sequenced using an Illumina MiSEQ to generate 2x151 bp reads.

Sequencing analysis

Quality and adapter trimming was performed using BBDuk (Bushnell). Salmon (v0.14.0) was used to quantify transcript expression using trimmed reads and the Ensembl GRCm38.p6 (version 92) transcriptome (Patro et al., 2017; Zerbino et al., 2018). Transcript expression was summarized at the gene level and input to R using TxImport (Soneson et al., 2016; R Core Team, 2019). A custom R script was used to visualize gene expression in the dataset. Samples were kept if expression of *Snap25* or *Uclh1* was > 5 TPM. Discarded samples were further interrogated for expression of neuron-specific genes, but they were rarely present.

Sequencing analysis of published datasets

Single cell sequencing FASTQ files were obtained from NCBI Gene Expression Omnibus (GSE102443

(Dvoryanchikov et al., 2017) and GSE135801 (Zhang et al., 2019)). Raw reads from Dvoryanchikov et al. were

filtered, trimmed, mapped, and summarized at the gene-level using the same tools as described above. A

processed gene count matrix was obtained from Zhang et al. Gene count matrices were used as the input for

Seurat (V3;(Butler et al., 2018; Stuart et al., 2019)) where samples were filtered, scaled, and clustered.

Meta analysis of datasets

Individual Seurat objects were imported to R and processed using Seurat following the "Multiple Dataset

Integration and Label Transfer" vignette available on the program website

(https://satijalab.org/seurat/v3.1/integration.html). Cluster markers of the integrated dataset were calculated

from the 'RNA' assay of the Seurat object using a logistic regression test.

Cluster marker correlation analysis

Cluster markers identified in Seurat were compared for overlapping markers across analyses using a custom R

script. Correlation for each comparison was calculated by dividing the number of overlapping markers by the

number of published cluster markers. See Table 1 of this manuscript and Table S2 of Zhang et al 2019 for

cluster markers used.

Data visualization

Data were visualized using the pheatmap package in R or with Seurat plotting features DimPlot and

DoHeatmap and FeaturePlot (Butler et al., 2018; Kolde, 2019; Stuart et al., 2019).

Results:

Characterization of Phox2b-cre mouse

To confirm that the *Phox2b*-cre driver faithfully drives CRE recombinase in PHOX2B-expressing cells,

geniculate ganglia and gustatory taste fields of *Phox2b*-cre/Rosa-tomato mice were examined using confocal

immunofluorescence microscopy. A subset of geniculate ganglion neurons expresses robust tdTomato

fluorescence consistent with previous reports (Ohman-Gault et al., 2017; Rios-Pilier and Krimm, 2019). The

majority of tdTomato-expressing neurons exhibited P2X3 immunoreactivity, and conversely, nearly all P2X3-

immunoreactive neurons exhibited tdTomato fluorescence (Figure 1). To confirm that tdTomato is being driven

in PHOX2B-expressing ganglion cells, sections were labeled with an antibody against PHOX2B. Indeed, every

tdTomato-expressing neuron showed positive nuclear labeling of the transcription factor (Figure 1).

Within taste papillae, tdTomato positive nerve fibers enter taste buds consistent with the *Phox2b* as a

marker of gustatory nerve fibers. Most tdTomato-expressing nerve fibers exhibit strong P2X3-immunoreactivity.

Additionally, these fibers are rarely outside of taste buds but when present are not immunoreactive for P2X3

(Figure 2)

scRNA-seq of Phox2b geniculate ganglion neurons

Single cell RNA sequencing was performed on manually isolated geniculate ganglion neurons (48 samples

from 6 total mice across 2 collection days; 40 tdTomato+, 6 tdTomato-, 2 blanks). Ganglion neurons were

enzymatically digested and plated on a Qiagen QiaScout Microraft array. Using the QiaScout device cells were

visually inspected for presence or absence of tdTomato, dislodged from the microraft array, and transferred to

cell lysis buffer where lysis was initiated followed by reverse transcription, whole transcriptome amplification,

and sequencing library preparation. Post processing of the sequencing data excluded samples that had < 50%

mapping rate, expression of *Snap25* or *Uchl1* < 5 TPM, and <2000 unique genes. Post processing resulted in

31 cells (17 tdTomato+, 4 tdTomato-). *Phox2b* expression correlated with tdTomato fluorescence (Figure 3)

and taste-related transcripts (*P2rx2, P2rx3*) were present in all *Phox2b*+ neurons. A subset of *Phox2b*+

neurons also expressed *Htr3a* (Figure 3). Additionally, genes such as *Penk and Spon1*, markers of sour-best

and sweet-best neurons (Zhang 2019) were identified in some neurons. Unfortunately, the low sample size

prevented use of unbiased clustering algorithms such as Seurat (Stuart et al., 2019).

Replication of published results

The first report of scRNA-seq on geniculate ganglion neurons used wildtype mice and the Fluidigm C1 platform

to perform expression profiling and demonstrated 4 cell clusters (3 gustatory, 1 non-gustatory). Reanalysis of

these 96 cells using Seurat reveals 6 clusters of cells (Figure 4A,B) including 1 somatosensory (Phox2b-) and

5 gustatory (Phox2b+) clusters. Cluster markers described by Dvoryanchikov et al. readily mark clusters in the

reanalyzed data (Figure 4C). Correlation analysis of cluster markers from the original study and the reanalyzed

data show strong concordance of data (Figure 4D). However, the main difference is that in the reanalyzed

data, three clusters (D1,D2,D3) correlate with the original TI cluster. We attribute the discrepancy among

cluster numbers likely arises due to 1) different clustering software/algorithms and 2) less conservative

clustering variables. Overall, we show that this independent analysis of data from Dvoryanchikov et al. yields

similar cell clustering results. See Tables 1 and 2 for specific cluster markers.

We next applied a similar processing pipeline to scRNA-seq data from Zhang et al 2019. The authors manually

collected over 800 *P2rx3*-tdTomato expressing geniculate ganglion cells and performed scRNA-sequencing

using the Cel-seq technology. Of the 454 neurons available from GEO: GSE135801, 372 had expression of

>2000 unique genes. Zhang et al originally reported 7 total clusters (labeled A-G). Gustatory clusters

(Phox2b+) were identified by *Spon1, Penk, Cdh4, Cdh13, or Egr2*. Reanalysis of these data using Seurat

reveals 7 molecularly distinct neuronal clusters (Z0-Z6) including those marked by *Spon1* (cluster Z2)*, Penk*

(cluster Z6)*, Cdh13* (cluster Z1)*, Egr2* (cluster Z3) and *Piezo2* (clusters Z0 and Z5; Figure 5B,C)*.* However,

Cdh4 was not detected as a significant marker and *Cdh13* was not significant by FDR-adjusted p value (see

Table 3 and 4 for a more comprehensive list of cluster markers). While these specific cluster markers were less

consistent between the original report and the reanalyzed data, pairwise correlation between original cluster

markers (A-G) and reanalyzed cluster markers (Z0-Z6) show that the reanalyzed clusters show overall

agreement with the original report (Figure 5D). Clusters Z1 and Z4 do not strongly correlate with any of the

clusters from the original report. Likewise, clusters A and B do not correlate strongly with any clusters identified

in the reanalyzed data. Interestingly, cells belonging to cluster Z4 had the least unique genes detected per cell,

the least number of reads per cell, and the fewest significant cluster markers (not shown). Thus cluster Z4

could represent a cluster of unhealthy cells and/or poorly sequenced cells. Interestingly, when the data are

reprocessed after removing cells with less than 4000 unique genes detected (as opposed to 2000), cluster Z4

disappears with overall preservation of the other clusters (not shown).

Comparison across datasets

Overall, independent reanalysis of both datasets using a parallel analysis pipeline agrees with the original

reports. We next asked whether clusters from one dataset correlate to the other dataset. First, we examined

how top cluster markers identified by the original reports correlate with cluster markers identified during

reanalysis of the other dataset (ie do the cluster markers identified by Zhang et al 2019 correlate with cluster

markers identified by reanalysis of Dvoryanchikov et al 2017?). Cluster markers identified by Zhang et al 2019

correlate strongly with some, but not all, markers identified by reanalysis of Dvoryanchikov et al 2017 (Figure

6A). For instance, clusters D0, D2, D4, and D5 from reanalysis of Dvoryanchikov et al 2017 correlate strongly

with clusters F, D, G, and E of Zhang et al 2019, respectively. Conversely, cluster markers identified by

Dvoryanchikov et al 2017 also correlate strongly with most markers identified by reanalysis of Zhang et al 2019

(Figure 6B). Clusters Z0, Z1, Z2, Z5, and Z6 from reanalysis of Zhang et al 2019 correlate strongly with

clusters somatosensory, TI, TI, TII, and TIII of Dvoryanchikov et al 2017, respectively. From these

comparisons, we can conclude with higher confidence that certain clusters are present and valid as they are

seen across both datasets and contain a high proportion of overlapping markers identified by all analysis. The

clusters identified in the original reports that we have the most confidence in (in descending order) are the

somatosensory cluster (SS, F, D0, Z0), *Penk*-expressing gustatory cluster (TIII, E, D5, Z6), *Calb1*-expressing

gustatory cluster (TII, G, D4, Z5), *Spon1*-expressing gustatory cluster (TI, D, D2, Z2), and *Pxmp2*-expressing

gustatory cluster (TI, B, Z1, D1). The remaining clusters show less correlation to the original reports.

Next, we compared the cluster markers identified by independent reanalysis of both datasets to find

overlapping markers. Indeed, there was a degree of concordance amongst the clusters (Figure 6C). We

conclude that 5 clusters from each dataset share many overlapping markers and thus are likely the same

identity.

Meta analysis of all datasets

Generally, in RNA-sequence analysis, it is against best-practices to combine datasets from multiple sources

given the range of variability in sample preparation, sequencing, and data processing. However, with the

recent release of Seurat V3 a method for combining datasets from single cell sequencing experiments across

multiple technologies has been developed (Stuart et al., 2019). This method finds integration anchors common

amongst the datasets, merges the data, and scales the data based on the anchors. This meta analysis allows
for integration of data on similar cell types from multiple sources.

The three datasets in this paper can be combined, scaled, and clustered together. Interestingly, UMAP
plotting of the integrated data shows a strong similarity among the three datasets, with cells from each dataset
being represented throughout the plot (Figure 7A,B). This suggests that despite three different methods and
technologies used to collect scRNA-seq data, the genetic information was similar between studies. Clustering
of the integrated data reveals 7 cell clusters. These clusters largely share markers identified by both the
original reports and the individually reanalyzed datasets. With the exception of cluster M4, all identified clusters
from the combined dataset show overlapping markers with clusters from the separate datasets (Figure 7F,G,
Table 6). For reasons discussed above, cluster M4 likely represents a population of unhealth of poorly
sequenced cells. When the datasets are reintegrated after removing cells with less than 4000 unique genes
detected, cluster M4 disappears with overall preservation of the other clusters (not shown).

Discussion:
Do neuron clusters represent specific taste modalities?

Single cell RNA-seq allows for the determination of heterogeneity within a cell population. By clustering
cells based on similarities, markers that are unique to those clusters can be identified, which can give strong
insight to their function. While Dvoryanchikov et al. were the first to reveal molecular heterogeneity by
identifying clusters of geniculate ganglion neurons, Zhang et al. were the first to attribute the role of specific cell
clusters to taste detection. Convincing evidence demonstrates that neurons in the *Penk* and *Spon1-*
expressing cell clusters represent sour- and sweet-best neurons. Thus, it is plausible that these neurons
receive specific input from Type III and Type II taste receptor cells, respectively.

Sour and sweet: All three datasets analyzed in the present study showed expression of *Penk* and
Spon1 in gustatory fibers and that they marked specific cell clusters. In Zhang et al 2019, *Penk* and *Spon1*
were cluster markers. Dvoryanchikov et al 2017 reported *Penk* as a cluster marker, but not *Spon1*.
Interestingly, secondary analysis reveals that *Spon1* neurons were a subset of the cluster labeled 'T1' in the
original report (Dvoryanchikov et al. 2017), which was subdivided in to three clusters in this current study
(Figure 5C). These findings together with the in vivo imaging of Zhang et al 2019 strongly suggest that sour-
best and sweet-best fibers are molecularly distinct from other ganglion neurons.

Bitter, salty, and umami: Zhang et al reports that *Cdh4*, *Egr2*, and *Cdh13* are cluster markers for bitter,
salty, and umami-best neurons. However, in the present analyses, these markers did not define clusters as
well as *Penk* and *Spon1*. Additionally, the behavioral assays performed by Zhang et al relied on constitutive
knockout of *Cdh4* and *Cdh13* and tetanus toxin inactivation of *Egr2* neurons. While the experiments showed
that umami, bitter, and salty taste preference rely on these transcripts, they do not show that this requirement
is at the level of the geniculate ganglion. Indeed, according to the Allen Brain Map, these three transcripts are
expressed in a variety of cell types throughout the central nervous system. Additionally, according to
secondary analysis of Qin et al, *Cdh13* is expressed in both Type II and Type III taste receptor cells and *Egr2*

is expressed in some Type III cells (not shown; Qin et al., 2018). Thus, any ganglion cell clusters responsible

for the detection of bitter, salty, and umami signals from taste receptor cells remains enigmatic.

Does Htr3a mark a specific cluster?

Stratford et al 2017 previously showed that nerve fibers expressing the serotonin receptor 5-HT3

preferentially contact Type III taste receptor cells in all taste fields (Stratford et al 2017) suggesting that these

fibers may be molecularly distinct within the geniculate ganglion. However, Dvoryanchikov et al. argued that

Htr3a did not delineate a specific cluster in their analyses. In the present analyses, all three datasets show

clustering of *Htr3a*-expressing neurons. While *Htr3a* was in multiple clusters, it had strongest expression in the

cluster marked by *Penk* – the sour-best fibers which support our findings that these cells wire to sour-sensing

Type III taste receptor cells. Conversely, *Htr3a* is absent in the *Spon1* cluster (sweet-best fibers). Interestingly,

in all datasets, *Htr3a* is also present in a gustatory (*Phox2b*+) cluster expressing *Piezo2*, a transcript for a

mechanosensitive channel. This cluster expressed low levels of *P2rx3*, and no *P2rx2*. Moayedi et al 2018

showed that *Piezo2*-GFP expressing nerve fibers are present in fungiform papillae, just adjacent to taste buds,

and that these fibers enter the papillae in a bundle with other gustatory fibers (Moayedi et al., 2018).

Interestingly, P2X2/3 double knockout mice (which lack nerve response to all tastants) still show nerve

responses to mechanical stimulation of the tongue (Finger et al., 2005). Perhaps this *Piezo2*/*Htr3a*/*Phox2b*

neuronal cluster is responsible for this ATP-independent response.

Uncorrelated clusters

We report that across datasets, there are some clusters that do not correlate with any other clusters (primarily

Z3, and Z4, and D3; Figure 6C). One possibility, as discussed above, is that some of these clusters could

represent unhealthy or poorly sequenced cells. This is primarily the case with cluster Z4. For the remaining

clusters, it could be that the sample size of the studies may not be large enough to encompass all neuron

clusters. It could be that each uncorrelated cluster represents a different cluster of cells. Thus, larger

geniculate ganglion scRNA-seq databases must be obtained.

Conclusions

Overall, the data analyzed in this paper confirm molecular heterogeneity among the neurons of the geniculate

ganglion. The datasets clearly separate gustatory vs non-gustatory neurons by expression of *Phox2b*. Within

the *Phox2b* expressing fibers, clear sub clusters are formed. Consistent among the datasets are clusters

defined by *Penk*, *Spon1*, and *Piezo2*. Zhang et al showed that *Penk* clusters represent sour-best fibers while

Spon1 clusters represent sweet-best fibers. The *Piezo2* cluster could be responsible for residual chorda

tympani nerve responses to lingual mechanical stimuli in the absence of purinergic signaling (Finger et al.

2005). The remaining clusters were more variable between the datasets and markers identified by Zhang et al

(*Cdh4*, *Cdh13*, *Egr2*) were not clearly observed in this study. While it is likely that neurons responsible for

detecting each taste modality are molecularly distinct, more work needs to be done to elucidate the molecular

phenotypes of bitter, salty, and umami sensitive fibers.

Figure Legends:

Figure 1. tdTomato expression in PHOX2B-immunoreactive neurons. Sections of Phox2bcre/tdTomato

geniculate ganglion labeled with antibodies against PHOX2B and P2X3. Nearly all tdTomato-expressing cells

are immunoreactive for PHOX2B and P2X3. Non-nuclear "green" signal is due to binding of anti-mouse

secondary antibodies to endogenous mouse IgG; this signal persists in no primary controls ,and no nuclear

signal is observed. Sections are maximal z-projections of ~12 μm image stacks.

Figure 2. tdTomato-expressing nerve fibers enter taste buds. Sections of lingual taste papillae showing

immunoreactivity to GNAT3 (α-gustducin; Type II cells) and P2X3 with endogenous tdTomato fluorescence.

Within taste buds, all tdTomato+ fibers exhibit P2X3 immunoreactivity. In some cases, tdTomato+ fibers were

found in the intergemmal area and were P2X3 negative. Images are maximal z-projections of ~12-16 μm

image stacks.

Figure 3. Expression of select transcripts in tdTomato-expressing neurons. Geniculate ganglion neurons were

collected using the QiaScout and processed for single cell RNA sequencing using the Qiagen FX single cell

RNA kit. Heatmap shows log(transcripts per million) of select transcripts. *Phox2b* expression correlates with

presence or absence of tdTomato.

Figure 4. Secondary analysis of Dvoryanchikov et al. 2017 scRNA-seq data. Analysis of cells publicly available

from GSE102443. A. Seurat UMAP clustering of cells expressing greater than 2000 genes. B. Heatmap

showing transcript expression of top 5 cluster markers identified using Seurat. C. Heatmap showing transcript

expression of cluster markers described in Dvoryanchikov et al. 2017. Note, that the Dvoryanchikov et al 2017

cluster T1 is further subdivided into 3 sub-clusters identified here as D1, D2, and D3. D. Correlation of cluster

markers described by Dvoryanchikov et al. 2017 and those identified in this study using Seurat.

Figure 5. Secondary analysis of Zhang et al. 2019 scRNA-seq data. Analysis of cells publicly available from

GSE135801. A. Seurat UMAP clustering of cells expressing greater than 2000 genes. B. Heatmap showing

transcript expression of top 5 cluster markers identified using Seurat. C. Heatmap showing transcript

expression of cluster markers described in Zhang et al. 2019. D. Correlation of cluster markers described by

Zhang et al. 20179 and those identified in this study using Seurat.

Figure 6. Comparison of published cluster markers and cluster markers identified from reanalysis of both

datasets. A. Heatmap showing reanalyzed data from Dvoryanchikov et al 2017 with markers identified in the

original report of Zhang et al 2019. Correlation heatmap shows pairwise comparisons between cluster markers.

B. Heatmap showing reanalyzed data from Zhang et al 2019 with markers identified in the original report of

Dvoryanchikov et al 2017. Correlation heatmapshows pairwise comparisons between cluster markers. C.

Correlation heatmap of cluster markers from both reanalyzed datasets

Figure 7. Combination of data from this paper, Dvoryanchikov et al 2017, and Zhang et al 2019. A. Seurat

UMAP clustering showing each dataset. B. Overall clusters of the combined datasets. C. Heatmap showing

transcript expression of top 5 markers of each cluster. D,E. Heatmaps of combined dataset analysis showing

expression of cluster markers identified in the original report of Dvoryanchikov et al 2017 (D) and Zhang et al

2019 (E). F,G. Correlation heatmap of cluster markers from the combined dataset analysis versus cluster

markers identified by reanalysis of Dvoryanchikov et al 2017 (F) or Zhang et al 2019 (G).

Table S1. Seurat-identified cluster markers for Zhang et al 2018.

Table S2. Seurat-identified cluster markers for Dvoryanchikov et al 2017.

Table S3. Seurat-identified cluster markers for meta analysis.

Conflict of Interest:

The authors have no financial conflicts of interest to declare.

Funding:

This work was supported by NIH/NIDCD R21DC015115-01 to Sue C. Kinnamon (University of Colorado

Anschutz Medical Campus).

Acknowledgments:

We thank Qiagen for allowing us to demo the QiaScout device and supplying samples of the QiaScout

Microraft Arrays. We also thank Drs. Tom Finger and Sue Kinnamon for their support and critique of the

manuscript.

Table 1. Comparison of cluster markers of Dvoryanchikov et al

Dvoryanchikov cluster	Gene	Seurat cluster	Confirmed marker	p <0.05, FDR-adjusted p > 0.05
Somatosensory	Fxyd2	D0	yes	
Phox2b-	Kcns3	D0	yes	
	Prrxl1	D0	yes	
	Ldb2	D0	yes	
	Pdlim1	D0	yes	
	Mef2c	D0	yes	
TI	Olfm3	D1,D2,D3	yes	* (D1,D2,D3)
Phox2b+	Itm2a	D2	yes	D2
	Hspb3	na	no	
	Foxg1	D1,D2	Yes	* (D1,D2)
	Irf6	D2,D3	yes	* (D2,D3)
	Eya2	D3	yes	
TII	Cd302	D4	yes	*
Phox2b+	Kcnip1	D4	Yes	*
	Slc38a11	D4	yes	
	Mafb	D4	yes	*
	Meis1	D4	Yes	*
	Tbx2	-	no	
	Dach2	D4	Yes	
	Runx3	-	No	
TIII	Lypd1	D5	yes	
Phox2b+	Penk	D5	yes	
	Trhr	D5	yes	
	Prox2	-	no	
	Phox2a	-	no	
	Twist2	-	no	

Table 2. Confirmation of cluster markers identified by Dvoryanchikov et al. 2017. Seurat cluster represents the cluster each marker was found in the reanalyzed data. Confirmed cluster is whether or not the cluster marker was found in the most-correlated cluster. Some markers were significant by p-value but not FDR-adjusted p-value, as denoted in the final column.

Table 2. Top Cluster markers from current analysis of Dvoryanchikov et al

cluster					
Phox2b-	Phox2b+				
D0	D1	D2	D3	D4	D5
Tafa3	Fam122b	Angpt4	Eya2	Mme	P2ry12
Hs6st2	Fxyd7	Naalad2	Ligp1	Dach2	Creg2
Antxr2	Pxmp2	Plpp4	Pcdh9	Ibsp	Pdlim3
Mob3b	Unc5c	Ptprk	Pon2	Draxin	Fstl4
Prmt8	Rassf9	Gm17950	Nnat	Col11a1	Rab3b
Hoxd1	Bambi	Kcnk2	Plxdc2	Slc38a11	Trhr
Nrp1	Thsd7b	Trappc3l	Gnas	Kcnip1	Htr3a
Ldb2	Cdh11	Spon1	Caprin1	Meis1	Sorbs2
Prrxl1	P2rx2	Flrt2	Cav1	Kif21b	Rgs17
Syt6	Foxg1	Fkbp11	Cd34	Hs3st2	Trim25

Table 2. Top cluster markers from reanalysis of Dvoryanchikov et al 2017. Markers ordered by ascending FDR-adjusted p value

Table 3. Comparison of cluster markers of Zhang et al

Zhang et al cluster	Gene	Seurat cluster	Confirmed marker?	p <0.05, FDR-adjusted p > 0.05
Somatosensory-F	Piezo2	Z0,Z5	yes	
Phox2b-	Pou2f4	na	no	
	Nrp1	Z0	yes	
	Fxyd2	Z0	yes	
	Pde1c	Z0	yes	
Somatosensory-G	Calb1	Z5	yes	
Phox2b+	Mafb	Z0,Z5	yes	
	Lrrc49	Z5	Yes	
	Pthlh	Z5	yes	
	Gfra2	Z5	Yes	*
Sweet-best- D	Spon1	Z2	yes	
Phox2b+	Itm2a	Z2	yes	
	Kcnk2	Z2	yes	
	Flrt2	Z2	yes	
	Adcyap1	Z2	yes	
Umami-best-A	Cdh4	na	no	
Phox2b+	Mdfic	Z2	no	
	Kcnip4	na	no	
	Nmb	na	no	
	Apold1	na	no	
Bitter-best- B	Cdh13	Z1	Yes	*
Phox2b+	Pxmp2	Z1,Z3	Yes	* (Z3)
	Fat1	na	No	
	Eya2	Z3	No	*
	Net1	na	No	
Salty-best-C	Egr2	Z3	yes	
Phox2b+	Sepp1	Z3	Yes	*
	Ednrb	Z3	Yes	
	Ncmap	na	No	
	Cyr61	Z3	Yes	*
Sour-best-E	Penk	Z6	yes	
Phox2b+	Lypd1	Z6	yes	
	Prlr	Z6	yes	
	Trhr	Z6	Yes	
	Cplx2	Z6	Yes	

Table 3. Confirmation of cluster markers identified by Zhang et al. 2019. Seurat cluster represents the cluster each marker was found in the reanalyzed data. Confirmed cluster is whether or not the cluster marker was found in the most-correlated cluster. Some markers were significant by p-value but not FDR-adjusted p-value, as denoted in the final column.

Table 4. Top Cluster markers from current analysis of Zhang et al data

Cluster						
Phox2b-	Phox2b+					
Z0	Z1	Z2	Z3	Z4	Z5	Z6
Kcnip3	P2rx2	Spon1	Ndrg1	Dnajc3	Runx3	Htr3a
Cadps2	Gabrb2	Kcnk2	Col5a1	Wdr13	Draxin	Trhr
Nrp1	P2rx3	Nbl1	Rttn	Tex2	Gabra4	Irx5
Lxn	Al504432	Prkcb	Mmd2	Pir	Spp1	Ncald
Rarres1	Thsd7b	Olfm3	Fbln2	Pik3c3	Cd302	Igf1
Pde1c	Pxmp2	Flrt2	Plxdc2	Zfr	Calb1	Cplx2
Prkar1a	Elmo1	Astn2	Pou3f1	2010012O05Rik	Dach2	Lypd1
Pla2g7	Atp2b4	S100a6	Gpm6b	Ap2b1	Rasgrp2	Creg2
Kcnj4	Efr3a	Fkbp11	Ednrb	Srrm2	Hs3st2	Penk
Acpp	Foxg1	Ifi27l2a	Cubn	Dcun1d1	Garem	Pdlim3

Table 4. Top cluster markers from reanalysis of Zhang et al 2018. Markers ordered by ascending FDR-adjusted p value

420 **Table 5. Comparison across datasets**

Cluster from combined analysis	corresponding cluster from Zhang et al	Corresponding cluster from Dvoryanchikov et al.	Corresponding cluster from Zhang This analysis	Corresponding cluster from Dvoryanchikov This analysis	Potential Function
M0	F-Somatosensory (Piezo2)	Somatosensory	Z0	D0	Somatosensory
M1	B- Cdh13/pxmp2 cluster	TI	Z1	D1	
M2	D- Spon1 cluster	TI	Z2	D2	Sweet-best
M3	*B- Cdh13/pxmp2 cluster*	*TI*	*Z3*	*D3*	
M4			*Z4*		*Poor quality cells?*
M5	G-Somatosensory2 (Piezo2)	TII	Z5	D4	Somatosensory?
M6	E- Penk cluster	TIII	Z6	D5	Sour-best

424 **Table 6. Top Cluster Markers combined dataset**

Cluster						
Phox2b-	Phox2b+					
M0	M1	M2	M3	M4	M5	M6
Fxyd2	P2rx2	Spon1	Pcdh9	App	Hopx	Lypd1
Rgs4	P2rx3	Prkcb	Igf1	Mlf2	Map7d2	Htr3a
Cadps2	Elmo1	Olfm3	Eya2	Cldn25	Calb1	Trhr
Sdcbp	Fxyd7	Tac1	Gnas	Mapk8ip3	S100b	Penk
Prkar1a	Cdh11	Nbl1	Nnat	Rpsa	AI593442	Prlr
Pde1c	Gabrb2	B3gnt2	Lmo2	Skint3	Piezo2	Cplx2
Kcns3	Unc5c	Mctp1	Tmem132b	Epdr1	Asic3	Rab3b
Tmsb4x	Gabra1	Kcnk2	Dnajb4	Mpz	Lrrc49	Ncald
Pla2g7	Efr3a	Itm2a	Cav1	Eif1	Cd302	Pdlim3
Nxpe2	Thsd7b	Flrt2	Serpinb1a	Sh3bp5l	Nefh	Cnr1

References:

Bushnell, B. BBMap.

Butler, A., Hoffman, P., Smibert, P., Papalexi, E., and Satija, R. 2018. Integrating single-cell transcriptomic data across different conditions, technologies, and species. Nature Biotechnology. 36:411–420.

Dvoryanchikov, G., Hernandez, D., Roebber, J.K., Hill, D.L., Roper, S.D., and Chaudhari, N. 2017. Transcriptomes and neurotransmitter profiles of classes of gustatory and somatosensory neurons in the geniculate ganglion. Nature Communications. 8:760.

Finger, T.E., Danilova, V., Barrows, J., Bartel, D.L., Vigers, A.J., Stone, L., Hellekant, G., and Kinnamon, S.C. 2005. ATP Signaling Is Crucial for Communication from Taste Buds to Gustatory Nerves. Science. 310:1495–1499.

Hashimshony, T., Wagner, F., Sher, N., and Yanai, I. 2012. CEL-Seq: Single-Cell RNA-Seq by Multiplexed Linear Amplification. Cell Reports. 2:666–673.

Huang, Y.-J., Maruyama, Y., Dvoryanchikov, G., Pereira, E., Chaudhari, N., and Roper, S.D. 2007. The role of pannexin 1 hemichannels in ATP release and cell–cell communication in mouse taste buds. PNAS. 104:6436–6441.

King, M.S., and Bradley, R.M. 2000. Biophysical properties and responses to glutamate receptor agonists of identified subpopulations of rat geniculate ganglion neurons. Brain Research. 866:237–246.

Koga, T., and Bradley, R.M. 2000. Biophysical Properties and Responses to Neurotransmitters of Petrosal and Geniculate Ganglion Neurons Innervating the Tongue. J Neurophysiol. 84:1404–1413.

Kolde, R. 2019. pheatmap: Pretty Heatmaps.

Larson, E.D., Vandenbeuch, A., Voigt, A., Meyerhof, W., Kinnamon, S.C., and Finger, T.E. 2015. The Role of 5-HT3 Receptors in Signaling from Taste Buds to Nerves. J Neurosci. 35:15984–15995.

Malin, S.A., Davis, B.M., and Molliver, D.C. 2007. Production of dissociated sensory neuron cultures and considerations for their use in studying neuronal function and plasticity. Nat Protocols. 2:152–160.

Moayedi, Y., Duenas-Bianchi, L.F., and Lumpkin, E.A. 2018. Somatosensory innervation of the oral mucosa of adult and aging mice. Sci Rep. 8:1–14.

Mtui, E., Gruener, G., and FitzGerald, M. 2011. Clinical Neuroanatomy and Neuroscience - 6th Edition.

Ohman-Gault, L., Huang, T., and Krimm, R. 2017. The transcription factor Phox2b distinguishes between oral and non-oral sensory neurons in the geniculate ganglion. J Comp Neurol. 525:3935–3950.

Patro, R., Duggal, G., Love, M.I., Irizarry, R.A., and Kingsford, C. 2017. Salmon: fast and bias-aware quantification of transcript expression using dual-phase inference. Nat Methods. 14:417–419.

Qin, Y., Sukumaran, S.K., Jyotaki, M., Redding, K., Jiang, P., and Margolskee, R.F. 2018. Gli3 is a negative regulator of Tas1r3-expressing taste cells. PLOS Genetics. 14:e1007058.

R Core Team. 2019. R: A Language and Environment for Statistical Computing. Vienna, Austria: R Foundation for Statistical Computing.

Rios-Pilier, J., and Krimm, R.F. 2019. TrkB expression and dependence divides gustatory neurons into three
subpopulations. Neural Dev. 14:3.

Romanov, R.A., Lasher, R.S., High, B., Savidge, L.E., Lawson, A., Rogachevskaja, O.A., Zhao, H.,
Rogachevsky, V.V., Bystrova, M.F., Churbanov, G.D., et al. 2018. Chemical synapses without synaptic
vesicles: Purinergic neurotransmission through a CALHM1 channel-mitochondrial signaling complex. Sci
Signal. 11:eaao1815.

Romanov, R.A., Rogachevskaja, O.A., Bystrova, M.F., Jiang, P., Margolskee, R.F., and Kolesnikov, S.S. 2007.
Afferent neurotransmission mediated by hemichannels in mammalian taste cells. The EMBO Journal. 26:657–
667.

Soneson, C., Love, M.I., and Robinson, M.D. 2016. Differential analyses for RNA-seq: transcript-level
estimates improve gene-level inferences. F1000Res. 4.

Stratford, J., Larson, E., Yang, R., Salcedo, E., and Finger, T. 2017. 5-HT3a-driven GFP delineates Gustatory
Fibers innervating Sour-responsive Taste Cells: A Labeled Line for Sour Taste? J Comp Neurol. 525:2358–
2375.

Stuart, T., Butler, A., Hoffman, P., Hafemeister, C., Papalexi, E., Mauck, W.M., Hao, Y., Stoeckius, M.,
Smibert, P., and Satija, R. 2019. Comprehensive Integration of Single-Cell Data. Cell. 177:1888-1902.e21.

Taruno, A., Vingtdeux, V., Ohmoto, M., Ma, Z., Dvoryanchikov, G., Li, A., Adrien, L., Zhao, H., Leung, S.,
Abernethy, M., et al. 2013. CALHM1 ion channel mediates purinergic neurotransmission of sweet, bitter and
umami tastes. Nature. 495:223–226.

Vandenbeuch, A., Larson, E.D., Anderson, C.B., Smith, S.A., Ford, A.P., Finger, T.E., and Kinnamon, S.C.
2015. Postsynaptic P2X3-containing receptors in gustatory nerve fibres mediate responses to all taste qualities
in mice. J Physiol. 593:1113–1125.

Zerbino, D.R., Achuthan, P., Akanni, W., Amode, M.R., Barrell, D., Bhai, J., Billis, K., Cummins, C., Gall, A.,
Girón, C.G., et al. 2018. Ensembl 2018. Nucleic Acids Res. 46:D754–D761.

Zhang, J., Jin, H., Zhang, W., Ding, C., O'Keeffe, S., Ye, M., and Zuker, C.S. 2019. Sour Sensing from the
Tongue to the Brain. Cell.

PHOX2B/**P2X3**   **PHOX2B**/**tdTomato**   **Merge**

100 μm

100 μm

Figure 1

GNAT3/P2X3 **GNAT3/tdTomato** **Merge**

Figure 2

Figure 3

Figure 4

Figure 5

A

Clusters from analysis of Dvoryanchikov

Clusters from analysis of Dvoryanchikov

markers/clusters identified by Zhang et al.

clusters identified by Zhang et al.

B

Clusters from analysis of Zhang

Clusters from analysis of Zhang

markers/clusters identified by Dvoryanchikov et al.

clusters identified by Dvoryanchikov et al.

C

Clusters from analysis of Zhang

Clusters from analysis of Dvoryanchikov