Quantitative properties of a feedback circuit predict frequency-dependent pattern

separation

Running Title: Feedback inhibition in dentate gyrus

Oliver Braganza[1*], Daniel Müller-Komorowska[1,2], Tony Kelly[1] and Heinz Beck[1,3 *]

[1]Institute for Experimental Epileptology and Cognition Research, University of Bonn

[2]International Max Planck Research School for Brain and Behavior, University of Bonn

[3]Deutsches Zentrum für Neurodegenerative Erkrankungen e.V., Bonn, Germany

* Correspondence should be addressed to:

Heinz Beck

Institute for Experimental Epileptology and Cognition Research, University of Bonn

Venusberg-Campus 1

53127 Bonn

Tel.: 0228 6885 270

Fax: 0228 6885 294

e-mail: Heinz.beck@ukbonn.de

Oliver Braganza (Lead Contact)

Institute for Experimental Epileptology and Cognition Research, University of Bonn

Sigmund-Freud Str. 25

53105 Bonn

Tel.: 0228 6885 157

Fax: 0228 6885 294

e-mail: oliver.braganza@ukbonn.de

Pages: 67

Figures: 6 (+11 Suppl.)

Tables: (2 Suppl.)

Abstract: 159 words; Introduction: 475 words; Results: 4366 words; Discussion: 1207 words

     ## Abstract

Feedback inhibitory motifs are thought to be important for pattern separation across species.

How feedback circuits may implement pattern separation of biologically plausible, temporally

structured input in mammals is, however, poorly understood. We have quantitatively

determined key properties of *net* feedback inhibition in the mouse dentate gyrus, a region

critically involved in pattern separation. Feedback inhibition is recruited steeply with a low

dynamic range (0 to 4% of active GCs), and with a non-uniform spatial profile. Additionally,

net feedback inhibition shows frequency-dependent facilitation, driven by strongly facilitating

mossy fiber inputs. Computational analyses show a significant contribution of the feedback

circuit to pattern separation of theta modulated inputs, even within individual theta cycles.

Moreover, pattern separation was selectively boosted at gamma frequencies, in particular for

highly similar inputs. This effect was highly robust, suggesting that frequency dependent

pattern separation is a key feature of the feedback inhibitory microcircuit.

Introduction

Efficiently discriminating similar percepts or experiences is a central capability common to invertebrate and vertebrate species. In general terms, such discrimination can be achieved by decreasing the overlap in representations by neuronal ensembles between input and output patterns, a process termed 'pattern separation' (Cayco-Gajic and Silver, 2019; Marr, 1971; McNaughton and Morris, 1987; Rolls, 2013). Numerous studies have proposed cellular and circuit mechanisms that support this computation. For instance, sparse divergent inputs, specialized intrinsic properties and feedforward inhibition are thought to generally contribute (Cayco-Gajic et al., 2017; Cayco-Gajic and Silver, 2019; Krueppel, 2010; Mircheva et al., 2019). Another common feature of most of these models and experimental studies is a critical role of feedback inhibition (Cayco-Gajic et al., 2017; Rolls, 2013). Feedback circuits differ from the above mechanisms in that they can i) implement direct competition between active cells through lateral inhibition and ii) integrate information about the actual global activity level in a population allowing efficient normalization (Braganza, 2018; Wick et al., 2010; Wiechert et al., 2010). Indeed, in the insect olfactory system a critical role of such a circuit has been causally demonstrated (Lin et al., 2014; Papadopoulou et al., 2011)

In mammals, substantial evidence points towards a role of the hippocampal dentate gyrus (DG) for pattern separation during memory formation and spatial discrimination (Bakker et al., 2008; Berron et al., 2016; Gilbert et al., 2001; Leal and Yassa, 2018; Leutgeb et al., 2007; McHugh et al., 2007; Neunuebel and Knierim, 2014; Stefanelli et al., 2016; van Dijk and Fenton, 2018). The DG is thought to subserve this task by converting different types of inputs to sparse, non-overlapping activity patterns of granule cells (GCs). However, in contrast to the insect olfactory system, the DG feedback circuit is extremely complex, comprising numerous interconnected interneuron types (Supplementary Table 1)(Bartos et al., 2002; Dasgupta and Sikdar, 2015; Espinoza et al., 2018; Ewell and Jones, 2010; Freund and Buzsáki, 1996; Geiger et al., 1997; Harney and Jones, 2002; Hefft and Jonas, 2005; Kraushaar and Jonas, 2000; Larimer and Strowbridge, 2008; Lee et al., 2016; Liu et al.,

2014; Lysetskiy et al., 2005; Sambandan et al., 2010; Savanthrapadian et al., 2014; Sik et

al., 1997; Yu et al., 2015; Yuan et al., 2017; Zhang et al., 2009). For instance, interneurons

subserving feedback inhibition are also incorporated into circuits mediating feedforward

inhibition (Ewell and Jones, 2010; Hsu et al., 2015; Lee et al., 2016) and disinhibition

(Savanthrapadian et al., 2014; Yuan et al., 2017). This makes it difficult to predict the *net*

inhibition arising from GC activity.

We reasoned that to assess if feedback inhibition is indeed suitable for the purpose of

pattern separation in the DG, it is necessary to determine how efficiently the activity of

sparse GC ensembles recruits *net* inhibition, i.e. the dynamic range and gain of the feedback

inhibitory microcircuit. It is furthermore necessary to quantify the spatial and temporal

properties of the elicited inhibition, in order to investigate its impact on biologically plausible,

temporally structured input. For instance, the DG shows prominent theta oscillations during

exploration and distinctive slow-gamma activity during associative memory encoding (Hsiao

et al., 2016; Pernía-Andrade and Jonas, 2014; Sasaki et al., 2018; Trimper et al., 2017).

Importantly, both sparsity and temporal oscillations will critically affect a proposed pattern

separation function. For instance, feedback inhibition must by definition occur with a delay, a

property frequently abstracted away in computational models (Myers and Scharfman, 2009;

Rolls, 2016), but potentially critical during oscillatory activity.

Here, we combine patch-clamp recordings, multiphoton imaging and optogenetics to provide

a first quantitative, empirical description of the *net* input-output function of a feedback

inhibitory microcircuit. This includes the spatiotemporal organization of *net* feedback

inhibition elicited by a spatially restricted GC population and the *net* short term dynamics

within the feedback microcircuit. Finally, we integrate our data into a biophysically realistic

computational model and probe its ability to perform pattern separation. We find a moderate

feedback inhibition mediated pattern separation effect during theta modulated input but a

substantial separation, particularly of highly similar inputs, during gamma oscillations.

Results

Input-output relation of the feedback inhibitory microcircuit

We reasoned that the ultimately relevant parameter for the putative pattern separation effect

of feedback inhibition is the *net* inhibition arriving at GCs. We therefore treated the feedback

microcircuit as a black-box striving to relate only its *net* input (fraction of GCs active) to its *net*

output (feedback inhibition in GCs). To this end we antidromically recruited feedback

inhibitory circuits, while simultaneously recording GC inhibition and population activity (see

schematic in **Fig. 1A**). Electrical stimulation reliably evoked graded IPSCs in dentate GCs,

that increased with stimulation strength (maximal amplitude of 324.1±99.2 pA, n=8; **Fig. 1B**).

Feedback IPSCs were completely blocked by 10µM GABAzine (to 1.5±0.9%, n=7 cells,

P(df=6, t=117.4)<.001, one-sided t-test), as expected (**Fig. 1C**). To ascertain that IPSCs

were mediated by synaptically activated interneurons rather than interneurons directly

recruited by electrical stimulation, we only included slices where inhibition was successfully

blocked by glutamatergic antagonists (25µM CNQX and 50µM D-APV, 8 of 21 experiments,

Fig. 1C). We also tested if inhibition of glutamate release from mossy fibers, which can be

specifically achieved via mGluR2/3 activation by DCG-IV(Doherty and Dingledine, 1998; Toth

et al., 2000), reduces feedback IPSCs. Indeed, we found that IPSCs were reduced to

16.3±6.1% by 0.5 µM DCG-IV (n=4 cells, P(df=3, t=13.73)<.001, one-sided t-test, **Fig. 1C**).

In order to relate the measured IPSCs to the fraction of GCs activated by a given stimulation

strength, we used population Ca^{2+} imaging with multibeam two-photon microscopy (**Fig. 1A,**

see Methods). After bolus-loading GCs with the Ca^{2+} indicator OGB-1-AM (see Methods),

antidromic stimulation caused action potential associated Ca^{2+} elevations in a subset of GCs

(**Fig. 1D,** transients indicated by *). Before quantifying population activity, we verified the

reliable detection of single action potentials under our conditions using simultaneous cell-

attached recordings from dentate GCs (**Fig. 1E, Supplementary Fig. 1**). Briefly, cells were

differentiated into true responders or non-responders on the basis of cell-attached recordings

(**Fig. 1E, F**; responders green, non-responders grey). A histogram of the peak ΔF/F of non-

responders upon a single stimulus was fitted with a Gaussian (**Fig. 1F** right, grey dots, grey

bars, n=33) and the threshold set to the quadruple standard deviation of this fit (0.94% ΔF/F,

dashed line in **Fig. 1F**). We estimated that this threshold would yield approximately equal

numbers of false positives and false negatives (**Supplementary Fig. 1F**). We additionally

controlled for possible errors through variable dye loading and the overestimation of the

active cell-fraction through accidental detection of adjacent active cells (**Supplementary Fig.**

1G, H, respectively).

Orientation of hippocampal slices may be a critical feature in determining the extent of

feedback connectivity. We therefore systematically assessed the magnitude of feedback

activation of GCs using imaging in slices obtained from different dorso-ventral levels of the

hippocampus (see inset of **Fig. 1 G**). We found a clear connectivity maximum within

horizontal slices obtained at a distance of ~1750 μm from the temporal pole (**Fig.**

1G,H)(Bischofberger et al., 2006). In these and all following experiments we therefore used

exclusively slices obtained at 1400-2100 μm from the temporal pole, where the orientation of

hippocampal slices matches the orientation of mossy fibers.

Combining the IPSC recordings with population Ca^{2+} imaging allowed us to probe the input-

output relationship of the feedback inhibitory microcircuit. Inhibition was recorded in a GC

within or immediately adjacent to the imaging field, and stimulation strength was increased

gradually (**Fig. 1I**). The IPSC saturated at 300μA stimulation strength, where the mean active

cell fraction was 2.2±0.7% and the mean IPSC reached 93.1±3.4% of the maximal IPSC

(**Fig. 1I**, **J,** n=20 for imaging, n=8 for IPSCs including 6 slices with both). Plotting the IPSC

magnitude vs. the cell fraction showed that the magnitude of feedback inhibition rises

steeply, reaching ~90% with less than 3% of GCs active and complete saturation at

3.7±1.7% of cells (**Fig. 1K**).

Fig. 1, Recruitment of feedback inhibition assessed using population Ca^{2+} imaging. Combined population Ca^{2+} imaging and IPSC recordings of GCs during antidromic electrical stimulation. **A)** Schematic illustration of the experimental setup. Dashed lines represent cuts to sever CA3 backprojections. **B)** Top: reconstruction of the dendritic tree of a representative GC. Bottom: Feedback IPSC at increasing stimulation strength during stratum lucidum stimulation. **C)** IPSCs were completely blocked by GABAzine and CNQX + D-APV and largely by DCG-IV. **D)** left: overlay of exemplary OGB1-AM loaded GC population (green) with a ΔF/F map (white). right: traces of ΔF/F over time of a subpopulation of cells depicted on the left. **E)** Simultaneous cell attached recording and calcium imaging to measure the action potential induced somatic calcium transient amplitude. **F)** Scatterplot and histogram of the calcium fluorescence peaks of cells which either did (green) or did not (grey) fire action potentials, as assessed by cell attached recordings. **G)** Illustration of the anatomical localization of maximum connectivity plane slices. Short black dashed lines indicate depth at which the slice plane is aligned to the dorsal brain surface. **H)** Antidromic stimulation elicited Ca^{2+} transients primarily at this depth (black bars). **I)** Normalized IPSC amplitude and activated cell fraction both increase with increasing stimulation strength (example from a single slice). **J)** Summary of all slices **K)** Summary data plotted to show the increase of inhibition as a function of the active GC fraction.

Optogenetic quantification of the recruitment of feedback inhibition

These experiments yielded a first quantitative estimate of the input-output relation of the

feedback-inhibitory microcircuit in the DG. We then decided to verify these findings using an

alternative method, which allowed spatially controlled and less synchronous GC activation.

Mice selectively expressing $ChR2^{(H134R)}$-eYFP in GCs were created by crossing Prox1-Cre

mice with Ai32-mice (**Fig. 2A**, see methods). Focal optogenetic stimulation was achieved

through a laser coupled into the microscope light path, yielding an 8 µm stimulation spot **(Fig.**

2B). Brief (20 ms, 473 nm) light pulses within the molecular layer approximately 40 µm from

the dentate GC layer elicited reliable IPSCs in GCs **(Fig. 2C)**. Increasing the light intensity

evoked larger IPSCs that showed clear saturation (**Fig. 2C, D,** Power=7 AU corresponding to

1.7 mW, see Methods). Inhibition was completely blocked by combined application of 40µM

CNQX and 50µM D-APV (**Fig. 2E**, n=9), confirming that it is recruited via glutamatergic

collaterals. The maximal IPSC amplitude obtained optically vs. electrically in experiments in

which both stimulations were performed were similar (**Fig. 2F**, paired t-test, P(df=3,

t=1.568)=.2148, n=4), indicating that similar maximal inhibition is recruited despite the

differences in the activated GC population (distributed vs local; synchronous vs. less

synchronous).

In order to relate feedback inhibition to the underlying GC activity levels, we performed

systematic cell attached recordings of GCs in the same slices in which inhibition was

recorded (~2 cells per slice, **Supplementary Fig. 2**). Briefly, we recorded the spatial firing

probability distribution in response to focal stimulation for each laser power. We then

estimated the mean firing probability of GCs throughout the section, which is equivalent to

the expected active GC fraction, by incorporating measurements of the light intensity

distribution throughout the slice (**Fig. 2G**, black). We additionally estimated an upper and

lower bound by assuming either no decay of firing probability with slice depth or isometric

decay (**Fig. 2G,** grey dashed lines). Combining the input-output relations of IPSCs **(Fig. 2D)**

and the estimated active cell fraction **(Fig. 2G)** again revealed that inhibition is recruited

steeply, saturating when approximately 4% of GC are active **(Fig. 2H).** Importantly, the

resulting recruitment function of inhibition is unlikely to be affected by voltage escape errors

(**Supplementary Fig. 3**). This is because such errors scale linearly with synaptic

conductance and will thus affect the absolute but not the relative amplitude of the somatically

measured IPSC. Next, we compared the focal light activation with global activation via a light

fiber positioned over the surface of the slice (with powers up to 50 mW, **Fig. 2l**). Under global

stimulation all cells tested fired APs with 100% reliability and independent of location, even

though focal stimulation in direct proximity to the cell led to much lower maximal firing

probabilities (**Fig. 2l, middle**, 100.0±0.0 versus 31.2±7.1% respectively, paired t-test, P(df=7,

t=9.74)<.001, n=8). At the same time, the maximal IPSC amplitude did not increase further

upon global stimulation (**Fig. 2l, right**, 356.9±76.2 versus 344.3±77.5 pA, paired t-test,

P(df=9, t=1.112)=.29, n=10). This implies that additional activation of remote GCs cannot

recruit interneurons beyond those activated by local GC populations. Thus the recruitment of

feedback inhibition in the DG is steep, with a dynamic range tuned to sparse populations of

GCs (up to 3-4% of cells).

Lower limit of feedback recruitment

Previous work has addressed the lower limit of the recruitment of feedback inhibition in

various cortical areas (Jouhanneau et al., 2018; Kapfer et al., 2007; Miles, 1990; Silberberg

and Markram, 2007). The authors report the ability of even a single principal cell to activate

feedback inhibitory interneurons and a supralinear increase of inhibition as the second and

third principal cells are co-activated (Kapfer et al., 2007). Given our findings so far we asked

whether single GCs might also suffice to elicit feedback inhibition in the DG. To this end we

performed dual patch clamp recordings and elicited short trains of ten action potentials at

100 Hz in one cell while monitoring inhibition in the other (**Supplementary Fig. 4,** n=15).

However, in contrast to the neocortex (Kapfer et al., 2007; Silberberg and Markram, 2007)

and area CA3 (Miles, 1990), we did not find single GC-induced feedback inhibition in any of

these experiments, consistent with a recent large scale study reporting that such connections

are extremely sparse (0.124%) (Espinoza et al., 2018).

Fig. 2, Recruitment of feedback inhibition assessed optogenetically. A) EYFP fluorescence in dentate GCs of Prox1/ChR2(H134R)-EYFP transgenic mice. The field of view for rapid focal optogenetic stimulation is indicated by a blue square. A typical stimulation site approx. 40 µm from the GC layer (two short black lines) is indicated by a blue dot. **B)** Schematic of the microscope setup used to achieve spatially controlled illumination. The inset shows the intensity profile of the laser spot. **C) Top left,** reconstruction of an Alexa594 filled GC.   **Left,** illustration of optical stimulation. **Right,** IPSCs following 20ms light pulses at increasing laser power (P = 1 to 7 AU). Each trace represents an average of 3 trials. **D)** Summary of IPSC amplitudes from cells in the superior blade (n=7 cells). IPSC amplitudes were normalized to the maximum amplitude within each cell. **E)** Optogenetically elicited IPSCs are abolished by glutamatergic blockers (40µM CNQX + 50µM D-APV, n=9). **F) Left,** Schematic of focal optical and electrical stimulation. Dashed lines indicate cuts to sever CA3 backprojections. **Middle,** Example traces for IPSCs following electrical or focal optogenetic stimulation. **Right,** maximal IPSC amplitude for the two stimulation paradigms (361±37 vs. 410±13 pA for electrical and optogenetic stimulation respectively, paired t-test, p=0.28, n=4) **G)** The optogenetically activated GC fraction responsible for recruiting the IPSC at the respective laser powers was estimated from systematic cell attached recordings (see Fig. S1 for details). The best estimate (black) incorporates measurements of the 3D light intensity profile in the acute slice. Upper and lower bounds were estimated by assuming no firing probability decay with increasing slice depth (upper grey dashed line) or isometric firing probability decay (lower grey dashed line). **H)** Data from (D) and (H, best estimate) plotted to show the recruitment of feedback inhibition. **I)** Comparison of focal optogenetic stimulation to global (light fiber mediated) optogenetic stimulation. **Left,** Schematic illustration. **Middle,** Comparison of the AP probability of individual GCs at maximal stimulation power for focal and global stimulation assessed by cell attached recordings. **Right,** Comparison of the maximal IPSC amplitude under focal and global stimulation for individual GCs.

Spatial distribution of feedback inhibition

Recent evidence indicates that inhibition by individual PV[+] fast spiking hilar border

interneurons is non-uniformly distributed over space, with decreasing connectivity and

inhibition at greater distances from the interneuron (Espinoza et al., 2018; Strüber et al.,

2015). To test whether feedback inhibition by the entire ensemble of feedback inhibitory

interneurons also displays a spatial gradient we activated cell populations at 100 µm intervals

along the GC layer while recording inhibition in individual GCs (**Fig. 3A**). Spatial profiles were

recorded for increasing laser powers in cells in the superior as well as inferior blade of the

DG (**Fig. 3B, C** respectively; n=8 cells for each blade). IPSC amplitudes across locations and

powers were normalized to the maximal IPSC amplitude in each respective cell. This

maximal amplitude did not differ between cells in different blades (366±40 vs 390±84 pA for

superior and inferior blades respectively; t-test, P(df=14, t=0.258)=.0686). Next, we

investigated the spatial organization of feedback inhibition at stimulation powers at which

inhibition had saturated (**Fig. 3D, E**). In all GCs tested, the inhibition was greatest when

stimulating in the direct vicinity of the recorded cell. Activating cells at increasing distances

led to monotonically decreasing IPSC amplitudes for both blades. Importantly, the term

distance here refers to the functional distance along the GC layer and not to Euclidean

distance. However, inhibition was observed even at the most remote stimulation sites,

indicating that even the most remote cells from the contralateral blade can contribute to the

activation of feedback inhibition in a given GC. In order to statistically compare the relation of

local versus remote inhibition between blades we defined a remote location in the

contralateral blade at 800 µm from the recorded cell (measured along the GC layer and

equidistant in all slices; **Fig. 3D, E**; grey lines) and compared it to the local IPSC (black

lines). Remote inhibition was significantly smaller than local inhibition while no difference

between blades or significant interaction was observed (**Fig. 3F**; two-way RM ANOVA;

Distance: $F_{(1,14)}=3.341$, P<.001; Blade: $F_{(1,14)}=2.615$, P=.128; Interaction: $F_{(1,14)}=3.341$,

P=.089). Posttests suggested inhibition of inferior GCs by superior activation might be

greater than vice versa. However, the difference was not significant (Sidak's multiple

comparison corrected posttest, P(df=28)=.932, P(df=28)=.051 for local and remote

respectively).

Next, we investigated whether there are differences in the steepness of recruitment of local

versus remote inhibition between blades (black and grey, respectively; **Fig. 3G, H**). To this

end, we calculated the active cell fraction which produces half-maximal inhibition during local

or remote stimulation for each individual slice. Comparison of the recruitment between the

four groups revealed no differences between blades (**Fig. 3I,** two-way RM ANOVA; Distance:

$F(1,14)=7.889$, P=.014; Blade: $F(1,14)=0.5506$, P=.470; Interaction: $F(1,14)=0.0976$,

P=.759). However, local inhibition was significantly more steeply recruited than remote

inhibition (1.99 ± 0.22% vs. 3.17 ± 0.57 % active cells for half-maximal inhibition).

Next, we tested if IPSCs elicited by increasing active GC populations differed between local

and remote activation with respect to their kinetic properties. Since all previous data showed

no indication of blade specific differences the analysis of the kinetics of feedback IPSCs were

performed on the pooled data for both blades. Interestingly, local and remote inhibition

differed in all tested respects (**Fig. 3J–M,** two-way RM ANOVAs with $df_{Distance}=1,183$, df_{cell}

$_{fraction}=6,183$ and $df_{interaction} =6,183$). Local IPSCs occurred with shorter latency and lower jitter

than remote IPSCs (**Fig. 3J, K;** Latency: P<.001, <.001 and =.031 for distance, cell fraction

and interaction, respectively; Jitter: P<.001, =.037 and =.707 for distance, cell fraction and

interaction, respectively). Furthermore, both latency and jitter decreased as larger

populations were activated. IPSCs were also significantly slower in remote versus local

inhibition. IPSC rise time was slightly shorter in the larger local IPSCs but did not correlate

with the active cell fraction (**Fig. 3L**: P=.010, =.633 and =.388 for distance, cell fraction and

interaction, respectively). Similarly, decay times were significantly shorter in local versus

remote inhibition while they progressively increased with increasing stimulation power (**Fig.**

3M; P<.001, <.001 and =.124 for distance, cell fraction and interaction, respectively). These

data demonstrate that remote inhibition shows greater delay, greater jitter and slower kinetics

than local inhibition.

Fig. 3, Spatial organization of feedback inhibition. Feedback IPSCs recorded from an individual GC while GCs at varying distances were activated. **A)** Schematic illustration of the stimulation paradigm and example IPSC traces of an individual trial (P=3). **B, C)** Distribution of normalized IPSC amplitudes as a function of laser power and distance from stimulation spot for superior and inferior blade GCs (n=8 for each blade). The relative location of the DG apex ± standard deviation is indicated by the black bar and grey area respectively. **D, E)** IPSC distribution over space at saturation (P≥5). Black and grey bars indicate a local and a remote location at 800 µm from the recorded cell respectively. **F)** Comparison of the amplitude of the locally and remotely activated IPSCs at saturation (two-way RM ANOVA, overall test significance indicated by §) **G, H)** Comparison of the recruitment curves during local (black) or remote (grey) stimulation for superior and inferior blade respectively. **I)** Comparison of the cell fraction required for halfmaximal IPSC activation between stimulation sites and blades (two-way RM ANOVA overall test significance indicated by §). **J-M)** Temporal properties of IPSCs between local (black) and remote (grey) stimulation. To test for systematic variations of kinetic parameters with increasing active cell fractions as well as stimulation site two-way RM ANOVAs with no post tests were performed. Overall significance indicated by §. **K)** Latency from beginning of light pulse to IPSC **L)** temporal jitter of IPSCs (SD of latency within cells) **M)** 20% to 80% rise time **N)** IPSC decay time constant.

Short term dynamics in the feedback inhibitory microcircuit

Different connections within the feedback inhibitory microcircuit have been shown to variably

facilitate or depress during trains of activity(Savanthrapadian et al., 2014) (**Supplementary**

Table 1). This makes it difficult to predict the net effect on the short term dynamics of GC

feedback inhibition. We therefore characterized the frequency-dependence of net feedback

inhibition using antidromic electrical stimulation as described above (**Fig. 4A-C**). In marked

contrast to the CA1 region of the hippocampus (Pothmann et al., 2014), feedback IPSCs

showed strong frequency dependent facilitation (**Fig. 4C**, n=10 cells, one-way RM ANOVA;

Frequency: $F_{(2.69, 29.54)}=13.99$, P<.001; Wilcoxon signed rank tests for deviation from

unity at each frequency with Bonferroni corrected p-values; P=1, =.004, .002 and .002 for 1,

10, 30 and 50Hz, respectively). We found no evidence for a spatial gradient of net feedback

inhibitory short term dynamics (**Supplementary Fig. 5**).

Because this unusual degree of facilitation may be important in allowing sparse activity of

GCs to recruit significant inhibition over time, we further examined the underlying circuit

mechanisms. Interestingly, previous literature on the DG feedback microcircuit almost

exclusively reports short term depression of dentate interneuron inputs to GCs

(**Supplementary Table 1**, blue rows), rendering our finding of pronounced facilitation at the

circuit level even more striking. We therefore measured feedback excitation of hilar neurons

by stimulating mossy fiber axons as described above (**Fig. 4D-L**). Mossy cells and

interneurons were classified according to their morpho-functional properties(Larimer and

Strowbridge, 2008) (**Fig. 4D, E, G, H, J, K**). Cell classification was confirmed using unbiased

k-means clustering (**Fig. 4K**). We found that feedback excitation of hilar cells displayed

marked facilitation, which was similar for both INs and MCs (**Fig. 4F, I, L;** n=9, 12

respectively, two-way RM-ANOVA, Frequency: $F_{(3,57)}=6.642$, P<.001; Cell type:

$F_{(1,19)}=0.0075$, P=.932; Interaction: $F_{(3,57)}=0.743$, P=.531). Facilitation indices of hilar cells

significantly deviated from one for all frequencies tested (**Fig. 4E, F**; n=23 cells; Wilcoxon

signed rank tests with Bonferroni corrected p-values; P<.001 for all frequencies). These data

demonstrate a pronounced frequency dependent net facilitation of the feedback inhibitory

microcircuit, which is supported by strongly facilitating mossy fiber inputs to hilar cells.

Fig. 4, Short term dynamics in the feedback inhibitory microcircuit. Trains of ten antidromic electrical stimulations at 1, 10, 30 or 50 Hz were applied to elicit disynaptic feedback inhibition or excitation of hilar cells (electrical stimulation artifacts were removed in all traces). **A, D, G)** Schematic illustration of the experimental setup and example traces of voltage responses to positive and negative current injections of GC and hilar cells (dashed lines indicate cuts to sever CA3 backprojections) **B)** Exemplary GC feedback IPSCs before (black) and after (grey) glutamatergic block (n = 7). **C)** Facilitation indices (mean of the last three IPSCs normalized to the first; n=10 cells) **D-L)** Hilar cells were manually classified into putative interneurons (blue) or mossy cells (green) based on their morpho-functional properties. **E)** Reconstruction of biocytin filled hilar interneuron (axon in red). **F)** Interneuron EPSCs in response to stimulation trains. **H)** Reconstruction of biocytin filled mossy cell (axon in red). **I)** Mossy cell EPSCs in response to stimulation trains. **J)** Quantification of intrinsic properties of hilar cells (see methods). **K)** k-means clustering based on intrinsic properties of hilar cells (coloring according to manual classification). **L)** Facilitation indices of classified hilar cells. (§ indicates significance in one-way RM ANOVA, * show significance in Bonferroni corrected Wilcoxon signed rank tests for deviation from 1).

Quantitative properties of the feedback circuit predict frequency dependent pattern

separation

Together, these data indicate that the dentate feedback circuit is able to deliver strong,

spatially graded inhibition with a high gain and the ability for temporal integration. To probe

how these quantitative properties of the feedback circuit affect the pattern separation

capability of the DG, we incorporated them into a biophysically realistic model of the lamellar

microcircuit (**Fig. 5**) based on Santhakumar et al. (2005)(Santhakumar et al., 2005; Yim et

al., 2015), making use of their carefully experimentally constrained DG cell-types (**Fig. 5A,**

Supplementary Fig. 6A). To maximize our models inferential value we clearly separated a

tuning phase, in which we constrained the model by our experimental data, and an

experimental phase, in which pattern separation was tested without further changes to the

model. In the tuning phase, we first scaled up the model four-fold to contain 400 perforant

path afferents (PPs), 2000 GCs, 24 basket cells (BCs), 24 hilar perforant path associated

cells (HC) and 60 MCs (**Fig. 5A, B**). BCs, HCs and MCs comprise the feedback inhibitory

circuit and BCs receive direct PP input thereby additionally mediating feedforward inhibition

(Ewell and Jones, 2010). We then adapted the spatial extent of the target pools of BC and

HC outputs to produce local and global inhibition respectively, reproducing the

experimentally determined spatial tuning of *net* feedback inhibition (**Fig. 5C**). We further

adjusted synaptic decay time constants and weights in order to reproduce the measured

PSCs of hilar neurons and GCs and the empirical recruitment curves (**Fig. 5D,**

Supplementary Fig. 6). Finally, we incorporated facilitation of the experimentally determined

magnitude into feedback excitatory mossy-fiber outputs, leading to GC IPSC facilitation in the

experimentally observed range (**Fig. 5E, Supplementary Fig. 6B**). Together, these minimal

adaptations resulted in a model with remarkably similar properties to our experimental

findings (**Fig. C-E**). We therefore concluded the tuning phase of the model and proceeded to

an in silico pattern separation experiment without further changes to the model.

Fig. 5, Computational model of the DG feedback circuit. A biophysically realistic model of DG was tuned to capture the key quantitative features of the feedback circuit. All analyses were performed as for the real data (including IPSC normalization to maximal IPSC over space and power within each respective cell) **A)** Schematic of the model circuit. GC: granule cell, BC: basket cell, HC: hilar perforant path associated cell, MC; mossy cell. **B)** Intrinsic responses of model cell types to positive and negative current injections. **C)** Spatially graded net feedback inhibition following simulated focal GC activation. **D)** Local and remote recruitment curves of the feedback inhibitory circuit (left) and the resulting saturated IPSC amplitudes and GC fractions recruiting halfmaximal inhibition (right). **E)** Facilitation indices resulting from simulated, 10 pulse, frequency stimulation of GCs as above.

To investigate the implications for pattern separation we probed the ability of this model to

separate PP input patterns with behaviorally relevant temporal structure and varying degrees

of overlap (Myers and Scharfman, 2009; Yim et al., 2015). Specifically, we created input

trains with constant mean rate, but with either theta (10 Hz) or slow-gamma (30 Hz)

modulation (**Supplementary Figs. 6C, 7A**), which are prominent during exploration and

novelty exposure respectively (Sasaki et al., 2018; Trimper et al., 2017). To model rapid

pattern separation in behaviorally relevant timescale we chose an input duration of

approximately five theta cycles (600ms, corresponding to the approximate duration of place

cell spiking during traversal of its place field). To obtain a range of input similarities, we

generated input patterns in which 24 of 400 PP afferents were activated (**Fig. 6A,**

Supplementary Fig. 6) and compared pairs of such patterns ranging from no overlap (two

separate sets of afferents) to complete overlap (identical trains in the same 24 afferents in

both patterns). Each model network was run with 25 input patterns leading to a total of 325

comparisons (data points in **Fig. 6C**). To quantify pattern separation we compared input

correlation (R_{in}) to output correlation (R_{out} ; **Fig. 6B**) both measured as Pearson's R between

the population rate vectors over the full 600ms time window (Leutgeb et al., 2007; Wiechert

et al., 2010). Our full, tuned model reliably decreased the population vector correlations for

similar patterns ($0 < R_{in} < 1$) thereby demonstrating robust pattern separation over the whole

range of input similarities ($R_{out} < R_{in}$; **Fig. 6C**, left). Next, we isolated the contribution of

feedback inhibition to pattern separation by rerunning the same input pattern combinations

on the network in which mossy fiber outputs to interneurons were removed (**Fig. 6C,** middle).

This eliminated HC activity and reduced BC spiking by nearly 50% (not shown). Note that

removing mossy fiber outputs also eliminates BC activity through cooperative activation by

summating feedforward and feedback inputs(Ewell and Jones, 2010). Removal of all

inhibitory outputs led to a further decrease in pattern separation, demonstrating the effect of

additionally removing feedforward inhibition (**Fig. 6C**, right). As expected these manipulations

increased both the fraction of active GCs and the activity per GC (**Supplementary Fig. 7B,**

C). In order to quantify the respective pattern separation effects over the full range of input

similarity, we computed the bin wise mean R_{out} (Fig. 6C, R_{in} bin-width: 0.1, dashed line) and

measured the area to the identity line (**Fig. 6C**, black lines). The resulting mean ΔR_{out} was

calculated for seven separate random networks, each challenged with theta as well as slow-

gamma modulated inputs in each of the three conditions. Both the frequency of the input

modulation as well as network manipulatons significantly affected pattern separation (**Fig.**

6D; two-way RM ANOVA with both factors matching, condition: $F_{(2,12)} = 145.1$, $P < .001$;

frequency: $F_{(1,6)} = 31.48$, $P = .001$; interaction: $F_{(2,12)} = 11.77$, $P = .002$; n=7 random network

seeds for these and all subsequent analyses). Specifically, both feedback and feedforward

inhibition significantly contributed to pattern separation (Sidak's multiple comparison posttest,

$P(df=12, t=11.33)<.001$ and $P(df=12, t=5.36)<.001$, respectively).

To quantify the isolated effects of feedback and feedforward inhibition we computed the

difference in R_{out} between the respective conditions for each individual comparison (i.e. data

point in Fig. 6C). For instance, the individual comparison shown in Fig. 6A, will lead to a

single R_{out} value in the network with MF inputs to interneurons (full model), which is

subtracted from the corresponding R_{out} value in the same network without this input (no FB).

This procedure isolates the effect of interest (ΔR_{out}) for each individual comparison,

controlling for other sources of variability. A single pattern separation measure was then

obtained as before, as the area under the curve of bin-wise means of these ΔR_{out} values

(**Fig. 6E**, bottom). We found a significant effect of both inhibitory motif and frequency domain

(**Fig. 6E**; two-way RM ANOVA with both factors matching, Motif: $F(1,6)=15.58$, $P=.008$;

Frequency: $F(1,6)=9.91$, $P=.020$; Interaction: $F(1,6)=76.37$, $P<.001$). Posttests revealed that

the frequency dependence of pattern separation was driven by feedback inhibition (Sidak's

multiple comparison posttest: FB: $P(df=6, t=13.68)<.001$; FF: $P(df=6, t=1.33)=.412$.

Interestingly, this frequency dependence of feedback inhibition mediated pattern separation

was particularly pronounced for highly similar input patterns ($0.9<R_{in}<1$; Fig. 6E, right; Motif:

$F(1,6)=261.7$, $P<.001$; Frequency: $F(1,6)=108.1$, $P<.001$; Interaction: $F(1,6)=109.5$, $P<.001$;

Sidak's multiple comparison posttest: FB: $P(df=6, t=15.78)<.001$; FF: $P(df=6, t=0.98)=.595$).

Indeed, feedback inhibitory pattern separation for highly similar input at 30Hz compared to

10Hz was more than doubled (from 0.04±0.01 to 0.09±0.01, mean±SD, Cohen's d=4.1, **Fig.**

6E, right).

It has recently been emphasized, that the assessment of pattern separation can depend

critically on the similarity measure used (Madar et al., 2019; Wick et al., 2010). Therefore we

tested the robustness of this result for two alternative similarity measures, namely normalized

dot product (NDP, also known as cosine similarity) and pattern overlap (# of coactive/ # of

totally active cells; **Supplementary Fig. 7).** The frequency dependence of feedback inhibition

mediated pattern separation, especially for highly similar inputs, proved robust for all three

similarity measures (see figure legend for statistics).

Effect of spatial tuning and facilitation of net feedback inhibition

Next we investigated the specific effects of two interesting empirical findings of the present

study, 1) the spatial tuning and 2) the facilitation of the feedback circuit (**Supplementary Fig.**

8). To this end we undertook two targeted, minimal manipulations of the full tuned network.

To probe the effect of spatially graded inhibition, we redistributed BC output synapses to a

global target pool (the whole GC population), leading to spatially uniform inhibition (global

FB; **Supplementary Fig. 8B; 8E, left**). To probe the effect of facilitation, we removed

facilitation from mossy fiber outputs (**Supplementary Fig. 8C; E, right**). We isolated the

effects of these manipulations by pairwise comparison to the corresponding full tuned

networks as described above (**Supplementary Fig. 8F-I**). The results showed a small but

significant contribution of facilitation (~20% of the isolated FB effect for both frequency

paradigms), but not spatial tuning to pattern separation (**Supplementary Fig. 8G, left**;

Wilcoxon signed rank test for deviation from 0, n=7, Bonferroni corrected P-values: P=.031

and P=1 respectively for 10Hz; P=.031 and P=1 respectively for 30Hz). We noted that while

spatial tuning did not affect mean pattern separation, it appeared to reduce its variability

(CoV) for a given input similarity (**Supplementary Fig. 8G, right**; Wilcoxon signed rank test

for deviation from 0, n=7, Bonferroni corrected P-values: P=.031 and P=.750 for tuning and

facilitation respectively at 10Hz; P=0.438 and P=1 respectively at 30Hz).

Fig. 6, Frequency dependent pattern separation of temporally structured inputs. The quantitative DG model was challenged with theta (10Hz) or slow gamma (30Hz) modulated input patterns with defined overlap to probe its pattern separation ability. **A)** Pair of theta modulated perforant path input patterns in which 50% of afferents overlap (grey area). **B)** Resulting pair of GC output patterns of the full tuned network. **Bottom:** Representative individual GC underlying the observed patterns. **C)** Comparison of 325 input pattern pairs and their resulting output pattern pairs. Each pair is characterized by its rate vector correlation for inputs (R_{in}) and outputs (R_{out}), where rates are measured over the full 600 ms time window. Dashed black lines represent the bin-wise mean R_{out} (in R_{in} bins of 0.1). **Left:** full tuned model, **middle:** model without mossy fiber inputs to interneurons, **right:** model without inhibitory synapses. **D)** Full pattern separation effects (mean ΔR_{out}) of all three conditions for both frequency domains quantified as the area enclosed by the dashed and unity lines in (C). Black lines represent individual network seeds. Two-way RM ANOVA indicated significance of condition, frequency and interaction, * indicate significance in Sidak's posttests between individual conditions. **E)** Isolated effects of feedback and feedforward motifs obtained by pairwise subtraction of R_{out} between conditions for each individual comparison. The inset shows the resulting ΔR_{out} for each R_{in} bin. The area under the curve quantifies the mean ΔR_{out} as in (D). Two-way RM ANOVA indicated significance of condition, frequency and interaction. *** indicate p<0.001 in Sidak's posttest. **F)**. 100ms time-resolved pattern separation effects of the full model, isolated FB or FF inhibition for theta modulated input (10Hz). All analyses were performed as above but with rate vector correlations computed for 100 ms time windows. The bottom insets show ΔR_{out} as a function of input similarity for each time window. The bottom right insets show the evolution of the mean ΔR_{out} over time. **G)** Same as (F) but for slow gamma (30Hz) modulated inputs. Arrow indicate the region of selectively increased pattern separation. Data in D-G represent mean ± SEM of n=7 random network seeds.

Frequency dependent pattern separation is robust over analysis scales

So far, all pattern separation analyses were conducted on the population rate vectors during

a 600 ms time window. However, many neural computations are likely to occur on shorter

timescales, such as within individual theta (~100 ms) and gamma (~10-33 ms) cycles

(Buzsáki, 2010; van Dijk and Fenton, 2018). Indeed, the time window in which correlation is

recorded can nontrivially affect the resulting correlation, depending on the timing of spikes

within it (Madar et al., 2019). We therefore first computed the networks pattern separation

ability within 100 ms time windows, revealing i) the pattern separation ability within such

short timescales and ii) the temporal evolution of pattern separation throughout a 600 ms

stimulus presentation (**Fig. 6F, G**). We find that pattern separation occurs even within a

single theta cycle, including a contribution of feedback inhibition in both frequency paradigms

(mean ΔR_{out} significantly different from 0 within the first 100 ms bin, Wilcoxon signed rank

test with Bonferroni corrected p-values: P=.031, =.031 for full and FB effect respectively in

both paradigms). While mean ΔR_{out} did not differ between frequency paradigms within this

first time window, it was significantly elevated in the 30 Hz paradigm in all subsequent time

windows (full model effect, two-way RM ANOVA, P<.001, <.001 and =.004 for time-bin,

frequency and interaction respectively, Sidak's posttest P=.234 for 1st bin and P<.001 for all

subsequent bins). Again, the selective increase during slow-gamma modulated inputs was

driven by feedback inhibition (isolated FB effect, two-way RM ANOVA, P=.007, <.001 and

=.041 for time-bin, frequency and interaction respectively, Sidak's posttest P=.708 for 1st bin

and P<.002 for all subsequent bins), including a contribution from MF facilitation

(**Supplementary Fig. 8**). As above, the effect was predominantly driven by the separation of

highly similar input patterns (isolated FB effect, $R_{in}>0.5$; two-way RM ANOVA on last time-

bin, P<.001, =.010 and <.001 for R_{in}-bin, frequency and interaction respectively, Sidak's

posttest on differences between frequency paradigms for each input similarity: P=1 for

$R_{in}<0.6$ and P=.032 to P<.001 for R_{in}= 0.6 to 0.9). These results were robust when analysis

time windows were decreased even further (to the duration of a slow gamma cycle, 33 ms,

Supplementary Fig. 9). This 33 ms resolved analysis additionally reveals that the pattern

separation effect, particularly of feedback inhibition, ramps up within a 100 ms window,

becoming effective only at the end of a theta cycle (**Supplementary Fig. 9B**).

Next we asked if the frequency dependence of feedback inhibitory pattern separation was

sensitive to variations of the inhibitory decay time constants and if there might be an

interaction between these decay time constants and the frequency range at which pattern

separation is most effective (**Supplementary Fig. 10**). Remarkably, we found the differential

effect between 10 & 30 Hz to be highly robust across a range of different decay time-

constants (0.5x to 5x the experimentally matched decay, **Supplementary Fig. 10A-C, Table

2**). Furthermore, the selective enhancement of feedback inhibitory pattern separation of

highly similar inputs was robust over the entire gamma range (up to 100 Hz, **Supplementary

Fig. 10D, E**).

Finally, we tested if our main results were robust to alterations in the relative strengths of

feedforward vs. feedback inhibition. Since, our model is closely constrained with respect to

the recruitment and functional properties of the feedback circuit, we are confident about the

resulting computational inferences concerning this circuit. However, the model does not allow

strong inferences about the relative roles of feedback and feedforward inhibition, and it is

thus necessary to probe if extremely powerful feedforward inhibition might occlude the

effects described here. We therefore selectively enhanced the feedforward inhibitory circuit in

our model by increasing the PP to BC circuit 2x (**Supplementary Fig. 11**). This robustly

increased the feedforward inhibitory contribution to pattern separation above that of feedback

inhibition (**Supplementary Fig. 11B**). However, it did not affect the frequency dependence of

the feedback inhibitory effect. Indeed, for highly similar input patterns, the feedback inhibitory

effect was so prominently enhanced during gamma input, as to again dominate the

feedforward inhibitory effect (**Supplementary Fig. 11C**).

Together these results suggest that frequency dependence is a key feature of the feedback

inhibitory microcircuits and predict that feedback inhibition selectively boosts the separation

of highly similar input patterns during gamma oscillations.

Discussion

Across brain regions and species, inhibitory circuits contribute critically to regulating the sparsity and overlap of neural representations (Cayco-Gajic and Silver, 2019; Lin et al., 2014; Papadopoulou et al., 2011; Stefanelli et al., 2016). In most, if not all brain regions, feedback inhibition is viewed as important in these capabilities, by directly mediating competition between active cell ensembles (de Almeida et al., 2009; Myers and Scharfman, 2009; Rolls, 2010). In the mammalian DG, feedback inhibition is implemented by an intricate network of interneurons that is capable of delivering spatiotemporally defined inhibition to the principal cell population. How *net* feedback inhibition is functionally organized in mammals, and how it may contribute to pattern separation of biologically relevant, temporally structured input patterns is, however, incompletely understood.

Quantitative physiological properties of DG feedback inhibition

We have therefore quantitatively described the recruitment of *net* feedback inhibition by defined GC population sizes in space and time in the hippocampal DG, a structure in which sparse activity and inhibition are thought to critically contribute to the function of pattern separation (Gilbert et al., 2001; Hunsaker et al., 2008; Leal and Yassa, 2018; McHugh et al., 2007; Stefanelli et al., 2016). The proposed role of the feedback inhibitory circuit depends critically on its dynamic range, i.e. the relation between the number of active principal cells and the resulting feedback inhibition. This property of the feedback circuit is determined by complex, mainly hilar cellular connectivity patterns (Espinoza et al., 2018; Savanthrapadian et al., 2014) (**Supplementary Table 1**). While delving into detailed cell-cell connectivities is clearly important, such studies do not allow the quantitative determination of the gain and dynamic range of *net* feedback inhibition (Kapfer et al., 2007; Silberberg and Markram, 2007). Using two complementary experimental approaches, we found that *net* feedback inhibition is steeply recruited by sparse populations of GCs (<4%). Accordingly, the gain and sensitivity of the circuit are well suited to strongly modulate feedback inhibition within the range of GC activity reported in vivo (Pernía-Andrade and Jonas, 2014; Schmidt et al.,

2012). Additionally, we have described the temporal and spatial distribution of inhibition

delivered by feedback circuits in the DG.

Frequency-dependent effects of feedback inhibition on pattern separation

How do the properties of feedback circuits influence the pattern separation capability of the

dentate gyrus? To address this question we adapted an established biophysically realistic

computational model of the DG circuitry (Santhakumar et al., 2005; Yim et al., 2015). We first

carefully constrained the model to match the spatial and temporal properties of *net* feedback

inhibition as assessed in our physiological data. We then fixed all model parameters, and

proceeded to probe the ability of this circuit to perform pattern separation on temporally

complex oscillatory inputs. The major, highly robust, result of this computational study was

that the impact of feedback inhibition on pattern separation is frequency-dependent.

Specifically, we find that the separation of input patterns during gamma oscillations >30 Hz is

powerfully and selectively enhanced by the feedback circuit. Remarkably, this mechanism

was particularly efficient for very similar input patterns. Such an effect has not been

discovered in earlier modeling studies, because most models have discretized time,

calculating the pre-inhibition population activity, the resulting inhibition, and the inhibition-

corrected population activity in a single time step, sometimes assuming an average corrected

population rate within this time step (Myers and Scharfman, 2009; Rolls and Treves, 1998;

Trappenberg, 2010). Thus, they do not capture temporal features of feedback circuits. On the

other hand, a number of spike based, temporally resolved models have considered only

temporally unstructured (Poisson) inputs (Chavlis et al., 2017; Hendrickson et al., 2015;

Hummos et al., 2014; Yim et al., 2015). We suggest that the precise spatiotemporal

organization of the feedback circuit, together with the temporal structure of DG inputs is a

crucial determinant of pattern separation. Indeed, the DG and its inputs have a strong,

behaviorally relevant, temporal structure (Mizuseki et al., 2009; Pernía-Andrade and Jonas,

2014; Skaggs et al., 1996). Novelty experience can induce increased gamma and beta range

activity (Berke et al., 2008; Rangel et al., 2015; Trimper et al., 2017), and explorative activity

with rearing is also associated with increased gamma oscillations (Barth et al., 2018). A

previous model has addressed how fast, rhythmic gamma-frequency feedback inhibition may

implement a type of 'k-winners-take-all' operation, a basic computational component of

pattern separation models (de Almeida et al., 2009), though this model relies on faster

synaptic timescales than we observed in our compound IPSCs. Perhaps most interestingly,

the occurrence of oscillations in the slow-gamma range has recently been reported to be

causally related to associative memory formation (Sasaki et al., 2018; Trimper et al., 2017), a

process thought to require pattern separation. Consistent with this finding, Hsiao et al. (Hsiao

et al., 2016) report DG driven gamma entrainment of CA3, the presumed primary storage

location of associative memories. Together, this suggests that the dentate pattern separator

may be optimized to rapidly detect subtle degrees of difference within the environment in

gamma-dominated exploratory brain states, a capability likely to support successful memory

encoding of novel environmental features, and potentially aiding in rapid discrimination

during recall.

Importantly, the frequency-dependency of pattern separation was driven by the feedback

circuit. This effect was highly robust when varying the decay time constants of the inhibitory

synaptic conductances, the time windows of analysis, or the similarity measure employed. By

contrast, feedforward inhibition and anatomical pattern separation was robustly independent

of frequency modulation. Together this suggests that frequency-dependent pattern

separation is a key property of the local inhibitory feedback circuit. Importantly, this does not

preclude that additional, long range projections may add further complexity (Szabo et al.,

2017). Also note that in addition to the instantaneous pattern separation mechanisms

investigated here, potentially complementary mechanisms at much longer time scales have

been proposed involving ongoing neurogenesis (Aimone et al., 2011; Clelland et al., 2009; Li

et al., 2017; Sahay et al., 2011; Severa et al., 2017; Temprana et al., 2015).

Spatiotemporal organization of inhibition and pattern separation

The model also allowed us to examine the impact of the spatiotemporal organization of

inhibition on pattern separation. Facilitation of feedback circuits produced a small but robust

enhancement of pattern separation, while spatial tuning of feedback inhibition did not. The

facilitation of feedback inhibition is a remarkable feature of the DG, which we to our

knowledge have described for the first time. It is in marked contrast to area CA1, where

somatically measured feedback inhibition shows strong depression (Pothmann et al., 2014;

Pouille and Scanziani, 2004) and is particularly surprising given the prevalence of depression

in the literature on pairwise connections (**Supplementary Table 1**). Our physiological and

modeling data suggest that the strong facilitation of the mossy fiber input to the feedback

circuit is the principal mediator of this net facilitation. The effect of facilitation on pattern

separation is intuitive, since this allows the feedback circuit to integrate GC activity over time,

and convert it to inhibition.

In our model, spatial tuning of feedback inhibition had no effects on pattern separation. This

may derive from the fact that PP inputs were spatially broad and random (Tamamaki, 1997;

Tamamaki and Nojyo, 1993). In general, the effect of localized inhibition could be more

relevant if synchronously activated populations of GCs are locally clustered (Feldt Muldoon

et al., 2013). For instance, GCs in the inferior and superior blades of the DG are known to be

differentially active (Alme et al., 2010; Chawla et al., 2005). Accordingly, localized inhibition

might be important for independent processing between the two blades. An alternative

function of spatially graded inhibition has been proposed by (Strüber et al., 2015), who

suggest that it is more effective in promoting synchronous gamma oscillations. Accordingly,

spatial tuning may play a role in creating the oscillatory dynamics, found here to critically

impact the feedback inhibitory pattern separation performance.

Together, this study provides the first comprehensive, quantitative description of the

spatiotemporal properties of the DG feedback inhibitory microcircuit, and predicts that these

properties will selectively enhance the separation of highly similar input patterns during

learning- related gamma oscillations. This mechanism may be relevant for understanding

disease states in which there is a coincidence of dentate gyrus-centered pathology with

abnormal oscillatory activity, and memory and pattern separation deficits such as temporal

lobe epilepsy, Alzheimer's disease or schizophrenia (Andrews-Zwilling et al., 2012; Gillespie

et al., 2016; Leal and Yassa, 2018; Verret et al., 2012).

Materials and Methods

Animals and Slice Preparation: All animals were treated according to the University of Bonn Animal Experiment Guideline, minimizing unnecessary pain and discomfort. Experiments were performed on horizontal hippocampal slices of 21 to 97 day old mice. Ca^{2+} imaging and a subset of dual recording experiments were performed in C57/Bl6 mice obtained from Charles River Laboratories (Wilmington, Massachusetts, USA). Optogenetic experiments and the remaining dual recording experiments were performed on double transgenic offspring of Tg(Prox1-cre)SJ39Gsat/Mmucd, MMRRC Cat# 036632-UCD, RRID: MMRRC_036632-UCD) obtained as cryopreserved sperm and rederived in the local facility (Gong et al., 2007, 2003) and Ai32-mice (B6;129S-Gt(ROSA)26Sor[tm32(CAG-COP4*H134R/EYFP)Hze]/J, IMSR Cat# JAX:012569, RRID: IMSR_JAX:012569). For preparation the animals were deeply anesthetized with Isoflurane (Abbott Laboratories, Abbot Park, USA) and decapitated. The head was instantaneously submerged in ice-cold carbogen saturated artificial cerebrospinal fluid (containing in mM: NaCl, 60; sucrose, 100; KCl, 2.5; NaH_2PO_4, 1.25; $NaHCO_3$, 26; $CaCl_2$, 1; $MgCl_2$, 5; glucose, 20) and the brain removed.

Horizontal 350 μm thick sections were cut with a vibratome (VT1200 S, Leica, Wetzlar, Germany, 300 μm sections for hilar recordings). To obtain maximum-connectivity-plane slices the brain was glued to its dorsal surface (compare Bischofberger et al., 2006). The slicing depth at which the temporal pole of the hippocampus first became visible was noted (depth = 0 μm). From here the first four sections were discarded (up to a depth of 1400 μm). The following two to three sections were secured such that one further section before the beginning of the dorsal hippocampus (approximately 2400 μm) could be discarded. Slices were incubated at 35 °C for 20 to 40 minutes and then stored in normal ACSF (containing in mM: NaCl, 125; KCl, 3.5; NaH_2PO_4, 1.25; $NaHCO_3$, 26; $CaCl_2$, 2.0; $MgCl_2$, 2.0; glucose, 15) at room temperature. Recordings were performed in a submerged recording chamber at 33-35 °C under constant superfusion with carbogen saturated ACSF (3 ml/min). Experiments were performed in the superior blade unless otherwise indicated.

Electrophysiological Recordings: Hippocampal dentate GCs were visually identified using

infrared oblique illumination contrast microscopy in a 20x or 60x water immersion objective

(Olympus, XLumPlanFl, NA0.95W or Nikon, N60X-NIR Apo, NA1.0W) on an upright

microscope (TriMScope®, LaVision Biotech, Bielefeld, Germany or Nikon Eclipse FN1,

Tokyo, Japan). For IPSC measurements the whole-cell patch-clamp configuration was

established with a low chloride cesium-methane-sulfonate based intracellular solution

(intracellular solution containing in mM: CH_3O_3SCs, 140; 4-(2-hydroxyethyl)-1-

piperazineethanesulfonic acid (HEPES-acid), 5; ethylene glycol tetraacetic acid (EGTA),

0.16; $MgCl_2$, 0.5; sodium phosphocreatine, 5; glucose, 10). For GC current clamp

experiments a low chloride solution (CC-intracellular solution containing in mM: K-gluconate,

140; 4-(2-hydroxyethyl)-1-piperazineethanesulfonic acid (HEPES-acid), 5; ethylene glycol

tetraacetic acid (EGTA), 0.16; $MgCl_2$, 0.5; sodium phosphocreatine, 5) was used. GCs with

input resistances greater than 300 MΩ were discarded in order to exclude immature GCs

(Schmidt-Hieber et al., 2004). Hilar cells were recorded with intracellular solution containing

in mM: K-gluconate, 140; KCL, 5; HEPES-acid, 10; EGTA, 0.16; Mg-ATP, 2; Na_2-ATP, 2; pH

adjusted to 7.25; 277mmol/kg without biocytin. 0.3% biocytin (Sigma-Aldrich, B4261). In all

imaging experiments and a subset of optogenetic experiments the intracellular solution

additionally contained 100 µM Alexa 594 hydrazide sodium salt (Life Technologies, Carlsbad,

USA). The identity of visually and electrophysiologically identified mature GC was confirmed

by their dendritic morphology after dye filling in every case tested. Pipette resistance of the

patch pipettes was 3 – 7 MΩ. Voltage-clamp recordings were performed with a Multiclamp

700B (Molecular Devices, Sunnyvale, USA) or a BVC-700A amplifier (Dagan Corporation,

Minneapolis, USA). Current-clamp recordings were performed with a Multiclamp 700B.

Voltage or current signals were digitized with a Digidata 1322A (Molecular Devices) or

(Instrutech ITC-16, Heka Electronics, Ludwigshafen, Germany) at 10 or 50 kHz and recorded

using Clampex 10.2 (Molecular Devices) or Igor Pro 6 (Wavemetrics, Lake Oswego, USA) on

a PC running Windows XP. For IPSC measurements cells were held at 0 mV including liquid-

junction potential correction (estimated at 16 mV). To aid the voltage clamp throughout the

cell, this depolarized membrane potential was slowly approached during a 15min pre-

equilibration period, during which Cs^+ entered the cell. For CC-recordings liquid junction

potential was not corrected. IPSCs were normalized to the maximally elicited IPSC over

space and power for each respective cell. Importantly, this normalization does not require

prespecification of the location or power at which a respective cell's maximum occurs. Note,

that due to this procedure all normalized IPSC values are by definition below 100%.

Chemicals for electrophysiological experiments were obtained from Sigma-Aldrich (St. Louis,

USA). All drugs were purchased from Tocris Bioscience (Bristol, UK).

Dual Patch Experiments: Two GCs within 100 µm of each other were recorded. To test for

single GC induced feedback inhibition 10 to 15 trains of 10 APs at 100 Hz were elicited by

brief (3 ms) current injections in one cell. Inhibition was monitored either in VC, while holding

the cell at 0 mV to allow the detection of small IPSCs (**Supplementary Fig. 2**, n=7 cell pairs,

7 directions) or current clamp while holding the cell at -60 mV, allowing to probe for inhibition

in both directions (not shown, n=4 cell pairs, 8 directions).

Ca^{2+} Imaging: Dye loading was modified from (Garaschuk et al., 2006) and performed in the

submerged chamber at 35°C under constant superfusion. Briefly, a dye solution containing: 1

mM Oregon Green® 488 BAPTA-1 acetoxy-methyl ester (OGB-1 AM); 2% pluronic F-

127;150mM; 2.5mM KCl; 10mM HEPES). The dye was injected into the slice along the

superior blade of the GC layer using standard patch pipettes (4-5 locations, 100 µm intervals,

30µm depth, 3 minutes at 500 mbar per location). Recordings were started at least 45

minutes after the staining procedure. Population Ca^{2+} Imaging was performed using a

multibeam two-photon fluorescence microscope (TriMScope®, LaVision Biotech, Bielefeld,

Germany) with excitation light at 810 nm. Images were acquired with a digital CMOS camera

(ORCA-Flash, Hamamatsu) through a high numerical aperture 20x water immersion

Objective (XLumPlanFl, NA-0.95, Olympus). This allowed imaging of a large field of view

(320 x 240 µm) with high spatial and temporal resolution (1920 x 1440 pixels, 20 Hz) at

acceptable signal to noise ratios. Time series were processed with ImageJ 1.48o and IGOR

Pro 6.3 in a semiautomatic manner. Regions of interest were manually placed onto all well

loaded cells which remained visible throughout the experiment. Ca^{2+} fluorescence increase

normalized to baseline ($\Delta F/F$) traces of individual cells were calculated without background

subtraction. The fraction of responders for each time series was extracted by automatic

thresholding at $\Delta F/F = 0.94\%$. The threshold was determined by combined cell-attached and

Ca^{2+} imaging experiments. Note, that for these experiments the stimulation electrode was

placed into the hilus in order to obtain a sufficient number of true positive responders. The

imaged cell population comprised on average 46 ± 18 (standard deviation) cells (n = 23

slices). The active cell fraction corresponds to the fraction of responders normalized to the

dye loaded population within each section. To assess the spatial distribution of cell activation

in imaging experiments, $\Delta F/F$ projections were created by averaging and smoothing four

frames during the transient and four frames at baseline fluorescence and then calculating the

pixel wise $\Delta F/F$.

Antidromic electrical stimulation was achieved using a bipolar cluster microelectrode (FHC,

Bowdoin, USA) connected to a digital stimulus isolator (AM-systems, Sequim, USA), placed

into stratum lucidum in the CA3 region. IPSCs at individual powers were elicited 5 to 13

times at 0.1 Hz and averaged (0.1 ms pulse time). The amplitude beyond which the stimulus

isolator could not pass the full current, determined the maximal stimulation amplitude for

each experiment.

In order to obtain the input-output relationships of the feedback inhibitory circuit data, each

variable was averaged over slices by power. This was necessary since only a small subset of

experiments in which inhibition was completely blocked could also be successfully imaged (6

of 8 sections). Due to the small numbers of active cells throughout the entire dataset with

sufficient dye loading (n = 23 slices) analysis of only these 6 slices leads to a very piecemeal

recruitment curve. A more accurate estimation of the recruitment of feedback inhibition was

obtained by averaging the cell activation and inhibition over all respectively appropriate slices

and relating them by power. Note that while the fraction of activated cells in non-MCP

sections (not included in the quantitative analysis) was mostly zero, IPSCs were almost

always present (in 28 of 29 cells in non-MCP sections).

Optogenetic Stimulation: Focal optogenetic stimulation was achieved through a

galvanometer driven spot illumination device coupled to a 473 nm DPSS Laser (UGA-40, DL-

473, Rapp Optoelectronics, Hamburg, Germany) on an upright microscope (Nikon Eclipse

FN1, Tokyo, Japan). The width of the resulting stimulation spot at the focal plane was 8.36 ±

0.04 µm (full width at half max; Nikon 10X Plan Fluor, NA 0.3 Laser powers are given in

arbitrary units from 1 to 7 corresponding to 15 ± 1 µW, 107 ± 14 µW, 292 ± 42 µW, 762 ± 105

µW, 1433 ± 49 µW, 1729 ± 165 µW and 1660 ± 163 µW at the objective (n=5

measurements). All illumination spots were placed at approximately 40 µm into the ML at the

slice surface. Stimulation pulses were of 20 ms duration.

Light Intensity distribution: To measure the light intensity distribution throughout a slice the

setup was modified to image the slice from below while the laser beam was focused to its

surface (**Fig. S1C-F**). This was achieved by focusing a surgical Microscope with 36x

magnification (M695, Leica Microsystems, Wetzlar, Germany) to the lower slice surface.

Images were taken with a CCD camera (Nikon D60). Acute sections of 100, 150, 200, 250,

300 and 350 µm thickness were cut from Prox1-ChR-eYPF mice as described above. The

laser was focused to the surface of the slice in the molecular layer and an image was taken

at every laser power (P = 1 to 7 AU). The stage was moved for every image to avoid

bleaching or phototoxicity. Linear profiles of the resulting isometric light distribution were

measured in several directions and averaged to obtain an x profile per section. The x-profiles

of slices of different thickness were then stacked to obtain the xz-profile. Values below 100

µm depth were obtained through fitting a Gaussian function in x-direction at 100 µm depth

and an exponential function in z-direction. Complete three-dimensional intensity profiles of

three different locations of two slices within the dentate molecular layer were averaged.

Calculation of the Optogenetically Activated Cell Fraction: To assess the active fraction of

GCs, approximately two GCs were recorded in cell-attached mode in each slice in which an

IPSC was recorded. Illumination spots were placed along the GC layer at 100 µm intervals

(**Fig. S1**). The entire profile was probed in triplicate with 1 s intervals between individual

locations. When the stimulation spot was in sufficient proximity to the recorded cell clear APs

were generally visible (in 25 of 26 cells), and otherwise could be induced through

simultaneous cell attached depolarization. Cell-attached spikes were detected by automatic

thresholding at 6x standard deviation of the baseline. The spatial profile of firing probabilities,

centered on the recorded cells, was averaged within each section. To test if cell activation

properties differed between blades the maximum firing probabilities (at $P = 7$) as well as the

slopes (increase in firing probability from $P = 1$ to 7) when simply averaging over all location

of a given cell were compared by t-test (n=7 sections per blade, $p = 0.490$ and 0.684 for max.

AP probability and slope, respectively). Since no difference was observed a single firing

probability distribution as a function of the distance along the GC layer (x – distance) was

calculated for each power (**Fig. S1B,** $n = 14$ sections, 7 per blade). However, the firing

probability of cells in the vicinity of the illumination spot is likely to increase not only as a

function of the laser power and spread at the surface, but also of the penetration depth of the

light cone. In order to calculate the firing probabilities throughout the slice, the firing

probability distribution at the surface was related to the measured light intensity distribution

throughout the slice (**Fig. S1C-F**; see above) utilizing a 'virtual distance' measure. Since cells

were measured at random distances from the molecular layer border, the light intensity

distribution, like the firing probabilities were collapsed to two dimensions, x-distance along

the GC layer and z-distance with increasing slice depth. The 'virtual distance' was calculated

as the mean distance from a given slice-surface pixel to all other pixels of the light intensity

distribution weighted by the intensity within those pixels (**Fig. S1G**). Assigning the firing

probabilities of pixels at the slice surface to their respective virtual distance yields the firing

probability distribution as a function of virtual distance, which was well approximated by a

gaussian fit (**Fig. S1H**). This fit was used to also calculate the firing probabilities of

pixels/cells deeper in the slice using the measured light intensity distribution as input. The

active cell fraction then corresponds simply to the mean firing probability throughout the slice.

This calculation is independent of the size and number of GC and was performed for every

power individually. We noted that a large fraction of the recorded spikes occurred with larger

latency than the typical IPSC following the beginning of the 20 ms stimulation pulse (**Fig. S1I**,

example from a single slice). Since only APs preceding the IPSC can participate in its

recruitment, we calculated the fraction of total spikes which preceded mean IPSC latency for

every power, and fitted the resulting relation with an exponential function (**Fig. S1J**). All

active cell fractions were corrected by this factor (**Fig. S1J**, bottom). Note that this does not

take account of the disynaptic delay between mossy fiber output and GC input, thereby

potentially slightly overestimating the true recruiting population. For comparison, the active

cell fraction was also computed with alternative assumptions about the decay of the firing

probability with increasing slice depth. If no firing probability decay with increasing depth is

assumed, the active cell fraction throughout the slice is given simply by the average of the

measured firing probabilities at the slice surface (**Fig. S1K**, upper grey dashed line).

Alternatively, the firing probability decay with depth was assumed to be identical to the

measured decay along the slice surface (isometric firing probability distribution; **Fig. S1K**,

lower grey dashed line). In this case, Gaussian functions were fit to the probability

distributions at the surface and these Gaussian functions were then assumed to extend also

in the z-dimension. The active GC fraction was then calculated by numerical integration

under the two dimensional Gaussian (with the bounds from 0 to 350 µm in z and -888 to 888

µm in x, which corresponds to the mean GC layer length) normalized to the same area with a

uniform firing probability of one. The best estimate of the active GC fraction, incorporating

light intensity measurements (**Fig. S1K**, black line), was within these upper and lower bound

estimates.

Comparison of Focal and Global Activation:, To globally activate the GC population a

multimode light fiber (BF-22, Thorlabs, New Jersey, USA) coupled to a 473 nm laser

(Omicron Phoxx, Rodgau-Dudenhofen, Germany) was placed above the slice surface, non-

specifically illuminating the entire hippocampus. Analogous to focal stimulations, the

activated cell fraction was calculated as the firing probability of individual cells following 20

ms pulses. Here no spatial normalization is necessary since cells were sampled from random

locations with respect to the light fiber. Firing probabilities for the focal stimulation in these

sections was calculated as the simple average of all stimulation locations.

Spatial distribution of feedback inhibition: The same stimulation paradigm which was used to

assess cell activation was used to assess the spatial distribution of feedback inhibition. For

individual cells, IPSCs at each location and power were averaged. The entire profile was

normalized to the largest measured IPSC of that cell, independent of the power and

stimulation location at which it occurred. For analysis, all IPSC profiles were spatially aligned

to the recorded cells. The mean distance to apex ± one standard deviation was 356 ± 163

μm and 322 ± 97 μm for cells from the superior and inferior blade respectively (n = 8 cells in

each blade). In order to test whether there were any distinct effects of the apex, such as a

steep decay of inhibition, which would be masked by alignment to the recorded cells, we also

aligned the profiles to the apex (not shown). However, no such effects were visible. To

analyze the saturated IPSC profiles, normalized IPSC amplitudes from P=5 to 7 were

averaged for each cell. In order to analyze the effects of local versus remote stimulation for

each blade a distance was chosen such that each remote location was still within the DG but

in the other blade (800 μm from the recorded cell). Normalized IPSCs of the three locations

surrounding the recorded cell or this remote location were averaged within each power to

obtain the IPSC amplitudes for further analysis. The cell fraction required for the activation of

a half-maximal IPSC in each section was assessed for each cell by linear interpolation

between the measured values. Since no differences were found between superior and

inferior inhibition, recordings of both blades were pooled to analyze the kinetic properties of

IPSCs. All parameters were calculated on the multiple trials of individual cells. The latency

was measured as the time from the beginning of the pulse to when the IPSC superseded 6

fold standard deviation of the baseline. The jitter was calculated as the standard deviation of

these latencies for individual cells. The rise time was calculated as the mean 20 to 80 rise

time of each cell and the decay time constant was obtained from an exponential fit to the

decaying phase of the compound IPSC.

Hilar recordings: Intrinsic properties of hilar cells were quantified based on 4.6s long

depolarizing current steps or 500ms hyperpolarizing current steps. AP threshold and fast

AHP amplitude were measured from the first AP in the first current step in which an AP

occurred within the first 10ms. Clustering fraction and mean AP time were calculated from

the current injection that elicited the maximum average AP frequency. The Clustering fraction

represents the fraction of APs that occur within 60ms before or after another AP (Larimer and

Strowbridge, 2008). Mean AP time was calculated as the mean AP time point normalized to

the duration of the current injection (4.6s). Input resistance was calculated as the slope of the

IO curve from the hyperpolarizing current ladder. Cells were manually classified as mossy

cells or interneurons based on these intrinsic properties. To objectively confirm classification

we performed unsupervised k-means clustering using scikit-learn (Pedregosa et al., 2011).

For clustering all six measures were normalized by mean and variance. Two cells with

conflicting classification were not included in further analysis.

After recording, slices were fixed for 1h in 4% PFA and stored overnight in 0.25% PBS-T at

room temperature. The following day they were transferred to PBS for short term storage or

immediately stained. For biocytin staining sections were washed with PBS and incubated

with Streptavidin-Alexa-Fluor-555 Conjugate (Invitrogen, S32355), 1:1000 in 0.25% PBS-T

overnight at 4°C. The following day they were co-stained with DAPI 1:1000 in PBS for 30

minutes and mounted with Aqua-Poly/Mount. Cells were imaged with the Leica SP8 Confocal

Microscope of the Microscopy Core Facility at the University Clinic Bonn using a 40x water

immersion objective.

Short term dynamics: Short term dynamics of GCs and hilar cells were assessed using

antidromic electrical or optogenetic stimulation at minimal power (the smallest stimulation

power that yielded reliable responses). Trains of 10 pulses at 1, 10, 30, 50Hz were delivered

in triplicate and averaged (excluding sweeps with action currents for hilar cells). In all GCs

and a subset of hilar cells we confirmed that PSCs could be blocked by at least 90% with

40µM CNQX + 50µM D-APV (n=12, 23 for GCs and hilar cells respectively). Facilitation

indices were obtained by normalizing the average of the last three PSC peaks to the first.

To test for differential dynamics between local and remote inhibition analogous trains of

optogenetic 20 ms pulses at powers below saturation (usually P=2 for local inhibition and

P=3 for remote inhibition) were delivered. For each power and frequency five repeats were

recorded and averaged. AP probabilities were assessed by cell-attached recordings with the

stimulation site close to the recorded cell. Cell-attached spikes were detected by automatic

thresholding as above.

Voltage escape estimation model.

A simple multicompartmental passive 'ball and stick' model with number of segments

following the d_lambda rule(Carnevale and Hines, 2006) and passive properties Ra = 181

Ωcm, Cm = 1 uFcm^{-2} and a leak conductance = 0.0002 Scm^{-2}, which gave an Rin of 165

MΩ, were adopted from (Carnevale and Hines, 2006; Krueppel et al., 2011). A soma (20 µm

diameter) contained one dendrite (3 µm diameter, 200 µm length) with an alpha synapse

point mechanism (Erev -90mV) placed at 180 µm from the soma. The range of synaptic

conductances (0.1 – 50 nS; adopted from (Williams and Mitchell, 2008)) elicited IPSC

amplitudes in the model, which covered the range of somatic IPSC amplitudes that were

experimentally measured (3pA – 1nA). Voltage clamp experiments were simulated using a

single electrode point mechanism at the soma (Rs 5 MOhms, to model a Rs of 15 MΩ

compensated 70%) with a holding potential of 0 mV. The transfer (Zc) and input impedance

(Zn) were determined from the model and used to calculate the actual peak IPSC amplitude

at the soma for a given synaptic conductance. Simulations were run in the Neuron 7.5

simulation environment.

Biophysically Realistic Dentate Gyrus Lamella Model: Simulations were run in python

2.7.with NEURON 7.4 (Carnevale and Hines, 2006) on Windows 7/10. We created a generic

python-NEURON interface (ouropy, Daniel Müller) which wraps NEURON's python module,

into which we ported the conductance based DG model by (Santhakumar et al., 2005).

Model code is available at https://github.com/danielmuellernai/pyDentate.

We first tuned the original model to capture our experimentally determined properties in the

most parsimonious way. During tuning we also updated some model properties to better

reflect current data and our experimental paradigm in an individual DG lamella:

We introduced a T-type Ca^{2+} channel mechanism into MCs to more realistically reflect the

depolarizing envelope at the onset of a positive current step observed in real MCs.

Furthermore, while the original model placed the perforant path input at the distal dendrite of

GCs, we moved all perforant path synapses to the middle compartment of the dendrite. In

order to be able to capture the results of convergent and divergent synaptic inputs in

sufficient resolution to produce the empirically observed activity gradations, we up-scaled cell

numbers by a factor of four. To model space, we assumed all cell types to be spread out on

a 2mm DG lamella. Since MCs project to GCs primarily outside the lamellar plane, we

removed the MC to GC connection. To allow patterned PP input we adapted PP input

specifications from (Yim et al., 2015).

We then proceeded in a first phase of model adjustment, and adapted several parameters to

reproduce our in vitro findings regarding spatial and temporal feedback inhibition

(**Supplementary Table 2**). To model frequency dependent facilitation on mossy fiber

outputs, we implemented a simple frequency dependent synapse model (tmgsyn) (Tsodyks

et al., 1998), and matched the facilitation time constant as well as the decay time constants

of individual PSCs to our experimental observations. As in the original model, each cell gives

rise to a fixed number of synaptic connections which are spatially restricted to a target pool of

adjacent cells. We tuned the size and spatial extent of this target pool to reproduce our

spatial data. To provide local inhibition we implemented a 'local' interneuron type (BC),

whose inputs and outputs were spatially restricted to an ~600μm area (as described by

(Strüber et al., 2015). To provide global inhibition we implemented a second class of

inhibitory interneurons (HC) whose inputs and outputs connect to GCs independent of space.

This simple formulation allowed us to reproduce the recruitment curves seen for local,

remote and global GC activation paradigms. To achieve plausible activity levels, we further

adapted synaptic weights similar to (Yim et al., 2015). We call the network incorporating both

spatially restricted BC synapses and mossy fiber facilitation the full tuned network. To isolate

the contribution of intrinsic GC properties to pattern separation we created a disinhibited

network by setting the synaptic weight from all interneurons to zero. We also isolated

feedforward inhibition by decreasing the mossy fiber to interneuron synaptic weight to zero.

To evaluate the effect of spatially constrained inhibition, we created a global network, where

the target pool of all interneuron was the entire GC population. To evaluate the effect of

mossy fiber facilitation we set the facilitation time constant to zero, effectively eliminating

facilitation. Details on the model parameters are summarized in **Supplementary Table 2**).

To study pattern separation, we generated 400 PP inputs. Each PP synapsed onto 100

randomly chosen GCs with the spatial connection probability being governed by a gaussian

probability distribution with standard deviation 1mm and random peak position, modeling a

full, nearly uniform input connectivity of individual afferents (Tamamaki and Nojyo, 1993). To

generate theta modulated spike patterns we used the inhomogeneous Poisson generator

from Elephant 0.5.0-Electrophysiology-Analysis-Toolkit with a 10 Hz (theta) sinusoidal rate

profile with a peak of 100 Hz, a minimum of 0 Hz and a duration of 600 ms. To generate

input patterns with varying overlap from PP afferents i = 1 to 400, we activated afferents i to

i+23 in increments of i = 1 per run. We performed 25 runs for each condition resulting in 300

unique comparisons, excluding self-comparisons. The random seed was held constant

between different runs of the same condition, resulting in differing input patterns being fed

into the same network. All randomness was generated with the python module

numpy.random.

To quantify pattern similarity we used Pearson's product moment correlation coefficient R of

the population rate vectors for input and output patterns. The population rate vector refers to

the vector of the mean firing rates of all cells in the population within the entire 600 ms

simulation, or 100 or 33 ms time windows for the time resolved analyses. All statistical

analyses of the model were performed with n=7 different random network seeds. During

Model development (tuning phase), we first ported the model by (Santhakumar et al., 2005)

with closely constrained DG cell-types, and further constrained it to reproduce our

physiological data. We then locked the model and proceeded to an (in silico) experimental

phase, in which pattern separation was investigated.

To compute full pattern separation effects (Fig. 6D) we calculated the mean R_{out} within R_{in}

bins of 0.1 and measured the area to the unity line (computed as the mean of the binwise R_{in}

– R_{out} differences). To compute isolated pattern separation effects of specific manipulations

we subtracted the respective R_{out} values with and without the manipulation, thereby obtaining

a ΔR_{out} value for each individual R_{in}. We then again computed the bin-wise mean and

quantified the area under the curve, yielding the mean ΔR_{out} analogous to the full effects.

Note, that the sequence of averaging and subtracting is irrelevant, and was inverted only to

match the figure panels. Data are displayed as mean ± SEM for each R_{in} bin (**Fig. 6E, G**).

The coefficient of variance (CoV) was calculated by normalizing the standard deviation of

ΔR_{out} within each bin by the mean of that bin, and then averaging over bins, analogous to the

previous analyses. However, only bins within $0.2 < R_{in} < 0.8$ were included, since at the borders

very small means led to unreliable results. ΔCoV represents the difference between the

mean CoV of the global (or nonfacilitating) and the tuned network models. For the temporally

resolved pattern separation analysis all measures were computed as above, but on

population vector correlations within 100 or 33 ms time bins.

Statistics and Data Analysis: Analyses were performed using ImageJ, Microsoft Excel,

Python and Igor Pro. Fits were performed using Igor Pro. Statistical analyses were performed

using GraphPad Prism 6 or Igor Pro. Comparisons were two-tailed whenever applicable.

Replicates refer to cells unless otherwise indicated (slices for imaging experiments and

network seeds for modeling data). Given the lack of previous information on effect sizes,

sample sizes were chosen according to field norms, such that only large effects can be

detected (e.g. Cohen's d>1 for paired tests). A single outlier facilitation index (Fig. 5E) during

model tuning was removed, as it was outside the triple standard deviation (due to a very

small initial IPSC). Statistical significance in Analysis of Variance (ANOVA) is indicated by §.

F-values and degrees of freedom are given as F(DFn, DFd). When ANOVAs were followed

by specific comparisons these are indicated by asterisks, where *P<.05, **P<.01 and

***P<0.001. Bargraphs and XY plots show means where error bars indicate standard error of

the mean. In boxplots error bars represent the data range and boxes the upper and lower

quartiles and the median.

References

Aimone JB, Deng W, Gage FH. 2011. Resolving new memories: a critical look at the dentate
gyrus, adult neurogenesis, and pattern separation. *Neuron* **70**:589–96.
doi:10.1016/j.neuron.2011.05.010

Alme CB, Buzzetti RA, Marrone DF, Leutgeb JK, Chawla MK, Schaner MJ, Bohanick JD,
Khoboko T, Leutgeb S, Moser EI, Moser M-B, McNaughton BL, Barnes CA. 2010.
Hippocampal granule cells opt for early retirement. *Hippocampus* **20**:1109–23.
doi:10.1002/hipo.20810

Andrews-Zwilling Y, Gillespie AK, Kravitz A V, Nelson AB, Devidze N, Lo I, Yoon SY, Bien-Ly
N, Ring K, Zwilling D, Potter GB, Rubenstein JLR, Kreitzer AC, Huang Y. 2012. Hilar
GABAergic interneuron activity controls spatial learning and memory retrieval. *PLoS
One* **7**:e40555. doi:10.1371/journal.pone.0040555

Bakker A, Kirwan CB, Miller M, Stark CEL. 2008. Pattern separation in the human
hippocampal CA3 and dentate gyrus. *Science* **319**:1640–2.
doi:10.1126/science.1152882

Barth AM, Domonkos A, Fernandez-Ruiz A, Freund TF, Varga V. 2018. Hippocampal
Network Dynamics during Rearing Episodes. *Cell Rep* **23**:1706–1715.
doi:10.1016/j.celrep.2018.04.021

Bartos M, Vida I, Frotscher M, Meyer A, Monyer H, Geiger JRP, Jonas P. 2002. Fast
synaptic inhibition promotes synchronized gamma oscillations in hippocampal
interneuron networks. *Proc Natl Acad Sci U S A* **99**:13222–7.
doi:10.1073/pnas.192233099

Berke JD, Hetrick V, Breck J, Greene RW. 2008. Transient 23-30 Hz oscillations in mouse
hippocampus during exploration of novel environments. *Hippocampus* **18**:519–529.
doi:10.1002/hipo.20435

Berron D, Schütze H, Maass A, Cardenas-Blanco A, Kuijf HJ, Kumaran D, Düzel E. 2016.

Strong Evidence for Pattern Separation in Human Dentate Gyrus. *J Neurosci* **36**.

Bischofberger J, Engel D, Li L, Geiger JRP, Jonas P. 2006. Patch-clamp recording from

mossy fiber terminals in hippocampal slices. *Nat Protoc* **1**:2075–81.

doi:10.1038/nprot.2006.312

Braganza O. 2018. Proxyeconomics, An agent based model of Campbell's law in competitive

societal systems.

Buzsáki G. 2010. Neural syntax: cell assemblies, synapsembles, and readers. *Neuron*

68:362–85. doi:10.1016/j.neuron.2010.09.023

Carnevale NT, Hines ML. 2006. The NEURON Book. Cambridge: Cambridge University

Press. doi:10.1017/CBO9780511541612

Cayco-Gajic NA, Clopath C, Silver RA. 2017. Sparse synaptic connectivity is required for

decorrelation and pattern separation in feedforward networks. *Nat Commun* **8**:1116.

doi:10.1038/s41467-017-01109-y

Cayco-Gajic NA, Silver RA. 2019. Re-evaluating Circuit Mechanisms Underlying Pattern

Separation. *Neuron* **101**:584–602. doi:10.1016/J.NEURON.2019.01.044

Chavlis S, Petrantonakis PC, Poirazi P. 2017. Dendrites of dentate gyrus granule cells

contribute to pattern separation by controlling sparsity. *Hippocampus* **27**:89–110.

doi:10.1002/hipo.22675

Chawla MK, Guzowski JF, Ramirez-Amaya V, Lipa P, Hoffman KL, Marriott LK, Worley PF,

McNaughton BL, Barnes C a. 2005. Sparse, environmentally selective expression of Arc

RNA in the upper blade of the rodent fascia dentata by brief spatial experience.

Hippocampus **15**:579–86. doi:10.1002/hipo.20091

Clelland CD, Choi M, Romberg C, Clemenson GD, Fragniere A, Tyers P, Jessberger S,

Saksida LM, Barker RA, Gage FH, Bussey TJ. 2009. A functional role for adult

hippocampal neurogenesis in spatial pattern separation. *Science* **325**:210–3.

doi:10.1126/science.1173215

Dasgupta D, Sikdar SK. 2015. Calcium permeable AMPA receptor-dependent long lasting

plasticity of intrinsic excitability in fast spiking interneurons of the dentate gyrus

decreases inhibition in the granule cell layer. *Hippocampus* **25**:269–285.

doi:10.1002/hipo.22371

de Almeida L, Idiart M, Lisman J. 2009. A second function of gamma frequency oscillations:

an E%-max winner-take-all mechanism selects which cells fire. *J Neurosci* **29**:7497–

7503. doi:10.1523/JNEUROSCI.6044-08.2009.A

Doherty J, Dingledine R. 1998. Differential Regulation of Synaptic Inputs to Dentate Hilar

Border Interneurons by Metabotropic Glutamate Receptors 2903–2910.

Espinoza C, Guzman SJ, Zhang X, Jonas P. 2018. Parvalbumin+ interneurons obey unique

connectivity rules and establish a powerful lateral-inhibition microcircuit in dentate gyrus.

Nat Commun **9**:4605. doi:10.1038/s41467-018-06899-3

Ewell L, Jones M. 2010. Frequency-tuned distribution of inhibition in the dentate gyrus. *J*

Neurosci **30**:12597–607. doi:10.1523/JNEUROSCI.1854-10.2010

Feldt Muldoon S, Soltesz I, Cossart R. 2013. Spatially clustered neuronal assemblies

comprise the microstructure of synchrony in chronically epileptic networks. *Proc Natl*

Acad Sci U S A **110**:3567–72. doi:10.1073/pnas.1216958110

Freund TF, Buzsáki G. 1996. Interneurons of the hippocampus. *Hippocampus* **6**:347–470.

doi:10.1002/(SICI)1098-1063(1996)6:4<347::AID-HIPO1>3.0.CO;2-I

Garaschuk O, Milos R, Konnerth A. 2006. Targeted bulk-loading of fluorescent indicators for

two-photon brain imaging in vivo. *Nat Protoc* **1**:380–6. doi:10.1038/nprot.2006.58

Geiger JR, Lübke J, Roth A, Frotscher M, Jonas P. 1997. Submillisecond AMPA receptor-

mediated signaling at a principal neuron-interneuron synapse. *Neuron* **18**:1009–23.

Gilbert PE, Kesner RP, Lee I. 2001. Dissociating hippocampal subregions: double

dissociation between dentate gyrus and CA1. *Hippocampus* **11**:626–36.

doi:10.1002/hipo.1077

1028 Gillespie AK, Jones EA, Lin Y-H, Karlsson MP, Kay K, Yoon SY, Tong LM, Nova P, Carr JS,

1029   Frank LM, Huang Y. 2016. Apolipoprotein E4 Causes Age-Dependent Disruption of

1030   Slow Gamma Oscillations during Hippocampal Sharp-Wave Ripples. *Neuron* **90**:740–

1031   51. doi:10.1016/j.neuron.2016.04.009

1032 Gong S, Doughty M, Harbaugh CR, Cummins A, Hatten ME, Heintz N, Gerfen CR. 2007.

1033   Targeting Cre recombinase to specific neuron populations with bacterial artificial

1034   chromosome constructs. *J Neurosci* **27**:9817–23. doi:10.1523/JNEUROSCI.2707-

1035   07.2007

1036 Gong S, Zheng C, Doughty ML, Losos K, Didkovsky N, Schambra UB, Nowak NJ, Joyner A,

1037   Leblanc G, Hatten ME, Heintz N. 2003. A gene expression atlas of the central nervous

1038   system based on bacterial artificial chromosomes. *Nature* **425**:917–25.

1039   doi:10.1038/nature02033

1040 Harney SC, Jones M V. 2002. Pre- and postsynaptic properties of somatic and dendritic

1041   inhibition in dentate gyrus. *Neuropharmacology* **43**:584–594. doi:10.1016/S0028-

1042   3908(02)00169-7

1043 Hefft S, Jonas P. 2005. Asynchronous GABA release generates long-lasting inhibition at a

1044   hippocampal interneuron-principal neuron synapse. *Nat Neurosci* **8**:1319–28.

1045   doi:10.1038/nn1542

1046 Hendrickson PJ, Yu GJ, Song D, Berger TW. 2015. Interactions between Inhibitory

1047   Interneurons and Excitatory Associational Circuitry in Determining Spatio-Temporal

1048   Dynamics of Hippocampal Dentate Granule Cells: A Large-Scale Computational Study.

1049   *Front Syst Neurosci* **9**:155. doi:10.3389/fnsys.2015.00155

1050 Hsiao Y-T, Zheng C, Colgin LL. 2016. Slow gamma rhythms in CA3 are entrained by slow

1051   gamma activity in the dentate gyrus. *J Neurophysiol* jn.00499.2016.

1052   doi:10.1152/jn.00499.2016

1053 Hsu T-T, Lee C-T, Tai M-H, Lien C-C. 2015. Differential Recruitment of Dentate Gyrus

1054   Interneuron Types by Commissural Versus Perforant Pathways. *Cereb Cortex*

26:bhv127-. doi:10.1093/cercor/bhv127

Hummos A, Franklin CC, Nair SS. 2014. Intrinsic mechanisms stabilize encoding and

retrieval circuits differentially in a hippocampal network model. *Hippocampus* **24**:1430–

1448. doi:10.1002/hipo.22324

Hunsaker MR, Rosenberg JS, Kesner RP. 2008. The role of the dentate gyrus, CA3a,b, and

CA3c for detecting spatial and environmental novelty. *Hippocampus* **18**:1064–1073.

doi:10.1002/hipo.20464

Jouhanneau J-S, Kremkow J, Poulet JFA. 2018. Single synaptic inputs drive high-precision

action potentials in parvalbumin expressing GABA-ergic cortical neurons in vivo. *Nat*

Commun **9**:1540. doi:10.1038/s41467-018-03995-2

Kapfer C, Glickfeld L, Atallah B, Scanziani M. 2007. Supralinear increase of recurrent

inhibition during sparse activity in the somatosensory cortex. *Nat Neurosci* **10**:743–53.

doi:10.1038/nn1909

Kraushaar U, Jonas P. 2000. Efficacy and Stability of Quantal GABA Release at a

Hippocampal Interneuron-Principal Neuron Synapse. *J Neurosci* **20**:5594–5607.

Krueppel R. 2010. Dendritic integration in hippocampal dentate gyrus granule cells.

Krueppel R, Remy S, Beck H. 2011. Dendritic integration in hippocampal dentate granule

cells. *Neuron* **71**:512–28. doi:10.1016/j.neuron.2011.05.043

Larimer P, Strowbridge BW. 2008. Nonrandom Local Circuits in the Dentate Gyrus. *J*

Neurosci **28**:12212–12223. doi:10.1523/JNEUROSCI.3612-08.2008

Leal SL, Yassa MA. 2018. Integrating new findings and examining clinical applications of

pattern separation. *Nat Neurosci* **21**:163–173. doi:10.1038/s41593-017-0065-1

Lee C-T, Kao M-H, Hou W-H, Wei Y-T, Chen C-L, Lien C-C. 2016. Causal Evidence for the

Role of Specific GABAergic Interneuron Types in Entorhinal Recruitment of Dentate

Granule Cells. *Sci Rep* **6**:36885. doi:10.1038/srep36885

Leutgeb JJKJ, Leutgeb S, Moser MM-BM, Moser EEI. 2007. Pattern separation in the

dentate gyrus and CA3 of the hippocampus. *Science (80-)* **315**:961–6.

doi:10.1126/science.1135801

Li L, Sultan S, Heigele S, Schmidt-Salzmann C, Toni N, Bischofberger J. 2017. Silent

synapses generate sparse and orthogonal action potential firing in adult-born

hippocampal granule cells. *Elife* **6**. doi:10.7554/eLife.23612

Lin AC, Bygrave AM, de Calignon A, Lee T, Miesenböck G. 2014. Sparse, decorrelated odor

coding in the mushroom body enhances learned odor discrimination. *Nat Neurosci*

17:559–68. doi:10.1038/nn.3660

Liu Y-C, Cheng J-K, Lien C-C. 2014. Rapid dynamic changes of dendritic inhibition in the

dentate gyrus by presynaptic activity patterns. *J Neurosci* **34**:1344–57.

doi:10.1523/JNEUROSCI.2566-13.2014

Lysetskiy M, Földy C, Soltesz I. 2005. Long- and short-term plasticity at mossy fiber

synapses on mossy cells in the rat dentate gyrus. *Hippocampus* **15**:691–6.

doi:10.1002/hipo.20096

Madar AD, Ewell LA, Jones M V. 2019. Temporal pattern separation in hippocampal neurons

through multiplexed neural codes. *PLOS Comput Biol* **15**:e1006932.

doi:10.1371/journal.pcbi.1006932

Marr D. 1971. Simple memory: a theory for archicortex. *Philos Trans R Soc Lond B Biol Sci*

262:23–81.

McHugh TJ, Jones MW, Quinn JJ, Balthasar N, Coppari R, Elmquist JK, Lowell BB,

Fanselow MS, Wilson MA, Tonegawa S. 2007. Dentate gyrus NMDA receptors mediate

rapid pattern separation in the hippocampal network. *Science* **317**:94–9.

doi:10.1126/science.1140263

McNaughton BL, Morris RGM. 1987. Hippocampal synaptic enhancement and information

storage within a distributed memory system. *Trends Neurosci* **10**:408–415.

doi:10.1016/0166-2236(87)90011-7

Miles R. 1990. Synaptic excitation of inhibitory cells by single CA3 hippocampal pyramidal

cells of the guinea-pig in vitro. *J Physiol* **428**:61–77.

Mircheva Y, Peralta MR, Tóth K. 2019. Interplay of Entorhinal Input and Local Inhibitory

Network in the Hippocampus at the Origin of Slow Inhibition in Granule Cells. *J Neurosci*

39:6399–6413. doi:10.1523/JNEUROSCI.2976-18.2019

Mizuseki K, Sirota A, Pastalkova E, Buzs?ki G. 2009. Theta Oscillations Provide Temporal

Windows for Local Circuit Computation in the Entorhinal-Hippocampal Loop. *Neuron*

64:267–280. doi:10.1016/j.neuron.2009.08.037

Myers CE, Scharfman HE. 2009. A role for hilar cells in pattern separation in the dentate

gyrus: a computational approach. *Hippocampus* **19**:321–337. doi:10.1002/hipo.20516.A

Neunuebel JP, Knierim JJ. 2014. CA3 Retrieves Coherent Representations from Degraded

Input: Direct Evidence for CA3 Pattern Completion and Dentate Gyrus Pattern

Separation. *Neuron* **81**:416–427. doi:10.1016/j.neuron.2013.11.017

Papadopoulou M, Cassenaer S, Nowotny T, Laurent G. 2011. Normalization for sparse

encoding of odors by a wide-field interneuron. *Science* **332**:721–5.

doi:10.1126/science.1201835

Pedregosa F, Varoquaux G, Gramfort A, Michel V, Thirion B, Grisel O, Blondel M,

Prettenhofer P, Weiss R, Dubourg V, Vanderplas J, Passos A, Cournapeau D, Brucher

M, Perrot M, Duchesnay É. 2011. Scikit-learn: Machine Learning in Python. *J Mach*

Learn Res **12**:2825–2830.

Pernía-Andrade AJ, Jonas P. 2014. Theta-gamma-modulated synaptic currents in

hippocampal granule cells in vivo define a mechanism for network oscillations. *Neuron*

81:140–52. doi:10.1016/j.neuron.2013.09.046

Pothmann L, Müller C, Averkin RG, Bellistri E, Miklitz C, Uebachs M, Remy S, Menendez de

la Prida L, Beck H. 2014. Function of inhibitory micronetworks is spared by Na+

channel-acting anticonvulsant drugs. *J Neurosci* **34**:9720–35.

doi:10.1523/JNEUROSCI.2395-13.2014

Pouille F, Scanziani M. 2004. Routing of spike series by dynamic circuits in the

hippocampus. *Nature* **429**:717–23.

Rangel LM, Chiba AA, Quinn LK. 2015. Theta and beta oscillatory dynamics in the dentate

gyrus reveal a shift in network processing state during cue encounters. *Front Syst*

Neurosci **9**:96. doi:10.3389/fnsys.2015.00096

Rolls ET. 2016. Pattern separation, completion, and categorisation in the hippocampus and

neocortex. *Neurobiol Learn Mem* **129**:4–28. doi:10.1016/j.nlm.2015.07.008

Rolls ET. 2013. The mechanisms for pattern completion and pattern separation in the

hippocampus. *Front Syst Neurosci* **7**:74. doi:10.3389/fnsys.2013.00074

Rolls ET. 2010. A computational theory of episodic memory formation in the hippocampus.

Behav Brain Res **215**:180–96. doi:10.1016/j.bbr.2010.03.027

Rolls ET, Treves A. 1998. Neural Networks and Brain Function - Chapter 4: Competitive

networks.

Sahay A, Scobie KN, Hill AS, O'Carroll CM, Kheirbek MA, Burghardt NS, Fenton AA,

Dranovsky A, Hen R. 2011. Increasing adult hippocampal neurogenesis is sufficient to

improve pattern separation. *Nature* **472**:466–70. doi:10.1038/nature09817

Sambandan S, Sauer J-FJ-F, Vida I, Bartos M. 2010. Associative plasticity at excitatory

synapses facilitates recruitment of fast-spiking interneurons in the dentate gyrus. *J*

Neurosci **30**:11826–37. doi:10.1523/JNEUROSCI.2012-10.2010

Santhakumar V, Aradi I, Soltesz I. 2005. Role of mossy fiber sprouting and mossy cell loss in

hyperexcitability: a network model of the dentate gyrus incorporating cell types and

axonal topography. *J Neurophysiol* **93**:437–53. doi:10.1152/jn.00777.2004

Sasaki T, Piatti VC, Hwaun E, Ahmadi S, Lisman JE, Leutgeb S, Leutgeb JK. 2018. Dentate

network activity is necessary for spatial working memory by supporting CA3 sharp-wave

ripple generation and prospective firing of CA3 neurons. *Nat Neurosci* **21**:258–269.

doi:10.1038/s41593-017-0061-5

Savanthrapadian S, Meyer T, Elgueta C, Booker S a, Vida I, Bartos M. 2014. Synaptic

properties of SOM- and CCK-expressing cells in dentate gyrus interneuron networks. *J*

Neurosci **34**:8197–209. doi:10.1523/JNEUROSCI.5433-13.2014

Schmidt-Hieber C, Jonas P, Bischofberger J. 2004. Enhanced synaptic plasticity in newly

generated granule cells of the adult hippocampus. *Nature* **429**:184–7.

doi:10.1038/nature02553

Schmidt B, Marrone DF, Markus EJ. 2012. Disambiguating the similar: the dentate gyrus and

pattern separation. *Behav Brain Res* **226**:56–65. doi:10.1016/j.bbr.2011.08.039

Severa W, Parekh O, James CD, Aimone JB. 2017. A Combinatorial Model for Dentate

Gyrus Sparse Coding. *Neural Comput* **29**:94–117. doi:10.1162/NECO_a_00905

Sik A, Penttonen M, Buzsaki G. 1997. Interneurons in the Hippocampal Dentate Gyrus□: an

In Vivo Intracellular Study. *Eur J Neurosci* **9**:573–588.

Silberberg G, Markram H. 2007. Disynaptic inhibition between neocortical pyramidal cells

mediated by Martinotti cells. *Neuron* **53**:735–46. doi:10.1016/j.neuron.2007.02.012

Skaggs WE, McNaughton BL, Wilson M a, Barnes C a. 1996. Theta phase precession in

hippocampal neuronal populations and the compression of temporal sequences.

Hippocampus **6**:149–72. doi:10.1002/(SICI)1098-1063(1996)6:2<149::AID-

HIPO6>3.0.CO;2-K

Stefanelli T, Bertollini C, Lüscher C, Muller D, Mendez P. 2016. Hippocampal Somatostatin

Interneurons Control the Size of Neuronal Memory Ensembles. *Neuron* **89**:1074–85.

doi:10.1016/j.neuron.2016.01.024

Strüber M, Jonas P, Bartos M. 2015. Strength and duration of perisomatic GABAergic

inhibition depend on distance between synaptically connected cells. *Proc Natl Acad Sci*

U S A **112**:1220–1225. doi:10.1073/pnas.1412996112

Szabo GG, Du X, Oijala M, Varga C, Parent JM, Soltesz I. 2017. Extended Interneuronal

Network of the Dentate Gyrus. *Cell Rep* **20**:1262–1268.

doi:10.1016/j.celrep.2017.07.042

Tamamaki N. 1997. Organization of the entorhinal projection to the rat dentate gyrus

revealed by Dil anterograde labeling. *Exp brain Res* **116**:250–8.

Tamamaki N, Nojyo Y. 1993. Projection of the entorhinal layer II neurons in the rat as

revealed by intracellular pressure-injection of neurobiotin. *Hippocampus* **3**:471–480.

doi:10.1002/hipo.450030408

Temprana SG, Mongiat LA, Yang SM, Trinchero MF, Alvarez DD, Kropff E, Giacomini D,

Beltramone N, Lanuza GM, Schinder AF. 2015. Delayed Coupling to Feedback

Inhibition during a Critical Period for the Integration of Adult-Born Granule Cells. *Neuron*

85:116–131. doi:10.1016/j.neuron.2014.11.023

Toth K, Suares G, Lawrence JJ, Philips-Tansey E, McBain CJ. 2000. Differential

mechanisms of transmission at three types of mossy fiber synapse. *J Neurosci*

20:8279–89.

Trappenberg TP. 2010. Fundamentals of computational neuroscience. *Oxford Univ Press.*

doi:10.5860/CHOICE.40-2778

Trimper JB, Galloway CR, Jones AC, Mandi K, Manns JR. 2017. Gamma Oscillations in Rat

Hippocampal Subregions Dentate Gyrus, CA3, CA1, and Subiculum Underlie

Associative Memory Encoding. *Cell Rep* **21**:2419–2432.

doi:10.1016/J.CELREP.2017.10.123

Tsodyks M, Pawelzik K, Markram H. 1998. Neural Networks with Dynamic Synapses. *Neural*

Comput **10**:821–835. doi:10.1162/089976698300017502

van Dijk MT, Fenton AA. 2018. On How the Dentate Gyrus Contributes to Memory

Discrimination. *Neuron* **98**:832-845.e5. doi:10.1016/j.neuron.2018.04.018

Verret L, Mann EO, Hang GB, Barth a M, Cobos I, Ho K, Devidze N, Masliah E, Kreitzer a

C, Mody I, Mucke L, Palop JJ. 2012. Inhibitory interneuron deficit links altered network

activity and cognitive dysfunction in Alzheimer model. *Cell* **149**:708–721.

doi:10.1016/j.cell.2012.02.046

Wick SD, Wiechert MT, Friedrich RW, Riecke H. 2010. Pattern orthogonalization via channel

decorrelation by adaptive networks. *J Comput Neurosci* **28**:29–45. doi:10.1007/s10827-

009-0183-1

Wiechert MT, Judkewitz B, Riecke H, Friedrich RW. 2010. Mechanisms of pattern

decorrelation by recurrent neuronal circuits. *Nat Neurosci* **13**:1003–1010.

doi:10.1038/nn.2591

Williams SR, Mitchell SJ. 2008. Direct measurement of somatic voltage clamp errors in

central neurons. *Nat Neurosci* **11**:790–798. doi:10.1038/nn.2137

Yim MY, Hanuschkin A, Wolfart J. 2015. Intrinsic rescaling of granule cells restores pattern

separation ability of a dentate gyrus network model during epileptic hyperexcitability.

Hippocampus **25**:297–308. doi:10.1002/hipo.22373

Yu J, Swietek B, Proddutur A, Santhakumar V. 2015. Dentate total molecular layer

interneurons mediate cannabinoid-sensitive inhibition. *Hippocampus* **25**:884–9.

doi:10.1002/hipo.22419

Yuan M, Meyer T, Benkowitz C, Savanthrapadian S, Ansel-Bollepalli L, Foggetti A, Wulff P,

Alcami P, Elgueta C, Bartos M. 2017. Somatostatin-positive interneurons in the dentate

gyrus of mice provide local- and long-range septal synaptic inhibition. *Elife* **6**.

doi:10.7554/eLife.21105

Zhang W, Yamawaki R, Wen X, Uhl J, Diaz J, Prince D a, Buckmaster PS. 2009. Surviving

hilar somatostatin interneurons enlarge, sprout axons, and form new synapses with

granule cells in a mouse model of temporal lobe epilepsy. *J Neurosci* **29**:14247–56.

doi:10.1523/JNEUROSCI.3842-09.2009

1235

1236

Acknowledgements:

This work was supported by the Deutsche Forschungsgemeinschaft (SFB 1089, ebGluNet), the BONFOR program of the University of Bonn Medical Center, the ERANET Neuron grant 'EpiNet' (to H.B.). We thank Olivia van Ray and Dominik Holtkamp for excellent technical assistance; Thoralf Opitz, Holger Dannenberg, Laura Ewell and Dirk Dietrich for valuable comments and Stefan Remy for access to multi-beam two-photon microscopy. We further acknowledge support by the Microscopy Core Facility of the Medical Faculty.

Author contributions

H.B. and O.B. conceived of the study and wrote the manuscript. D.M. performed hilar experiments, created the computational model and contributed to the manuscript. T.K. contributed technical and analytical expertise and modeled voltage escape errors. All other experiments and analyses were performed by O.B.

Competing interests

The authors declare no competing interests.

1252 **Supplementary Materials**

1253

Fig. S1 (associated to Fig. 1), Detection of single action potential induced calcium transients, A
section of the dentate gyrus was loaded with OGB1-AM and imaged with multibeam two-photon
microscopy while antidromically eliciting action potentials and recording from individual cells in cell-
attached mode. **A)** A schematic illustration of the experimental setup. **B)** Example of OGB1-AM loaded
GCs. Scale bar: 10 μm **C)** Cells were stimulated with a single pulse (left) or bursts of 5 pulses at 30 Hz
(middle) or 100 Hz (right). Cell attached recordings revealed the exact number of induced action
potentials (bottom) which could then be correlated with the intracellular calcium signal (middle). **D)**
Superposition of the calcium fluorescence traces of 49 recorded cells constituted of cells identified as
responders (green) or non-responders (grey) by cell attached recordings. **E)** Peak ΔF/F for single APs
of identified responders and non-responders plotted against their respective baseline fluorescence
(left). A histogram of the peak ΔF/F of both groups fitted with a Gaussian distribution of the non-
responders (right, scale bar = 5 cells). The dashed line indicates detection threshold at the quadruple
standard deviation of this fit (0.94% ΔF/F). **F)** False positive (gray) and false negative (green) rates
were plotted as a function of the detection threshold and fitted with sigmoidal functions. A detection
threshold of 0.94% leads to exactly equal numbers of false positives and false negatives at a true
positive rate of 3% (inset, dashed lines). **G)** To test for potential effects of variable dye loading on
detection efficacy, we tested for a correlation between peak ΔF/F of responders and baseline
fluorescence intensity (p>0.05). **H)** -To test if increasing numbers of responders at increasing
stimulation power led to increases of false positives in the densely packed GC layer we correlated
peak ΔF/F of non-responders with stimulation power (p>0.05). Dashed lines in (G) and (H) represent
the 95% confidence intervals of linear regressions.

1275

1276

Fig. S2 (associated to Fig. 2), Optogenetically activated cell fraction. A) Schematic illustration of the experimental setup. Cells were recorded in cell attached mode (2 per slice), while systematically stimulating at varying distances. Traces from a representative trial at P = 3. **B)** Mean firing probability of every location over trials and cells for each laser power (3 example powers shown). **C)** Schematic of the modified setup to record the 3 dimensional light intensity profile in an acute slice. In order to avoid saturation a neutral density filter (ND4) was inserted into the light path. **D)** Cross section of the light intensity profile of the laser spot at increasing slice depth. The dashed white lines indicate the location of the cross sections shown in (E) and (F). Depths below 100 μm were extrapolated from fits to (E) and (F). **G) top,** Illustration of the calculation of the virtual distance for a particular cell/pixel 440μm lateral to the laser focus. The distances between the given cell/pixel and all other pixels (individual xz-distances) were weighted by the intensity at those pixels. **Bottom,** This weighting is illustrated by a histogram displaying the intensities for each respective xz-distance. The virtual distance corresponds to the intensity weighted mean of xz-distances. **H)** The measured firing probabilities were assigned to the respective virtual distances. The resulting firing probability distribution was well approximated by a Gaussian fit (black lines). **I)** Example of the IPSC and AP latencies upon a stimulation pulse from an individual slice. Laser Powers are color coded. **J) Top**, Example Histogram of the distribution of all AP latencies for P = 3 (blue). The black bar indicates the mean IPSC latency ± standard deviation at that power. **Bottom,** The fraction of action potentials that precede the mean IPSC for each power was well approximated by an exponential fit (black line). Light stimulation in (I) and (J) was from 0 to 20 ms. **K) black,** Estimated active cell fraction in the slice calculated from the light intensity profiles in (D) and the virtual firing probability distributions in (H) and corrected by the fraction of APs occurring after the mean IPSC (J). The estimated active cell fraction is identical to the mean firing probability throughout the slice. For comparison the cell fraction was also estimated assuming no firing probability decay with increasing depth (upper grey dashed line) or assuming isometric decay

1277

1278

Fig. S3 (associated to Figs. 1,2), Error in somatic IPSC measurements with increasing inhibitory conductance. A simple ball and stick model was used to estimate the impact of voltage escape errors for dendritic IPSCs (soma diameter 20um; dendrite diameter and length 3 µm and 200 µm, respectively). To estimate the maximum errors the inhibitory synapse was placed at a distal site (180 µm from the soma) and inhibitory currents were measured using a single electrode voltage-clamp at the soma. **A)** Illustration of the model and an attenuated somatic IPSC measurement. **B,** Peak amplitudes of the measured IPSC over a range of distal synaptic conductances (measured peak IPSC), as well as the actual peak IPSC in the absence of voltage errors, calculated from the transfer and input impedances of the model. **C)** Error in somatically measured peak IPSC as percentage of the actual peak IPSC (I syn, % error) at a given synaptic conductance. Errors in estimating synaptic inhibitory currents were linear. **D,** Illustration of corrected and uncorrected recruitment curves of absolute IPSC amplitudes for varying degrees of voltage error (using data from the recruitment curve in **Fig. 2H**). Note that due to the linearity of voltage escape errors, absolute IPSC amplitudes change, but the saturation point does not. **E)** This is illustrated by the normalized recruitment curves, as shown throughout the manuscript (see i.e. **Fig. 2H**). Note that normalized curves are practically unaffected by voltage escape errors.

1279

1280

Fig. S4 (associated to Figs. 1,2), Absence of single GC induced feedback inhibition. Pairs of juxtaposed GCs (< 100µm distance) were recorded to test for single GC induced feedback inhibition. **A)** Schematic illustration of the experimental setup. **B)** Example of a paired recording where cell 1 is fired at 100 Hz in current clamp mode while cell 2 is recorded in voltage clamp mode in order to detect IPSCs. (gray, 10 individual trials; black, average).

1281

1282

1283

Fig. S5 (associated to Fig. 3,4), Frequency dependence of feedback inhibition over space.
Trains of 10 focal optic stimulations (20ms duration) were applied either locally (1) or remotely (2)
to elicit feedback inhibition. **A)** Schematic of the experimental paradigm and example traces of
elicited cell attached spikes or IPSCs. **B)** Example traces for stimulation at 1, 10, 30 Hz or
continuously for 200 ms of a local or remote GC populations (black and grey, respectively). **C)** The
AP probability index (mean probability during the last three pulses normalized to the first pulse
(one-way RM ANOVA, P<.001 for frequency, Bonferroni corrected Wilcoxon signed rank test for
deviation from 1; P=.024, =.008, =.008 and =.008 for 1Hz, 10Hz, 30Hz and continuous stimulation
respectively). **D)** Facilitation indices of local (dark grey) and remote (light grey) stimulation (two-
way RM ANOVA; P=.635, =.314 and =.687 for location, frequency and interaction respectively).

1284

1285

Fig. S6 (associated to Fig. 5,6), Model Tuning and Validation. A) Frequency responses to somatic current injections of model celltypes (GC: granule cells, BC: basket cells, HC: hilar perforant path associated cells, MC: mossy cells). All model cells were matched to data by Santhakumar et al., (2005). **B)** Simulation of synchronous frequency stimulation of GCs and the resulting PSCs in modelled cell types, analogous to Fig. 4. **C)** Representative theta modulated PP input and population responses (scatterplots) of all modelled cell types. Following each scatterplot are three examples of spiking cells of the respective type. **D)** Same as (C) but for slow gamma modulated inputs.

1286

Fig. S7 (associated to Fig. 6), Robustness over different Similarity Metrics, To test if the main finding of frequency dependent pattern separation, particularly for highly similar inputs, depended on the similarity metric used, the original data was reanalyzed with two alternative similarity metrics. As in the main figures data points in bar graphs represent the 7 independent network seeds. For each network (seed) we ran sets of patterns for the full model, a network without feedback inhibition (no FB) and a model with no inhibition (no inh.). **A)** Examples of the theta (10Hz) and gamma (30Hz) modulated perforant path (PP) input spike patterns. **B)** Mean number of APs per active PP afferent. **C)** Fraction of active GCs across the different conditions. **D)** Mean number of APs per active GC across conditions. **E)** Pearson's correlation coefficient R (as in Fig. 6) **left:** exemplary scatterplots of pattern separation effects. **right:** bargraphs of the mean pattern separation effect over the full input similarity range (0<Rin1) or only highly similar input patterns (0.9<R<1). Full effects measure the mean pattern separation effect for each network condition: full model, no feedback (no FB) and no inhibition (no inh.). Isolated effects measure the pattern separation contribution of individual circuit motifs: feedback inhibition (FB) and feedforward inhibition (FF). **F)** same as (E) but using normalized dot product (NDP) as similarity metric for input and output comparisons. **G)** Same as (E) but using population overlap as similarity metric. Overlap is defined as the number of cells active in both patterns (logical and) divided by the number of cells active in either pattern (logical or). The full effects were tested with 2x3 ANOVAs followed by Sidak's postests for differences between conditions. Isolated effects were tested with 2x2 ANOVAs followed by Tukey posttests for differences between frequencies. Asterisks indicate significance in posttests given significant overall effects (*P<.05,**P<0.01,***P<0.001).

1289

1290

1291

Fig. S8 (associated to Fig. 6), Isolated pattern separation effects of spatial tuning and MF facilitation, Effects of isolated manipulations were computed for the DG model as in Fig 6. **A)** Schematic of the full tuned network and the resulting spatial profile of inhibition (as in Fig. 5). **B)** Schematic of the global network (with unrestricted BC target pool) and the resulting spatial profile of inhibition. **C)** Schematic of the non-facilitating network. **D)** Isolated mean feedback effects of the global, tuned and non-facilitating models. Two-way RM ANOVA showed: p<0.001, =0.020, =0.402 for frequency, condition and interaction respectively with * indicating significance in Dunnett's posttest against the full tuned effect. p=0.742 and 0.020 for global and non-facilitating, respectively at 10Hz; p=0.650 and 0.001 for global and non-facilitating, respectively at 30Hz. **E)** Exemplary pattern separation plots of theta modulated inputs when spatial tuning (left) or MF facilitation (right) was removed. **F)** Isolated pattern separation effects of the given manipulation for theta (blue) or gamma (green) modulated inputs as a function of input similarity. **G)** Isolated effect of the given manipulation on mean ΔR_{out} (left) and the coefficient of variance (ΔCoV) of pattern separation between individual comparisons (right). **H, I)** Time-resolved analyses of isolated effects of spatial tuning (left) and MF facilitation (right) for theta (top row) and slow gamma (bottom row) modulated inputs. In each subpanel the bottom left and middle insets show ΔR_{out} as a function of input similarity of the first and last time windows respectively. The bottom right insets show the evolution of the mean ΔR_{out} over time.

1292

1293

Fig. S9 (associated to Fig. 6), Robustness for shorter analysis time-window, A). 33ms time-resolved pattern separation effects of the full model, isolated feedback (FB) or feedforward (FF) inhibition for theta modulated input (10Hz). All analyses were performed as above but with rate vector correlations computed for 33 ms time windows (instead of 100 ms or 600 ms, as in Fig.6). The bottom insets show ΔR_{out} as a function of input similarity for the first and last three time windows. The bottom right insets show the evolution of the mean ΔR_{out} over time. **B)** Same as (A) but for slow gamma (30Hz) modulated inputs. Data represent mean ± SEM of n=7 random network seeds.

1294

1295

Fig. S10 (associated to Fig. 6), Robustness over various IPSC decay time-constants and over the full gamma range, A-C) To test if the frequency dependence of feedback inhibitory pattern separation remained robust for different IPSC decay time constants we probed a range of altered time constants (our experimentally matched time constant x0.5, x1, x2 and x5) while maintaining total inhibitory conductance in the network constant by complementary adjustment of IPSC amplitude. As we expected a potential interaction between IPSC decay and modulation frequency, we probed model runs for each factor with 10Hz, 30Hz and 60Hz modulation. The isolated feedback inhibitory effects were computed and impacts of decay and frequency were examined with 4x3 ANOVAs followed by Tukey posttests for differences between frequencies. Asterisks indicate significance in posttests given significant overall effects (*P<.05,**P<0.01,***P<0.001).**A)** Illustration of modified IPSC time-courses. **B)** Mean pattern separation effect of isolated feedback inhibition over the full input similarity range ($0<R_{in}<1$). **C)** Same as (B) but only for highly similar input patterns. Analyses in A-C were performed on 7 new network seeds with simulation and analysis otherwise identical to Fig.6. **D-E)** To probe the robustness of frequency dependent feedback inhibitory pattern separation over an even larger range of frequency modulation we next simulated the effects over a range from 10 to 100Hz in 5Hz steps. To provide computational tractability we performed only 8 runs per frequency (instead of 24 runs as in all other simulations) leading to fewer pattern comparisons, and somewhat noisier readouts. For the majority of frequencies, no input comparisons with R>0.9 occurred so we defined ($0.8<R_{in}<1$) as highly similar input patterns, potentially leading to a slight underestimation of our

1296   effects.

1297

1298

Fig. S11 (associated to Fig. 6), Robustness for increased feedforward inhibition, To test if the frequency dependent enhancement of feedback inhibitory pattern separation of highly similar inputs was sensitive to the changes in the relative strengths of feedforward and feedback inhibition, we increased the perforant path (PP) to basket cell (BC) synapse weight 2x. **A)** Illustration of the network alteration. **B)** The resulting full pattern separation effects (left) and isolated feedback (FB) and feedforward (FF) effects (right) as mean over all input similarities. **C)** Same as (B) but only for highly similar input patters. Full effects were tested with 2x3 ANOVAs followed by Sidak's postests for differences between conditions. Isolated effects were tested with 2x2 ANOVAs followed by Tukey posttests for differences between frequencies. Asterisks indicate significance in posttests given significant overall effects (*P<.05,**P<0.01,***P<0.001).

1299

1300

	From	To	IN location	Method	Freq.	short term dynamics	pulses	species	Authors	
GC to IN	GC	BC	GC-border	dual patch	50 Hz	PPD	2	juv. wistar rats	Geiger and Jonas (1997)	
	GC	BC	not stated	dual patch	20 & 50 Hz	depression	10 & 5	C57/BL6 mice	Espinoza et al. (2019)	
	GC	SST	not stated	dual patch	20 Hz	facilitation	10	C57/BL6 mice	Espinoza et al. (2019)	
e-stim to IN	PP & C/A	FS	GC-border	e-stim/ spike prob.	50 Hz	decreased spike prob.	10	juv. SD rats	Liu et al. (2014)	q
	PP	FS	GC-border	e-stim	10 Hz	depression to 40%	20	juv. C57/BL6 mice	Ewell and Jones (2010)	q
	PP & C/A	non-FS	GC-border	e-stim/ spike prob.	50 Hz	increased spike prob.	10	juv. SD rats	Liu et al. (2014)	q
	PP & MF	FS	GC-border	e-stim	30 Hz	facilitation		juv. wistar rats	Sambandan et al. (2010)	
	MF	FS	GC-border	e-stim (close)	30 Hz	facilitation, PPF		juv. wistar rats	Dasgupta et al (2015)	
e-stim to MC	MF	MC	Hilus	e-stim	20 Hz	facilitation to 500%	5	juv. wistar rat	Lysetskiy (2005)	q
IN to GC	BC	GC	GC-border	dual patch	10 & 50 Hz	depression to ~25% & 0%	> 850	juv. wistar rats	Kraushaar and Jonas (2000)	q
	CCK & PV	GC	GC-border	dual patch	50 Hz	depression to 20%	10	juv. wistar rats	Hefft and Jonas (2005)	q
	SST	GC	Hilus	dual patch	50 Hz	depression to 20%	20	adult GIN mice	Zhang et al. (2009)	q
	BC	GC	GC-border	dual patch	20 Hz	PPD to 39%	2	juv. C57/BL6 mice	Bartos et al. (2002)	q
	FS	GC	GC-border	dual patch	10 Hz	PPD to 85%	2	juv. SD rats	Harney and Jones (2002)	q
	non-FS	GC	GC-border	dual patch	10 Hz	PPD to 77 %	2	juv. SD rats	Harney and Jones (2002)	q
	PP	GC	GC-border	e-stim	10 Hz	depression to 40%	20	juv. C57/BL6 mice	Ewell and Jones (2010)	q
	FS	GC	GC-border	dual patch	25 Hz	depression	5	juv. SD rats	Liu et al. (2014)	
	non-FS	GC	GC-border	dual patch	25 Hz	facilitation	5	juv. SD rats	Liu et al. (2014)	
	CCK	GC	not stated	dual patch	20 Hz	depression	10	C57/BL6 mice	Espinoza et al. (2019)	
IN to IN	BC	BC	GC-border	dual patch	50 Hz	depression to 50%	10	juv. wistar rats	Savanthrapadian et al. (2014)	q
	HICAP	HICAP	GC-border	dual patch	50 Hz	facilitation to 160%	10	juv. wistar rats	Savanthrapadian et al. (2014)	q
	HIPP	HIPP	GC-border	dual patch	50 Hz	facilitative envelope	10	juv. wistar rats	Savanthrapadian et al. (2014)	q
	HICAP	BC	GC-border	dual patch	50 Hz	no change	10	juv. wistar rats	Savanthrapadian et al. (2014)	q
	HIPP	BC	GC-border	dual patch	50 Hz	no change	10	juv. wistar rats	Savanthrapadian et al. (2014)	q
	BC	BC	GC-border	dual patch	20 Hz	PPD to 26%	2	juv. C57/BL6 mice	Bartos et al. (2002)	q
	TML (non-FS)	TML	GC-border	dual patch	50 Hz	facilitation , PPF to ~220%	8	adult wistar rats	Yu et al. (2015)	
	IN	IN	Hilus	dual patch	20 Hz	PPD to ~80%	2	juv. SD rats	Larimer and Strowbridge (2008)	q
MC	IN	MC	Hilus	dual patch	20 Hz	PPD to ~80%	2	juv. SD rats	Larimer and Strowbridge (2008)	q
	MC	IN / MC	Hilus	dual patch	20 Hz	no change	2	juv. SD rats	Larimer and Strowbridge (2008)	q

cells			other	
BC	basket cell		blue	depression, quantitatively described
CCK	cholecystokinin expressing cell		green	facilitation, quantitatively described
FS	fast spiking cell (> 50Hz peak firing frequency)		PPD	paired pulse depression
GC	granule cell		PPF	paired pulse facilitation
HICAP	hilar commissural/associational path associated cell		e- stim	extracellular electrical stimulation
HIPP	hilar perforant path associated cell		C/A	commissural/ associational
IN	not further categorized interneuron		MF	mossy fiber
MC	mossy cell		PP	perforant path
non- FS	non- FS (< 50Hz peak firing frequency)		SD rat	spraque dawley rat
SST	somatostatin expressing cell		juv.	juvenile (12 to 25 days postnatally)
TML	total molecular layer cell		q	quantitatively described

1301

Supplementary Table 1, Literature review for DG circuit short term dynamics, Studies reporting short term dynamics within the DG circuit were reviewed with a main focus on facilitation or depression of synaptic connections defined by pre and postsynaptic cell types. Note the abundance of depressing synapses (quantitative descriptions of depression blue). Also note the complexity of direct connections between Interneurons (lower third of the table).

1302

1303

	To			
PP	**GC, 2000**	**MC, 60**	**BC, 24**	**HC, 24**
Weight (nS)	1		1 (2)	
Decay Tau (ms)	10		6.3	
Facilit. Tau (ms)	0		0	
Delay (ms)	0	/	0	/
Target pool	2000		24	
Divergence	100		1 BC/PP	
Target segments	mid. dendrite		distal dend	
GC				
Weight (nS)		2	2.5	2.5
Facilitation Max.		10x	10x	10x
Decay Tau (ms)		7.6	8.7	8.7
Facilit. Tau (ms)		500	500	500
Delay (ms)	/	1.5	0.8	1.5
Target pool		12	8	24
Divergence		1:1	1:1	1:1
Target Segments		prox. dendrite	prox. dendrite	prox. dendrite
MC				
Weight (nS)		0.5	0.3	0.2
Decay Tau (ms)		2.2	2	6.2
Facilit. Tau (ms)		0	0	0
Delay (ms)	/	2	3	3
Target pool		24	12	20
Divergence		1:3	1:1	1:2
Target segments		prox. dendrite	prox. dendrite	mid. dendrite
BC				
Weight (nS)	1.2	1.5	7.6	
Decay Tau (ms)	20 (10, 40, 100)	3.3	1.8	
Facilit. Tau (ms)	0	0	0	
Delay (ms)	0.85	1.5	0.8	/
Target pool	560	28	12	
Divergence	1:400	1:3	1:2	
Target segments	soma	prox. dendrite	prox. dendrite	
HC				
Weight (nS)	6	1.5	0.5	
Decay Tau (ms)	20 (10, 40, 100)	6	5.8	
Facilit. Tau (ms)	0	0	0	
Delay (ms)	3.8	1	1.6	/
Target pool	2000	60	24	
Divergence	1:640	1:4	1:4	
Target segments	distal dend	mid dendrite 1, 2	distal dend	

From

1304

Supplementary Table 2, Model Parameters, Overview of synaptic and intrinsic parameters between model cell-types. First row includes modeled cell number per type. **PP**: perforant path, **GC**: granule cell, **MC**: mossy cell, **BC**: basket cell, **HC**: Hilar perforant path associated cell; Weight: maximal synaptic conductance, Facilitation Max.: maximal fold increase of synaptic conductance, Decay Tau: synaptic decay time constant, Facilit. Tau: facilitation time constant, Delay: latency to postsynaptic event after presynaptic action potential, Target pool: range of n closest cells potentially receiving an output, Divergence: number of output synapses per cell stochastically picked from target pool, Target segments: cellular compartment receiving the synapse. Values in brackets are values for robustness analyses in Figs. S10, S11.

1305