Title: GPI-anchored SKU5/SKS are maternally required for integument

development in Arabidopsis

Running title: SKU5/SKS are required for integument development

Authors: Ke ZHOU[1,3] (zhouke0525@gmail.com), Elisa FIUME[2] (Elisa.Fiume@gmail.com), Delphine DE-

VOS[2] (delphine.de-vos@inra.fr) and Enrico MAGNANI[2] (enrico.magnani@inra.fr)

Affiliations: 1. FAFU-UCR Joint Center for Horticultural Biology and Metabolomics, Haixia Institute of

Science and Technology, Fujian Agriculture and Forestry University, Fuzhou, 350002, China

2. Institut Jean-Pierre Bourgin, INRA, AgroParisTech, CNRS, University of Paris-Saclay, 78026 Versailles

Cedex, France

3. Institut des Sciences du Végétal, UPR2355, CNRS, Saclay Plant Sciences, 91198 Gif sur Yvette Cedex,

France.

Corresponding author: Ke ZHOU, (zhouke0525@gmail.com), FAFU-UCR Joint Center for Horticultural

Biology and Metabolomics, Haixia Institute of Science and Technology, Fujian Agriculture and Forestry

University, Fuzhou, 350002, China

CRCID: https://orcid.org/0000-0002-5883-2550

ABSTRACT

Glycosylphosphatidylinositol-anchored proteins (GPI-APs) play crucial roles in various processes in eukaryotes. In *Arabidopsis*, *SKS1*, *SKS2*, *SKS3* and *SKU5* from *SKU5/SKS* gene family could encode GPI-anchored proteins, and they were recently reported to regulate cell polar expansion and cell wall synthesis redundantly in roots. Here, we report that, they are also redundantly crucial for seed production and seed morphogenesis in *Arabidopsis* through regulating maternal integument development. Their *loss-of-functions* resulted in disrupted development of integuments that failed to protect embryo sacs from exposure to external space due to physical restriction, and lead to female gametophytic abortion. Interestingly, those less defective ovules could be fertilized and develop into seeds normally, however, their seed morphogenesis was largely affected.

Our research made SKS1, SKS2, SKS3 and SKU5 be not only the first class of GPI-anchored proteins that could regulate maternal integument development, but also the first class of proteins that could regulate cell polar expansion in both root and integument cells besides several MAPK cascade components. Our study also underlined the regulation by integument development in reproductive processes.

KEY WORDS

Glycosylphosphatidylinositol; GPI-anchored protein; GPI-AP; Integument Development; Seed Morphogenesis; *Arabidopsis*

INTRODUCTION

In eukaryotes, various protein precursors could be modified during maturation in endoplasmic reticulum and Golgi bodies, where their special hydrophobic carboxyl-terminal signal polypeptides could be cleaved and catalyzed to be replaced by a common glycosylphosphatidylinositol (GPI) oligosaccharide structure. It covalently links the lipid bilayer and the carboxyl-terminus of glycosylphosphatidylinositol-anchored proteins (GPI-APs), which allows these proteins to be attached on outer surface of plasma membrane (Kinoshita and Fujita, 2016; Oxley and Bacic, 1999; Stevens, 1995; Zhou, 2019b). Disruption of GPI moieties biosynthesis resulted in broad and severe defects during various developmental processes (Bundy et al., 2016; Gillmor et al., 2005; Kawagoe et al., 1996; Kinoshita, 2014).

In *Arabidopsis*, more than 300 GPI-APs from diverse families have been identified or predicted by bioinformatics or proteomic tools (Borner et al., 2003; Takahashi et al., 2016), among which, only a small proportion of them has been characterized of their involvement in cell wall components synthesis, catalytic processes, signaling transduction, stress and pathogen response, and so on (Zhou, 2019b). But to date, no particular GPI-AP has been reported to regulate integument development in plants.

SKU5-Similar (*SKU5/SKS*) gene family, named after *SKU5*, which has been reported to be involved in twisting bias of root growth in *Arabidopsis*, contains 19 members (Rutherford and Masson, 1996; Sedbrook et al., 2002). *SKS1*, *SKS2*, *SKS3* and *SKU5* (*GPI-SKU5/SKS*) within the same subgroup of this family could encode GPI-anchored proteins, and they were reported by us to be redundant essential for root development through affecting anisotropic cell expansion in root and cell wall synthesis, and *SKS1* made significant contribution during this process (Zhou, 2019a; Zhou, 2019c).

In this study, we further investigated the involvement of this subgroup of *GPI-SKU5/SKS* genes in reproductive processes of *Arabidopsis*. *Loss-of-functions* redundantly and drastically reduced their seed production, which was demonstrated to be due to maternal-dependent ovule abortions. In those aborted ovules, the extension of their maternal integuments was disturbed and resulted in physical restriction to female gametophytes, which lead to their exposure to out space and consequent female gametophytic abortion. Although those less defective ovules could be fertilized and develop into seeds, their seed morphogenesis was largely affected as the aberrant integument/seed coat development.

The interference on integument development seemed due to similar defective polar expansion of integument cells as we described in their root cells previously, which is very surprising, as to our knowledge, none of gene has been reported to interfere polar expansion on both root and integument cells besides a series of MAPK components.

Our study firstly reported the crucial involvement of particular GPI-APs, GPI-SKU5/SKS proteins, in reproductive processes through regulating maternal integument development, which increased functional

diversity of GPI-APs in *Arabidopsis*. Surprisingly, this group of GPI-APs could regulate cell polar

expansion in both root and integument the same as a series of MAPK components.

MATERIALS AND METHODS

Pseudo-Schiff Propidium Iodide Staining

Ovules and seeds were harvested and then fixed in greenfix solution (Diapath) for 1 to 3 h at room

temperature. Fixed samples were transferred into water and kept at 4°C for up to 7 d. The samples were

rinsed once more with water and incubated in a 1%SDSand 0.2 N NaOH solution at 37°C for 1 h. The

samples were rinsed in water and incubated in a a-amylase solution (0.4 mg/mL a-amylase, 20 mM $NaPO_4$,

and 6.7mMNaCl, pH 6.9) at 37°C for 2 h. The samples were rinsed in water and incubated in a 12.5%

bleach solution (1.25% active Cl⁻) for 10 to 15 min. The samples were rinsed in water and then transferred

to a 1% periodic acid solution at room temperature for 20 min. The samples were rinsed in water and

incubated in a Schiff reagent and propidium iodide solution (100 mM sodium metabisulfite, 0.15 N HCl,

and 100 mg/mL propidium iodide) until plants were visibly stained. The samples were rinsed in water and

then incubated in achloral hydrate solution (4gchloral hydrate, 1mL glycerol, and 2 mL water) at room

temperature for 30min (Fiume et al., 2017; Xu et al., 2016). Finally, the samples were transferred onto

microscope slides for confocal laser scanning microscopy analysis (Leica SP5)

Mature embryo sac observation

The oldest closed flower buds of a given inflorescence was emasculated and harvested two days later. The

collected pistils were fixed in ethanol: acetic acid (9:1) solution overnight, and followed by two washes in

90% and 70% ethanol. Then they were cleared by a chloral hydrate:glycerol:water solution (8:1:2, w:v:v)

shortly before observed under DMI6000 microscope system with white light (Gross-Hardt et al., 2007;

Yadegari et al., 1994).

Plant materials and growth conditions

Arabidopsis.Thaliana Wassilewskija (*Ws*) and columbia accession was acquired as wild type. Mutants

acquired in the study were all in *Ws* background. *T-DNA* insertion lines, *sks1* (FLAG_521F09), *sks2*

(FLAG_415F09), *sku5* (FLAG_386B03) were obtained from Arabidopsis Thaliana Resource Center of

Institute Jean-Pierre Bourgin (IJPB) and *sks3* (SALK_067925). Seeds were surface sterilized and plated on

1/2MS medium containing 1% agar. After 48 hours vernalization at 4°C in dark, they were plated at 22°C

under continuous light. About 7 days growth, seedlings were transferred into soil in green house. Primers

for genotyping has been show in Table S1.

RESULTS

Maternally affected seed production in *gpi-sku5/sks* mutants

Characterization of various combinations of *null* mutations of members of the *GPI-SKU5/SKS* genes revealed their severe defects in seed production, and this group of genes seemed play redundant roles in this process (Fig.S1). Among *gpi-sku5/sks* triple and quadruple mutants, those with *SKS1* mutation generated shorter mature siliques (Fig.1A) containing a large proportion of small, wrinkled and white ovules that failed to develop into seeds (Fig.1B and 1C). Interestingly, together with quadruple mutant, *sks1;sks2;sks3* triple mutant was nearly infertile due to those failures of ovule development, and bare seed was produced.

To determine the origin of such a phenotype, we analyzed the segregation of the quadruple mutant allele in the progeny of *sks1+/-;sks2;sks3;sku5* plants. A gametophytic- or zygotic-caused fertility due to *GPI-SKU5/SKS* mutations would result in rare detection of quadruple mutant seeds. By contrast, 20.69% (n=116) seedlings germinated from seeds generated by *sks1+/;sks2;sks3;sku5* plants were quadruple homozygous, which suggested the major contribution made by maternal tissue in this process.

According to our previous quantitative PCR (Zhou, 2019c) and *in silico* data (Fig.S2 and S3), unlike the specific expression of *SKS2* in reproductive organs, *SKS1*, *SKS3* and *SKU5* ubiquitously expressed at different level among the whole developmental progresses. Surprisingly, in spite of the significant contribution for reproduction, *SKS1* was expressed at relatively low level in reproductive organs.

Altogether, these data indicated that, *GPI-SKU5/SKS* genes played redundant and crucial roles in seed production, potentially through regulating ovule or seed development, and *SKS1* made significant contribution. By genetic analyses, defective maternal tissues were determined to be majorly responsible in this process.

Fig1. Dramatic decreased seed production in *gpi-sku5/sks* mutant. (A), Mature siliques generated by wild-type and *gpi-sku5/sks* mutant lines. From Left to Right: Wild-type, *sks1,sks2,sks3*; *sks1,sks2,sku5*; *sks1,sks3,sku5*; *sks2,sks3,sku5* and *sks1,sks2,sks3,sku5*; bar, 2cm. **(B)**, Dissected siliques from *wild-type* and *gpi-sku5/sks* mutants. From Top to Bottom: wild-type, *sks1,sks2, sks3*; *sks1,sks2,sku5*; *sks1,sks3,sku5*; *sks2,sks3,sku5* and *sks1,sks2,sks3,sku5*; bar, 1mm. **(C)**, Proportion of white, small, and wrinkled undeveloped ovules among all ovules/seeds in mature green siliques from triple and quadruple *gpi-sku5/sks* mutants were shown. N=15 siliques, standard errors were indicated.

Drastic decreased seed production was due to ovule abortion in *gpi-sku5/sks* mutants

To further investigate the maternal origin of the *gpi-sku5/sks* mutant phenotype, the ovules from flowers just blooming were collected and observed using the modified pseudo-Schiff propidium iodide (mPS-PI) technique (Fiume et al., 2017; Xu et al., 2016). Surprisingly, more than 90% of the *sks1;sks2;sks3;sku5* ovules failed to develop an embryo sac (Fig.2B, and 2I), and the rest showed an exposed embryo sac that was not covered by the outer integument (Fig.2C and 2I), whereas both of them displayed irregular maternal integuments. The same defective ovules were also observed, at lower frequency, in *gpi-sku5/sks* triple mutant ovules (Fig.2D-2I) and the proportion of *sku5/sks* ovules that failed to form an embryo sac matched that of aborted ovules in mature siliques in their respective mutant backgrounds (Fig.1C and Fig. 2I).

It demonstrated that, the drastic decreased seed production observed in *gpi-sku5/sks* mutants maternally resulted from increased irregular ovules that failed to develop an embryo sac and the subsequent ovule abortion, and *GPI-SKU5/SKS* genes were redundantly involved in these processes.

Fig2. Defective ovules generated by *gpi-sku5/sks* mutants with false colors. Ovules were collected from just flowering siliques from (**A**) wild-type; *(***B***) sks1,sks2,sks3,sku5* that failed to form an embryo sac; (**C**) *sks1,sks2,sks3,sku5* that generated an aberrant embryo sac; (**D**) *sks1,sks2,sks3* that failed to form an embryo sac; (**E**) *sks1,sks2,sks3* that generated an aberrant embryo sac; (**F**) *sks1,sks2,sks5*; (**G**) *sks1,sks3,sks5*; and (**H**) *sks2,sks3,sks5*. To better visualized, inner integuments are colored in yellow, and outer integuments are colored in purple; bar: 50µM. (**I**) Proportion of ovules with embryo sac formation failures generated by various combinations of *sku5/sks* mutants. N=3 siliques, and the standard error bars were indicated.

Female gametophytic abortion due to physical restriction resulted in ovule abortion in *gpi-sku5/sks* mutants

To follow the processes of their ovule abortion, the triple *null* mutants, *sks1;sks2;sku5*, which exhibited less defective ovules but be capable of generating reasonable seeds, were carefully studied. Our results showed a series of ovules and embryo sacs with various degrees of defections: slightly affected ovule of which integument could basically cover the embryo sac (Fig.3B); embryo sac were not well covered and partly exposed to external space (Fig.3C); embryo sac could barely covered by integument, but be exposed to external space as an outstretching bubble (Fig.3D); collapsing embryo sac at external space without protect by integument (Fig.3E); and ovule loss of embryo sac (Fig.3F).

It indicated that, the female gametophytic abortion, which resulted from physical restriction due to undeveloped maternal integument, should be the main reason of ovule abortion in *gpi-sku5/sks* mutants.

Fig3. Female gametophytic abortion in *gpi-sku5/sks* triple null mutant line. Ovules with mature embryo sacs were collected and observed as described in "materials and methods". **(A)**, wild-type; and a series of ovules and embryo sacs with various degrees of defections **(B-F)**; o.i: outer integuments, i.i: inner integuments, f.g: female gametophytes; bar, 50μm;

Fertilization, embryonic development and seed formation in *gpi-sku5/sks* mutants

Interestingly, those less affected ovules of *sks1;sks2;sku5* could be fertilized normally, however, due to defective integument development, inner integument could not be fully covered and normally curved by asymmetrically elongated outer integuments, which lead their embryos appeared mis-oriented (Fig.4A and 4B). Integuments develop into seed coats. Those mis-oriented embryos might develop at wrong position and break through the limit of defective seed coat and grow partially at external space (Fig.4C-4E). These defections certainly affected their seed morphogenesis largely, and exhibited irregularly shaped seed coat epidermal cells (Fig.4C-4E).

It demonstrated that, the defections occurred on integuments of *gpi-sku5/sks* mutants also largely affect their seed formation and seed morphogenesis.

Fig.4 Embryonic development and seed formation in *gpi-sku5/sks* triple null mutant line. Embryos at pre-globular stage of **(A)** Wild-type and **(B)** *sks1;sks2;sku5* stained by mPS-PI technique and observed under confocal SP8; bar:50μm; **(C)** aberrant seed and **(D)** embryo separated from this seed generated by *sks1;sks2;sku5* mutant at mature embryo stage; s.c: seed coat; em: embryo; hy: hypocotyl; co: cotyledon; bar: 100μm; dry seeds from wild-type **(E)** and *sks1;sks2;sku5* mutant **(F** and **G)** under Scanning Electronic Microscopy; bar: 100μm.

Maternal regulation of seed morphogenesis in *gpi-sku5/sks* mutant lines

The same affected seed formation and morphogenesis were also found in other *gpi-sku5/sks* mutants that generated defective integuments (Fig.5A-5F), which are irregularly shaped, and some time with tissue protrusion (Fig.5H-5J). Interestingly, the affection on seed formation and morphogenesis was so strong that all seeds generated by mutants with aberrantly developed integuments were irregularly shaped (Fig.5K).

To determine the contribution of maternal integument development regulated by *GPI-SKU5/SKS* genes in this process, the segregation of *sks1* mutant allele in the progeny of *sks1+/-;sks2;sks3;sku5* plants was characterized, which showed that, the aberrant seed formation and morphogenesis were only determined by the affected integuments, but not embryonic reason (Fig.5L).

These data indicated that, integument development regulated by *GPI-SKU5/SKS* genes were crucial during reproductive processes from gametophytic development to seed formation.

Fig5. Aberrant seed morphogenesis in *gpi-sku5/sks* mutant and seeds segregation generated by *sks1+/-,sks2,sks3,sku5* heterozygotes. Mature seeds generated by **(A)**,wild-type **(B)**, *sks1,sks2,sks3* **(C)**, *sks1,sks2,sku5*; **(D)**, *sks2,sks3,sku5* **(E)**, *sks1,sks2,sks3,sku5* **(F)**, *sks1+/-,sks2,sks3,sku5* **(G)**, *sks1+/-,sks2,sks3,sku5*, bar: 1mm; typically seed morphogenesis of **(H)**, wild-type and **(I-J)**, *sku5/sks* quadruple mutant, bar,0.2mm; **(K)**, proportion of aberrant seeds generated by all *sku5/sks* triple and quadruple mutants; **(L)**, proportion of aberrant seeds generated by triple, heterozygous and quadruple *sku5/sks* mutants. wild-type (n=375); *sks1,sks2,sks3*(n=11); *sks1,sks2,sku5* (n=234)*;* *sks1,sks3,sku5* (n=322); *sks2,sks3,sku5* (n= 226), *sks1+/-,sks2,sks3,sku5* (n-347) and *sks1,sks2,sks3,sku5*(n=10).

DISCUSSION

GPI-SKU5/SKS regulate maternal integument development in *Arabidopsis*

Integuments were totally maternal tissues and composed by special double-layer cylinder structures, inner- and outer- integument; after initiation on chalaza, the outer integument asymmetrically grew and eventually covered the inner integument and embryo sac, which determined the curvature of inner integument and shaped the ovule (Gallagher and Gasser, 2008; Gasser and Skinner, 2019; Kunieda et al., 2008; Villanueva et al., 1999). Integument development was believed regulated by a series of signaling pathways, including cell surface receptors (Jorda et al., 2016; Lee et al., 2015; Pillitteri et al., 2007; Terpstra et al., 2010), their intracellular signaling MAP Kinases (Bush and Krysan, 2007; Lopez-Bucio et al., 2014; Meng et al., 2012; Wang et al., 2008; Wang et al., 2007), and nucleus transcriptional factors (Elliott et al., 1996; Gaiser et al., 1995; Gallagher and Gasser, 2008; Gasser and Skinner, 2019; Kunieda et al., 2008; Nole-Wilson and Krizek, 2000; Villanueva et al., 1999). Although fertilization and seed production was largely affected when GPI moieties biosynthesis was interrupted in *Arabidopsis* (Bundy et al., 2016; Gillmor et al., 2005), the mechanism has not been revealed. Here, we firstly reported the involvement of particular GPI-anchored proteins in seed production and seed morphogenesis through regulating integument development.

Defective integument generated by *gpi-sku5/sks* mutants

Through our genetic and microscopy analyses, both drastic decrease of seed production resulting from ovule abortion and aberrant seed morphogenesis resulting from defective seed coat formation found in *gpi-sku5/sks* mutants were determined to be due to irregularly developed integument. In those mutants, both inner and outer integument were affected, however, defective outer integument, which was supposed to cover the inner integuments, to enclose the embryo sac and to shape the ovules, contributed more in this process. It made inner integument growing symmetrically, and embryo sac exposing to external space, which resulted in loss of female gametophytic due to physical restriction and consequent ovule abortion. Interestingly, those less affected ovules could be fertilized and develop into seeds, however, mis-localization of embryos, or insufficient seed coat that both due to defective integuments would result in post-embryonic abortion or aberrant seed morphogenesis. Here, our study clearly demonstrated the crucial roles *GPI-SKU5/SKS* genes played during integument development, and how they affected seed production and morphogenesis in *Arabidopsis*.

Disturbed cell polar expansion in *gpi-sku5/sks* mutants

In *gpi-sku5/sks* mutants, defective integuments seemed due to their disturbed cell polar expansion (Coen and Magnani, 2018; Endress, 2011; Enugutti et al., 2012; Enugutti and Schneitz, 2013; Gaiser et al., 1995; Scholz et al., 2019), where cells from both inner and outer integument seemed shorter, thicker, and aberrantly shaped. It largely disturbed their elongation and extension, where *GPI-SKU5/SKS* played redundant roles and *SKS1* made significant contribution. The same redundancy and significant contribution of *SKS1* in root cell polar expansion were reported by us recently, which exhibited similar shorter, thicker and aberrantly shaped cells (Zhou, 2019c).

But surprisingly, to our knowledge, none of gene has been reported to interfere polar expansion on both root and integument cells besides a series of MAPK components, YODA(MAPKKK)-MEK4/5(MAPKK)-MPK3/6(MAPK) cascade. And interruption of this cascade resulted in similar defective polar expansion in both roots and integuments (Bush and Krysan, 2007; Lopez-Bucio et al., 2014; Smekalova et al., 2014; Wang et al., 2008; Wang et al., 2007; Zhang et al., 2017).

The potential involvement of GPI-SKU5/SKS proteins in signaling pathway

As been reviewed recently, GPI-Anchored Proteins in *Arabidopsis* broadly participate in various biological processes, including signaling transduction through associating with cell surface receptors and/or extracellular ligands, to mediate developmental processes, stress and immune response, and so on (Zhou, 2019b). Interestingly, SKU5 was reported to directly interact with the synthesized functional Carboxyl-terminal region of Auxin Binding Protein 1 (Shimomura, 2006), that was recently believed to be a potential extracellular ligand that could bind Auxin and mediate extracellular auxin signaling through binding auxin and being recognized by cell surface reporter TMK1, then activate quick auxin response (Cao et al., 2019; Xu et al., 2014). Additionally, exogenous auxin could rapidly activate YODA(MAPKKK)-MEK4/5(MAPKK)-MPK3/6(MAPK) cascade (Mockaitis and Howell, 2000; Pillitteri et al., 2007; Smekalova et al., 2014), of which interruptions were reported to exhibit similar defective roots and integuments as *gpi-sku5/sks* mutants.

Altogether, GPI-SKU5/SKS associate with Auxin-related extracellular polypeptide/ligand ABP1, which could be recognized by cell surface receptor kinase TMK1 and activate intracellular signaling components; and their *loss-of-functions* exhibit similar defections as interrupted intracellular MAPK cascade components. It exhibits us a hypothesis that, GPI-SKU5/SKS may work as typical GPI-APs in signaling transduction during regulation of cell polar expansion in both root and integument cells.

Supplemental Materials

Fig.S1. Seed sets of all *gpi-sku5/sks* mutants. Proportion of mature seeds among all ovules/seeds in

mature green siliques from wild-type and *gpi-sku5/sks* mutants were shown. N=15 siliques, and standard

errors were indicated.

Fig.S2 Expression level of *GPI-SKU5/SKS* genes in reproductive processes of *Arabidopsis.* The expression patterns of *GPI-*

SKU5/SKS genes in reproductive organs based on microarray data from AtGenExpress Visualization Tool (AVT), were shown.

Different genes were show with different colors, different developmental stages were shown at X-axis and the relative expression

levels were shown at Y-axis.

Fig.S3. Exprssion level of *GPI-SKU5/SKS* genes during embryo development of *Arabidopsis*. Expression pattern of *GPI-SKS* genes during different embryo developmental stages based on microarray data from database http://seedgenenetwork.net/ were shown in different colors, which represented their expression levels. PEN, Peripheral Endosperm; GSC, General Seed Coat; CZE, Chalazal Endosperm; CZEC, Chalazal Seed Coat; MCE, Micropylar Endosperm; S, Suspendsor; EP, Embryo Proper;

	Primers for T-DNA insertion genotyping
SKS1-LP	GAATGGAGACTCAACACAGGC
SKS1-RP	TCGATTGAGAGATTCAATGGC
SKS2-LP	TTTGTTGCTGTCATTGCTTTG
SKS2-RP	TGAGAGTTTACCTCCACCGTG
SKS3-LP	GCAGTTTTGCTACAATCAGGC
SKS3-RP	TGGGAAAATGCTGTTTGGTAG
SKU5-LP	CTCGTGAATCCAATGCTCTTC
SKU5-RP	CAAAACGCGAGCTCTGATTAC
LB4	CGTGTGCCAGGTGCCCACGGAATAGT
RB4	TCACGGGTTGGGGTTTCTACAGGAC
LBa1	TGGTTCACGTAGTGGGCCATCG

Table.S1 Primers being utilized for genotyping T-DNA insertions in *gpi-sku5/sks* mutants were shown.

References

Borner, G.H., K.S. Lilley, T.J. Stevens, and P. Dupree. 2003. Identification of glycosylphosphatidylinositol-anchored proteins in Arabidopsis. A proteomic and genomic analysis. *Plant Physiol* 132:568-577.

Bundy, M.G., P.Z. Kosentka, A.H. Willet, L. Zhang, E. Miller, and E.D. Shpak. 2016. A Mutation in the Catalytic Subunit of the Glycosylphosphatidylinositol Transamidase Disrupts Growth, Fertility, and Stomata Formation. *Plant Physiol* 171:974-985.

Bush, S.M., and P.J. Krysan. 2007. Mutational evidence that the Arabidopsis MAP kinase MPK6 is involved in anther, inflorescence, and embryo development. *J Exp Bot* 58:2181-2191.

Cao, M., R. Chen, P. Li, Y. Yu, R. Zheng, D. Ge, W. Zheng, X. Wang, Y. Gu, Z. Gelova, J. Friml, H. Zhang, R. Liu, J. He, and T. Xu. 2019. TMK1-mediated auxin signalling regulates differential growth of the apical hook. *Nature* 568:240-243.

Coen, O., and E. Magnani. 2018. Seed coat thickness in the evolution of angiosperms. *Cell Mol Life Sci* 75:2509-2518.

Elliott, R.C., A.S. Betzner, E. Huttner, M.P. Oakes, W.Q. Tucker, D. Gerentes, P. Perez, and
D.R. Smyth. 1996. AINTEGUMENTA, an APETALA2-like gene of Arabidopsis with
pleiotropic roles in ovule development and floral organ growth. *Plant Cell* 8:155-
168.

Endress, P.K. 2011. Angiosperm ovules: diversity, development, evolution. *Ann Bot*
107:1465-1489.

Enugutti, B., C. Kirchhelle, M. Oelschner, R.A. Torres Ruiz, I. Schliebner, D. Leister, and K.
Schneitz. 2012. Regulation of planar growth by the Arabidopsis AGC protein
kinase UNICORN. *Proc Natl Acad Sci U S A* 109:15060-15065.

Enugutti, B., and K. Schneitz. 2013. Genetic analysis of ectopic growth suppression
during planar growth of integuments mediated by the Arabidopsis AGC protein
kinase UNICORN. *BMC Plant Biol* 13:2.

Fiume, E., O. Coen, W. Xu, L. Lepiniec, and E. Magnani. 2017. Developmental patterning
of sub-epidermal cells in the outer integument of Arabidopsis seeds. *PLoS One*
12:e0188148.

Gaiser, J.C., K. Robinson-Beers, and C.S. Gasser. 1995. The Arabidopsis SUPERMAN Gene
Mediates Asymmetric Growth of the Outer Integument of Ovules. *Plant Cell*
7:333-345.

Gallagher, T.L., and C.S. Gasser. 2008. Independence and interaction of regions of the
INNER NO OUTER protein in growth control during ovule development. *Plant*
Physiol 147:306-315.

Gasser, C.S., and D.J. Skinner. 2019. Development and evolution of the unique ovules of
flowering plants. *Curr Top Dev Biol* 131:373-399.

Gillmor, C.S., W. Lukowitz, G. Brininstool, J.C. Sedbrook, T. Hamann, P. Poindexter, and C.
Somerville. 2005. Glycosylphosphatidylinositol-anchored proteins are required
for cell wall synthesis and morphogenesis in Arabidopsis. *Plant Cell* 17:1128-
1140.

Gross-Hardt, R., C. Kagi, N. Baumann, J.M. Moore, R. Baskar, W.B. Gagliano, G. Jurgens,
and U. Grossniklaus. 2007. LACHESIS restricts gametic cell fate in the female
gametophyte of Arabidopsis. *PLoS Biol* 5:e47.

Jorda, L., S. Sopena-Torres, V. Escudero, B. Nunez-Corcuera, M. Delgado-Cerezo, K.U.
Torii, and A. Molina. 2016. ERECTA and BAK1 Receptor Like Kinases Interact to
Regulate Immune Responses in Arabidopsis. *Front Plant Sci* 7:897.

Kawagoe, K., D. Kitamura, M. Okabe, I. Taniuchi, M. Ikawa, T. Watanabe, T. Kinoshita, and
J. Takeda. 1996. Glycosylphosphatidylinositol-anchor-deficient mice: implications
for clonal dominance of mutant cells in paroxysmal nocturnal hemoglobinuria.
Blood 87:3600-3606.

Kinoshita, T. 2014. Biosynthesis and deficiencies of glycosylphosphatidylinositol. *Proc Jpn*
Acad Ser B Phys Biol Sci 90:130-143.

Kinoshita, T., and M. Fujita. 2016. Biosynthesis of GPI-anchored proteins: special
emphasis on GPI lipid remodeling. *J Lipid Res* 57:6-24.

Kunieda, T., N. Mitsuda, M. Ohme-Takagi, S. Takeda, M. Aida, M. Tasaka, M. Kondo, M.
Nishimura, and I. Hara-Nishimura. 2008. NAC family proteins NARS1/NAC2 and
NARS2/NAM in the outer integument regulate embryogenesis in Arabidopsis.

Plant Cell 20:2631-2642.

Lee, J.S., M. Hnilova, M. Maes, Y.C. Lin, A. Putarjunan, S.K. Han, J. Avila, and K.U. Torii. 2015. Competitive binding of antagonistic peptides fine-tunes stomatal patterning. *Nature* 522:439-443.

Lopez-Bucio, J.S., J.G. Dubrovsky, J. Raya-Gonzalez, Y. Ugartechea-Chirino, J. Lopez-Bucio, L.A. de Luna-Valdez, M. Ramos-Vega, P. Leon, and A.A. Guevara-Garcia. 2014. Arabidopsis thaliana mitogen-activated protein kinase 6 is involved in seed formation and modulation of primary and lateral root development. *J Exp Bot* 65:169-183.

Meng, X., H. Wang, Y. He, Y. Liu, J.C. Walker, K.U. Torii, and S. Zhang. 2012. A MAPK cascade downstream of ERECTA receptor-like protein kinase regulates Arabidopsis inflorescence architecture by promoting localized cell proliferation. *Plant Cell* 24:4948-4960.

Mockaitis, K., and S.H. Howell. 2000. Auxin induces mitogenic activated protein kinase (MAPK) activation in roots of Arabidopsis seedlings. *Plant J* 24:785-796.

Nole-Wilson, S., and B.A. Krizek. 2000. DNA binding properties of the Arabidopsis floral development protein AINTEGUMENTA. *Nucleic Acids Res* 28:4076-4082.

Oxley, D., and A. Bacic. 1999. Structure of the glycosylphosphatidylinositol anchor of an arabinogalactan protein from Pyrus communis suspension-cultured cells. *Proc Natl Acad Sci U S A* 96:14246-14251.

Pillitteri, L.J., S.M. Bemis, E.D. Shpak, and K.U. Torii. 2007. Haploinsufficiency after successive loss of signaling reveals a role for ERECTA-family genes in Arabidopsis ovule development. *Development* 134:3099-3109.

Rutherford, R., and P.H. Masson. 1996. Arabidopsis thaliana sku mutant seedlings show exaggerated surface-dependent alteration in root growth vector. *Plant Physiol* 111:987-998.

Scholz, S., J. Plessmann, B. Enugutti, R. Huttl, K. Wassmer, and K. Schneitz. 2019. The AGC protein kinase UNICORN controls planar growth by attenuating PDK1 in Arabidopsis thaliana. *PLoS Genet* 15:e1007927.

Sedbrook, J.C., K.L. Carroll, K.F. Hung, P.H. Masson, and C.R. Somerville. 2002. The Arabidopsis SKU5 gene encodes an extracellular glycosyl phosphatidylinositol-anchored glycoprotein involved in directional root growth. *Plant Cell* 14:1635-1648.

Shimomura, S. 2006. Identification of a glycosylphosphatidylinositol-anchored plasma membrane protein interacting with the C-terminus of auxin-binding protein 1: a photoaffinity crosslinking study. *Plant Mol Biol* 60:663-677.

Smekalova, V., I. Luptovciak, G. Komis, O. Samajova, M. Ovecka, A. Doskocilova, T. Takac, P. Vadovic, O. Novak, T. Pechan, A. Ziemann, P. Kosutova, and J. Samaj. 2014. Involvement of YODA and mitogen activated protein kinase 6 in Arabidopsis post-embryogenic root development through auxin up-regulation and cell division plane orientation. *New Phytol* 203:1175-1193.

Stevens, V.L. 1995. Biosynthesis of glycosylphosphatidylinositol membrane anchors. *Biochem J* 310 (Pt 2):361-370.

Takahashi, D., Y. Kawamura, and M. Uemura. 2016. Cold acclimation is accompanied by

445   complex responses of glycosylphosphatidylinositol (GPI)-anchored proteins in
446   Arabidopsis. *J Exp Bot* 67:5203-5215.

447 Terpstra, I.R., L.B. Snoek, J.J. Keurentjes, A.J. Peeters, and G. van den Ackerveken. 2010.
448   Regulatory network identification by genetical genomics: signaling downstream
449   of the Arabidopsis receptor-like kinase ERECTA. *Plant Physiol* 154:1067-1078.

450 Villanueva, J.M., J. Broadhvest, B.A. Hauser, R.J. Meister, K. Schneitz, and C.S. Gasser.
451   1999. INNER NO OUTER regulates abaxial- adaxial patterning in Arabidopsis
452   ovules. *Genes Dev* 13:3160-3169.

453 Wang, H., Y. Liu, K. Bruffett, J. Lee, G. Hause, J.C. Walker, and S. Zhang. 2008. Haplo-
454   insufficiency of MPK3 in MPK6 mutant background uncovers a novel function of
455   these two MAPKs in Arabidopsis ovule development. *Plant Cell* 20:602-613.

456 Wang, H., N. Ngwenyama, Y. Liu, J.C. Walker, and S. Zhang. 2007. Stomatal development
457   and patterning are regulated by environmentally responsive mitogen-activated
458   protein kinases in Arabidopsis. *Plant Cell* 19:63-73.

459 Xu, T., N. Dai, J. Chen, S. Nagawa, M. Cao, H. Li, Z. Zhou, X. Chen, R. De Rycke, H.
460   Rakusova, W. Wang, A.M. Jones, J. Friml, S.E. Patterson, A.B. Bleecker, and Z.
461   Yang. 2014. Cell surface ABP1-TMK auxin-sensing complex activates ROP GTPase
462   signaling. *Science* 343:1025-1028.

463 Xu, W., E. Fiume, O. Coen, C. Pechoux, L. Lepiniec, and E. Magnani. 2016. Endosperm and
464   Nucellus Develop Antagonistically in Arabidopsis Seeds. *Plant Cell* 28:1343-1360.

465 Yadegari, R., G. Paiva, T. Laux, A.M. Koltunow, N. Apuya, J.L. Zimmerman, R.L. Fischer, J.J.
466   Harada, and R.B. Goldberg. 1994. Cell Differentiation and Morphogenesis Are
467   Uncoupled in Arabidopsis raspberry Embryos. *Plant Cell* 6:1713-1729.

468 Zhang, M., H. Wu, J. Su, H. Wang, Q. Zhu, Y. Liu, J. Xu, W. Lukowitz, and S. Zhang. 2017.
469   Maternal control of embryogenesis by MPK6 and its upstream MKK4/MKK5 in
470   Arabidopsis. *Plant J* 92:1005-1019.

471 Zhou, K. 2019a. The alternative splicing of SKU5-Similar3 in Arabidopsis. *Plant Signal*
472   *Behav* 1-4.

473 Zhou, K. 2019b. Glycosylphosphatidylinositol-Anchored Proteins in Arabidopsis and One
474   of Their Common Roles in Signaling Transduction. *Frontiers in Plant Science* 10:

475 Zhou, K. 2019c. GPI-anchored SKS proteins regulate root development through
476   controlling cell polar expansion and cell wall synthesis. *Biochem Biophys Res*
477   *Commun* 509:119-124.

Parsed Citations

Borner, G.H., K.S. Lilley, T.J. Stevens, and P. Dupree. 2003. Identification of glycosylphosphatidylinositol-anchored proteins in Arabidopsis. A proteomic and genomic analysis. Plant Physiol 132:568-577.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Bundy, M.G., P.Z. Kosentka, A.H. Willet, L. Zhang, E. Miller, and E.D. Shpak. 2016. A Mutation in the Catalytic Subunit of the Glycosylphosphatidylinositol Transamidase Disrupts Growth, Fertility, and Stomata Formation. Plant Physiol 171:974-985.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Bush, S.M., and P.J. Krysan. 2007. Mutational evidence that the Arabidopsis MAP kinase MPK6 is involved in anther, inflorescence, and embryo development. J Exp Bot 58:2181-2191.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Cao, M., R. Chen, P. Li, Y. Yu, R. Zheng, D. Ge, W. Zheng, X. Wang, Y. Gu, Z. Gelova, J. Friml, H. Zhang, R. Liu, J. He, and T. Xu. 2019. TMK1-mediated auxin signalling regulates differential growth of the apical hook. Nature 568:240-243.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Coen, O., and E. Magnani. 2018. Seed coat thickness in the evolution of angiosperms. Cell Mol Life Sci 75:2509-2518.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Elliott, R.C., A.S. Betzner, E. Huttner, M.P. Oakes, W.Q. Tucker, D. Gerentes, P. Perez, and D.R. Smyth. 1996. AINTEGUMENTA, an APETALA2-like gene of Arabidopsis with pleiotropic roles in ovule development and floral organ growth. Plant Cell 8:155-168.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Endress, P.K. 2011. Angiosperm ovules: diversity, development, evolution. Ann Bot 107:1465-1489.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Enugutti, B., C. Kirchhelle, M. Oelschner, R.A. Torres Ruiz, I. Schliebner, D. Leister, and K. Schneitz. 2012. Regulation of planar growth by the Arabidopsis AGC protein kinase UNICORN. Proc Natl Acad Sci U S A 109:15060-15065.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Enugutti, B., and K. Schneitz. 2013. Genetic analysis of ectopic growth suppression during planar growth of integuments mediated by the Arabidopsis AGC protein kinase UNICORN. BMC Plant Biol 13:2.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Fiume, E., O. Coen, W. Xu, L. Lepiniec, and E. Magnani. 2017. Developmental patterning of sub-epidermal cells in the outer integument of Arabidopsis seeds. PLoS One 12:e0188148.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Gaiser, J.C., K. Robinson-Beers, and C.S. Gasser. 1995. The Arabidopsis SUPERMAN Gene Mediates Asymmetric Growth of the Outer Integument of Ovules. Plant Cell 7:333-345.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Gallagher, T.L., and C.S. Gasser. 2008. Independence and interaction of regions of the INNER NO OUTER protein in growth control during ovule development. Plant Physiol 147:306-315.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Gasser, C.S., and D.J. Skinner. 2019. Development and evolution of the unique ovules of flowering plants. Curr Top Dev Biol 131:373-399.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Gillmor, C.S., W. Lukowitz, G. Brininstool, J.C. Sedbrook, T. Hamann, P. Poindexter, and C. Somerville. 2005. Glycosylphosphatidylinositol-anchored proteins are required for cell wall synthesis and morphogenesis in Arabidopsis. Plant Cell 17:1128-1140.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Gross-Hardt, R., C. Kagi, N. Baumann, J.M. Moore, R. Baskar, W.B. Gagliano, G. Jurgens, and U. Grossniklaus. 2007. LACHESIS restricts gametic cell fate in the female gametophyte of Arabidopsis. PLoS Biol 5:e47.
Pubmed: Author and Title

Google Scholar: Author Only Title Only Author and Title

Jorda, L., S. Sopena-Torres, V. Escudero, B. Nunez-Corcuera, M. Delgado-Cerezo, K.U. Torii, and A. Molina. 2016. ERECTA and BAK1 Receptor Like Kinases Interact to Regulate Immune Responses in Arabidopsis. Front Plant Sci 7:897.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Kawagoe, K., D. Kitamura, M. Okabe, I. Taniuchi, M. Ikawa, T. Watanabe, T. Kinoshita, and J. Takeda. 1996. Glycosylphosphatidylinositol-anchor-deficient mice: implications for clonal dominance of mutant cells in paroxysmal nocturnal hemoglobinuria. Blood 87:3600-3606.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Kinoshita, T. 2014. Biosynthesis and deficiencies of glycosylphosphatidylinositol. Proc Jpn Acad Ser B Phys Biol Sci 90:130-143.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Kinoshita, T., and M. Fujita. 2016. Biosynthesis of GPI-anchored proteins: special emphasis on GPI lipid remodeling. J Lipid Res 57:6-24.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Kunieda, T., N. Mitsuda, M. Ohme-Takagi, S. Takeda, M. Aida, M. Tasaka, M. Kondo, M. Nishimura, and I. Hara-Nishimura. 2008. NAC family proteins NARS1/NAC2 and NARS2/NAM in the outer integument regulate embryogenesis in Arabidopsis. Plant Cell 20:2631-2642.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Lee, J.S., M. Hnilova, M. Maes, Y.C. Lin, A. Putarjunan, S.K. Han, J. Avila, and K.U. Torii. 2015. Competitive binding of antagonistic peptides fine-tunes stomatal patterning. Nature 522:439-443.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Lopez-Bucio, J.S., J.G. Dubrovsky, J. Raya-Gonzalez, Y. Ugartechea-Chirino, J. Lopez-Bucio, L.A. de Luna-Valdez, M. Ramos-Vega, P. Leon, and A.A. Guevara-Garcia. 2014. Arabidopsis thaliana mitogen-activated protein kinase 6 is involved in seed formation and modulation of primary and lateral root development. J Exp Bot 65:169-183.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Meng, X., H. Wang, Y. He, Y. Liu, J.C. Walker, K.U. Torii, and S. Zhang. 2012. A MAPK cascade downstream of ERECTA receptor-like protein kinase regulates Arabidopsis inflorescence architecture by promoting localized cell proliferation. Plant Cell 24:4948-4960.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Mockaitis, K., and S.H. Howell. 2000. Auxin induces mitogenic activated protein kinase (MAPK) activation in roots of Arabidopsis seedlings. Plant J 24:785-796.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Nole-Wilson, S., and B.A. Krizek. 2000. DNA binding properties of the Arabidopsis floral development protein AINTEGUMENTA. Nucleic Acids Res 28:4076-4082.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Oxley, D., and A. Bacic. 1999. Structure of the glycosylphosphatidylinositol anchor of an arabinogalactan protein from Pyrus communis suspension-cultured cells. Proc Natl Acad Sci U S A 96:14246-14251.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Pillitteri, L.J., S.M. Bemis, E.D. Shpak, and K.U. Torii. 2007. Haploinsufficiency after successive loss of signaling reveals a role for ERECTA-family genes in Arabidopsis ovule development. Development 134:3099-3109.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Rutherford, R., and P.H. Masson. 1996. Arabidopsis thaliana sku mutant seedlings show exaggerated surface-dependent alteration in root growth vector. Plant Physiol 111:987-998.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Scholz, S., J. Plessmann, B. Enugutti, R. Huttl, K. Wassmer, and K. Schneitz. 2019. The AGC protein kinase UNICORN controls planar growth by attenuating PDK1 in Arabidopsis thaliana. PLoS Genet 15:e1007927.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Sedbrook, J.C., K.L. Carroll, K.F. Hung, P.H. Masson, and C.R. Somerville. 2002. The Arabidopsis SKU5 gene encodes an extracellular glycosyl phosphatidylinositol-anchored glycoprotein involved in directional root growth. Plant Cell 14:1635-1648.

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Shimomura, S. 2006. Identification of a glycosylphosphatidylinositol-anchored plasma membrane protein interacting with the C-terminus of auxin-binding protein 1: a photoaffinity crosslinking study. Plant Mol Biol 60:663-677.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Smekalova, V., I. Luptovciak, G. Komis, O. Samajova, M. Ovecka, A. Doskocilova, T. Takac, P. Vadovic, O. Novak, T. Pechan, A. Ziemann, P. Kosutova, and J. Samaj. 2014. Involvement of YODA and mitogen activated protein kinase 6 in Arabidopsis post-embryogenic root development through auxin up-regulation and cell division plane orientation. New Phytol 203:1175-1193.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Stevens, V.L. 1995. Biosynthesis of glycosylphosphatidylinositol membrane anchors. Biochem J 310 (Pt 2):361-370.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Takahashi, D., Y. Kawamura, and M. Uemura. 2016. Cold acclimation is accompanied by complex responses of glycosylphosphatidylinositol (GPI)-anchored proteins in Arabidopsis. J Exp Bot 67:5203-5215.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Terpstra, I.R., L.B. Snoek, J.J. Keurentjes, A.J. Peeters, and G. van den Ackerveken. 2010. Regulatory network identification by genetical genomics: signaling downstream of the Arabidopsis receptor-like kinase ERECTA. Plant Physiol 154:1067-1078.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Villanueva, J.M., J. Broadhvest, B.A. Hauser, R.J. Meister, K. Schneitz, and C.S. Gasser. 1999. INNER NO OUTER regulates abaxial-adaxial patterning in Arabidopsis ovules. Genes Dev 13:3160-3169.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Wang, H., Y. Liu, K. Bruffett, J. Lee, G. Hause, J.C. Walker, and S. Zhang. 2008. Haplo-insufficiency of MPK3 in MPK6 mutant background uncovers a novel function of these two MAPKs in Arabidopsis ovule development. Plant Cell 20:602-613.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Wang, H., N. Ngwenyama, Y. Liu, J.C. Walker, and S. Zhang. 2007. Stomatal development and patterning are regulated by environmentally responsive mitogen-activated protein kinases in Arabidopsis. Plant Cell 19:63-73.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Xu, T., N. Dai, J. Chen, S. Nagawa, M. Cao, H. Li, Z. Zhou, X. Chen, R. De Rycke, H. Rakusova, W. Wang, A.M. Jones, J. Friml, S.E. Patterson, A.B. Bleecker, and Z. Yang. 2014. Cell surface ABP1-TMK auxin-sensing complex activates ROP GTPase signaling. Science 343:1025-1028.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Xu, W., E. Fiume, O. Coen, C. Pechoux, L. Lepiniec, and E. Magnani. 2016. Endosperm and Nucellus Develop Antagonistically in Arabidopsis Seeds. Plant Cell 28:1343-1360.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Yadegari, R., G. Paiva, T. Laux, A.M. Koltunow, N. Apuya, J.L. Zimmerman, R.L. Fischer, J.J. Harada, and R.B. Goldberg. 1994. Cell Differentiation and Morphogenesis Are Uncoupled in Arabidopsis raspberry Embryos. Plant Cell 6:1713-1729.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Zhang, M., H. Wu, J. Su, H. Wang, Q. Zhu, Y. Liu, J. Xu, W. Lukowitz, and S. Zhang. 2017. Maternal control of embryogenesis by MPK6 and its upstream MKK4/MKK5 in Arabidopsis. Plant J 92:1005-1019.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Zhou, K. 2019a. The alternative splicing of SKU5-Similar3 in Arabidopsis. Plant Signal Behav 1-4.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Zhou, K. 2019b. Glycosylphosphatidylinositol-Anchored Proteins in Arabidopsis and One of Their Common Roles in Signaling Transduction. Frontiers in Plant Science 10:
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Zhou, K. 2019c. GPI-anchored SKS proteins regulate root development through controlling cell polar expansion and cell wall synthesis. Biochem Biophys Res Commun 509:119-124.

Pubmed: [Author and Title](#)
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)