TITLE

Developing a transcriptomic framework for testing testosterone-mediated handicap hypotheses

AUTHORS

Daniel J. Newhouse[a,d,e], Ben J. Vernasco[b,c]

[a]Department of Biology, East Carolina University, Greenville, NC, USA

[b]Department of Biological Sciences, Virginia Tech, Blacksburg, VA, USA

[c]Current Address: School of Biological Sciences, Washington State University, Pullman, WA, USA

[d]Current Address: Department of Immunology, University of Washington, Seattle, WA, USA

[e]Corresponding author: newhoused12@students.ecu.edu

KEYWORDS RNA-seq; testosterone; handicap hypothesis; immunocompetence; oxidative stress; co-expression network

ABSTRACT

Sexually selected traits are hypothesized to be honest signals of individual quality due to the costs associated with their maintenance, development, and/or production. Testosterone, a sex steroid associated with the development and/or production of sexually selected traits, has been proposed to enforce the honesty of sexually selected traits via its immunosuppressive effects (i.e., the Immunocompetence Handicap Hypothesis) and/or by influencing an individual's exposure/susceptibility to oxidative stress (i.e., the Oxidation Handicap Hypothesis). Previous work testing these hypotheses has primarily focused on physiological measurements of immunity or oxidative stress, but little is known about the molecular pathways by which testosterone could influence immunity and/or oxidative stress pathways. To further understand the transcriptomic

consequences of experimentally elevated testosterone in the context of handicap hypotheses, we

used previously published RNA-seq data from studies that measured the transcriptome of

individuals treated with either a testosterone-filled or an empty (i.e., control) implant. Two

studies encompassing three species of bird and three tissue types fit our selection criteria and we

reanalyzed the data using weighted gene co-expression network analysis. Our results show that

testosterone-treated individuals exhibited signatures of immunosuppression and we provide some

evidence to suggest that the transcriptomic signature of immunosuppression is evolutionarily

conserved between the three species. While our results provide no evidence to suggest

testosterone mediates handicaps via pathways associated with oxidative stress, they do support

the hypothesis that testosterone enforces the honesty of sexually-selected traits by influencing an

individual's immunocompetence. Overall, this study develops a framework for testing

testosterone-mediated handicap hypotheses and provides guidelines for future integrative and

comparative studies focused on the proximate mechanisms mediating sexually selected traits.

1. INTRODUCTION

There is a long-standing interest in understanding why sexually selected traits have evolved and

one hypothesis suggests that mates have selected for traits that are costly to develop or bear (i.e.,

the handicap hypothesis; Zahavi, 1975). Under this framework, an individual's investment in

sexually selected traits is thought to correlate with their investment in traits that influence their

survival (Grafen, 1990; Andersson, 1994). Individuals face tradeoffs when fitness-related traits

exhibit negative correlations and, because of these tradeoffs, individuals can incur survival costs

from their reproductive investments (Stearns, 1992). These costs are thought to arise because the

development and/or expression of traits important for reproduction (e.g., sexually selected traits)

and traits important for survival (e.g., immune function) are dependent on the same mechanism

(Zera and Harshman, 2001). Therefore, our understanding of the evolution of sexually selected

traits is dependent upon our understanding of the mechanisms that underlie their production and

the pleiotropic effects of such mechanisms (Kokko et al., 2003).

Testosterone is a sex steroid that is known for its influence on a diverse suite of

behavioral, morphological, and physiological traits associated with reproduction (e.g.,

metabolism and courtship behaviors; Ketterson et al. 2009), including the development and/or

expression of sexually selected traits (Hau, 2007; Fusani, 2008; Ball and Balthazart, 2009). In

combination with its effects on traits associated with survival (e.g., immune function, Segner et

al., 2017), testosterone is thought to enforce the honesty of sexually selected traits (Ketterson and

Nolan, Jr., 1999, Buchanan et al., 2001, Wingfield et al., 2001, Reed et al., 2006). Two

prominent hypotheses have been proposed to explain how testosterone maintains the honesty of

sexually selected traits: the Immunocompetence Handicap Hypothesis (Folstad and Karter, 1992)

and the Oxidation Handicap Hypothesis (Alonso-Alvarez et al., 2007). The Immunocompetence

Handicap Hypothesis proposes that sexually selected traits remain honest because of

testosterone's antagonistic effects on an individual's immune function. Therefore, poor quality or

low condition individuals cannot maintain high levels of circulating testosterone due to

testosterone's immunosuppressive effects (Folstad and Karter, 1992). An initial meta-analysis

revealed weak support for this hypothesis (Roberts et al., 2004), but a subsequent meta-analysis

found that experimentally increasing testosterone results in suppression of both cell-mediated

and humoral immunity (Foo et al., 2017). Foo et al. (2017) also found positive trends between

multiple measures of immune function and naturally occurring levels of circulating testosterone.

These results fit the predictions of the Immunocompetence Handicap Hypothesis because

individuals naturally expressing high of testosterone represent high quality or high condition

individuals that can invest in sexually selected traits without compromising their immune system

(Peters, 2000). The Oxidation Handicap Hypothesis, on the other hand, states that sexually

selected traits remain honest because testosterone increases an individual's susceptibility and/or

exposure to oxidative stress (Alonso-Alvarez et al., 2007). In other words, testosterone may

influence an individual's ability to protect or repair cellular machinery from oxidative damage

(e.g., an individual's antioxidant defenses) or testosterone may influence the rate that reactive

oxygen species are produced (Alonso-Alvarez et al., 2007). Importantly, either one of these

consequences may occur independent of the other. Of the few studies that have directly tested the

Oxidation Handicap Hypothesis, some have found support (Mougeot et al., 2009; Hoogenboom

et al., 2012) while other studies did not find support for this hypothesis (Isaksson et al., 2011;

Casagrande et al., 2012; Taff and Freeman-Gallant, 2014; Baldo et al., 2015). Nonetheless, both

hypotheses have primarily been tested using physiological measurements of oxidative stress and

immunity, but less is known about the underlying molecular pathways. Given that sex steroids

partly function by binding to intracellular receptors and acting as transcription factors (Ketterson

and Nolan, Jr., 1999; Nelson, 2011), measuring the relationship between testosterone and

transcription can shed light on the proximate pathways that testosterone influences.

Modern sequencing approaches, like RNA sequencing (RNA-seq), allow for

comprehensive measurements of whole transcriptomes and the relative abundance of each

transcript (Wang et al., 2009). This approach assesses coordinated, large-scale transcriptional

responses rather than focusing on targeted candidate genes (e.g., via qPCR). RNA-seq

approaches have been used to investigate the role of androgens on gene expression, particularly

in the context of sex differences (Gao et al., 2015; Cox et al., 2017) and gonadal development

(Monson et al., 2017; Zheng et al., 2019). Similarly, RNA-seq based studies have been crucial in

providing a more comprehensive understanding of the complex and dynamic immune and stress

responses (e.g., Barshis et al., 2013; Huang et al., 2013; Kim et al., 2018). In the context of mate

choice, measuring the relationship between testosterone and transcription can shed light on the

pathways that testosterone influences to potentially enforce the honesty of sexually selected traits

(e.g., immune or oxidative stress pathways). Such studies can ultimately inform our

understanding of the effects of circulating testosterone on the expression of genes related to

immune function and/or oxidative stress. To date, these hypotheses have rarely been tested using

genome scale approaches. However, in red grouse (*Lagopus lagopus scoticus*), testosterone

treatment had little effect on overall gene expression in the liver and spleen but did result in the

down-regulation of genes related to immune function in caecal tissue (Wenzel et al., 2013).

Given that Wenzel et al. (2013) used a microarray-based approach, testing these handicap

hypotheses using RNA-seq represents a more modern, robust test as RNA-seq provides many

advantages over microarray technologies, including higher sensitivity and no hybridization

biases (Wang et al., 2009).

Here, we use published RNA-seq datasets to further examine the effects of testosterone

on the transcriptome. We re-analyze studies that measured the transcriptome of testosterone-

treatment and control subjects in three species of bird: zebra finch (*Taenopygia guttata*), golden-

collared manakin (*Manacus vitellinus*), and Japanese quail (*Coturnix japonica*). In the zebra

finch, male vocal behavior as well as bill color (a sexually-selected trait) have been found to be

sensitive to testosterone (Cynx et al., 2005, McGraw et al. 2006). Similarly, male golden-

collared manakin engage in elaborate courtship behaviors during the breeding season and

experimentally blocking androgen receptors decreases the expression of male courtship

behaviors (Day et al., 2007; Schlinger et al., 2013). Male Japanese quail also produce brightly

colored cheek feathers to attract females (Hiyama et al., 2018). Castrating males influences the

color of a male's cheek feathers and administering testosterone to castrated males causes cheek

patches of castrated males to match those of males that have not been castrated. In this study, we

focus on transcriptomic data from tissues that are known to be express androgen receptors: the

foam gland of quail and muscular tissues from the zebra finch and golden-collared manakin,

(Adkins-Regan, 1999; Fuxjager et al., 2016). We constructed co-expression networks to identify

gene networks that show correlated expression patterns following testosterone treatment. If

testosterone influences the honesty of sexually selected traits via immune mechanisms, relative

to the control group, the group treated with testosterone treatment will exhibit lower expression

of genes with annotated immune function (i.e. immunosuppression). Alternatively, if testosterone

acts via mechanisms associated with an individual's susceptibility or exposure to oxidative

stress, then we predict that testosterone treatment will cause a decrease in the expression of genes

with annotated functions in antioxidant protection and/or an increase in genes that are expressed

in response to oxidative stress. An important caveat is that no support for either hypothesis does

not exclude the possibility that these pathways maintain the honestly of sexually selected traits

independent of testosterone's effects, as has been suggested before (Metcalfe and Alonso-

Alvarez 2010, Weaver et al. 2017). Lastly, given the limited taxonomic diversity of our dataset,

we acknowledge that future studies including more diverse taxon and tissues are ultimately

needed to robustly examine if testosterone mediates the honesty of sexually selected traits via

transcriptomic pathways.

2. METHODS

2.1 Study Selection

To identify studies of interest, we first performed a literature search on both Scopus and Google

Scholar with the following search terms: "testosterone" AND "RNA-seq" or "transcriptome" or

"transcriptomics". This literature search produced 260 results. From this list of 260 studies, we

retained RNA-seq studies that measured gene expression in adult males from both testosterone-

manipulated and control groups. This process resulted in one study for re-analysis and we also

identified an additional dataset by searching within NCBI's Sequence Read Archive

(Supplemental Figure 1). Finseth and Harrison (2018) experimentally increased testosterone in

Japanese quail (*Coturnix japonica*, "quail") experiencing short days and performed RNA-seq on

the foam gland (n=6 each testosterone and control). Fuxjager et al. (2016) experimentally

increased testosterone in male golden-collared manakins (*Manacus vitellinus*, "manakin") and

performed RNA-seq on pectoralis and scapulohumeralis caudalis tissue (n = 3 individuals per

treatment group and 2 tissue types per individual). In an effort to understand the role of androgen

receptors in mediating the gene expression patterns observed in the manakin samples, Fuxjager

et al. (2016) also administered 2 different combinations of hormone implants to adult zebra

finches (*Taeniopygia guttata*): one group received testosterone-filled implants and an empty

implant while the second group was administered a testosterone implant and an implant filled

with flutamide, a drug that blocks the androgen receptor. Fuxjager et al. (2016) then measured

the transcriptome of the pectoralis and scapulohumeralis caudalis muscles (n = 3 individuals per

treatment group and 2 tissues types per individual) using RNA-seq. Thus, no unmanipulated,

control zebra finch samples were included in the zebra finch RNA-seq data generated by

Fuxjager et al. (2016). Because the current study is not focused on understanding the role of the

androgen receptor in mediating handicap hypotheses and due to the limited sample size available

for testing the role of androgen receptor in mediating handicap hypotheses, only samples from

the zebra finches treated with just testosterone were included in our analysis. While the zebra

finch samples lack an appropriate control (i.e., individuals not treated with testosterone), the

zebra finch samples can nonetheless be used in the shared orthologous gene expression analysis

(see below for more details).

2.2 Data Re-analysis

We downloaded the raw sequencing data from SRA with sratoolkit fastq-dump (quail:

PRJNA397592; manakin & zebra finch: PRJNA297576) and adaptor trimmed all reads with

Trim Galore! v0.3.8 (https://github.com/FelixKrueger/TrimGalore). We aligned trimmed reads to

the respective reference genome (*M. vitellinus* v2, *C. japonica* v2, *T. guttata* bTaeGut1_v1.p) for

each species with STAR v2.5.3 (Dobin et al., 2013) and quantified expression with htseq-count

v0.6.0 (Anders et al., 2015), specifying strand 'no'. We normalized counts to sequencing depth

and variance stabilizing transformed counts with DEseq2 (Love et al., 2014). Transformed

counts were visualized with a principal component analysis (PCA) using pcaExplorer v2.8.1

(Marini & Binder 2019) and PCAtools v2.0.0 (Blighe & Lun 2020).

To test for the effect of testosterone treatment on transcription, we used the weighted

gene co-expression network analysis (WGCNA) tool (Langfelder et al., 2011; Langfelder and

Horvath, 2008). We created modules independently for quail and manakin with the following

shared parameters: network type=signed, minimum module size=30, and module

dissimilarity=0.2. We used $\beta=12$ for quail and $\beta=18$ for manakin, which represents the point the

network reached scale free topology. We then tested for correlations between modules and

testosterone treatment using a $p < 0.05$ cutoff. We identified the hub genes of each module by

selecting the top five genes with the highest module membership (MM) score. Since we created a

signed network in WGCNA and correlated modules to a binary trait of Testosterone or Control,

each module correlated with testosterone treatment shows opposite expression patterns between

the two treatment groups. In other words, genes showing low expression in testosterone treated

animals are relatively highly expressed among control samples.

As each of these studies was relatively underpowered for gene expression analyses

(reviewed in Schurch et al. 2016), we used a subsampling approach to test the robustness of each

WGCNA analysis. In other words, we were interested if the results obtained in the full analysis

were reproducible using subsets of the data. Gene signatures observed in both the full analysis

and subset analysis would thus help confidently identify meaningful pathways responding to

testosterone treatment. While both studies contain the same number of RNA-seq samples, the

manakin dataset represents 6 total individuals and the quail dataset represents 12 individuals. We

divided the quail data into two sets of six individuals (n=3 control, n=3 testosterone). For

manakin, we divided the data in two by muscle type (n=3 control, n=3 testosterone each). We

then re-ran WGCNA on each of the four subsets with the same parameters, except specifying

β=16 for both quail subsets and β=18 for both manakin subsets. We then performed functional

enrichment analyses on each module as described below.

We were also interested if there was a shared response to testosterone among all species.

Therefore, using all 30 RNA-seq samples (n=12 quail, n=12 manakin, n=6 zebra finch), we

identified 8,454 one-to-one (i.e. single copy, non-duplicated) orthologs among all samples using

Ensembl BioMart (Kinsella et al. 2011). This filtering process restricts the analyses to genes that

are evolutionarily conserved between species. We normalized and variance stabilized

transformed all samples using DESeq2. Since the data come from two independent studies, we

then used the removeBatchEffect function from limma (Ritchie et al., 2015) to control for batch

effects. Lastly, we used these conserved, expressed orthologs to create a single co-expression

network with WGCNA for all samples. Module reconstruction parameters were identical to the

species-specific analysis described above, except we used $\beta=12$.

To identify functional pathways in each of the WGCNA modules, we performed an

Overrepresentation Analysis with WebGestaltR (Liao et al., 2019) using the Gene Ontology

Biological Processes database. For each module, we tested the genes found in each module

against a background list of all genes used to perform the WGCNA analysis. GO categories were

significantly enriched if the FDR value < 0.05. We then interrogated significant GO categories,

with an emphasis on immune and oxidative stress related pathways, in modules significantly

correlated with testosterone treatment. To find support for the Immunocompetence Handicap

Hypothesis, immune related categories (e.g., "immune system process") had to be significantly

enriched among genes expressed lower in testosterone treated animals. To find support for the

Oxidation Handicap Hypothesis, oxidative stress related GO categories had to be significantly

enriched among genes highly expressed (e.g., "response to oxidative stress") or gene exhibiting

lower expression levels (e.g., "antioxidant activity") following testosterone treatment.

3. RESULTS

After filtering, we used 13,509 manakin genes and 13,946 quail genes for PCA and WGCNA

network construction. Testosterone treatment had pronounced effects on gene expression and

individuals clustered by treatment in both species-specific comparisons (Figure 1) and in the

analysis ran on orthologous genes (Supplemental Figure 2).

3.1 WGCNA – Quail

WGCNA constructed 18 modules for quail, six of which were correlated with testosterone

treatment (Supplemental Figure 3). The yellow module (925 genes, r=-0.74) was strongly

enriched for broad immune related GO categories spanning both the innate (e.g. "inflammatory

response") and adaptive (e.g. "T cell activation") arms of the immune system (Table 1,

Supplemental Table 1). This represents a significant decrease in immune gene expression

following treatment (Figure 2A). The yellow module hubs were *SASH3*, *ITGB2*, *SLAMF8*

(LOC107324444), *TRAF3IP3*, and *EVI2A*.

The purple and black modules were also negatively correlated with testosterone treatment

and were enriched for translation and muscle process GO categories respectively. Additionally,

we found two modules containing genes highly expressed following testosterone treatment. The

turquoise module was the most strongly correlated with testosterone treatment (4423 genes,

r=0.98). GO enrichment was largely driven by genes involved in the Golgi apparatus and

endoplasmic reticulum functions (Supplemental Table 1). The green module (795 genes, r=0.61)

was primarily enriched for broad transcriptional regulation and protein modification processes.

Lastly, we found no modules that were associated with either antioxidant defenses or oxidative

damage.

3.2 WGCNA – Manakin

WGCNA constructed 34 modules for manakin, 12 of which were correlated with testosterone

treatment (Supplemental Figure 3). Seven modules were correlated with muscle type but, given

none of these seven modules were also correlated with testosterone treatment, our results provide

no support for a tissue specific response to testosterone at the network level. Of the 12 modules,

7 were negatively correlated and 5 positively correlated with testosterone treatment. Like the

quail, manakins treated with testosterone also exhibited a significant decrease in immune gene

expression relative to control individuals (Figure 2B, Supplemental Table 2). The dark turquoise

module (198 genes, r=-0.71) was strongly enriched for a broad range of immune related GO

categories spanning both the innate and adaptive components of the immune system (Table 1,

Supplemental Table 2). The dark turquoise hub genes were *MHC1A* (LOC108639055), *INPPL1*

(LOC103767762), *CCL14* (LOC103758017), *CCL3L* (LOC103757995), and an uncharacterized

non-coding RNA (LOC108640668).

The remaining negatively correlated modules were primarily enriched for glycolysis and

mitochondria related categories (steel blue, pale turquoise). Among the positively correlated

modules, we only observed significant GO enrichment in the light cyan module, which

encompassed transcriptional regulation (Supplemental Table 2). Lastly, like the quail, we found

no modules with significant enrichment of pathways related to oxidative damage or protection.

3.3 WGCNA – Subset Analysis

Our subsampling approach resulted in 43 modules for quail subset 1, 56 modules for

quail subset 2, 55 modules for manakin subset 1, and 53 modules for manakin subset 2

(Supplemental Figure 4). Four modules in quail subset 1, six modules in quail subset 2, and three

each in the manakin subsets were correlated with testosterone treatment and, of these modules, a

single immune-related module was correlated with testosterone-treatment in both quail subsets

and one manakin subset. Both the greenyellow module (n=518 genes) in quail subset 1 and

antiquewhite1 module (n=1280 genes) in quail subset 2 were primarily enriched for lymphocyte

activation and T cell signaling pathways. Two hundred & fifty-two genes were shared between

these two quail modules and these genes were enriched for broad immune system processes,

including lymphocyte and T cell activation. Combined, the greenyellow and antiquewhite1

modules contain 688/925 genes found in the quail yellow module from the full analysis. The

blanchedalmond module (n=102 genes) in manakin subset 1, which represents pectoralis tissue,

was primarily enriched for interleukin and cytokine signaling (Supplemental Table 3). This

module contained 41/198 genes found in the dark turquoise module from the full analysis of

manakin data. In each case, the module consisted of immune related genes exhibiting lower

expression levels in the testosterone treated group relative to the control group (Supplementary

Table 3). Manakin subset 2, which represents scalpulohumeralis caudalis tissue, did not have an

immunosuppression signature. None of the subsets exhibited an oxidative stress signature

correlated with testosterone treatment.

3.4 WGCNA – Shared response among Quail, Manakin, & Zebra Finch

Our integration of all three species resulted in 18 modules, 8 of which were correlated

with testosterone treatment in all three species (i.e., $p < 0.05$; Supplemental Figure 5;

Supplemental Table 4). Of these testosterone responsive modules, the purple module (195 genes,

$r = -0.58$) was significantly enriched for broad immune related categories (Figure 3A). This

result represents lower expression of these module genes in testosterone treated animals (relative

to control animals) and reflects the shared response of immunosuppression among all three

species. Interestingly, the black module (1,331 genes, $r = 0.58$) also showed significant

enrichment of neutrophil mediated immunity, which represents increased expression of these

module genes in testosterone treated animals (Figure 3B). This analysis reveals that while

testosterone leads to widespread immunosuppression, a small component of the immune system

is also activated.

4. DISCUSSION

In this study, we examined transcriptomic responses to experimentally increased circulating

testosterone in three species of bird. As expected based on the pleiotropic nature of sex steroids

(Ketterson and Nolan, Jr., 1999, Hau 2007), our gene network analyses demonstrate that, in all

three species, individuals treated with testosterone exhibited altered expression patterns of a

diverse suite of genes relative to controls (e.g., regulation of immune response, hemopoiesis,

endosomal transport). In the context of handicap hypotheses, the expression genes associated

with immune function was significantly lower in the testosterone-treated group lower relative to

controls, providing transcriptomic support for the Immunocompetence Handicap Hypothesis.

These results also begin to elucidate the molecular pathways (e.g., lymphocyte activation,

cytokine signaling, T cell activation) potentially underlying the testosterone-induced

immunosuppression previously documented using physiological measures of immune function in

Foo et al. (2017). Moreover, testosterone is necessary to produce the secondary sexual

characteristics associated with mating in all three species (Cynx et al., 2005, McGraw et al. 2006,

Schlinger et al., 2013, Hiyama et al., 2018). Our results therefore begin to detail the potential

molecular pathways underlying the trade-off between the expression of sexually selected traits

and immune function in these species. We did not find support for the Oxidation Handicap

Hypothesis as there was no enrichment of genes expressed related to oxidative damage, nor

suppression of genes related to antioxidant defenses in either species. These results suggest that

costs associated with maintaining high levels of circulating testosterone are not via pathways

associated with an individual's susceptibility or exposure to oxidative stress. Importantly,

oxidative stress could still be involved in enforcing the costs of reproduction or sexually selected

traits, but our results suggest that this cost is not borne out via molecular pathways that are

sensitive to testosterone, at least in the limited number of tissues and species examined here.

Our analyses revealed that transcriptomic immunosuppression was broad, encompassing

aspects of both innate immunity (e.g., interferon signaling and cytokine signaling) as well as

adaptive immunity (e.g., antigen processing and presentation; Table 1, Supplemental Tables 1 &

2). While the observed effect of testosterone could occur through non-genomic pathways,

previous work examining the relationship between androgen receptors and the immune system

suggests genomic pathways are involved (Trigunaite et al., 2015; Segner et al., 2017; Gubbels

Bupp and Jorgensen, 2018). More specifically, while testosterone exposure and subsequent

androgen receptor activity can promote innate immune cell differentiation and development,

testosterone also reduces activity of these cells (Gubbels Bupp and Jorgensen, 2018). Indeed, the

hub genes of the immune related modules highlight broad suppression of innate immune

signaling (quail yellow: *SASH3*, *SLAMF8*, *TRAF3IP3*; manakin dark turquoise: *INPPL1*, *CCL14*,

CCL3L, ncRNA; (Beer et al., 2005; Veillette, 2010; Dauphinee et al., 2013; Sokol and Luster,

2015; Zou et al., 2015; Thomas et al., 2017; Wang et al., 2018). Similarly, testosterone impacted

expression of genes involved in the adaptive immune system. Testosterone exposure reduces T

cell activity, a crucial component of proper adaptive immune signaling, (Lin et al., 2010; Kissick

et al., 2014), and this is a prominent signature in both quail (Supplemental Table 1) and manakin

(Table 1). In addition to suppression of T cell activity in manakin, we also identified MHC class

IA as a hub gene in the manakin dark turquoise module. MHC class IA binds and presents viral

peptides to CD8+ T cells, which is a critical component of the adaptive immune response

(Neefjes et al., 2011). Previous work has shown suppressive effects of testosterone on CD4+ T

cells/MHC class IIB (Lin et al., 2010) and CD8+ T cells (Page et al., 2006). However, our study

is the first to describe suppression of genes involved in T cell activity as well as MHC class I.

The subset analysis demonstrated that small sample sizes can influence both module

construction and functional enrichment. For quail, we identified a single module in each subset

that was significantly correlated with testosterone treatment and enriched for immune pathways.

As with the original full quail analysis, both immune related modules from the quail subsets

exhibited enrichment for T cell related activity, suggesting that testosterone has a robust effect

on T cell signaling in the quail foam gland (Supplemental Table 3). In contrast, we identified a

single immune related module correlated with testosterone treatment only in the manakin subset

containing pectoralis tissue. Whereas the full manakin analysis revealed a T cell signaling

signature, the subset analysis primarily exhibited a cytokine production and signaling signature.

Individual variation in gene expression, tissue-specific expression, responses to the treatment, as

well as the limited sample size likely influenced our ability to detect an immunosuppression

signature in both manakin subsets. Indeed, the manakin pectoralis subset exhibited a cytokine

signature not observed in the full analysis, suggesting individuals and/or muscle types exhibited

quite variable responses to testosterone treatment. Overall, the results of the subset analysis

highlight the sensitivity of WGCNA to small sample sizes and underscore the importance of

adequate sample sizes for future studies using this analytical approach.

As with each species-specific analysis, the WGCNA conducted on orthologous genes

(i.e., evolutionary conserved genes) also identified a broad signature of immunosuppression in

the testosterone-treated group but not the control groups. This result provides preliminary insight

into the evolutionary conserved transcriptomic response to testosterone and provides additional

support for the immunocompetence handicap hypothesis. The increased power of the WGCNA

conducted on the orthologous genes (n=30 samples) may have reduced the influence of pre-

existing individual differences and allowed us to detect more subtle changes in gene expression

that were missed in the species-specific analyses (n=12 samples). For instance, a module

encompassing genes related to neutrophil-mediated immunity was highly expressed following

testosterone treatment. Neutrophil activity, a hallmark of innate immunity, is associated with

inflammation predominantly in lymphoid tissues, but their broader tissue-specific functions

remain uncertain (reviewed in Ng et al., 2019). Interestingly, we also observed an inflammation

signature within the purple module (Supplemental Table 4), which contains genes exhibiting

lower expression levels in the testosterone treatment relative to controls. Overall, these results

suggest the changes in inflammation-related gene expression following testosterone treatment are

dynamic and that studies with larger sample size are better able to detect more complex

responses.

Our results were based on the transcriptomes from skeletal muscles or the quail foam

gland, tissues which are not traditionally studied in avian immunology (Rose, 1979; Schat et al.,

2014). A growing body of research on fish and mammals does however suggest that muscle

tissues play an important role in immune responses (reviewed in Valenzuela et al. 2017).

Nonetheless, in combination with the testosterone-related patterns of immunosuppression found

in Foo et al. (2017), our study provides some additional evidence to suggest that RNA-seq can

detect functional signatures in non-traditional tissues (e.g., Louder et al., 2018). As such, public

repositories of genomic data will become increasingly valuable to test evolutionary hypotheses in

novel ways (e.g. Shultz & Sackton 2019).

5. CONCLUSION AND FUTURE DIRECTIONS

Testosterone-mediated handicap hypotheses have captivated biologists since their development

but have primarily been tested using physiological measures of oxidative stress or immune

function (e.g., Roberts et al. 2004, Foo et al. 2017). Here we develop a transcriptomic approach

for testing these hypotheses. Our findings, while limited by sample size and taxonomic diversity,

document species-specific and evolutionary conserved signatures of immunosuppression in three

species of bird. These results provide a foundation for future transcriptomic studies

encompassing additional tissue types and/or more taxonomically diverse species. Future studies

should prioritize measuring testosterone's effect on gene expression using a within-individual

sampling approach to mitigate the influence of pre-existing individual differences in gene

expression. Studies should also strive for sample sizes > 20, as recommend by the developers of

WGCNA (Langfelder & Horvath 2017). The importance of larger sample sizes is exemplified

here by the subset analysis which shows that modules built using smaller sample sizes

recapitulate an immunosuppression signature most but not all the times (i.e., 3 out of 4 subsets

included immune-related modules significantly correlated with treatment in this study).

Additionally, the immune modules within the 3 subsets exhibited variable immunological

pathway enrichment compared to the full analysis potentially due to individual variation

influencing the genes found in each module. When terminal sampling is the only option for

collecting samples, a common scenario for transcriptomic studies, larger sample sizes are needed

to reduce the influence of such pre-existing individual differences. Furthermore, future works

should also consider conducting experimental infections and/or immune challenges in

combination with RNA-seq analyses to examine how transcriptomic signatures relate to immune

function. Novel endocrine-based experiments similar to Goymann et al. (2015) and Goymann

and Dávila (2017) paired with RNA-seq analyses can also shed light on the transcriptomic

responses to acute stimulation of the hypothalamic-pituitary-gonadal axis (i.e., the endocrine

cascade associated with gonadal testosterone production). Future transcriptomic studies should

also consider the best practices for experimental design to ensure appropriate sample sizes are

selected and that any results are biologically meaningful (Schurch et al. 2016). Overall, these

integrative, comparative, and mechanistic approaches will ultimately provide novel insights into

the mechanisms underlying handicap hypotheses and, more broadly, the evolution of sexually

selected traits.

ACKNOWLEDGEMENTS

We would like to thank Ignacio Moore, Dana Hawley, Heather Watts, Christopher Balakrishnan,

and three anonymous reviewers for providing helpful feedback on earlier versions of this

manuscript.

COMPETING INTERESTS

The authors declare no competing interests.

FUNDING

BJV was supported by a grant from the U.S. National Science Foundation (IOS-1353093).

TABLES AND FIGURES

Table 1. GO enrichment for modules (from both species) that are associated with immune

function. The top 5 gene ontology (GO) categories are presented, along with FDR adjusted p-

value and WebGestaltR enrichment score.

GO ID	Description	FDR	Enrichment
Quail, Yellow Module			
GO:0050778	positive regulation of immune response	0	3.52
GO:0001816	cytokine production	0	3.36
GO:0046649	lymphocyte activation	0	4.52
GO:0045087	innate immune response	0	3.02
GO:0019221	cytokine-mediated signaling pathway	0	3.47
Manakin, Dark Turquoise Module			
GO:0030217	T cell differentiation	6.56×10^{-12}	12.78
GO:0002521	leukocyte differentiation	6.56×10^{-12}	7.83
GO:0030098	lymphocyte differentiation	2.27×10^{-11}	9.69
GO:0042110	T cell activation	1.57×10^{-10}	8.00
GO:0030097	hemopoiesis	6.11×10^{-09}	4.99

Figure 1. PCA of (A) manakin and (B) quail. Samples separate by treatment along PC3 for

manakin and PC1 for quail. Each circle represents a sample and is color-coded by treatment.

Manakin samples are labeled by muscle type. Ellipses represent 95% confidence intervals.

Figure 2. Expression heatmaps of the (A) Manakin Dark Turquoise Module and (B) Quail

Yellow Module, which represent low expression of immune related genes in testosterone, but not

control, treated animals. Each column represents a sample color coded by treatment or muscle

type. Each row represents a module gene. High expression is indicated by orange colors and low

expression is represented by blue colors.

Figure 3. PCA and functional enrichment of the (A) purple and (B) black modules from the orthologous gene WGCNA. PCAs were constructed using genes belonging to each module. The shape of the points represents species and the color represents treatment. The dashed line in the enrichment bar plots represents FDR=0.05.

REFERENCES

Adkins-Regan, E., 1999. Foam Produced by Male Coturnix Quail: What Is Its Function? The Auk 116, 184–193. https://doi.org/10.2307/4089465

Alonso-Alvarez, C., Bertrand, S., Faivre, B., Chastel, O., Sorci, G., 2007. Testosterone and oxidative stress: the oxidation handicap hypothesis. Proceedings of the Royal Society B: Biological Sciences 274, 819–825. https://doi.org/10.1098/rspb.2006.3764

Anders, S., Pyl, P.T., Huber, W., 2015. HTSeq-A Python framework to work with high-throughput sequencing data. Bioinformatics 31, 166–169. https://doi.org/10.1093/bioinformatics/btu638

Andersson, M.B., 1994. Sexual selection, Monographs in behavior and ecology. Princeton University Press, Princeton, N.J.

Baldo, S., Mennill, D.J., Guindre-Parker, S., Gilchrist, H.G., Love, O.P., 2015. The Oxidative Cost of Acoustic Signals: Examining Steroid Versus Aerobic Activity Hypotheses in a Wild Bird. Ethology 121, 1081–1090. https://doi.org/10.1111/eth.12424

Ball, G.F., Balthazart, J., 2009. Neuroendocrine Regulation of Reproductive Behavior in Birds, in: Hormones, Brain and Behavior. Elsevier, pp. 855–897. https://doi.org/10.1016/B978-008088783-8.00025-5

Barshis, D.J., Ladner, J.T., Oliver, T.A., Seneca, F.O., Traylor-Knowles, N., Palumbi, S.R., 2013. Genomic basis for coral resilience to climate change. Proceedings of the National Academy of Sciences 110, 1387–1392. https://doi.org/10.1073/pnas.1210224110

Beer, S., Scheikl, T., Reis, B., Huser, N., Pfeffer, K., Holzmann, B., 2005. Impaired Immune Responses and Prolonged Allograft Survival in Sly1 Mutant Mice. Molecular and Cellular Biology 25, 9646–9660. https://doi.org/10.1128/MCB.25.21.9646-9660.2005

Blighe K., Lun A., 2020. PCAtools: PCAtools: Everything Principal Components Analysis. R package version 2.0.0, https://github.com/kevinblighe/PCAtools.

Breschi, A., Djebali, S., Gillis, J., Pervouchine, D.D., Dobin, A., Davis, C.A., Gingeras, T.R., Guigó, R., 2016. Gene-specific patterns of expression variation across organs and species. Genome Biology 17. https://doi.org/10.1186/s13059-016-1008-y

Buchanan, K.L., Evans, M.R., Goldsmith, A.R., Bryant, D.M., Rowe, L.V., 2001. Testosterone influences basal metabolic rate in male house sparrows: a new cost of dominance signalling? Proc. R. Soc. Lond. B 268, 1337–1344. https://doi.org/10.1098/rspb.2001.1669

Casagrande, S., Costantini, D., Groothuis, T.G.G., 2012. Interaction between sexual steroids and immune response in affecting oxidative status of birds. Comparative Biochemistry and Physiology Part A: Molecular & Integrative Physiology 163, 296–301. https://doi.org/10.1016/j.cbpa.2012.07.018

Cox, R.M., Cox, C.L., McGlothlin, J.W., Card, D.C., Andrew, A.L., Castoe, T.A., 2017. Hormonally Mediated Increases in Sex-Biased Gene Expression Accompany the Breakdown of Between-Sex Genetic Correlations in a Sexually Dimorphic Lizard. The American Naturalist 189, 315–332. https://doi.org/10.1086/690105

Cynx, J., Bean, N., Rossman, I., 2005. Testosterone implants alter the frequency range of zebra finch songs. Hormones and Behavior 47, 446–451. https://doi.org/10.1016/j.yhbeh.2004.11.018

Dauphinee, S.M., Clayton, A., Hussainkhel, A., Yang, C., Park, Y.-J., Fuller, M.E., Blonder, J., Veenstra, T.D., Karsan, A., 2013. SASH1 Is a Scaffold Molecule in Endothelial TLR4

Signaling. The Journal of Immunology 191, 892–901.
https://doi.org/10.4049/jimmunol.1200583

Day, L.B., Fusani, L., Hernandez, E., Billo, T.J., Sheldon, K.S., Wise, P.M., Schlinger, B.A., 2007. Testosterone and its effects on courtship in golden-collared manakins (Manacus vitellinus): Seasonal, sex, and age differences. Hormones and Behavior 51, 69–76. https://doi.org/10.1016/j.yhbeh.2006.08.006

Dobin, A., Davis, C.A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., Batut, P., Chaisson, M., Gingeras, T.R., 2013. STAR: ultrafast universal RNA-seq aligner. Bioinformatics 29, 15–21.

Finseth, F.R., Harrison, R.G., 2018. Genes Integral to the Reproductive Function of Male Reproductive Tissues Drive Heterogeneity in Evolutionary Rates in Japanese Quail. G3: Genes|Genomes|Genetics 8, 39–51. https://doi.org/10.1534/g3.117.300095

Folstad, I., Karter, A.J., 1992. Parasites, Bright Males, and the Immunocompetence Handicap. The American Naturalist 139, 603–622. https://doi.org/10.1086/285346

Foo, Y.Z., Nakagawa, S., Rhodes, G., Simmons, L.W., 2017. The effects of sex hormones on immune function: a meta-analysis: Sex hormones and immune function. Biological Reviews 92, 551–571. https://doi.org/10.1111/brv.12243

Fusani, L., 2008. Testosterone control of male courtship in birds. Hormones and Behavior 54, 227–233. https://doi.org/10.1016/j.yhbeh.2008.04.004

Fuxjager, M.J., Lee, J.-H., Chan, T.-M., Bahn, J.H., Chew, J.G., Xiao, X., Schlinger, B.A., 2016. Research Resource: Hormones, Genes, and Athleticism: Effect of Androgens on the Avian Muscular Transcriptome. Molecular Endocrinology 30, 254–271. https://doi.org/10.1210/me.2015-1270

Gao, J., Liu, S., Zhang, Y., Yang, Y., Yuan, C., Chen, S., Wang, Z., 2015. Effects of 17 α-methyltestosterone on transcriptome, gonadal histology and sex steroid hormones in rare minnow Gobiocypris rarus. Comparative Biochemistry and Physiology Part D: Genomics and Proteomics 15, 20–27. https://doi.org/10.1016/j.cbd.2015.05.001

Goymann, W., Flores Dávila, P., 2017. Acute peaks of testosterone suppress paternal care: evidence from individual hormonal reaction norms. Proceedings of the Royal Society B: Biological Sciences 284, 20170632. https://doi.org/10.1098/rspb.2017.0632

Goymann, W., Villavicencio, C.P., Apfelbeck, B., 2015. Does a short-term increase in testosterone affect the intensity or persistence of territorial aggression? — An approach using an individual's hormonal reactive scope to study hormonal effects on behavior. Physiology & Behavior 149, 310–316. https://doi.org/10.1016/j.physbeh.2015.06.029

Grafen, A., 1990. Biological signals as handicaps. Journal of Theoretical Biology 144, 517–546. https://doi.org/10.1016/S0022-5193(05)80088-8

Gubbels Bupp, M.R., Jorgensen, T.N., 2018. Androgen-Induced Immunosuppression. Frontiers in Immunology 9. https://doi.org/10.3389/fimmu.2018.00794

Hau, M., 2007. Regulation of male traits by testosterone: implications for the evolution of vertebrate life histories. BioEssays 29, 133–144. https://doi.org/10.1002/bies.20524

Hiyama, G., Mizushima, S., Matsuzaki, M., Tobari, Y., Choi, J.-H., Ono, T., Tsudzuki, M., Makino, S., Tamiya, G., Tsukahara, N., Sugita, S., Sasanami, T., 2018. Female Japanese quail visually differentiate testosterone-dependent male attractiveness for mating preferences. Scientific Reports 8. https://doi.org/10.1038/s41598-018-28368-z

Hoogenboom, M.O., Metcalfe, N.B., Groothuis, T.G.G., de Vries, B., Costantini, D., 2012. Relationship between oxidative stress and circulating testosterone and cortisol in pre-

spawning female brown trout. Comparative Biochemistry and Physiology Part A:
Molecular & Integrative Physiology 163, 379–387.
https://doi.org/10.1016/j.cbpa.2012.07.002
Huang, Y., Li, Y., Burt, D.W., Chen, H., Zhang, Y., Qian, W., Kim, H., Gan, S., Zhao, Yiqiang,
Li, J., Yi, K., Feng, H., Zhu, P., Li, B., Liu, Q., Fairley, S., Magor, K.E., Du, Z., Hu, X.,
Goodman, L., Tafer, H., Vignal, A., Lee, T., Kim, K.-W., Sheng, Z., An, Y., Searle, S.,
Herrero, J., Groenen, M.A.M., Crooijmans, R.P.M.A., Faraut, T., Cai, Q., Webster, R.G.,
Aldridge, J.R., Warren, W.C., Bartschat, S., Kehr, S., Marz, M., Stadler, P.F., Smith, J.,
Kraus, R.H.S., Zhao, Yaofeng, Ren, L., Fei, J., Morisson, M., Kaiser, P., Griffin, D.K.,
Rao, M., Pitel, F., Wang, J., Li, N., 2013. The duck genome and transcriptome provide
insight into an avian influenza virus reservoir species. Nature Genetics 45, 776–783.
https://doi.org/10.1038/ng.2657
Isaksson, C., While, G., McEvoy, J., van de Crommenacker, J., Olsson, M., Groothuis, T.,
Komdeur, J., Wapstra, E., 2011. Aggression, but not testosterone, is associated to
oxidative status in a free-living vertebrate. Behaviour 148, 713–731.
https://doi.org/10.1163/000579511X574204
Ketterson, E.D., Nolan, Jr., V., 1999. Adaptation, Exaptation, and Constraint: A Hormonal
Perspective. The American Naturalist 154, S4–S25. https://doi.org/10.1086/303280
Ketterson, E.D., Atwell, J.W., McGlothlin, J.W., 2009. Phenotypic integration and
independence: Hormones, performance, and response to environmental change.
Integrative and Comparative Biology 49, 365–379. https://doi.org/10.1093/icb/icp057
Kim, J., Kim, H., Son, C., 2018. Tissue-Specific Profiling of Oxidative Stress-Associated
Transcriptome in a Healthy Mouse Model. International Journal of Molecular Sciences
19, 3174. https://doi.org/10.3390/ijms19103174
Kinsella, R.J., Kahari, A., Haider, S., Zamora, J., Proctor, G., Spudich, G., Almeida-King, J.,
Staines, D., Derwent, P., Kerhornou, A., Kersey, P., Flicek, P., 2011. Ensembl BioMarts:
a hub for data retrieval across taxonomic space. Database 2011, bar030–bar030.
doi:10.1093/database/bar030
Kissick, H.T., Sanda, M.G., Dunn, L.K., Pellegrini, K.L., On, S.T., Noel, J.K., Arredouani, M.S.,
2014. Androgens alter T-cell immunity by inhibiting T-helper 1 differentiation.
Proceedings of the National Academy of Sciences 111, 9887–9892.
https://doi.org/10.1073/pnas.1402468111
Kokko, H., Brooks, R., Jennions, M.D., Morley, J., 2003. The evolution of mate choice and
mating biases. Proc. R. Soc. Lond. B 270, 653–664.
https://doi.org/10.1098/rspb.2002.2235
Langfelder, P., & Horvath, S. 2017. WGCNA package: Frequently Asked Questions.
https://horvath.genetics.ucla.edu/html/CoexpressionNetwork/Rpackages/WGCNA/faq.ht
ml (accessed 20 July 2020).
Langfelder, P., Horvath, S., 2008. WGCNA: an R package for weighted correlation network
analysis. BMC Bioinformatics 9, 559–559. https://doi.org/10.1186/1471-2105-9-559
Langfelder, P., Luo, R., Oldham, M.C., Horvath, S., 2011. Is My Network Module Preserved and
Reproducible? PLoS Computational Biology 7, e1001057.
https://doi.org/10.1371/journal.pcbi.1001057
Liao, Y., Wang, J., Jaehnig, E.J., Shi, Z., Zhang, B., 2019. WebGestalt 2019: gene set analysis
toolkit with revamped UIs and APIs. Nucleic Acids Research 47, W199–W205.
https://doi.org/10.1093/nar/gkz401

Lin, A.A., Wojciechowski, S.E., Hildeman, D.A., 2010. Androgens suppress antigen-specific T cell responses and IFN-γ production during intracranial LCMV infection. Journal of Neuroimmunology 226, 8–19. https://doi.org/10.1016/j.jneuroim.2010.05.026

Love, M.I., Huber, W., Anders, S., 2014. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. Genome biology 15, 550–550. https://doi.org/10.1186/s13059-014-0550-8

Marini, F., Binder, H., 2019. pcaExplorer: an R/Bioconductor package for interacting with RNA-seq principal components. BMC Bioinformatics 20, 331. https://doi.org/10.1186/s12859-019-2879-1

McGraw, K.J., Correa, S.M., Adkins-Regan, E., 2006. Testosterone upregulates lipoprotein status to control sexual attractiveness in a colorful songbird. Behav Ecol Sociobiol 60, 117–122. https://doi.org/10.1007/s00265-005-0135-3

Metcalfe, N.B., Alonso-Alvarez, C., 2010. Oxidative stress as a life-history constraint: the role of reactive oxygen species in shaping phenotypes from conception to death: Oxidative stress as a life-history constraint. Functional Ecology 24, 984–996. https://doi.org/10.1111/j.1365-2435.2010.01750.x

Monson, C., Forsgren, K., Goetz, G., Harding, L., Swanson, P., Young, G., 2017. A teleost androgen promotes development of primary ovarian follicles in coho salmon and rapidly alters the ovarian transcriptome†. Biology of Reproduction 97, 731–745. https://doi.org/10.1093/biolre/iox124

Mougeot, F., Martínez-Padilla, J., Webster, L.M.I., Blount, J.D., Pérez-Rodríguez, L., Piertney, S.B., 2009. Honest sexual signalling mediated by parasite and testosterone effects on oxidative balance. Proceedings of the Royal Society B: Biological Sciences 276, 1093–1100. https://doi.org/10.1098/rspb.2008.1570

Neefjes, J., Jongsma, M.L.M., Paul, P., Bakke, O., 2011. Towards a systems understanding of MHC class I and MHC class II antigen presentation. Nature Reviews Immunology 11, 823–836. https://doi.org/10.1038/nri3084

Nelson, R.J., 2011. An introduction to behavioral endocrinology, 4th ed. ed. Sinauer Associates, Sunderland, MA.

Ng, L.G., Ostuni, R., Hidalgo, A., 2019. Heterogeneity of neutrophils. Nat Rev Immunol 19, 255–265. https://doi.org/10.1038/s41577-019-0141-8

Page, S.T., Plymate, S.R., Bremner, W.J., Matsumoto, A.M., Hess, D.L., Lin, D.W., Amory, J.K., Nelson, P.S., Wu, J.D., 2006. Effect of medical castration on CD4 [+] CD25 [+] T cells, CD8 [+] T cell IFN-γ expression, and NK cells: a physiological role for testosterone and/or its metabolites. American Journal of Physiology-Endocrinology and Metabolism 290, E856–E863. https://doi.org/10.1152/ajpendo.00484.2005

Peters, A., 2000. Testosterone treatment is immunosuppressive in superb fairy–wrens, yet free–living males with high testosterone are more immunocompetent. Proceedings of the Royal Society of London. Series B: Biological Sciences 267, 883–889. https://doi.org/10.1098/rspb.2000.1085

Reed, W.L., Clark, M.E., Parker, P.G., Raouf, S.A., Arguedas, N., Monk, D.S., Snajdr, E., Nolan Jr., V., Ketterson, E.D., 2006. Physiological Effects on Demography: A Long☐Term Experimental Study of Testosterone's Effects on Fitness. The American Naturalist 167, 667–683. https://doi.org/10.1086/503054

Ritchie, M.E., Phipson, B., Wu, D., Hu, Y., Law, C.W., Shi, W., Smyth, G.K., 2015. limma powers differential expression analyses for RNA-sequencing and microarray studies. Nucleic Acids Research 43, e47–e47. https://doi.org/10.1093/nar/gkv007

Roberts, M.L., Buchanan, K.L., Evans, M.R., 2004. Testing the immunocompetence handicap hypothesis: a review of the evidence. Animal Behaviour 68, 227–239. https://doi.org/10.1016/j.anbehav.2004.05.001

Rose, M.E., 1979. The Immune System in Birds. Journal of the Royal Society of Medicine 72, 701–705. https://doi.org/10.1177/014107687907200914

Schat, K., Kaspers, B., Kaiser, P. (Eds.), 2014. Avian Immunology. Elsevier. https://doi.org/10.1016/C2011-0-07243-5

Schlinger, B.A., Barske, J., Day, L., Fusani, L., Fuxjager, M.J., 2013. Hormones and the neuromuscular control of courtship in the golden-collared manakin (Manacus vitellinus). Frontiers in Neuroendocrinology 34, 143–156. https://doi.org/10.1016/j.yfrne.2013.04.001

Schurch, N.J., Schofield, P., Gierliński, M., Cole, C., Sherstnev, A., Singh, V., Wrobel, N., Gharbi, K., Simpson, G.G., Owen-Hughes, T., Blaxter, M., Barton, G.J., 2016. How many biological replicates are needed in an RNA-seq experiment and which differential expression tool should you use? RNA 22, 839–851. https://doi.org/10.1261/rna.053959.115

Segner, H., Verburg-van Kemenade, B.M.L., Chadzinska, M., 2017. The immunomodulatory role of the hypothalamus-pituitary-gonad axis: Proximate mechanism for reproduction-immune trade offs? Developmental & Comparative Immunology 66, 43–60. https://doi.org/10.1016/j.dci.2016.07.004

Shultz, A.J., Sackton, T.B., 2019. Immune genes are hotspots of shared positive selection across birds and mammals. eLife 8, e41815. https://doi.org/10.7554/eLife.41815

Sokol, C.L., Luster, A.D., 2015. The Chemokine System in Innate Immunity. Cold Spring Harbor Perspectives in Biology 7, a016303. https://doi.org/10.1101/cshperspect.a016303

Stearns, S.C., 1992. The evolution of life histories. Oxford University Press, Oxford ; New York.

Taff, C.C., Freeman-Gallant, C.R., 2014. An experimental test of the testosterone mediated oxidation handicap hypothesis in a wild bird. Hormones and Behavior 66, 276–282. https://doi.org/10.1016/j.yhbeh.2014.05.006

Thomas, M.P., Erneux, C., Potter, B.V.L., 2017. SHIP2: Structure, Function and Inhibition. ChemBioChem 18, 233–247. https://doi.org/10.1002/cbic.201600541

Trigunaite, A., Dimo, J., Jørgensen, T.N., 2015. Suppressive effects of androgens on the immune system. Cellular Immunology 294, 87–94. https://doi.org/10.1016/j.cellimm.2015.02.004

Valenzuela, C.A., Zuloaga, R., Poblete-Morales, M., Vera-Tobar, T., Mercado, L., Avendaño-Herrera, R., Valdés, J.A., Molina, A., 2017. Fish skeletal muscle tissue is an important focus of immune reactions during pathogen infection. Developmental & Comparative Immunology 73, 1–9. https://doi.org/10.1016/j.dci.2017.03.004

Veillette, A., 2010. SLAM-Family Receptors: Immune Regulators with or without SAP-Family Adaptors. Cold Spring Harbor Perspectives in Biology 2, a002469–a002469. https://doi.org/10.1101/cshperspect.a002469

Wang, M., Jiang, S., Wu, W., Yu, F., Chang, W., Li, P., Wang, K., 2018. Non-coding RNAs Function as Immune Regulators in Teleost Fish. Frontiers in Immunology 9. https://doi.org/10.3389/fimmu.2018.02801

Wang, Z., Gerstein, M., Snyder, M., 2009. RNA-Seq: a revolutionary tool for transcriptomics. Nature Reviews Genetics 10, 57–63. https://doi.org/10.1038/nrg2484

Weaver, R.J., Koch, R.E., Hill, G.E., 2017. What maintains signal honesty in animal colour displays used in mate choice? Phil. Trans. R. Soc. B 372, 20160343. https://doi.org/10.1098/rstb.2016.0343

Wenzel, M.A., Webster, L.M.I., Paterson, S., Mougeot, F., Martínez-Padilla, J., Piertney, S.B., 2013. A transcriptomic investigation of handicap models in sexual selection. Behavioral Ecology and Sociobiology 67, 221–234. https://doi.org/10.1007/s00265-012-1442-0

Williams, T.D., 2008. Individual variation in endocrine systems: moving beyond the 'tyranny of the Golden Mean.' Phil. Trans. R. Soc. B 363, 1687–1698. https://doi.org/10.1098/rstb.2007.0003

Wingfield, J.C., Lynn, S.E., Soma, K.K., 2001. Avoiding the 'Costs' of Testosterone: Ecological Bases of Hormone-Behavior Interactions. Brain Behav Evol 57, 239–251. https://doi.org/10.1159/000047243

Zahavi, A., 1975. Mate selection—A selection for a handicap. Journal of Theoretical Biology 53, 205–214. https://doi.org/10.1016/0022-5193(75)90111-3

Zera, A.J., Harshman, L.G., 2001. The Physiology of Life History Trade-Offs in Animals. Annual Review of Ecology and Systematics 32, 95–126. https://doi.org/10.1146/annurev.ecolsys.32.081501.114006

Zheng, Y., Yuan, J., Meng, S., Chen, J., Gu, Z., 2019. Testicular transcriptome alterations in zebrafish (Danio rerio) exposure to 17β-estradiol. Chemosphere 218, 14–25. https://doi.org/10.1016/j.chemosphere.2018.11.092

Zou, Q., Jin, J., Xiao, Y., Hu, H., Zhou, X., Jie, Z., Xie, X., Li, J.Y.H., Cheng, X., Sun, S.-C., 2015. T cell development involves TRAF3IP3-mediated ERK signaling in the Golgi. The Journal of Experimental Medicine 212, 1323–1336. https://doi.org/10.1084/jem.20150110