Exploring the niche concept in a simple metaorganism

Peter Deines[1]*, Katrin Hammerschmidt[2] and Thomas CG Bosch[1]

[1]Zoological Institute, Christian-Albrechts-University Kiel, 24118 Kiel, Germany
[2]Institute of General Microbiology, Christian-Albrechts-University Kiel, 24118 Kiel, Germany

*Correspondence:
Peter Deines
pdeines@zoologie.uni-kiel.de

Abstract
Organisms and their resident microbial communities - the microbiome - form a complex and mostly stable ecosystem. It is known that the specific composition and abundance of certain bacterial species have a major impact on host health and *Darwinian fitness*, but the processes that lead to these microbial patterns have not yet been identified. We here apply the niche concept and trait-based approaches as a first step in understanding the patterns underlying microbial community assembly and structure in the simple metaorganism *Hydra*. We find that the carrying capacities in single associations do not reflect microbiota densities as part of the community, indicating a discrepancy between the fundamental and realized niche. Whereas in most cases, the realized niche is smaller than the fundamental one, as predicted by theory, the opposite is observed for *Hydra's* two main bacterial colonizers. Both, *Curvibacter sp.* and *Duganella sp.* benefit from association with the other members of the microbiome and reach higher fractions as in single colonisations. This cannot be linked to any particular trait that is relevant for interacting with the host or by the utilization of specific nutrients but is most likely determined by metabolic interactions between the individual microbiome members.

Keywords
fundamental niche, realized niche, microbiome, species abundance, microbial traits, *Hydra*

Running title
Microbial traits, niches, and the metaorganism

Number of words: 4590
Number of figures: 6

Introduction

Microbiomes contribute to ecosystems as key engines that power system-level processes (Falkowski et al., 2008). This also applies to host ecosystems, where they are critical in maintaining host health, survival, and function (Kau et al., 2011; McFall-Ngai et al., 2013). Despite their importance, the mechanisms governing microbiome assembly and composition are largely unknown. This is different for macroscopic communities, thanks to the application of niche (Holt, 2009; Leibold, 1995; Whittaker et al., 1973) and trait-based theories, which might also provide a useful framework for studying the ecology and evolution of microbiomes in metaorganisms (Kopac and Klassen, 2016).

The niche concept is one of the core concepts in ecology and has been rediscovered by modern ecology for explaining biodiversity and species coexistence patterns (Pocheville, 2015). The niche-based theory states that an ecological community is made up of a limited number of niches, each occupied by a single species. Hutchinson (Hutchinson, 1957) defined the *fundamental niche* as the needs of a species for it to maintain a positive population growth rate, disregarding biotic interactions (Hutchinson, 1957; Pearman et al., 2008). The fundamental niche therefore represents an idealized situation exclusive of interspecific interactions. The effect of biological interactions is taken into account in the definition of the *realized niche* (Hutchinson, 1957). This is the portion of the fundamental niche in which a species has a positive population growth rate, despite the constraining effects of biological interactions, such as inter-specific competition (Hutchinson, 1957; Pearman et al., 2008).

In the last two decades, the shift from taxonomy to function by using trait-based approaches has provided a detailed understanding of biodiversity-ecosystem functioning (Louca et al., 2018). Recently, this framework is also being used by microbial ecologists to study microbial biogeography (Green et al., 2008), or to unravel microbial biodiversity-ecosystem functioning relationships (Krause et al., 2014). Further, this approach allows studying microbiomes in the light of coexisting traits/ functions rather than of coexisting microbes (Martiny et al., 2015). A recent study successfully used this approach and analysed trait-based patterns to understand the mechanisms of community assembly and succession of the infant gut microbiome (Guittar et al., 2019). Microbial traits cover a range of phenotypic characteristics ranging from simple to complex, for example organic phosphate utilization, bacteriophage host range, cellulose degradation, biofilm formation, nitrogen fixation, methanogenesis, and salinity preference (Martiny et al., 2015). Potential microbial traits can be measured directly by laboratory assays (as in this study) or can be indirectly inferred based on genomic information.

The aim of this study is to apply the niche concept and trait-based theory to the metaorganism *Hydra* to gain insight into the mechanisms underlying the microbial community composition. We thus specifically extend the niche-assembly perspective, classically used for assessing species assembly and coexistence in abiotic environments, to a host-associated microbiome, thus a biotic environment.

The freshwater polyp *Hydra* and its microbiome have become a valuable model system for metaorganism research as it provides a bridge between the simplicity of synthetic communities and the complex mouse model (Deines and Bosch, 2016). The ectoderm is covered by a multi-layered glycocalyx, which is the habitat for a highly stable species-specific microbiome of low complexity (Bosch, 2013; Deines et al., 2017; Franzenburg et al., 2013). The most abundant bacterial colonizers of *Hydra vulgaris* (strain AEP) can be cultured and manipulated *in vitro*

(Bosch, 2013; Fraune et al., 2015; Wein et al., 2018), allowing the measurement of
phenotypic microbial traits and fitness. Fitness, as defined by niche theory, is the
positive population growth of the focal species, which in our study is that of the six
isolated microbiome members. Measurements of the performance of the bacterial
populations when grown singly, i.e. in the absence of the other microbial competitors
in vitro and *in vivo* (on germ-free *Hydra* polys), specify the fundamental niche. For
each species, we compare the fundamental niche to the realized niche, which we
calculated based on published data on the microbiome composition of wild-type and
conventionalized polys (germ-free polyps incubated with tissue homogenates of wild-
type animals) (Franzenburg et al., 2013; Murillo-Rincon et al., 2017). We also
measure phenotypic traits that might be connected to the success of the various
microbial species in occupying the fundamental niche; these are essentially traits that
might play a role in successfully populating their environment, the host, such as
biofilm formation, surface hydrophobicity (bacterial cells are more likely to attach to
surfaces with the same hydrophobicity), and nutrient utilisation patterns. As the
realized niche is determined by biological interactions of one species with its
associate microbial community, we focus on traits that are important when competing
with other species, such as growth rate, niche overlap, and niche breadth. Ultimately,
we take the traits and the ecological niches as determinants of species interactions,
which may infer the assembly and structure of the host-associated microbiome.
117

Materials and Methods
Animals used, culture conditions, and generation of germ-free animals
Hydra vulgaris (strain AEP) was used in the experiments and cultured according to standard procedures at 18°C in standardized *Hydra* culture medium (Lenhoff and Brown, 1970). Animals were fed three times a week with 1st instar larvae of *Artemia salina*. Germ-free polyps were obtained as previously described (Franzenburg et al., 2013; Murillo-Rincon et al., 2017). After two weeks of treatment, polyps were transferred into antibiotic-free sterile *Hydra* culture medium for recovery (four days). Sterility was confirmed by established methods (Franzenburg et al., 2013). During antibiotic treatment and re-colonization experiments, polyps were not fed.

Bacterial species and media
The bacterial species used in this study are *Curvibacter sp*. AEP1.3, *Duganella sp*. C1.2, *Undibacterium sp*. C1.1, *Acidovorax sp*. AEP1.4, *Pseudomonas sp*. C2.2, *Pelomonas sp*. AEP2.2., which are species isolated from the *Hydra vulgaris* (strain AEP) microbiome (Fraune et al., 2015). These bacteria were cultured from existing isolate stocks in R2A medium at 18°C, shaken at 250 r.p.m for 72 h before use in the different experiments.

Fundamental and realized niche
Germ-free polys were inoculated with single bacterial species using 5×10^3 cells in 1.5 ml Eppendorf tubes containing 1 ml of sterile *Hydra* culture medium. After 24 h of incubation, all polyps were washed with, and transferred to sterile *Hydra* culture medium, incubated at 18°C. After three days of incubation individual polyps were homogenized in an Eppendorf tube using a sterile pestle, after which serial dilutions of the homogenate were plated on R2A agar plates to determine colony-forming units (CFUs) per polyp (Deines et al., 2020).

  The carrying capacities of mono-associations provide information of the occupied niche space on the host in the absence of other microbial species that are part of the microbiome, and thus specifies the fundamental niche for each (as calculated from the proportion of each species from the sum of all).
Estimates of the realized niche are based on underlying original data of previously published values of community composition and relative abundances (as estimated based on OTUs) of wild-type and conventionalized polys that have been reported to be remarkably stable over time (Bosch, 2013; Franzenburg et al., 2013; Fraune et al., 2015; Murillo-Rincon et al., 2017).

  Note that carrying capacities of the members when part of the full community could not be assessed via plating of the community as was done for the mono-associations as not all bacterial species can be differentiated based on colony morphology, which is why we based calculations of the realized niche on data that was previously published (Franzenburg et al., 2013; Fraune et al., 2015; Murillo-Rincon et al., 2017). Data from another study (Deines et al., 2020) demonstrates that germ-free polyps, which were inoculated with the natural community of species (conventionalized animals), harbour an equally dense microbiome as wild-type polyps. This indicates that the generation and re-exposure of germ-free animals does not lead to an overall change in carrying capacity.

Cell surface hydrophobicity (CSH)
The BATH assay was performed as described previously (Borecká-Melkusová and Bujdáková, 2008; Rosenberg, 1984). It uses a biphasic separation method to measure

cell surface hydrophobicity. In short, for each species tested, 4 ml of bacterial
suspension (OD_{600} = 0.1; $OD_{initial}$) was placed into a class tube, overlaid with 1 ml of
n-hexadecane (Sigma Aldrich), and vortexed for 3 min. The phases were then allowed
to separate for 15 min, after which the ODs of the aqueous (lower) phase containing
hydrophilic cells was measured at OD_{600} ($OD_{residual}$). The hydrophobic cells are found
in the n-hexadecane overlay (upper phase). OD values were compared to the bacterial
suspension before mixing with n-hexadecane. The relative hydrophobicity (RH) was
calculated as follows: RH = (($OD_{initial} - OD_{residual}$)/$OD_{initial}$) x 100%. The experiment
was performed in triplicate with independent bacterial overnight cultures.
177
Biofilm quantification by use of crystal violet (CV)
Biofilm formation was assayed and quantified as previously described (Ren et al.,
2015). Briefly, exponential growth phase cultures of the six species were adjusted to
an optical density at 600nm (OD_{600}) of 0.1 in R2A medium. Biofilm formation was
assayed in a 96 well plate using eight replicates for each treatment. For single isolates
an inoculation volume of 180 µl was used and for the six-species biofilm 30 µl per
species. After 48 h of incubation at 18°C with shaking (200 r.p.m) biofilm formation
was quantified by a modified crystal violet (CV) assay (Peeters et al., 2008).
186
Characterizing nutrient utilisation
To characterise the nutrient profiles (niches) of *Hydras* microbiota, we measured the
carbon metabolism profile for each species using BIOLOG GN2 plates. BIOLOG
GN2 plates are 96-well microwell plates containing 95 different carbon sources plus a
carbon-absent water control well. Species were grown from isolate stocks in R2A
medium (18°C, shaken at 250 r.p.m.), centrifuged at 3000 r.c.f. for 5 min, re-
suspended in S medium and adjusted to an OD_{600} of 0.1. Each well of the BIOLOG
plate was inoculated with 150 µl of bacterial suspension and incubated for three days
at 18°C in a humid chamber. Growth on each of the 95 nutrients was determined as
OD_{600} of each well using a TECAN plate reader. For each plate, the OD of the water
control was subtracted from the reading of all other wells prior to analysis, and
differenced OD values below 0.005 were considered as no growth (Vaz Jauri et al.,
2013). Nutrient use was evaluated on three replicate plates. Nutrient (niche) overlap
(NO) was calculated using the formula: NO = (number of nutrients used by both A
and B)/((number of nutrients used by A + number of nutrients used by B)/ 2) (Vaz
Jauri et al., 2013). A value of 1 indicates the use of the same nutrients (100% overlap)
and 0 indicates no nutrient overlap among the 95 substrates tested. We also calculated
the relative use of the eleven functional groups (carbohydrates, carboxylic acids,
amino acids, polymers, aromatic chemicals, amines, amides, phosphorylated
chemicals, esters, alcohols and bromide chemicals) according to Daou et al. (2017). In
brief, the relative use of C substrates was calculated as absorption values in each well
divided by the total absorption in the plate.
209
Measurement of bacterial growth rates *in vitro*
Cultures of the all six species were produced in R2A microcosms (grown for 72 h at
18°C, at 250 r.p.m). Aliquots of each culture were first washed in S medium and then
re-suspended in fresh R2A medium to an optical density of 0.025 at 600 nm (OD_{600}).
Growth kinetics of all species were determined in 96-well microtiter plates. A 100 µl
aliquot of each re-suspension was pipetted into 100 µl of fresh R2A medium. The
microtiter plate was then placed in a microplate reader (TECAN Spark 10M, Tecan
Group Ltd., Switzerland), and the OD_{600} of each well was measured at 30 min

intervals for 96 cycles (with 10 sec shaking at 150 r.p.m. prior to each read). The
growth of each species was determined in five well locations on an individual 96-well
plate, which was replicated six times. The maximum growth rate (V_{max}) was
calculated from the maximum slope of the absorbance over time.
222
Statistical analysis
Analysis of variance (ANOVA) and subsequent post hoc Tukey-Kramer tests were
used to test for differences in the carrying capacity of the six *Hydra* colonizers. To
meet the requirements for the model, the variable was Box-Cox transformed. A
Welch ANOVA (and subsequent Wilcoxon posthoc tests) was used to test for
differences in the fraction of the different species in the community, differences in
biofilm formation capacity, and *in vitro* growth rates between the species.
Analysis of variance (ANOVA) and subsequent post hoc Tukey-Kramer tests were
used to test for differences in the cell surface hydrophobicity of the six *Hydra*
colonizers.
Sample size was chosen to maximise statistical power and ensure sufficient
replication. Assumptions of the tests, that is, normality and equal distribution of
variances, were visually evaluated. Non-significant interactions were removed from
the models. Effects were considered significant at the level of $P < 0.05$. All statistical
analyses were performed with JMP 9. Graphs were produced with GraphPad Prism
5.0, and RStudio (RStudio Team, 2015).
239

Results
Fundamental and realized niche occupation of the different bacterial species
In mono-colonisations, the six bacterial species differ significantly in their carrying capacity on *Hydra* (Figure 1A; ANOVA: $F_{5,12}$=12.696, P=0.0002). The most extreme cases are *Acidovorax sp.* that reaches the highest numbers with $2.6*10^5$ CFUs/polyp, and *Duganella sp.* the lowest with $1.7*10^4$ CFUs/polyp. Based in the carrying capacity of the single species in mono-colonisations, we estimated the fundamental niche of each of the six species. The realized niche of the six bacterial species is calculated based on previously published data on the composition of the extremely stable microbial community. The species differed in their relative abundance as part of the microbial community (Figure 1B; Welch ANOVA: $F_{5,14}$=86.722, P<0.0001). The most dominant species is *Curvibacter sp.* representing 65% of the microbial community, followed by *Duganella sp.* that reaches about 16%. All other four species reach only comparatively low fractions (around 1%), with *Acidovorax sp.* being the lowest. When comparing the fundamental to the realized niche (Figure 2), we find that the realized niche of *Curvibacter sp.* and *Duganella sp.* is larger than their fundamental niche. This is in contrast to all other species, where as expected by theory, competition with other microbes leads to a smaller realized than the fundamental niche.

Bacterial traits
Associated with occupation of the fundamental niche
The BATH assay was conducted with six species of the *Hydra* microbiome to measure cell surface hydrophobicity (CSH). The CSH of the bacterial species spanned a medium wide range; the values ranged from 0% to 42% and differs significantly between species (ANOVA: $F_{5,12}$=26.869; P<0.0001). *Curvibacter sp.* and *Pelomonas sp.* were the only two species that didn't show any affinity to the hexadecone; thus their cell population can be considered homogeneous consisting of only hydrophilic cells, which significantly differs from the others, except for *Pseudomonas sp.* (Figure 3A). *Pseudomonas sp.* and *Undibacterium sp.* show a mixed cell population, where 10 to 20% of the cells are hydrophobic. The species with the highest percentage of hydrophobic cells are *Acidovorax sp.* and *Duganella sp.*, between 30 and 35%.

The species differed in their biofilm formation (Welch ANOVA: $F_{5,18}$=350.723, P<0.0001). All species formed biofilms (Figure 3B), with *Pelomonas sp.* producing the largest biomass amount, which significantly differed from all other species. The biofilm amount of *Acidovorax sp.* was also significantly different from all other species but only roughly a third of the mass that *Pelomonas sp.* produced. All other species didn't differ and are comparatively weak biofilm producers.

Nutrient utilisation of all species was determined using a BIOLOG assay. The 94 carbon substrates are organized into eleven functional groups (Figure 4). Results showed that all six species actively oxidize carbon compounds such as carbohydrates (30-50% relative use), carboxylic acids (15-35% relative use) and amino acids (15-35% relative use) (Figure 4; Figure 5A). Carbohydrates are being used to an equal extent between all species, except for *Undibacterium sp.*, which uses the highest amount of around 50% (relative use). Turanose is the compound most highly utilised, followed by a-D-lactose, L-rhamnose, and D-cellobiose. The use of carboxylic acids increases in the species, which are characterized by low frequencies in the *Hydra* microbiome, with the exception of *Duganella sp.* (with a relative use of 25-30%). Here D-galactonic acid lactone is the substrate with the highest usage, followed by different forms of hydroxyl butyric acids. Amino acids are most excessively used by

290   *Acidovorax sp.*, *Curvibacter sp.* and *Pseudomonas sp.,* whereas the other species use
291   amino acids to a lesser extent. Polymers are being used very differently between the
292   species with *Pseudomonas sp.* showing the highest and *Undibacterium sp.* the lowest
293   values. Amines are being used more frequent by the dominant species in the
294   microbiome and are utilized to a lesser extent by the low abundant species.
295
Associated with occupation of the realized niche
When comparing the *in vitro* growth rates we find species perform differently (Figure
3C; Welch ANOVA: $F_{5,174}$=223.856, P<0.0001). The fastest species, *Undibacterium*
sp., grows twice as fast as compared to the slowest one, *Pelomonas sp.*.
The overlap in carbon substrate usage between all six microbiome members is
displayed as a Venn diagram (Figure 5B). There are only two substrates, which are
not utilized by any species, whereas 20 substrates are used by all species. There are
only two species that can metabolize substrates that none of the other species is using.
While *Pseudomonas sp.* uses two substrates: i-erythritol and lactulose, *Curvibacter*
sp. is able to utilise eight substrates: D-arabitol, D-mannose, D-trehalose, mono-
methyl succinate, formic acid, glucuronamide, L-pyroglutamic acid, and D-serine.
The number of substrates shared exclusively between two species only is between one
and two.
Niche overlap (NO) among all pairwise species combinations ranged from 60
to 80% (Figure 5C). *Curvibacter sp.* shares the highest overlap (80%) with
Pseudomonas sp. and *Duganella sp.*. For the other species the overlap ranges between
60 and 70%. *Duganella sp.* displays the highest overlap with *Pseudomonas sp.* and
Undibacterium sp. around 80%, whereas the overlap between *Pelomonas sp.* and
Acidovorax sp. reaches almost 70%. *Undibacterium sp.* exhibits an overlap of 60 to
70% with *Acidovorax sp.*, *Pseudomonas sp.* and *Pelomonas sp.*. *Acidovorax sp.* a
80% overlap with *Pseudomonas sp.* and 70% overlap with *Pelomonas sp.*.
Pseudomonas sp. and *Pelomonas sp.* show a 75% overlap of the nutrients used. Mean
niche overlap was determined as the mean of all pairwise niche overlap values for
each species. Comparing the number of nutrients being used by the individual species
we find that *Curvibacter sp.* and *Pseudomonas sp.* are able to use 70% of the provided
substrates. *Duganella sp.* uses 57%, *Acidovorax sp.* 54% and *Pelomonas sp.* 51%.
The lowest substrate utilization was measured for *Undibacterium sp.* with 41% of the
available substrates.
Overall, we find that neither the occupation of the fundamental or the realized
niche can be linked to a specific bacterial trait or substrate utilization pattern (Figure
6).
327

Discussion

Microbial communities residing in abiotic environments typically comprise numerous interacting species. Such communities have been studied with traditional approaches, for example the niche-assembly concept, which is an extension of the classical niche theory (Hutchinson, 1957). The niche-assembly perspective proposes that any ecosystem is made up of a limited number of niches, each occupied by a single species (Wennekes et al., 2012). Thus, the partitioning of these niches leads to the stable coexistence of competing species within an ecosystem. To assess the rules of assembly and coexistence of microbiota in host-associated microbiomes, we here apply the niche-assembly perspective to a metaorganism, and thus specifically extend the concept to biotic environments.

We find that the fundamental niche (here defined by the absence of interspecific microbial interactions) differs considerably from the realized niche of *Hydras* associated microbes (Figure 2). This reflects the difference in performance between the species when they individually occupy *Hydra* (mono-association) as to when they occur as part of their microbial community on the host. As predicted by niche theory, we find for the majority of the species that the realized niche is smaller than the fundamental one, most likely caused by interspecific microbial competition, as has also been observed in other systems, e.g. *Vibrios* in their marine environment (Materna et al., 2012). In our study, the best colonizer in the mono-colonisations, *Acidovorax sp.* (as also observed by Fraune et al. (2015)), is the least abundant species as part of the microbial community. This is different for the two main colonizers in the community, *Curvibacter sp.* and *Duganella sp.*, where the realized niche is six times the size of the fundamental niche for *Curvibacter sp.*, and about ten times the size for *Duganella sp.*. This finding is very interesting and indicates that the two species benefit from interactions when part of the microbiome. This can happen directly through positive interactions with the other members of the microbiome or indirectly by benefitting from the interactions between the other microbiome members and the host. We draw attention to the fact that the latter aspect differs from the classical Hutchinson niche concept, in that in our case the environment, i.e. the host, has the potential to change its interactions depending on the specific bacterial colonizers. Our finding also highlights the importance of the low frequency community members in shaping the overall community composition, as has recently been suggested for *Hydra* (Deines et al., 2020).

For linking the community composition in *Hydra's* microbiome to specific characteristics, we used a trait-based approach focussing on traits potentially involved in microbiome assembly and stability. A first step in microbiome assembly is the attachment to host surfaces, which can happen in a multitude of ways. In the human intestine, for example, microbes have been found to bind to mucin, a major component of the human mucosa (de Vos, 2015). Adhesion is thus thought to be a powerful mechanism for exerting both, positive and negative selection for or against specific microbes (McLoughlin et al., 2016; Schluter et al., 2015). Amongst others (van Loosdrecht et al., 1987), bacterial cell surface hydrophobicity has been shown to play a crucial role in surface attachment (Krasowska and Sigler, 2014). In general, hydrophobic cells adhere more strongly to hydrophobic surfaces and vice versa (Giaouris et al., 2009; Kochkodan et al., 2008). Nevertheless, the heterogeneity of a bacterial population needs to be taken into account. For example, the presence of both, hydrophilic and hydrophobic cells, have been observed in planktonic bacteria cell populations, implying that only part of the population participates in an adhesion process to substrates (Krasowska and Sigler, 2014). We also observe mixed cell

populations for most of *Hydra* microbial associates, except for two species, *Curvibacter sp.* and *Pelomonas sp.*, which only consist of hydrophobic cells. They seem to be perfectly adapted to *Hydras* epithelial cells, which are coated with a carbohydrate-rich layer, the glycocalyx (Ouwerkerk et al., 2013; Schröder and Bosch, 2016). The microbiome inhabits the outer mucus-like layer of the glycocalyx (Fraune et al., 2015), which is hydrophilic. Thus, hydrophilic bacterial cells should adhere more strongly to *Hydra* than hydrophobic cells. Both, *Curvibacter sp.* and *Pelomonas sp.*, have been shown to be of particular importance to the host. *Curvibacter sp.* shows signs of coevolution with its host and contributes to fungal resistance against the filamentous fungus *Fusarium* (Fraune et al., 2015). *Pelomonas sp.* has been shown to be of central importance in modulating the spontaneous body contractions in *Hydra* (Murillo-Rincon et al., 2017). So both species contribute to host fitness, providing the opportunity for the speculation that the host actively selects for specific microbes. This could happen, for example, by controlling the production and release of adhesive molecules from the host epithelium as suggested by (McLoughlin et al., 2016).

After successful attachment, bacteria need to colonise the habitat. In most cases, this happens through the formation of biofilms, as has been reported for the gut (de Vos, 2015; Kania et al., 2007). The biofilm succeeds the planktonic phase in the bacterial life cycle (McDougald et al., 2012) and represents a key ecological process for the colonization of different habitats. Thus, the difference in the ability to form biofilms could provide an explanation for why one species outcompetes the other species or has got a higher chance of persistence in the *Hydra* ecosystem. Further, biofilms have been shown to protect bacterial cells from various environmental stressors (Flemming and Wingender, 2010). Interestingly, from the six species tested here, the one with the highest ability to form biofilms is *Pelomonas sp.*, whereas the two main colonizers, *Curvibacter sp.* and *Duganella sp.* show a reduced capacity to form biofilms. Our finding indicates that the capability of biofilm formation is not a good predictor of the bacterial performance in the *Hydra* habitat. Nevertheless, it might be of importance for the establishment and persistence of some of the low abundance species, such as *Pelomonas sp.* and *Acidovorax sp.*.

Importantly, microbiomes on external surfaces of metaorganisms, such as the skin, have been reported to be highly stable despite their constant exposure to extrinsic factors (Oh et al., 2016). Whereas bacterial diversity is widely recognized in leading to temporal stability of ecosystem processes (Bell et al., 2009; Griffin et al., 2009; Prosser et al., 2007), the influence of resource niche breadth has received little scientific attention (Hunting et al., 2015). Recent work studying the decomposition of organic matter in experimental microcosms found that the higher the overlap in resource niches, the higher the stability of the microbial community. It is reasonable to assume that the same underlying principles govern stability in host-associated microbial communities. We therefore measured the niche overlap and resource use of the six species isolated from the *Hydra* microbiome. Interestingly, we find the niche overlap between all pairwise combinations to be between 60 and 80%, with about 20% of the carbon sources being metabolized by all species. This suggests that metabolic overlap could be involved in promoting the extreme temporal stability of *Hydra's* microbiome (Fraune and Bosch, 2007). We also found the two main colonisers, *Curvibacter sp.* and *Duganella sp.*, to possess the widest resource niche breadth of all species, and that five out of six species were able to metabolize more than 50% of the 95 offered carbon substrates. Overall, the relative niche breadth observed in the tested species can serve as a proxy of the metabolic diversity of the *Hydra* microbiome.

The metabolic overlap, i.e. redundancy, within *Hydra's* microbial community
indicates that the individual species are not occupying a specific metabolic niche.
Nevertheless, the only one for which we observed a specific carbon usage pattern is
the main colonizer *Curvibacter sp.*, which utilizes eight carbon sources that are not
metabolized by any other tested microbiome members. Whether this hints at the
occupation of a specific niche within *Hydra's* microbial community and can be linked
to the observation that its realized is bigger than its fundamental niche when part of
the community as compared to single occupation on *Hydra*, is currently open to
speculation. An alternative option might be that *Curvibacter sp.* is auxotrophic in
producing certain amino acids, as are 98% of all sequenced microbes (Zengler and
Zaramela, 2018), and thus relies on the uptake of external substrates that might not be
secreted by the host but by its fellow community members. Analysing the metabolic
interactions within this microbial network will be essential for understanding
community assembly, composition, and maintenance.
In summary we find that the here measured bacterial traits vary across
microbiome members. Further, the dominant species in the microbiome do not
necessarily perform best in all of the measured traits. We rather observe that all
species, independent of their density, perform well in a subset of traits, likely
facilitating the coexistence of several niches within the host ecosystem. Whether a
change in the realized niche of microbes can be linked to potential for dysbiosis is an
interesting aspect, which warrants further investigation.
449

Ethics statement

Ethical restrictions do not apply to cnidarian model organisms such as *Hydra*.

Author contributions

PD and KH designed the experiments. PD performed the experiments. PD and KH analysed the data. PD, KH and TB wrote the paper.

Funding

PD received funding from the European Union's Framework Programme for Research and Innovation Horizon 2020 (2014–2020) under the Marie Skłodowska-Curie Grant Agreement No. 655914 and KH under the Marie Skłodowska-Curie Grant Agreement No. 657096. Both also received a Reintegration Grant from the Deutscher Akademischer Austausch Dienst (DAAD). This work was further supported by the Deutsche Forschungsgemeinschaft (DFG) Collaborative Research Centre (CRC) 1182 ("Origin and Function of Metaorganisms").

Acknowledgements

TB gratefully appreciates support from the Canadian Institute for Advanced Research (CIFAR) and thanks the Wissenschaftskolleg (Institute of Advanced Studies) in Berlin for a sabbatical leave.

Competing Interests

The authors declare no conflict of interest.

References

Bell, T., Gessner, M. O., Griffiths, R. I., McLaren, J. R., Morin, P. J., and van der Heijden, M. (2009). "Microbial diversity and ecosystem functioning under controlled conditions and in the wild," in *Biodiversity, Ecosystem Functioning, and Human Wellbeing*, eds. S. Naeem, D. E. Bunker, A. Hector, M. Loreau, and C. P. Perring (Oxford, UK), 121–133.

Borecká-Melkusová, S., and Bujdáková, H. (2008). Variation of cell surface hydrophobicity and biofilm formation among genotypes of *Candida albicans* and *Candida dubliniensis* under antifungal treatment. *Can.J. Microbiol.* 54, 718–724. doi:10.1139/w08-060.

Bosch, T. C. G. (2013). Cnidarian-microbe interactions and the origin of innate immunity in metazoans. *Annu. Rev. Microbiol.* 67, 499–518. doi:10.1146/annurev-micro-092412-155626.

Daou, L., Luglia, M., Périssol, C., Calvert, V., and Criquet, S. (2017). Sporulation and physiological profiles of bacterial communities of three Mediterranean soils affected by drying-rewetting or freezing-thawing cycles. *Soil Biol. Biochem.* 113, 116–121. doi:10.1016/j.soilbio.2017.06.008.

de Vos, W. M. (2015). Microbial biofilms and the human intestinal microbiome. *NPJ Biofilms Microbiomes* 1, 15005. doi:10.1038/npjbiofilms.2015.5.

Deines, P., and Bosch, T. C. G. (2016). Transitioning from microbiome composition to microbial community interactions: the potential of the metaorganism *Hydra* as an experimental model. *Front. Microbiol.* 7, 1610. doi:10.3389/fmicb.2016.01610.

Deines, P., Hammerschmidt, K., Bosch, T. C. B. (2020). Microbial species coexistence depends on the host environment. *bioRxiv https://doi.org/10.1101/609271*

Deines, P., Lachnit, T., and Bosch, T. C. G. (2017). Competing forces maintain the *Hydra* metaorganism. *Immunol. Rev.* 279, 123–136. doi:10.1111/imr.12564.

Falkowski, P. G., Fenchel, T., and Delong, E. F. (2008). The microbial engines that drive Earth's biogeochemical cycles. *Science* 320, 1034–1039. doi:10.1126/science.1153213.

Flemming, H. C., and Wingender, J. (2010). The biofilm matrix. *Nat. Rev. Microbiol.* 8, 623–633. doi:10.1038/nrmicro2415.

Franzenburg, S., Walter, J., Künzel, S., Wang, J., Baines, J. F., Bosch, T. C. G., et al. (2013). Distinct antimicrobial peptide expression determines host species-specific bacterial associations. *Proc. Natl. Acad. Sci. U.S.A.* 110, E3730–8. doi:10.1073/pnas.1304960110.

Fraune, S., and Bosch, T. C. G. (2007). Long-term maintenance of species-specific bacterial microbiota in the basal metazoan *Hydra. Proc. Natl. Acad. Sci. U.S.A* 104, 13146–13151. doi:10.1063/1.1599624.

Fraune, S., Anton-Erxleben, F., Augustin, R., Franzenburg, S., Knop, M., Schröder,
K., et al. (2015). Bacteria-bacteria interactions within the microbiota of the
ancestral metazoan *Hydra* contribute to fungal resistance. *ISME J.* 9, 1543–1556.
doi:10.1038/ismej.2014.239.

Giaouris, E., Chapot-Chartier, M.-P., and Briandet, R. (2009). Surface
physicochemical analysis of natural *Lactococcus lactis* strains reveals the
existence of hydrophobic and low charged strains with altered adhesive
properties. *Int. J. Food Microbiol.* 131, 2–9.
doi:10.1016/j.ijfoodmicro.2008.09.006.

Green, J. L., Bohannan, B. J. M., and Whitaker, R. J. (2008). Microbial biogeography:
from taxonomy to traits. *Science* 320, 1039–1043. doi:10.1126/science.1153475.

Griffin, J. N., O'Gorman, E. J., Emmerson, M. C., Jenkins, S. R., Klein, A. M., and
Loreau, M. (2009). "Biodiversity and the stability of ecosystem functioning," in
Biodiversity, Ecosystem Functioning, and Human Wellbeing, eds. S. Naeem, D.
E. Bunker, A. Hector, M. Loreau, and C. P. Perring (Oxford, UK: Oxford
University Press), 78–93.

Guittar, J., Shade, A., and Litchman, E. (2019). Trait-based community assembly and
succession of the infant gut microbiome. *Nat. Commun.* 10, 512.
doi:10.1038/s41467-019-08377-w.

Holt, R. D. (2009). Bringing the Hutchinsonian niche into the 21st century: ecological
and evolutionary perspectives. *Proc. Natl. Acad. Sci. U.S.A.* 106, 19659–19665.
doi:10.1073/pnas.0905137106.

Hunting, E. R., Vijver, M. G., van der Geest, H. G., Mulder, C., Kraak, M. H. S.,
Breure, A. M., et al. (2015). Resource niche overlap promotes stability of
bacterial community metabolism in experimental microcosms. *Front. Microbiol.*
6, 11512. doi:10.3389/fmicb.2015.00105.

Hutchinson, G. E. (1957). Concluding remarks. *Cold Spring Harb. Sym.* 22, 415–427.

Kania, R. E., Lamers, G. E. M., Vonk, M. J., Huy, P. T. B., Hiemstra, P. S.,
Bloemberg, G. V., et al. (2007). Demonstration of bacterial cells and glycocalyx
in biofilms on human tonsils. *Arch. Otolaryngol.* 133, 115–121.
doi:10.1001/archotol.133.2.115.

Kau, A. L., Ahern, P. P., Griffin, N. W., Goodman, A. L., and Gordon, J. I. (2011).
Human nutrition, the gut microbiome and the immune system. *Nature* 474, 327–
336. doi:10.1038/nature10213.

Kochkodan, V., Tsarenko, S., Potapchenko, N., Kosinova, V., and Goncharuk, V.
(2008). Adhesion of microorganisms to polymer membranes: a photobactericidal
effect of surface treatment with TiO_2. *Desalination* 220, 380–385.
doi:10.1016/j.desal.2007.01.042.

Kopac, S. M., and Klassen, J. L. (2016). Can they make it on their own? Hosts,
microbes, and the holobiont niche. *Front. Microbiol.* 7, 1647.
doi:10.3389/fmicb.2016.01647.

Krasowska, A., and Sigler, K. (2014). How microorganisms use hydrophobicity and
what does this mean for human needs? *Front. Cell. Infect. Microbiol.* 4, 653.
doi:10.3389/fcimb.2014.00112.

Krause, S., Le Roux, X., Niklaus, P. A., Van Bodegom, P. M., Lennon, J. T.,
Bertilsson, S., et al. (2014). Trait-based approaches for understanding microbial
biodiversity and ecosystem functioning. *Front. Microbiol.* 5, 2019.
doi:10.3389/fmicb.2014.00251.

Leibold, M. A. (1995). The niche concept revisited: mechanistic models and
community context. *Ecology* 76, 1371–1382. doi:10.2307/1938141.

Lenhoff, H. M., and Brown, R. D. (1970). Mass culture of *Hydra*: an improved
method and its application to other aquatic invertebrates. *Lab. Anim.* 4, 139–154.
doi:10.1258/002367770781036463.

Louca, S., Polz, M. F., Mazel, F., Albright, M. B. N., Huber, J. A., O'Connor, M. I.,
et al. (2018). Function and functional redundancy in microbial systems. *Nat. Ecol.*
Evol. 2, 936–943. doi:10.1038/s41559-018-0519-1.

Martiny, J. B. H., Jones, S. E., Lennon, J. T., and Martiny, A. C. (2015). Microbiomes
in light of traits: a phylogenetic perspective. *Science* 350, aac9323–aac9323.
doi:10.1126/science.aac9323.

Materna, A. C., Friedman, J., Bauer, C., David, C., Chen, S., Huang, I. B., et al.
(2012). Shape and evolution of the fundamental niche in marine *Vibrio*. *ISME J.*
6, 2168–2177. doi:10.1038/ismej.2012.65.

McDougald, D., Rice, S. A., Barraud, N., Steinberg, P. D., and Kjelleberg, S. (2012).
Should we stay or should we go: mechanisms and ecological consequences for
biofilm dispersal. *Nat. Rev. Microbiol.* 10, 39–50. doi:10.1038/nrmicro2695.

McFall-Ngai, M., Hadfield, M. G., Bosch, T. C. G., Carey, H. V., Domazet-Loso, T.,
Douglas, A. E., et al. (2013). Animals in a bacterial world, a new imperative for
the life sciences. *Proc. Natl. Acad. Sci. U.S.A.* 110, 3229–3236.
doi:10.1073/pnas.1218525110.

McLoughlin, K., Schluter, J., Rakoff-Nahoum, S., Smith, A. L., and Foster, K. R.
(2016). Host selection of microbiota via differential adhesion. *Cell Host Microbe*
19, 550–559. doi:10.1016/j.chom.2016.02.021.

Murillo-Rincon, A. P., Klimovich, A., Pemöller, E., Taubenheim, J., Mortzfeld, B.,
Augustin, R., et al. (2017). Spontaneous body contractions are modulated by the
microbiome of *Hydra*. *Sci. Rep.* 7, 15937. doi:10.1038/s41598-017-16191-x.

Oh, J., Byrd, A. L., Park, M., NISC Comparative Sequencing Program, Kong, H. H.,
and Segre, J. A. (2016). Temporal stability of the human skin microbiome. *Cell*
165, 854–866. doi:10.1016/j.cell.2016.04.008.

Ouwerkerk, J. P., de Vos, W. M., and Belzer, C. (2013). Glycobiome: bacteria and
mucus at the epithelial interface. *Best Pract. Res. Cl. Ga.* 27, 25–38.
doi:10.1016/j.bpg.2013.03.001.

Pearman, P. B., Guisan, A., Broennimann, O., and Randin, C. F. (2008). Niche
dynamics in space and time. *Trends Ecol. Evol.* 23, 149–158.
doi:10.1016/j.tree.2007.11.005.

Peeters, E., Nelis, H. J., and Coenye, T. (2008). Comparison of multiple methods for
quantification of microbial biofilms grown in microtiter plates. *J. Microbiol.*
Meth. 72, 157–165. doi:10.1016/j.mimet.2007.11.010.

Pocheville, A. (2015). "The ecological niche: history and recent controversies," in
Handbook of Evolutionary Thinking in the Sciences, eds. T. Heams, P. Huneman,
G. Lecointre, and M. Silberstein (Dordrecht: Springer, Dordrecht), 547–586.
doi:10.1007/978-94-017-9014-7_26.

Prosser, J. I., Bohannan, B. J. M., Curtis, T. P., Ellis, R. J., Firestone, M. K.,
Freckleton, R. P., et al. (2007). The role of ecological theory in microbial
ecology. *Nat. Rev. Microbiol.* 5, 384–392. doi:10.1038/nrmicro1643.

Ren, D., Madsen, J. S., Sørensen, S. J., and Burmølle, M. (2015). High prevalence of
biofilm synergy among bacterial soil isolates in cocultures indicates bacterial
interspecific cooperation. *ISME J.* 9, 81–89. doi:10.1038/ismej.2014.96.

Rosenberg, M. (1984). Bacterial adherence to hydrocarbons: a useful technique for
studying cell surface hydrophobicity. *FEMS Microbiol Lett* 22, 289–295.
doi:10.1111/j.1574-6968.1984.tb00743.

RStudio Team (2015). *RStudio: Integrated Development for R (RStudio, Inc., Boston,*
MA). Available at: Available at https://www.rstudio.com.

Schluter, J., Nadell, C. D., Bassler, B. L., and Foster, K. R. (2015). Adhesion as a
weapon in microbial competition. *ISME J.* 9, 139–149.
doi:10.1038/ismej.2014.174.

Schröder, K., and Bosch, T. C. G. (2016). The origin of mucosal immunity: lessons
from the holobiont *Hydra*. *mBio* 7, e01184–16. doi:10.1128/mBio.01184-16.

van Loosdrecht, M. C., Lyklema, J., Norde, W., Schraa, G., and Zehnder, A. J.
(1987). The role of bacterial cell wall hydrophobicity in adhesion. *Appl. Environ.*
Microb. 53, 1893–1897.

Vaz Jauri, P., Bakker, M. G., Salomon, C. E., and Kinkel, L. L. (2013). Subinhibitory
antibiotic concentrations mediate nutrient use and competition among soil
streptomyces. *PLoS ONE* 8, e81064. doi:10.1371/journal.pone.0081064.

Wein, T., Dagan, T., Fraune, S., Bosch, T. C. G., Reusch, T. B. H., and Hülter, N. F.
(2018). Carrying capacity and colonization dynamics of *Curvibacter* in the *Hydra*
host habitat. *Front. Microbiol.* 9, 185. doi:10.3389/fmicb.2018.00443.

Wennekes, P. L., Rosindell, J., and Etienne, R. S. (2012). The neutral-niche debate: a
philosophical perspective. *Acta Biotheor.* 60, 257–271. doi:10.1007/s10441-012-
9144-6.

Whittaker, R. H., Levin, S. A., and Root, R. B. (1973). Niche, habitat, and ecotope.

Am. Nat. 107, 321–338. doi:10.1086/282837;subPage:string:Access.

Zengler, K., and Zaramela, L. S. (2018). The social network of microorganisms - how
auxotrophies shape complex communities. *Nat. Rev. Microbiol.* 16, 383–390.
doi:10.1038/s41579-018-0004-5.

638
639

 Figures

641
Figure 1. Performance of the six main bacterial colonizers isolated from the *Hydra*
microbiome. (**A**) Carrying capacity of the *Hydra* ecosystem during mono-associations
of germ-free polyps with individual bacterial species. Error bars are s.e.m., based on
n=3. (**B**) Relative abundances of the different bacterial colonizers in wild-type (open
circles) and conventionalized polys (filled circles), compiled from previously
published studies (left: (Murillo-Rincon et al., 2017), right: (Franzenburg et al.,
2013)).
649
650
651
652

653
654 **Figure 2.** *Hydra* functions as an ecosystem, which allows for the niche allocation.
655 Shown are the fundamental and realized niches of six microbiome members (based on
656 mono-colonisations and microbial community composition). The realized niche
657 includes additional constrains arising from inter-specific competition between
658 microbiome members.

659
660 **Figure 3.** Trait measures of bacterial species isolated from the *Hydra* microbiome.
661 (**A**) Cell surface hydrophobicity (Error bars are s.e.m., n=3), and (**B**) Biofilm
662 formation capacity of six bacterial isolates. Error bars are s.e.m., n=4. (**C**) Bacterial
663 growth rates of individual species measured *in vitro*. Error bars are s.e.m., based on
664 n=30.

665
666 **Figure 4.** Substrate utilization pattern of six bacterial isolates from the *Hydra*
667 microbiome measured with a BIOLOG assay. Colours indicate the relative magnitude
668 of substrate utilization.

669
Figure 5. Metabolic similarity between *Hydra's* microbiome members. (**A**) Number
of substrates utilized and their relative use (%). (**B**) Venn diagram showing the
distribution of shared substrates among the microbiome members. (**C**) Niche overlap
among all pairwise combinations of six *Hydra* microbiome members. A value of 1
indicates the use of the same nutrients (100% overlap) and 0 indicates no nutrient
overlap.
676

677
Figure 6. Association of traits with the occupation of the fundamental and realized
niches in *Hydra's* six microbiome members.
680