A sparse observation model to quantify species interactions in time and space

Sadoune. Ait Kaci Azzou[a,b,*], Liam. Singer[a,b,*], Thierry. Aebischer[a,b], Beat. Wolf[c], Daniel. Wegmann[a,b,**]

[a]*Department of Biology, Université de Fribourg, Chemin du Musée 10, CH-1700 Fribourg, Switzerland*
[b]*Swiss Institute of Bioinformatics, 1700 Fribourg, Switzerland*
[c]*iCoSys, University of Applied Sciences Western Switzerland, Fribourg, Switzerland*

Summary

1. Camera traps and acoustic recording devices are essential tools to quantify the distribution, abundance and behavior of mobile species. Varying detection probabilities among device locations must be accounted for when analyzing such data, which is generally done using occupancy models.

2. We introduce a Bayesian Time-dependent Observation Model for Camera Trap data (`Tomcat`), suited to estimate relative event densities in space and time. `Tomcat` allows to learn about the environmental requirements and daily activity patterns of species while accounting for imperfect detection. It further implements a sparse model that deals well will a large number of potentially highly correlated environmental variables.

3. By integrating both spatial and temporal information, we extend the notation of overlap coefficient between species to time and space to study niche partitioning.

4. *Synthesis and applications.* We illustrate the power of `Tomcat` through an application to camera trap data of eight sympatrically occurring duiker *Cephalophinae* species in the the savanna - rainforest ecotone in the Central African Republic and show that most species pairs show little overlap. Exceptions are those for which one species is very rare, likely as a result of direct competition.

Keywords: Biodiversity monitoring, Camera trap data, Species distribution, Niche partitioning, Daily activity, Sparsity, Overlap coefficient

---

[*]These authors contributed equally
[**]Corresponding author
Email address: `daniel.wegmann@unifr.ch` (Daniel. Wegmann)

1. Introduction

Camera traps, acoustic recorders and other devices that allow for continuous recording of animal observations have become an essential part of many wildlife monitoring efforts that aim at quantifying the distribution, abundance and behavior of mobile species. However, the inference of these biological characteristics is not trivial due to the confounding factor of detection, which may vary greatly among recording locations. Hence, variation in the rates at which a species is recorded (e.g. the photographic rate) may indeed reflect differences in local abundance, but might just as well reflect differences in the probabilities with which individuals are detected, or more likely a combination of both (see Burton et al., 2015; Sollmann, 2018, for two excellent reviews).

Since local detection rates are generally not known, both processes have to be inferred jointly. The most often used methods are variants of occupancy models that treat the detection probability explicitly (MacKenzie et al., 2002). The basic quantity of interest in these models is whether or not a particular site is occupied by the focal species. While the detection of a species implies that it is present, the absence of a record does not necessarily imply the species is absent. Since the probabilities of detection and occupation are confounded, they can not be inferred for each site individually. It is therefore common to use hierarchical models that express detection probabilities as a function of environmental variables (MacKenzie et al., 2002).

Here we introduce `Tomcat`, a Time-dependent Observation Model for Camera Trap data, that extends currently used occupancy models in three important ways:

First, we propose to quantify a measure of relative species density, namely the rate at which animals pass through a specific location, rather than occupancy. Quantifying occupancy assumes there exists a well defined patch or site that is either occupied by a species or not (closure assumption). The notation of a discrete patch is, however, often difficult when analyzing camera trap or similar data of mobile species, which complicates interpretation (Efford and Dawson, 2012; Steenweg et al., 2018). In addition, summarizing such data by a simple presence-absence matrix ignores the information about differences in population densities at occupied sites. Occupancy is therefore not necessarily a good surrogate for abundance (Efford and Dawson, 2012; Steenweg et al., 2018; MacKenzie and Royle, 2005),

even though it has been advocated for birds (MacKenzie and Nichols, 2004).

By quantifying relative densities as an alternative measure to occupancy, this limitation can be overcome as we do not need to make strict assumptions about the independence of recording locations. Furthermore, `Tomcat` quantifies measures of relative species densities while accounting for imperfect detection of unmarked species, which has previously been identified as a key challenge in wildlife surveys (Burton et al., 2015). However, we note that relative densities do not easily allow for an absolute quantification of density because it is not possible to distinguish mobility from abundance. On the other hand, they readily allow for the comparison of densities in space and time and hence to identify habitat important for a particular species. It therefore appears more useful than occupancy to monitor changes in species abundances over time, as for many species, changes in population size will be reflected in the rate at which a species is detected prior to local extinction.

Second, we explicitly model daily activity patterns. Several models have been proposed to estimate such patterns from continuous recording data (Frey et al., 2017), including testing for non-random distributions of observation events in predefined time-bins (Bu et al., 2016) and circular kernel density functions (Oliveira-santos et al., 2013; Rowcliffe et al., 2014), with latter allowing for the quantification of activity overlap between species (Ridout and Linkie, 2009). Jointly inferring activity patterns with relative densities allows us not only account for imperfect detection, but also extent the idea of overlap to space, shedding additional light on species interactions.

Third, we explicitly account for the sparsity among environmental coefficients. This is relevant since many environmental variables are generally available and it is usually not known which ones best explain the variation in abundance of a species. Enforcing sparsity on the vector of coefficients avoids the problem of over-fitting in case the number of recording locations is smaller or on the same order as the number of environmental coefficients.

In this article, we begin by describing the proposed model in great details. We then verify its performance using extensive simulations and finally illustrate this idea by inferring spatio-temporal overlap of eight duiker species *Cephalophinae* within the forest-savanna ecotone of Central Africa.

2. Material and Methods

2.1. A Time-dependent Observation Model

We present a Bayesian Time-dependent Observation Model for Camera Trap data (`Tomcat`), suited to estimate relative event densities in space and time. Let us denote by $\Lambda_j(\tau)$ the rate at which a device at location $j = 1, \ldots, J$ takes observations of a particular species (or guild) at the time of the day $\tau \in [0, T], T = 24h$. We assume that this rate is affected by three processes: 1) the average rate $\overline{\lambda_j}$ at which individuals pass through location j, 2) the daily activity patterns $\mathcal{T}(\tau)$, and 3) the probability p_j with which an individual passing through location j is detected by the device and the downstream processing of the records:

$$\Lambda_j(\tau) = p_j \overline{\lambda_j} \mathcal{T}(\tau), \quad \int_0^T \mathcal{T}(\tau) d\tau = T.$$

The number of records $W_j(d, \tau_1, \tau_2)$ taken by a device at location j within the interval $[\tau_1, \tau_2)$ on day d is then given by the non-homogeneous Poisson process

$$W_j(d, \tau_1, \tau_2) \sim \text{Poisson}(\Lambda_j(\tau_1, \tau_2)),$$

with intensity function

$$\Lambda_j(\tau_1, \tau_2) = p_j \overline{\lambda_j} \int_{\tau_1}^{\tau_2} \mathcal{T}(\tau) d\tau. \tag{1}$$

As in all models dealing with imperfect detection, the parameters $\overline{\lambda_j}$ and p_j related to species densities and detection, respectively, are confounded and can not be estimated individually for each location without extra information. However, it is possible to estimate relative densities between locations using a hierarchical model. Following others (e.g. Tobler et al., 2015), we assume that both parameters are functions of covariates (e.g. the environment), and hence only attempt to learn these hierarchical parameters. Here, we use

$$\log(\overline{\lambda_j}) = a + X_j A, \tag{2}$$

$$\text{logit}(p_j) = Y_j B, \tag{3}$$

where X_j and Y_j are known (environmental) covariates at location j and a, A and B are species specific coefficients. Note that to avoid non-identifiability issues, we did not include

78 an intercept for p_j, and hence we set the average detection probability across locations
79 $\frac{1}{J} \sum_j^J p_j = 0.5$ (assuming X and Y have mean zero). Also, X_j and Y_j should not contain
80 strongly correlated covariates.

2.1.1. Non-independent events

Another issue specific to data from continuous recordings is that not every record is
necessarily reflective of an independent observation as the same individual might trigger
multiple observations while passing (or feeding, resting, ...) in front of a recording device. It
is often difficult and certainly laborious to identify such recurrent events. We thus propose
to account for non-independent events by dividing the day into n_o intervals of equal length
$h_o = \frac{T}{n_o}$ (o for observation), and then only consider whether or not at least one recording was
taken within each interval $[c_{m-1}, c_m), m = 1, \ldots, n_o$, where $c_0 = c_M = T$. Specifically, for an
interval m,

$$\mathbb{P}(W_j(d, c_{m-1}, c_m) = w) = \begin{cases} e^{-\Lambda_j(c_{m-1}, c_m)} & \text{if } w = 0 \\ 1 - e^{-\Lambda_j(c_{m-1}, c_m)} & \text{if } w > 0 \end{cases}, \tag{4}$$

90 where $\Lambda_j(c_{m-1}, c_m)$ is given by (1).

2.1.2. Daily activity patterns

Here we assume that $\mathcal{T}(\tau)$ is a piece-wise constant function with n_a activity intervals of
equal length $h_a = \frac{T}{n_a}$ (a for activity). While activity patterns are unlikely strictly piece-wise
constant, we chose this function over a combination of periodic functions (e.g. Oliveira-santos
et al., 2013) as they fit to complicated, multi-peaked distributions with fewer parameters.
Since the best tiling of the day is unknown, we allow for a species-specific shift δ such
that the first interval is $[\delta, h_a + \delta)$ and the last overlaps midnight and becomes $[T - h_a + \delta, \delta)$
(Figure 1). We therefore have

$$\mathcal{T}(\tau) = \sum_{i=1}^{n_{h_a}} k_i \mathbb{1}_{[(i-1)h_a, ih_a)}(\tau - \delta), \tag{5}$$

99 where the indicator function $\mathbb{1}_{[\tau_0, \tau_1)}(\tau)$ is 1 if $t \in [\tau_0, \tau_1)$ and zero otherwise and k_i reflects
100 the relative activity of the focal species (or guild) in interval $i = 1, \ldots, n_h$ with $k_i = 0$
101 implying no activity and $k_i = 1$ implying average activity. Note that

$$\int_0^T \mathcal{T}(\tau)d\tau = h_a \sum_{l=1}^{n_a} k_l = T,$$

and hence

$$\frac{1}{n_a} \sum_{l=1}^{n_a} k_l = 1.$$

102 *2.1.3. Bayesian inference*

103 We conduct Bayesian inference on the parameter vector $\boldsymbol{\theta} = \{a, A, B, \delta, \boldsymbol{k}\}$, where $\boldsymbol{k} =$
104 $\{k_1, \ldots, k_{n_a}\}$, by numerically evaluating the posterior distribution $\mathbb{P}(\boldsymbol{\theta}|\boldsymbol{W}) \propto \mathbb{P}(\boldsymbol{W}|\boldsymbol{\theta})\mathbb{P}(\boldsymbol{\theta})$,
105 where $\boldsymbol{W} = \{W_1, \ldots, W_J\}$ denotes the full data from all locations $j = 1, \ldots, J$.

106 The likelihood $\mathbb{P}(\boldsymbol{W}|\theta)$ is calculated as

$$\mathbb{P}(\boldsymbol{W}|\boldsymbol{\theta}) = \prod_{j=1}^J \prod_{d=D_{j1}}^{D_{j2}} \prod_{m=1}^M \mathbb{P}(W_j(d, c_{m-1}, c_m)|\boldsymbol{\theta}),$$

where $\mathbb{P}(W_j(d, c_{m-1}, c_m)|\boldsymbol{\theta})$ is given by equation (4) and D_{j1} and D_{j2} denote the first and
last day of recording at location j.

Since it is usually not known which covariates X_j and Y_j are informative, nor at which
spatial scale they should be evaluated, the potential number of covariates to be considered
may be large. To render inference feasible, we enforce sparsity on the vectors of coefficients
A and B. Specifically, we assume that $\mathbb{P}(A_i \neq 0) = \pi_\lambda$ and, correspondingly, $\mathbb{P}(B_i \neq 0) = \pi_p$.

We chose uniform priors on all other parameters, namely $\mathbb{P}(a) \propto 1$, $\mathbb{P}(\boldsymbol{k}) \propto 1$ for all
vectors of $\boldsymbol{k}$ that satisfy $\sum_{l=1}^{n_h} k_l = n_h$, and $\mathbb{P}(\delta) \propto 1$ for all $0 \leq \delta < h_a$. For simplicity, we
only consider cases in which h_a, the length of the activity intervals, is a multiple of h_o, the
length of the observation intervals, and allow only for discrete $\delta \in \{0, \ldots, h_a/h_o\}$. Finally,
we set $\pi_p = \pi_\lambda = 0.1$.

We use a reversible-jump MCMC algorithm (Green, 1995) to generate samples from the
posterior distribution $\mathbb{P}(\boldsymbol{\theta}|\boldsymbol{W})$. The update $\boldsymbol{k} \to \boldsymbol{k}'$ is noteworthy. We begin by picking a
random activity interval i and proposing a move $k_i \to k_i'$ according to a symmetric transition
kernel. We then scale all other entries of $\boldsymbol{k}'$ to satisfy the constraint on the sum. Specifically,
we set $k_j' = \alpha k_j$ for all $j \neq i$, where

$$\alpha = \frac{n - k_i'}{\sum_{j \neq i} k_j}.$$

 2.1.4. Prediction

124   Using a set of S posterior samples $\boldsymbol{\theta}_1, \ldots, \boldsymbol{\theta}_S \sim \mathbb{P}(\boldsymbol{\theta}|\boldsymbol{W})$, we project event densities to
125 a not-surveyed location ι with covariates X_ι by calculating the mean $\hat{\overline{\lambda}}_\iota$ of the posterior
126 $\mathbb{P}(\overline{\lambda}_\iota|X_\iota, \boldsymbol{W})$ as

$$\hat{\overline{\lambda}}_\iota \approx \frac{1}{S} \sum_{s=1}^{S} e^{a^{(s)} + X_\iota A^{(s)}}, \tag{6}$$

where $a^{(i)}$ and $A^{(i)}$ denote the i-th posterior sample of these parameters.

2.1.5. Species overlap in space and time

An important interest in ecology is to compare activity patterns among species and to
see how overlapping patterns may relate to competition or predation (e.g. Ridout and Linkie,
2009; Rowcliffe et al., 2014).

We can quantify overlapping patterns of animal activity by estimating the coefficient of
overlap Δ (Ridout and Linkie, 2009). This quantitative measure ranges from 0 (no overlap)
to 1 (identical activity patterns) and is the area lying under two activity density curves (see
Figure 4). For two known density functions $f(x)$ and $g(x)$, Δ is given by:

$$\Delta(f, g) = \int \min\{f(x), g(x)\} \tag{7}$$

136   The overlap measure $\Delta(f, g)$ can be related to the well known measure of distance between
137 two densities L_1 as

$$\Delta(f, g) = 1 - \frac{1}{2} \int |f(x) - g(x)| dx, \tag{8}$$

which justifies the visualization of overlap coefficients between k species in a n-dimensional
space using a Multidimensional Scaling (MDS) by considering $1 - \hat{\Delta}(f, g)$ as a measure of
dissimilarity.

In practice, the true density functions $f(x)$ and $g(x)$ are usually not known. Here we
obtain an estimate of Δ numerically from posterior samples. We distinguish three types of
overlap coefficients, Δ_T for overlap in time, Δ_S for overlap in space and Δ_{ST} for overlap in
time and space.

Overland coefficient $\mathbf{\Delta_T}$. For a large number n_T of equally spaced time values $\tau_1, \tau_2, \ldots, \tau_{n_T} \in$ $(0, T)$, we sample $\Delta_T^{(m)}$ from the posterior distribution $\mathbb{P}(\Delta_T | \boldsymbol{W}^{(1)}, \boldsymbol{W}^{(2)})$ where $\boldsymbol{W}^{(l)} = \{W_1^{(l)}, \cdots, W_J^{(l)}\}$ denotes the full data for a species $l = 1, 2$.

$$\widehat{\Delta}_T^{(m)} = \frac{1}{n_T} \sum_{i=1}^{n_T} \min \left\{ \mathcal{T}_1^{(m)}(\tau_i), \mathcal{T}_2^{(m)}(\tau_i) \right\}, \tag{9}$$

where $\mathcal{T}_l^{(m)}$ is computed according to equation (5) with species specific parameters $\delta_l^{(m)}$ and $\boldsymbol{k}_l^{(m)} = (k_{1,l}^{(m)}, k_{2,l}^{(m)}, \cdots, k_{n_{h_a},l}^{(m)})$ sampled from $\mathbb{P}(\boldsymbol{\theta}_l | \boldsymbol{W}^{(l)})$.

Overlap coefficient $\mathbf{\Delta_S}$. For a given number n_S of sites reflecting the habitat in a region, we sample $\Delta_S^{(s)}$ from the posterior distribution $\mathbb{P}(\Delta_S | \boldsymbol{W}^{(1)}, \boldsymbol{W}^{(2)})$ as

$$\widehat{\Delta}_S^{(s)} = \frac{1}{n_S} \sum_{j=1}^{n_S} \min \left\{ \overline{\lambda}_{1j}^{(s)}, \overline{\lambda}_{2j}^{(s)} \right\}, \tag{10}$$

where $\overline{\lambda}_{lj}^{(s)}, l = 1, 2$ is computed according to equation (2) and normalized such as $\sum_{j=1}^{n_S} \overline{\lambda}_{lj}^{(s)} = 1$ with species specific parameters $a_l^{(s)}$ and $A_l^{(s)}$ sampled from $\mathbb{P}(\boldsymbol{\theta}_l | \boldsymbol{W}^{(l)})$.

Overlap coefficient $\mathbf{\Delta_{ST}}$. For n_T time values and n_S number of sites, we sample $\Delta_{ST}^{(f)}$ from the posterior distribution $\mathbb{P}(\Delta_{ST} | \boldsymbol{W}^{(1)}, \boldsymbol{W}^{(2)})$ as

$$\widehat{\Delta}_{ST}^{(f)} = \frac{1}{n_T \times n_S} \sum_{i=1}^{n_T} \sum_{j=1}^{n_S} \min \left\{ \overline{\lambda}_{1j}^{(f)} \mathcal{T}_1^{(f)}(\tau_i), \overline{\lambda}_{2j}^{(f)} \mathcal{T}_2^{(f)}(\tau_i) \right\}, \tag{11}$$

where for species $l = 1, 2$ we calculate $\mathcal{T}_l^{(f)}$ according to equation (5) and $\overline{\lambda}_{lj}^{(f)}$ according to equation (2) with species specific parameters $a_l^{(f)}$, $A_l^{(f)}$, and $\boldsymbol{k}_l^{(f)}$ sampled from $\mathbb{P}(\boldsymbol{\theta}_l | \boldsymbol{W}^{(l)})$, but normalized such that

$$\sum_{i=1}^{n_T} \sum_{j=1}^{n_S} \overline{\lambda}_{1j}^{(f)} \mathcal{T}_1^{(f)}(\tau_i) = 1.$$

2.1.6. Implementation

All methods were implemented in the `C++` program `Tomcat`, available through a git repository at `https://bitbucket.org/WegmannLab/tomcat/`.

 2.2. Simulations

We assessed the performance of our algorithm using 100 replicates for each combination

of $J = 20$ or 100 device locations and $D = 1, 2, 5, 10, 20, 50$ or 100 days at which data was

collected. All simulations were conduced with one-dimensional $X_j \sim N(0, 1)$ and $Y_j \sim N(0, 1)$

and parameter choices such that on average one observation per species per location was

expected, and hence the expected number of observations per species was JD.

To evaluate the accuracy of our estimates we then estimated the overlap between two

species, since errors in parameter estimates directly translate into biases in overlap coefficients.

We thus simulated data for two species with little overlap ($\Delta_T = 0.2$), moderate overlap

($\Delta_T = 0.5$) or large overlap in time ($\Delta_T = 0.8$) as described in Table 1 as well as for two

species with varying overlap in space ($\Delta_S = 0.2$, 0.5, and 0.8) and with varying overlap in

space and time ($\Delta_{ST} = 0.2$, 0.5, and 0.8) as described in the Supporting Information.

2.3. Application to Central African duikers

We applied `Tomcat` to camera trapping data obtained during the dry seasons from 2012

to 2018 from a region in the eastern Central African Republic (CAR), a wilderness exceeding

100,000 km^2 without permanent settlements, agriculture or commercial logging (Aebischer

et al., 2017). The available data was from 1,059 camera traps set at 532 distinct locations

that cover the Aire de Conservation de Chinko (ACC), a protected area of about 20,000 km^2.

For more information about camera deployment and sampling design, see Aebischer et al.

(2017). Here, we focus on duikers *Cephalophinae*, which are a diverse mammalian group

common in the data set and for which near-perfect manual annotation was available.

To infer habitat preferences for these species, we benefited from an existing land cover

classification at a 30m resolution that represents the five major habitat types of the Chinko

region: moist closed canopy forest (CCF), open savanna woodland (OSW), dry lakéré

grassland (DLG), wet marshy grassland (WMG) and surface water (SWA) (Aebischer et al.,

2017). Around every camera trap location and at 10,200 regular grid points spaced 2.5 km

apart and spanning the entire ACC, we calculated the percentage of each of these habitats

in 11 buffers of sizes 30; 65; 125; 180; 400; 565; 1260; 1785; 3,990; 5,640 and 17,840 meters.

We complemented this information with the average value within every buffer for each of 15

additional environmental and bioclimatic variables from the WorldClim database version 2

(Table S.1 Fick and Hijmans, 2017) that we obtained at a resolution of 30 seconds, which

translates into a spatial resolution of roughly 1km^2 per grid cell. To aid in the interpretation,

we then processed our environmental data by 1) keeping only the additional effect of each

variables after regressing out the habitat variables CCF and OSW at the same buffer, and by

2) keeping only the additional effect of every variable after regressing out the information

contained in the same variable but at smaller buffers (see Supporting Information for details).

To avoid extrapolation, we restricted our analyses to 2,639 grid locations that exhibited

similar environments to those at which camera traps were placed as measured by the

Mahanalobis distance between each grid point and the average across all camera trap

locations (see Supporting Information for details). For each location, we further used the

binary classification of the four most common habitat types (CCF, OSW, MWG, DLG) and

determined the presence or absence of six additional habitat characteristics: animal path,

road, salt lick, mud hole, riverine zone and bonanza.

3. Results

3.1. Performance against simulations

We first assessed the performance of our algorithm by inferring overlap coefficients between

two species from simulated data. We chose to focus on overlap coefficients as they are directly

affected by any error or bias in parameter estimates. As shown in Figure 2, the posterior means

$\widehat{\Delta_T}$, $\widehat{\Delta_S}$ and $\widehat{\Delta_{ST}}$ were unbiased and highly accurate for all overlap coefficients regardless of

the true values, if sufficient data is provided, i.e. if at least several hundred pictures were

available ($J \times D \geq 500$). If less data was available, estimates were biased towards the prior

expectations, which are 0.5 and 1.0 for Δ_T and Δ_S, respectively.

4. Application to Central African duikers

215 We used `Tomcat` to study the spatio-temporal distribution and overlap of duikers *Cephalophi-*

216 *nae* in the eastern Central African Republic (CAR). We benefited from existing camera

217 trapping data (Aebischer et al., 2017), collected between 2012 and 2018 at 532 distinct

218 locations in the Aire de Conservation de Chinko (ACC), a protected area of about 20,000

219 km^2 that was established in 2014 by the government of the CAR in former hunting zones.

The ACC consists of an ecotone of tropical moist closed canopy forests of the northeastern Congolian lowland rain forest biome and a Sudanian-Guinean woodland savanna that is interspersed with small patches of edaphic grasslands on rocky ground or swampy areas (Boulvert, 1985; Olson and Dinerstein, 1998).

Duikers are common in the data set and often observed in sympatry, i.e. several species were captured by the same camera trap within a few hours. We detected a total of eight species in the data set (Table 2): Eastern Bay Duiker *Cephalophus dorsalis castaneus*, Uele White Bellied Duiker *Cephalophus leucogaster arrhenii*, Black Fronted Duiker *Cephalophus nigrifrons*, Red Flanked Duiker *Cephalophus rufilatus*, Western Yellow Backed Duiker *Cephalophus silvicultor castaneus*, Weyns Duiker *Cephalophus weynsi*, Eastern Blue Duiker *Philantomba monticola aequatorialis* and Bush Duiker *Sylvicapra grimmia*.

The eight duiker species varied greatly in their habitat preferences (Figures 3, S.2) as inferred by `Tomcat`. As shown in Figure 3, *C. dorsalis* and *C. weynsi* both have a strong preference for CCF over OSW habitat at the smallest buffers, in contrast to *S. grimmia* that shows a strong preference for OSW. At higher buffers the signal is less clear, probably owing to the heterogeneous nature of the habitat in which both CCF and OSW correlated negatively with WMG and DLG, habitats not well suited for any of these species. Interestingly, *P. monticola* and *C. silvicultor* seem to be true ecotone species preferring a mixture of the canonical habitats CCF and OSW (Figure S.2). Our analysis further suggests that the two rarely studied species *C. weynsi* and *C. leucogaster* not only occur in large blocks of CCF as suggested in the literature (Kingdon et al., 2013), but also in narrow gallery forests within the forest-savanna ecotone several kilometers away from the next extensive forest block (Figures 3 and S.2).

As shown in Figure 4 and S.1, the species also varied greatly in their daily activity patterns, with some being almost exclusively nocturnal (*C. dorsalis* and *C. silvicultor*), some almost exclusively diurnal (*C. leucogaster*, *P. monticola*, *C. nigrifons*, *C. rufilatus*, *C. weynsi*) and one crepuscular (*S. grimmia*).

To better understand how these closely related duiker species of similar size and nutritional needs can occur sympatrically, we estimated pairwise overlap coefficients in space and time (Figure 4, Table S.2). Not surprisingly, most species pairs differed substantially either in their habitat preferences or daily activity patterns. Of the two forest dwellers *C. dorsalis* and

$C.\ weynsi$ ($\widehat{\Delta_S} = 0.36$) for instance, one is almost exclusively nocturnal and the other almost exclusively diurnal ($\widehat{\Delta_T} = 0.18$), resulting in a small overlap in space and time ($\widehat{\Delta_{ST}} = 0.11$). Similarly, the nocturnal $C.\ dorsalis$ and the crepuscular $S.\ grimmia$ that share a lot of temporal overlap ($\widehat{\Delta_T} = 0.57$) use highly dissimilar habitats ($\widehat{\Delta_S} = 0.02$), resulting in a very small overlap in time and space ($\widehat{\Delta_{ST}} = 0.01$).

A visualization using Multidimensional Scaling (MDS) of the pair-wise overlap coefficients of all six species with observations from at least 50 independent camera trap locations is shown in Figure 5. For these species, 84.6% of variation in the temporal overlap can be explained by a single axis separating nocturnal from diurnal species. In contrast, only 44.5% of the variation in the spatial overlap is explained by the first axis distinguishing forest dwellers from savanna species. When using both temporal and spatial information, the two first axis explain 32.3% and 25.5%, respectively, suggesting that a single axis is not sufficient to explain both temporal and spatial differences between species.

A striking observation is that frequently observed and therefore evidently abundant species within a certain community tend to differ in their habitat preference and/or daily activity. In contrast, infrequently observed and therefore putative rare taxa seem to have large overlap with co-occurring species. $C.\ leucogaster$, for instance, which is rather rare and was only observed at 11 distinct locations (Table 2), has similar habitat preferences and is active at the same time (Table S.2) as $C.\ weynsi$, which is among the most common forest duikers within the ACC. In contrast, $C.\ dorsalis$, which is strictly nocturnal, seems to co-exist with the $C.\ weynsi$ at higher densities (Figure 4, Table S.2).

Conclusion

Despite world-wide efforts, there are still major areas for which almost no information on biodiversity is available (e.g. Hickisch et al., 2019). But technological advances, in particular camera traps and acoustic recorders, make it possible to obtain a first glimpse on the presence and distribution of larger or highly vocal animals like mammals and birds in relatively short time and with a reasonable budget. Thanks to increased battery life, larger media to store data and other technical advances, such data sets can now be produced with comparatively little manual labor, even under the demanding conditions of large and remote areas. In addition, the annotation of these data sets on the species level is now aided by machine learning algorithms

that automatize the detection of common species and recordings without observations (e.g. Norouzzadeh et al., 2018). Thanks to these developments, existing knowledge gaps may now be increasingly addressed, allowing for a re-evaluation of conservation strategies and optimization of conservation management.

Here we introduce `Tomcat`, an model that infers habitat preferences and daily activities from spatio-temporal observations and hence allows to learn about the ecological requirements of animals, including rare, elusive and unmarked species. Unlike many previous methods that estimate the presence or absence of a species, `Tomcat` estimates relative species densities, from which overlap coefficients between species can be estimated in space and time while accounting for variation in detection probabilities between locations. While estimates of overlap coefficients require larger data sets than the inference of pure occupancy, we believe they constitute a major step forward in understanding the complex species interactions in an area, which is particularly relevant for conservation planning in heterogeneous or fragmented habitats.

Acknowledgments

This work was supported by a Swiss National Science Foundation grant (31003A_173062) to DW.

Author contributions

TA and DW conceived the idea; SA, BW and DW developed and implemented the method; TA collected the data; SA, LS, TA analyzed the data; SA, LS and DW led the writing of the manuscript. All authors contributed critically to the drafts and gave final approval for publication.

References

Aebischer, T., Siguindo, G., Rochat, E., 2017. First quantitative survey delineates the distribution of chimpanzees in the eastern central african republic. Biological Conservation 213, 84–94.

Boulvert, Y., 1985. Carte phytogéographique de la république centrafricaine. ORSTOM
(Office de la recherche scientifique et technique Outre-Mer) .

Bu, H., Wang, F., Mcshea, W.J., Lu, Z., Wang, D., Li, S., 2016. Spatial Co-Occurrence and
Activity Patterns of Mesocarnivores in the Temperate Forests of Southwest China. PLOS
ONE , 1–15doi:10.1371/journal.pone.0164271.

Burton, A.C., Neilson, E., Moreira, D., Ladle, A., Steenweg, R., Fisher, J.T., Bayne, E.,
Boutin, S., 2015. Review: Wildlife camera trapping: a review and recommendations
for linking surveys to ecological processes. Journal of Applied Ecology 52, 675–685.
doi:10.1111/1365-2664.12432.

Efford, M.G., Dawson, D.K., 2012. Occupancy in continuous habitat. Ecosphere 3, art32.
doi:10.1890/ES11-00308.1.

Fick, S.E., Hijmans, R.J., 2017. Worldclim 2: new 1-km spatial resolution climate surfaces
for global land areas. International Journal of Climatology 37, 4302–4315. doi:10.1002/
joc.5086.

Frey, S., Fisher, J.T., Burton, A.C., Volpe, J.P., 2017. Investigating animal activity patterns
and temporal niche partitioning using camera-trap data: challenges and opportunities.
Remote Sensing in Ecology and Conservation 3, 123–132. doi:10.1002/rse2.60.

Green, P., 1995. Reversible jump markov chain monte carlo computation and bayesian model
determination. Biometrika 82, 711–732.

Hickisch, R., Hodgetts, T., Johnson, P.J., Sillero-Zubiri, C., Tockner, K., Macdonald, D.W.,
2019. Effects of publication bias on conservation planning. Conservation Biology 33,
1151–1163. doi:10.1111/cobi.13326.

Kingdon, J., Happold, D., Butynski, T., Hoffmann, M., Happold, M., Kalina, J., 2013.
Mammals of Africa. v. 1-6, Bloomsbury Publishing.

MacKenzie, D.I., Nichols, J.D., 2004. Occupancy as a surrogate for abundance estimation.
Animal biodiversity and conservation 27, 461–467.

MacKenzie, D.I., Nichols, J.D., Lachman, G.B., Droege, S., Andrew Royle, J., Langtimm, C.A., 2002. Estimating site occupancy rates when detection probabilities are less than one. Ecology 83, 2248–2255. doi:10.1890/0012-9658(2002)083[2248:ESORWD]2.0.CO;2.

MacKenzie, D.I., Royle, J.A., 2005. Designing occupancy studies: general advice and allocating survey effort. Journal of Applied Ecology 42, 1105–1114. doi:10.1111/j.1365-2664.2005.01098.x.

Norouzzadeh, M.S., Nguyen, A., Kosmala, M., Swanson, A., Palmer, M.S., Packer, C., Clune, J., 2018. Automatically identifying, counting, and describing wild animals in camera-trap images with deep learning. Proceedings of the National Academy of Sciences 115, E5716–E5725. doi:10.1073/pnas.1719367115.

Oliveira-santos, L.G.R., Zucco, C.A., Agostinelli, C., 2013. Using conditional circular kernel density functions to test hypotheses on animal circadian activity. Animal Behaviour 85, 269–280. URL: http://dx.doi.org/10.1016/j.anbehav.2012.09.033, doi:10.1016/j.anbehav.2012.09.033.

Olson, D.M., Dinerstein, E., 1998. The global 200: A representation approach to conserving the earth's most biologically valuable ecoregions. Conserv. Biol. 12, 502–515.

Ridout, M.S., Linkie, M., 2009. Estimating overlap of daily activity patterns from camera trap data. Journal of Agricultural, Biological, and Environmental Statistics 14, 322–337. URL: https://doi.org/10.1198/jabes.2009.08038, doi:10.1198/jabes.2009.08038.

Rowcliffe, J.M., Kays, R., Kranstauber, B., Carbone, C., Jansen, P.A., 2014. Quantifying levels of animal activity using camera trap data. Methods in Ecology and Evolution 5, 1170–1179.

Sollmann, R., 2018. A gentle introduction to camera-trap data analysis. African Journal of Ecology 56, 740–749. doi:10.1111/aje.12557.

Steenweg, R., Hebblewhite, M., Whittington, J., Lukacs, P., McKelvey, K., 2018. Sampling scales define occupancy and underlying occupancy–abundance relationships in animals. Ecology 99, 172–183. doi:10.1002/ecy.2054.

360 Tobler, M.W., Hartley, A., Carrillo-Percastegui, S.E., Powell, G.V.N., 2015. Spatiotemporal

361 hierarchical modelling of species richness and occupancy using camera trap data. Journal

362 of Applied Ecology 43, 413–421.

Tables

Table 1: Values of k used for the simulation of the daily activity patterns $\mathcal{T}(\tau)$.

Δ_T	Species	k_1	k_2	k_3	k_4	k_5	k_6	k_7	k_8
0.2	1	0.2	0.2	1.8	1.8	1.8	1.8	0.2	0.2
	2	1.8	1.8	0.2	0.2	0.2	0.2	1.8	1.8
0.5	1	0.5	0.5	1.5	1.5	1.5	1.5	0.5	0.5
	2	1.5	1.5	0.5	0.5	0.5	0.5	1.5	1.5
0.8	1	0.8	0.8	1.2	1.2	1.2	1.2	0.8	0.8
	2	1.2	1.2	0.8	0.8	0.8	0.8	1.2	1.2

Table 2: Available data on the eight detected species of duikers.

Species	Common Name	Pictures	Camera Traps
Cephalophus dorsalis castaneus	Eastern Bay Duiker	1,631	66
Cephalophus leucogaster arrhenii	Uele White Bellied Duiker	432	11
Cephalophus nigrifrons	Black Fronted Duiker	102	7
Cephalophus rufilatus	Red Flanked Duiker	5,762	168
Cephalophus silvicultor castaneus	Western Yellow Backed Duiker	10,321	222
Cephalophus weynsi	Weyns Duiker	10,037	146
Philantomba monticola aequatorialis	Eastern Blue Duiker	50,979	212
Sylvicapra grimmia	Bush duiker	5,626	124

 Figures

Figure 1: The solid line plot represents the piecewise-constant function $\mathcal{T}$ with $h_a = 3$ hours, and the dashed line plot represents $\mathcal{T}$ shifted with $\delta = 1$ hour.

Figure 2: Distribution of bias in the estimated overlap coefficients $\hat{\Delta}_T$, $\hat{\Delta}_S$ and $\hat{\Delta}_{ST}$ for different sample sizes.

Figure 3: Habitat preference of the three duiker species *C. dorsalis*, *C. weynsi* and *S. grimmia*. Left: Distribution of closed canopy forest (CCF, top, green) and open savanna woodland (OSW, bottom, yellow) across the study region with the ACC borders and camera trap locations (black dots). Top right: Relative densities d_{sj} of the three duikers predicted at 2,639 grid points. For each species the colors indicates $d_{sj} = \log_{10}\left(\bar{\lambda}_{sj}/\mathrm{median}(\bar{\lambda}_s)\right)$, where $\mathrm{median}(\bar{\lambda}_s)$ is the median value over all the grid points j. Red shades indicate $d_{sj} > 0$, blue shades $d_{sj} < 0$. Bottom right: Posterior inclusion probabilities for the CCF (green) and OSW (yellow) habitat variables for each buffer. Values above the dashed line indicate the posterior probability that the habitat correlates positively with the relative species density, values below the dashed line imply a negative correlation.

Figure 4: Interaction in space and time between the duiker species *C. dorsalis*, *C. weynsi*, and *S. grimma*. Top row: interactions in space quantified as $\log_{10}\left(\bar{\lambda}_{s_1 j}/\bar{\lambda}_{s_2 j}\right)$ between species 1 and 2. Bottom: posterior mean (solid line) and 90% credible intervals (shades) of temporal activity patterns. The area shaded in gray represents the overlap coefficient Δ_T.

Figure 5: Illustration of the overlap coefficients in time and space between six duiker species visualized in two dimensions using the multidimensional scaling.

A sparse observation model to quantify species interactions in time and space

—

Supporting Information

Sadoune. Ait Kaci Azzou[a,b,*], Liam. Singer[a,b,*], Thierry. Aebischer[a,b], Beat. Wolf[c],

Daniel. Wegmann[a,b,**]

[a]*Department of Biology, Université de Fribourg, Chemin du Musée 10, CH-1700 Fribourg, Switzerland*
[b]*Swiss Institute of Bioinformatics, 1700 Fribourg, Switzerland*
[c]*iCoSys, University of Applied Sciences Western Switzerland, Fribourg, Switzerland*

S1. Simulating data with specific overlap coefficients

Simulating data with specific Δ_S

For the simulation of scenarios with a given Δ_S, we have simulated for each site j, an environmental variable X_j from $N(0,1)$. Δ_S will be estimated with $\hat{\Delta}_S$, where

$$\hat{\Delta}_S = \mathbb{E}\left(\min\left(\bar{\lambda}_1(X), \bar{\lambda}_2(X)\right)\right),$$

where $\bar{\lambda}_i = \{\bar{\lambda}_{i1}, \cdots, \bar{\lambda}_{ij}, \cdots, \bar{\lambda}_{in}\}$ in presence of n sites. For a species i and site j, $\bar{\lambda}_{ij}$ is given by equation (2), with the constraint :

$$\int \bar{\lambda}_{ij}(x_j) f_{X_j}(x_j) dx_j = 1. \tag{S.12}$$

We have for species 1 :

$$\begin{aligned}
\text{(S.12)} \quad &\Leftrightarrow \int_{-\infty}^{+\infty} \exp(\mu_1 + A_1 x_j) \times \frac{1}{\sqrt{2\pi}} e^{-\frac{1}{2}x_j^2} dx_j = 1 \\
&\Leftrightarrow \exp(\mu_1 + \frac{1}{2}A_1^2) = 1 \\
&\Leftrightarrow \mu_1 = -\frac{1}{2}A_1^2.
\end{aligned} \tag{S.13}$$

Equivalently, we have for species 2, $\mu_2 = -\frac{1}{2}A_2^2$. Therefore,

$$\bar{\lambda}_{i1} = \exp(-\frac{1}{2}A_i^2 + A_i x_j), \; i = 1, 2. \tag{S.14}$$

We have

$$\hat{\Delta}_S = \int \min\left(\bar{\lambda}_1(x_j), \bar{\lambda}_2(x_j)\right) f_{X_j}(x_j) dx_j. \tag{S.15}$$

We have

$$\bar{\lambda}_1(x_j) < \bar{\lambda}_2(x_j) \iff \exp(-\tfrac{1}{2}A_1^2 + A_1 x_j) < \exp(-\tfrac{1}{2}A_2^2 + A_2 x_j)$$
$$\iff x_j < \tfrac{A_1 + A_2}{2} = B.$$

The integral in equation (S.15) is given by:

$$
\begin{aligned}
\hat{\Delta}_S &= \int_{-\infty}^{B} \exp(-\tfrac{1}{2}A_1^2 + A_1 x_j) \times \tfrac{1}{\sqrt{2\pi}} e^{-\frac{1}{2}x_j^2} dx_j + \int_{B}^{+\infty} \exp(-\tfrac{1}{2}A_2^2 + A_2 x_j) \times \tfrac{1}{\sqrt{2\pi}} e^{-\frac{1}{2}x_j^2} dx_j \\
&= \Phi(\tfrac{1}{2}(A_2 - A_1)) + (1 - \Phi(\tfrac{1}{2}(A_1 - A_2))) \\
&= 2 - 2\Phi(\tfrac{1}{2}(A_1 - A_2)),
\end{aligned}
$$

where Φ represents the cumulative distribution function (CDF) of the standard normal distribution.

Finally, we have

$$\frac{2 - \hat{\Delta}_S}{2} = \Phi(\frac{1}{2}(A_1 - A_2)) \tag{S.16}$$

It is possible using equation (S.16) to simulate a model for which $\Delta_S = a$.

For example, for $\Delta_S = \delta_s$, and for $A_1 = a_1$, it is possible to get the value of A_2 from (S.16), which gives

$$\Phi^{-1}\left(\frac{2 - \delta_S}{2}\right) = \frac{1}{2}(a_1 - A_2) \iff A_2 = a_1 - 2\Phi^{-1}\left(\frac{2 - \delta_S}{2}\right). \tag{S.17}$$

After computing A_2, it is easy to deduce the value of μ_2 using equation (S.13).

Simulating data with specific Δ_{ST}

To simulate scenarios for a given Δ_{ST}, we have simulated for each site j an environmental variable X_j from $N(0, 1)$, and $T \sim U_{[0,24]}$. Δ_{ST} is estimated using equation (11).

For a species i and site j, we have the constraint

$$\int_{-\infty}^{+\infty} \int_0^{24} \bar{\lambda}_{ij}(x_j)\mathcal{T}_i(\tau)f_T(\tau)f_{X_j}(x_j)dt\, dx_j \ = \ \mathbb{E}\left(\mathcal{T}_i(\tau) \times \bar{\lambda}_{ij}(X_j)\right) = 1,$$

$$= \ \mathbb{E}\left(\mathcal{T}_i(\tau)\right) \times \mathbb{E}\left(\bar{\lambda}_{ij}(X_j)\right),$$

where $\mathcal{T}_i(\tau)$ is the daily activity for a species i, $i = 1, 2$ observed at time t.

We have

$$\begin{aligned}
\mathbb{E}\left(\mathcal{T}_i(\tau)\right) &= \int_0^{24} \mathcal{T}_i(\tau)\tau_T(\tau)d\tau \\
&= \tfrac{1}{24} \int_0^{24} \mathcal{T}_i(\tau)d\tau \\
&= \tfrac{1}{24} \int_0^{24} \sum_{l=1}^{n_a} K_l \mathbb{1}_{[(l-1)h_a,\, l\times h_a]}d\tau \ , \\
&= \tfrac{1}{24}h_a \sum_{l=1}^{n_a} K_l \\
&= \tfrac{1}{24}h_a \times n_a = 1
\end{aligned}$$

and

$$\begin{aligned}
\mathbb{E}\left(\bar{\lambda}_{ij}(X_j)\right) &= \int_{-\infty}^{+\infty} \exp(\mu_i + A_i x_j)f_{X_j}(x_j)dx_j \\
&= \int_{-\infty}^{+\infty} \exp(\mu_i + A_i x_j) \times \tfrac{1}{\sqrt{2\pi}}e^{-\frac{1}{2}x_j^2}dx_j \quad , \\
&\Leftrightarrow \exp(\mu_i + \tfrac{1}{2}A_i^2) \int_{-\infty}^{+\infty} \tfrac{1}{\sqrt{2\pi}}e^{-\frac{1}{2}(x_j - A_i)^2}dx_j
\end{aligned}$$

which gives

$$\mathbb{E}\left(\mathcal{T}_i(T)\right) \times \mathbb{E}\left(\bar{\lambda}_{ij}(X_j)\right) = 1 \Leftrightarrow \mu_i = -\frac{1}{2}A_i^2. \tag{S.18}$$

We can therefore simulate the scenario $\Delta_{ST} = \delta_{ST}$ with known activity patterns $\mathcal{T}_1, \mathcal{T}_2$ respectively for species 1 and 2 as follows:

1. Simulate n_s $X_j \sim N(0,1)$, $j = 1, 2, \cdots n_s$.

2. Propose a value of A_1. (Should not be very large. Namely between -2 and 2).

3. Compute the value of μ_1 using equations (S.18).

4. Compute numerically the value of A_2 by solving the equation :

$$\delta_{ST} - \frac{1}{n_T \times n_S} \sum_{i=1}^{n_T} \sum_{j=1}^{n_S} \min\left\{\bar{\lambda}_{1j}^{(f)}\mathcal{T}_1^{(f)}(\tau_i), \bar{\lambda}_{2j}^{(f)}\mathcal{T}_2^{(f)}(\tau_i)\right\} = 0. \tag{S.19}$$

5. Using `Tomcat` for the two species, and by fixing the value A_i, $i = 1, 2$, we can sample from the posterior distribution of Δ_{ST} and compute the posterior mean of the values Δ_{ST} using equation (11).

S2. Decorrelation environmental variables

While `Tomcat` readily handles correlated environmental variables, we chose to decorrelate specific variables to aid in interpretation. Specifically, we processed our environmental data as follow (two steps):

Step 1: A major interest in our application was to study the impact of the prevalence of closed canopy forest (f) and savanna (s) habitat on species densities. For each scale (buffer) b, we therefore regress each environmental variable $V_{ib}, i \neq f, s$:

$$V_{ib} = \alpha_{fb}V_{fb} + \alpha_{sb}V_{sb} + \epsilon_{ib}, \tag{S.20}$$

where V_{fb} and V_{sb} represents, respectively, the forest and the Savannah habitat for a buffer $b = 1, \cdots, B$, $i = 1, \cdots, n_{env}$, and ϵ_{ib} is the error from the linear model described by equation (S.20), which captures the information of the variable V_{ib} independent of V_{fb} and V_{sb} at buffer b. We therefore replace the V_{ib} variables by $\widetilde{V}_{ib} = \epsilon_{ib}$ in our model, but kept $\widetilde{V}_{fb} = V_{fb}$ and $\widetilde{V}_{sb} = V_{sb}$.

Step 2: To evaluate relevant spatial scale of environmental variables, we also regressed out the larger buffers from the smaller one as:

$$\widetilde{V}_{ib} = \sum_{j=1}^{b-1} \beta_j \widetilde{V}_{i,j} + e_{ib}, \tag{S.21}$$

In the second step, and for a given environmental variable $\widetilde{V}_{ib}$ at the scale, we give priority to the smaller scales by keeping only the additional explanation by the studied variable to what we already know by replacing $\widetilde{V}_{ib}$ by $\widetilde{\widetilde{V}}_{ib} = e_{ib}$.

S3. Restricting analysis to environmentally homogeneous regions

Let $\overrightarrow{X} = (x_1, x_2, \cdots, x_n)^T$ be a matrix where the rows represent the observed camera traps, and the columns the environmental variables, and let $\overrightarrow{Y} = (y_1, \cdots, y_m)^T$ a matrix containing the environmental variables for the locations for which we want to predict $\bar{\lambda}_s$, $s = 1, 2, \cdots, m$.

We define the Mahanalobis distance D_O of $\overrightarrow{X}$ given by

$$D_O = \sqrt{(\overrightarrow{X} - \overrightarrow{\mu})^T S(\overrightarrow{X} - \overrightarrow{\mu})},$$

436 where $\overrightarrow{\mu} = (\mu_1, \cdots, \mu_n)$, and S is the variance covariance matrix.

437 We define a second Mahalanobis distance D_P which measures the distance of $\overrightarrow{Y}$ from the the

438 mean $\overrightarrow{\mu}$ of the camera traps environmental variables. D_P is given by

$$D_P = \sqrt{(\overrightarrow{Y} - \overrightarrow{\mu})^T S (\overrightarrow{Y} - \overrightarrow{\mu})}.$$

439 After computing D_O and D_P, we decided to remove the grid points for which we have

440 $D_P > 1.2 \times \max(D_O)$.

 ## S4. Supplementary figures

Figure S.1: Estimation of the daily activity of the eight observed duiker species. The solid line represents the posterior mean and the shades, 90% credible intervals of the temporal activity patterns.

Figure S.2: Relative densities d_{sj} of the eight duiker species predicted at 2,639 grid points. For each species the colors indicates $d_{sj} = \log_{10}\left(\bar{\lambda}_{sj}/\mathrm{median}(\bar{\lambda}_s)\right)$, where $\mathrm{median}(\bar{\lambda}_s)$ is the median value over all the grid points j. Red shades indicate $d_{sj} > 0$, blue shades $d_{sj} < 0$. Posterior inclusion probabilities for the CCF (green) and OSW (yellow) habitat variables for each buffer. Values above the dashed line indicate the posterior probability that the habitat correlates positively with the relative species density, values below the dashed line imply a negative correlation.

 S5. Supplementary tables

Table S.1: Environmental and bioclimatic variables obtained from the WorldClim database version 2.

Variable name	Description
dem	Digital elevation model
slope	Slope of a point
aspect	Direction of slope
TRI	Terrain Ruggedness Index
TWI	Topographic Wetness Index
wind	wind speed (m s^{-1})
bio2	Mean Diurnal Range (Mean of monthly (max temp - min temp))
bio5	Max Temperature of Warmest Month
bio6	Min Temperature of Coldest Month
bio12	Annual Precipitation
bio15	Precipitation Seasonality (Coefficient of Variation)
NDVI	Normalized Difference Vegetation Index
GC	General Curvature
MPI	Morphometric Protection Index
srad	Solar radiation (kJ m^{-2} day^{-1})

Table S.2: Summary of the overlap coefficients between the six duiker species.

	C. dorsalis	C. leucogaster	C. nigrifrons	C. rufilatus	C. silvicultor	C. weynsi	P. monticola	S. grimmia
$\bar{\Delta}_T$: Overlap coefficient in time								
C. dorsalis	0	0.285	0.338	0.235	0.752	0.176	0.202	0.571
C. leucogaster	0.285	0	0.482	0.632	0.384	0.632	0.595	0.524
C. nigrifrons	0.338	0.482	0	0.524	0.406	0.515	0.483	0.495
C. rufilatus	0.235	0.632	0.524	0	0.350	0.774	0.703	0.566
C. silvicultor	0.752	0.384	0.406	0.350	0	0.261	0.333	0.717
C. weynsi	0.176	0.632	0.515	0.774	0.261	0	0.747	0.427
P. monticola	0.202	0.595	0.483	0.703	0.333	0.747	0	0.530
S. grimmia	0.571	0.524	0.495	0.566	0.717	0.427	0.530	0
$\bar{\Delta}_S$: Overlap coefficient in space								
C. dorsalis	0	0.164	0.137	0.048	0.280	0.365	0.233	0.016
C. leucogaster	0.164	0	0.044	0.013	0.103	0.158	0.081	0.006
C. nigrifrons	0.137	0.044	0	0.072	0.158	0.171	0.194	0.045
C. rufilatus	0.048	0.013	0.072	0	0.308	0.065	0.131	0.307
C. silvicultor	0.280	0.103	0.158	0.308	0	0.350	0.332	0.202
C. weynsi	0.365	0.158	0.171	0.065	0.350	0.000	0.273	0.028
P. monticola	0.233	0.081	0.194	0.131	0.332	0.273	0	0.099
S. grimmia	0.016	0.006	0.045	0.307	0.202	0.028	0.099	0
$\bar{\Delta}_{ST}$: Overlap coefficient in space and time								
C. dorsalis	0	0.082	0.074	0.023	0.257	0.111	0.089	0.013
C. leucogaster	0.082	0	0.032	0.011	0.065	0.132	0.066	0.004
C. nigrifrons	0.074	0.032	0	0.054	0.099	0.123	0.131	0.033
C. rufilatus	0.023	0.011	0.054	0.000	0.166	0.062	0.117	0.233
C. silvicultor	0.257	0.065	0.099	0.166	0.000	0.147	0.168	0.185
C. weynsi	0.111	0.132	0.123	0.062	0.147	0	0.247	0.021
P. monticola	0.089	0.066	0.131	0.117	0.168	0.247	0	0.074
S. grimmia	0.013	0.004	0.033	0.233	0.185	0.021	0.074	0