BIOLOGICAL SCIENCES

CLATHRIN LIGHT CHAIN DIVERSITY REGULATES BUDDING EFFICIENCY IN VITRO

AND SYNAPTIC VESICLE FORMATION IN VIVO

Lisa Redlingshöfer[1, 2], Faye McLeod[3], Marine D. Camus[1, 2], Yu Chen[1, 2], Jemima J. Burden[4],

Ernest Palomer[3], Kit Briant[1, 2], Philip N. Dannhauser[1, 2], Patricia C. Salinas[3] and Frances M.

Brodsky[1, 2]*

[1] Research Department of Structural and Molecular Biology, Division of Biosciences,

University College London, Gower Street, London WC1E 6BT, UK

[2] Institute of Structural and Molecular Biology, Birkbeck and University College London,

Malet Street, London WC1E 7HX, UK

[3] Research Department of Cell and Developmental Biology, Division of Biosciences,

University College London, Gower Street, London WC1E 6BT, UK

[4] Medical Research Council Laboratory for Molecular Cell Biology, University College

London, Gower Street, London WC1E 6BT, UK

*to whom correspondence should be addressed:

Professor Frances M. Brodsky

f.brodsky@ucl.ac.uk

ABSTRACT

Clathrin light chain (CLC) subunits in vertebrates are encoded by paralogous genes *CLTA*

and *CLTB* and both gene products are alternatively spliced in neurons. To understand how

this CLC diversity influences neuronal clathrin function, we characterised the biophysical

properties of clathrin comprising individual CLC variants for correlation with neuronal

phenotypes of mice lacking either CLC-encoding gene. CLC splice variants differentially

influenced clathrin knee conformation within assemblies, and clathrin with neuronal CLC

mixtures was more efficient in membrane budding than clathrin with single neuronal isoforms

nCLCa or nCLCb. Correspondingly, electrophysiological recordings revealed that neurons

from mice lacking nCLCa or nCLCb were both defective in synaptic vesicle replenishment.

Mice with only nCLCb had a reduced synaptic vesicle pool and impaired neurotransmission

compared to wild-type mice, while nCLCa-only mice had increased synaptic vesicle

numbers, restoring normal neurotransmission. These findings highlight functional differences

between the CLC isoforms and show that isoform mixing influences tissue-specific clathrin

function in neurons.

KEY WORDS

Clathrin, membrane traffic, coated vesicle formation, protein isoforms, neuronal synapse

SIGNIFICANCE STATEMENT

This study reveals that diversity of clathrin light chain (CLC) subunits alters clathrin

properties and demonstrates that the two neuronal CLC subunits work together for optimal

clathrin function in synaptic vesicle formation. Our findings establish a role for CLC diversity

in synaptic transmission and illustrate how CLC variability expands the complexity of clathrin

to serve tissue-specific functions.

INTRODUCTION

Clathrin mediates vesicle formation from the plasma membrane and endosomal compartments (1). Recruited by cargo-recognising adaptor proteins, triskelion-shaped clathrin proteins assemble into polyhedral lattices to capture membrane-associated cargo and promote membrane bending into clathrin-coated vesicles (CCVs). Through sequestration of a variety of cargo, CCVs play fundamental roles in general cellular physiology including regulation of nutrient uptake and signalling, as well as in tissue-specific membrane traffic such as synaptic vesicle (SV) generation (2). This range of clathrin functions has been attributed to adaptor and accessory protein variation (3). However, functional diversity is also generated by variability of clathrin subunits. Vertebrates have two types of clathrin heavy chains with distinct functions (4). The major vertebrate clathrin is formed from clathrin heavy chain CHC17 (herein referred to as CHC) associated with clathrin light chains (CLCs), which do not bind the minor CHC22 clathrin isoform. Vertebrate CLCs are encoded by two different genes *CLTA* and *CLTB* (in humans), producing CLCa and CLCb isoforms of about 60% sequence identity (5, 6). Their sequence differences were conserved throughout evolution, after their encoding genes arose through duplication, suggesting the two isoforms can mediate distinct functions (7, 8). Expression levels of the CLCa and CLCb isoforms are tissue-specific (9), and further variation is created by alternative gene splicing during development (10) and in brain (5, 6). Here we address how CLC diversity affects the biophysical properties of clathrin and how the resulting variation affects the specialised function of clathrin in synaptic vesicle (SV) replenishment.

Limited functional differences between CLCa and CLCb have been observed in cell culture with respect to clathrin dynamics (11), during focal adhesion formation (12) and during cancer cell migration (13), with mechanisms attributed to possible isoform-specific differences in CLC binding proteins and post-translational modification (14). Loss of CLCs in culture and *in vivo* affects CCV uptake of some, but not all cargo, possibly reflecting

variability in mechanical demand for packaging different cargo into CCVs (9, 14, 15). The

capacity for biophysical differences in clathrin comprising different CLC isoforms and their

splicing variants to influence tissue-specific clathrin functions has not yet been considered.

CLC variability has potential to affect clathrin function as CLCs interact with key domains of

the clathrin triskelion that contribute to clathrin-mediated membrane bending. They bind the

trimerisation domain (TxD) of the triskelion vertex where the three component CHCs interact

(16, 17), and extend along the triskelion leg to the bend at the knee where CLCs regulate

conformation (18). Like various other endocytic proteins (3), CLCs undergo neuron-specific

splicing (19), which introduces one exon in CLCb (encoding 18 residues) and two exons

(encoding 30 residues) in CLCa at equivalent positions near the CLC C-termini, resulting in

higher molecular weight forms nCLCb and nCLCa (5, 6), and suggests neuron-specific

functions for these variants. *In vitro*, CLCs are required for efficient clathrin-mediated

membrane vesiculation at low temperature (20). Thus, CLCs could affect clathrin-mediated

membrane deformation directly through their influence on clathrin lattice properties (20, 21),

as well as indirectly through cellular interaction with proteins that influence the actin

cytoskeleton (22-25).

SV replenishment by membrane traffic following neuronal degranulation is critical for

sustained neurotransmission. Endocytosis is required to regulated synaptic plasma

membrane surface area and to retrieve SV proteins (26), and SVs are generated from the

plasma membrane and endosome-like compartments (27). The essential sites of clathrin

function in these pathways are debated (2, 28), but dysfunction of clathrin-associated

endocytic proteins is associated with neurological defects observed in Parkinson's disease

(29), Alzheimer's disease (3) and in numerous animal models in which such proteins have

been genetically deleted (30-35).

Here, we assess how clathrin lattices formed with neuronal and non-neuronal CLC variants

could differentially affect function by correlating their *in vitro* biophysical properties with *in*

vivo neuronal phenotypes of mice lacking CLC-encoding genes. We found that CLC

composition significantly influenced clathrin lattice properties and their ability to form vesicles

from liposome membrane *in vitro*. CLC splice variation influenced assembly curvature by

regulating CHC knee conformation, resulting in decreased ability of individual neuronal splice

variants to deform membrane *in vitro*. This effect was ameliorated for lattices formed from a

mixture of neuronal CLC isoforms. Mice lacking either CLCa or CLCb isoforms showed

corresponding differences in SV replenishment and synaptic neurotransmission compared to

their wild-type (WT) littermates expressing both isoforms, and to each other. Collectively,

these functional differences between CLC isoforms observed *in vitro* and *in vivo* establish a

role for CLC diversity in regulating clathrin function in neurons.

RESULTS

CLC splice variants differentially influence lattice curvature via the triskelion knee

Within clathrin lattices, the knee bending angle determines whether a hexagon (non-

curvature inducing) or pentagon (curvature-inducing) is formed (36). In addition, the

characteristic pucker at the triskelion vertex and the crossing angle of interacting CHC legs

can further modulate lattice curvature (37, 38) (Fig. 1a). This versatility of clathrin assemblies

allows clathrin to sequester cargo of various sizes as well as to form stable, flat assemblies

that serve as signalling hubs (3, 39). For each closed cage, there is a fixed number of 12

pentagons, but a varying number of hexagons (40) (Fig. 1a). Thus, larger cages (i.e. lattices

of lower curvature) have a smaller pentagon to hexagon ratio. CLCs maintain the puckered

conformation of triskelia in flat assemblies (20) and structural studies showed that nCLCb

can influence conformation of the triskelion knee up to a degree that inhibits assembly (18).

Considering these effects on CHCs, we hypothesized that sequence differences between the

CLC isoforms and alterations due to splicing might modulate CLC influence on triskelion

conformation and lattice curvature.

To address this, we produced clathrin triskelia of defined CLC composition by reconstituting

tissue-derived CHC triskelia with recombinantly expressed CLCa, CLCb, nCLCa or nCLCb

(Fig. S1a-d) (41). We then induced assembly of these different clathrins into closed cages

and analysed their diameters as a measure of lattice curvature by electron microscopy.

Previous studies showed that tissue-derived, CLC-bound clathrin (native) forms two major

size classes, while CHC-only triskelia predominantly form small-sized cages (42-45). Our

results confirmed that without CLCs, CHC formed predominantly cages of less than 90 nm in

diameter with an average size of around 70 nm, representing cages of up to 60 triskelia (44).

Clathrin with each of the CLC variants was further able to form larger cages with an average

diameter of around 110 nm (140 triskelia) for those cages. The degree to which larger cages

were formed varied significantly between splice variants (Fig. 1b, c), indicating that the

neuronal splice inserts regulate the influence of CLCs on lattice curvature.

To establish whether differences in larger cage formation resulted from CLC influence on the

triskelion vertex or on the CHC knee conformation or both, we dissected the effect of CLCs

separately for each domain. To test whether CLC diversity exerts influence through varying

CHC knee conformation, we produced clathrin cages from two CHC fragments that together

constitute a full-length clathrin triskelion "cut" at the knee (Fig. 1d) (46). The Hub fragment

(residues 1074-1675) was reconstituted with different CLCs (47) and combined with the

terminal-distal leg segment (TDD, residues 1-1074, Fig. S1e) under conditions promoting

their co-assembly (46). Cages produced when TDD was combined with CLC-reconstituted

Hub fragments were similar in size to fragment cages without CLCs, and the larger cages

observed for CLC variants associated with intact CHC did not form (Fig. 1d, e). This result

demonstrates that for CLC diversity to exert an effect on lattice curvature, the CHC knee

must be intact, and shows that CLC splice variants differ in their influence on CHC knee

conformation. Formation of larger cages by neuronal CLCs suggests that the neuronal splice

inserts reduce the likelihood of knee conformation favouring pentagon formation and support

hexagon formation. To establish whether cage size differences also correlated with CLC

effects at the TxD, the different CLCs were compared for their ability to stabilise the TxD (Fig.

S2) in an assay that measures triskelion dissociation. While dissociation differences were

observed between triskelia reconstituted with the different CLC isoforms and their splice

variants, these effects did not correlate with cage size differences, suggesting that CLC

influence at the vertex is not a major factor in lattice curvature.

CLC diversity modulates mechanical properties of the clathrin lattice

We next investigated whether the influence of CLCs on the CHC knee had mechanical

consequences for their ability to support membrane deformation. Mechanical properties of

clathrin lattices were previously shown to be CLC-dependent and reduced ability of CLC-free

clathrin to bend membranes *in vitro* was correlated with poorer planar lattice quality (20). To

establish whether CLC diversity influences lattice quality, clathrins with different CLC

composition were assembled on electron microscopy (EM) grids coated with a clathrin-

binding fragment of epsin1 (H$_6$-ΔENTH-epsin[144-575], Fig. S1f) (48). The flat lattices formed

were visualized by EM and their periodicity (quality) assessed by Fourier transform analysis.

Lattice quality was significantly reduced for clathrin with the neuronal splice variants of either

CLCa or CLCb compared to clathrin reconstituted with their respective non-neuronal variants

(Fig. 2a, b). While lattice quality was not significantly different between clathrin with nCLCa or

nCLCb, nCLCb clathrin showed a tendency to form lattices of poorer quality (Fig. 2c) and

lattice quality was significantly improved if formed from a 1:1 mix of nCLCa clathrin and

nCLCb clathrin (Fig. 2b). To visualise the nature of CLC distribution in these mixtures,

lattices were generated using His-tagged nCLCa clathrin or nCLCb clathrin combined with

the cognate untagged CLC and labelled with Ni-NTA-gold (Figure S3), confirming a uniform

distribution of each neuronal CLC within the mixed lattice. In contrast to the behaviour of the

neuronal CLCs, lattice quality was reduced in mixtures of CLCa clathrin and CLCb clathrin

compared to lattices formed by clathrin with only CLCa or CLCb. Notably, the lattice quality

resulting from neuronal and non-neuronal CLC compositions correlated with their influence

on cage size (Table 1), and suggests potential differential influence on membrane budding.

To determine whether CLC-related variation in lattice quality affects membrane bending, as

seen for complete absence of CLCs, we tested the function of the CLC-reconstituted

clathrins using a low temperature *in vitro* budding assay (20). In this assay, H$_6$-ΔENTH-

epsin[144-575], coupled to liposomes via modified Ni-NTA lipids, captures clathrin at the

liposome surface. Lattice assembly on these liposomes generates buds, which remain

attached to the liposome due to lack of dynamin, needed for scission (20, 48). The efficiency

of clathrin with different CLC composition to form coated buds was assessed at 15°C, a

temperature permissive for budding by native clathrin but reduced for CLC-free clathrin (20).

For each clathrin, we measured the diameter of clathrin-coated membrane profiles in thin-

section electron micrographs and assessed budding efficiency by the percentage of clathrin-

coated buds (defined by a fitted diameter < 200 nm) compared to all clathrin-coated

membrane profiles analysed (Fig. 3a). In line with previous findings, we found that at 15°C

native clathrin was about twice as efficient in membrane deformation as CLC-free clathrin

(20), and that budding efficiencies varied with the CLC composition of clathrin tested (Fig.

3b). CLCa or CLCb clathrin lattices vesiculated membrane more efficiently than either

neuronal clathrin variant (Fig. 3b), demonstrating that CLC splicing affects budding efficiency.

We found no statistically significant differences between CLCa and CLCb isoforms of the

neuronal or non-neuronal splice variants over the whole data set, yet differences within each

experiment indicate that in budding, clathrin with CLCb may be slightly more efficient than

CLCa clathrin, and nCLCb clathrin potentially less efficient than nCLCa clathrin (Fig. 3c).

Visualising the clathrin-coated liposomes absorbed to EM grids confirmed that all clathrins

tested assembled into intact lattices or buds on the liposome surface (Fig. S4), suggesting

that the observed variation in budding efficiency was not due to impaired lattice formation,

but CLC-dependent lattice properties. Notably, mixing reconstituted neuronal clathrins

significantly improved budding efficiency (Fig. 3d), overall reflecting the same pattern as

observed for both lattice quality and curvature (Table 1), suggesting all three parameters

result from differential CLC effects at the triskelion knee. Furthermore, all three assays

demonstrated that neuronal clathrin benefits from cooperative co-assembly of clathrin with

both isoforms.

Membrane deformation could result either from lattice formation at constant curvature or

through transitioning from flat to curved lattices (Fig. 3e). To determine how budding was

generated in our *in vitro* system and how it was influenced by CLCs, we measured the angle

between the convex side of the coat and the coat-free membrane (Fig. 3e), coat curvature

(i.e. diameter of the clathrin-coated structure), coat length and neck width for all coated

membrane profiles (49) and then correlated these measurements. We observed a variety of

coat curvatures in our system (Fig. 3f) and found that while neck width decreased with

budding angle, as would be expected for both modes of deformation, coat curvature and

neck width correlated with the budding angle of the coated to the coat-free membrane (Fig

3g, i). This indicates that lattices transition from flat to curved in our system, rather than

assembling with constant curvature, which would be independent of budding angle (Fig. 3e).

In further support of this, we found no correlation between coat length and budding angle

(Fig. 3h). Thus, we conclude that in our *in vitro* system, clathrin initially assembles into flat

lattices on the liposome membrane and then gradually deforms the underlying membrane

into coated buds, as observed by others in human SK-MEL-2 cells (49, 50). Our results

suggest that CLC-mediated regulation is required for the transition of flat to curved lattices

and that CLC variants differ in their ability to promote this transition. Thus, collectively from

these *in vitro* studies, we conclude that CLC variants differentially regulate triskelion knee

conformation within the lattice, affecting the mechanical ability to deform membrane into

coated buds through varying the probability of pentagon formation via influence on the CHC

knee. In particular, single neuronal splice variants were less functional in these properties

and lattice quality compared to their non-neuronal splice variants, while a mixture of neuronal

variants appeared to improve their functionality to non-neuronal levels.

CLC composition affects SV pool size and replenishment

Our *in vitro* experiments indicate that clathrin lattices formed from a mixture of clathrin with

each neuronal CLC isoform efficiently support membrane deformation, while clathrin

comprising only one neuronal CLC isoform is attenuated for membrane budding. Therefore,

we predicted that neurons of animals with only one CLC isoform might be defective in

clathrin-mediated pathways that rely on SV formation. This hypothesis was investigated in

knock-out (KO) mice lacking the *Clta* or *Cltb* genes in all tissues (CLCa KO and CLCb KO

mice). We previously generated CLCa KO mice (9) and have now produced CLCb KO mice.

Loss of *Cltb* in CLCb KO mice was confirmed by PCR, and no CLCb or nCLCb protein was

detected in all tissues analysed (Fig. S5). Whereas wild-type (WT) mice express a mixture of

CLCs in most tissues (9), CLCa KO mice express only CLCb or nCLCb, and CLCb KO mice

express only CLCa or nCLCa, enabling functional analysis of clathrin with only one type of

CLC in neurons (Fig. 4a).

Hippocampal neurons have frequently been used to study the function of endocytic proteins

in synaptic transmission (27, 30, 31, 33, 34, 51). H&E staining of the hippocampus showed

no morphological abnormalities in either KO strain, indicating that absence of either CLC

isoform does not affect gross hippocampus development in the CLC KO mice (Fig. S6). We

therefore used this system to evaluate the function of mature excitatory synapses in the CA1

region of the adult hippocampus, a region of well-defined neuronal architecture and

neurophysiological circuitry, in the CLC KO and WT mice. For electrophysiological recordings

at Schaffer collateral (SC)-CA1 synapses in acute hippocampal slices, a stimulating

electrode was placed in the SC fibres of the CA3 region and responses were then recorded

in the CA1 pyramidal cell layer (Fig. 4b).

To determine the role of CLC diversity in SV pool replenishment *in vivo*, we recorded the

responses to a prolonged high-frequency stimulus, which maximally depletes presynaptic

terminals of the readily releasable pool (RRP) of SVs (52-54). Under these conditions, initial

responses would draw from the pre-existing SV pool, while sustained neurotransmission

would rely on the efficiency of SV pool replenishment (53). Using this approach, we were

able to assess CLC-dependency for different stages of SV exo- and endocytosis from these

recordings. We found that the initial fusion rate of SVs in either KO strain was similar to that

of their WT littermates, indicating that correct packaging of fusion-mediating cargo was not

affected by changes in CLC composition (Fig. 4c, d and Fig. S7). Instead, both KO models

had a decreased SV recycling rate characteristic of a defect in acute SV pool replenishment

(Fig. 4c, d). This finding correlated with our expectation from *in vitro* properties of

reconstituted clathrin isoforms that showed clathrin lattices with only one type of neuronal

CLC were different from mixed lattices in their assembly properties (Table 1).

Further calculations based on data obtained from sustained trains of action potentials

indicated that the RRP was larger in the CLCb KO mice but reduced in the CLCa KO mice

when compared to WT littermates (Fig. 4c, d and Fig. S7), revealing a phenotype that we did

not expect from our *in vitro* data. To establish whether indeed the SV formation was altered

in the CLC KO mice rather than the ability to mobilise SVs from other SV pools, we assessed

the ultrastructure of excitatory synapses in the CLC KO mice and their WT littermates (Fig.

4e). Indeed, SV density was significantly reduced in hippocampal neurons in the CLCa KO

animals. In contrast, SV density in equivalent neurons of the CLCb KO mice was significantly

increased compared to their WT littermates (Fig. 4f, j), in line with our RRP estimates from

electrophysiological recordings (Fig. 4c, d). Other parameters such as individual SV size

(Fig. 4g, k), postsynaptic density (PSD) length (Fig. 4h, l) and excitatory synapse density

(Fig. 4i, m) were similar between both KO strains and respective WT littermates. Together,

these data show that CLCs are required for efficient and sustained synaptic function and

suggest that loss of either CLC decreases acute SV replenishment by affecting immediate

formation, correlating with expectations from reduced in vitro budding efficiency, but that the

KO strains differ in their ability to generally maintain SV pools. Thus, CLC composition

influences SV density in excitatory neurons of the hippocampus and indicates differential

functions for nCLCa and nCLCb in SV generation, such that mice with only nCLCa can

sustain a compensatory pathway for SV generation and mice with only nCLCb cannot.

CLCa KO and CLCb KO mice have distinct neurological and behavioural defects

To further investigate the differential consequences of CLCa or CLCb KO for synapse

function, we conducted additional electrophysiological experiments to characterise synaptic

transmission in our KO animals. Analyses of the evoked excitatory postsynaptic current

(EPSC) responses to stimuli of increasing magnitude revealed decreased EPSC response

amplitude in the CLCa KO mice compared to WT littermates, suggesting impaired basal

synaptic transmission in the CLCa KO mice (Fig. 5a). In contrast, the evoked EPSC

response amplitude in hippocampal slices from the CLCb KO mice was of similar or higher

magnitude than that of their WT littermates, suggesting that basal synaptic transmission was

intact and even enhanced (Fig. 5b). These differences in synaptic connectivity between

hippocampal function in the two KO strains could arise from defects in presynaptic

neurotransmitter release correlating with differences in SV pool size in the KO animals

relative to their WT littermates (Fig. 4).

Further, we performed paired-pulse ratio (PPR) recordings experiments, an approach that

allows assessment of defects in neurotransmitter release (55, 56). In excitatory neurons such

as those analysed here, a second stimulus pulse (P_2) fired in short succession after an initial

pulse (P_1) results in a larger response than the first stimulus. This facilitation is dependent on

changes in the probability of neurotransmitter release. PPR is inversely correlated with

release probability, as lower initial release probability leaves more fusion-competent vesicles

remaining at the terminal which are then released after the second stimulus (55, 56). We

observed that, compared to WT littermates, the PPR was increased in CLCa KO mice,

consistent with a reduction in release probability (Fig. 5c). In contrast, no differences in PPR

were observed in CLCb KO mice compared to WT littermates (Fig. 5d). Thus, specific loss of

nCLCa impairs presynaptic function compromising neurotransmitter release at the SC-CA1

synapse. Collectively, our electrophysiological data supports the morphological data (Fig. 4),

indicating that neurons with only nCLCa clathrin can compensate for defective SV

regeneration by expanding their SV pool from another pathway, while neurons with only

nCLCb clathrin cannot compensate and show decimated SV pools from impaired

replenishment.

Ongoing breeding of the CLCa KO colony confirmed a 50% survival rate compared to that

expected for homozygous CLCa KO mice (9), whereas homozygous CLCb KO mice had no

survival defects (Fig. 5e). Survival phenotypes similar to the CLCa KO animals have been

reported for other genetic deletions in endocytic pathways, which also displayed neuronal

phenotypes (30) in the hippocampus and other regions of the brain (32). Thus, we assessed

performance in the rotarod test for neuro-motor coordination for both CLC KO mice (57).

Compared to their WT littermates, surviving CLCa KO mice displayed defects in rotarod

balance, whereas CLCb KO mice did not (Fig. 5f). Further assessment of sensorimotor

function revealed that CLCa KO mice exhibited defects in a grid-walking test, (Fig. 5g), but

not in grip strength (Fig. 5h), suggesting neurological rather than muscular dysfunction in the

CLCa KO animals, supporting the notion that loss of nCLCa is less tolerable than loss of

nCLCb. Together, the phenotypes of the KO animals indicate that neuronal CLC isoforms are

differentially able to support compensatory mechanisms for adjusting SV pools hippocampal

synapses with impaired acute SV replenishment, and suggest that clathrin with only nCLCa

is more functional than clathrin with only nCLCb, while a mix of both neuronal CLCs is

necessary for optimal clathrin function in neurons, in keeping with their synergistic

contribution to clathrin function *in vitro.*

DISCUSSION

To understand the consequences of CLC diversity for clathrin function, we characterised the

biophysical properties of *in vitro* assemblies formed from clathrin comprising single CLC

isoforms and correlated these properties with neuronal phenotypes observed in KO animals

expressing single CLC isoforms. We found that neuronal CLC splicing affects lattice

properties (Fig. 1b and Fig. 2b) by diversifying the CLC influence on the CHC knee to

regulate lattice curvature (Fig. 1c, d) and deform membrane (Fig. 3b). In particular, CLC

splicing apparently influences clathrin's ability to introduce pentagons in order to transition

from flat to curved lattices (Fig. 1a and Fig. 3e-i). Our *in vitro* studies further suggest that

lattices formed from mixtures of clathrin with nCLCa and nCLCb have different assembly

properties and are more efficient in membrane budding compared to clathrin with only one

type of neuronal CLC (Fig. 2b, 3d). Consistent with this observation, neurons from both

CLCa and CLCb KO mice showed defects in synaptic transmission that indicated acute

clathrin-dependent SV replenishment was impaired (Fig. 4c, d). However, the CLCa KO mice

had reduced numbers of SVs in their hippocampal synapses, while the CLCb KO mice had

more SVs than their WT littermates (Fig. 4f, j). Thus, although acute SV replenishment was

impaired in both KO strains, clathrin with only nCLCa (in the CLCb KO mice) was able to

support a compensatory pathway of SV formation, while clathrin with only nCLCb (in the

CLCa KO) was not (Fig. 5a-d). These findings establish functional differences between CLC

isoforms *in vitro* and *in vivo* and demonstrate how resulting clathrin diversity is important for

clathrin function in neurons.

Neuronal splicing predominates for CLCb in neurons (58), while neuronal CLCs are not

present in other brain cells such as glial or Schwann cells (19), indicating that the CLC splice

variants segregate within brain cell types to fulfil distinct functions. Neuronal CLC splicing

had a significant influence on lattice curvature and budding efficiency, decreasing the

likelihood of forming pentagons (Table 1), which was attributable to effects on the CHC knee

domain (Fig. 1d). Given that the inserted sequences are located near the TxD where the C-

termini of CLCs are bound (arrowheads, Fig. 6a), we propose that the splice inserts affect

the conformation of the adjacent knee of a neighbouring triskelion, which is closer than the

knee of the triskelion to which the CLC is bound (Fig. 6b). This intermolecular influence could

involve splice inserts at the C-terminus (stars, Fig. 6b) interacting with the neighbouring CHC

or with the N-terminal domain of a CLC bound to the neighbouring knee (Ns, Fig. 6b).

Considering the average 40% sequence differences between CLCa and CLCb isoforms and

their even greater variation at the N-terminus, inter-CLC interactions could vary depending

on which splice isoform interacts with which CLC isoform N-terminus, thereby influencing

overall lattice properties. This imputed interaction may be lost from clathrin with non-neuronal

CLCs, which would explain why mixing non-neuronal CLC isoforms does not affect *in vitro*

budding to the same extent as mixing neuronal CLC isoforms (Fig. 3d).

Here, we characterise the properties of homo- and hetero-assemblies of clathrin comprising

single CLC isoforms. In brain, CLC isoforms are apparently randomly distributed on triskelia

(59), in which case neuronal inserted sequences could still influence lattice curvature via

interactions with an adjacent triskelion knee. The presence of nCLCb would serve as an

attenuator of nCLCa-specific interactions and vice versa. Further, in cells where CLCb

expression is transiently increased (60), newly synthesized triskelia would be generally

occupied by a single CLC isoform as clathrin subunit turnover is slow (61). The resulting

homotypic clathrins would then participate mixed in lattices, as we have studied here. Thus,

neuronal CLC isoform expression ratios (and/or local abundance within cells) could tailor

clathrin lattice properties specifically to particular needs.

There are several pathways involved in SV recycling in neurons and the role of clathrin has

been widely debated, possibly because the contribution of each pathway to SV formation is

variable between organisms, types of neurons and stimulus (51, 62, 63). SV recycling can

either be achieved by direct regeneration of SVs through clathrin-mediated endocytosis from

the plasma membrane at low-frequency stimulus, or through clathrin-independent

mechanisms, which predominantly mediate retrieval of SV components from the plasma

membrane under high stimulus (2, 27, 28, 33, 51). Rapid clathrin-dependent trafficking of SV

proteins from the endosomal pathway then leads to re-sorting and formation of SVs at high-

frequency stimulus (2), which can occur in a timeframe of 1-3 seconds after stimulation (27).

In our analysis of neurotransmission, SV recycling defects were detected by

electrophysiology within three seconds at high stimulus (20 Hz for 3s), suggesting the

defects detected were due to impaired clathrin-mediated regeneration of SVs from

endosomes. These findings are consistent with previously observed roles for CLCs in

recycling from non-neuronal endosomal compartments (60, 64).

That an acute SV pool replenishment defect was shared by neurons in both CLC KO strains

fits with our *in vitro* biophysical data that demonstrate single neuronal CLCs are not as

efficient in membrane deformation as a mixture of clathrin with both neuronal CLCs (Fig. 3d).

In response to this defect, we found that clathrin with only nCLCa generated an increased

steady-state SV pool in CLCb KO neurons, while clathrin with only nCLCb could not (Fig.

6b). CLCa seems to have preferential connection with the actin cytoskeleton compared to

CLCb (12, 24), which might account for the ability of nCLCa clathrin to function on its own

without nCLCb and benefit from involvement of actin and its accessory molecules to

compensate for defects in acute SV pool replenishment. It is possible that in the (normal)

presence of nCLCb, the actin interactions of nCLCa may be "diluted down", so that nCLCb

works as an attenuating balancing mechanism to control SV pools at steady state. The

presence of nCLCb would simultaneously create lattice properties for efficient CCV budding

during neurotransmission. In its absence, the acute replenishment pathway would decrease

in efficiency but loss of the nCLCb attenuation effect enables a compensatory pathway with

nCLCa only. Thus, the budding defect resulting from a change in biophysical properties of

clathrin is detectable only under acute demand, and when demand is less acute, there are

mechanisms by which nCLCa clathrin can generate SVs that are not supported by nCLCb

clathrin.

Considering the whole animal phenotypes, including the increased mortality rate for CLCa

but not CLCb KO animals, CLCa seems to be the more functional CLC, while CLCb functions

mainly in conjunction with CLCa. This is in line with the observation that CLC expression

ratios vary with tissue, and while lymphoid cells almost exclusively express CLCa, no tissue

has been found to express exclusively CLCb (9, 61). In addition, CLC isoform splicing

changes during cell differentiation, development (10, 58) and ratios of CLCb relative to CLCa

are modulated in tumour progression (13) and cell migration (60). These observations, in

combination with the *in vitro* and *in vivo* data reported here, support the concept that, in

addition to influencing accessory protein interaction, the conserved CLC isoform and splicing

differences characteristic of vertebrates facilitate tissue-specific clathrin functions by directly

modulating clathrin lattice properties. In particular, CLC diversity regulates clathrin budding

efficiency, a property that we show here affects SV formation under acute demand, and is

likely to influence other pathways that rely on rapid clathrin-mediated membrane traffic.

MATERIALS AND METHODS

Protein expression and purification

CLCs in pET28a were expressed with a thrombin-cleavable, N-terminal His-tag and purified

by standard nickel affinity chromatography and further polished by size exclusion

chromatography on a Superdex 200 10/300 GL column (GE Healthcare). His-tags of CLC

constructs were removed by thrombin cleavage. His-tagged epsin1 in pET32c and His-

tagged ΔENTH-epsin$^{144-575}$ in pQE32 were expressed and purified as previously described by

standard nickel affinity chromatography and size exclusion chromatography on a Superdex

200 10/300 GL column (GE Healthcare) (65).

CCVs were purified from porcine brain as described previously (45) and clathrin triskelia

purified by tris-extraction and size exclusion chromatography on a Superose 6 Increase

10/300 GL column (GE Healthcare). To generate CHC-only triskelia, endogenous CLCs were

removed using NaSCN and separated from CHC by size exclusion chromatography (41).

For reconstitution of clathrin of defined CLC composition, CHC in buffer C (50 mM Tris-HCl

pH 8.0, 50 mM NaCl, 2mM EDTA, 1mM DTT) at 0.8-1.0 mg/ml was incubated with purified

CLCs at 0.7 – 1.0 mg/ml in in the same buffer or in 20 mM Tris-HCl pH 8.0, 200 mM NaCl at

a 1:6.6 ratio (w/w) for 1 h on ice.

Electron microscopy

Freshly glow-discharged, homemade, carbon-coated formvar films on copper grids (Agar

Scientific) were used for all EM applications. All specimens were observed using a Tecnai

G2 (FEI) electron microscope at an acceleration voltage of 120 kV. Images were obtained

using a SIS Morada digital camera and TIA software (FEI).

Cage size and lattice curvature determination

Samples were dialysed in assembly-promoting buffer A (100 mM MES, 1 mM EGTA, 0.5 mM MgCl$_2$, 2 mM CaCl$_2$). Assembled cages were adsorbed to EM grids by placing grids on 10 µl droplets of samples at 0.2 mg/ml on Parafilm for 90 s. Excessive liquid was removed using filter paper and grids washed twice by transferring them sequentially onto two 15 µl droplets of buffer A. Samples were then stained with 2% uranyl acetate in water for 1 min and air-dried before subjected to EM analysis. Diameters of clathrin cages were measured from electron micrographs using ImageJ software (NIH). Distribution of diameters was averaged from the size distribution of 200 cages from three independent sets of experiments. For quantification of the proportion of the two cage populations, grouped into cages smaller or larger than 90nm in diameter, the proportions of the area under the curve for each population and experiment were averaged and tested for statistically significant differences between reconstituted clathrin isoforms using Prism 6 (GraphPad).

Lattice assembly and quality assessment

For incubation, EM grids were placed on 5-15 µl droplets on Parafilm at room temperature. Grids were firstly incubated with 0.04 mg/ml tag-free epsin 1 or H$_6$-ΔENTH-epsin[144-575] in buffer G (25 mM HEPES pH 7.2, 125 mM potassium acetate, 5 mM magnesium acetate) for 30 min. Unbound protein was removed by transferring grids to two droplets of buffer G before incubating grids with clathrin at 0.05 mg/ml in buffer G containing 0.1% BSA for 30 min. Lattices were fixed with 3% glutaraldehyde in buffer G for 15 min and stained with 2% or 5% uranyl acetate in water for 1 min. Lattice regularity was assessed from electron micrographs of similar quality and various focuses for each specimen as a measure of lattice quality (20). For this, 2D Fast Fourier Transform (FFT) was produced using ImageJ software (NIH). The height of the peak corresponding to the periodicity of the lattice (at ~0.036 nm^{-1}) served as a measure for lattice quality – the higher the peak, the higher the quality of the lattice. 5-12 images were analysed for each reconstituted clathrin isoform within each of three independent sets of experiments.

In vitro budding assay

In vitro budding was performed as previously described (48). In brief, H_6-ΔENTH-epsin$^{144-575}$

were bound to liposomes made from brain polar lipid extracts containing 5% DGS-Ni-NTA

lipids (Avanti). H_6-ΔENTH-epsin$^{144-575}$-coated liposomes were then chilled to 15°C and mixed

with clathrin triskelia stock solutions at 15°C, incubated for 30 min at 15°C and then

transferred on ice. Samples were then fixed at 4°C overnight and processed for EM analysis.

Diameter, coat length, budding angle and neck width of clathrin-coated membrane profiles

were measured from electron micrographs using ImageJ (NIH) and data processed using

Prism (GraphPad). Between 150 and 200 coat profiles, randomly sampled across four thin

sections, were analysed for each sample (> 60 000 nm total coat length per experiment per

sample). The proportion of curved lattices, defined by a diameter of less than 200 nm, of all

lattice profiles examined, was used as a measure of budding efficiency. Results from 3-5

independent sets of experiments were tested for statistical significance.

Generation of CLC KO mice

The *Cltb*$^{ko/ko}$ mouse strain used for this research project was created from ES cell clone

19159A-F4, generated by Regeneron Pharmaceuticals, Inc., and obtained from the KOMP

Repository (www.komp.org). Methods used to create the CLCb-null ES cell clone have

previously been published (66). In brief, the complete coding region of the *Cltb* gene was

completely deleted by homologous recombination using a large BAC-based targeting vector.

Targeted ES cells were then injected into albino C57BL/6J-N blastocytes and transferred into

foster mothers. Chimeric offspring were mated with C57BL/6J females (Charles River), and

germ-line transmission of the *Cltb*-null allele (*Cltb*^{ko}) was established. Heterozygote *Cltb*$^{ko/+}$

mice were backcrossed on the C57BL/6 background and bred to produce *Cltb*$^{ko/ko}$

homozygous mice. CLCa KO mice were derived from C57BL/6 WT mice (9).

Analysis of protein expression in brain

Mouse brains were harvested from six 9-12-month old C57BL/6 WT and homozygous CLCa and CLCb KO mice and quickly snap-frozen in liquid nitrogen and stored at −80°C until further use. Tissue were quickly thawed and homogenised in lysis buffer (50 mM HEPES pH 8.0, 50 mM NaCl, 1% Triton-X 100, 5 mM EDTA, 2 mM $CaCl_2$, 1 mM PMSF, cOmplete™ Protease Inhibitor Cocktail mix (Roche)). The homogenate was further incubated on ice for 45 min before centrifugation at 21 000 g and 4°C for 2 x 10 min in an Eppendorf 5424 R benchtop centrifuge (Eppendorf). Protein content of the lysate was determined by BCA assay (Thermo Fisher). For analysis, 25 µg of sample were loaded on 4-15% acrylamide gels (Bio-Rad) and subjected to SDS-PAGE and western blotting. Primary antibodies used for immunoblotting were T.D1 (anti-CHC, made in-house), X16 (anti-CLCa, made in-house), CLTB (anti-CLCb, Proteintech) and anti-beta-actin (Sigma-Aldrich).

Electrophysiology

Experiments were performed in 8-10-month-old mice. Both male and female mice were used for electrophysiological experiments. Acute transverse hippocampal slices (300µm) of homozygous CLCa or CLCb KO and control mice were cut on a Leica VT-1000 vibratome in ice-cold artificial cerebrospinal fluid (ACSF) bubbled with 95% O_2/5% CO_2 containing (in mM): NaCl (125), KCl (2.4), $NaHCO_3$ (26), NaH_2PO_4 (1.4), D-(+)-Glucose (20), $CaCl_2$ (0.5) and $MgCl_2$ (3). At 5-minute intervals, slices were then transferred into a series of 3 different chambers oxygenated (95% O_2/5% CO_2) in the same base ACSF but with the following temperature and component (in mM) variations: **1.** 21°C initially with $MgCl_2$ (1) and $CaCl_2$ (0.5) then allowed to heat gradually to 36°C; **2.** 36°C with $MgCl_2$ (1) and $CaCl_2$ (1); and **3.** 36°C initially with $MgCl_2$ (1) and $CaCl_2$ (2) before cooling to 21°C. Slices were then left for at least 1 hr before recordings commenced.

Evoked recordings were performed on an upright microscope continually perfused with oxygenated recording solution at room temperature containing the same ACSF composition

as the third chamber and supplemented with 10 µM bicuculline. Pyramidal cells in the CA1

region were held at -60mV in whole-cell voltage-clamp configuration using glass

microelectrodes (resistance 3-8 MΩ) filled with caesium gluconate intracellular solution

containing (in mM): D-gluconic acid lactone (130), HEPES (10), EGTA (10), NaCl (10), $CaCl_2$

(0.5), $MgCl_2$(1), ATP (1) and GTP (0.5), QX314 (5), pH to 7.2 with CsOH. To evoke

postsynaptic EPSCs, a bipolar concentric stimulation electrode (FHC) was placed in the SC

fibres of the CA3 region. For input-output recordings, the stimulus pulse was varied between

0.2 and 1mA with a pulse width of 0.1 ms and stimuli were delivered at a rate of 0.1 Hz.

Paired pulse stimuli were given at rate of 0.2 Hz with different inter-stimulus intervals,

ranging from 50 ms to 200 ms and a stimulation strength set to approximately 50% of the

maximal response for each cell. PPR was calculated as the ratio of the peak amplitude of the

second response over the first response. Calculation of RRP size, initial fusion efficiency and

SV recycling rate was done on ESPC recordings that underwent 3 s duration trains of

stimulation at 20Hz and estimated as previously described (53, 67). Briefly, RRP, fusion

efficiency (*fe*) and vesicle recycling rate (α) were evaluated from the cumulative charge

during the stimulation train using the following two equations:

Equation 1: $$fe = \frac{r(1)}{r(\infty)}\left(1 - \exp(-\alpha\Delta t)\right)$$

$$fe = \frac{r(1)}{\sum_{i=1}^{s} r(i)\exp(-\alpha(S-i)\Delta t)}$$

Equation 2: $$fe = \frac{r(1)}{\sum_{i=1}^{s} r(i)\exp(-\alpha(S-i)\Delta t)}$$

$r(1)$ is the charge of the first EPSC in the train

r(i) the charge passed by the ith EPSC

$r(\infty)$ was calculated from the average charge of the last 10 EPSCs in the train

Δt is the stimulus interval in the train

The RRP was estimated as RRP=r(1)/*fe*.

All currents were recorded using an Axopatch 200B amplifier, filtered (1 kHz) and digitised

(10 kHz). Data were monitored online and analysed offline using WinEDR and WinWCP

software (available free online at http://spider.science.strath.ac.uk/sipbs/software_ses.htm).

Stimulus artefacts in representative traces were digitally removed for clarity.

Electron microscopy of synapse ultrastructure

Experiments were performed in 4-12-month-old mice. Female mice were used for

ultrastructural experiments. Brain hemispheres were collected from homozygous KO mice

and WT littermates and cut in sagittal orientation. Slices were fixed in 2% paraformaldehyde

and 1.5% glutaraldehyde and stained sequentially with 1.5% potassium ferricyanide and 1%

OsO_4, 1% thiocarbohydrazide, 2% OsO4, 1% uranyl acetate, 0.66% lead nitrate in aspartic

acid. Samples were then gradually dehydrated and embedded for electron microscopy. The

ultrastructure of excitatory synapses in four electron micrographs of the VI lobule of the

cerebellum or CA1 hippocampus perinuclear region per genotype were analysed for the

ultrastructure of excitatory synapses using ImageJ (NIH) and Prism 6 (GraphPad) software

(n = 3). To determine SV density, SVs in 300 nm vicinity to the centre of the PSD in each

synapse were counted.

Mouse behavioural tests

All experiments were performed in accordance with the Animals Scientific procedures Act UK

(1986). Heterozygous $Cltb^{ko/+}$ mice were crossed to obtain homozygous $Cltb^{ko/ko}$ mice and

WT littermates. Heterozygous Clta$^{ko/+}$ mice were crossed to obtain homozygous $Clta^{ko/ko}$ mice

and WT littermates. Experiments were performed in 2-8-month-old mice. Male mice only

were used for all behavioural experiments.

Rotarod latency (Accelerating Rotarod; Ugo-Basile 7650 model) was used to assess motor

coordination. Mice were placed on the Rotarod, which was slowly accelerated from 3 to 30

rounds per minute over 5 minutes. Mice were given 3 trials per day for 2 days with at least 10

min inter-trial intervals. Time to fall from the rotarod (latency) was recorded for each trial.

Mice that remained on the Rotarod for the whole 5 min trial were assigned a 300 second

latency.

Grip strength (grip strength meter; Columbus Instruments) was used to measure muscle

strength in the forepaws. The grip strength meter was positioned horizontally, and mice were

held by the tail and allowed to grasp the metal pull-bar. The animals were then pulled back

and the force applied to the bar and the moment the grasp was released was recorded. Mice

were given 5 trials with 1 min inter-trial intervals and were tested on 2 days. The mean of

results from 10 trials was assigned for the grip strength of each animal.

Grid-walking was used to assess spontaneous motor deficits and limb movements involved

in precise stepping, coordination, and accurate paw placement. Adult mice were required to

navigate over a wire mesh grid. After each trial, 70% ethanol was used to clean the

apparatus. Mice were given 3 trials with 1 min inter-trial intervals and were tested on 2 days.

Behaviour on the grid was recorded on camera and was analysed later by an experimenter

who was blind to the genotype. A foot-slip was scored when the paw completely missed a

rung and the limb fell between the rungs. Counts of foot-slips for right forelimb, left forelimb,

right hindlimb and left hindlimb were obtained. Total footsteps were also recorded. The mean

of results from 10 trials was used for each animal.

Statistical analysis

All experiments were performed at least three times. All calculations and graphs were

performed with ImageJ, Microsoft Excel and GraphPad Prism software. P-values were

calculated using two-tailed Student's unpaired or paired (*in vitro* budding) t-tests, two-way

ANOVA with repeated measures (electrophysiology), one-way ANOVA followed by Holm-

Sidak correction for multiple comparison or one-way ANOVA for preselected pairs without

correction for multiple comparison (clathrin biophysical properties). Detailed statistical

information including statistical tests used, number of independent experiments, p values and

definition of error bars is listed in individual figure legends.

ACKNOWLEDGMENTS

This work was supported by grants to F. M. B. from the Wellcome Trust (107858/Z/15/Z), to

P.C.S. from the Medical Research Council (MR/M024083/1) and to P.C.S. and E. P. from

Alzheimer's Research UK (ARUK-PG2018A-002). L. R. was supported by a Wellcome Trust

4-year interdisciplinary PhD studentship. J. J. B. was supported by MRC funding to the MRC

Laboratory of Molecular Cell Biology at UCL, award code MC_U12266B. P. N. D. was

supported by a UCL Excellence Fellowship. The authors would like to acknowledge UCL

IQPath, UCL Institute of Neurology, Queen Square London, WC1N 3BG for processing

tissue slices for H&E staining and Massimo Signore (ICH) and Anna Crowley (KLB

Transgenic facility, UCL) for the generation of the *Cltb*[ko] mouse strain.

AUTHOR CONTRIBUTIONS

The study was conceived by L. R. and F. M. B. with expert input from P. N. D., F. M., E. P. V.

and P. C. S. and the project was managed by F.M.B. Electrophysiology experiments were

conducted and analysed by F. M. All *in vitro* reconstitution, protein level and EM experiments

and analyses were carried out by L. R. with expert assistance from J. J. B., F. M., E. P. V., P.

N. D. and K. B. Mouse behaviour data and survival characteristics were collected by M. D. C.

and Y. C. The paper was written by L. R., F. M. B., F. M. and P. C. S. and then all authors

read and commented on the paper.

COMPETING INTEREST

The authors declare no competing interest.

REFERENCES

1. F. M. Brodsky, Diversity of clathrin function: new tricks for an old protein. *Annu Rev Cell Dev Biol* **28**, 309-336 (2012).

2. T. Soykan, T. Maritzen, V. Haucke, Modes and mechanisms of synaptic vesicle recycling. *Curr Opin Neurobiol* **39**, 17-23 (2016).

3. H. T. McMahon, E. Boucrot, Molecular mechanism and physiological functions of clathrin-mediated endocytosis. *Nat Rev Mol Cell Biol* **12**, 517-533 (2011).

4. P. N. Dannhauser *et al.*, CHC22 and CHC17 clathrins have distinct biochemical properties and display differential regulation and function. *J Biol Chem* **292**, 20834-20844 (2017).

5. T. Kirchhausen *et al.*, Clathrin heavy chain: molecular cloning and complete primary structure. *Proc Natl Acad Sci U S A* **84**, 8805-8809 (1987).

6. A. P. Jackson, H. F. Seow, N. Holmes, K. Drickamer, P. Parham, Clathrin light chains contain brain-specific insertion sequences and a region of homology with intermediate filaments. *Nature* **326**, 154-159 (1987).

7. D. E. Wakeham *et al.*, Clathrin heavy and light chain isoforms originated by independent mechanisms of gene duplication during chordate evolution. *Proc Natl Acad Sci U S A* **102**, 7209-7214 (2005).

8. M. Fumagalli *et al.*, Genetic diversity of CHC22 clathrin impacts its function in glucose metabolism. *eLife* **8** (2019).

9. S. Wu *et al.*, Clathrin light chains' role in selective endocytosis influences antibody isotype switching. *Proc Natl Acad Sci U S A* **113**, 9816-9821 (2016).

10. J. Giudice *et al.*, Alternative splicing regulates vesicular trafficking genes in cardiomyocytes during postnatal heart development. *Nat Commun* **5**, 3603 (2014).

11. M. Mettlen *et al.*, Endocytic Accessory Proteins Are Functionally Distinguished by Their Differential Effects on the Maturation of Clathrin-coated Pits. *Mol Biol Cell* **20**, 3251-3260 (2009).

12. O. M. Tsygankova, J. H. Keen, A unique role for clathrin light chain a in cell spreading and migration. *J Cell Sci* 10.1242/jcs.224030 (2019).

13. P. H. Chen *et al.*, Crosstalk between CLCb/Dyn1-Mediated Adaptive Clathrin-Mediated Endocytosis and Epidermal Growth Factor Receptor Signaling Increases Metastasis. *Dev Cell* **40**, 278-288.e275 (2017).

14. H. Maib, F. Ferreira, S. Vassilopoulos, E. Smythe, Cargo regulates clathrin-coated pit invagination via clathrin light chain phosphorylation. *J Cell Biol* **217**, 4253 (2018).

15. J. C. Stachowiak, F. M. Brodsky, E. A. Miller, A cost-benefit analysis of the physical mechanisms of membrane curvature. *Nat Cell Biol* **15**, 1019-1027 (2013).

16. J. A. Ybe *et al.*, Light Chain C-Terminal Region Reinforces the Stability of Clathrin
Heavy Chain Trimers. *Traffic* **8**, 1101-1110 (2007).

17. D. S. Chu, B. Pishvaee, G. S. Payne, The Light Chain Subunit Is Required for
Clathrin Function in Saccharomyces cerevisiae. *J Biol Chem* **271**, 33123-33130
(1996).

18. J. D. Wilbur *et al.*, Conformation switching of clathrin light chain regulates clathrin
lattice assembly. *Dev Cell* **18**, 841-848 (2010).

19. D. H. Wong *et al.*, Neuron-specific expression of high-molecular-weight clathrin light
chain. *J Neurosci* **10**, 3025-3031 (1990).

20. P. N. Dannhauser *et al.*, Effect of clathrin light chains on the stiffness of clathrin
lattices and membrane budding. *Traffic* **16**, 519-533 (2015).

21. M. Lherbette, L. Redlingshofer, F. M. Brodsky, I. A. T. Schaap, P. N. Dannhauser,
The AP2 adaptor enhances clathrin coat stiffness. *FEBS J* **286**, 4074-4085 (2019).

22. C. Y. Chen, F. M. Brodsky, Huntingtin-interacting protein 1 (Hip1) and Hip1-related
protein (Hip1R) bind the conserved sequence of clathrin light chains and thereby
influence clathrin assembly in vitro and actin distribution in vivo. *J Biol Chem* **280**,
6109-6117 (2005).

23. J. D. Wilbur *et al.*, Actin binding by Hip1 (huntingtin-interacting protein 1) and Hip1R
(Hip1-related protein) is regulated by clathrin light chain. *J Biol Chem* **283**, 32870-
32879 (2008).

24. M. Biancospino *et al.*, Clathrin light chain A drives selective myosin VI recruitment to
clathrin-coated pits under membrane tension. *Nat Commun* 10.1038/s41467-019-
12855-6 (2019).

25. A. M. Schreij *et al.*, LRRK2 localizes to endosomes and interacts with clathrin-light
chains to limit Rac1 activation. *EMBO Rep* **16**, 79-86 (2014).

26. J. E. Heuser, T. S. Reese, Evidence for recycling of synaptic vesicle membrane
during transmitter release at the frog neuromuscular junction. *Journal of Cell Biology*
57, 315-344 (1973).

27. S. Watanabe *et al.*, Clathrin regenerates synaptic vesicles from endosomes. *Nature*
515, 228-233 (2014).

28. S. Watanabe, E. Boucrot, Fast and ultrafast endocytosis. *Curr Opin Cell Biol* **47**, 64-
71 (2017).

29. S. F. Soukup, R. Vanhauwaert, P. Verstreken, Parkinson's disease: convergence on
synaptic homeostasis. *EMBO J* **37**, e98960 (2018).

30. S. J. Koo *et al.*, Vesicular Synaptobrevin/VAMP2 Levels Guarded by AP180 Control
Efficient Neurotransmission. *Neuron* **88**, 330-344 (2015).

31. M. Cao *et al.*, Parkinson Sac Domain Mutation in Synaptojanin 1 Impairs Clathrin
Uncoating at Synapses and Triggers Dystrophic Changes in Dopaminergic Axons.
Neuron **93**, 882-896.e885 (2017).

32.   I. Milosevic *et al.*, Recruitment of Endophilin to Clathrin-Coated Pit Necks Is Required
750   for Efficient Vesicle Uncoating after Fission. *Neuron* **72**, 587-601 (2011).

751   33.   Natalia L. Kononenko *et al.*, Clathrin/AP-2 Mediate Synaptic Vesicle Reformation
752   from Endosome-like Vacuoles but Are Not Essential for Membrane Retrieval at
753   Central Synapses. *Neuron* **82**, 981-988 (2014).

754   34.   Y.-I. Yim *et al.*, Endocytosis and clathrin-uncoating defects at synapses of auxilin
755   knockout mice. *Proceedings of the National Academy of Sciences* **107**, 4412 (2010).

756   35.   T. Mitsunari *et al.*, Clathrin adaptor AP-2 is essential for early embryonal
757   development. *Mol Cell Biol* **25**, 9318-9323 (2005).

758   36.   A. Musacchio *et al.*, Functional organization of clathrin in coats: combining electron
759   cryomicroscopy and X-ray crystallography. *Mol Cell* **3**, 761-770 (1999).

760   37.   W. K. den Otter, M. R. Renes, W. J. Briels, Self-assembly of three-legged patchy
761   particles into polyhedral cages. *J Phys: Condens Matter* **22**, 104103 (2010).

762   38.   A. Fotin *et al.*, Molecular model for a complete clathrin lattice from electron
763   cryomicroscopy. *Nature* **432**, 573-579 (2004).

764   39.   F. Baschieri *et al.*, Frustrated endocytosis controls contractility-independent
765   mechanotransduction at clathrin-coated structures. *Nat Commun* **9**, 3825 (2018).

766   40.   S. Schein, Architecture of clathrin fullerene cages reflects a geometric constraint--the
767   head-to-tail exclusion rule--and a preference for asymmetry. *J Mol Biol* **387**, 363-375
768   (2009).

769   41.   F. K. Winkler, K. K. Stanley, Clathrin heavy chain, light chain interactions. *EMBO*
770   *Journal* **2**, 1393-1400 (1983).

771   42.   P. K. Nandi, H. T. Pretorius, R. E. Lippoldt, M. L. Johnson, H. Edelhoch, Molecular
772   properties of the reassembled coat protein of coated vesicles. *Biochemistry* **19**, 5917-
773   5921 (1980).

774   43.   P. P. Van Jaarsveld, P. K. Nandi, R. E. Lippoldt, H. Saroff, H. Edelhoch,
775   Polymerization of clathrin protomers into basket structures. *Biochemistry* **20**, 4129-
776   4135 (1981).

777   44.   J. Heuser, T. Kirchhausen, Deep-etch views of clathrin assemblies. *J Ultrastruct Res*
778   **92**, 1-27 (1985).

779   45.   E. Ungewickell, H. Ungewickell, Bovine brain clathrin light chains impede heavy chain
780   assembly in vitro. *J Biol Chem* **266**, 12710-12714 (1991).

781   46.   B. Greene, S. H. Liu, A. Wilde, F. M. Brodsky, Complete reconstitution of clathrin
782   basket formation with recombinant protein fragments: adaptor control of clathrin self-
783   assembly. *Traffic* **1**, 69-75 (2000).

784   47.   S.-H. Liu, M. L. Wong, C. S. Craik, F. M. Brodsky, Regulation of Clathrin Assembly
785   and Trimerization Defined Using Recombinant Triskelion Hubs. *Cell* **83**, 257-267
786   (1995).

787   48.   P. N. Dannhauser, E. J. Ungewickell, Reconstitution of clathrin-coated bud and
788   vesicle formation with minimal components. *Nat Cell Biol* **14**, 634-639 (2012).

49. O. Avinoam, M. Schorb, C. J. Beese, J. A. G. Briggs, M. Kaksonen, Endocytic sites
mature by continuous bending and remodeling of the clathrin coat. *Science* **348**,
1369-1372 (2015).

50. B. L. Scott *et al.*, Membrane bending occurs at all stages of clathrin-coat assembly
and defines endocytic dynamics. *Nat Commun* **9**, 419 (2018).

51. S. Watanabe *et al.*, Ultrafast endocytosis at mouse hippocampal synapses. *Nature*
504, 242-247 (2013).

52. L. E. Dobrunz, Release probability is regulated by the size of the readily releasable
vesicle pool at excitatory synapses in hippocampus. *International Journal of*
Developmental Neuroscience **20**, 225-236 (2002).

53. J. F. Wesseling, D. C. Lo, Limit on the role of activity in controlling the release-ready
supply of synaptic vesicles. *J Neurosci* **22**, 9708-9720 (2002).

54. C. F. Stevens, J. H. Williams, Discharge of the Readily Releasable Pool With Action
Potentials at Hippocampal Synapses. *Journal of Neurophysiology* **98**, 3221-3229
(2007).

55. L. E. Dobrunz, C. F. Stevens, Heterogeneity of release probability, facilitation, and
depletion at central synapses. *Neuron* **18**, 995-1008 (1997).

56. R. S. Zucker, W. G. Regehr, Short-term synaptic plasticity. *Annu Rev Physiol* **64**,
355-405 (2002).

57. H. Shiotsuki *et al.*, A rotarod test for evaluation of motor skill learning. *J Neurosci*
Methods **189**, 180-185 (2010).

58. S. Stamm *et al.*, Clathrin light chain B: gene structure and neuron-specific splicing.
Nucleic Acids Res **20**, 5097-5103 (1992).

59. T. Kirchhausen, S. C. Harrison, P. Parham, F. M. Brodsky, Location and distribution
of the light chains in clathrin trimers. *Proc Natl Acad Sci U S A* **80**, 2481-2485 (1983).

60. S. R. Majeed *et al.*, Clathrin light chains are required for the gyrating-clathrin recycling
pathway and thereby promote cell migration. *Nat Commun* **5**, 3891 (2014).

61. S. L. Acton, Brodsky, F. M., Predominance of clathrin light chain LCb correlates with
the presence of a regulated secretory pathway. *J Cell Biol* **111**, 1419-1426 (1990).

62. H. Heerssen, R. D. Fetter, G. W. Davis, Clathrin Dependence of Synaptic-Vesicle
Formation at the Drosophila Neuromuscular Junction. *Curr Biol* **18**, 401-409 (2008).

63. K. Sato *et al.*, Differential requirements for clathrin in receptor-mediated endocytosis
and maintenance of synaptic vesicle pools. *Proc Natl Acad Sci U S A* **106**, 1139-1144
(2009).

64. V. Poupon *et al.*, Clathrin light chains function in mannose phosphate receptor
trafficking via regulation of actin assembly. *Proceedings of the National Academy of*
Sciences **105**, 168-173 (2008).

65. C. Kalthoff, J. Alves, C. Urbanke, R. Knorr, E. J. Ungewickell, Unusual Structural
Organization of the Endocytic Proteins AP180 and Epsin 1. *J Biol Chem* **277**, 8209-
8216 (2002).

66.   D. M. Valenzuela *et al.*, High-throughput engineering of the mouse genome coupled
830   with high-resolution expression analysis. *Nat Biotechnol* **21**, 652-659 (2003).

831   67.   L. Ciani *et al.*, Wnt signalling tunes neurotransmitter release by directly targeting
832   Synaptotagmin-1. *Nat Commun* **6**, 8302 (2015).
833

835 **TABLES**

837 *Table 1: Regulation of clathrin lattice properties by CLC isoforms*

	high					low
Lattice curvature (pentagon formation)	CLCb	CLCa	nCLCa/b	CLCa/b	nCLCa	nCLCb
Lattice quality	CLCb	CLCa	nCLCa/b	CLCa/b	nCLCa	nCLCb
Budding efficiency	CLCb	CLCa	nCLCa/b	CLCa/b	nCLCa	nCLCb

    FIGURE LEGENDS

Fig. 1: CLCs differentially affect lattice curvature via regulation of the CHC knee. **a** Pucker

(1) and knee (2) angles of the clathrin triskelion (black: CHC, blue: CLC) dictate lattice

architecture. Different knee angles (encircled, black straight and dashed lines) are adopted

for hexagon (non-curvature inducing) or pentagon (curvature inducing) formation (3, whole

triskelion in black and parts of others in grey). Lattice curvature is further amended through

changes in pucker angle (4, dashed and black lines), or changes in proximal leg-crossing

angle (5). CLC subunits are omitted for simplicity. Different sizes of closed cages are

achieved by varying numbers of hexagons and a fixed number of 12 pentagons (filled black,

based on (44)). **b** Quantification of the percentage of small cages (< 90 nm in diameter) of >

200 cages, generated by *in vitro* assembly of clathrin reconstituted with indicated CLCs, 1:1

mixtures of reconstituted clathrins (CLCa/b or nCLCa/b) or CHC only (CHC), determined

from electron micrographs (mean ± SEM, *$P < 0.05$, ***$P < 0.001$ one-way ANOVA followed

by Holm-Sidak correction for multiple comparison, n = 3). **c** Cage size distributions from

assemblies as described in **b**. **d** Size distributions of cages formed from TDD (black) and

Hub (grey) co-assemblies, with Hub fragments reconstituted or not with the different CLC

isoforms or 1:1 mixtures thereof. **e** Representative EM images of cages formed from full-

length CHC (top) or Hub/TDD fragments (bottom) without or following reconstitution with CLC

isoforms. Scale bar: 100 nm.

Fig. 2: CLC splicing and mixing affects *in vitro* lattice properties. **a** Clathrin reconstituted with

indicated CLC isoforms and mixtures thereof were assembled into flat lattices and visualised

by negative stain for EM analysis. Scale bar: 200 nm. **b** Quality (regularity quantified by

Fourier transform) of lattices generated as in **a**, (mean ± SEM, *$P < 0.05$, **$P < 0.01$, one-

way ANOVA followed by Holm-Sidak correction for multiple comparison, n = 5). **c** Differences

in lattice quality between indicated reconstituted clathrins in individual experiments as in **b**

(data points and mean $\pm$ SEM, n = 5).

Fig. 3: CLC splicing and mixing affects *in vitro* lattice budding efficiency. **a** Representative

EM images of clathrin reconstituted with indicated CLC isoforms, 1:1 mixtures thereof, CHC

only (CHC) and tissue-derived clathrin (native) assemblies on H_6-ΔENTH-epsin[144-575]-coated

liposomes. Scale bar: 200 nm. **b** Quantification of the percentage of curved lattices (defined

as coats with < 200 nm fitted diameter) of > 60 000 nm total coat profiles generated as in **a**

(mean $\pm$ SEM, *P < 0.05, **P < 0.01, paired Student's t-test for native and CHC, n = 3, one-

way ANOVA with repeated measures for CLC-reconstituted clathrin, n = 4). **c** Differences in

percentage of curved lattices between indicated reconstituted clathrins in individual

experiments in **b** (data points and mean $\pm$ SEM, n = 4). **d** Quantification of budding efficiency

(% curved lattices determined as in **b**, normalised to native) of reconstituted clathrins and

mixtures thereof (*P < 0.05, one-way ANOVA for sets of neuronal or non-neuronal samples,

followed by Holm-Sidak correction for multiple comparison, n = 3). **e** Parameters

characterising coated buds (top); coat length *c*, membrane bud diameter *d*, neck width *n*, and

budding angle θ. Models of clathrin-mediated membrane deformation by transitional

curvature generation and lattice rearrangement (middle) or lattice growth under constant

lattice curvature (bottom). **f** Representative coat profiles for flat (left), curved (middle) and

budded (right) structures. **g** to **i** Analysis of the dataset generated as shown in **a** for bud

parameters *d* (**g**), *c* (**h**) and *n* (**i**) as shown in **e** in relation to the θ of each structure measured

for all reconstituted clathrins within the same experiment. Inserts in **g** and **h** show the

different correlations of these parameters as predicted according to the curvature transition

(red, straight line) constant curvature models (red, dashed line) shown in **e**.

Fig. 4: CLC composition regulates SV pool replenishment in hippocampal neurons. **a**

Immunoblot of CLC KO and WT brain lysate for CHC (TD.1 antibody), CLCa (X16 antibody),

CLCb (CLTB antibody) and actin (anti-beta-actin antibody). **b** Schematic illustration of

stimulating and recording electrode setup in hippocampal slices. The synapses investigated

are formed by a Schaffer-collateral (SC) axon (black) from a pyramidal neuron (black

triangles, cell bodies) in the CA3 region synapsing with a receiving pyramidal neuron (grey

triangle, cell body) in the CA1 region from which responses are recorded. Black circles

denote dentate gyrus (DG) granule cell bodies. **c** Representative traces of EPSCs following a

20 Hz electrical stimulation for 3 s in CLCa KO and WT hippocampal slices (left). Graphs

(right) show the mean ($\pm$ SEM) initial fusion efficiency, recycling rate and readily releasable

pool (RRP) size calculated for all cells (n = 9-11 slices from 3 animals/genotype; *P < 0.05

and **P < 0.01, unpaired Student's t-test). **d** Representative traces of EPSCs following a 20

Hz electrical stimulation for 3 s in CLCb KO and WT hippocampus slices (left). Graphs (right)

show the mean ($\pm$ SEM) initial fusion efficiency, recycling rate and RRP size calculated for all

cells (n = 11-12 slices from 4 animals/genotype; *P < 0.05, unpaired Student's t-test). **e**

Representative EM images of excitatory synapses in the CA1 region of the hippocampus of

CLCa WT, CLCa KO, CLCb WT and CLCb KO. Scale bars; 300 nm. **f** to **m** Quantification of

data extracted from EM images as in **e**. Graphs show SV density within 300 nm of the PSD

(**e** and **i**), cumulative frequency distribution of SV size (**g** and **k**), postsynaptic density (PSD)

length (**h** and **l**) and synapse density (**i** and **m**) expressed as mean $\pm$ SEM (*P < 0.05,

unpaired Student's t-test, n = 3).

Fig. 5: Neuronal defects are compensated for in CLCb KO but not CLCa KO mice. **a** Input-

output relationship of evoked excitatory postsynaptic currents (EPSCs) at CA1 synapses of

WT and CLCa KO hippocampus slices. Traces show responses of representative cells at

increasing stimulus intensity with an average of 3 responses for each stimulus strength

(mean $\pm$ SEM; n = 9-11 slices from 3 animals/genotype; *P < 0.05, two-way ANOVA with

repeated measures). **b** Input-output relationship of evoked EPSCs at CA1 synapses of WT

and CLCb KO hippocampus slices. Traces show response of representative cells at

increasing stimulus intensity with an average of 3 responses for each stimulus strength

(mean $\pm$ SEM; n = 12-15 slices from 3 animals/genotype; $*P < 0.05$, two-way ANOVA with

repeated measures). **c** Paired-pulse ratio (PPR) of evoked EPSCs at CA1 synapses of WT

and CLCa KO hippocampal slices. Traces show responses of representative cells. Graph

displays the mean PPR ($P_2/P_1 \pm$ SEM) from all cells (n = 9-11 slices from 3

animals/genotype) at different time intervals between paired pulses ($***P < 0.001$, unpaired

Student's t-test). **d** PPR of EPSCs at CA1 synapses of WT and CLCb KO hippocampal

slices. Traces show responses of representative cells. Graph displays the mean PPR (P_2/P_1

$\pm$ SEM) from all cells (n = 12-15 slices from 4 animals/genotype) at different time intervals

between paired pulses. **e** Genotype distribution for CLCa KO (n = 590) and CLCb KO (n =

359) mice after weaning compared to expected distribution ($****P < 0.0001$, *Chi*-square test).

f Performance in the accelerated rotarod test (latency to fall) by CLC KO (CLCa KO = 8,

CLCb KO = 8) and control wild-type littermates (CLCa WT = 8, CLCb WT = 7) expressed as

data points and mean $\pm$ SEM ($****P < 0.0001$, unpaired Student's t-test). **g** Time to cross grid

for CLCa KO (n = 10) and control mice (n = 10) expressed as data points and mean $\pm$ SEM

($*P < 0.05$, unpaired Student's t-test). **h** Grip strength of CLCa KO (n = 10) and control mice

(n = 11, gram-force relative grip strength over body weight) expressed as data points and

mean $\pm$ SEM.

Fig. 6: Model of how neuronal CLC diversity regulates synaptic vesicle recycling and lattice

properties. **a** CLC (cyan) neuronal splice inserts (arrowheads) are located near the TxD of

the bound CHC (green, left, PDB: 3LVH). **b** CLC (cyan) neuronal splice inserts (stars) are

near the CHC knee of the neighbouring triskelion (red or orange) within lattices (right, PDB:

3IYV). Interaction between a neuronal splicing insert and the knee of an adjacent triskelion

and/or adjacent CLC N-terminus (N) could promote conformational change in the CHC knee.

As the formation of pentagons and hexagons requires different knee angles, this interaction

could consequently influence lattice curvature (see Fig. 1a). **c** In WT mice (centre), a mix of

nCLCa and nCLCb clathrin creates the appropriate biophysical properties to mediate SV

generation from endosomal compartments (and possibly the plasma membrane, blue-green

arrows). Loss of nCLCa (left) creates nCLCb clathrin lattices that are defective in efficient SV

regeneration (blue arrows), resulting in decimated SV pools. Loss of nCLCb (right) creates

nCLCa clathrin lattices, which are also less efficient in SV regeneration than WT, but able

maintain an overall increased SV pool by excess budding to compensate for reduced acute

SV pool replenishment efficiency (green arrows).

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6