Host gill attachment enables blood-feeding by the salmon louse (*Lepeophtheirus salmonis*) chalimus larvae and alters parasite development and transcriptome

Erna Irene Heggland[1], Michael Dondrup[2], Frank Nilsen[1] & Christiane Eichner[1]

[1]Department of Biological Sciences & Sea Lice Research Centre (SLRC), University of Bergen, Norway

[2]Department of Informatics & Sea Lice Research Centre (SLRC), University of Bergen, Norway

KEYWORDS

Salmon louse; ectoparasite; blood-feeding; hematophagy; gills; RNA-sequencing

ABSTRACT

Blood-feeding is a common strategy among parasitizing arthropods, including the

ectoparasitic salmon louse (*Lepeophtheirus salmonis*), feeding off its salmon host's skin and

blood. Blood is rich in nutrients, among these iron and heme. These are essential molecules

for the louse, yet their oxidative properties render them toxic to cells if not handled properly.

Blood-feeding might therefore alter parasite gene expression. We infected Atlantic salmon

with salmon louse copepodids and sampled the lice in two different experiments at day 10 and

18 post infestation. Parasite development and presence of host blood in their intestines were

determined. We find that lice start feeding on blood when becoming mobile preadults if

sitting on the fish body, however they may initiate in blood-feeding at the chalimus I stage if

attached to gills. Lice attached to gills develop at a slower rate. Lice of similar instar age from

gills versus lice from skin epidermis were analyzed for gene expression by RNA-sequencing

in samples taken at day 10 for both experiments and at day 18 for one of the experiments. By

differential expression analysis, we found 355 transcripts elevated in lice sampled from gills

and 202 transcripts elevated in lice sampled from skin consistent in all experiments. Genes

annotated with "peptidase activity" are among the ones elevated in lice sampled from gills,

while in the other group genes annotated with "phosphorylation" and "phosphatase" is

pervasive. Transcripts elevated in lice sampled from gills are often genes relatively highly

expressed in the louse intestine compared with other tissues, while this was not the case for

transcripts found elevated in lice sampled from skin. In both groups, more than half the

transcripts are from genes higher expressed after attachment. In conclusion, blood-feeding

results in an alteration in gene expression, and a premature onset of blood-feeding likely

causes the parasite to develop at a slower pace.

INTRODUCTION

The salmon louse, *Lepeophtheirus salmonis* (Krøyer, 1837) (Crustacea: Caligidae) and its Atlantic subspecies *L. salmonis salmonis* (Skern-Mauritzen et al., 2014), is an obligate ectoparasite of salmonid fish, such as the Atlantic salmon (*Salmo salar*). The parasite is of major concern for the aquaculture sector in the Northern Hemisphere, as it causes challenges for the industry with its high fecundity and resistance towards several chemotherapeutants (Aaen et al., 2015). The parasite life cycle consists of both planktonic and parasitic stages (Hamre et al., 2013; Johnson and Albright, 1991a). Upon hatching from a fertilized egg, the parasite is in the nauplius I stage. Thereafter, the salmon louse molts into the nauplius II stage, and further to the infective copepodid stage. Successive molting occurs on the host, first to the parasitic chalimus I and II. These stages are attached to the host by their elongated frontal filament (Bron et al., 1991; Gonzalez-Alanis et al., 2001), and are therefore immobile. Another molting renders the parasite mobile, as it is no longer secured by the frontal filament, but holds itself by using its cephalothorax as a suction cup. These stages are the preadult I and II and adult lice. Now, the parasite grazes on larger parts of its host, selecting its preferred feeding site and is causing greater damage to the fish (Bjørn and Finstad, 1998; Grimnes and Jakobsen, 1996). Progression of the salmon louse life cycle is temperature dependent, and at 10 °C, the time from fertilization to mature adult is approximately 40 (male) to 52 (female) days (Johnson and Albright, 1991b), or 38 (male) to 44 (female) days for the fastest developers (Hamre et al., 2019).

The alimentary canal of the salmon louse develops during the copepodid stage (Bron et al., 1993). The alimentary canal is composed of a mouthpart, an esophagus, a midgut, and a hindgut ending in a short rectum (Bron et al., 1993; Nylund et al., 1992). Vertebrate blood is a highly nutritious tissue fluid that is constantly renewed. Hematophagy (blood-feeding habit) is therefore a common strategy among parasitizing arthropods. The salmon louse diet is

considered to consist of the skin and blood of its host (Brandal et al., 1976), and the blood-

filled intestine is visible as a red line throughout the salmon louse body. Upon ingestion of

blood, hematophagous parasites need to express genes encoding proteins that can manage the

blood components. Blood is particularly enriched in proteins that contain the pro-oxidant

molecules heme and iron. These are essential cofactors for the salmon louse, yet also highly

toxic if not bound and detoxified by chaperones. Therefore, the alimentary canal needs to

withstand, digest and absorb components of the food bolus. Trypsin-like enzymes (Johnson et

al., 2002; Kvamme et al., 2004), a lipid transfer protein (Khan et al., 2017), a putative heme

scavenger receptor (Heggland et al., 2019a) and the iron storage units of ferritin (Heggland et

al., 2019b) are all expressed in the salmon louse midgut.

The distribution of copepodids on wild and farmed hosts shows that the preferred settlement

site is on the fins and scaled body of the host (Bron et al., 1991). Some groups have reported

the settlement of lice on gills as well, however this is considered rather uncommon (reviewed

by Treasurer and Wadsworth (2004)). In laboratory trials, on the other hand, lice are often

found on gills, although there still seems to be a higher preference for the fins and body

(Bjørn and Finstad, 1998; Treasurer and Wadsworth, 2004). Copepodid gill settlement is

therefore often considered an experimental artefact due to an altered host behavior during

laboratory infestations (Treasurer and Wadsworth, 2004). Gill tissue in teleost fish is highly

vascular, whereas skin epidermis is not. The chalimus frontal filament, appendages and mouth

tube have been shown to not breach the basement membrane within the salmon skin (Jones et

al., 1990), thus not reaching the dermal vascular layer. Salmon lice settling on gills might

therefore be more prone to ingest a blood meal than those lice elsewhere on the host during

early stages of attachment.

The genome of the Atlantic salmon louse is fully sequenced and high-throughput

transcriptomics studies have been conducted under various experimental conditions using

microarrays as well as sequencing. Examples of such experimental settings include host-parasite interactions on different hosts (Braden et al., 2017), hosts fed different diets (Sutherland et al., 2017), response to drugs (Sutherland et al., 2014), larval stress response (Sutherland et al., 2012), parasite sex differences (Poley et al., 2016), and development (Eichner et al., 2008). Recently, we have used RNA-sequencing (RNA-seq) to investigate patterns of gene expression during molting in parasitic larval stages of *L. salmonis* (Eichner et al., 2018). Transcriptome plasticity in response to hematophagy has been investigated in various arthropods for which controlled blood-feeding is possible. Arthropod species subjected to such controlled feeding trials include mosquitos (*Aedes* species (Bonizzoni et al., 2011; Bottino-Rojas et al., 2015; Huang et al., 2015), *Anopheles gambiae* (Marinotti et al., 2005)), the biting midge *Culicoides sonorensis* (Nayduch et al., 2014), and ticks (*Ixodes* species (Kotsyfakis et al., 2015; Perner et al., 2016). However, investigating transcriptional changes induced by a blood meal within the salmon louse is challenging, as no protocol for feeding lice *in vitro* exists. To overcome this limitation, equally developed lice of the same batch, infecting the same fish, were sampled from host body attachment sites with predicted differing access to blood.

In this study, we infected Atlantic salmon with salmon louse copepodids and sampled the lice on the 10th and 18th day post infestation (dpi), when the lice were in the chalimus I and chalimus II stage or had recently molted to the preadult I stage. Parasite settlement site and visible presence of host blood in louse intestines were determined. Transcriptomes of equally developed lice sampled from different locations (gills and the body/fins), representing lice with access to blood versus lice without access at 10 and 18 dpi, were examined by RNA-sequencing. Specific aims of this study were to investigate i) visible blood ingestion from various sampling locations, ii) development of lice from locations differing in blood access,

and iii) differences in gene expression of immobile lice from locations with unequal access to

blood.

MATERIAL AND METHODS

Animals

Atlantic salmon lice (*L. salmonis salmonis*) (Skern-Mauritzen et al., 2014) were raised on

Atlantic salmon in tanks with seawater (salinity 34.5‰ and temperature 10 °C) (Hamre et al.,

2009). A laboratory strain of *L. salmonis* called LsGulen (Hamre et al., 2009) was used. Fish

were daily handfed commercial dry pellets and maintained according to Norwegian animal

welfare regulations. Experiments conducted herein were approved by the governmental

Norwegian Animal Research Authority (ID7704, no 2010/245410). Fish were anesthetized by

a mixture of methomidate (5 mg/l) and benzocaine (60 mg/l) prior to handling. For sampling

of early developmental stages of lice, fish were killed by a swift blow to the head. Salmon

louse egg string pairs were incubated and hatched in incubators in a seawater flow through

system (Hamre et al., 2009). Emerging copepodids were used to infect fish in 500-liter tanks.

Copepodids between 4 and 14 days post hatching were used. A total of 34 or 37 fish in the

two experiments respectively were infected with about 70 copepodids per fish. The amount of

copepodids used was estimated as described by Hamre et al. (2009). Prior to infestation, the

tank water was lowered and copepodids spread on the surface.

Sampling of lice

At 10 and 18 dpi, fish were sacrificed and lice were removed with forceps and photographed.

The gills were cut out and observed under a microscope. Any lice present were sampled,

photographed and placed on RNAlater in individual tubes. Measurements of lice were done

on photographs. Total length (TL) and cephalothorax length (CL) were measured as earlier

described (Eichner et al., 2018, 2015) and determination of the developmental status as well

as sex differentiation for chalimus II larvae was done using TL and CL measurements as

described previously (Eichner et al., 2018, 2015). For RNA isolation prior to RNA-seq, lice

were sorted into groups of equal developmental status within each group as described by

Eichner et al. (2018), and between groups from different sampling locations (gills, skin). Five

chalimus I or four chalimus II lice respectively were pooled together in one sample. Eight

(Ex1 10 dpi), six (Ex2 10 dpi) or five (Ex2 18 dpi) replicates were sampled per group (lice

sampled from gills, lice sampled from skin). RNA from both experiments sampled at 10 dpi

and from one experiment sampled at 18 dpi was analyzed by RNA-seq.

RNA isolation and sequencing

RNA was isolated as described before (Eichner et al., 2014). In brief, pools of four or five

chalimus larvae were homogenized in TRI reagent and mixed with chloroform (both Merck).

The water phase was withdrawn and further used with the RNeasy micro kit (Qiagen) for

RNA isolation according to the manufacturer's instructions. RNA was stored at -80 °C until

being used. Library preparation and RNA-sequencing were conducted by the Norwegian

Sequencing Centre, Oslo as previously described (Eichner et al., 2018). Briefly, sequencing

libraries were prepared from 0.5 µg total RNA using the TruSeq stranded mRNA reagents

(Illumina). Indexed libraries were blended into a single pool and sequenced during three runs

of a NextSeq 500 instrument (Illumina) using 76 base pair (bp) single end reads. Image

analysis and base calling were performed using Illumina's RTA software version 2.4.11, and

data were converted to fastq format using bcl2fastq version 2.17.1.14. Raw sequencing data

has been deposited in the NCBI database under BioProject ID PRJNA577842.

Data processing of RNA-seqencing data

Obtained sequences were quality controlled by FastQC v0.11.5 (Andrews, 2016). Reports

were summarized using MultiQC 1.0 (Ewels et al., 2016). As reference genomes, we used a

combination of the Ensembl Metazoa reference assembly of the nuclear genome (LSalAtl2s,

http://metazoa.ensembl.org/Lepeophtheirus_salmonis) and the mitochondrial genome RefSeq

sequence NC_007215.1 (Tjensvoll et al., 2005). The gene models from Ensembl Metazoa

were further augmented with gene models derived from full-length sequences of LsFer1 and

LsFer4 obtained by rapid amplification of cDNA ends (RACE) (Heggland et al., 2019b), by

aligning the RACE consensus sequences against the nuclear assembly with GMAP (Wu and

Watanabe, 2005). RNA-seq reads were aligned against the reference using the STAR aligner

(Dobin et al., 2013). Then, alignments were sorted and indexed using SAM-tools (Li et al.,

2009) and saved in BAM format. Technical replicates were merged prior to counting using

the merge function in SAM-tools. RNA-seq reads and their overlap with annotated nuclear

and mitochondrial transcripts were counted using the software featureCounts (Liao et al.,

2014) with settings for strand-specific reverse stranded libraries.

Differential expression (DE) analysis was done with DESeq2 (Love et al., 2014) on raw

counts using Galaxy (Giardine et al., 2005) under the Norwegian e-Infrastructure for Life

Sciences (NeLS) platform (Tekle et al., 2018). Prior to DE analysis, all transcripts with less

than four counts in all samples were removed. Venn diagrams were prepared using the

BioVenn platform (http://www.biovenn.nl/) (Hulsen et al., 2008). Hierarchical clustering as

well as GO annotation enrichment were performed in J-Express (Dysvik and Jonassen, 2001;

Stavrum et al., 2008). GO terms were summarized in REVIGO (Supek et al., 2011).

Transcript annotation

Protein-coding transcripts were annotated by running NCBI-Blast+, BlastP version 2.6.0+

(Altschul et al., 1990; Camacho et al., 2009) of their corresponding predicted Ensembl protein

sequences against the GenBank (NR) (https://www.ncbi.nlm.nih.gov/) and SwissProt

(Bairoch and Apweiler, 2000) databases. GO terms (full terms and GOslim annotation) and

protein families (Pfam) were automatically assigned by InterProScan 5 (Jones et al., 2014).

RESULTS

Distribution and characteristics of lice

Upon termination all lice were removed from the salmon, and their settlement site and

developmental stage and the visibility of an intestine filled with blood were assessed. Figure 1

depicts the distribution of different developmental stages and instar ages of lice on the host at

10 and 18 dpi. At 10 dpi, most lice are attached to the fins, but there is also a high proportion

of lice on the body and gills. At 18 dpi, however, the highest proportion of lice is found on the

body of the salmon. Different stages are distributed differently between sites.

At 10 dpi, the highest proportion of lice is in the chalimus I stage (83% and 95% in Ex1 and

Ex2 respectively) while the remaining lice are chalimus II. The chalimus II larvae (mainly

males) are found on the body and fins, but none on gills. Here, we rather find chalimus I

(higher proportion of middle than old) larvae (Figure 1, Table 1).

At 18 dpi, we find mostly preadult I lice (75% or 74% in Ex1 and Ex2 respectively). On the

body and fins, most lice were at the preadult I stage, while on the gills, most lice were in the

chalimus II stage. Preadult I females are found on the body and fins, but nearly none on the

gills (<1%). On gills, we rather find a higher proportion of old chalimus II females. There is a

higher proportion of preadult I male lice found on the body compared with the fins. On the

fins, most of the preadult I lice are young females. (Figure 1, Table 1)

For preadult I lice from 10 or 9 fish in Ex1 and Ex2 respectively also the presence of a frontal

filament was investigated. Presence of a frontal filament and a visible blood filled intestine

with respect to the settlement site are summarized in Table 2. The majority of the preadult I

lice were located on the host body, and the minority was located on the gills. The preadult I

lice on the gills, however, were more often secured by their frontal filament. Of the preadult I

lice still attached by the filament, only the ones on the gills had a blood-filled intestine (Figure

2c). None of the lice on the fins had a blood-filled intestine, and on the host body, only the

mobile lice had apparently fed on blood. Additionally, we found both chalimus I (Figure 2a)

(at 10 dpi) and chalimus II (Figure 2b) larvae attached to the gills that had fed on blood,

whereas lice of the same age on the fins and body had no visible blood in the intestine.

Effect of gill settlement on gene expression in chalimus larvae

In order to determine the effect of gill settlement on the gene expression in chalimus larvae,

RNA-sequencing of pooled individuals of equal development was performed. All counts per

million (CPM) values can be found in Supplementary Table S1. The overall gene expression

of the individual samples in comparison with chalimus I and chalimus II larvae of different

instar age (data taken from Eichner et al. 2018) is shown in a correspondence analysis (CA)

plot (Figure 3c).

All lice from this study sampled at 10 dpi are clustering together with chalimus I larvae

sampled directly before molting as well as molting ones (old, molt) from Eichner et al. (2018)

and all lice sampled at 18 dpi from this study are clustering with chalimus II lice sampled

directly before molting (Eichner et al., 2018). Lice sampled from gills and lice sampled from

skin differed also slightly in their overall gene expression. Lice from Ex1 at 10 dpi cluster

together with lice from the respective group (from gills or from skin) of Ex2, showing lice

sampled at 10 dpi in the different experiments are composed of comparable batches of lice.

DE analyses were performed for each sampling separately. MA plots as well as a principle

component analysis (PCA) for each sampling are shown in Figure 3a and b. A list of all genes

with log2 fold changes and false discovery rate (FDR) adjusted p-values (padj) for each

sampling can be found in Supplementary Table S2. 5878 genes were differentially expressed

in at least one of the samplings (Supplementary Table S3).

Most DE genes were found in Experiment 1, 10 dpi (2188 or 2015 up-regulated in gill or skin

samples respectively). In Experiment 2, 10 dpi only 1112 or 1081 transcripts respectively

were found, of which 79% or 68% respectively are overlapping with the ones found in Ex1 10

dpi. DE genes found at 18 dpi are less overlapping with DE genes found in Ex1 at 10 dpi.

Only 43% or 32% respectively of the genes found here are overlapping with genes from the

respective groups in Ex1 10 dpi and 35% or 28% respectively are overlapping with Ex2 10

dpi (Figure 4). 616 genes are DE in all three samplings, of which 355 are elevated in lice

samples from gills and 202 are elevated in lice samples from skin (Supplementary Table S6

and S7), while 59 are significant different, but regulation directions are differing between time

points (31 elevated in lice from gills at 10 dpi but lower at 18 dpi, 24 the other way around

and 4 are differing between lice sampled at day 10 of the two different experiments)

(Supplementary Table S8). Transcripts solely regulated at either 10 dpi or 18 dpi are listed in

Supplementary Table S4 and S5, respectively.

Among DE genes found at 10 dpi or at 18 dpi we found a high number (51) of genes

annotated with Pfam domain PF00040: "Fibronectin type II domain". Mostly, these are

elevated in lice sampled from gills at 10 dpi. However, a smaller number of PF00040 is under

the DE genes which are elevated in lice sampled from skin than from gills. Additionally, 85

transcripts with Pfam domain PF00089: "Trypsin" were under the DE genes and are mostly

found in the group of lice sampled from gills.

The sizes and overlaps of gene sets that were DE in each experiment separated by expression

pattern (elevated in lice sampled from gills or elevated in lice sampled from skin respectively)

are depicted in Venn diagrams together with most representative GO terms in Figure 4. All

significantly enriched GO terms are listed in Supplementary Table S9. Additional

summarized GO annotations belonging to biological process are visualized in a TreeMap

(REVIGO) (Supplementary Figure S1). "Peptidase activity" is an enriched GO term in lice

sampled from gills across all groups (except the ones exclusively found in Ex2 10 dpi), and in

particular "serine type endopeptidase activity", whereas "serine-type endopeptidase inhibitor

activity" is enriched in lice sampled from skin. Notably, "glycolysis" as well as

"oxidoreductase activity" are GO terms highly enriched in genes elevated in lice sampled

from gills. GO terms containing "phosphorylation" as well as "phosphatase" are found

enriched in nearly all groups in genes elevated in lice sampled from skin (also here, the

exception is the ones exclusively found in Ex2 10 dpi).

Equal gene expression changes throughout all three analyses

To determine which genes may be important in relation to the blood meal in general,

independent of the stage of the lice at the different time points, we investigated the transcripts

which were either significantly elevated in lice sampled from gills or significantly elevated in

lice sampled from skin in all three samplings (Ex1 10 dpi, Ex2 10 dpi, Ex2 18 dpi). We found

355 transcripts that were elevated in all three samplings in lice from gills, and 202 from skin.

Of the 355 genes elevated in lice sampled from gills, 60% had predicted Pfam domains, and

of the 202 elevated in lice sampled from skin, 82% had predicted Pfam domains. A highly

prevalent Pfam domain in the DE genes found in all 3 samplings elevated in lice sampled

from gills is PF00089: Trypsin. Other more frequent found domains are PF01400: Astacin

(Peptidase family M12A), PF02469: Fasciclin domain, PF05649: Peptidase family M13,

PF00171: Aldehyde dehydrogenase family, as well as different Zinc finger domains. In the

group of DE genes which were elevated in lice sampled from skin, prevalent domains are

PF00040: Fibronectin type II domain; PF00069: Protein kinase domain, PF00096: Zinc

finger, PF00135: Carboxylesterase family, PF01391: Collagen triple helix repeat. A full list

can be found in Supplementary Table S6 and S7. GOslim was used to minimize GO

categories. DE genes in lice sampled from gills fall under fewer GOslim categories than DE

genes elevated in lice from skin, even though there are more genes in former DE group

(Figure 5). Often, genes in the group found elevated in lice sampled from gills fall under the

enriched GO group "catalytic activity" (all are shown in Figure 5a). Remarkably strong

enriched (factor 30) DE genes elevated in lice from skin, are genes belonging to "extracellular

matrix". However, only five genes are in that group. More than 30 genes were found in GO

categories "catalytic activity", "hydrolase activity", "binding" and "ion binding" (Figure 5b).

All enriched GOslim terms, numbers of genes found in each category, and enrichment factor

for the two groups are shown in Figure 5.

42% of the 355 genes elevated in lice sampled from gills have a fold-change more than two

compared to skin. When also taking the p-value into account we find 21%, which are strongly

regulated (average fold-change over two and average adjusted p-value ≤ 0.005). In this group,

the strongest elevated transcript is most similar to a nematode astacin

(EMLSAT00000010457). Among the other strong ones regulated are three transcripts with

FNII domains, another transcript with an astacin domain, as well as nine transcripts with

trypsin domains, as well as a chemosensory protein (EMLSAT00000005105), yet also many

transcripts with no annotation or known protein domains. There are several transcripts with

similarity to various proteases that are up-regulated in lice from gills. All genes are listed in

Supplementary Table S6. The expression patterns of the ten strongest regulated genes over all

samples are shown in Supplementary Figure S2a.

In the group of genes elevated in lice sampled from skin, we find fewer genes highly

differentially expressed than in lice sampled from gills. Only 16% have a fold-change of two

or higher compared to samples from gills. Most up-regulated in this group is a transcript with

no predicted annotation or Pfam domains (EMLSAT00000009920). Among the strongest

elevated genes in all three samplings (average fold-change over two and p-value $\leq$ 0.005) in

lice sampled from skin are several transcripts with predicted FNII domains. Moreover, we

find several genes with no annotation or known protein domains. All genes are listed in

Supplementary Table S7. Expression patterns for the ten strongest regulated ones are shown

in Supplementary Figure S2b.

We further looked at the expression profile of these detected DE genes during the course of

development, as well as in various tissues (Edvardsen et al., 2014; Eichner et al., 2018;

http://licebase.org) . We were particularly interested in determining if these transcripts were

also elevated in the louse intestine compared with other tissues, or if these transcripts are up-

or down-regulated after attachment or after molting to preadult, the expected time point for

accessing host blood. Of the 202 genes elevated in lice from the skin, 124 were also

investigated in an oligo microarray regarding to expression in different tissues (gut adult

female, gut adult male, ovaries, testis, subcuticular tissue and brain) and from the 355 genes

elevated in lice sampled from gills 209 were represented in that study (Edvardsen et al.,

2014).

Among transcripts elevated in lice sampled from gills, 94 (26%) are higher expressed in the

intestine (only 5% of transcripts were lowest in intestine compared with other tissues

investigated) (LiceBase). Moreover, 39 of these were very highly expressed in the intestine,

compared to other tissues (more than 100 times higher) (LiceBase). 77 of these elevated in

intestine were also analyzed in the microarray study investigating different tissues of *L.*

salmonis and 52 were also found highest expressed in the intestine there (Edvardsen et al.,

2014). 66% of the transcripts are elevated after attachment and 55% are higher expressed in

preadult lice than in chalimus II when comparing to the time series data (Eichner et al., 2018).

The genes higher expressed in intestine, higher expressed after attachment or higher expressed

in pad1 respectively are marked in Supplementary Table S7.

Only 17 (8%) of the transcripts in the DE gene group elevated in lice sampled from skin are

higher expressed in the intestine than in other investigated tissues (12% lowest of all tissues

investigated) (LiceBase). Only two were much higher (more than 100 times) expressed in

intestine than other tissues (LiceBase). However, the ten of these also found on the oligo

microarray were not highest expressed in the intestine, except one (EMLSAT00000008355)

(Edvardsen et al., 2014). Here two others were found highest expressed compared with the

other tissues analyzed. 57% of the transcripts are elevated after attachment. Nearly all of the

strongest regulated transcripts are elevated after attachment when comparing with LiceBase

data. 53% are higher expressed in preadult lice than chalimus II (Eichner et al., 2018). The

genes higher expressed in intestine, higher expressed after attachment or higher expressed in

pad1 respectively are marked in Supplementary Table S7.

DISCUSSION

In this study, we have investigated the biology of blood-feeding in the marine ectoparasitic

salmon louse with a special focus on gene expression of immobile lice situated on host gills.

We chose immobile lice, because this allowed us to focus on those individuals that had stayed

at one location at least since extruding the frontal filament in the late copepodid stage. Being

attached to the gills allowed the lice to initiate blood-feeding prior to becoming mobile.

Samples from two experiments terminated at 10 days post infestation, and one sample

terminated at 18 days post infestation of one the experiments were included in the RNA-seq

and subsequent gene expression analyses.

At 10 dpi, the amount of lice is rather evenly distributed between the investigated body parts

(32 or 29% on body, 43 or 39% on fins and 24 or 30% on gills in Ex1 and Ex2 respectively).

The favored site at 18 dpi is the body with 66 or 61% in Ex1 or Ex2 respectively (15 or 14%

on fins and 17 or 23% on gills) (Figure 1). Lice at day 10 are in chalimus I or chalimus II

stage and attached by the frontal filament. On the more closely analysed fish at day 18 (Table

2), we found 68 or 66% preadult lice in Ex1 or Ex2 respectively. Of these, 10 or 21%

respectively are attached with a filament, while the others are mobile and can freely move on

the fish. The finding that lice are differently distributed when being mobile rather than

attached, and mainly found on the body of the fish, suggests that the mobile preadult lice

choose the general body surface as a preferred feeding site and migrate there from host fins

and gills when becoming mobile. The majority (78 or 87% respectively) of the preadult I lice

on fins are females. Female lice are known to develop slower than males (Hamre et al., 2019,

2013), and this as well indicates that the lice tend to leave the fins for other host feeding areas

when becoming mobile. A preadult I louse that is still attached to its host by its frontal

filament has recently molted from the chalimus II stage, and has stayed at that feeding site

since attachment. There are no preadult I lice with a visible blood-filled intestine on the fins,

whereas we find this on the gills and the body. Interestingly, of the lice still on their filament,

only those on gills have apparently fed on blood. Moreover, already in the chalimus I stage,

we find lice with blood-filled guts on the gills, but not on any other feeding site. As the

preadult I lice on the body with a blood-filled intestine are mobile, these lice have either

started with blood-feeding in the mobile preadult I stage or are preadult lice migrated from the

gills, meaning that blood-feeding is initiated from the mobile preadult I stage and onwards in

the development of the salmon louse occurring under natural conditions.

Development of lice on the gills was delayed, compared to development of lice on body or

fins. At 10 dpi, no chalimus II lice were found and a higher proportion of chalimus I lice was

of less developed instar age on gills. Comparing only the attached chalimus at 18 dpi (25% of

all lice), 51 or 62% in Ex1 or 2 respectively are found on gills. In addition, on gills, there is a

higher proportion of male chalimus II lice, which develop faster than females. Developing on

host gills caused the salmon lice to develop slower than those developing on other locations.

There have been contradicting results about this in the past (Johnson, 1993; Johnson and

Albright, 1992), however here we have determined instar ages, and not only the

developmental stages, which adds more confidence to our results.

We conclude that during the normal development on the body or the fins, the salmon louse

does not start to feed on blood until reaching the mobile preadult I stage. By that reasoning,

we wanted to compare gene expression of chalimus larvae located on the vascular gills with

access to blood with that of chalimus larvae of equal development from the rest of the body.

The salmon louse has approximately 13,000 protein encoding genes

(http://metazoa.ensembl.org/Lepeophtheirus_salmonis), and we find in our RNA-seq analyses

that over 5800 genes had an altered expression in at least one of our samplings. As expected,

we found a higher amount of overlapping DE genes in the two samplings at 10 dpi. These are

chalimus I larvae, which are soon molting to chalimus II, while lice sampled at 18 dpi are

chalimus II larvae, shortly prior to molting to preadult I lice. As such, all lice are sampled at a

similar instar age. However, phenotype and lifestyle differ in preadult lice and one can expect

expression of genes in preparation for this stage in the lice sampled at 18 dpi. The high

amount of DE genes exclusively found in Ex1 10 dpi could be caused by batch differences

between Ex1 10 dpi and Ex2 10 dpi, or could be as a result of more powerful statistics due to

a higher amount of parallel samples (8 versus 6 biological parallels of each group in Ex1 10

dpi and Ex2 10 dpi, respectively). However, we know also that minor differences in

development have a high impact on gene expression (Eichner et al., 2018), and individual

differences occurring within groups, with possible consequences between groups, could bias

the results.

To investigate gene expression caused by nutrition differences, we mainly concentrated on the

DE genes found in all 3 samplings. Transcripts over-expressed in lice sampled from gills

could be important for hematophagy. However, many (70 of 74) of the strongest DE genes in

this group are not highest expressed in the intestine, but rather in other tissues, suggesting

these contribute to other functions in the louse that may be modified by hematophagy. Genes

elevated in lice from gills show a more homogenous GO annotation (fewer GOslim

categories) than the ones elevated in lice from skin, suggesting that several DE genes are

involved in the same processes. There are also more genes with a greater fold change within

the group of DE genes elevated in lice sampled from gills (42% over 2-fold change, whereas

only 16% in lice sampled from skin), pointing towards a high demand of these gene products

when feeding on blood. However, as GO terms can be unspecific or general, the following

discussion deals with selected groups of transcripts.

Iron and heme

Among the regulated transcripts, the iron storage units of ferritin (*LsFer1* (RACE sequence)

and *LsFer2*: EMLSAT00000006305) are both elevated in chalimus larvae sampled from gills

compared with other settlement sites (Supplementary Figure S3 a, b). We have previously

established that these genes are important for the adult female salmon louse blood-feeding and

reproductive success, as the parasite had a clear gut and failed to produce viable eggs upon

silencing the two genes (Heggland et al., 2019b). Blood contains several iron-proteins, and

when initiating blood-feeding, the salmon louse needs to obtain a way of storing and

detoxifying iron absorbed from the blood. Up-regulating ferritin when ingesting a blood meal

is therefore an important defense mechanism for a blood-feeding parasite. The putative heme

scavenger receptor, *LsHSCARB* (EMLSAT00000005382), is elevated in lice on gills at 18 dpi

compared to lice on skin (Supplementary Figure S3d). We recently found that upon silencing

LsHSCARB by RNA interference, adult female lice had absorbed less heme and produced

fewer viable eggs and less offspring (Heggland et al., 2019a). Lacking early (10 dpi)

transcriptional elevation of *LsHSCARB* could indicate alternative mechanisms of absorption

during the earlier developmental stages, or the existence of a post-transcriptional mode of

regulating the LsHSCARB protein. Alternatively, the lack of early regulation might serve to

maintain homeostasis of heme levels when feeding on the vascular gills.

Detoxification

A Glutathione S-transferase (GST) (PF02798) transcript (EMLSAT00000009830) was

elevated in lice on gills in all samplings. GSTs are major detoxification enzymes. A GST in

the hard tick *Ixodes ricinus* (IrGST1) (GenBank ID: MF984398) was also found to be

elevated in the midgut of blood-fed ticks compared with serum-fed ticks (Perner et al., 2016).

Further characterization of IrGST1 showed that it was heme-inducible and the recombinant

protein was able to bind heme *in vitro* (Perner et al., 2018). The authors speculated that

IrGST1 is important for detoxifying excess heme to avoid cytotoxicity in the tick (Perner et

al., 2018). Recombinant GSTX2 (GenBank ID: AAK64286.1) of *A. aegypti* also binds heme

(Lumjuan et al., 2007), and was elevated in a heme-incubated *A. aegypti* Aag2 cell line

(Bottino-Rojas et al., 2015). Of the six different predicted salmon louse proteins with the GST

domain (PF02798), EMLSAP00000009830 is the most similar to both IrGST1 and *A. aegypti*

GSTX2. The connection of GST and blood-feeding in the salmon louse is an interesting topic

for future studies, as we per today do not know what mechanisms the salmon louse depends

on to detoxify heme.

Digestion

Food protein hydrolysis is a fundamental step of digestion, and is mediated by peptidases that

enzymatically cleave peptide bonds. Blood is highly enriched in protein, and one of the most

abundant ones is the gas transporter hemoglobin. Investigating changes in the salmon louse

transcriptome upon initiating blood-feeding could thus give clues as to which enzymes are

essential for breakdown of blood components. Trypsin is a digestive enzyme belonging to the

S1A subfamily of serine endopeptidases, and five main trypsin-encoding transcripts in the

salmon louse intestine have previously been characterized (Johnson et al., 2002; Kvamme et

al., 2004). Trypsins and other proteins involved in protein degradation were found elevated

e.g. in blood-fed mosquito *A. aegypti* (Bonizzoni et al., 2011). Twenty-eight transcripts with

trypsin as the only predicted protein domain (PFAM: PF00089) (29 in total with trypsin +

other domains) were found to be elevated in lice on host gills at day 10 (Ex1 and 2) and 18

dpi. Of these, 11 are predicted to be highest expressed in the intestine compared with other

tissues investigated in the salmon louse (LiceBase; Edvardsen et al., 2014; Supplementary

Table S6). A heat map showing the expression patterns for all transcripts with trypsin

domains found DE in all 3 samplings in data taken from LiceBase and from the time-series

study are shown in a hierarchical cluster in Figure 6. *LsTryp1* (GenBank ID: AY294257, best

blast hit: EMLSAT00000004828) is elevated in all three samplings in lice on gills. One

transcript with a trypsin domain only (EMLSAT00000004988) is found to be elevated in lice

on host skin at both 10 dpi (Ex1 and Ex2) and at 18 dpi. RNA-seq data in LiceBase as well as

microarray data from Edvardsen et al (2014) however show that this transcript has a low

expression in the louse intestine and is rather expressed in antenna and feet (LiceBase) or

subcuticular tissue and brain (microarray). It might therefore be of importance for other

purposes than blood meal digestion.

Peptidases other than trypsins are also regulated in lice on host gills. There are 17 transcripts

with Pfam domains "peptidase" other than trypsins elevated in lice on gills in all samplings.

Among these are four transcripts with Astacin-domains (Peptidase family M12A) and five are

M13 peptidases (Figure 6). Both groups are metallopeptidases and are enriched in arthropods.

Astacin-like metallopeptidases are implicated in digestive processes, but are also reported to

have anticoagulative effects, as they are found to have fibrinogenolytic activity in spider

venoms (Trevisan-Silva et al., 2010). M13 metallopeptidases are widely distributed in

animals, and e.g. make up the major group of the hematophagous tick degradome (Mulenga

and Erikson, 2011). Furthermore, we also find many of the same types of peptidases elevated

rather in lice from skin (one with Astacin domain, two with Peptidase family M13 domain).

This could indicate different modes of digesting a blood meal versus digesting components of

ingested salmon skin. Further investigation into the elevated trypsins and other peptidases

expressed in the salmon louse gut should be conducted.

Putative anti-coagulation

Blood coagulation is a key mechanism in maintaining homeostasis in vertebrates if a blood

vessel were to rupture. A parasite feeding on vertebrate blood would therefore require

mechanisms in order to counteract this to maintain its feeding habit. Anti-coagulation factors

targeting host proteins could thus be vital for the successful blood-feeding in the parasitizing

arthropod. A thrombin (coagulation factor) inhibitor, hemalin, was found to be important to

avoid clotting of the blood meal in the bush tick *Haemaphysalis longicornis* (Liao et al.,

2009). A salmon louse transcript (EMLSAT00000003009) encoding two Kunitz/Bovine

pancreatic trypsin inhibitor domains (PF00014), as also found in the tick hemalin, was

elevated here in all three samplings in lice on gills. However, four other transcripts with the

same domain were elevated in lice from skin in all samplings (EMLSAT00000000152,

EMLSAT00000007907, EMLSAT00000008877 and EMLSAT00000009255).

We also find serine protease inhibitors (serpins, PF00079) regulated. From the 15 predicted

serpin transcripts in the louse, four were DE in all three samplings. Two were elevated in lice

on gills (EMLSAT00000010931 and EMLSAT00000001743), one elevated in lice on skin

(EMLSAT00000011353), while the last (EMLSAT00000005224) was expressed lower at 10

dpi, but elevated at 18 dpi in the lice sampled from gills. One transcript

(EMLSAT00000000552) was elevated in lice on gills at 10 dpi (Ex1 and Ex2). Anti-

coagulation factors could be targets for pest control as they are likely secreted and in contact

with the host, and thus probably vital for the host-parasite interaction.

Fibronectin type II

The approximately 60 amino acid long fibronectin type II (FNII) domain (PF00040) is a

protein domain found within the glycoprotein fibronectin. It contains four conserved cysteine

residues that form disulfide bridges. These residues are important for e.g. fibronectin's

collagen binding properties (Guidry et al., 1990). The FNII domain is also found within the

vertebrate blood coagulation protein Factor XII (McMullen and Fujikawa, 1985). The FNII

domain is the most expanded protein domain of the salmon louse with over 200 copies within

over 80 genes identified so far (http://metazoa.ensembl.org/Lepeophtheirus_salmonis). Some

of these genes (*LsFNII1*, *2* and *3*) have been characterized, and are expressed in tegumental

type 1 (teg1) glands of the salmon louse (Harasimczuk et al., 2018; Øvergård et al., 2016).

Teg1 glands are exocrine and their secretory ducts are extending to the dorsal and ventral side

of the salmon louse (Øvergård et al., 2016). The functions of FNII-containing proteins have

not been determined in the salmon louse, however it has been suggested that proteins with the

domain may be of importance for lubricating the integument and functioning as an anti-

fouling agent, or as part of the salmon louse fuzzy coat (acid mucopolysaccharide layer (Bron

et al., 2000)) (Harasimczuk et al., 2018). Genes with FNII domains expressed in teg1 glands

have also been suggested to be of importance for host immune modulation by the parasite

(Øvergård et al., 2016).

We find that several FNII-containing genes are significantly regulated in chalimus larvae in

our dataset. Five transcripts with predicted FNII domains are elevated in all three samplings in

lice on gills. Of these, all but one (EMLSAT00000011958) are predicted to be up-regulated

after louse attachment (http://licebase.org; Supplementary Table S6) (Figure 6). Seven

transcripts with FNII are rather found elevated in lice on skin in all three experiments. Here as

well, all but one transcript (EMLSAT00000006178) are up-regulated after louse attachment

(http://licebase.org; Supplementary Table S7). The FNII encoding genes characterized by

Øvergård et al. (2016) and Harasimczuk et al. (2018) are not among the transcripts regulated

in all samplings here (*LsFNII1*: EMLSAT00000012082, *LsFNII2*: EMLSAT00000007294,

LsFNII3: EMLSAT00000009744), however LsFNII1 was elevated in both experiments at 10

dpi, and LsFNII3 at 18 dpi as well as in Ex1 10 dpi. The transcripts elevated in lice on gills

should be further characterized, in order to elucidate a possible role of FNII in blood-feeding.

Given the earlier reports that FNII domains in vertebrates may be important for blood clotting,

one hypothesis is that proteins with FNII domains only could have an anti-coagulant effect.

CONCLUSIONS

Blood is a major dietary component for the ectoparasitic salmon louse, which the parasite has

access to when attached to a salmonid host. We find that the salmon louse initiates blood-

feeding during the mobile preadult I stage. However, if the parasite is attached to host gills, it

may start feeding on blood already at the chalimus I stage or even earlier. Blood can be found

even in copepodids sampled from gills (personal observation, Supplementary Figure S4). The

premature onset of blood-feeding caused lice on gills to develop at a slower pace than lice that

were attached to host fins and general body surfaces. Chalimus lice of equivalent age on gills

versus other attachment sites were therefore analyzed for gene expression comparisons.

Several genes are elevated in lice attached to the gills, and among these, we find e.g. genes of

importance for the absorption, storage and/or transportation of the pro-oxidative molecules

iron and heme, digestive and detoxification enzymes, genes that could be important for anti-

clotting of host blood, and several genes with FNII domains. The results of this study raise a

number of new gene targets to investigate further in order to elucidate the blood-feeding habit

of the infamous salmon louse.

ACKNOWLEDGEMENTS

This research has been funded by The Research Council of Norway, SFI-Sea Lice Research

Centre, grant number 203513/O30 and 226266. Further, this work was funded by the

ELIXIR2 (270068) infrastructure grant from the Research Council of Norway to MD. The

sequencing service was provided by the Norwegian Sequencing Centre

(www.sequencing.uio.no); a national technology platform hosted by the University of Oslo

and supported by the "Functional Genomics" and "Infrastructure" programs of the Research

Council of Norway and the South-Eastern Regional Health Authorities. We would like to

thank Lars Are Hamre and Per Gunnar Espedal for help in the wet-labs and Madeleine

Carruthers for helpful comments on the manuscript.

AVAILABILITY OF DATA

The datasets supporting the conclusions of this article are included within the article and its

additional files. Raw RNA-sequencing data files have been deposited to NCBI BioProject

under the accession number PRJNA577842. A preprint of this manuscript and its

supplementary files have been made publicly available on bioRxiv:

https://doi.org/10.1101/815316.

REFERENCES

Aaen, S.M., Helgesen, K.O., Bakke, M.J., Kaur, K., Horsberg, T.E., 2015. Drug resistance in sea lice:
a threat to salmonid aquaculture. Trends Parasitol. 31, 72–81.

Altschul, S.F., Gish, W., Miller, W., Myers, E.W., Lipman, D.J., 1990. Basic local alignment search
tool. J. Mol. Biol. 215, 403–410.

Andrews, S., 2016. FastQC: A quality control tool for high throughput sequence data
https://www.bioinformatics.babraham.ac.uk/projects/fastqc/

Bairoch, A., Apweiler, R., 2000. The SWISS-PROT protein sequence database and its supplement
TrEMBL in 2000. Nucleic Acids Res. 28, 45–48.

Bjørn, P.A., Finstad, B., 1998. The development of salmon lice (*Lepeophtheirus salmonis*) on
artificially infected post smolts of sea trout (*Salmo trutta*). Can. J. Zool. 76, 970–977.

Bonizzoni, M., Dunn, W.A., Campbell, C.L., Olson, K.E., Dimon, M.T., Marinotti, O., James, A.A.,
2011. RNA-seq analyses of blood-induced changes in gene expression in the mosquito vector
species, *Aedes aegypti*. BMC Genomics 12, 1–13. doi:10.1186/1471-2164-12-82

Bottino-Rojas, V., Talyuli, O.A.C., Jupatanakul, N., Sim, S., Dimopoulos, G., Venancio, T.M., Bahia,
A.C., Sorgine, M.H., Oliveira, P.L., Paiva-Silva, G.O., 2015. Heme signaling impacts global
gene expression, immunity and Dengue virus infectivity in *Aedes aegypti*. PLoS One 1–19.

Braden, L.M., Sutherland, B.J.G., Koop, B.F., Jones, S.R.M., 2017. Enhanced transcriptomic
responses in the Pacific salmon louse *Lepeophtheirus salmonis* oncorhynchi to the non-native
Atlantic Salmon *Salmo salar* suggests increased parasite fitness. BMC Genomics 18, 1–14.

Brandal, P.O., Egidius, E., Romslo, I., 1976. Host blood: a major food component for the parasitic
copepod *Lepeophtheirus salmonis* Kröyeri, 1838 (Crustacea: Caligidae). Nor. J. Zool. 24, 341–
343.

Bron, J.E., Shinn, A.P., Sommerville, C., 2000. Ultrastructure of the cuticle of the chalimus larva of
the salmon louse *Lepeophtheirus salmonis* (Krøyer, 1837) (Copepoda: Caligidae). Contrib. to
Zool. 69, 39–49.

Bron, J.E., Sommerville, C., Jones, M., Rae, G.H., 1991. The settlement and attachment of early stages
of the salmon louse, *Lepeophtheirus salmonis* (Copepoda: Caligidae) on the salmon host, *Salmo
salar*. J. Zool. 224, 201–212.

Bron, J.E., Sommerville, C., Rae, G.H., 1993. The functional morphology of the alimentary canal of
larval stages of the parasitic copepod *Lepeophtheirus salmonis*. Zool. Soc. London 203, 207–
220.

Camacho, C., Coulouris, G., Avagyan, V., Ma, N., Papadopoulos, J., Bealer, K., Madden, T.L., 2009.
BLAST+: architecture and applications. BMC Bioinformatics 9, 1–9.

Dobin, A., Davis, C.A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., Batut, P., Chaisson, M.,
Gingeras, T.R., 2013. STAR: ultrafast universal RNA-seq aligner. Bioinformatics 29, 15–21.

Dysvik, B., Jonassen, I., 2001. J-Express: exploring gene expression data using Java. Bioinformatics
17, 369–370.

Edvardsen, R.B., Dalvin, S., Furmanek, T., Malde, K., Mæhle, S., Kvamme, B.O., Skern-Mauritzen, R., 2014. Gene expression in five salmon louse (*Lepeophtheirus salmonis*, Krøyer 1837) tissues. Mar. Genomics 18, 39–44.

Eichner, C., Dondrup, M., Nilsen, F., 2018. RNA sequencing reveals distinct gene expression patterns during the development of parasitic larval stages of the salmon louse (*Lepeophtheirus salmonis*). J. Fish Dis. 41, 1005–1029.

Eichner, C., Frost, P., Dysvik, B., Jonassen, I., Kristiansen, B., Nilsen, F., 2008. Salmon louse (*Lepeophtheirus salmonis*) transcriptomes during post molting maturation and egg production, revealed using EST-sequencing and microarray analysis. BMC Genomics 9, 1–15.

Eichner, C., Hamre, L.A., Nilsen, F., 2015. Instar growth and molt increments in *Lepeophtheirus salmonis* (Copepoda: Caligidae) chalimus larvae. Parasitol. Int. 64, 86–96.

Eichner, C., Nilsen, F., Grotmol, S., Dalvin, S., 2014. A method for stable gene knock-down by RNA interference in larvae of the salmon louse (*Lepeophtheirus salmonis*). Exp. Parasitol. 140, 44–51.

Ewels, P., Magnusson, M., Lundin, S., Max, K., 2016. MultiQC: summarize analysis results for multiple tools and samples in a single report. Bioinformatics 32, 3047–3048.

Giardine, B., Riemer, C., Hardison, R.C., Burhans, R., Elnitski, L., Shah, P., Zhang, Y., Blankenberg, D., Albert, I., Taylor, J., Miller, W., Kent, W.J., Nekrutenko, A., 2005. Galaxy: a platform for interactive large-scale genome analysis. Genome Res. 15, 1451–1455.

Gonzalez-Alanis, P., Wright, G.M., Johnson, S.C., Burka, J.F., 2001. Frontal filament morphogenesis in the salmon louse *Lepeophtheirus salmonis*. J. Parasitol. 87, 561–574.

Grimnes, A., Jakobsen, P.J., 1996. The physiological effects of salmon lice infection on post-smolt of Atlantic salmon. J. Fish Biol. 48, 1179–1194.

Guidry, C., Miller, E.J., Hook, M., 1990. A second fibronectin-binding region is present in collagen α chains. J. Biol. Chem. 265, 19230–19236.

Hamre, L.A., Bui, S., Oppedal, F., Skern-Mauritzen, R., Dalvin, S., 2019. Development of the salmon louse *Lepeophtheirus salmonis* parasitic stages in temperatures ranging from 3 to 24 °C. Aquac. Environ. Interact. 11, 429–443.

Hamre, L.A., Eichner, C., Caipang, C.M.A., Dalvin, S.T., Bron, J.E., Nilsen, F., Boxshall, G., Skern-Mauritzen, R., 2013. The salmon louse *Lepeophtheirus salmonis* (Copepoda: Caligidae) life cycle has only two chalimus stages. PLoS One 8, 1–9.

Hamre, L.A., Glover, K.A., Nilsen, F., 2009. Establishment and characterisation of salmon louse (*Lepeophtheirus salmonis* (Krøyer 1837)) laboratory strains. Parasitol. Int. 58, 451–460.

Harasimczuk, E., Øvergård, A.-C., Grotmol, S., Nilsen, F., Dalvin, S., 2018. Characterization of three salmon louse (*Lepeophtheirus salmonis*) genes with fibronectin II domains expressed by tegumental type 1 glands. Mol. Biochem. Parasitol. 219, 1–9.

Heggland, E.I., Eichner, C., Støve, S.I., Martinez, A., Nilsen, F., Dondrup, M., 2019a. A scavenger receptor B (CD36)-like protein is a potential mediator of intestinal heme absorption in the hematophagous ectoparasite *Lepeophtheirus salmonis*. Sci. Rep. 9, 1–14.

Heggland, E.I., Tröße, C., Eichner, C., Nilsen, F., 2019b. Heavy and light chain homologs of ferritin are essential for blood-feeding and egg production of the ectoparasitic copepod *Lepeophtheirus*

salmonis. Mol. Biochem. Parasitol. 232, 1–10.

Huang, X., Poelchau, M.F., Armbruster, P.A., 2015. Global transcriptional dynamics of diapause
induction in non-blood-fed and blood-fed *Aedes albopictus*. PLoS Negl. Trop. Dis. 1–28.

Hulsen, T., de Vlieg, J., Alkema, W., 2008. BioVenn – a web application for the comparison and
visualization of biological lists using area-proportional Venn diagrams. BMC Genomics 9, 1–6.

Johnson, S.C., 1993. A comparison of development and growth rates of *Lepeophtheirus salmonis*
(Copepoda: Caligidae) on naive Atlantic (*Salmo salar*) and chinook (*Oncorhynchus tshawytscha*)
salmon, in: Boxshall, G.A., Defaye, D. (Eds.), Pathgens of Wild and Farmed Fish: Sea Lice. Ellis
Horwood, pp. 68–80.

Johnson, S.C., Albright, L.J., 1992. Comparative susceptibility and histopathology of the response of
naive Atlantic, chinook and coho salmon to experimental infection with *Lepeophtheirus salmonis*
(Copepoda: Caligidae). Dis. Aquat. Organ. 14, 179–193.

Johnson, S.C., Albright, L.J., 1991a. The developmental stages of *Lepeophtheirus salmonis* (Krøyer,
1837) (Copepoda: Caligidae). Can. J. Zool. 69, 929–950.

Johnson, S.C., Albright, L.J., 1991b. Development, growth, and survival of *Lepeophtheirus salmonis*
(Copepoda: Caligidae) under laboratory conditions. J. Mar. Biol. Assoc. United Kingdom 71,
425–436.

Johnson, S.C., Ewart, K. V., Osborne, J.A., Delage, D., Ross, N.W., Murray, H.M., 2002. Molecular
cloning of trypsin cDNAs and trypsin gene expression in the salmon louse *Lepeophtheirus*
salmonis (Copepoda: Caligidae). Parasitol. Res. 88, 789–796.

Jones, M.W., Sommerville, C., Bron, J., 1990. The histopathology associated with the juvenile stages
of *Lepeophtheirus salmonis* on the Atlantic salmon, *Salmo salar* L. J. Fish Dis. 13, 303–310.

Jones, P., Binns, D., Chang, H.-Y., Fraser, M., Li, W., McAnulla, C., McWilliam, H., Maslen, J.,
Mitchell, A., Nuka, G., Pesseat, S., Quinn, A.F., Sangrador-Vegas, A., Scheremetjew, M., Yong,
S.-Y., Lopez, R., Hunter, S., 2014. Sequence analysis InterProScan 5: genome-scale protein
function classification. Bioinformatics 30, 1236–1240.

Khan, M.T., Dalvin, S., Nilsen, F., Male, R., 2017. Microsomal triglyceride transfer protein in the
ectoparasitic crustacean salmon louse (*Lepeophtheirus salmonis*). J. Lipid Res. 58, 1613–1623.

Kotsyfakis, M., Schwarz, A., Erhart, J., Ribeiro, J.M.C., 2015. Tissue- and time-dependent
transcription in *Ixodes ricinus* salivary glands and midguts when blood feeding on the vertebrate
host. Sci. Rep. 5, 1–10.

Kvamme, B.O., Skern, R., Frost, P., Nilsen, F., 2004. Molecular characterisation of five trypsin-like
peptidase transcripts from the salmon louse (*Lepeophtheirus salmonis*) intestine. Int. J. Parasitol.
34, 823–832.

Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G., Abecasis, G., Durbin,
R., 1000 Genome Project Data Processing Subgroup, 2009. The sequence alignment/map format
and SAMtools. Bioinformatics 25, 2078–2079.

Liao, M., Zhou, J., Gong, H., Boldbaatar, D., Shirafuji, R., Battur, B., Nishikawa, Y., Fujisaki, K.,
2009. Hemalin, a thrombin inhibitor isolated from a midgut cDNA library from the hard tick
Haemaphysalis longicornis. J. Insect Physiol. 55, 165–174.

Liao, Y., Smyth, G.K., Shi, W., 2014. featureCounts: an efficient general purpose program for
assigning sequence reads to genomic features. Bioinformatics 30, 923–930.

Love, M.I., Huber, W., Anders, S., 2014. Moderated estimation of fold change and dispersion for
RNA-seq data with DESeq2. Genome Biol. 15, 1–21.

Lumjuan, N., Stevenson, B.J., Prapanthadara, L., Somboon, P., Brophy, P.M., Loftus, B.J., Severson,
D.W., Ranson, H., 2007. The *Aedes aegypti* glutathione transferase family. Insect Biochem. Mol.
Biol. 37, 1026–1035.

Marinotti, O., Nguyen, Q.K., Calvo, E., James, A.A., Ribeiro, J.M.C., 2005. Microarray analysis of
genes showing variable expression following a blood meal in *Anopheles gambiae*. Insect Mol.
Biol. 14, 365–373.

McMullen, B.A., Fujikawa, K., 1985. Amino acid sequence of the heavy chain of human a-Factor
XIIa (activated Hageman factor). J. Biol. Chem. 260, 5328–5341.

Mulenga, A., Erikson, K., 2011. A snapshot of the *Ixodes scapularis* degradome. Gene 482, 78–93.

Nayduch, D., Lee, M.B., Saski, C.A., 2014. The reference transcriptome of the adult female biting
midge (*Culicoides sonorensis*) and differential gene expression profiling during teneral, blood,
and sucrose feeding conditions. PLoS One 9, 1–15.

Nylund, A., Økland, S., Bjørknes, B., 1992. Anatomy and ultrastructure of the alimentary canal in
Lepeophtheirus salmonis (Copepoda: Siphonostomatoida). J. Crustac. Biol. 12, 423–437.

Perner, J., Kotál, J., Hatalová, T., Urbanová, V., Bartosova-Sojkova, P., Brophy, P.M., Kopácek, P.,
2018. Inducible glutathione S-transferase (IrGST1) from the tick *Ixodes ricinus* is a haem-
binding protein. Insect Biochem. Mol. Biol. 95, 44–54.

Perner, J., Provazník, J., Schrenková, J., Urbanová, V., Ribeiro, J.M.C., Kopáček, P., 2016. RNA-seq
analyses of the midgut from blood- and serum-fed *Ixodes ricinus* ticks. Sci. Rep. 6, 1–18.

Poley, J.D., Sutherland, B.J.G., Jones, S.R.M., Koop, B.F., Fast, M.D., 2016. Sex-biased gene
expression and sequence conservation in Atlantic and Pacific salmon lice (*Lepeophtheirus*
salmonis). BMC Genomics 17, 1–16.

Skern-Mauritzen, R., Torrissen, O., Glover, K.A., 2014. Pacific and Atlantic *Lepeophtheirus salmonis*
(Krøyer, 1838) are allopatric subspecies: *Lepeophtheirus salmonis salmonis* and *L. salmonis*
oncorhynchi subspecies novo. BMC Genet. 15, 1–9.

Stavrum, A.K., Petersen, K., Jonassen, I., Dysvik, B., 2008. Analysis of gene-expression data using
J-Express. Curr. Protoc. Bioinforma. 21, 7.3.1-7.3.25.

Supek, F., Bosnjak, M., Skunca, N., Smuc, T., 2011. REVIGO summarizes and visualizes long lists of
gene ontology terms. PLoS One 6, 1–9.

Sutherland, B.J.G., Covello, J.M., Friend, S.E., Poley, J.D., Koczka, K.W., Purcell, S.L., MacLeod,
T.L., Donovan, B.R., Pino, J., González-Vecino, J.L., Gonzalez, J., Troncoso, J., Koop, B.F.,
Wadsworth, S.L., Fast, M.D., 2017. Host-parasite transcriptomics during immunostimulant-
enhanced rejection of salmon lice (*Lepeophtheirus salmonis*) by Atlantic salmon (*Salmo salar*).
Facets 2, 477–495.

Sutherland, B.J.G., Jantzen, S.G., Yasuike, M., Sanderson, D.S., Koop, B.F., Jones, S.R.M., 2012.

Transcriptomics of coping strategies in free-swimming *Lepeophtheirus salmonis* (Copepoda)
larvae responding to abiotic stress. Mol. Ecol. 21, 6000–6014.

Sutherland, B.J.G., Poley, J.D., Igboeli, O.O., Jantzen, J.R., Fast, M.D., Koop, B.F., Jones, S.R.M.,
2014. Transcriptomic responses to emamectin benzoate in Pacific and Atlantic Canada salmon
lice *Lepeophtheirus salmonis* with differing levels of drug resistance. Evol. Appl. 8, 133–148.

Tekle, K.M., Gundersen, S., Klepper, K., Bongo, L.A., Raknes, I.A., Li, X., Zhang, W., Andreetta, C.,
Mulugeta, T.D., Kalaš, M., Rye, M.B., Hjerde, E., Samy, J.K.A., Fornous, G., Azab, A., Våge,
D.I., Hovig, E., Willassen, N.P., Drabløs, F., Nygård, S., Petersen, K., Jonassen, I., 2018.
Norwegian e-Infrastructure for Life Sciences (NeLS) [version 1; peer review: 2 approved].
F1000Research 7, 1–15.

Tjensvoll, K., Hodneland, K., Nilsen, F., Nylund, A., 2005. Genetic characterization of the
mitochondrial DNA from *Lepeophtheirus salmonis* (Crustacea; Copepoda). A new gene
organization revealed. Gene 353, 218–230.

Treasurer, J.W., Wadsworth, S.L., 2004. Interspecific comparison of experimental and natural routes
of *Lepeophtheirus salmonis* and *Caligus elongatus* challenge and consequences for distribution
of chalimus on salmonids and therapeutant screening. Aquac. Res. 35, 773–783.

Trevisan-Silva, D., Gremski, L.H., Chaim, O.M., Silveira, R.B. da, Meissner, G.O., Mangili, O.C.,
Barbaro, K.C., Gremski, W., Veiga, S.S., Senff-Ribeiro, A., 2010. Astacin-like metalloproteases
are a gene family of toxins present in the venom of different species of the brown spider (genus
Loxosceles). Biochimie 92, 21–32.

Wu, T.D., Watanabe, C.K., 2005. GMAP: a genomic mapping and alignment program for mRNA and
EST sequences. Bioinformatics 21, 1859–1875.

Øvergård, A.-C., Hamre, L.A., Harasimczuk, E., Dalvin, S., Nilsen, F., Grotmol, S., 2016. Exocrine
glands of *Lepeophtheirus salmonis* (Copepoda: Caligidae): distribution, developmental
appearance, and site of secretion. J. Morphol. 277, 1616–1630.

Figures

749 **Figure 1.** Distribution of different stages and instar ages of lice on fish at 10 days post infestation
750 (dpi) and at 18 dpi sampled from the fish body, fins and gills in both experiments (Ex1 and Ex2). Lice
751 instar ages were defined on photographs as described in the main text. Transcriptome sequencing
752 was performed for lice from Ex1 10 dpi, Ex2 10 dpi (both chalimus I old), and Ex2 18 dpi (chalimus II
753 old) marked with *.
754

Figure 2. Photographs of salmon lice with (right) and without (left) a blood-filled intestine. Chalimus I larvae sampled 10 days post infestation (a). Chalimus II larvae sampled 18 days post infestation (b). Preadult I lice on frontal filament sampled 18 days post infestation (c). The lice with blood-filled intestine were sampled from the gills, and the others were sampled from the skin of their host. Scale bars = 1 mm.

Figure 3 a) MA plot comparing the different conditions (lice sampled from gills versus lice sampled from skin at 10 days post infestation (dpi) and 18 dpi respectively) in the samplings: Experiment (Ex) 1, sampled at 10 dpi, Ex2 sampled at 10 dpi and Ex2 sampled at 18 dpi. The average log intensity for a dot in the plot (A) is shown on the x-axis and the binary logarithm of the intensity ratio (M) is shown on the y-axis. b) Principle component analysis (PCA) plots of the same data. c) Correspondence analysis (CA) plot showing the overall gene expression of the samples analyzed in this study (pink and turquoise dots) in comparison with other chalimus I and chalimus II larvae divided into various instar ages taken from Eichner et al. (2018).

Sampling	Higher in lice sampled from gills	Higher in lice sampled from skin
all 3 samplings	protein metabolism glycolytic process	protein phosphorylation
Ex1 10 dpi and Ex2 10 dpi	protein metabolism carbohydrate metabolism	phosphorus metabolic process
Ex1 10 dpi and Ex2 18 dpi	protein metabolism carbohydrate metabolism	cellular response to chemical stimulus
Ex2 10 dpi and Ex2 18 dpi	protein metabolism carbohydrate metabolism	protein phosphorylation response to stress
Ex1 10 dpi only	translation chromosome organization	transport regulation of intracellular signal transduction
Ex2 10 dpi only	second-messenger-mediated signaling	chitin metabolism regulation of cellular component organization
Ex2 18 dpi only	transport carbohydrate metabolism protein metabolism localization response to stress	transport dephosphorylation regulation of cellular process biological regulation

Figure 4. Scaled Venn diagrams showing the number of differentially expressed genes between
chalimus larvae sampled from gills versus those sampled from skin, for Experiment 1, sampled at 10
days post infestation (dpi), Experiment 2, sampled at 10 dpi and Experiment 2 sampled at 18 dpi. The
numbers of genes elevated in chalimus larvae sampled from gills are shown in (a) and the ones
elevated in chalimus larvae sampled from fish skin are shown in (b). Representative GO terms for
each group are given in the table in the bottom panel. A full list of enriched GO terms is shown in
Supplementary Table S9 which are summarized as Tree maps in supplementary Figure S1.

Figure 5. GOslim annotations of genes elevated in lice sampled from gills (a) and genes elevated in
lice sampled from skin (b) in all three sampling points. Columns indicate the number of genes in each
GOslim category, while black dots show the enrichment of genes in that category in relation to
number of genes of the specific category in the whole dataset.

Figure 6. Hierarchical clustering of the expression profiles of genes found DE in all three sampling points with trypsin, other peptidase and FNII domains after expression profiles from LiceBase (various tissues and stages) and from the time series study (average values of biological parallels) by Eichner et al. (2018). A blue stippled line is separating planktonic and parasitic stages. Stable IDs with font in blue = domains other than trypsin predicted as well, font in green = predicted M13 peptidases, font in red = predicted Astacin peptidases.

1. Supplementary Fig S1

Higher in lice from gills   **Higher in lice from skin**

a) Enriched GO terms for genes found higher expressed in samples from Ex1_10dpi, Ex2_10dpi and Ex2_18dpi

b) Enriched GO terms for genes found higher expressed in samples from Ex1_10dpi and Ex2_10dpi

c) Enriched GO terms for genes found higher expressed in samples from Ex1_10dpi and Ex2_18dpi

d) Enriched GO terms for genes found higher expressed in samples from Ex2_10dpi and Ex2_18dpi

Higher in lice from gills   **Higher in lice from skin**

e) Enriched GO terms for genes found higher expressed in samples from Ex1_10dpi only

f) Enriched GO terms for genes found higher expressed in samples from Ex2_10dpi only

g) Enriched GO terms for genes found higher expressed in samples from Ex2_18dpi only

799   Supplementary Fig S1. Tree maps (Revigo) of GO annotation belonging to biological process
800   for a) enriched GO terms for genes found higher expressed in samples from all three
801   samplings; b) of the genes DE in both samplings at 10 dpi; and in 2 samplings sampled at
802   different time point (c and d). In e- g are enriched GO annotations shown for genes
803   differentially expressed in one of the three samplings only.

2. Supplementary Fig S2

807
808
Supplementary Fig S2 Expression profiles of the ten strongest genes upregulated in lice sampled from
gills (a) and in lice sampled from skin (b) over all samples.

3. Supplementary Fig S3

Supplementary Figure S3. Expression of a) Ferritin 1, b) Ferritin 2, c) Ferritin 4, d) LsHSCARB, e) Lipid
transfer protein (EMLSAT00000001530)(Khan et al., 2017) in the samples from lice sampled from skin
(light grey) and gills (dark grey). Shown are average values with standard deviation. * = Significant
different (T-test) between lice samples from different location in the respective sampling point.
817

818   4. Supplementary Fig S4

820   Supplementary Fig S4. Copepodite with blood in intestine sampled from gills (a) and without

821   blood (b) at 3 days post infestation (samples not included in this study).

Tables

823   5. Table 1

	Chalimus I	Chalimus II		Preadult I		NN
		male	female	male	female	
Ex1 10 dpi						
body	75.6	14.0	1.2	-	-	9.3
fin	72.1	16.6	-	-	-	11.4
gill	98.4	-	-	-	-	1.6
Ex2 10 dpi						
body	88.6	8.0	-	-	-	3.4
fin	89.9	7.2	-	-	-	3.0
gill	100.0	-	-	-	-	-
Ex1 18 dpi						
body	-	1.0	5.5	61.9	30.5	1.0
fin	-	3.6	40.1	9.5	37.2	9.5
gill	-	14.8	52.3	27.5	1.3	4.0
Ex2 18 dpi						
body	-	2.0	5.5	61.5	27.3	3.7
fin	-	1.9	27.9	5.8	51.0	13.5
gill	-	13.5	48.8	34.4	0.3	3.0

825   Table 1 proportion of different stages and sexes (for Chalimus II and preadult I) from all fish
826   sampled at 10 days post infestation (dpi) and 18 dpi in Experiment (Ex) 1 and 2. NN =
827   undefined

	Experiment 1				Experiment 2			
	all	body	fin	gills	all	body	fin	gills
Pad I of total	68 (294)	88 (228)	49 (40)	30 (26)	66 (329)	87 (206)	63 (55)	38 (62)
female of pad I	33 (102)	30 (68)	78 (31)	8 (2)	35 (115)	30 (62)	87 (48)	0 (0)
Pad I on filament	10 (29)	5 (12)	25 (10)	27 (7)	21 (69)	6 (12)	38 (21)	58 (36)
Pad I with visible blood	16 (48)	13 (30)	0 (0)	69 (18)	24 (80)	13 (27)	0 (0)	85 (53)
Pad I with visible blood on filament	1 (2)	0 (0)	0 (0)	8 (2)	9 (30)	0 (0)	0 (0)	48 (30)

837
838 Table 2. Preadult (pad) I lice were investigated in more detail of 10 and 9 fish in Experiment 1 and 2
839 respectively. The proportion and absolute numbers (in brackets) of preadult I lice on the different
840 body parts, the proportion of females of these as well as the proportion of preadult I lice with visible
841 blood in the intestine or with filament or with both are shown.
842
843