Disrupted remapping of place cells and grid cells in knock-in model of Alzheimer's disease

Heechul Jun[1,2,3], Shogo Soma[1,2,3] †, Takashi Saito[4] ‡, Takaomi C Saido[4], Kei M Igarashi[1,2,3,5]*

[1]Department of Anatomy and Neurobiology, University of California, Irvine.

[2]Center for the Neurobiology of Learning and Memory, University of California, Irvine.

[3]Institute for Memory Impairments and Neurological Disorders, University of California, Irvine

[4]Lab for Proteolytic Neuroscience, RIKEN Center for Brain Science, Japan

[5]Japan Science and Technology Agency

*Correspondence to: kei.igarashi@uci.edu

†Current address: Department of Molecular Cell Physiology, Graduate School of Medical Science, Kyoto Prefectural University of Medicine

‡Current address: Department of Neurocognitive Science, Institute of Brain Science, Nagoya City University Graduate School of Medical Science

Abstract:

Patients with Alzheimer's disease (AD) frequently suffer from spatial memory impairment and wandering behavior, but brain circuits causing such symptoms remain largely unclear. In healthy brains, spatially-tuned hippocampal place cells and entorhinal grid cells represent distinct spike patterns in different environments, a circuit function called "remapping" that underlies pattern separation of spatial memory. We investigated whether a knock-in manipulation to mutated amyloid precursor protein affects the remapping of place cells and grid cells. We found that the remapping of hippocampal place cells was severely disrupted although their spatial tuning was only mildly diminished. Entorhinal grid cells were severely impaired, sending disrupted remapping signals to the hippocampus. These results point to the link between grid cell impairment and remapping disruption as a circuit mechanism causing spatial memory impairment in AD.

Main Text:

Spatial memory impairment such as wandering behavior is one of the most troublesome symptoms in Alzheimer's disease (AD), occurring in more than 60% of AD patients (Hope et al., 1994). Despite recent molecular and cellular findings in AD research, it is still largely unclear how deterioration of brain circuit function causes spatial memory loss in AD. In healthy brains of both humans and rodents, grid cells in the medial entorhinal cortex (MEC) and place cells in the hippocampus form a neural circuit that is critical for spatial memory and navigation (**Fig. S1**) (Scoville and Milner, 1957; O'Keefe and Dostrovsky, 1971; Morris et al., 1982; Ekstrom et al., 2003; Fyhn et al., 2004; Steffenach et al., 2005; Doeller et al., 2010). In this circuit, spatially-tuned grid cells in the superficial layer of MEC send spatial signals to the hippocampus to form spatially-tuned place cells (Moser et al., 2015). These grid cells and place cells have an ability to store multiple combinations of firing patterns when subjects enter into different environments, a function called "remapping" (Muller and Kubie, 1987; Fyhn et al., 2007; Alme et al., 2014; Kyle et al., 2015). Because remapping of place cells and grid cells can provide subjects with distinct combinatorial codes for multiple environments, it provides support for the circuit mechanism for the pattern separation of spatial memory (Colgin et al., 2008; Yassa and Stark, 2011). Although impairments in the spatial tuning of place cells or grid cells were reported (Cacucci et al., 2008; Kunz et al., 2015; Fu et al., 2017; Mably et al., 2017), no previous studies have investigated remapping of place cells and grid cells in AD subjects.

In this study, we specifically asked two questions: (1) Is remapping of hippocampal place cells deteriorated in AD subjects? (2) If so, is place cell deterioration linked to impairment of entorhinal grid cells? Alternatively, does place cell deterioration occur autonomously inside the hippocampus and independent from grid cells? To this end, we investigated the remapping of hippocampal place cells and entorhinal grid cells in a novel amyloid precursor protein knock-in (APP-KI) mouse we recently developed (Saito et al., 2014). Previous transgenic AD models express excessive amount of mutated genes from multiple copies of transgenes in both the entorhinal cortex and the hippocampus, making

it difficult to compare pathophysiology between the two brain regions. By contrast, in our knock-in model, mutated APP protein is expressed from only two copies of mutated human APP gene, as is the case for human AD patients. Therefore, relative pathological progressions between the hippocampus and the entorhinal cortex of APP-KI mice are expected to recapitulate those from AD patients, allowing us to compare the pathophysiology between the hippocampus and the entorhinal cortex. Because APP-KI mice show severe spatial memory impairments (Masuda et al., 2016), we investigated the remapping of place cells and grid cells underlying the spatial memory impairments.

Results

Hippocampal CA1 neurons showed severely disrupted remapping, but mildly diminished spatial tuning in APP-KI mice.

In AD patients, pathophysiology starts with amyloid-β (Aβ) plaque deposition from the expression of APP, which slowly leads to Tau pathology and deterioration of brain circuit functions in the course of more than 20 years (Sasaguri et al., 2017). The circuit deterioration ultimately leads to memory impairment, with which patients are diagnosed with AD. To replicate the slow and cumulative effect of pathological APP on brain circuit functions in mice, we recently generated a knock-in mouse model by manipulating the endogenous murine APP gene into the mutated human APP^{NL-F-G} gene (Saito et al., 2014). This knock-in manipulation causes Aβ deposition starting at 4-mo and severe spatial memory impairment at 6-mo (Saito et al., 2014; Masuda et al., 2016). To investigate place cells after memory impairment, we implanted a recording device containing 64 channels of electrodes in APP-KI mice and recorded place cells after 7-mo (7-13 mo, **Fig. 1A-B** and **S2-S3**; no change observed in place cells and grid cells from APP-KI mice across this age period, **Fig. S4**). Because the APP-KI mouse is on the C57BL/6 background, we used 7-13 mo C57BL/6 mice as a control (referred to as WT mice). No behavioral difference was observed between APP-KI and WT mice (**Fig. S5**). To avoid biased sampling toward spatially tuned cells that can be fewer in APP-KI mice, we collected all recorded principal neurons in our data pool (n=105 neurons from n = 10 APP-KI mice; n=113 neurons from n = 10 WT mice). No difference was observed in the distribution of

putative principal neurons and interneurons between WT and APP-KI mice (**Fig. S6**). Neural activities of CA1 neurons were first recorded in a 1x1 m open field to evaluate their spatial tuning and place cell property (**Fig. 1C**). Although our WT mice were between the age of 7-13 months, a good amount of CA1 neurons of WT mice showed spatially tuned spike activities (**Fig. 1D**). CA1 neurons in APP-KI mice also showed spatially tuned spike activities, but exhibited less apparent firing peaks (**Fig. 1E**). The degree of spatial tuning for individual CA1 neurons was quantitatively assessed using *spatial information* score (Skaggs et al., 1993) (**Fig. 1F-G**). The analysis revealed that APP-KI mice have neurons with less spatial tuning ($p < 0.001$, Kolmogorov-Smirnov (KS) test), with lower mean spatial information than those from the CA1 neurons recorded in the WT mice ($p < 0.001$, Wilcoxon rank sum test; detailed statistic results are shown in Supplementary Tables S2-S14). However, mean firing rates, field size and firing field number of CA1 neurons did not differ between WT and APP-KI mice, suggesting the spatial tuning of CA1 neurons in APP-KI mice was only mildly deteriorated (**Fig. S7**). We then defined place cells using the spatial information score (Langston et al., 2010). Neurons were classified as place cells if their spatial information score was larger than 0.63, which was the 95th percentile of a distribution of spatial information score based on shuffled data from WT mice (**Fig. 1F and S8**). This revealed that proportion of place cells decreased down from 50% in WT mice to 31% in APP-KI mice ($p < 0.001$, binomial test; **Fig 1G**). Among these place cells, we did not observe any differences in their properties recorded from the open field task between APP-KI and WT mice **(Fig. S8)**.

To investigate the remapping of CA1 neurons in APP-KI mice, we then trained animals to run back and forth in two 1-m linear tracks (Tracks A and B) having distinct colors and textures (**Fig. 1H**, see experimental protocol). In an everyday session, animals made 10 raps for each track in a following series: Track A, Track B, Track B and then back again in Track A. On the fifth day of the training, CA1 neurons in WT mice showed remapping across Track A and Track B, with non-overlapping firing patterns between the two tracks (**Fig. 1I,** also known as "global remapping" (Leutgeb et al., 2005b)). However, with the same training condition, CA1 neurons in APP-KI mice showed unchanged firing patterns between Track A and Track B (**Fig. 1J**). Population vector (PV) correlation analysis (Leutgeb et al., 2005b) was used to assess the degree of remapping between

Track A and Track B (**Fig. 1K**). PV correlation of -1 indicates that recorded population of CA1 neurons showed completely distinct firing patterns (that is, perfect remapping), whereas PV correlation of 1 indicates exactly same firing patterns (that is, no remapping at all). PV correlation analysis revealed that WT mice showed good remapping with low PV correlation between Track A and Track B (**Fig. 1L**). However, APP-KI mice exhibited PVs with significantly higher correlation, indicating that the remapping of CA1 neurons was significantly disrupted in APP-KI mice (P<0.001, KS test, **Fig. 1L**). Mean PV correlation in APP-KI mice was significantly higher than that of WT mice (p<0.001, Wilcoxon rank sum test; **Fig. S9G**). Disrupted remapping was observed independent of the direction of animals' runs in the tracks (**Fig. S10**). Remapping of the neurons defined as place cells was also significantly disrupted (**Fig. S8**). We also assessed the degree of remapping in individual neurons using *spatial correlation* score between Track A and Track B (Leutgeb et al., 2005b), confirming that CA1 neurons showed disrupted remapping in APP-KI mice compared to that from WT mice (**Fig. S11**). By contrast, activity patterns of place cells between the two recordings in Track A were similar, with no difference in PV correlations between WT and APP-KI mice, indicating that stability of spatial tuning in CA1 neurons was spared in APP-KI mice (p>0.05, KS test, **Fig. 1L**; p>0.05, Wilcoxon rank sum test, **Fig. S9F**). Together, these results demonstrate that CA1 principal neurons from APP-KI mice exhibit disrupted remapping for different spatial environments, while the spatial tuning of CA1 neurons in is only mildly deteriorated.

Diminished fast gamma oscillations, but not slow gamma oscillations in the hippocampal CA1 of APP-KI mice

CA1 neurons in the hippocampus receive spatial inputs from hippocampal CA3 via the Schaeffer collateral as well as from the MEC via temporoammonic pathway (Witter and Amaral, 2004). We next asked whether these inputs contribute to disrupted remapping in CA1. We leveraged findings from previous healthy animal studies that identified two distinct types of gamma oscillations (Bragin et al., 1995; Colgin et al., 2009) (**Fig. 2A**). When fast gamma oscillations occur in the hippocampal CA1, the MEC also exhibits fast gamma oscillations. The oscillations from these two brain regions are highly synchronized, suggesting that fast gamma oscillations support signal transfer from the

MEC to CA1. By contrast, synchronization of slow gamma oscillations suggests signal transfer from CA3 to CA1 (Colgin et al., 2009). We thus analyzed gamma oscillations in CA1 of WT and APP-KI mice. In WT mice, we observed gamma episodes across slow and fast gamma range (20 – 90 Hz; **Fig. 2B**). However, in APP-KI mice, gamma oscillations were observed only in low frequency ranges, and marginal gamma episodes were observed at a frequency range above 40Hz. Comparison of oscillatory power normalized by gross 1-100 Hz frequency power (Deshmukh et al., 2010) showed reduction of power at 62 – 100 Hz and 34 – 36 Hz ranges, and increase of power at 5 – 8 Hz theta oscillation range in APP-KI mice (false discovery rate (FDR) corrected for 100 multiple comparisons at 1–100 Hz bins, q<0.05; **Fig. 2C**).

To test whether fast and slow gamma oscillations are deteriorated in CA1, we investigated temporal structures of gamma oscillations. Previous healthy animal studies showed that gamma oscillations occur mainly at the peak to falling phase of co-existing theta oscillations, a property termed theta-gamma coupling (Bragin et al., 1995; Colgin et al., 2009). The theta-gamma cross-frequency coupling correlates with animals' memory performance and is suggested to be a mechanism for local information processing during memory demands (Tort et al., 2009; Axmacher et al., 2010). Cross-frequency coherence plot from WT mice revealed peaks at two gamma frequency ranges at 20 – 40 Hz and 40 – 90 Hz that were coupled to specific phases of theta oscillations (**Fig. 2D**). We thus defined slow gamma oscillations at 20 – 40 Hz frequency range, and fast gamma oscillations at 40 – 90 Hz range in our experiment. In APP-KI mice, only the slow gamma oscillations from 20 – 40 Hz range was coherent to co-existing theta oscillations (**Fig. 2D**). The occurrence rate of fast gamma oscillation was diminished in APP-KI mice (p<0.001, Wilcoxon rank sum test, **Fig. 2E**), whereas the occurrence rate of slow gamma remained similar (p >0.05 **Fig. 2E**). Cross-frequency coupling spectrogram revealed fast and slow gammas in the peak to falling phase of theta oscillations in WT mice, whereas fast gamma oscillations were significantly lost in APP-KI mice (**Fig. 2F**). The degree of theta-gamma cross-frequency coupling was assessed by measuring the phase-locking vector length (Colgin et al., 2009). The analysis revealed diminished theta-fast gamma coupling, while theta-slow gamma coupling was spared (p<0.05 for fast gamma; p>0.05 for slow gamma, Wilcoxon ranks sum test; **Fig. 2G**). To test whether spike timing is entrained by fast and

slow gamma oscillations, we examined the temporal organization of spike activities during slow and fast gamma oscillations (Colgin et al., 2009). Phase-locking of spikes to fast gamma oscillations was diminished in APP-KI mice ($p<0.001$, Wilcoxon rank sum test; **Fig. 2H-I**). Altogether, our results demonstrate that fast gamma oscillations were disrupted in the hippocampal CA1 of APP-KI mice, while slow gamma oscillations were relatively spared. These results suggest that MEC→CA1 signal transfer via fast gamma oscillations is deteriorated in APP-KI mice, whereas slow gamma-mediated CA3→CA1 signal transfer remains relatively intact.

MEC neurons in APP-KI mice showed severe impairment in spatial tuning

Disrupted fast gamma oscillations observed in hippocampal CA1 implies that neural activity in the MEC is affected in APP-KI mice. To test this possibility, we next investigated spatial representation in the MEC. We implanted a recording device into the MEC, and performed unbiased recording of principal neurons in layers 2 and 3 of the dorsal MEC (n=61 neurons from n = 13 WT mice; n= 65 neurons from n = 13 APP-KI mice; **Fig. 3A-C and Fig. S12-S13**; no difference in the sampling positions between WT and APP-KI mice, **Fig. S13**). To test for their spatial tuning, MEC neurons were recorded while animals were randomly foraging through 1x1m open field task. Among the recorded MEC neurons from WT mice, a portion of cells exhibited spatially tuned multiple firing fields as grid cells (**Fig. 3D**). By contrast, most of the recorded MEC neurons in APP-KI mice showed scattered firing patterns around the open field (**Fig. 3E**). To evaluate grid cell property, we analyzed *gridness score* by computing the autocorrelation of firing rate map (Sargolini et al., 2006) (**Fig. 3D-E**). A cumulative distribution curve of the gridness score for all recorded MEC neurons indicate that, compared to WT mice, APP-KI mice significantly lost a group of neurons with gridness score of more than ~0.4 ($p<0.05$, KS test; **Fig. 3F**). To define grid cells, we used a cut-off of gridness score at 0.41, which was the 95th percentile of the distribution of gridness score based on the shuffled WT data (Langston et al., 2010) (**Fig. 3F and S14**). We found that, among recoded MEC neurons, 20% was grid cells in WT mice, whereas only 2% was grid cells in APP-KI mice ($p<0.001$, binomial test; **Fig. 3F-G**). The scattered firing of APP-KI neurons resulted in significantly

reduced spatial tuning (p< 0.001, Wilcoxon rank sum test) and increased firing field area (p <0.01, Wilcoxon rank sum test, **Fig. 3G**). However, we did not observe any difference in their mean or peak firing rates, suggesting that MEC neurons in APP-KI mice did not show hyperactivity reported previously in severe transgenic AD models (Busche et al., 2008) (**Fig. 3G and S14**). The distribution of spatial information score for MEC neurons indicate that APP-KI mice lost significant portion of neurons with high spatial information score, suggesting that aperiodic spatially tuned MEC neurons (Miao et al., 2017) were also impaired in APP-KI mice (**Fig. S15**). These data indicate that spatial tuning of MEC neurons, including grid cells, was severely disrupted in APP-KI mice.

MEC neurons in APP-KI mice could not send remapping signals to the hippocampus

We then tested whether remapping signal is property signaled by these impaired MEC neurons. Previous healthy animal study showed that MEC grid cells exhibit remapping across distinct environments (Fyhn et al., 2007). The remapping of MEC grid cells occurs coherently with the remapping of hippocampal place cells, suggesting that MEC grid cells feed remapping signals into hippocampal place cells. Consistent with the previous finding, we observed that MEC neurons in WT mice showed non-overlapping grid fields between Track A and Track B (**Fig. 3H**). However, MEC neurons in APP-KI mice showed scattered firing along both Track A and Track B, showing marginal difference in the firing patterns between the two tracks (**Fig. 3I**). PV correlation analysis revealed that APP-KI mice show larger PV correlations than that from WT mice, indicating that remapping property of MEC neurons is disrupted in APP-KI mice (P<0.001, KS text; **Fig. 3J and S16-S17**). We also assessed spatial correlation of individual neurons between Track A and Track B, confirming that MEC neurons showed disrupted remapping in APP-KI (**Fig. S18**). However, we did not observe significant difference between WT and APP-KI mice in the stability of MEC neurons from two Track A recordings (p > 0.05, KS test; **Fig. 3J**). These data clearly demonstrate that remapping signals encoded by MEC neurons are severely impaired in APP-KI mice.

Remapping impairment co-existed with the disrupted spatial tuning in MEC neurons, but not in CA1 neurons

To examine the causality between impairment of spatial tuning and disruption of remapping in CA1 and MEC neurons, we measured the correlation between spatial information score and spatial correlation between Track A and Track B in individual neurons (**Fig. 3K**). In WT mice, neither CA1 neurons nor MEC neurons showed correlation between spatial tuning and remapping, suggesting that these two functional properties are independent in healthy WT mice (P>0.05, Pearson correlation). However, in APP-KI mice, we found that MEC neurons show significant negative correlation between spatial information score and spatial correlation (p<0.001 and r(65) = -0.45, Pearson correlation). These results suggest that the impairment of spatial tuning causes remapping disruption in individual MEC neurons in a cell autonomous manner. By contrast, CA1 neurons in APP-KI mice did not show significant correlation (p>0.05, Pearson correlation), suggesting that the remapping disruption of individual CA1 neurons is not autonomously caused by the impairment of spatial tuning.

Fast gamma coupling between CA1 and MEC were impaired in APP-KI mice.

To test if the impairment of MEC neurons is accompanied by the deterioration of local gamma oscillations, we examined gamma oscillations in the MEC (**Fig. 4A**). Because previous healthy animal studies found only fast gamma but not slow gamma oscillations in the MEC (Chrobak and Buzsaki, 1998; Colgin et al., 2009), we focused our analysis on fast gamma oscillations (no difference was observed for MEC slow gamma oscillations between WT and APP-KI mice, **Fig. S19**). The analysis of theta-fast gamma cross-frequency coupling revealed the absence of fast gamma oscillations coupled to theta oscillations (**Fig. 4B**). However, we did not see any difference in the normalized oscillatory power nor in the occurrence rate of fast gamma oscillation between WT and APP-KI mice (q>0.05 for normalized power, FDR corrected, **Fig. 4C**; p>0.05 for gamma occurrence, Wilcoxon rank sum test; **Fig. 4D**). Theta-phase spectrogram showed tendency of fast gamma oscillations being weakly modulated by theta oscillations (p=0.051, Wilcoxon rank sum test, **Fig. 4E-F**). The spikes from MEC neurons in WT mice

were phase-locked to local fast gamma oscillations, as reported in previous study (Quilichini et al., 2010) (**Fig. 4G-H**). However, the spike-fast gamma phase-locking was markedly disrupted in APP-KI mice (p<0.01, Wilcoxon rank sum test, **Fig. 4H and S20**).

To directly test the idea that the impairment of fast gamma oscillations affects MEC→CA1 signal transfer in APP-KI mice, we implanted two bundles of electrodes and simultaneously recorded from the hippocampal CA1 and the MEC (n = 6 WT mice and n = 6 APP-KI mice) (**Fig. 4I**). To test the degree of synchronization between the gamma oscillations in the MEC and CA1, we assessed coherence of oscillatory activity (Igarashi et al., 2014). Analysis revealed that the gamma oscillations between the MEC and CA1 have weaker coherence at 58 – 75 Hz range in APP-KI mice (q<0.05, FDR corrected; **Fig. 4J**). We then collected oscillatory activity in the MEC when fast gamma oscillations were detected from the hippocampal CA1 (Colgin et al., 2009) (**Fig. 4K**). The MEC fast gamma oscillations were significantly desynchronized from the CA1 fast gamma oscillations in APP-KI mice. This was confirmed by the decreased fast gamma power ratio in APP-KI mice (P<0.05, Wilcoxon rank sum test; **Fig. 4L**). Altogether, these results demonstrate that the temporal coordination of the gamma oscillations between the MEC and the CA1 is diminished, suggesting that MEC→CA1 signal transfer mediated by the fast gamma oscillations is deteriorated in APP-KI mice.

Discussion

In this study, using the novel APP-KI mice, we investigated the effect of mutated APP on the remapping of MEC neurons and CA1 neurons in AD for the first time. We found that the remapping of CA1 neurons was disrupted in post-plaque APP-KI mice. Spatial tuning of MEC neurons, including grid cells, was severely impaired, sending disrupted remapping signals to the hippocampus. Fast gamma oscillations were deteriorated in both CA1 and MEC, suggesting impaired signal transfer in the MEC→CA1 circuit. This line of evidence points to the impairment of MEC neurons as an element of cause for remapping disruption in the entorhinal-hippocampal circuit of APP-KI mice (**Fig. S1**). Previous anatomical studies in AD patients showed that neurons in the entorhinal cortex exhibits the earliest neuronal deterioration (Van Hoesen et al., 1991; Gomez-Isla

et al., 1996), but the link between deterioration of entorhinal neurons and spatial memory impairment has been unclear. Our study provides the first circuit-level evidence that identifies the link between the impairment of MEC neurons and the remapping disruption of hippocampal CA1 neurons. In early-stage AD patients, deterioration of entorhinal neuron may trigger remapping impairment in hippocampal neurons. Because the remapping of hippocampal neurons is one of the most plausible circuit functions for spatial pattern separation to distinguish multiple environments (Colgin et al., 2008; Yassa and Stark, 2011), disruption of remapping would ultimately lead to spatial memory loss and wandering symptom in AD. Although the hippocampus has been used as the main model brain region in AD research for decades, our finding strengthens the importance of entorhinal cortex as a potential area of disease origin. Together with a previous study showing diminished grid cell activities in humans at AD risk (Kunz et al., 2015), our results also point to entorhinal cortex as a potential target for future diagnostic and therapeutic endeavor against AD.

In healthy animals, neurons in the entorhinal-hippocampal circuit exhibit two forms of remapping known as "global remapping" and "rate remapping" (Leutgeb et al., 2005b). When animals move across environments with distinct features, entorhinal grid cells and hippocampal place cells show distinct patterns of firing (global remapping). Global remapping of place cells and grid cells occur simultaneously (Fyhn et al., 2007), and inhibition of entorhinal grid cells scrambles remapping of hippocampal place cells (Miao et al., 2015). These evidences suggest that MEC input plays a crucial role in global remapping of hippocampal place cells. By contrast, when there is only a slight change in an environment, hippocampal place cells show change in their firing rate, while keeping the position of firing field constant (rate remapping). Because grid cells do not exhibit rate remapping (Fyhn et al., 2007), and because neurons in the hippocampal CA3 and dentate gyrus exhibit the most pronounced rate remapping (Leutgeb et al., 2005a; Leutgeb et al., 2007), rate remapping signal is suggested to be mainly generated inside dentate gyrus/CA3. Although it is likely that both global remapping and rate remapping are needed for high-precision spatial memory, disruption of global remapping would completely deprive AD patients of spatial information about which environment they are currently in. Our results showing the disruption of global remapping in the MEC and CA1 of APP-KI

mice clearly explain the devastating impairment of spatial memory in AD patients, who often exhibit wandering symptoms and cannot discriminate distinct features of surrounding environments.

Circuit components that cause the impairments of MEC neurons still remain to be elucidated. A previous study in heathy animals showed that inhibition of fast-spiking parvalbumin (PV)-expressing interneurons modifies firing patterns of grid cells to scattered patterns (Miao et al., 2017), a result very similar to what is observed in MEC neurons from our APP-KI mice. PV interneurons are vulnerable in the entorhinal cortex of AD patients (Solodkin et al., 1996). Thus, disruption of grid cells in our APP-KI mice might be caused by deterioration of PV interneurons. Future circuit-level investigations are expected to further pinpoint the circuit components causing memory loss in AD.

Acknowledgments: We thank Tomoko Viaclovsky, Kaori Shiraiwa, Anish Reddy, Allen Bramian and Ayushi Patel for technical assistance, and Mathew Blurton-Jones, Kim Green, Masashi Kitazawa, Robert Hunt and members of Igarashi lab for helpful discussion.

Funding: The work was supported by NIH R01 grant (R01AG063864), BrightFocus Foundation Research grant (A2019380S), Alzheimer's Association Research Grant (AARG-17-532932), Brain Research Foundation Fay-Frank Seed Grant (BRFSG-2017-04), Donors Cure Foundation New Vision Award (CCAD201902), PRESTO grant from Japan Science and Technology Agency (JPMJPR1681) and Whitehall Foundation Research Grant (2017-08-01) to K.M.I., and research grants from RIKEN Center for Brain Science, the Ministry of Education, Culture, Sports, Science and Technology, the Ministry of Health and Welfare, and AMED (JP18dm0207001, Brain Mapping by Integrated Neurotechnologies for Disease Studies (Brain/MINDS)) to T.C.S.

Author contributions: K.M.I. and H.J. designed experiments and analyses; H.J. and K.M.I. performed the experiments; H.J. and K.M.I. performed the analyses, with input from S.S.; T.S and T.C.S generated the APP-KI mice; H.J. and K.M.I wrote the paper with input from all authors.

Competing interests: The authors declare that they have no competing financial interests.

Data and materials availability: APP-KI mouse is available at RIKEN Bio Resource Center (RBRC06344). Neurophysiological data and analytical codes are available upon request.

Supplementary Materials:

Materials and Methods

Figures S1-S20

Tables S1-S14

FIGURE LEGENDS

Fig. 1. Hippocampal CA1 neurons showed severely disrupted remapping, but mildly diminished spatial tuning in APP-KI mice.

(**A**) A 64-channel recording device was targeted to the CA1 region of the hippocampus. (**B**) Sagittal sections with anti-Aβ immunostaining from 12-mo WT mouse (left) and 12-mo APP-KI mouse (middle), and with Nissl staining showing a representative recording position in CA1 (right, red arrow). A, anterior; P, posterior; D, dorsal; V, ventral. Scale bars, 500 μm. (**C**) 1x1 m open field for testing spatial tuning of CA1 neurons. (**D**) Four representative CA1 neurons in WT mice. Top: Red dots and gray lines denote spike position and animal trace in the 1x1 m open field, respectively. Bottom: Firing rate map. Color is scaled with maximum firing rate (Hz) shown at top right of each rate map. Spatial information (SI) score is shown at top left. (**E**) Four representative CA1 neurons as in (D), but from APP-KI mice. (**F**) Cumulative distribution plot for spatial information score of CA1 neurons in WT mice (blue, N=105) and APP-KI mice (red, N=113). A threshold of 0.63, obtained from 95th percentile of shuffled WT data, was used to define place cells. P<0.001, KS test. (**G**) Top: Mean spatial information score of CA1 neurons in WT mice (N=105) and APP-KI mice (N=113). P<0.001, Wilcoxon rank sum test. Bottom: Percentage of neurons defined as place cells. P<0.001, binomial test. (**H**) Remapping was tested in two 1-m linear Track A and Track B with distinct colors and textures. (**I**) Three representative CA1 neurons in WT mice recorded in Tracks A, B, B, and A (from Top to Bottom). Path (gray), spike position (red) and firing rate map are shown as in (D). Maximum firing rate is shown at top right. Spatial correlation between Track A and Track B (SC_{AB}) is shown at top left. (**J**) Three representative CA1 neurons from APP-KI mice are shown as in (I). (**K**) For evaluating the degree of remapping, population vector (PV) was constructed for each spatial bin (green) in Track A and Track B, and their correlation was analyzed (PV correlation). (**L**) Cumulative distribution plots for PV correlation between Track A and Track B (left) and between two recordings in Track A (right). Lower PV correlation denotes stronger remapping (black arrow). P<0.001, KS test.

Fig. 2. Fast gamma oscillations, but not slow gamma oscillations, were diminished in the hippocampal CA1 of APP-KI mice.

(**A**) Fast and slow gamma oscillations that reflect inputs from MEC and CA3, respectively, were recorded in CA1 (see text). (**B**) A representative time-resolved spectrogram showing gamma oscillation episodes in a WT mouse (left) and APP-KI mouse (right). Corresponding raw local field potential traces are shown at the bottom. (**C**) Normalized power spectra from WT and APP-KI mice (N=10 mice each). Significant difference in power for each 1Hz bin is shown with green dot at the top (q<0.05 at ranges of 5-8 Hz, 34-36 Hz and 62-100 Hz, FDR corrected for multiple comparison). (**D**) Example cross-frequency coherence plots showing that oscillatory powers at 20–40 Hz and 40–80 Hz

frequency (y-axis) were modulated by theta phase (x-axis) in a WT mouse (left), but the theta modulation of 40–80 Hz oscillatory power was absent in an APP-KI mouse (right). Dotted line denotes 40-Hz border between fast and slow gamma. (**E**) Occurrence rates of fast gamma episodes (top) and slow gamma episodes (bottom) for WT (N=10) and APP-KI mice (N=10). P<0.001, Wilcoxon ranks sum test. (**F**) Normalized power spectrograms averaged across all theta cycles in three representative WT mice (left) and APP-KI mice (right). Theta phase is shown at the bottom. Dotted line denotes 40-Hz border between fast and slow gammas. (**G**) Mean vector length for the distribution of fast gamma episodes (top) and slow gamma episodes (bottom) across theta phase. P<0.05, Wilcoxon ranks sum test. (**H**) Fast gamma phase distribution of spikes from example three CA1 neurons from WT mice (left) and APP-KI mice (right). Schematics of two cycles of gamma waveform are shown in the bottom. (**I**) Mean vector length for the distribution of spikes across fast gamma phase (top) and slow gamma phase (bottom) for WT CA1 neurons (N=104 cells) and APP-KI CA1 neurons (N=105 cells). P<0.05, Wilcoxon ranks sum test.

Fig. 3. Parallel disruptions of spatial tuning and remapping in MEC neurons of APP-KI mice.

(**A**) A 64-channel recording device was targeted to the layers 2/3 of the dorsal part of medial entorhinal cortex (MEC). (**B**) Sagittal sections with anti-Aβ immunostaining for 12-mo WT mouse (left) and 12-mo APP-KI mouse (right). (**C**) Sagittal sections with Nissl staining showing a representative recording position in the dorsal MEC (right, red arrow). Dotted lines denote the MEC. A, anterior; P, posterior; D, dorsal; V, ventral. Scale bars, 500 μm. (**D**) Four representative MEC neurons from WT mice recorded in the 1x1 m open field. Top: Red dots and gray lines denote spike position and animal trace, respectively. Middle: Firing rate map. Color is scaled with maximum firing rate (Hz) shown at top right of each rate map. Spatial information (SI) score is shown at top left. Bottom: Autocorrelograms of the firing rate maps shown for 2x2 m range. Color is scaled with correlation from r = -1 to 1. Gridness score (GS) is shown at top right. (**E**) Four representative MEC neurons as in (D), but from APP-KI mice. (**F**) Cumulative distribution plot for gridness score of MEC neurons in WT mice (blue, N=61) and APP-KI mice (red, N=65). A threshold of 0.41, obtained from 95th percentile of shuffled WT data, was used to define grid cells. P<0.05, KS test. (**G**) From left to right: Percentage of neurons defined as grid cells (P<0.001, binomial test), spatial information score (P<0.001, Wilcoxon rank sum test), firing field size (P<0.01, Wilcoxon rank sum test), and mean firing rate (P>0.05, Wilcoxon rank sum test). (**H**) Three representative MEC neurons from WT mice recorded in Tracks A, B, B, and A (from Top to Bottom). Path (gray), spike position (red) and firing rate map are shown as in (D). Maximum firing rate is shown at top right. Spatial correlation between Track A and B (SC$_{AB}$) is shown at top left. (**I**) Three representative MEC neurons from APP-KI mice are shown as in (H). (**J**) Cumulative distribution plots for PV correlation

between Track A and Track B (Top) and between two recordings in Track A (Bottom). Lower PV correlation denotes stronger remapping (black arrow). P<0.001, KS test. (**K**) Correlation between spatial tuning (spatial information score) and remapping (spatial correlation for Tracks A vs Track B). Each dot represents recorded neurons in CA1 (left) and MEC (right) of WT (top) and APP-KI mice (bottom). Only MEC neurons in APP-KI mice showed significant correlation (P<0.001, r (65) = -0.45, Pearson correlation).

Fig. 4. Impaired coordination of fast gamma oscillations between the MEC and CA1 in APP-KI mice.

(**A**) Fast gamma oscillations were recorded in the MEC. (**B**) Example cross-frequency coherence plots showing that 40–100 Hz oscillatory power (y-axis) was modulated by theta phase (x-axis) in a WT mouse (left), but this modulation was absent in an APP-KI mouse (right). Dotted line denotes 40-Hz border between fast and slow gamma. (**C**) Normalized power spectra for WT and APP-KI mice (N=13 mice each). No difference was observed in the power spectra (q>0.05, FDR corrected for multiple comparison). (**D**) Occurrence rates of fast gamma episodes in WT and APP-KI mice. (**E**) Normalized power spectrograms averaged across all theta cycles in three representative WT mice (left) and APP-KI mice (right). Theta phase is shown at the bottom. Dotted line denotes 40-Hz border between fast and slow gammas. (**F**) Mean vector length for the distribution of fast gamma episodes across theta phase. P=0.051, Wilcoxon ranks sum test. (**G**) Fast gamma phase distribution of spikes from example three MEC neurons in WT (left) and APP-KI (right) mice. Schematics of two cycles of gamma waveform are shown in the bottom. (**H**) Mean vector length for the distribution of spikes across fast gamma phase for WT CA1 neurons (N=60 cells) and APP-KI CA1 neurons (N=62 cells). P<0.01, Wilcoxon ranks sum test. (**I**) Fast gamma oscillations were recorded simultaneously in the MEC and hippocampal CA1. (**J**) Mean coherence plot of oscillatory activity as a function of frequency in WT and APP-KI mice (N=6 mice each). Significant difference in coherence for each 1Hz bin is shown with green dot at the top (q<0.05 at 58-75 Hz range, FDR corrected for multiple comparison). (**K**) Representative averaged local field potential traces from the MEC of WT mouse (top) and APP-KI mouse (bottom) during fast gamma episodes detected in CA1. t = 0 corresponds to the peak amplitudes of the fast gamma episodes in CA1. (**L**) MEC:CA1 power ratio during fast gamma episodes detected in CA1 (P<0.05, Wilcoxon rank sum test).

REFERENCES

Alme, C.B., Miao, C., Jezek, K., Treves, A., Moser, E.I., and Moser, M.B. (2014). Place cells in the hippocampus: eleven maps for eleven rooms. Proceedings of the National Academy of Sciences of the United States of America *111*, 18428-18435.

Axmacher, N., Henseler, M.M., Jensen, O., Weinreich, I., Elger, C.E., and Fell, J. (2010). Cross-frequency coupling supports multi-item working memory in the human hippocampus. Proceedings of the National Academy of Sciences of the United States of America *107*, 3228-3233.

Bragin, A., Jando, G., Nadasdy, Z., Hetke, J., Wise, K., and Buzsaki, G. (1995). Gamma (40-100 Hz) oscillation in the hippocampus of the behaving rat. J Neurosci *15*, 47-60.

Busche, M.A., Eichhoff, G., Adelsberger, H., Abramowski, D., Wiederhold, K.H., Haass, C., Staufenbiel, M., Konnerth, A., and Garaschuk, O. (2008). Clusters of hyperactive neurons near amyloid plaques in a mouse model of Alzheimer's disease. Science *321*, 1686-1689.

Cacucci, F., Yi, M., Wills, T.J., Chapman, P., and O'Keefe, J. (2008). Place cell firing correlates with memory deficits and amyloid plaque burden in Tg2576 Alzheimer mouse model. Proceedings of the National Academy of Sciences of the United States of America *105*, 7863-7868.

Chrobak, J.J., and Buzsaki, G. (1998). Gamma oscillations in the entorhinal cortex of the freely behaving rat. J Neurosci *18*, 388-398.

Colgin, L.L., Denninger, T., Fyhn, M., Hafting, T., Bonnevie, T., Jensen, O., Moser, M.B., and Moser, E.I. (2009). Frequency of gamma oscillations routes flow of information in the hippocampus. Nature *462*, 353-357.

Colgin, L.L., Moser, E.I., and Moser, M.B. (2008). Understanding memory through hippocampal remapping. Trends in neurosciences *31*, 469-477.

Deshmukh, S.S., Yoganarasimha, D., Voicu, H., and Knierim, J.J. (2010). Theta modulation in the medial and the lateral entorhinal cortices. J Neurophysiol *104*, 994-1006.

Doeller, C.F., Barry, C., and Burgess, N. (2010). Evidence for grid cells in a human memory network. Nature *463*, 657-661.

Ekstrom, A.D., Kahana, M.J., Caplan, J.B., Fields, T.A., Isham, E.A., Newman, E.L., and Fried, I. (2003). Cellular networks underlying human spatial navigation. Nature *425*, 184-188.

Fu, H., Rodriguez, G.A., Herman, M., Emrani, S., Nahmani, E., Barrett, G., Figueroa, H.Y., Goldberg, E., Hussaini, S.A., and Duff, K.E. (2017). Tau Pathology Induces

Excitatory Neuron Loss, Grid Cell Dysfunction, and Spatial Memory Deficits Reminiscent of Early Alzheimer's Disease. Neuron *93*, 533-541 e535.

Fyhn, M., Hafting, T., Treves, A., Moser, M.B., and Moser, E.I. (2007). Hippocampal remapping and grid realignment in entorhinal cortex. Nature *446*, 190-194.

Fyhn, M., Molden, S., Witter, M.P., Moser, E.I., and Moser, M.B. (2004). Spatial representation in the entorhinal cortex. Science *305*, 1258-1264.

Gomez-Isla, T., Price, J.L., McKeel, D.W., Jr., Morris, J.C., Growdon, J.H., and Hyman, B.T. (1996). Profound loss of layer II entorhinal cortex neurons occurs in very mild Alzheimer's disease. J Neurosci *16*, 4491-4500.

Hope, T., Tilling, K.M., Gedling, K., Keene, J.M., Cooper, S.D., and Fairburn, C.G. (1994). The Structure of Wandering in Dementia. Int J Geriatr Psych *9*, 149-155.

Igarashi, K.M., Lu, L., Colgin, L.L., Moser, M.B., and Moser, E.I. (2014). Coordination of entorhinal-hippocampal ensemble activity during associative learning. Nature *510*, 143-147.

Kunz, L., Schroder, T.N., Lee, H., Montag, C., Lachmann, B., Sariyska, R., Reuter, M., Stirnberg, R., Stocker, T., Messing-Floeter, P.C.*, et al.* (2015). Reduced grid-cell-like representations in adults at genetic risk for Alzheimer's disease. Science *350*, 430-433.

Kyle, C.T., Stokes, J.D., Lieberman, J.S., Hassan, A.S., and Ekstrom, A.D. (2015). Successful retrieval of competing spatial environments in humans involves hippocampal pattern separation mechanisms. Elife *4*.

Langston, R.F., Ainge, J.A., Couey, J.J., Canto, C.B., Bjerknes, T.L., Witter, M.P., Moser, E.I., and Moser, M.B. (2010). Development of the spatial representation system in the rat. Science *328*, 1576-1580.

Leutgeb, J.K., Leutgeb, S., Moser, M.B., and Moser, E.I. (2007). Pattern separation in the dentate gyrus and CA3 of the hippocampus. Science *315*, 961-966.

Leutgeb, J.K., Leutgeb, S., Treves, A., Meyer, R., Barnes, C.A., McNaughton, B.L., Moser, M.B., and Moser, E.I. (2005a). Progressive transformation of hippocampal neuronal representations in "morphed" environments. Neuron *48*, 345-358.

Leutgeb, S., Leutgeb, J.K., Barnes, C.A., Moser, E.I., McNaughton, B.L., and Moser, M.B. (2005b). Independent codes for spatial and episodic memory in hippocampal neuronal ensembles. Science *309*, 619-623.

Mably, A.J., Gereke, B.J., Jones, D.T., and Colgin, L.L. (2017). Impairments in spatial representations and rhythmic coordination of place cells in the 3xTg mouse model of Alzheimer's disease. Hippocampus *27*, 378-392.

Masuda, A., Kobayashi, Y., Kogo, N., Saito, T., Saido, T.C., and Itohara, S. (2016). Cognitive deficits in single App knock-in mouse models. Neurobiol Learn Mem *135*, 73-82.

Miao, C., Cao, Q., Ito, H.T., Yamahachi, H., Witter, M.P., Moser, M.B., and Moser, E.I. (2015). Hippocampal Remapping after Partial Inactivation of the Medial Entorhinal Cortex. Neuron *88*, 590-603.

Miao, C., Cao, Q., Moser, M.B., and Moser, E.I. (2017). Parvalbumin and Somatostatin Interneurons Control Different Space-Coding Networks in the Medial Entorhinal Cortex. Cell *171*, 507-521 e517.

Morris, R.G., Garrud, P., Rawlins, J.N., and O'Keefe, J. (1982). Place navigation impaired in rats with hippocampal lesions. Nature *297*, 681-683.

Moser, M.B., Rowland, D.C., and Moser, E.I. (2015). Place cells, grid cells, and memory. Cold Spring Harb Perspect Biol *7*, a021808.

Muller, R.U., and Kubie, J.L. (1987). The effects of changes in the environment on the spatial firing of hippocampal complex-spike cells. J Neurosci *7*, 1951-1968.

O'Keefe, J., and Dostrovsky, J. (1971). The hippocampus as a spatial map. Preliminary evidence from unit activity in the freely-moving rat. Brain research *34*, 171-175.

Quilichini, P., Sirota, A., and Buzsaki, G. (2010). Intrinsic circuit organization and theta-gamma oscillation dynamics in the entorhinal cortex of the rat. J Neurosci *30*, 11128-11142.

Saito, T., Matsuba, Y., Mihira, N., Takano, J., Nilsson, P., Itohara, S., Iwata, N., and Saido, T.C. (2014). Single App knock-in mouse models of Alzheimer's disease. Nature neuroscience *17*, 661-663.

Sargolini, F., Fyhn, M., Hafting, T., McNaughton, B.L., Witter, M.P., Moser, M.B., and Moser, E.I. (2006). Conjunctive representation of position, direction, and velocity in entorhinal cortex. Science *312*, 758-762.

Sasaguri, H., Nilsson, P., Hashimoto, S., Nagata, K., Saito, T., De Strooper, B., Hardy, J., Vassar, R., Winblad, B., and Saido, T.C. (2017). APP mouse models for Alzheimer's disease preclinical studies. EMBO J *36*, 2473-2487.

Scoville, W.B., and Milner, B. (1957). Loss of recent memory after bilateral hippocampal lesions. Journal of neurology, neurosurgery, and psychiatry *20*, 11-21.

Skaggs, W.E., McNaughton, B.L., Gothard, K.M., and Markus, E.J. (1993). An information-theoretic approach to deciphering the hippocampal code. . In Advances in

Neural Processing Systems, Vol 5 Eds: SJ Hanson, JD Cowan & CL Giles (San Mateo, Morgan Kaufmann), pp. 1030-1037.

Solodkin, A., Veldhuizen, S.D., and Van Hoesen, G.W. (1996). Contingent vulnerability of entorhinal parvalbumin-containing neurons in Alzheimer's disease. J Neurosci *16*, 3311-3321.

Steffenach, H.A., Witter, M., Moser, M.B., and Moser, E.I. (2005). Spatial memory in the rat requires the dorsolateral band of the entorhinal cortex. Neuron *45*, 301-313.

Tort, A.B., Komorowski, R.W., Manns, J.R., Kopell, N.J., and Eichenbaum, H. (2009). Theta-gamma coupling increases during the learning of item-context associations. Proceedings of the National Academy of Sciences of the United States of America *106*, 20942-20947.

Van Hoesen, G.W., Hyman, B.T., and Damasio, A.R. (1991). Entorhinal cortex pathology in Alzheimer's disease. Hippocampus *1*, 1-8.

Witter, M.P., and Amaral, D.G. (2004). Hippocampal Formation. In The Rat Nervous System 3rd edn (ed G Paxinos), G. Paxinos, ed. (Amsterdam, The Netherlands: Elsevier).

Yassa, M.A., and Stark, C.E. (2011). Pattern separation in the hippocampus. Trends in neurosciences *34*, 515-525.

Fig1

Fig. 2

Fig. 3

A CA3 CA1 Temporoammonic Pathway DG Hippocampus Perforant Pathway L3 L2 Medial Entorhinal Cortex (MEC) L2/3

B Aβ (WT) 500 µm MEC Aβ (APP-KI) 500 µm MEC

C Nissl MEC 500 µm D A P V

F % MEC Neurons — Gridness Score — 0.41 APP-KI WT *p<0.05 Grid cells

D WT
Path/Spike SI: 0.71 5.1 Hz SI: 0.43 7.4 Hz SI: 0.33 4.7 Hz SI: 1.1 5.1 Hz
Firing Rate Max(FR) 0
Autocorrelation GS: 0.63 GS: 0.81 GS: 0.54 GS: 0.83 r = 1 r = -1

E APP-KI
Path/Spike SI: 0.042 5.1 Hz SI: 0.090 5.9 Hz SI: 0.15 1.8 Hz SI: 0.46 1.0 Hz
Firing Rate Max(FR) 0
Autocorrelation GS: -0.12 GS: 0 GS: 0.23 GS: 0.28 r = 1 r = -1

G
% Grid Cells	Spatial Information	Field Size (cm²)	Mean Firing Rate (Hz)
*p<0.001	*p<0.001	*p<0.01	N.S.
20	0.75	x10³ 3 2 1	1.5 1 0.5
WT APP-KI	61 65	61 65	61 65

H WT
A SC_AB: -0.43 3.1Hz SC_AB: -0.11 13Hz SC_AB: -0.29 6.5Hz
B
B
A

I APP-KI
A SC_AB: 0.03 2.1Hz SC_AB: 0.03 3.7Hz SC_AB: 0.55 5.1Hz
B
B
A

J Tracks A vs B — % Population Vector — WT APP-KI Stronger Remapping *p<0.001
Tracks A vs A — WT APP-KI N.S. — PV Correlation

K
WT CA1 $r(105) = -0.15$ $P = 0.14$
WT MEC $r(61) = -0.22$ $P = 0.09$
APP-KI CA1 $r(113) = -0.14$ $P = 0.22$
APP-KI MEC $r(65) = -0.45$ $P < 0.001$
Spatial Correlation (Tracks A vs B) — Spatial Information Score

Fig 4

Supplementary materials for

Disrupted remapping of place cells and grid cells

5 # in knock-in model of Alzheimer's disease

Heechul Jun, Shogo Soma, Takashi Saito, Takaomi C Saido, Kei M Igarashi

Correspondence to: kei.igarashi@uci.edu

This file includes:

15   Materials and Methods
        Supplementary Figs. S1 to S20
        Tables S1 to S14

Materials and Methods

Subjects

Mice were maintained in standard housing conditions on a 12h dark/light cycle with food and water provided ad libitum. All procedures were conducted in accordance with the
5 guidelines of the National Institutes of Health and approved by the Institutional Animal Care and Use Committee at the University of California, Irvine. We previously generated $App^{NL-G-F/NL-G-F}$ maintained on a C57BL/6 background (Saito et al., 2014). For all experiments, eighteen $App^{NL-G-F/NL-G-F}$ mice (fourteen males and four females) and seventeen $App^{WT/WT}$ mice (eleven males and six females) between 7 months and 13
10 months of age were used. We noted no difference in our data between sexes. Detailed information about ages when the recording was performed is in Supplementary Table S1. Animals were housed in a reversed 12h light - dark cycle, and all testing occurred during the dark phase. If the animals died or were sick during, or before the final test, their recording positions could not be validated and therefore were removed from the data
15 analysis.

Surgery and Electrode Preparation

All mice received a custom-built 64-channel drive modified from Liang et al (Liang et al., 2017) targeting either CA1 or MEC. For a group of animals, eight tetrodes targeted CA1 of hippocampus and other eight tetrodes targeted superficial layer of medial entorhinal
20 cortex. Tetrodes were constructed from four twisted 17 µm polyimide-coated platinum-iridium (90%–10%) wires (California Fine Wire). The electrode tips were plated with gold to reduce electrode impedances to between 150 and 300 kΩ at 1 kHz. Animals were anesthetized with isoflurane (air flow: 0.8–1.0 l/min, 1% isoflurane, adjusted according to physiological condition). The mice received subcutaneous injections of buprenorphine at
25 the start of the surgery. Depth of anesthesia was examined by testing tail and pinch reflexes as well as breathing. Upon induction of anesthesia, the animal was fixed in a Kopf stereotaxic frame for implantation. A craniotomy was performed centering at AP 2.5 mm ML 2.5 mm from Bregma for hippocampus (approximately 1 x 1mm) and 0.2-0.4mm anterior to transverse sinus and ML 3.5mm from midline for medial entorhinal
30 cortex (approximately 1 x 1mm). The dura was carefully removed, and the tetrodes were implanted. A stainless-steel screw fixed to skull above the cerebellum served as the ground electrode. The recording drive was secured to the scratched skull using dental cement.

Apparatus and Training Procedures

Behavioral training started at least 5 days after the surgery and data collection was performed exactly on the fifth day of the training. All experiments were conducted in a room containing 1m-square box made of black acryl and a polarizing white cute card (216 mm x 280mm) and two 1-m long linear tracks. In the open field task, running was motivated in the 1m-square box with cookie crumbs thrown randomly into the enclosure. Each session lasted 10 to 20 min. Upon finishing the open field task, animals were rested for 5 minutes and they were tested in the remapping task. The remapping task training required mice to run in 1-m long linear tracks. Two 1-m long linear tracks were used: (Track A) Black acryl enclosure with inner wall decorated with patterns of white cue cards across the track and black rubber floor. (Track B) White acryl enclosure with inner wall decorated with distinct patterns of black cue signs across the track and white sandpaper floor. During remapping task, animals were trained to run in successive series of Track A → Track B → Track B → Track A. Between the sessions, animals were given 5 minutes rest. Running was motivated by placing cookie crumbs on the end positions of the linear tracks. Each session lasted 5 to 10 min. On the linear tracks, the mice ran 10 full laps (back and forth).

Data Collection

LFPs were recorded single-ended using a ground in the skull above the cerebellum. One mouse without single-ended recording was excluded for LFP analyses. The tetrodes were connected to a multichannel, impedance matching, unity gain headstage (Neuralynx). The output of the headstage was conducted to a data acquisition system (Neuralynx). Unit activity was amplified by a factor of 3000–5000 and band-pass filtered from 600 Hz to 6000 Hz. Spike waveforms above a threshold set by the experimenter (~55 µV) were time-stamped and digitized at 32 kHz for 1 ms. LFP signals, 1 per tetrode, were recorded in the 0–475-Hz frequency band at a sampling rate of 2000 Hz. Notch filters were not applied. The recording system tracked the position of two light-emitting diodes (LEDs), one large and one small, on the head stage (sampling rate 50 Hz) by means of an overhead video camera

Data Analysis

Unless indicated otherwise, analyses were performed using MATLAB codes kindly provided by Dr. Edvard Moser, or written by the authors (Igarashi et al., 2014)

Spike Analysis

Spike Sorting and Cell Classification. Spike sorting was performed offline using graphical cluster-cutting software, MClust (A.D. Redish). Putative excitatory cells were

distinguished from putative interneurons by spike peak-valley width and average rate. (Bartho et al., 2004). All putative excitatory cells with spike peak-valley width of more than 230 μs and mean firing rates of more than 0.1 Hz (mean firing rate during all left and right cue sampling intervals) were included for further analysis.

Firing rate maps. Position estimates were based on tracking of the LEDs on the head stage connected to the recording drive. For rate maps in the random foraging task, data were speed-filtered; only epochs with instantaneous running speeds of 1 cm/s or more were included.

10 To characterize firing fields, the position data were sorted into one-dimensional 1cm bins for the linear tracks and 2.5 cm × 2.5 cm bins for the open field. For the open field map, the path was smoothed with a 21-sample boxcar window filter (400 ms; 10 samples on each side). Firing rate distributions were then determined by counting the number of spikes in each bin as well as the time spent per bin. Maps for number of spikes and time

15 were smoothed individually using a boxcar average over the surrounding 5 × 5 bins. Weights were distributed as follows:

box =   [0.0025 0.0125 0.0200 0.0125 0.0025;…

0.0125 0.0625 0.1000 0.0625 0.0125;…

20 0.0200 0.1000 0.1600 0.1000 0.0200;…

0.0125 0.0625 0.1000 0.0625 0.0125;…

0.0025 0.0125 0.0200 0.0125 0.0025;]

Spatial information score. For each cell, the spatial information score in bits per spike was
25 calculated from the recordings in the open field task as

$$ \textit{Spatial Information Score} = \sum_i p_i \frac{\lambda_i}{\lambda} \log_2 \frac{\lambda_i}{\lambda} $$

where λ_i is the mean firing rate of a unit in the ith bin, λ is the overall mean firing rate,
30 and p_i is the probability of the animal being in the ith bin (occupancy in the ith bin / total recording time) (Skaggs et al., 1993). An adaptive smoothing method, introduced by

Skaggs *et al.* (Skaggs et al., 1996), was used before the calculation of information scores (Henriksen et al., 2010).

<u>Defining Place Cells.</u> Place cells were defined as cells with spatial information score above the chance level (Langston et al., 2010). The chance level was determined in each brain region by a random permutation procedure using all cells recorded at that time-point in that region. One hundred permutations were performed for each cell in the sample. For each permutation trial, the entire sequence of spikes fired by the cell was time-shifted along the animal's path by a random interval between 20 s and 20 s less than the total length of the trial (usually 600 −20 = 580 s), with the end of the trial wrapped to the beginning to allow for circular displacements. This procedure allowed the temporal firing structure to be retained in the shuffled data at the same time as the spatial structure was lost. Spatial information was then calculated for each shuffled map. The distribution of spatial information values across all 100 permutations of all cells in the sample was computed and finally the 95th percentile was determined. Place cells were defined as cells with spatial information scores above the 95th percentile of the distribution from shuffled data for the relevant group.

<u>Gridness score</u>. The structure of all rate maps was evaluated for all cells by calculating the spatial autocorrelation for each smoothed rate map (Sargolini et al., 2006; Fyhn et al., 2007). Autocorrelograms were based on Pearson's product moment correlation coefficient with corrections for edge effects and unvisited locations. With λ *(x, y)* denoting the average rate of a cell at location *(x, y)*, the autocorrelation between the fields with spatial lags of τ_x and τ_y was estimated as:

$$r(\tau_x, \tau_y) = \frac{n \sum \lambda(x, y) \lambda(x - \tau_x, y - \tau_y) - \sum \lambda(x, y) \sum \lambda(x - \tau_x, y - \tau_y)}{\sqrt{n \sum \lambda(x, y)^2 - \left(\sum \lambda(x, y)\right)^2} \sqrt{n \sum \lambda(x - \tau_x, y - \tau_y)^2 - \left(\sum \lambda(x - \tau_x, y - \tau_y)\right)^2}}$$

where the summation is over all n pixels in λ *(x, y)* for which rate was estimated for both λ *(x, y)* and λ *(x − τ_x, y − τ_y)*. Autocorrelations were not estimated for lags of τ_x, τ_y where $n < 20$.

The degree of spatial periodicity ('gridness score') was determined for each recorded cell by taking a circular sample of the autocorrelogram, centered on the central peak but with the central peak excluded, and comparing rotated versions of this sample. The Pearson correlation of this circle with its rotation in α degrees was obtained for angles of 60° and 120° on one side and 30°, 90° and 150° on the other. The cell´s gridness score was

defined as the minimum difference between any of the elements in the first group and any of the elements in the second. The radius of the excluded central peak was defined as either the first local minimum in a curve showing correlation as a function of average distance from the center, or as the first incidence where the correlation was negative, whichever occurred first.

Defining Grid Cells. Grid cells were defined as cells in which rotational symmetry-based gridness scores exceeded the 95th percentile of a distribution of grid scores for shuffled recordings from the entire population of cells. Shuffling was performed in the same way as for place cells, i.e. for each permutation trial, the entire sequence of spikes fired by the cell was time-shifted along the animal's path by a random interval between 20 s and 20 s less than the length of the trial (usually 600 – 20 = 580 s), with the end of the trial wrapped to the beginning. The shuffling procedure was repeated 100 times for each of the cells in the sample. For each permutation, a firing rate map and an autocorrelation map were constructed and a grid score was calculated. Grid scores were computed for all permutations for all cells and the 95th percentile was read out from the overall distribution. Grid cells were defined as cells with grid scores higher than the 95th percentile of the grid scores of the distribution for the shuffled data.

Remapping Analysis. Remapping analysis was performed as described previously in Miao et al (Miao et al., 2015). For experiments on the linear track, instantaneous firing rates on individual laps were estimated using a Gaussian kernel on the spike data for temporal smoothing:

$$r(t) = \sum_{i=1}^{N} g\left(\frac{t_i - t}{n}\right)$$

where g is a 1D Gaussian kernel, h is a bandwidth, N is the total number of spikes, and t_i is the time of the i^{th} spike. The bandwidth was set at 100 ms. Due to minimal coverage per bin, temporal smoothing gives more robust rate estimates compared to those based on spatial bins during fast running on the track. Spike rate at each track position (1 cm bin) on each lap was estimated using a linear interpolation method applied to the temporally-smoothed spike rates. Because the same cells often had different firing fields on left and right runs, firing fields of the same cell in each run direction were analyzed as two distinct data sets. A segment of 5 cm was excluded from each end of the linear track for analysis.

Spatial correlation was obtained by calculating the Pearson correlation coefficient for mean firing rates across 1 cm wide bins from the track on a pair of sessions. For population vector analysis, population vectors were defined for each 3.6-cm bin of rate maps (25 bins in total) from all cells in the experimental group. To maintain independence of neighboring spatial bins, we estimated mean firing rates in spatial bins without smoothing. For lap-by-lap population vector analyses, a total of 10 laps (defined as pairs of forward and backward runs) were taken from each recording session. Population vector correlations were determined for each spatial bin of each lap.

LFP analysis

Animals with single-ended LFP recording were used for the analysis.

Time–frequency analysis of power. Time-resolved power across frequencies was computed using a wavelet transform method. Signals were convolved by a family of complex Morlet's wavelets $w(t, f)$, one for each frequency, as a function of time:

$$w(t, f) = A \exp(- t^2/\sigma_t^2) \exp (2i\pi ft)$$

in which the family of wavelets was characterized by a constant ratio f/σ_f, which was set to 7, and with $\sigma_f = \frac{1}{2}\pi\sigma_t$. The coefficient A was set at:

$$(\sigma_t\sqrt{\sigma})^{-1/2}$$

in order to normalize the wavelets such that their total energy was equal to 1.

Normalization of LFP power. To control for impedance differences between tetrodes, LFP power was normalized, for each tetrode, to the total power in 1–100 Hz band.

Bandpass filtering. An acausal (zero phase shift), frequency domain, finite impulse response bandpass filter was applied to the signals. For theta filtering, 5 and 6 Hz were chosen for the stopband and passband, respectively, for the low cut-off frequencies; 10 and 11 Hz were chosen for passband and stopband high cut-off frequencies. For filtering of gamma oscillations, 19 and 20 Hz were chosen for the stopband and passband, respectively, for the low cut-off frequencies; 90 and 91 Hz were chosen for passband and stopband high cut-off frequencies. For filtering of slow gamma oscillations, 19 and 20 Hz were chosen for the stopband and passband, respectively, for the low cut-off frequencies;

40 and 41 Hz were chosen for passband and stopband high cut-off frequencies. For filtering of fast gamma oscillations, 39 and 40 Hz were chosen for the stopband and passband, respectively, for the low cut-off frequencies; 90 and 91 Hz were chosen for passband and stopband high cut-off frequencies.

Selection of theta cycles. Theta cycles were selected by bandpass filtering the signal from 6-10 Hz and selecting local minima in the filtered signal (i.e., $\theta(t-1) > \theta(t)$ and $\theta(t+1) > \theta(t)$). Segments of the recording were collected and defined as a theta cycle if the time between detected points fell within a range criterion that corresponded to the period of an ~8 Hz theta cycle (i.e. 125 ± 25 ms). Local minima of detected theta cycles were required to be separated by at least 100 ms.

Detection of oscillation episodes. To extract periods of gamma oscillatory activity in the LFP, we first computed time-varying power within the frequency bands for each recording. Power at each time point was averaged across the frequency range to obtain time-varying estimates of oscillatory power. Time points were collected when the power exceeded 2 SD of the time-averaged power. Time windows, 160 ms in length, were cut around the identified time points. In each 160-ms segment, the maxima of gamma oscillatory amplitude were determined from the gamma bandpass filtered versions of the recordings. Duplicated gamma oscillatory periods, a common consequence of extracting overlapping time windows, were avoided by discarding identical maxima values within a given gamma oscillatory subtype and further requiring that maxima of a given subtype be separated by at least 100 ms. Individual gamma oscillatory windows were finally constructed from the original, non-bandpass filtered recordings as 400 ms long windows centered around the gamma oscillatory amplitude maxima.

Relationship of gamma to theta phase. LFP recordings were bandpass-filtered in the theta range (6–10 Hz), and theta phases for each time point were estimated using the Hilbert transform function from the Signal Processing Toolbox in MATLAB. Theta phases at the time points associated with gamma maxima (determined as described above) as well as theta phases for spikes were collected. Theta phases for each gamma oscillatory event were sorted into 30° bins, allowing the phase distribution of each event to be determined. For a given recording, the distributions of gamma oscillations were normalized by dividing the bins by the total number of gamma oscillatory episodes within a given recording. In this analysis, and in all analyses involving oscillation phase, the oscillation peak was defined as 0°.

Time–frequency representation of power across individual theta cycles. Time-varying power in 2-Hz-wide frequency bands, from 2 Hz to 140 Hz, was obtained for individual theta cycles using the wavelet transform method described above. Time frequency representations for multiple theta cycles recorded from the same site and session within the same animal were then averaged.

Strength of theta-gamma coupling. The theta phase at the time of gamma oscillation maxima occurrence was calculated by bandpass-filtering the LFP in the theta range, performing a Hilbert transform on the filtered signal, and then locating theta phase of individual gamma oscillation maxima. Resultant vectors were calculated from the phase distributions gamma maxima. Lengths of resultant vectors were used as an index for the strength of theta-gamma coupling.

Phase-locking of spikes to gamma oscillations and theta oscillations. Neurons recorded in animals with single-ended LFP recording were used for the analysis. The oscillatory phase at the time of spike occurrence was estimated. This was performed by first bandpass-filtering the LFP, performing a Hilbert transform on the filtered signal, and then extracting the phase component at the spike times. Cells were considered to be phase-locked if their phase distribution differed significantly from a uniform distribution ($P < 0.05$, Rayleigh test). Resultant vectors were calculated from the distributions of phase component. Lengths of resultant vectors were used as an index for the strength of phase-locking.

Histology and Reconstruction of Recording Positions

Electrode positions were confirmed by anaesthetizing the drive-implanted mice using isoflurane and performing small electrolytic lesions through passing the current (10 µA for 20 s) through the electrodes. Immediately after this, the mice received an overdose of isoflurane and were perfused intracardially with saline followed by 4% freshly depolymerized paraformaldehyde in phosphate buffer (PFA). The brains were extracted and stored in the same fixative overnight. After overnight cryoprotection in phosphate buffer saline with 30% sucrose at 4°C, tissue samples were embedded in O.C.T. mounting medium and sagittal sections (40 µm) were cut and stained with cresyl violet. All tetrodes were identified, and the tip of each electrode was found by comparison with adjacent sections. Only data from tetrodes in the hippocampal CA1 or the layer 2/3 of dorsal MEC was collected for analysis.

For immunostaining, sections were rinsed 3 times for 10 min in 1 x PBS (pH 7.6) at room temperature, preincubated for 1 hr in 10% normal goat serum in PBST (1 x PBS with 0.5% Triton X-100). Between incubation steps, sections were rinsed in PBST. Sections

5 were incubated with antibodies against Aβ, raised in mouse (MCSA1, Medimabs, 1:1000) for 24 hr in antibody-blocking buffer at 4°C. After three times of 15 min washing in PBST at room temperature, sections were incubated in a goat-anti mouse antibody conjugated with Alexa Fluor 568nm (Invitrogen, 1:250) for 2 hr at room temperature. After rinsing in PBS, sections were mounted onto glass slides with 40 ,60 -diamidino-2-phenylindole

10 (DAPI)-containing Vectashield mounting medium (Vector Laboratories), and a coverslip was applied. Digital photomicrographs were acquired with a Olympus BX53 fluorescent microscope equipped with a digital camera.

Statistical Analysis

15 All statistical testing assumed a non-parametric distribution and Wilcoxon rank sum test was used. For comparing the distribution of place cells and grid cells, Kolmogorov-Smirnov tests were applied. All statistical methods used are summarized in Supplementary Tables S2-S14.

Fig. S1. Schematics for parallel disruptions of entorhinal grid cells and hippocampal place cells

(**A**) In the entorhinal-hippocampal circuit of healthy brain, neurons in layers 2 and 3 of the medial entorhinal cortex (MEC) show spatially tuned neuronal activity, and exhibit remapping across distinct environments. Pyramidal cells (Pyr) in the layers 2 and 3 send these signals to neurons in the CA1 region of the hippocampus. In the hippocampal CA1, neurons show spatial activity as place cells and exhibit remapping. Stellate cells (St) in MEC layer 2 send information to dentate gyrus and CA3. (**B**) In this study, we determined whether (1) remapping of hippocampal place cells is deteriorated in AD subjects, and if so, (2) whether place cell deterioration is linked to impairment of entorhinal grid cells, or independent from grid cells. We found that hippocampal CA1 neurons showed severely disrupted remapping, but mildly diminished spatial tuning in APP-KI mice. We also found that entorhinal grid cells were severely impaired, sending disrupted remapping signals to the hippocampus. These results point to the link between grid cell impairment and remapping disruption as an underlying circuit mechanism causing spatial memory loss in AD patients, who often cannot discriminate distinct features of surrounding environments.

Fig. S2. Electrophysiological recording of principal neurons in CA1 and MEC

(**A**) Multi-electrode 64-channel drive used for CA1 and MEC neural recordings. Inset, magnification image showing two separate bundles containing tetrodes. (**B**) Surgical scheme illustrating the location of tetrodes implanted for CA1 of hippocampus and layer II/III of medial entorhinal cortex. (**C**) Example of spike sorting in energy space to identify single units across the sessions, demostrating stable chronic recording across the sessions. (**D**) Two cluster of spikes and their corresponding spike wave forms from identified clusters representing two single units.

Fig. S3. Histological validation of recording position in CA1 of WT and APP-KI mice

Brightfield images of cresyl violet stained sections of all animals recorded. Arrowhead points to the tip of tetrode where the cells were recorded with electrolytic lesions made before sacrificing the animals.

Fig. S4. Spatial Tuning and remapping property across age (7 - 13 months old)

(**A-D**), Spatial information score and spatial correlation between Track A and Track B were plotted for CA1 neurons and MEC neurons across age (7-13 months old). No significant effects of aging were observed in APP-KI mice (P>0.05, Pearson correlation).

A weak positive correlation was observed for spatial correlation in WT MEC neurons (r(61) = -0.30, P<0.05, Pearson correlation).

Fig. S5. Running behaviors did not differ between WT and APP-KI mice.

(**A**) Speed analysis during open field exploration. No significant difference between WT and APP-KI mice (q>0.05, FDR corrected for multiple comparison). (**B**) Speed analysis during linear track experiment. No significant difference between WT and APP-KI mice (q>0.05, FDR corrected for multiple comparison).

Fig. S6. Electrophysiological features for classifying putative principal neurons in CA1 and MEC

(**A-B**) Peak to valley time of spike waveform was used to distinguish putative interneurons from principal neurons. Dashed line (230μs) represents cut off for interneuron to principal neuron classification (Bartho et al., 2004). (**A**) Peak to valley time histogram for neurons recorded from CA1. No difference was observed for the distribution of peak to valley time between WT and APP-KI mice (Kolmogorov Smirnov test, p >0.05). (**B**) Peak to valley time histogram for neurons recorded from MECs. No difference was observed for the distribution of peak to valley time between WT and APP-KI mice (Kolmogorov Smirnov test, p >0.05).

Fig. S7. Additional properties of CA1 neurons during the open Field task

(**A**) Representative CA1 neuron recorded from WT and APP-KI animal during open field task. Spatial information (SI) and average firing rate (Hz) for each cell is shown above rate map. (**B**) Field area (cm^2) measured from CA1 neurons in open field task. (Wilcoxon-rank-sum, p>0.05) (**C**) Number of fields measured from CA1 neurons in open field task. (Wilcoxon-rank-sum, p>0.05) (**D**) Spike width measured from CA1 neurons in open field task. (Wilcoxon-rank-sum, p>0.05) (**E**) Average firing rate measured from CA1 neurons in open field task. (Wilcoxon-rank-sum, p>0.05) (**F**) Peak firing rate measured from CA1 neurons in open field task. (Wilcoxon-rank-sum, p>0.05)

Fig. S8. Properties of CA1 neurons defined as place cells in WT and APP-KI mice

(**A**) Top, distribution of spatial information scores calculated from firing rate distributions of CA1 neurons in the open field task (real data). Bottom, distribution of shuffled data based on 100 permutations per cell. Red dashed line indicates 95th percentile value (chance level) for a distribution based on all permutations. Percentage of cells that

exceeded chance level is shown on top. 50% of CA1 neurons in WT mice and 31% of CA1 neurons in APP-KI mice passed this chance level, and they were considered as place cells.

(**B**) Field area (cm^2) measured from place cells of CA1 (Wilcoxon-rank-sum test, p>0.05).

(**C**) Number of fields measured from place cells (Wilcoxon-rank-sum test, p>0.05). (**D**) Average firing rate measured from place cells (Wilcoxon-rank-sum test, p>0.05). (**E**) Peak firing rate measured from place cells (Wilcoxon-rank-sum test, p>0.05). (**F**) Spatial information measured from place cells. (Wilcoxon-rank-sum test, p>0.05). (**G-J**) Properties in the linear tracks. (**G**) Field area (cm^2) measured from place cells (Wilcoxon-rank-sum test, p>0.05). (**H**) Number of fields measured from place cells (Wilcoxon-rank-sum test, P>0.05). (**I**) Average firing rate measured from place cells (Wilcoxon-rank-sum test, p>0.05). (**J**) Peak firing rate measured from place cells (Wilcoxon-rank-sum test, p>0.05) (**K**) Cumulative distribution plot of PV correlation between Track A and Track B from place cells. Significant increase in PV correlation was observed compared to that of place cells in APP-KI mice (Wilcoxon-rank-sum test, p <0.001). (**L**) Cumulative distribution plot of PV correlation between two recordings in Track A (Wilcoxon-rank-sum test, p <0.01). (**M**) Cumulative distribution plot of Spatial Correlation between Track A and Track B (Wilcoxon-rank-sum test, p <0.01). (**N**) Cumulative distribution plot of Spatial Correlation between two recordings in Track A (Wilcoxon-rank-sum test, p <0.01).

Fig. S9. Additional properties of CA1 neurons during the remapping task

(**A**) Representative CA1 neuron recorded from WT and APP-KI mice during remapping task. Spatial correlation between Track A and Track B (SC$_{AB}$) and average firing rate for each cell is shown above for each cell. (**B-G**) Properties of CA1 neurons in the linear tracks (**B**) Field area (cm^2) (Wilcoxon-rank-sum, p>0.05). (**C**) Number of fields (Wilcoxon-rank-sum, p>0.05). (**D**) Average firing rate (Wilcoxon-rank-sum, p>0.05). (**E**) Peak firing rate (Wilcoxon-rank-sum, p>0.05) (**F**) Averaged PV correlation between two recordings in Track A (Wilcoxon-rank-sum, p>0.05) (**G**) Averaged PV correlation between Track A and Track B. APP-KI mice showed significant increase (that is, weaker remapping) compared to that of WT mice (Wilcoxon-rank-sum, p<0.001).

Fig. S10. Disrupted remapping of CA1 neurons was observed regardless of animals' running directions

(**A-D**) Cumulative distribution plot of PV Correlation between Track A and Track B from CA1 neurons during remapping task. Four possible combinations of directions have been analyzed. (**A**) Track A right vs Track B right. (**B**) Track A right vs Track B left. (**C**) Track A left vs Track B right. (**D**) Track A left vs Track B left. All combinations showed significant increase in PV correlation from CA1 neurons of APP-KI mice compared to that of WT

mice indicating significant impairment in remapping regardless of directions (Wilcoxon rank sum test, p <0.001 for **A-D**).

Fig. S11. Spatial correlation of CA1 neurons in the linear tracks

(**A**) Cumulative distribution plot of spatial correlation between Track A and Track B (spatial correlation AB) from CA1 neurons during remapping task. Significant increase in spatial correlation AB from CA1 neurons of APP-KI mice compared to that of WT mice indicates disrupted remapping (Wilcoxon rank sum test, p <0.001). (**B**) Cumulative distribution plot of spatial correlation between two recordings in Track A (Wilcoxon rank sum test, p >0.05). (**C-F**) Cumulative distribution plot of Spatial Correlation AB (different tracks). Four possible combinations of directions have been analyzed. (**C**) Track A right vs Track B right. (**D**) Track A right vs Track B left. (**E**) Track A left vs Track B right. (**F**) Track A left vs Track B left. All combinations showed significant increase in spatial correlation AB from CA1 neurons of APP-KI mice compared to that of WT mice, indicating significant impairment in remapping regardless of directions. (KS test, p <0.05 or better for **C-F**)

Fig. S12. Histological validation of recording position in MEC of WT and APP-KI mice Brightfield images of cresyl violet stained sections of all animals recorded. Arrowhead points to the tip of tetrode where the cells were recorded with electrolytic lesions made before sacrificing the animals. Dashed line indicates the border between the MEC and the postrhinal cortex.

Fig. S13. Recording positions in the MEC did not differ between WT and APP-KI mice

Recorded positions along dorsoventral axis of MEC. Distances were calculated from entorhinal border. The mean recording position did not differ between WT and APP-KI mice. (Wilcoxon ranks sum test, p>0.05).

Fig. S14. Additional properties of MEC neurons during the open Field task

(**A**) Two representative MEC neurons recorded from WT mouse and APP-KI mouse during open field task. Spatial information (SI) and average firing rate (Hz) for each cell is shown above rate map. Gridness score (GS) for each cell is shown above autocorrelation map. (**B-D**) Properties of MEC neurons in open field task: (**B**) Number of fields (Wilcoxon-rank-sum, p>0.05). (**C**) Spike width (Wilcoxon-rank-sum, p>0.05). (**D**) Peak firing rate (Wilcoxon-rank-sum, p>0.05). (**E**) Top, distribution of gridness scores calculated from firing rate distributions of MEC neurons in the open field task (real data). Bottom, distribution of shuffled data based on 100 permutations per cell (shuffled data). Red dashed line indicates 95th percentile value (chance level) for a distribution based on

all permutations. Percentage of cells that exceeded chance level is shown on top. 20% of MEC neurons in WT mice and 2% of MEC neurons in APP-KI mice passed this chance level, and they were considered as grid cells.

Fig. S15. Properties of MEC neurons defined as spatially tuned cell

(**A**) We first analyzed cumulative distribution plot of spatial information score from all recorded MEC neurons from WT mice and APP-KI mice. (Wilcoxon rank sum test, p <0.001) (**B**) Top, distribution of spatial information scores in all recorded MEC neurons in the open field task (real data). Bottom, distribution of shuffled data based on 100 permutations per cell (shuffled data). Red dashed line indicates 95th percentile value (chance level) for a distribution based on all permutations. Percentage of cells that exceeded chance level is shown on top. 56% of MEC neurons in WT mice and 20% of MEC neurons in APP-KI mice passed this chance level, and they were considered as "spatially tuned cells".

(**C-H**) Properties of MEC spatially tuned cells in the open field task. (**C**) Field area (cm^2) (Wilcoxon-rank-sum, p>0.05). (**D**) Number of fields (Wilcoxon-rank-sum, p>0.05). (**E**) Average firing rate (Wilcoxon-rank-sum, p>0.05). (**F**) Peak firing rate (Wilcoxon-rank-sum, p>0.05). (**G**) Spatial information score (Wilcoxon-rank-sum, p>0.05). (**H**) Gridness score (Wilcoxon-rank-sum, p>0.05).

(**I-P**) Properties of MEC spatially tuned cells in the linear tracks. (**I**) Field area (cm^2) (Wilcoxon-rank-sum, p>0.05). (**J**) Number of fields (Wilcoxon-rank-sum, p>0.05). (**K**) Average firing rate (Wilcoxon-rank-sum, p>0.05). (**L**) Peak firing rate (Wilcoxon-rank-sum, p>0.05). (**M**) Cumulative distribution plot of PV correlation between Track A and Track B (Wilcoxon rank sum test, p >0.05). (**N**) Cumulative distribution plot of PV correlation between two recordings in Track A (Wilcoxon rank sum test, p >0.05). (**O**) Cumulative distribution plot of spatial correlation between Track A and Track B (Wilcoxon rank sum test, p >0.05). (**P**) Cumulative distribution plot of between two recordings in Track A (Wilcoxon rank sum test, p>0.05).

Fig. S16. Additional properties of MEC neurons during the remapping task.

(**A**) Two representative MEC neurons recorded from WT and APP-KI animal during the remapping task. Spatial correlation between Track A and Track B (SC$_{AB}$) and average firing rate (Hz) for each cell is shown above for each cell. (**B**) Field area (cm^2) (Wilcoxon-rank-sum, p>0.05). (**C**) Number of fields (Wilcoxon rank sum test, p>0.05). (**D**) Average firing rate (Wilcoxon-rank-sum, p>0.05). (**E**) Peak firing rate (Wilcoxon rank sum test, p>0.05). (**F**) Mean PV correlation of MEC neurons between two recordings in Track A (Wilcoxon rank sum test, p>0.05) (**G**) Mean PV correlation of MEC neurons between Track A and Track B. Significant increase in PV correlation of MEC neurons of APP-KI

mice compared to that of WT mice indicates impairment in remapping between different environments. (Wilcoxon rank sum test, p<0.001)

Fig. S17. Disrupted remapping of MEC neurons was observed regardless of animals' running directions

(**A-D**) Cumulative distribution plot of PV Correlation between Track A and Track B from MEC neurons during remapping task. Four possible combinations of directions have been analyzed. (**A**) Track A right vs Track B right. (**B**) Track A right vs Track B left. (**C**) Track A left vs Track B right. (**D**) Track A left vs Track B left. All combinations showed significant increase in PV correlation from MEC neurons of APP-KI mice compared to that of WT mice, indicating the impairment in remapping regardless of directions (Wilcoxon rank sum test, p <0.001 for **A-D**).

Fig. S18. Spatial correlation of MEC neurons during the remapping task.

(**A**) Cumulative distribution plot of spatial correlation between Track A and Track B (spatial correlation AB) from MEC neurons during remapping task. Significant increase in spatial correlation AB from MEC neurons of APP-KI mice compared to that of WT mice indicates disrupted remapping (Wilcoxon rank sum test, p <0.001). (**B**) Cumulative distribution plot of spatial correlation between two recordings in Track A (Wilcoxon rank sum test, p >0.05). (**C-F**) Cumulative distribution plot of spatial correlation AB (different tracks). Four possible combinations of directions have been analyzed. (**C**) Track A right vs Track B right. (**D**) Track A right vs Track B left. (**E**) Track A left vs Track B right. (**F**) Track A left vs Track B left. All combinations showed significant increase in Spatial Correlation AB from MEC neurons of APP-KI mice compared to that of WT mice, indicating significant impairment in remapping regardless of directions. (Wilcoxon rank sum test, p <0.001 for **C-F**).

Fig. S19. Slow gamma oscillations in MEC

(**A**) Occurrence rates of slow gamma episodes in MEC for WT (N=13 mice) and APP-KI mice (N=13 mice) (Wilcoxon rank sum test, p >0.05). (**B**) Mean vector length for the distribution of slow gamma episodes across theta phase (Wilcoxon rank sum test, p>0.05). (**C**) Mean vector length for the distribution of spikes from WT MEC neurons (N=64 cells) and APP-KI MEC neurons (N=62 cells) across slow gamma phase (Wilcoxon ranks sum test, p>0.05). (**D**) MEC:CA1 ratio of slow gamma powers during slow gamma episodes detected in CA1 (Wilcoxon rank sum test, p >0.05).

Fig. S20. Percentage of neurons showing spike phase locking to slow and fast gammas

(**A**) CA1 neurons spikes phase locking to fast gamma oscillations (Binomial test, p<0.001) (**B**) CA1 neurons spikes phase locking to slow gamma oscillations (Binomial test, p<0.05) (**C**) MEC neurons spikes phase locking to fast gamma oscillations (Binomial test, p<0.001) (**D**) MEC neurons spikes phase locking to slow gamma oscillations (Binomial test, p>0.05)

Table S1. Distribution of recorded principal cells across all animals

The total number of isolated CA1, layer II and layer III MEC cells recorded from each animal.

Table S2. Statistics used for properties of CA1 neurons from open field

Summary of statistics applied on analysis from CA1 neurons recorded during open field task. (WT CA1 neurons=105 and APP-KI CA1 neurons=113)

Table S3. Statistics used for properties of CA1 neurons from Linear Track

Summary of statistics applied on analysis from CA1 neurons recorded during open field task. (WT CA1 neurons=93 and APP-KI CA1 neurons=83)

Table S4. Statistics used for properties of place cells from open field and Linear Track

Summary of statistics applied on analysis from place cells recorded during open field task (WT place cells=52 and APP-KI place cells=35) and Linear Track (WT place cells=39 and APP-KI place cells=26)

Table S5. Statistics used for properties of MEC neurons from open field

Summary of statistics applied on analysis from MEC neurons recorded during open field task. (WT MEC neurons=61 and APP-KI MEC neurons=65)

Table S6. Statistics used for properties of MEC neurons from Linear Track

Summary of statistics applied on analysis from MEC neurons recorded during open field task. (MEC CA1 neurons=62 and APP-KI MEC neurons=69)

Table S7. Statistics used for properties of spatially modulated MEC neurons from open field and Linear Track

Summary of statistics applied on analysis from spatially modulated MEC neurons recorded during open field task (WT place cells=34 and APP-KI place cells=13) and Linear Track (WT place cells=30 and APP-KI place cells=12)

Table S8. Statistics used for LFP analysis on CA1

Summary of statistics applied on LFP analysis from CA1 recording during open field task

Table S9. Statistics used for LFP analysis on MEC

Summary of statistics applied on LFP analysis from MEC recording during open field task

Table S10. Statistics used for LFP analysis on CA1-MEC

Summary of statistics applied on LFP analysis from simultaneous recording of CA1 and MEC regions during open field task

Table S11. Statistics used for spatial tuning vs Spatial Correlation AB

Summary of statistics applied on Spatial tuning vs Spatial Correlation AB analysis

Table S12. Statistics used for spatial tuning and Spatial Correlation AB across age

Summary of statistics applied on Spatial tuning and Spatial Correlation AB across age.

Table S13. Shuffled values

Summary of 95th percentile shuffled value obtained for spatial information and gridness score.

Table S14. Statistics used for animal Behavior and Recording Positions

Summary of statistics applied on speed analysis and recording positions from entorhinal border.

Supplementary Reference

Bartho, P., Hirase, H., Monconduit, L., Zugaro, M., Harris, K.D., and Buzsaki, G. (2004). Characterization of neocortical principal cells and interneurons by network interactions and extracellular features. J Neurophysiol *92*, 600-608.

Henriksen, E.J., Colgin, L.L., Barnes, C.A., Witter, M.P., Moser, M.B., and Moser, E.I. (2010). Spatial representation along the proximodistal axis of CA1. Neuron *68*, 127-137.

Liang, L., Oline, S.N., Kirk, J.C., Schmitt, L.I., Komorowski, R.W., Remondes, M., and Halassa, M.M. (2017). Scalable, Lightweight, Integrated and Quick-to-Assemble (SLIQ) Hyperdrives for Functional Circuit Dissection. Front Neural Circuits *11*, 8.

Miao, C., Cao, Q., Ito, H.T., Yamahachi, H., Witter, M.P., Moser, M.B., and Moser, E.I. (2015). Hippocampal Remapping after Partial Inactivation of the Medial Entorhinal Cortex. Neuron *88*, 590-603.

Fig. S1

Schematics for parallel disruptions of entorhinal grid cells and hippocampal place cells

WT

A

APP-KI

B

Fig. S2

Electrophysiological recording of principal neurons in CA1 and MEC

Fig. S3

Histological validation of recording position in CA1 of **WT mice**

Fig. S3 (continued)

Histological validation of recording position in CA1 of APP-KI mice

Mouse00144 Mouse00178 Mouse00180 Mouse00217

Mouse00333 Mouse00335 Mouse00373 Mouse00374

Mouse00384 Mouse00385

Table S1

Distribution of recorded principal cells across all animals

Animal ID	Genotype	Drive Used	Recorded Age	Recorded Area	Number of Principal Cells Recorded (OPF)	Number of Principal Cells Recorded (Linear Track)
mouse00113	APP-KI	MEC only	252	MEC Layer II/III	1	1
mouse00115	APP-KI	MEC only	237	MEC Layer II/III	5	5
mouse00144	APP-KI	HPC only	212	CA1	12	N/A
mouse00178	APP-KI	Split drive	353	CA1 MEC Layer II/III	11 4	10 3
mouse00179	APP-KI	Split drive	236	MEC Layer II/III	3	3
mouse00180	APP-KI	Split Drive	236	CA1 MEC Layer II/III	7 4	7 4
mouse00217	APP-KI	Split Drive	377	CA1 MEC Layer II/III	26 5	24 5
mouse00230	APP-KI	Split Drive	365	MEC Layer II/III	3	3
mouse00271	APP-KI	MEC only	368	MEC Layer II/III	6	4
mouse00282	APP-KI	MEC only	368	MEC Layer II/III	6	8
mouse00283	APP-KI	MEC only	371	MEC Layer II/III	3	2
mouse00333	APP-KI	HPC only	380	CA1	6	4
mouse00335	APP-KI	HPC only	379	CA1	13	7
mouse00372	APP-KI	Split drive	358	MEC Layer II/III	1	5
mouse00373	APP-KI	Split drive	359	CA1	10	9
mouse00374	APP-KI	Split drive	357	CA1 MEC Layer II/III	9 5	7 7
mouse00384	APP-KI	Split drive	371	CA1 MEC Layer II/III	6 7	6 6
mouse00385	APP-KI	Split drive	371	CA1 MEC Layer II/III	13 13	9 13

Table S1 (continued)

Distribution of recorded principal cells aross all animals

Animal ID	Genotype	Drive Used	Recorded Age	Recorded Area	Number of Principal Cells Recorded (OPF)	Number of Principal Cells Recorded (Linear Track)
mouse00090	WT	Split Drive	362	MEC Layer II/III	5	2
mouse00106	WT	HPC only	286	CA1	6	N/A
mouse00177	WT	Split Drive	327	CA1 MEC Layer II/III	8 4	6 7
mouse00227	WT	Split Drive	259	CA1 MEC Layer II/III	15 3	13 4
mouse00228	WT	Split Drive	259	CA1 MEC Layer II/III	22 2	21 2
mouse00229	WT	Split Drive	390	CA1	11	14
mouse00280	WT	MEC only	371	MEC Layer II/III	3	3
mouse00281	WT	MEC only	369	MEC Layer II/III	5	5
mouse00295	WT	MEC only	370	MEC Layer II/III	7	7
mouse00296	WT	MEC only	353	MEC Layer II/III	14	14
mouse00331	WT	HPC only	374	CA1	11	9
mouse00332	WT	HPC only	375	CA1	18	19
mouse00369	WT	Split Drive	339	CA1 MEC Layer II/III	3 4	1 4
mouse00370	WT	Split Drive	356	CA1 MEC Layer II/III	5 5	5 6
mouse00371	WT	Split Drive	327	MEC Layer II/III	1	1
mouse00382	WT	Split Drive	322	MEC Layer II/III	6	6
mouse00383	WT	Split Drive	320	CA1 MEC Layer II/III	6 2	5 2

Fig. S4

Spatial Tuning and Spatial Correlation AB
across age (7 - 13 months old)

Fig. S5

Running behaviors did not differ between WT and APP-KI mice

Fig. S6

Electrophysiological features of classifying principal neurons in CA1 and MEC

Fig. S7

Additional properties of CA1 neurons- Open Field

A

WT CA1

SI: 1.3   1.46Hz

APP-KI CA1

SI: 0.5   1.05Hz

Max

B Field Area (cm²)   N.S.

C # of Fields   N.S.

D Spike Width (ms)   N.S.

E Average Firing Rate (Hz)   N.S.

F Peak Firing Rate (Hz)   N.S.

Fig. S8

Properties of CA1 neurons defined as place cells in WT and APP-KI mice

Open Field

Linear Track

Fig. S9

Additional properties of CA1 neurons - Linear Tracks

Disrupted remapping of CA1 neurons was observed regardless of animals' running direction

Fig. S11

Spatial Correlation of CA1 neurons in the Linear Tracks

Fig. S12

Histological validation of recording position in MECII/III of **WT mice**

Histological validation of recording position
in MECII/III of APP-KI mice

Mouse00113   Mouse00115   Mouse00178   Mouse00179

Mouse00180   Mouse00217   Mouse00230   Mouse00271

Mouse00282   Mouse00283   Mouse00372   Mouse00374

Mouse00384   Mouse00385

Fig. S13

Recording positions in the MEC did not differ between WT and APP-KI mice

Fig. S14

Additional properties of MEC neurons - Open Field

A

WT MEC

SI: 1.1   0.58Hz   GS: 0.83

APP-KI MEC

SI: 0.042   3.2Hz   GS: -0.12

Max

B   # of Fields

N.S.

61   65

WT   APP-KI

C   Spike Width (ms)

N.S.

61   65

WT   APP-KI

D   Peak Firing Rate (Hz)

N.S.

61   65

WT   APP-KI

E   Gridness Score (Obseved frequency)

Observed frequency

WT (20%)

APP-KI (2%)

Gridness Score (Shuffled Frequency)

1000

Shuffled frequency

95th percentile value: 0.41

-0.5   0   0.5   1

Gridness Score

Properties of MEC neurons defined as spatially tuned cell

Open Field

Linear Track

Additional properties of MEC neurons - Linear Tracks

Fig. S17

Disrupted remapping of MEC neurons was observed regardless of animals' running directions

A   Black Right vs White Right

B   Black Right vs White Left

C   Black Left vs White Right

D   Black Left vs White Left

Fig. S18

Spatial Correlation of MEC neurons in the Linear Tracks

Fig. S19

Slow gamma oscillations in MEC

A

B · Theta-Gamma Coupling

C · Spike-Gamma Coupling

D

Fig. S20

Percentage of neurons showing spike phase locking to slow and fast gammas

Table S2
Basic properties of CA1 neurons from OPF

WT CA1 neurons (n=105) vs APP-KI CA1 neurons (n=113)

Experimental measure	Figure	Test Used	P-Value	degree of freedom & F/T/z/R/ETC value
Spatial Information Index (cumulative)	Fig. 1F	Two sample Kolmogorov–Smirnov test	p=1.4e-04	
Spatial Information Index (bar)	Fig. 1G	Wilcoxon Ranksum	p=7.5e-05	Z=-0.57, WT SEM=0.053 APP-KI SEM= 0.041
Percentage of Place Cells	Fig. 1G	Binomial test	p=6.3e-5	
Field Area (cm2)	SFig. 7B	Wilcoxon Ranksum	p=0.54	Z=-0.57, WT SEM=73, APP-KI SEM=93
# of Fields	SFig. 7C	Wilcoxon Ranksum	p=0.69	Z=-0.57, WT SEM=0.12, APP-KI SEM=0.13
Spike Width	SFig. 7D	Wilcoxon Ranksum	p=0.093	Z=-0.57, WT SEM=9.3, APP-KI SEM=8.5
Average Firing Rate	SFig. 7E	Wilcoxon Ranksum	p=0.60	Z=-0.57, WT SEM=0.087, APP-KI SEM=0.081
Peak Firing Rate	SFig. 7F	Wilcoxon Ranksum	p=0.27	Z=-0.57, WT SEM=0.26, APP-KI SEM=0.28

Table S3
Basic properties of CA1 neurons from Linear Track

WT CA1 neurons (n=93) vs APP-KI CA1 neurons (n=83)

Experimental measure	Figure	Test Used	P-Value	degree of freedom & F/T/z/R/ETC value
Population Vector Correlation AB	Fig. 1L	Wilcoxon Ranksum	p=4.1e-29	Z=-11
Population Vector Correlation AA	Fig. 1L	Wilcoxon Ranksum	p=0.42	Z=0.81
Field Area	SFig. 9B	Wilcoxon Ranksum	p=0.10	Z=-0.79, WT SEM=2.0, APP-KI SEM=2.5
# of Fields	SFig. 9C	Wilcoxon Ranksum	p=0.070	Z=-0.79, WT SEM=0.032, APP-KI SEM=0.039
Average Firing Rate	SFig. 9D	Wilcoxon Ranksum	p=0.52	Z=-0.79, WT SEM=0.13, APP-KI SEM=0.17
Peak Firing Rate	SFig. 9E	Wilcoxon Ranksum	p=0.43	Z=-0.79, WT SEM=0.37, APP-KI SEM=0.57
Population Vector Correlation AA	SFig. 9F	Wilcoxon Ranksum	p=0.42	Z=0.81
Population Vector Correlation AB	SFig. 9G	Wilcoxon Ranksum	p=4.1e-20	Z=-11
Population Vector Correlation ArBr	SFig. 10A	Wilcoxon Ranksum	p=1.4e-24	Z=-10
Population Vector Correlation ArBl	SFig. 10B	Wilcoxon Ranksum	p=1.4e-25	Z=-11
Population Vector Correlation AlBr	SFig. 10C	Wilcoxon Ranksum	p=2.2e-21	Z=-9.5
Population Vector Correlation AlBl	SFig. 10D	Wilcoxon Ranksum	p=1.3e-17	Z=-8.5
Spatial Correlation AB	SFig. 11A	Wilcoxon Ranksum	p=0.0013	Z=-3.2
Spatial Correlation AA	SFig. 11B	Wilcoxon Ranksum	p=0.63	Z=0.49
Spatial Correlation ArBr	SFig. 11C	Wilcoxon Ranksum	p=0.0028	Z=-3.0
Spatial Correlation ArBl	SFig. 11D	Wilcoxon Ranksum	p=0.032	Z=-2.1
Spatial Correlation AlBr	SFig. 11E	Wilcoxon Ranksum	p=0.0081	Z=-2.6
Spatial Correlation AlBl	SFig. 11F	Wilcoxon Ranksum	p=0.0028	Z=-3.0

Table S4
Basic properties of place cells from OPF and Linear Track

OPF: WT Place Cells (n=52) vs APP-KI Place Cells (n=35)
Linear Track: WT Place Cells (n=39) vs APP-KI Place Cells (n=26)

Experimental measure	Figure	Test Used	P-Value	degree of freedom & F/T/z/R/ETC value
OPF Field Area	SFig. 8B	Wilcoxon Ranksum	p=0.54	Z=1.9, WT SEM=62, APP-KI SEM=71
OPF # of Fields	SFig.8C	Wilcoxon Ranksum	p=0.46	Z=1.9, WT SEM=0.14, APP-KI SEM=0.12
OPF Average Firing Rate	SFig. 8D	Wilcoxon Ranksum	p=0.79	Z=1.9, WT SEM=0.078, APP-KI SEM=0.084
OPF Peak Firing Rate	SFig. 8E	Wilcoxon Ranksum	p=0.94	Z=1.9, WT SEM=0.37, APP-KI SEM=0.47
OPF Spatial Information Index	SFig. 8F	Wilcoxon Ranksum	p=0.19	Z=1.9, WT SEM=0.072, APP-KI SEM=0.074
Linear Track Field Area	SFig. 8G	Wilcoxon Ranksum	p=0.63	Z=1.6, WT SEM=85, APP-KI SEM=96
Linear Track # of Fields	SFig. 8H	Wilcoxon Ranksum	p=0.52	Z=1.6, WT SEM=0.18, APP-KI SEM=0.16
Linear Track Average Firing Rate	SFig. 8I	Wilcoxon Ranksum	p=0.78	Z=1.6, WT SEM=0.094, APP-KI SEM=0.11
Linear Track Peak Friing Rate	SFig. 8J	Wilcoxon Ranksum	p=0.88	Z=1.6, WT SEM=0.44, APP-KI SEM=0.53
Linear Track Population Vector Correlation AB	SFig. 8K	Wilcoxon Ranksum	p=3.7e-32	Z=-12
Linear Track Population Vector Correlation AA	SFig. 8L	Wilcoxon Ranksum	p=0.0020	Z=-3.1
Linear Track Spatial Correlation AB	SFig. 8M	Wilcoxon Ranksum	p=0.0044	Z=2.9
Linear Track Spatial Correlation AA	SFig. 8N	Wilcoxon Ranksum	p=0.0032	Z=3.0

Table S5
Basic properties of MEC neurons from OPF

WT MEC neurons (n=61) vs APP-KI MEC neurons (n=65)

Experimental measure	Figure	Test Used	P-Value	degree of freedom & F/T/z/R/ETC value
Gridness Score (cumulative)	Fig. 3F	Two sample Kolmogorov–Smirnov test	p=0.032	
Percentage of grid cells	Fig. 3G	Binomial test	p=1.5e-10	
Spatial Information Index	Fig. 3G	Wilcoxon Ranksum	p=2.2e-07	Z=-0.37, WT SEM=0.043, APP-KI SEM=0.037
Field Area	Fig. 3G	Wilcoxon Ranksum	p=0.0047	Z=-0.37, WT SEM=140, APP-KI SEM=330
Average Firing Rate	Fig. 3G	Wilcoxon Ranksum	p=0.29	Z=-0.37, WT SEM=0.11, APP-KI SEM=0.14
# of Fields	SFig. 14B	Wilcoxon Ranksum	p=0.24	Z=-0.37, WT SEM=0.22, APP-KI SEM=0.23
Spike Width	SFig. 14C	Wilcoxon Ranksum	p=0.62	Z=-0.37, WT SEM=11, APP-KI SEM=12
Peak Firing Rate	SFig. 14D	Wilcoxon Ranksum	p=0.74	Z=-0.37, WT SEM=1.1, APP-KI SEM=2.0

Table S6
Basic properties of MEC neurons from Linear Track

MEC neurons (n=62) vs APP-KI CA1 neurons (n=69)

Experimental measure	Figure	Test Used	P-Value	degree of freedom & F/T/z/R/ETC value
Population Vector Correlation AB	Fig. 3J	Wilcoxon Ranksum	p=5.5e-27	Z=-11
Population Vector Correlation AA	Fig. 3J	Wilcoxon Ranksum	p=0.41	Z=0.83
Field Area	SFig. 16B	Wilcoxon Ranksum	p=1.7e-5	Z=-0.63, WT SEM=3.6, APP-KI SEM=4.0
# of Fields	SFig. 16C	Wilcoxon Ranksum	p=2.9e-5	Z=-0.63, WT SEM=3.6, APP-KI SEM=4.0
Average Firing Rate	SFig. 16D	Wilcoxon Ranksum	p=0.48	Z=-0.63, WT SEM=0.20, APP-KI SEM=0.16
Peak Firing Rate	SFig. 16E	Wilcoxon Ranksum	p=0.47	Z=-0.63, WT SEM=0.40, APP-KI SEM=0.37
Population Vector Correlation AA	SFig. 16F	Wilcoxon Ranksum	p=0.41	Z=0.82
Population Vector Correlation AB	SFig. 16G	Wilcoxon Ranksum	p=5.5e-27	Z=-11
Population Vector Correlation ArBr	SFig. 17A	Wilcoxon Ranksum	p=3.5e-14	Z=-7.6
Population Vector Correlation ArBl	SFig. 17B	Wilcoxon Ranksum	p=2.8e-15	Z=-7.9
Population Vector Correlation AlBr	SFig. 17C	Wilcoxon Ranksum	p=5.7e-19	Z=-8.9
Population Vector Correlation AlBl	SFig. 17D	Wilcoxon Ranksum	p=2.8e-9	Z=-5.9
Spatial Correlation AB	SFig. 18A	Wilcoxon Ranksum	p=6.5e-7	Z=-5.0
Spatial Correlation AA	SFig. 18B	Wilcoxon Ranksum	p=0.31	Z=1.0
Spatial Correlation ArBr	SFig. 18C	Wilcoxon Ranksum	p=5.4e-7	Z=-5.0
Spatial Correlation ArBl	SFig. 18D	Wilcoxon Ranksum	p=4.1e-5	Z=-4.1
Spatial Correlation AlBr	SFig. 18E	Wilcoxon Ranksum	p=5.2e-7	Z=-5.0
Spatial Correlation AlBl	SFig. 18F	Wilcoxon Ranksum	p=5.4e-7	Z=-5.0

Table S7
Basic properties of Spatially modulated MEC neurons form OPF and Linear Track

OPF: WT Spatially Modulated Cells (n=34) vs APP-KI Spatially Modulated Cells (n=13)
Linear Track: WT Spatially Modulated Cells (n=30) vs APP-KI Spatially Modulated Cells (n=12)

Experimental measure	Figure	Test Used	P-Value	degree of freedom & F/T/z/R/ETC value
Spatial Information Index (cumulative)	SFig. 15A	Two sample Kolmogorov–Smirnov test	p=2.2e-6	
OPF Field Area	SFig. 15C	Wilcoxon Ranksum	p=0.23	Z=3.1, WT SEM=75, APP-KI SEM=160
OPF # of Fields	SFig. 15D	Wilcoxon Ranksum	p=0.84	Z=3.1, WT SEM=0.23, APP-KI SEM=0.48
OPF Average Firing Rate	SFig. 15E	Wilcoxon Ranksum	p=0.11	Z=3.1, WT SEM=0.10, APP-KI SEM=0.12
OPF Peak Firing Rate	SFig. 15F	Wilcoxon Ranksum	p=0.32	Z=3.1, WT SEM=1.9, APP-KI SEM=0.71
OPF Spatial Information Index	SFig. 15G	Two sample Kolmogorov-Smirnov test	p=0.55	
OPF Gridness Score	SFig. 15H	Two sample Kolmogorov-Smirnov test	p=0.066	
Linear Track Field Area	SFig. 15I	Wilcoxon Ranksum	p=0.36	Z=2.8, WT SEM=90, APP-KI SEM=110
Linear Track # of Fields	SFig. 15J	Wilcoxon Ranksum	p=0.13	Z=2.8, WT SEM=0.24, APP-KI SEM=0.39
Linear Track Average Firing Rate	SFig. 15K	Wilcoxon Ranksum	p=0.24	Z=2.8, WT SEM=0.10, APP-KI SEM=0.27
Linear Track Peak Friing Rate	SFig. 15L	Wilcoxon Ranksum	p=0.75	Z=2.8, WT SEM=2.1, APP-KI SEM=0.79
Linear Track Population Vector Correlation AB	SFig. 15M	Wilcoxon Ranksum	p=0.91	Z=0.12
Linear Track Population Vector Correlation AA	SFig. 15N	Wilcoxon Ranksum	p=0.13	Z=-1.3
Linear Track Spatial Correlation AB	SFig. 15O	Wilcoxon Ranksum	p=0.56	Z=-0.58
Linear Track Spatial Correlation AA	SFig. 15P	Wilcoxon Ranksum	p=0.75	Z=1.8

Table S8
LFP analysis on CA1

Experimental measure	Figure	Test Used	Sample	P-Value	degree of freedom & F/T/z/R/ETC value
Normalized oscillatory power (Slow and Fast gamma)	Fig. 2C	unpaired t-test with FDR corrected for 100 multiple comparison	WT=10 mice, APP-KI = 10 mice	q=0.023 (average)	
Occurance rate of fast gamma	Fig. 2E	Wilcoxon Ranksum	WT=10 mice, APP-KI = 10 mice	P= 1.8e-4	Z=3.74
Occurance rate of slow gamma	Fig. 2E	Wilcoxon Ranksum	WT=10 mice, APP-KI = 10 mice	P=0.38	Z=0.87
Theta to slow gamma coupling	Fig. 2G	Wilcoxon Ranksum	WT=10 mice, APP-KI = 10 mice	p=0.35	Z=0.95
Theta to fast gamma coupling	Fig. 2G	Wilcoxon Ranksum	WT=10 mice, APP-KI = 10 mice	p=0.045	Z=2.00
Phase locking of spikes to slow gamma	Fig. 2I	Wilcoxon Ranksum	WT=104 neurons, APP-KI=105 neurons	p=0.40	Z=0.84
Phase locking of spikes to fast gamma	Fig. 2I	Wilcoxon Ranksum	WT=104 neurons, APP-KI=105 neurons	p=8.5e-10	Z=6.13
Phase locking of spikes to fast gamma	SFig. 20A	binomial test	WT=(64/104),APP-KI=(9/105)	p=1.5e-12	
Phase locking of spikes to slow gamma	SFig. 20B	binomial test	WT=(48/104),APP-KI=(38/105)	p=0.041	

Table S9
LFP analysis on MEC

Experimental measure	Figure	Test Used	Sample	P-Value	degree of freedom & F/T/z/R/ETC value
Normalized oscillatory power (Slow and Fast gamma)	Fig. 4C	unpaired t-test with FDR corrected for 100 multiple comparison	WT=13 mice, APP-KI = 13 mice	q=0.73 (average)	
Occurance rate of fast gamma	Fig. 4D	Wilcoxon Ranksum	WT=13 mice, APP-KI = 13 mice	p=0.31	Z=1.02
Theta to fast gamma coupling	Fig. 4F	Wilcoxon Ranksum	WT=13 mice, APP-KI = 13 mice	p=0.051	Z=1.95
Phase locking of spikes to fast gamma	Fig. 4H	Wilcoxon Ranksum	WT=60 neurons, APP-KI=62 neurons	p=0.0061	Z=2.74
Occurance rate of slow gamma	SFig. 19A	Wilcoxon Ranksum	WT=13 mice, APP-KI = 13 mice	p=0.38	Z=0.87
Theta to slow gamma couplnig	SFig. 19B	Wilcoxon Ranksum	WT=13 mice, APP-KI = 13 mice	p=0.15	Z=1.4
Phase locking of spikes to slow gamma	SFig. 19C	Wilcoxon Ranksum	WT=60 neurons, APP-KI=62 neurons	p=0.58	Z=0.55
Phase locking of spikes to fast gamma	SFig. 20C	binomial test	WT=(51/60), APP-KI=(35/62)	p=5.1e-6	
Phase locking of spikes to slow gamma	SFig. 20D	binomial test	WT=(29/60), APP-KI=(31/62)	p=0.90	

Table S10
LFP analysis on CA1-MEC

Experimental measure	Figure	Test Used	Sample	P-Value	degree of freedom & F/T/z/R/ETC value
MEC-CA1 Coherence	Fig. 4J	unpaired t-test with FDR corrected for 100 multiple comparison	WT=6 mice, APP-KI = 6 mice	q=0.0089 (average)	
MEC/CA1 fast gamma power ratio	Fig. 4L	Wilcoxon Ranksum	WT=6 mice, APP-KI = 6 mice	p=0.039	Z=2.08
MEC/CA1 slow gamma power ratio	SFig. 19D	Wilcoxon Ranksum	WT=6 mice, APP-KI=6 mice	p=0.24	Z=1.28

Table S11
Spatial tuning vs Spatial Correlation AB

Experimental measure	Figure	Test Used	Sample #	P-Value	degree of freedom & F/T/z/R/ETC value
Spatial Tuning vs Spatial Correlation (WT CA1)	Fig. 3K	Pearson Correlation	WT=105 neurons	p=0.14	r(105)=-0.15
Spatial Tuning vs Spatial Correlation (WT MEC)	Fig. 3K	Pearson Correlation	WT=61 neurons	p=0.09	r(61)=-0.22
Spatial Tuning vs Spatial Correlation (APP-KI CA1)	Fig. 3K	Pearson Correlation	APP-KI=113 neurons	p=0.22	r(113)=-0.14
Spatial Tuning vs Spatial Correlation (APP-KI MEC)	Fig. 3K	Pearson Correlation	APP-KI=65 neurons	p=1e-04	r(66)=-0.45

Table S12
Spatial tuning and Spatial Correlation AB across age

Experimental measure	Figure	Test Used	Sample #	P-Value	degree of freedom & F/T/z/R/ETC value
Spatial Information across age (CA1)	SFig. 4A	Pearson Correlation	WT=105, APP-KI=113	p(WT)=0.13, p(APP-KI)=0.58	r(105,WT)= -0.15 , r(116,APP-KI)=0.053
Spatial Information across age (MEC)	SFig. 4B	Pearson Correlation	WT=61,APP-KI=65	p(WT)=0.25, p(APP-KI)=0.26	r(61,WT)= 0.15 , r(66,APP-KI)=0.14
Spatial Correlation AB across age (CA1)	SFig. 4C	Pearson Correlation	WT=105,APP-KI=113	p(WT)=0.21, p(APP-KI)=0.21	r(105,WT)= 0.13 , r(113,APP-KI)=0.14
Spatial Correlation AB across age (MEC)	SFig. 4D	Pearson Correlation	WT=61,APP-KI=65	p(WT)=0.017, p(APP-KI)=0.39	r(61,WT)= -0.30 , r(65,APP-KI)=0.11

Table S13
Shuffled Values

Experimental measure	Figure	Test Used	Sample #	degree of freedom & F/T/z/R/ETC value
Spatial Information Shuffle (WT CA1)	SFig. 8A	Shuffling (time shift)	105 neurons	95th percentile value= 0.63
Spatial Information Shuffle (APP-KI CA1)	N/A	Shuffling (time shift)	113 neurons	95th percentile value= 0.53
Spatial Information Shuffle (CA1)	N/A	Shuffling (time shift)	218 neurons	95th percentile value= 0.57
Spatial Information Shuffle (WT MEC)	SFig. 15B	Shuffling (time shift)	61 neurons	95th percentile value= 0.43
Spatial Information Shuffle (APP-KI MEC)	N/A	Shuffling (time shift)	65 neurons	95th percentile value= 0.37
Spatial Information Shuffle (MEC)	N/A	Shuffling (time shift)	126 neurons	95th percentile value= 0.41
Gridness Score Shuffle (WT MEC)	SFig. 14E	Shuffling (time shift)	61 neurons	95th percentile value= 0.41
Gridness Score Shuffle (APP-KI MEC)	N/A	Shuffling (time shift)	65 neurons	95th percentile value= 0.40
Gridness Score Shuffle (MEC)	N/A	Shuffling (time shift)	126 neurons	95th percentile value= 0.41

Table S14
Animal Behavior and Recording Positions

Experimental measure	Figure	Test Used	Sample #	P-Value	degree of freedom & F/T/z/R/ETC value
Speed across OPF	SFig. 5A	unpaired t-test with FDR corrected for 25 multiple comparison	WT=17 mice, APP-KI = 18 mice	q=0.84 (average)	
Speed across Linear Track	SFig. 5B	unpaired t-test with FDR corrected for 25 multiple comparison	WT=17 mice, APP-KI = 18 mice	q=0.77 (average)	
Recording positions from entorhinal border	SFig. 13	T-test	WT=61 neurons, APP-KI= 65 neurons	p=0.59	t(125)=0.55