Broad-Spectrum Antifungal Activities and Mechanism of Drimane

Sesquiterpenoids

Edruce Edouarzin[1], Connor Horn[2], Anuja Paduyal[2], Cunli Zhang[1], Jianyu Lu[1], Zongbo Tong[1],

Guri Giaever[3], Corey Nislow[3], Raja Veerapandian[2], Duy H. Hua[1] and Govindsamy

Vediyappan[2*]

[1]Department of Chemistry, 1212 Mid Campus Drive North, Kansas State University, Manhattan,

KS 66506 U. S. A.

[2]Division of Biology, 1717 Claflin Road, Kansas State University, Manhattan, KS 66506

U. S. A.

[3]Faculty of Pharmaceutical Sciences, University of British Columbia,Vancouver, BC Canada

V6T 1Z3

12
13
14
15
16   **Running title:** Drimenol, a broad-spectrum fungicidal antifungal agent
17
18
19

20   **Corresponding author**
21
22   * gvediyap@ksu.edu
23
24

ABSTRACT

Eight drimane sesquiterpenoids including (-)-drimenol and (+)-albicanol were synthesized from (+)-sclareolide and evaluated for their antifungal activities. Three compounds, (-)-drimenol, (+)-albicanol, and (1*R*,2*R*,4a*S*,8a*S*)-2-hydroxy-2,5,5,8a-tetramethyl-decahydronaphthalene-1-carbaldehyde (**4**) showed strong activity against *C. albicans*. (-)-Drimenol, the strongest inhibitor of the three, (at concentrations of 8 – 64 μg/ml, causing 100% death of fungi), acts not only against *C. albicans* as a fungicidal manner, but also inhibits other fungi such as *Aspergillus*, *Cryptococcus*, *Pneumocystis*, *Blastomyces*, *Fusarium*, *Rhizopus*, *Saksenaea* and FLU resistant strains of *C. albicans*, *C. glabrata, C. krusei, C. parapsilosis* and *C. auris.* These observations suggest drimenol is a broad-spectrum antifungal agent. At high concentration (100 μg/ml), drimenol caused a rupture of the fungal cell wall/membrane. In a nematode model of *C. albicans* infection, drimenol rescued the worms from *C. albicans*-mediated death, indicating drimenol is tolerable and bioactive in a metazoan. Genome-wide fitness profiling assays of both *S. cerevisiae* (nonessential homozygous and essential heterozygous) and *C. albicans* (Tn-insertion mutants) collections revealed putative genes and pathways affected by drimenol. Using a *C. albicans* mutants spot assay, the Crk1 kinase associated gene products, Ret2, Cdc37, and novel putative targets orf19.759, orf19.1672, and orf19.4382 were revealed to be the potential targets of drimenol. Further, computational modeling results suggest possible modification of the structure of drimenol including the A ring for improving antifungal activity.

INTRODUCTION

Life-threatening fungal infections are an important cause of morbidity and mortality, particularly for patients with immune deficiency and those who are undergoing chemotherapeutic treatments. Some of the leading invasive fungal pathogens include *Candida* sp., *Aspergillus* sp. and *Cryptococcus* sp. Currently, the antifungal therapeutic options are limited, especially when compared to available antibacterial agents (1-4). Among the five classes of antifungals, azoles, echiocandins, polyenes, allylamines, and pyrimidine derivatives, only three are used clinically; azoles, echiocandins, and polyenes. Azole drugs, such as fluconazole (FLU), inhibit ergosterol synthesis through inhibition of lanosterol 14-α-demethylase, impairing formation of the fungal cell wall. Echocandins, such as caspofungin (CAS), block 1,3-β-glucan synthase and lead to depletion of glucan in fungal cell wall. Polyenes, including amphotericin B (AMB), bind to ergosterol in fungal cell membrane and change the cell membrane transition temperature, resulting in leakage of ions and small organic molecules, and eventual cell death. Allylamines, such as amorolfin, affect ergosterol synthesis by inhibition of squalene epoxidase. Pyrimidines, such as flucytosine (or 5-fluorocytosine), block nucleic acid synthesis, leading to the inhibition of protein synthesis (5, 6). No new antifungal agents have been approved by Food and Drug Administration since 2006 (2) and the latest antimycotic agents, echinocandins, were developed over 30 years ago (7, 8). Thus, new broad-spectrum antifungal agents are sorely needed to overcome the increasing emergence of antifungal drug resistance.

C. albicans is the most frequently found fungal pathogen in humans and costs the US health care system around $3 billion annually in treatment and lost productivity. *C. albicans*, a polymorphic fungus, exists as yeast, pseudohyphal and hyphal forms, with each contributing to its virulence. While the yeast form is essential for dissemination, the hyphal form is critical for

invasion of cells, immune evasion, and biofilm formation. Furthermore, the ability to switch

between forms is also essential for pathogenicity. Traditional antimycotics have many drawbacks,

including toxicity to human cells, a limited range of cellular targets, the development of antifungal

resistance (9-12), and the failure to successfully control pathogenesis. To develop new antifungal

agents based on drimane sesquiterpenes, we have investigated synthetic drimane terpenes, (-)-

drimenol (**1**) and (+)-albicanol (**2**), along with six analogs, **3 – 8** (**Fig. 1**), for their antifungal

activities and identified (-)-**1** as a potent broad-spectrum fungicidal agent. Moreover, we

determined their mechanism of action through forward genetic screening of mutant libraries of *C.*

albicans and baker's yeast and found that (-)-**1** affects the fungal activities of protein secretion,

vacuolar biogenesis, chromation remodeling and cyclin dependent protein kinases (CDK) activity.

RESULTS

Several drimane sesquiterpenoids were synthesized in our laboratory during the total synthesis of

(+)-chloropuupehenone, a natural product from marine sponges (13). Based on the

antimycobacterial activity of sesquiterpene natural product, cinnamodial, isolated from the

liverwort plant, *Warburgia salutaris* (14), we anticipated that drimane sesquiterpenes and closely

related compounds (15) could be effective antimycotics for *C. albicans*. Five representative

drimane terpenes, **1 – 5**, along with their derivatives, **6 – 8** (**Fig. 1**), were screened for their ability

to inhibit *C. albicans* growth. It was assumed that additional hydroxyl group(s) or oxygen atoms

in the molecule enhances water solubility and may improve bioactivity (16). Molecules **3 – 7**

possess extra hydroxyl, aldehyde, or acetoxy function and molecule **8** contains an acetonide moiety

in the drimane structure.

FIG. 1. Drimane sesquiterpenoids bioevaluated for their antifungal activities.

Compounds **1** – **5** were prepared by following previously reported methods (13). Molecule **6** was prepared from a mono-acetylation of **5** with acetic anhydride and pyridine in dichloromethane (Scheme 1). Molecule **7** was made by a sequence of three reactions: (i) silylation of the primary alcohol with *t*-butyldimethylsilyl chloride and imidazole in dichloromethane; (ii) acetylation of the tertiary alcohol with acetyl chloride and pyridine; and (iii) removal of the silyl ether group with tetra-*n*-butylammonium fluoride in THF. Compound **8** was produced from the treatment of triol **3** with 2,2-dimethoxylpropane and a catalytic amount of *p*-toluenesulfonic acid in toluene. The experimental procedures were described in Materials and Methods section.

Scheme 1. Synthesis of drimane sesquiterpenoids **3** – **8**.

Examinataion of the antifungal activities of these molecules along with the mechanistic study of the most active molecule may allow future improvement in bioactivity and reduction in toxicity. We used *C. albicans* strain SC5314 for our initial screening of antifungal activities of drimane sesquiterpenoids **1** – **8**. The compounds were solubilized in dimethyl sulfoxide (DMSO), 10 mg/ml, as stock solutions and stored at -20°C. Prior to assays, stock solutions of compounds were diluted to 200 – 12.5 µg/ml in the growth media for yeast antifungal assays using the Clinical and Laboratory Standards Institute (CLSI) M38-A2 method (17). Fortuitously, we found (-)-drimenol (**1**) and (+)-albicanol (**2**) along with compound **4**, inhibit *C. albicans* SC5314 growth (**Table 1**). Among these three compounds, we identified **1** being more potent (with MIC value

116   ~30 µg/ml) than other compounds (~60 µg/ml); therefore **1** was used for further studies including

117   mechanistic investigation.

119   **Table 1**. Antifungal activities of drimane sesquiterpenoids **1 – 8** against *C. albicans* SC5314.

120   Highest activity: ++++; medium activity: +++; and low activity: ++.

Molecule	**1**	**2**	**3**	**4**	**5**	**6**	**7**	**8**
Antifungal activity	++++	++	Inactive	+++	Inactive	Inactive	Inactive	Inactive

Antifungal activities of drimenol against various pathogenic fungi

Since our initial assay with *C. albicans* confirmed the antifungal activities of **1**, we

extended the susceptibility assays to other pathogenic fungi including FLU resistant *C. albicans*,

various species of candida, *Cryptococcus*, *Aspergillus* and a dermatophyte fungus. The CLSI broth

dilution methods of M27-A3 for yeasts and M38-A2 for filamentous fungi (17) were used to

determine the susceptibility. Molecule **4** was not investigated due to the presence of an aldehyde

function, which may react with biological molecules.

Briefly, yeast cells or conidia (for filamentous fungi) were suspended in RPMI 1640

medium to a final concentration of 10^5 cfu/ml and distributed in 96-well microplate to a total

volume of 100 µl/well. Molecules **1** was added to the wells and two-fold serial dilutions prepared.

Duplicates were used for each concentration and wells with or without DMSO served as controls.

Plates were incubated without shaking at 37°C for 24 – 48 h for yeasts and 30°C for 4 days for

filamentous fungi (*Aspergillus* and *Trichophyton* sp). The MIC was defined as the lowest

compound concentration at which no growth occurred, as determined visually and microscopically

(an inverted microscope). The results are shown in **Fig. 2** and some (*C. glabrata* [BG2], *C.*

albicans [FLU resistant], *C. auris, T. equinum*) in **Table 2**.

FIG. 2. Antifungal activity of **1** against various human pathogenic fungi. The CLSI broth dilution

methods of M27-A3 for yeasts and M38-A2 for filamentous fungi were used. Drimenol (**1**) was

dissolved in DMSO and a two fold serial dilution was used between 200 - 12.5 μg/ml.

Representative microscopic images (Leica, inverted microscope) taken at 200X magnification

from **1** with cells exposed 50 μg/ml (MIC), except *T. equinum* which was exposed to 15 μg/ml at

30 °C, are shown. The MIC for **1** is 50 μg/ml under these assay conditions except mentioned

otherwise. *C. a* (SC), *Candida albicans* (SC5314); *C. a* (FL-R), *C. albicans* FLU-resistant; *C. g*

(BG2), *C. glabrata* (BG2); *C. k.* (MHF2), *C. krusei* (MHF2); *C. n* (H99), *Cryptococcus*

neoformans (H99); *T. e, Trichophyton equinum.*

Results summarized in **Fig. 2** show that **1** has broad-spectrum fungicidal activity against various

fungi including FLU resistant *C. alibcans* and *Cryptococcus* sp albeit, at a higher (50 µg/ml) MIC

concentration. However, for a dermatophyte fungus, the MIC value was lower, 15 µg/ml. When

compared to DMSO controls, fungi exposed to **1** showed absence of fungal growth. At higher

concentration of **1** (100 µg/ml), *C. albicans* yeast cells lysed and released their cellular contents

(**Fig. 3**, arrow). Consistent with this observation, **1** inhibited the germination of *A. nidulans* spores

and appeared to cause swelling of germinating spores (**Fig. 3**, lower right).

FIG. 3. Antifungal activity of **1** against *C. albicans* and *A. nidulans*. The CLSI broth dilution

methods of M27-A3 for yeasts and M38-A2 for filamentous fungi were used. *C. albicans*

(SC5314) showed lysis of yeast cells (arrow) at 100 µg/ml of **1** compared to the control where a

network of hyphal growth was observed**.** Similarly, the germination of *Aspergillus nidulans* spores

was inhibited by **1** (lower, right panel). Representative microscopic images (Leica, inverted

microscope) were shown. Scale bar, 20 μm.

To extend our antifungal screening with **1** against additional human pathogenic fungi, we have

used the non-clinical and pre-clinical service program offered by the NIH NIAID supported fungus

testing center at the University of Texas Health Sciences Center, San Antonio. The fungus testing

center used the CLSI M38-A2 method (17) to determine the MIC of **1** after 24 - 72 hours incubation

with concentrations ranging from 0.125 - 64 μg/ml. Positive control antifungals (FLU,

posaconazole and voriconazole) were also included in parallel. The results are summarized in

Table 2.

Table 2. Drimenol (**1**) activities against various pathogenic fungi. The MIC (100% growth

inhibition) of **1** ranges from 8 μg/ml to 64 μg/ml. Fluconazole, posaconazole, and voriconazole

were used as controls.

Antifungal		Drimenol		Fluconazole	Posaconazole	Voriconazole
Species	Isolate No.	50%	100%	50%	100%	100%
C. parapsilosis	CLSI QC	32	32	1	-	-
C. krusei	CLSI QC	32	>64	16	-	-
P. variotii	CLSI QC	16	32	-	≤0.03	0.125
C. albicans	CA1	32	32	0.125	-	-
	CA2	32	>64	0.25	-	-
	CA3	32	32	>64	-	-
C. neoformans	CN1	16	32	4	-	-
	CN2	8	64	64	-	-
	CN3	16	32	64	-	-
A. fumigatus	AF1	16	32	-	-	0.5
	AF2	8	32	-	-	2
	AF3	16	32	-	-	4
Fusarium	FO1	>64	>64	-	-	4

	FO2	>64	>64	-	-	8
	FS1	32	64	-	-	8
Scedosporium	LP1	16	>64	-	-	>16
	SA1	16	>64	-	-	1
	SB1	16	>64	-	-	2
Rhizopus	RA1	32	32	-	0.25	-
	RA2	32	32	-	0.25	-
	RA3	64	>64	-	0.25	-
Apophysomyces	AP01	32	>64	-	≤0.03	-
	AP02	32	>64	-	0.25	-
Saksenaea	SAK1	16	16	-	≤0.03	-
	SAK2	32	64	-	0.06	-
	SAK3	4	32	-	≤0.03	-
Blastomyces	BD1	8	8	-	-	0.5
	BD2	4	16	-	-	0.06
	BD3	4	8	-	-	≤0.03
C. glabrata	BG2	30	50	-	-	-
C. albicans	95-98-flu resistant	30	50	-	-	-
C. auris		30	50	-	-	-
Trichophyton equinum		-	15	-	-	-

Next, we determined the viability of fungal cells that were exposed to **1**. Cells exposed to

50 μg/ml (MIC) for 24 h or at 100 μg/ml for 48h were used. To determine the viability of treated

fungal cells, small volumes (1 - 5 μg/ml) of mixed cell suspensions were removed from wells and

spotted on YPD agar medium. The agar plates were incubated at 30°C for 24 h - 72 h and the

growth of fungi were recorded. Growth of yeasts occur in 24 h and filamentous fungi in 48 - 72 h

for control (without **1**) but not for those treated with **1** suggesting that it acts as a fungicidal

compound (data not shown).

Drimenol acts better than fluconazole against *Candida auris* growth

C. auris is an emerging multidrug resistant fungal pathogen that is known to cause nosocomial

infections with 'superbug"-like traits (18). *C. auris* was first discovered in 2009 in Southeast Asia

and now it is present in 33 countries across 6 continents. Since this fungus is resistant to all

antifungals and is invasive, the mortality rate is high (19). Recently, CDC has issued a clinical

health emergency warning about this fungus. Since **1** showed a broad-spectrum fungicidal activity,

we determined its effect against *C. auris* growth using a bioscreen-C growth monitoring system.

C. auris was grown in the presence or absence of **1** in RPMI medium (CLSI method) (17) for 24

h at 37° C. Negative controls (solvent) and positive controls (FLU) were included in parallel.

Results depicted in **Fig. 4** indicate that **1** inhibited *C. auris* growth completely at 60 μg/ml. In

contrast, FLU at the same concentration (60 μg/ml) showed poor inhibition of growth. Thus, **1**

could be useful as a broad-spectrum fungicidal compound.

FIG. 4. Drimenol (1) inhibits *C. auris* growth better than fluconazole. *C. auris* was grown in

honeycomb microtiter wells containing RPMI medium in the presence and absence of **1** for 20 h

at 37º C. Fungus growth was measured by absorbance at OD 600 nm using Bioscreen-C growth

monitor. Growth curves showed the mean of triplicates and experiments were repeated at least

twice. Error bars are SD and were too short to appear in the line graphs.

Drimenol (1) is tolerated by *C. elegans* and protects it from fungal mediated death

Invertebrate animal models provide an inexpensive and powerful platform to test antifungal

compounds for their efficacy and toxicity simultaneously. We evaluated **1** for its antifungal

activity and tolerance in *C. elegans* infection model of candidiasis as described before (20).

Results shown in **Fig. 5** indicate that **1** can protect worms from *C. albicans* mediated death and

that the worms were not adversely affected by **1**, as judged by their motility and viability following

compound exposure.

−Dri   +Dri

FIG. 5. Protection of *Caenorhabditis elegans* worms from *C. albicans* mediated death by

drimenol. *C. albicans* (yeast cells) fed larvae were incubated in RPMI medium without and with

1 (+Dri) (50 µg/ml) in a 96 well microtiter plate and incubated at 30 °C for 2 – 3 days. Left panel

without drimenol shows died worms (straight and immobile, thick arrows) due to *C. albicans*

growth. Thin arrow shows a weakly moving worm. Right panel shows **1** containing well where

worms are alive as judged by their movements and the lack of fungal growth.

Mechanism of drimenol (1) antifungal activity

To understand the compound's mechanisms of action (MOA), researchers have used pooled library

of genome-wide barcoded mutant collections of *Saccharomyces cerevisiae* or *C. albicans* for drug-

induced sensitivity assay or the haploinsufficiency (HIP) assay (21-23). For example, if **1** can

inactivate partially or completely its protein target in the heterozygous mutant pool, the resulting

growth defect of that mutant(s) can be measured quantitatively by sequencing the tagged unique

barcodes. This approach will help narrowing down the putative target(s). Similarly, a

homozygous nonessential mutant library can be used as a complementary approach to the

heterozygous essential mutant collection to verify the target pathway/genes of compounds. In this

case, if the homozygous mutant of a gene is sensitive to the compound, then that gene may not be

the drug target (21) as the homozygous mutant lacks the gene product. This imples that the

compound may exert its effect via drug-induced synthetic lethality. Thus, by combining data from

both heterozygote and homozygote screens one may determine the compound's MOA.

In this study, we used *S. cerevisiae* barcoded homozygous nonessential and heterozygous essential,

and *C. albicans* barcoded heterozygous Tn mutant (23) libraries. Briefly, IC-50 of **1** for *C.*

albicans and *S. cerevisiae* was determined in yeast growth (YPD) condition (**Fig. 6**). Based on

this assay results, IC-50 of 25 μg/ml for *C. albicans* and 15 μg/ml for *S. cerevisiae* was calculated

for **1**. Two different sub-MIC concentrations of **1** were selected for determining the mechanism

of action against mutant libraries.

FIG. 6. Determination of IC-50 for drimenol (1) activity against *C. albicans* and *S. cerevisiae*.
Yeast growth conditions (YPD medium at 30ºC) were used to determine the IC-50 values. The red line in each panel indicates growth of the DMSO reference. An IC-50 of ~25 μg/ml for *C. albicans* and ~15 μg/ml for *S. cerevisiae* were calculated for **1**. Albicanol (**2**) showed weaker activity against both fungi and was not considered for further analysis.

Next, pooled *S. cerevisiae* and *C. albicans* mutant collections were grown separately in the presence or absence of compounds (with DMSO) for 20-generations, barcodes from genomic DNA were amplified and relative strain abundance were quantified based on TAG microarray signals. The $\log_2$ ratio of tag signals between DMSO control and **1** exposed samples were presented in scatter plots as the "fitness defect" (**Figs. 7A & B**). Mutants that were depleted from the growth pool due to **1** are indicated by circles. Mutants that were highly susceptible to **1** are shown with high log ratio (*e.g.* SEC66 in **Fig. 7A**) (highly depleted in the pool) and considered putative targets.

Lists of *S. cerevisiae* and *C. albicans* mutants that are highly sensitive to **1** were given in

Supplementary Tables 1 and 2, respectively.

FIG. 7. **Genome-wide screens of** *S. serevisiae* **(A) and** *C. albicans* **(B) mutant libraries against**

drimenol (1) for drug induced hypersensitivity. Pooled collections of *S. cerevisiae* nonessential

homozygous and essential heterozygous mutants were grown in the presence and absence of **1** at

the concentration of 0.025 mg/ml for the indicated number of generations before profiling for their

abundance (DNA barcodes). Twenty generations for essential heterozygous and five generations

for non essential homozygous mutants were used which gave an optimum of ~20% growth

inhibition. Similarly, *C. albicans* Tn-insertion mutants (heterozygous, 20 generations) was used

at 0.025 mg/ml of **1.** Each spot represents single mutant. The log ratio of each mutant (**1** exposed

vs no drug control) was calculated and presented in scatter plots where greater the number the

more sensitive that strain is.

Our results of forward genetic screening from mutant libraries of *C. albicans* and *S. cerevisiae*

with **1** indicate that it affects cellular activities involved in protein secretion, vacuolar functions,

chromation remodeling and cyclin dependent protein kinases (CDK) (**Fig. 7** and **Supplementary**

Tables 1 & 2). For example, SEC66, highly sensitive to **1** ($\log_2$ >4.5, **Fig. 7A**), is a component of

Sec63 SECretary complex in *S. cerevisiae* involved in protein targeting and importing into the ER.

Similarly, VMA22 is a vacuolar membrane ATPase required for vacuolar H+-ATPase function

and localized to the yeast ER (Saccharomyces Genome Database).

The *ISN1* ($\log_2$ 4.4, **Fig. 7B**) gene product is involved in inosine 5'-monophosphate 5'-

nucleotidase activity in *C. albicans* (Candida Genome Database). This gene product is

uncharacterized and it is present only in fungi and not in human or murine, suggesting that Isn1p

a suitable antifungal drug target. *IFG3* is a putative D-amino acid oxidase, which is

uncharacterized, and *CRK1* is a protein kinase of the Cdc2 subfamily involved in hyphal

development and virulence in *C. albicans* (24). The *CRK1* ortholog in *S. cerevisiae* is *SGV1,*

which is a part of BUR2 kinase complex and plays a major role in transcriptional regulation.

Yeasts spot assay to validate drimenol (1) mechanism of action

296 Based on the forward genetic library screening assay results (**Fig. 7**) and their functions inferred

297 from the available literature, we selected few heterozygous mutants of *C. albicans* that had high

298 to medium positive $\log_2$ ratio (hypersensitive, *CRK1* and its putative interacting partners proteins

FIG. 8. Validation of *C. albicans* Tn-insertion mutant screen data by yeast spot assay. A four-

fold serially diluted yeast cultures of indicated *C. albicans* heterozygous mutants (GRACE) were

spot tested on YPD agar containing **1** (30 μg/ml) or DMSO (- **1**). Heterozygous mutants (*RET2/-*

, *Orf19.759/-, Orf19.1672/-*, and *Orf19.4382/-*) affected by **1** directly or indirectly were

hypersensitive and showed lack of growth.

CDC37, Orf19.759, Orf19.1672 and Orf19.4382) and few with negative $\log_2$ ratio (resistance,

VPS53, *TSC11* & *PHO89*) to verify the genetic screening data. Agar medium containing sub-MIC

concentration of **1** (30 μg/ml) or solvent was used to spot test suspensions of various mutants

(GRACE mutant collection) (25).

Results shown in **Fig. 8** confirm the findings of forward genetic screening assays of *C. albicans*

mutants. For example, Crk1 Kinase is predicted to associate with *RET2*, which involves in

retrogradetion of vesicle transport for protein signaling and secretion (*SEC*66), and plays a role in

protein translation with G1/S cell-cycle transition (26). Products encoded by Ret2, orf19.759

(*SEC21*), orf19.1672 (*COP*1) and orf19.4382 (*RET3*) are uncharacterized and are likely targets of

Crk1 kinase, which are defective in growth on agar medium containing **1** (**Fig. 8**). Molecule **1**

induced hypersensitivity of these mutants, which represent candidate targets. The orf19.759

(*SEC21* ortholog of *S. cerevisiae*) is uncharacterized in *C. albicans*. Sec21 involves in transport

from endoplasmic reticulum (ER) to Golgi vsicle-mediated transport (anterograde), Golgi to ER

(retrograde) and COPI vesicle coat, and endosome localization (27). *RET2* is also uncharacterized

in *C. albicans* and the ortholog is a subunit of the coatomer complex (COPI), which coats Golgi-

derived transport vesicles, involves in retrograde transport between Golgi and ER, and interacts

with Crk1 kinase in the two-hybrid system (28). Crk1 is known to play a role in regulating

trafficking and secretion of effectors by interacting with the early endosome during *Ustilago*

maydis (corn smut fungus) infection in corn plants (29, 30).

Since Crk1 kinase may interact with multiple targets (Ret2, orf19.759, orf19.1672 and orf19.4382)

and because Crk1 represents an important antifungal drug target (24), we performed computational

molecular docking of **1** with *Cryptosporidium parvum* Crk1 crystal structure (2QKR-A) (31) using

AutoDock Vina software (32). Results showed in **Fig. 9**, suggest that **1** can interact with the *N*-

terminal catalytic domain of *C. albicans* Crk1 (which has 61% similarity and 40% identical to the

C. parvum Crk1). Particularly, noteworthy from our computational docking studies is that **1** has

close interactions with Gly 31, Val 37, Gln 148, Leu 151, and Phe 98 amino acid residues. The

docked structure shows that an available open space in Crk1 for incorporation of an additional

function group onto the cyclohexane A ring of **1 (Fig. 1)**, signifying a possible modification of **1**

for future improvement of biological activity. Thus, this CDK member may comprise the target

of **1** and it is notable that this conserved gene is present in many of the tested pathogenic fungi

(**Table 2**).

353
354 **FIG. 9. Molecular docking of drimenol with *C. parvum* Crk1 kinase**.
355
356
357

DISCUSSION

Fungi have emerged in the last two decades as major causes of human disease. *C. albicans* is a

major fungal pathogen affecting at all ages and fourth leading cause of nosocomial bloodstream

infections in the US (33). *C. albicans* and other *Candida* spp cause mucosal, disseminated and

invasive candidiasis, especially among patients who are immunocompromised or hospitalized with

serious underlying diseases. The overall mortality for invasive diseases caused by *Candida* spp.

and *Aspergillus* spp. is around 50% (33, 34). While there are more than 150 species of Candida,

about 15 species are recognized as frequent human pathogens (34, 35). Some of them are: *C.*

albicans, C. glabrata, C. krusei, C. tropicalis and *C. parapsilosis*. Among these, *C. albicans* is by

far the most common species isolated from humans and is a frequent denizen of the oropharynx,

mucousal surfaces, gastrointestinal and genitourinary tracts. In the developing world, there are ~1

million cases of cryptococcal diseases per year resulting in 675,000 deaths (9, 36). *Cryoptococcus*

neoformans is an opportunistic fungal pathogen that causes meningitis in immunocompromised

individuals. Often found in soils contaminated with bird feces, *C. neoformans* enters its host

through the lungs via inhalation of spores. Some of the cryptococcal species are hypervirulent

(37) and have drawn a considerable public attention due to their causative role in the

cryptococcosis outbreak throughout the Pacific Northwest (38, 39). Only few antifungals can be

useful to treat cryptococcosis and drug resistant strains are emerging.

Aspergillus spp are ubiquitous molds found widely in the environment as saprophytes and produce

microscopic spores or conidia which upon inhalation, cause invasive pulmonary disease. In

immunocompromised patients such as hematopoietic stem cell transplantation, solid organ

transplantation and undergoing chemotherapeutic agents or immunomodulatory agents, invasive

aspergillosis remains the most important cause of infection-related mortality (40, 41). Among

several species of Aspergillus, *A. fumigatus* and *A. flavus* are frequent pathogens. Dermatophytes

are another group of keratinophilc pathogenic fungi that causes variety of infections in humans

and animals (42). Some of these fungi include *Trichophyton tonsurans* (scalp ring-worm), *T.*

equinum, and *Microsporum gypseum* (garderner's ringworm). Emerging fungal disease such as

zygomycosis is life-threatening particularly during natural calamity (*e.g.* the 2004 tsunami, the

2008 Katrina and May-2011 Joplin tornado). Novel compounds with broad-spectrum antifungal

activity are highly desirable to treat various fungal pathogens.

Because fungi are eukaryotes, the development of antifungal therapeutics that are nontoxic to

human is challenging due to the availability of relatively few targets. In the last thirty years, only

one new class of antimycotic (β-glucan synthase inhibitor, the echinocandins) was introduced into

clinical practice. Although this drug is an important addition, it has a number of limitations

including ineffectiveness against *Cryptococcus sp* and poor oral bioavailability (43). New drugs

are needed to be discovered and developed.

In our search for novel antifungal small molecules from our available synthetic terpenoids, we

have identified two compounds, (-)-drimenol (**1**) and (+)-albicanol (**2**) (**Fig. 1**), that showed strong

activity against *C. albicans*. Among these two compounds, **1** showed stronger bioactivity. It acts

not only against *C. albicans* as fungicidal but also against *Aspergillus nidulans*, FLU resistant

strains of *C. albicans*, *C. glabrata, C. krusei, Cryptococcus* spp and other dermatophytes,

suggesting that **1** is a broad-spectrum antifungal agent (**Figs. 2, 3** and **Table 2**). At an increased

concentration (100 μg/ml), **1** caused rupturing of the fungal cell wall/membrane, *e.g*. *C. albicans*

(**Fig. 3**) and *Cryptococcus sp.* (data not included). *Candida auris* is an emerging and multi

antifungal resistant strain that causes nosocomial infection and has been reported recently across

the world (18). Our bioscreen-based growth curve monitoring assay with **1** showed better activity

than clinical antifungal drug FLU (**Fig. 4**) indicating a potential use of **1** against *C. auris* and other

drug-resistant fungal pathogens.

Molecule **1** is a natural product presents in liverworts and higher plants (44, 45), and its antifungal

mechanism against human pathogenic fungi has not been reported previously. A recent study has

shown that **1** has antifungal activity against *Botrytis cinerea*, a plant fungal pathogen and the

mechanisms appear to act by fungal membrane damage and reactive oxygen species (ROS)

production (46).

In order to develop broad-spectrum novel antifungal compounds, we evaluated **1** against various

fungi that are pathogenic to humans and determined its mechanisms of action in *C. albicans* and

S. cerevisiae. Based on our yeasts mutant screening data and subsequent spot assay results, we

found that **1** acts as a fungicidal compound by affecting cellular activities targeting protein

trafficking between Golgi to ER, protein secretion (Sec system) and cell signaling, possibly

through cell division related kinase 1, Crk1 (**Figs. 7 & 8**). Genetic methods have been used to

determine the mechanism of antifungal compounds by drug-induced hypersensitivity assay (22,

47). Using similar approaches, we showed that **1**-mediated inhibition of *C. albicans* heterozygous

mutants of *CDC37, Orf19.759, Orf19.1672 and Orf19.4382*, the known or putative targets of Crk1

kinase, at sub-MIC concentration. In support of this observation, computational molecular

docking of **1** with the crystal structure of a fungal (*C. parvum*) Crk1 kinase showed interactions of

1 with the key residues in the catalytic domain (*N*-terminal) of Crk1 (**Fig. 9**).

Cinnamodial is a closely related compound belong to the drimane sesquiterpenoid family with

potent antifungal activity (48), but its chemical structure (containing dialdehyde groups; **Fig. 1**)

and physiological properties are quite different from **1**. For example, the antifungal activity of

cinnamodial was shown to abolish by amine compounds (likely due to a coupling reaction from

the aldehyde functions of cinnamodial with the amino group of amine compounds) or when

cinnamodial was incubated in YPD medium (19, 49). In contrast, molecule **1**'s bioactivity was

not affected by amines or YPD medium (**Fig. 5**). Thus, the antifungal mechanisms of **1** could be

different from cinnamodial. Since the synthetic route for **1** and its analogs are well established,

improvements of its antifungal properties are possible through medicinal chemistry approaches.

In summary, we have synthesized a focused library of drimane sesquiterpenoid compounds and

identified **1** as a broad-spectrum fungicidal compound against various human pathogenic fungi

including *C. albicans, C. auris, Cryp. neoformans*, *Aspergillus, Blastomyces, Scedosporium,*

Fusarium, Pneumocystis, and dermatophytes at 8 - 64 µg/ml. By employing the libraries of bar-

coded *C. albicans* and *S. cerevisiae* genome-wide mutants, the mechanism of action of **1** was

determined. Further evaluation of **1** in animal models of fungal diseases would help develop **1** as

an antifungal agent.

MATERIALS AND METHODS

Synthesis of drimane molecules. (1*R*,2*R*,4a*S*,8a*S*)-2-Hydroxy-2,5,5,8a-tetramethyl-

decahydronaphthalene-1-carbaldehyde (4). To a solution of 0.20 g (0.74 mmol) of triol **3** (13)

in 10 ml of THF and 2.5 ml of water was added 0.19 g (0.81 mmol) of potassium periodate. The

resulting mixture was stirred at 25°C for 4 hours, diluted with water (50 ml) and extracted three

times with ethyl acetate (50 ml each). The combined extract was washed with water and brine,

dried (anhydrous Na_2SO_4), concentrated, and column chromatographed on silica gel using a

mixture of hexane and ethyl acetate (20:1) as an eluent to give 0.16 g (91% yield) of compound **4**,

whose spectral data is in agreement with that reported (13).

(1*S*,2*R*,4a*S*,8a*S*)-1-(Hydroxymethyl)-2,5,5,8a-tetramethyl-decahydronaphthalen-2-ol (5). To

a cold (0°C) solution of 1.0 g (4.2 mmol) of aldehyde **4** in 80 mL of diethyl ether under argon, 80

mg (2.1 mmol) of lithium aluminum hydride was added in portions. The resulting solution was

stirred at 0°C for 30 minutes, diluted with aqueous NH_4Cl, and extracted with diethyl ether three

times (50 ml each). The combined extract was washed with water and brine, dried ($MgSO_4$), and

concentrated to give 0.98 g (97% yield) of diol **5**, whose spectral data are in agreement with that

reported (13).

[(1*S*,2*R*,4a*S*,8a*S*)-2-Hydroxy-2,5,5,8a-tetramethyl-decahydronaphthalen-1-yl]methyl

acetate (6). To a cold (0°C) solution of 0.10 g (0.40 mmol) of diol **5** in 2 ml of dichloromethane

and 0.32 g (4.0 mmol) of pyridine under argon, was added 49 μl (0.48 mmol) of acetic anhydride,

and the resulting solution was stirred at 0°C for 30 minutes and 25°C for 1 h. It was diluted with

30 ml of aqueous NH_4OH, extracted twice with diethyl ether (30 ml each), and the combined

extracts were washed with water and brine, dried (anhydrous Na_2SO_4), concentrated, and column

chromatographed on silica gel using a gradient mixture of hexane and diethyl ether as eluents to

give 90 mg (80% yield) of acetate **6**. Mp. 64 – 67 °C; $[\alpha]^D_{22}$ = -8.2 (c = 0.55, $CHCl_3$); [1]H NMR

($CDCl_3$; 400 MHz) δ 4.35 (dd, *J* = 12, 4 Hz, 1 H), 4.24 (dd, *J* = 12, 4 Hz, 1 H), 2.05 (s, 3 H), 1.88

(dt, *J* = 12, 2 Hz, 1 H), 1.70 – 0.93 (a series of m, 11 H), 1.17 (s, 3 H), 0.88 (s, 3 H), 0.86 (s, 3 H),

0.80 (s, 3 H) ppm; ^{13}C NMR (CDCl₃; 100 MHz) δ 171.4, 72.6, 62.6, 60.0, 55.7, 44.0, 41.7, 39.7,

38.1, 33.5, 33.2, 24.6, 21.6, 21.3, 20.3, 18.4, 15.8 ppm. MS (electrospray ionization), m/z 283.1

(M+H⁺). HRMS-ESI: m/z [M + H]⁺ calcd for $C_{17}H_{31}O_3^+$: 283.2268, found: 283.2273.

(1*S*,2*R*,4a*S*,8a*S*)-1-(Hydroxymethyl)-2,5,5,8a-tetramethyl-decahydronaphthalen-2-yl

acetate (7). Compound **7** was prepared by a sequence of three reactions: (i) silylation of the

primary alcohol function of **5** with *t*-butyldimethylsilyl chloride; (ii) acetylation of the tertiary

alcohol function with acetyl chloride and pyridine; and (iii) removal of the *t*-butyldimethylsilyl

ether protecting group with tetra-*n*-butylammonium fluoride in THF.

To a solution of 9.5 mg (40 µmol) of compound **5**, 11 mg (150 µmol) of imidazole, and 6 mg (49

µmol) of 4-(dimethylamino)pyridine in 2 ml of dichloromethane under argon at 25°C, was added

14.3 mg (95 µmol) of *t*-butyldimethylsilyl chloride, and the solution was stirred for 4 h. The

reaction mixture was diluted with 10 ml of aqueous ammonium chloride and extracted with diethyl

ether three times (10 ml each). The combined extracts were washed with water (10 ml) and brine

(10 ml), dried (anhydrous Na₂SO₄), concentrated to give 12.5 mg of the mono-silylated product.

This crude product was used in the subsequent step without purification. To a solution of the

above mono-silylated product and 0.1 ml of pyridine in 0.5 ml of dichloromethane under argon at

0°C, was added 10 µl (0.13 mmol) of acetyl chloride. The reaction mixture was stirred at 25°C for

2 h, diluted with aqueous ammonium chloride (10 ml), and extracted three times with diethyl ether

(10 ml each).   The combined extracts were washed with brine, dried (anhydrous Na₂SO₄),

concentrated to give the crude product, which was used in the following step without purification.

The above crude product was dissolved in 1 ml of dried THF (distilled over sodium/benzophenone)

and 0.3 ml (0.3 mmol) of tetra-*n*-butylammonium fluoride (1 M solution in THF) and stirred at

25°C under argon for 1 h. The reaction solution was diluted with 0.1 N ammonium hydroxide (10

ml) and extracted with diethyl ether three times (10 ml each). The combined extracts were washed

with water (10 ml) and brine (10 ml), dried (anhydrous Na_2SO_4), concentrated, and column

chromatographed on silica gel using a gradient mixture of hexane and diethyl ether to give 4.2 mg

(38% overall yield from diol **5**) of compound **7**. Compound **7**: Mp. 101 – 103 ºC; $[\alpha]^D_{22}$ = +0.35

(c = 0.23, CHCl$_3$); ^{1}H NMR (CDCl$_3$; 400 MHz) δ 3.91 (dd, J = 12, 2 Hz, 1 H), 3.84 (dd, J = 12, 2

Hz, 1 H), 2.95 – 2.90 (m, 1 H), 1.98 (s, 3 H), 1.88 (dt, J = 12, 2 Hz, 1 H), 1.62 (s, 3 H), 1.70 – 0.88

(a series of m, 10 H), 0.94 (s, 3 H), 0.87 (s, 3 H), 0.82 (s, 3 H) ppm; ^{13}C NMR (CDCl$_3$; 100 MHz)

δ 169.9, 84.9, 63.8, 59.8, 55.8, 41.8, 39.5, 38.2, 36.1, 33.5, 33.2, 25.8, 22.8, 21.7, 18.3 (2 C), 16.1

ppm. MS (electrospray ionization), m/z 305.1 (M+Na$^+$). HRMS-ESI: m/z [M + Na]$^+$ calcd for

$C_{17}H_{30}NaO_3{}^+$: 305.2087, found: 305.2082.

(1S,2R,4aS,8aS)-1-(2,2-Dimethyl-1,3-dioxolan-4-yl)-2,5,5,8a-tetramethyl-

decahydronaphthalen-2-ol (8). A solution of 18 mg (67 μmol) of triol **3**, 50 μl of 2,2-

dimethoxypropane and 3 mg of anhydrous p-toluenesulfonic acid in 1 ml of toluene was stirred

under argon at 55ºC for 1 h. The solution was cooled to room temperature, neutralized with sodium

bicarbonate (~3 mg), diluted with 10 ml of water, and extracted with ethyl acetate three times (15

ml each). The combined extract was washed with brine, dried (MgSO$_4$), concentrated and column

chromatographed on silica gel using a gradient mixture of hexane and diethyl ether as eluent to

give 14 mg (71% yield) of compound **8** as a mixture of two stereoisomers: (the major isomer was

partially purified and reported) Mp. 114 – 117 ºC; $[\alpha]^D_{22}$ = -25.1 (c = 1.0, CHCl$_3$); ^{1}H NMR

(CDCl$_3$; 400 MHz) δ 4.96 (s, 1 H, OH), 4.24 – 4.20 (m, 2 H), 3.59 (td, J = 8, 4 Hz, 1 H), 1.84 (dt,

J = 12, 2 Hz, 1 H), 1.70 – 0.83 (a series of m, 11 H), 1.45 (s, 3 H), 1.41 (s, 6 H, 2 CH$_3$), 0.97 (s, 3

H), 0.90 (s, 3 H), 0.83 (s, 3 H) ppm; ^{13}C NMR (CDCl$_3$; 100 MHz) δ 107.4, 73.5, 72.8, 62.2, 55.7,

42.8, 41.5, 40.4, 37.3, 33.6, 33.3, 26.5, 26.2, 25.8, 21.7, 19.7, 18.4 (2 C), 16.1 ppm. MS

(electrospray ionization), m/z 333.1 (M+Na$^+$). HRMS-ESI: m/z [M + Na]$^+$ calcd for $C_{19}H_{34}NaO_3^+$:

333.2406, found: 333.2411.

Determination of Antifungal activity of synthetic compounds

Synthetic pure drimenol or albicanol was dissolved in DMSO (10 mg/ml as stock solution) and

used for determining their antifungal activities (minimum inhibitory concentration, MIC) against

various fungi according to the microdilution assay of CLSI (17). The CLSI broth dilution methods

of M27-A3 for yeasts and M38-A for filamentous fungi were used to determine the susceptibility.

Since our initial assay with *C. albicans* confirmed the antifungal activity of drimenol and albicanol,

we extended the susceptibility assay to other pathogenic fungi including FLU resistant *C. albicans*,

various species of candida, *Cryptococcus*, *Aspergillus* and a dermatophyte fungus (strains were

generously provided by Dr. Ted C. White at The University of Missouri Kansas City (UMKC). *C.*

auris was obtained from Dr. Baha Abdalhamid at The University of Nebraska Medical Center,

Omaha NE). Briefly, yeast cells or conidia (for filamentous fungi) were suspended in RPMI 1640

medium to a final concentration of 10^5 cfu/ml and distributed in 96-well microplate to a total

volume of 100 μl/well. Drimenol or albicanol was added into the wells and a two-fold serial

dilution was made. Duplicates were used for each concentration and wells with or without DMSO

served as controls. Plates were incubated without shaking at 37^{o}C for 24 - 48h for yeasts and 30^{o}C

for 4 days for filamentous fungi (*Aspergillus* sp and *Trichophyton* sp). The MIC was defined as

the lowest compound concentration at which no growth occurred, as determined visually and

microscopically (inverted microscope).

Determination of *C. auris* growth inhibition by drimenol

The effect of drimenol on the growth of *C. auris* was determined by Bioscreen-C real time growth

monitoring system (Oy Growth Curves Ab Ltd, Finland) as described earlier (50). Briefly, 200 µl

of RPMI medium containing exponentially growing *C. auris* yeast cells (each at 0.07 OD600)

were added into the honeycomb wells with or without compound (control) and measured their

growth rates for 20 hours at 37ºC. Compound treatment was done at two different concentrations

for drimenol (50 and 60 µg/ml). The absorbance was measured at 600 nm at 30 min intervals for

24 h at 37°C with shaking for 10 s before each reading. Solvent negative control (DMSO) and

FLU (60 µg/ml; antifungal drug) positive control were included in the study. The experiments

were repeated at least two times with three technical replicates.

Yeast spot assay

Yeast Peptone Dextrose (YPD) agar containing a sub-MIC concentration of drimenol (30 µg/ml)

or an equal volume of DMSO was used to spot test the *C. albicans* heterozygous mutants (GRACE

library (25)). Yeast suspensions of various mutants and the wild type *C. albicans* were used. Five

µl of a four-fold serially diluted suspension was spotted on the agar plates and incubated at 30ºC

for yeast growth for 24 h, and photographed. Experiments were repeated at least three times and

a representative result was shown.

Genome-wide fitness assay

The Saccharomyces yeast deletion collection was comprised of approximately 5,900 individually

bar-coded heterozygous diploid strains (HIP [haploinsufficiency profiling]) and ~4,800

homozygous diploid strains (HOP [homozygous deletion profiling]). Pools of approximately

equal strain abundance were generated by robotically pinning (S and P Robotics, Ontario, Canada)

each strain (from frozen stocks) onto YPD agar plates as arrays of 384 strains/plate (21, 51, 52).

After 2 days of growth at 30°C, colonies were collected from plates by flooding with YPD, and

cells were adjusted to an optical density at 600 nm (OD_{600}) of 2. The fitness of each strain in each

experimental pool was assessed as described previously (21). The dose that resulted in 15%

growth inhibition in *S. cerevisiae* BY4743 (the parent strain of the yeast deletion collection) was

determined by analyzing dose response over the course of 16 h of growth at 30°C. Screens of the

homozygous deletion collection were performed for 5 generations of growth and screens of the

heterozygous deletion collection were collected after 20 generations of growth. Cells were

processed as described previously (21). Genomic DNA was extracted from each sample and

subjected to PCR to amplify the unique bar code identifiers. The abundance of each bar code was

determined by quantifying the microarray signal as previously described (21).

Candida albicans pooled screens used the tn-transposon collection (23). Growth assays were

performed in duplicate and samples were recovered at 20 generations of growth. Genomic DNA

extraction, tag amplification, and hybridization were performed as described above.

Conflict of interest: A US patent (US 8,980,951 B2) on synthetic drimenol was approved in 2015

to Kansas State University Research Foundation (KSURF) with authors GV and DHH.

Supplementary data.

Supplementary Table 1. *S. cerevisiae* genetic screening data

Supplemenary Table 2. *C. albicans* genetic screening data

Acknowledgement. We thank Ted C. White, University of Missouri Kansas City (UMKC),

Division of Cell Biology and Biophysics, Kansas City for various species of candida including

fluconazole-resistant strains, *Cryptococcus*, *Aspergillus* and dermatophytes. Our sincere thanks to

Terry Roemer, Merck Sharp and Dohme Corp, NJ for providing *C. albicans* GRACE and DBC

mutant libraries. *C. auris* was obtained from Baha Abdalhamid, University of Nebraska Medical

Center, Omaha NE. Kansas State University has utilized the non-clinical and pre-clinical service

program offered by the National Institute of Allergy and Infectious Diseases at the Fungus Testing

Laboratory, UTHSC San Antonio. Our sincere thanks to Nathan Wiederhold, Hoja Patterson, and

April Torres for their assistance in testing drimenol for antifungal activity through the NIH service

program. We thank Johnson Cancer Research Center, KSU for Innovative Research Award and

Kansas Idea Network of Biomedical Research Excellence (K-INBRE) for CORE Facility support

award to GV. CN is a CRC Tier 1 Chair in Translational Genomics and support for GG and CN

is provided by the CIHR. Research reported in this publication was, in part, supported by the

National Institute of General Medical Sciences of the National Institutes of Health under award

number NIH R01GM128659 (to DHH). The content is solely the responsibility of the authors and

does not necessarily represent the official views of the National Institutes of Health. This material

was based upon work in part supported by the National Science Foundation under 1826982 (to

DHH) for the purchase of a 400-MHz NMR spectrometer.

References

1.   Fuentefria AM, Pippi B, Dalla Lana DF, Donato KK, de Andrade SF. 2018. Antifungals

discovery: an insight into new strategies to combat antifungal resistance. Lett Appl

Microbiol 66:2-13.

2.   Perfect JR. 2017. The antifungal pipeline: a reality check. Nat Rev Drug Discov 16:603-

616.

3.   Perlin DS, Rautemaa-Richardson R, Alastruey-Izquierdo A. 2017. The global problem of

antifungal resistance: prevalence, mechanisms, and management. Lancet Infect Dis

17:e383-e392.

4.   Nami S, Aghebati-Maleki A, Morovati H, Aghebati-Maleki L. 2019. Current antifungal

drugs and immunotherapeutic approaches as promising strategies to treatment of fungal

diseases. Biomed Pharmacother 110:857-868.

5.   Nett JE, Andes DR. 2016. Antifungal Agents: Spectrum of Activity, Pharmacology, and

Clinical Indications. Infect Dis Clin North Am 30:51-83.

6.   Waldorf AR, Polak A. 1983. Mechanisms of action of 5-fluorocytosine. Antimicrob Agents

Chemother 23:79-85.

7.   Denning DW. 2002. Echinocandins: a new class of antifungal. J Antimicrob Chemother

49:889-91.

8.   Roemer T, Krysan DJ. 2014. Antifungal drug development: challenges, unmet clinical

needs, and new approaches. Cold Spring Harb Perspect Med 4:pii: 4/5/a019703. doi:

10.1101/cshperspect.a019703.

9.   Denning DW, Hope WW. 2010. Therapy for fungal diseases: opportunities and priorities.

Trends Microbiol 18:195-204.

10.   Pfaller MA, Diekema DJ. 2007. Epidemiology of invasive candidiasis: a persistent public

health problem. Clin Microbiol Rev 20:133-63.

11.   Marie C, White TC. 2009. Genetic Basis of Antifungal Drug Resistance. Curr Fungal Infect

Rep 3:163-169.

12.   Sanglard D, Coste A, Ferrari S. 2009. Antifungal drug resistance mechanisms in fungal

pathogens from the perspective of transcriptional gene regulation. FEMS Yeast Res

9:1029-50.

13.   Hua DH, Huang X, Chen Y, Battina SK, Tamura M, Noh SK, Koo SI, Namatame I,

Tomoda H, Perchellet EM, Perchellet JP. 2004. Total syntheses of (+)-chloropuupehenone

and (+)-chloropuupehenol and their analogues and evaluation of their bioactivities. J Org

Chem 69:6065-78.

14.   Madikane VE, Bhakta S, Russell AJ, Campbell WE, Claridge TD, Elisha BG, Davies SG,

Smith P, Sim E. 2007. Inhibition of mycobacterial arylamine N-acetyltransferase

contributes to anti-mycobacterial activity of *Warburgia salutaris*. Bioorg Med Chem

15:3579-86.

15.   Cortes M, Delgado V, Saitz C, Armstrong V. 2011. Drimenol: A versatile synthon for

compounds with trans-drimane skeleton. Nat Prod Commun 6:477-90.

16.   Liu Y, Liu Z, Cao X, Liu X, He H, Yang Y. 2011. Design and synthesis of pyridine-

substituted itraconazole analogues with improved antifungal activities, water solubility and

bioavailability. Bioorg Med Chem Lett 21:4779-83.

17.   CLSI. 2008. Reference method for broth dilution antifungal susceptibility testing of yeasts

and filamentous fungi; Approved Standard-Third Edition. Clinical and Laboratory

659   Standards Institute, 940 West Valley Road, Wayne, Pa. CLSI document M27-A3 & M38-

660   A 28.

661 18. Kean R, Ramage G. 2019. Combined Antifungal Resistance and Biofilm Tolerance: the

662   Global Threat of *Candida auris*. mSphere 4:DOI: 10.1128/mSphere.00458-19.

663 19. de Jong AW, Hagen F. 2019. Attack, Defend and Persist: How the Fungal Pathogen

664   *Candida auris* was Able to Emerge Globally in Healthcare Environments. Mycopathologia

665   184:353-365.

666 20. Vediyappan G, Dumontet V, Pelissier F, d'Enfert C. 2013. Gymnemic acids inhibit hyphal

667   growth and virulence in *Candida albicans*. PLoS One 8:e74189.

668 21. Pierce SE, Davis RW, Nislow C, Giaever G. 2007. Genome-wide analysis of barcoded

669   *Saccharomyces cerevisiae* gene-deletion mutants in pooled cultures. Nat Protoc 2:2958-

670   74.

671 22. Xu D, Jiang B, Ketela T, Lemieux S, Veillette K, Martel N, Davison J, Sillaots S, Trosok

672   S, Bachewich C, Bussey H, Youngman P, Roemer T. 2007. Genome-wide fitness test and

673   mechanism-of-action studies of inhibitory compounds in *Candida albicans*. PLoS Pathog

674   3:e92.

675 23. Oh J, Fung E, Schlecht U, Davis RW, Giaever G, St Onge RP, Deutschbauer A, Nislow C.

676   2010. Gene annotation and drug target discovery in *Candida albicans* with a tagged

677   transposon mutant collection. PLoS Pathog 6:e1001140.

678 24. Chen J, Zhou S, Wang Q, Chen X, Pan T, Liu H. 2000. Crk1, a novel Cdc2-related protein

679   kinase, is required for hyphal development and virulence in *Candida albicans*. Mol Cell

680   Biol 20:8696-708.

25.   Roemer T, Jiang B, Davison J, Ketela T, Veillette K, Breton A, Tandia F, Linteau A,

Sillaots S, Marta C, Martel N, Veronneau S, Lemieux S, Kauffman S, Becker J, Storms R,

Boone C, Bussey H. 2003. Large-scale essential gene identification in *Candida albicans*

and applications to antifungal drug discovery. Mol Microbiol 50:167-81.

26.   An T, Liu Y, Gourguechon S, Wang CC, Li Z. 2018. CDK Phosphorylation of Translation

Initiation Factors Couples Protein Translation with Cell-Cycle Transition. Cell Rep

25:3204-3214 e5.

27.   CGD. 2019. Candida Genome Database, http://www.candidagenome.org/cgi-

bin/locus.pl?locus=Sec21&organism=C_albicans_SC5314.

28.   CGD. 2019. Candida Genome Database, (http://www.candidagenome.org/cgi-

bin/locus.pl?locus=ret2&organism=C_albicans_SC5314.

29.   Higuchi Y. 2015. Initial fungal effector production is mediated by early endosome motility.

Commun Integr Biol 8:e1025187.

30.   Bielska E, Higuchi Y, Schuster M, Steinberg N, Kilaru S, Talbot NJ, Steinberg G. 2014.

Long-distance endosome trafficking drives fungal effector production during plant

infection. Nat Commun 5:5097.

31.   Wernimont AK, Dong A, Lew J, Lin YH, Hassanali A, Arrowsmith CH, Edwards AM,

Weigelt J, Sundstrom M, Bochkarev A, Hui R, Artz JD, (Structural Genomics

Consortium). 2007. *Cryptosporidium parvum* cyclin-dependent kinase cgd5_2510 with

indirubin 3'-monoxime bound. https://wwwrcsborg/structure/2qkr (database release).

32.   Trott O, Olson AJ. 2010. AutoDock Vina: improving the speed and accuracy of docking

with a new scoring function, efficient optimization, and multithreading. J Comput Chem

31:455-61.

33. Pfaller MA, Diekema DJ. 2010. Epidemiology of invasive mycoses in North America. Crit

Rev Microbiol 36:1-53.

34. Pappas PG. 2006. Invasive candidiasis. Infect Dis Clin North Am 20:485-506.

35. Pfaller MA, Diekema DJ, Rinaldi MG, Barnes R, Hu B, Veselov AV, Tiraboschi N, Nagy

E, Gibbs DL. 2005. Results from the ARTEMIS DISK Global Antifungal Surveillance

Study: a 6.5-year analysis of susceptibilities of Candida and other yeast species to

fluconazole and voriconazole by standardized disk diffusion testing. J Clin Microbiol

43:5848-59.

36. Park BJ, Wannemuehler KA, Marston BJ, Govender N, Pappas PG, Chiller TM. 2009.

Estimation of the current global burden of cryptococcal meningitis among persons living

with HIV/AIDS. AIDS 23:525-30.

37. Byrnes EJ, 3rd, Li W, Lewit Y, Ma H, Voelz K, Ren P, Carter DA, Chaturvedi V, Bildfell

RJ, May RC, Heitman J. 2010. Emergence and pathogenicity of highly virulent

Cryptococcus gattii genotypes in the northwest United States. PLoS Pathog 6:e1000850.

38. Kidd SE, Hagen F, Tscharke RL, Huynh M, Bartlett KH, Fyfe M, Macdougall L, Boekhout

T, Kwon-Chung KJ, Meyer W. 2004. A rare genotype of *Cryptococcus gattii* caused the

cryptococcosis outbreak on Vancouver Island (British Columbia, Canada). Proc Natl Acad

Sci U S A 101:17258-63.

39. NPR-News. 2010. Fungal disease spreads through Pacific Northwest;

http://www.npr.org/templates/story/story.php?storyId=126198896.

40. Alangaden GJ. 2011. Nosocomial fungal infections: epidemiology, infection control, and

prevention. Infect Dis Clin North Am 25:201-25.

41.   Neofytos D, Horn D, Anaissie E, Steinbach W, Olyaei A, Fishman J, Pfaller M, Chang C,

727   Webster K, Marr K. 2009. Epidemiology and outcome of invasive fungal infection in adult

728   hematopoietic stem cell transplant recipients: analysis of Multicenter Prospective

729   Antifungal Therapy (PATH) Alliance registry. Clin Infect Dis 48:265-73.

730   42.   Achterman RR, Smith AR, Oliver BG, White TC. 2011. Sequenced dermatophyte strains:

731   growth rate, conidiation, drug susceptibilities, and virulence in an invertebrate model.

732   Fungal Genet Biol 48:335-41.

733   43.   Butler MS. 2008. Natural products to drugs: natural product-derived compounds in clinical

734   trials. Nat Prod Rep 25:475-516.

735   44.   Jansen BJ, de Groot A. 2004. Occurrence, biological activity and synthesis of drimane

736   sesquiterpenoids. Nat Prod Rep 21:449-77.

737   45.   Toyota M, Ooiso Y, Kusuyama T, Asakawa Y. 1994. Drimane-type sesquiterpenoids from

738   the liverwort *Diplophyllum serrulatum.* Phytochemistry 35:1263-1265.

739   46.   Robles-Kelly C, Rubio J, Thomas M, Sedan C, Martinez R, Olea AF, Carrasco H, Taborga

740   L, Silva-Moreno E. 2017. Effect of drimenol and synthetic derivatives on growth and

741   germination of *Botrytis cinerea*: Evaluation of possible mechanism of action. Pestic

742   Biochem Physiol 141:50-56.

743   47.   Giaever G, Shoemaker DD, Jones TW, Liang H, Winzeler EA, Astromoff A, Davis RW.

744   1999. Genomic profiling of drug sensitivities via induced haploinsufficiency. Nat Genet

745   21:278-83.

746   48.   Kubo I, Fujita K, Lee SH. 2001. Antifungal mechanism of polygodial. J Agric Food Chem

747   49:1607-11.

49. Fujita K, Kubo I. 2005. Multifunctional action of antifungal polygodial against

Saccharomyces cerevisiae: involvement of pyrrole formation on cell surface in antifungal

action. Bioorg Med Chem 13:6742-7.

50. Veerapandian R, Vediyappan G. 2019. Gymnemic acids inhibit adhesive nanofibrillar

mediated *Streptococcus gordonii-Candida albicans* mono-species and dual-species

biofilms. Front Microbiol 10: 2328.

51. Mor V, Rella A, Farnoud AM, Singh A, Munshi M, Bryan A, Naseem S, Konopka JB,

Ojima I, Bullesbach E, Ashbaugh A, Linke MJ, Cushion M, Collins M, Ananthula HK,

Sallans L, Desai PB, Wiederhold NP, Fothergill AW, Kirkpatrick WR, Patterson T, Wong

LH, Sinha S, Giaever G, Nislow C, Flaherty P, Pan X, Cesar GV, de Melo Tavares P,

Frases S, Miranda K, Rodrigues ML, Luberto C, Nimrichter L, Del Poeta M. 2015.

Identification of a New Class of Antifungals Targeting the Synthesis of Fungal

Sphingolipids. MBio 6:e00647.

52. Proctor M, Urbanus ML, Fung EL, Jaramillo DF, Davis RW, Nislow C, Giaever G. 2011.

The automated cell: compound and environment screening system (ACCESS) for

chemogenomic screening. Methods Mol Biol 759:239-69.