Interaction of contexts in context-dependent orientation estimation

Ron Dekel[1] and Dov Sagi[1,*]

Department of Neurobiology, The Weizmann Institute of Science, Rehovot 7610001, Israel.

*Corresponding author. Email: Dov.Sagi@Weizmann.ac.il

Abstract

The processing of a visual stimulus is known to be influenced by the statistics in recent visual history and by the stimulus' visual surround. Such contextual influences lead to perceptually salient phenomena, such as the tilt aftereffect and the tilt illusion. Despite much research on the influence of an isolated

10   context, it is not clear how multiple, possibly competing sources of contextual influence interact. Here, using psychophysical methods, we compared the combined influence of multiple contexts to the sum of the isolated context influences. The results showed large deviations from linear additivity for adjacent or overlapping contexts, and remarkably, clear additivity when the contexts were sufficiently segregated. Specifically, for adjacent or overlapping contexts, the combined effect was often lower than the sum of

15   the isolated component effects (sub-additivity), or was more influenced by one component than another (selection). For contexts that were segregated in time (600 ms), the combined effect measured the exact sum of the isolated component effects (in degrees of bias). These results imply that the "encoding" of the context is a compressive non-linear transformation, followed by a non-linear selection, manifested by strong interactions between the encoded components. For selected encoded representations, the

20   contextual influence is a linear sum of the individual effects, as predicted from a theory of optimally combining reference cues.

Keywords

Visual adaptation, Visual aftereffects, Tilt Illusion, Orientation processing, Natural vision, Spatial

25   integration, Cue combination

Highlights

- Non-linear sub-additivity for increased context contrast or area
- Non-linear selection between overlapping or adjacent, dissimilar contexts
- Linear additivity for combinations of temporally segregated contexts
- This suggests non-linear encoding and selection, but a linear influence of the context

Introduction

Visual processing is continuously adjusted based on previous visual stimulations, a process termed visual adaptation (Clifford et al., 2007; Kohn, 2007). Similarly, the processing of a visual region can be influenced by the surrounding visual stimulations (Clifford, 2014; Lamme, 1995). Changes in processing due to context are potentially useful, allowing for adjusting the dynamic range, calibrating transmitted information, and enhancing the saliency of novelty. Correspondingly, context, both temporal and spatial, gives rise to salient perceptual effects, often referred to as visual illusions (Clifford & Rhodes, 2005; Gibson, 1937; Gibson & Radner, 1937; Webster, 2011).

In the case of orientation features (Fig. 1), the influence of spatial context can be measured by the tilt illusion (TI), and the influence of temporal context can be measured by the tilt aftereffect (TAE), with clear qualitative similarities between space and time (Schwartz, Hsu, & Dayan, 2007). In both cases, the oriented context leads to a change (bias) in perception, such that the perceived orientation is shifted in the opposite direction from the context orientation (though for very large orientation differences between context and target, the direction of the effect is inverted, which is known as the "indirect effect", Clifford, 2014; Schwartz et al., 2007).

Traditionally, these context effects were thought to be mediated by shifts of internal "norms" for vertical and horizontal orientations (Gibson & Radner, 1937), or, according to later theories, by early visual adaptation (in time) and lateral inhibition (in space) of orientation-selective units (Blakemore & Campbell, 1969; Blakemore, Carpenter, & Georgeson, 1971; Coltheart, 1971). However, more recent evidence implies the involvement of more complex mechanisms, such as learning of statistical regularities (Nakashima & Sugita, 2014; Pinchuk-Yacobi, Dekel, & Sagi, 2016), task experience (Dong, Gao, Lv, & Bao, 2016; Pinchuk-Yacobi, Harris, & Sagi, 2016; Yehezkel, Sagi, Sterkin, Belkin, & Polat,

2010), and perceptual grouping (He, Kersten, & Fang, 2012; Suzuki, Clifford, & Rhodes, 2005). Such observations are important because natural vision is clearly more complex than the testing conditions of most laboratory experiments, raising the question of how findings about context obtained with reduced stimuli can be extended to actual natural stimulation (Snow, Coen-Cagli, & Schwartz, 2017; Solomon & Kohn, 2014). Specifically, most research about the TAE and TI is performed using simplified stimuli, such as lines (Gibson, 1937; Gibson & Radner, 1937), gratings (Campbell & Maffei, 1971; Mitchell & Muir, 1976), or Gabor patches (Knapen, Rolfs, Wexler, & Cavanagh, 2010). However, natural stimulation has high-order statistical regularities in space and time, leading to differences in the operation of the visual system for simplified vs. natural input (Alam, Vilankar, Field, & Chandler, 2014; Carandini et al., 2005; Movshon & Simoncelli, 2014; Olshausen & Field, 2005). It is therefore very important to understand how contextual influence differs between ecological vision and the laboratory (Coen-Cagli, Kohn, & Schwartz, 2015; Snow et al., 2017; Solomon & Kohn, 2014).

Recently, we used unmodified natural images that were selected to match the oriented low-level properties of the reduced stimulation, and found TAE due to image exposure (Dekel & Sagi, 2015; see also, Goddard, Clifford, & Solomon, 2008; Ismail, Solomon, Hansard, & Mareschal, 2016). This finding is consistent with a simple view of contextual influence: that it follows the sum over all the contrast in the image, regardless of a higher-order structure. Here, to address this simplified hypothesis in a systematic fashion, we considered different ways of combining contexts (in space/time, overlapping/non-overlapping), and compared the bias from a combined context, to the sum of the biases from the isolated components (additivity). This is a crucial first step towards more complex combinations, and natural vision.

Specifically, we performed the following experiments:

- First, we investigated the additivity of TAE for combinations of contexts presented at different times (segregated). The results showed additivity, under both fixation and periphery conditions. This is the main finding of this work. To account for the additivity, we offer an explanation in terms of an optimal combination of independent reference cues (see Fig. 3 and the Discussion).
- Next, we investigated the additivity of combinations of contexts presented at the same time. We mostly used TI, because TAE magnitudes for non-retinotopic combinations were too small (data not shown). The results generally showed sub-additivity (Fig. 4), and interestingly, suggest spatial

anisotropy of the sub-additivity. In addition, we considered experiments with different adaptor

85   contrasts, which showed sub-linearity of the TAE vs. contrast function (Fig. 5). This finding rules

out the possibility that the additivity of segregated contexts is due to measuring a TAE with a small

magnitude. To interpret findings of sub-additivity, we considered compression in the early visual

encoding (see the Discussion).

- Finally, we investigated combinations of dissimilar contexts. Such experiments are important,

90   because natural stimuli are composed of dissimilar orientations. We considered combinations that

are adjacent in space (Fig. 6), overlapping (i.e., plaids, Fig. 7), or that use natural images (Fig. 8).

The results generally showed sub-additivity, showing a strong dependence on the spatial

arrangement of the components. To interpret these findings, we considered selection between

competing encoded alternatives (see the Discussion).

95   Overall, the results suggest that in the absence of interactions during encoding or selection, context-

dependent biases are approximately additive (see the Discussion).

Fig. 1 – Tilt illusion and the tilt aftereffect. *(**A**) In the tilt illusion (TI), an oriented surround (here, an*
100 *annulus) leads to a change in the perceived orientation of a target (here, the center circle). The target*
*without surround, shown on the left, is provided as a reference and was not used in the experiments. (**B**)*
In the tilt aftereffect (TAE), exposure to an oriented adaptor leads to a change in the perceived
orientation of a subsequently viewed target. Figure reproduced from (Dekel & Sagi, 2019).

Methods

Observers

The experiments reported here were performed in a laboratory or with a web-based interface through the Amazon Mechanical Turk (MTurk). In the laboratory experiments, $N = 23$ observers (20 females, 3 males, aged 22-40) with normal or corrected-to-normal vision participated, and provided their written informed consent. In the MTurk experiments, $N = 270$ observers participated (reported age of 38 ± 10, Mean $\pm$ STD, median of 35, in the range 20-70; gender information was usually not collected). The web-based study was reviewed by the Institutional Review Board (IRB) of the Weizmann Institute of Science and deemed exempt from the collection of informed consent forms. The typical selection criteria for MTurk observers were as follows: 'PercentAssignmentsApproved' $\geq$ 99, 'NumberHITsApproved' $\geq$ 2000, 'LocaleCountry' = 'US'. All observers were naïve to the purpose of the experiments. The work was carried out in accordance with the Code of Ethics of the World Medical Association (Declaration of Helsinki), and was approved by the Institutional Review Board (IRB) of the Weizmann Institute of Science.

Apparatus

Laboratory setup. The stimuli were presented using dedicated software on a 22" HP p1230 monitor operating at 85Hz with a resolution of 1600x1200 that was gamma-corrected (linearized). The mean luminance of the display was 26.06 cd/m^2 (TAE experiments, excluding the conditions with image adaptors) or 48.96 cd/m^2 (TI experiments, and TAE experiment with image adaptors), in an otherwise dark environment. The monitor was viewed at a distance of 100 cm.

Amazon Mechanical Turk setup. To display time-accurate full-screen stimuli in arbitrary MTurk worker machines, we developed a dedicated web-browser client. The client was implemented in Javascript using WebGL, and achieved approximately single-frame accuracy on standard machines that run the Chrome, Mozilla, or Opera browsers and have a GPU-enabled WebGL implementation (this hardware support is indicated by the 'failIfMajorPerformanceCaveat' Javascript API flag). In addition to the browser and WebGL requirements, clients were required to use a PC rather than a tablet or a smartphone (as indicated by standard API flags), and the client machine had to pass a simple performance test (intended to ensure that the machine can achieve the required time accuracy). During an experiment, timing information estimations were collected (obtained using the 'requestAnimationFrame' callback). Timing diagnostics showed that most trials, of most observers, measured nearly perfect single-frame

accuracy. The worst-case 95% presentation duration error (compared to the observers' average) was less than 20% (e.g., 100 ms ± 20 ms) for practically all observers (99%). Similarly, inter-observer differences were negligible (SD of ~5 ms). As such, we deemed the timing accuracy of the setup to be more than sufficient for the purpose of the current study. The display luminance, gamma calibration, sitting distance, and environmental lightning conditions were only controlled to the degree reflected in the task performance (see below). We note that all statistical comparisons we report are within-observer (paired), hence reflect measurements from the same display. Moreover, in all comparable experiments, we found the crowdsourcing setup to be nearly as reliable as that of the lab.

Stimuli and tasks

All stimuli were presented using dedicated software on a uniform gray background. To begin stimulus presentation in a trial, observers pressed the spacebar (self-initiated trials). In the laboratory, observers were instructed to fixate on the center of the display before stimulus initiation. Responses were provided using the left and right arrow keys. We measured the change in the perceived orientation of a target stimulus following exposure to an adapting oriented stimulus (TAE) or due to a surrounding oriented stimulus (TI). In MTurk, the stimuli, defined in pixels, were resized using linear interpolation to maintain a fixed proportion relative to the full-screen monitor resolution. To minimize resizing artifacts, the resizing multiplier was rounded to the nearest lower multiple of 0.25 (e.g., resizing by a multiplier of 1.25 instead of 1.33). Under all TAE conditions (and separately under all TI conditions), the stimuli used to measure the perceived orientation (targets) were nearly identical.

1- and 2-back experiments (TAE). The following presentation sequence was used (Fig. 3A): a blank screen (600 ms), 2-back Gabor adaptor (oriented -20° or +20° to vertical, 50 ms), a blank screen (600 ms), 1-back Gabor adaptor (±20°, 50 ms), a black screen (600 ms), and a near-vertical Gabor "test" (50 ms). Under conditions with additional adaptor presentations (see Table 1, where "AAAAT" trials had four adaptors, and "AAAT" trials had three), the additional presentations were added in a similar way (600 ms padding of blank screen). All presentations were randomized independently (location and orientation). Gabor patches were 50% Michelson contrast with Gaussian envelope σ = 0.42°, spatial frequency λ = 0.3°, and random phase. Two versions of the experiment were run: fixation and periphery. In fixation, adaptors and targets were presented at the fixated center of the display, and targets were oriented -9° to +9° (in steps of 1°). In the periphery, adaptors and targets were presented either at the left or right of the fixation (at ±1.8°). The target was presented either at the same side as the adaptors

(retinotopic) or at the opposite side (non-retinotopic), randomly. Targets were oriented from -12° to +12° (in steps of 2°). Observers were instructed to inspect the adaptors and the target presentations, and then to report whether the orientation of the near-vertical target was tilted clockwise or counter-clockwise relative to vertical (no feedback, except during a brief pre-experiment practice). Responses were provided using the left and right arrow keys. Four peripheral crosses co-appeared with the target to facilitate discrimination between the adaptor and the target.

Note that our analyses were only for the 1- and 2-back presentations, whereas the design of some conditions permitted further conditioning of the analysis to the 3- and 4-back presentations. These data were not shown, because insufficient trials per condition due to a combinatorial explosion, which prevented drawing meaningful conclusions from the analysis. In addition, the effect from the 3- and 4-back adaptors was weaker and less consistent than that observed for the 1- and 2-back adapters.

Img+Phs experiments (TAE). In these experiments, we measured the TAE caused by an adaptor presentation using a subsequently presented Gabor target (Fig. 8).

'Biased' image adaptors were unmodified natural images depicting animals and plants that were selected to maximize the orientation content around a particular combination of orientation and spatial frequency. The images were obtained using a procedure similar to that of (Dekel & Sagi, 2015). Briefly, sub-images of 256×256 pixel size were cropped from ImageNet images (Deng et al., 2009), were windowed using a circle with smoothed edges, and normalized to have an RMS contrast of 23% and an average pixel value of 127 (i.e., background). Importantly, the sub-images were selected based on the maximization of the response of an orientation filter at a specific frequency band, leading to a "biased" set of unmodified images having similarly oriented content. The orientation filter had an orientation profile of a Gaussian around the orientation 25° counter-clockwise to vertical, with a standard deviation of 15°. The frequency band had a frequency profile of a second-order Butterworth, with half-responses at 6.75 and 33.75 cycles/image. Manual pruning was used to ensure that only intelligible images of animals and plants were used. Here, the process resulted in a pool of 47 animal and 21 no-animal images (out of ~6 million sub-image candidates, cropped with overlapping from 279,827 ImageNet images). By mirroring images, the orientation of the content could be switched from +25° to -25°. During a session, presentations of images of both orientations were randomly mixed (+25° and -25°), and each image could be presented multiple times, but always for the same orientation (randomized between observers by using a pool of 8 random assignments).

Phase scramble adaptors were obtained by applying the following procedure to a 'biased' image:

applying a circular window, randomizing the phase content of the image, applying the circular window again, and then normalizing the image to have an RMS contrast of 23% and an average pixel value of 127. With color images, each RGB color channel was scrambled separately (in the standard RGB color space, which appeared to produce fewer artifacts than scrambling at an HSV or HSL color space).

Mixture adaptors were a mixture between a biased image and a phase scramble of a biased image, each at half amplitude, and each corresponding to an opposite orientation content (biased +25° and scrambled -25°, or vice versa; note that at MTurk, the contrast remained identical for both components, even if it was not at 50%).

Gabor targets were Gabor patches, oriented from -6° to +6° (in steps of 0.75°), with a random phase. In the laboratory setting, Gabor targets were 50% Michelson contrast with $\sigma = 0.47°$ and $\lambda = 0.33°$. In the MTurk setting, Gabor targets had an amplitude of 64 gray levels (out of 256, i.e., corresponding to a contrast of 50% in a linearized display), $\sigma = \sqrt{2}\lambda$, and $\lambda = \sim2\%$ of screen height.

The trial structure used the following presentation sequence: a blank screen (350 ms), adaptor presentation (content oriented -25° or +25° to vertical; 300 or 100 ms; equal width and height, height in lab: 5.64°, height in MTurk: approx. 30% of the total monitor height), a blank screen (600 ms), and a near-vertical Gabor "target" (50 ms). A double-task was used. In the first reply, observers reported the orientation of the target (presented second). In the second reply, observers either reported the content of the image adaptor (animal/no-animal, each 50% of trials; the task showed ~90% and ~78% accuracy for natural and mixture adaptors, respectively), or reported whether the adaptor appeared or not (adaptor-present in 75% of trials; when the adaptor was absent, a blank screen was presented instead; the task showed ~97% accuracy). There was no feedback in any of the tasks, except during a brief pre-experiment practice.

Contrast experiments (TAE). The contrast experiments (Fig. 5) had a Gabor adaptor version and an image adaptor version. In the Gabor adaptor version, the structure of trials in the "1- and 2-back" fixation experiments was used, without the 2-back presentation and its preceding ISI, and with varying Michelson contrast of the 1-back adaptor (11%, 33%, and 100%). In the image adaptor version, the structure of trials from the "Img+Phs" experiment was used, with only biased images (no phase scrambles), with detection as the second task in the double-task design, and with varying contrasts of the image adaptor (10%, 30%, 100%, and technically also 0% for the detection task).

TI experiments. Stimuli (e.g., Fig. 1A right) were presented at the center of the display, and consisted of a sine-wave circular "target" (oriented near-vertical from -9° to +9° in steps of 1°), a full, partial, or empty "near-surround" sine-wave annulus (oriented -20°, +20°, -90°, or 0°, or combinations), and a full or empty "far-surround" sine-wave annulus (oriented -20°, +20°, -90°, or 0°). A partial near-surround was either a single half (up or down), a single quadrant (left, right, up, or down), two quadrants (left and right, or up and down), or four quadrants (left, right, up, and down with smoothed edges). Under laboratory conditions (MTurk conditions), the following sizes were used: target circle radius = 0.6° (~3.3% of screen height), gap between target and near surround = 0.15° (~1%), near surround radius = 1.2° (~6.6%), gap between near and far surrounds = 0.15° (~1%), far surround radius = 3° (~16.6%), and spatial frequency $\lambda = 0.3°$ (~1.6%) for all sine-wave gratings. The phase was separately randomized for each stimulus component (target, near surround, quadrants of the near surround, and far surround). The sine-wave gratings had a contrast of 100%, except in the experiments with overlapping combinations (2nd-order stimuli, Fig. 7), where the "A", "H", and "V" components (-20°/+20°, -90°, and 0° orientation, respectively) each had a Michelson contrast of 50% in the lab, and an amplitude of 64 gray levels in MTurk (out of 256, i.e., corresponding to a contrast of 50% in a linearized display; note that contrast remained identical for both components even if it was not at 50%). In the experiments, observers were instructed to ignore the surrounding annuli, inspect the target (central circle), and report whether the target is tilted clockwise or counter-clockwise relative to vertical (no feedback, except during a brief pre-experiment practice). The stimuli were presented starting from 350 ms after the trial initiation, or, under a few conditions, randomly 350 to 550 ms after the trial initiation ("onset jitter", see Table 1; this design was intended to enable a reaction time-based analysis, as in Dekel & Sagi, 2019). The stimulus was presented for 100 or 400 ms.

Procedure

Most laboratory observers participated in multiple experiments in this study, with at least three days break between sessions (within and between experiments). All MTurk observers participated in a single experiment, for a single daily session (as indicated by having a unique 'Worker ID'). In all experiments, we required a low rate of finger errors, achieved by persistent coaxing of laboratory observers, and pruning of MTurk observers (see below). Each daily session was preceded by a brief practice block with easy stimuli (this practice was repeated until close-to-perfect accuracy was achieved, manually in the laboratory, and automatically in the MTurk experiments). Each daily session consisted of either four or five blocks, with breaks lasting at least two minutes between blocks. The number of trials per block was

255 determined such that the block duration was approximately five minutes (fewer trials per block when the trial duration was longer, see Table 1). When different trial types were mixed in a block, the number of trials per type was approximately equal, with the exception of the "quadrants" experiments where 2/3 of the trials were used for the single quadrant types, and 1/3 of the trials were used for the two or four quadrant types. The intended number of observers per condition was ~10 in the lab, and ~15 in the

260 MTurk, which seems more than sufficient for the type of effects considered here. Under a few conditions, we doubled the number of observers following the data analysis, to ensure the validity of the found results.

Analysis

Fitting perceived orientation. The magnitude of TAE and TI was calculated based on the reported

265 orientation (clockwise vs. counterclockwise) of the near-vertical target (the Gabor patch for TAE, and central sine-wave circle for TI). First, for each context condition, a cumulative normal distribution (with lapse rates) was fitted to the psychometric function of %clockwise reports as a function of target orientation. Then, using of the fitted function, the perceived vertical orientation (PV) of the target was calculated as the interpolated orientation having an equal probability for clockwise and counter-

270 clockwise reports (50%). This was achieved with Psignifit 3.0 software (Fründ, Haenel, & Wichmann, 2011). Then, the TAE and TI magnitudes were calculated as half the shift in PV between two opposing adaptor or surround orientations (e.g., +20° vs. -20°). When combining measurements across experimental days or spatial positions (left and right, or up and down), the above calculation was performed separately for each measurement, which was then averaged.

275 Observer and trial pruning. For the MTurk observers, the sessions were pruned, based on the parameter thresholds described in Table 2; the counts of pruned observers are presented in Table 1. It can be seen that the thresholds used were relatively lenient, and nearly none of the cooperative observers who performed the task(s) as instructed should have been pruned. Importantly, most observers were either similar to the laboratory observers in terms of measured performance, or nearly at chance level

280 performance, so pruning, which aimed to disqualify the uncooperative observers, was usually mostly straightforward. In addition, for both MTurk and the laboratory data, trials were pruned if the orientation discrimination task RT was faster than 300 ms, or slower than 10 seconds (see Table 1 for the counts of pruned trials).

Statistics. All relevant statistical tests were two-tailed. All *t*-tests were paired (i.e., repeated-measures).

285 **Table 1 – Procedural information.**

	#Obs. used	#Obs. pruned	Sessions per obs.	Blocks per session	Trials per block	Total used trials	Total pruned trials	Fig.
1- and 2-back fixation	9	0	1-2	5	93	4,633	17	3B
1- and 2-back periphery (AAAAT)	5	0	5-10	5	65	13,271	39	3C
1- and 2-back periphery (AT/AAT/AAAT/AAAAT)	6	1	7-9	5	~95[**]	15,608	15	3C
Area near/far 400ms	21[*]	7	1	4	228	19,119	33	4A
Area up/down 100ms	19[*]	7	1	4	228	17,293	35	4B
Area up/down 400ms	26[*]	3	1	4	228	23,670	42	4B
Area quadrants Q1/Q2 100ms	20[*]	5	1	4	228	18,220	20	4C
Area quadrants Q1/Q2 100ms, onset jitter	13[*]	7	1	4	228	11,806	50	4C
Area quadrants Q1/Q4 100ms	10[*]	5	1	4	228	9,118	2	4C
Area quadrants Q1/Q4 400ms	17[*]	5	1	4	228	15,489	15	4C
Contrast Gabor	9	0	1	5	125	5,560	65	5
Contrast 'biased' images	10	0	1	4	102	4,077	3	5
Far=V/H, 100ms	9[*]	6	1	4	190[***]	5,444	28	6
Far=V/H, 400ms	14[*]	14	1	4	190[***]	8,510	2	6
Far=H/V/E, 100 ms	26[*]	6	1	4	228	23,629	83	6
Far=H/V/E, 400 ms, onset jitter	15[*]	1	1	4	228	13,655	25	6
Near=45°, Far=V/H, 400ms	12[*]	11	1	4	190[***]	7,279	17	6
2nd order 400ms	10	0	1	4	228	9,119	1	7
2nd order 400ms	7[*]	2	1	4	228	6,384	0	7
2nd order 100ms	12[*]	2	1	4	228	10,896	48	7
Img+Phs common	9[*]	3	1	4	102	3,635	37	8
Img+Phs laboratory	10	0	1	4	102	4,079	1	8
Img+Phs color	10[*]	0	1	4	102	4,072	8	8
Img+Phs 100ms	11[*]	2	1	4	102	4,484	4	8
Img+Phs detection task	19[*]	6	1	4	102	7,749	3	8

(*) MTurk observers.
(**) In the "AT/AAT/AAAT/AAAAT" experiment, ~25% of the trials were "AT", i.e., adaptor-test trials lacking a 2-back adaptor, and were ignored in the analysis.
(***) In the "Far=V/H" experiments, 20% of the trials had an orientation of 0° or 90° (V or H) in both near and far surround annuli, and were ignored in the analysis.

Table 2 – Observer pruning for MTurk.

MTurk observer pruning	"Img+Phs" Orientation disc. (first task)	"Img+Phs" Recognition task (second task)	"Img+Phs" Detection task (second task)	TI experiments Orientation disc. (single task)
Mean RT	>430 ms	>650 ms	>650 ms	>450 ms
Mean RT in the fastest block	>430 ms	>450 ms	>450 ms	>450 ms
Mean accuracy	>65%	>60%	>90%	>65%
Mean accuracy in the worst block	>65%	>60%	>85%	>65%
Reduction in the mean accuracy from the best to the worst block	<0.2	<0.2	<0.2	<0.2
Reply heterogeneity	<12% with Det. <20% with Rec.	-	-	<17% for Q1/Q2 <25% otherwise

RT in the second task refers to the combined RT of both the first and second tasks. Accuracy is the percentage of trials that measured a correct reply, excluding the impossible trials (the target oriented 0° or content-recognition for "Phs"). The chance level is 50% in all tasks except the detection task, where it is 75%. Reply heterogeneity is the mean probability that the observer provided a different reply for identical stimuli, where the identity requires all the following: the same target orientation, the same context type (e.g., up vs. down), and the same context orientation(s). Specifically, given a reply probability, p, measured for a set of identical stimuli, of size n, the reply heterogeneity was 2p(1-p), averaged across all sets of identical stimuli. The threshold for heterogeneity that appears in the table was determined by the number of identical stimuli, n, where n = 3 for "Img+Phs" with the detection task, n = 4 for "Img+Phs" with the recognition task, and n = 8 for all TI experiments with the exception of the Q1 and Q1 quadrant trial types for which n = 4 (see Table 1). The heterogeneity heuristic was typically more sensitive than the performance accuracy for identifying uncooperative observers.

Modeling

Additive model.

In the TAE experiment, we found additivity (see the Results). To model this, we describe each adaptor's influence by a scalar indicating the weight of its influence; the sign of the weight is the sign of the adaptor (clockwise, counterclockwise, and absent, indicated by +, -, and 0, respectively), and the predicted PV is the sum. This can be written as:

$$\text{PV}_{\text{fixation}} = o_0 + w_1 o_1 + w_2 o_2 \tag{1}$$

$$\text{PV}_{\text{periphery}} = o_0 + w_1^R o_1^R + w_2^R o_2^R + w_1^N o_1^N + w_2^N o_2^N, \tag{2}$$

where PV is the perceived orientation, o_0 is the reference (unadapted) orientation, o_i is the i-back adaptor orientation (R and N indicate the retinotopic vs. non-retinotopic adaptor, respectively), and w_i is the weight of the i-back adaptor.

Cue combination model.

To interpret biases in the perceived orientation, we considered the idea that a reference frame is updated based on using the contextual orientations as cues. We modelled this idea as a Bayesian cue combination of independent Gaussian cues (Oruç, Maloney, & Landy, 2003):

$$\tilde{\theta} = \frac{\sum_i \frac{1}{\sigma_i^2} \theta_i}{\sum_i \frac{1}{\sigma_i^2}}, \tag{3}$$

where θ_i and σ_i^2 are the means and variances of the Gaussians, and $\tilde{\theta}$ is the posterior mean. The quantity $\frac{1}{\sigma_i^2}$ is often referred to as *reliability*. The model described each adaptor, as well as the unadapted vertical reference, act as independent Gaussian cues (Fig. 2). Specifically, the unadapted reference cue had an unknown orientation, θ_0, each adaptor cue had a known orientation, θ_i, and the relative reliability between each adaptor and the unadapted reference, $\frac{\sigma_0^2}{\sigma_i^2}$, was also unknown.

To model the fixation experiment using a cue combination, let θ_0 be the initial (unadapted) internal reference orientation, θ_1 be the 1-back adaptor orientation, and θ_2 be the 2-back adaptor orientation. The updated internal reference, inferred using Bayes' rule and by assuming that the cues are Gaussian with variances σ_i^2, is (see Eq. (3)):

$$\tilde{\theta} = \frac{\frac{1}{\sigma_0^2}\theta_0 + \frac{1}{\sigma_1^2}\theta_1 + \frac{1}{\sigma_2^2}\theta_2}{\frac{1}{\sigma_0^2} + \frac{1}{\sigma_1^2} + \frac{1}{\sigma_2^2}} \tag{4}$$

The predicted vertical orientation is minus (opposite) the change in the internal reference, which is,

335 $PV = -\tilde{\theta}$. Rearranging, we get:

$$PV = -\frac{\theta_0 + \frac{\sigma_0^2}{\sigma_1^2}\theta_1 + \frac{\sigma_0^2}{\sigma_2^2}\theta_2}{1 + \frac{\sigma_0^2}{\sigma_1^2} + \frac{\sigma_0^2}{\sigma_2^2}} \tag{5}$$

When an i-back adaptor is missing, we can set $\sigma_i = \infty$, which removes the relevant terms from the equation. Of the model parameters, the adaptor orientations are known and equal to $\pm 20°$; therefore, three unknown parameters remain: θ_0 (the initial internal reference), $\frac{\sigma_0^2}{\sigma_1^2}$, and $\frac{\sigma_0^2}{\sigma_2^2}$.

340 Similarly, the equation for the periphery experiment can be obtained (five unknown parameters). Note that in both cases, the number of parameters is the same as in an additive model (Eqs. (1) and (2)). When fitting behavior to the model, nearly identical predictions were found when fitting the average observer, compared to when fitting each observer separately, and then averaging. We therefore used the latter, which is the stricter alternative in the sense that it can be more non-additive.

345 Of note, based on the above cue-combination model equations, the TAE, which is the change in the perceived orientation, can be derived for the case of a single adaptor as follows (TI is calculated similarly):

$$TAE = -\frac{\theta_0 + \frac{\sigma_0^2}{\sigma_1^2}\theta_1}{1 + \frac{\sigma_0^2}{\sigma_1^2}} - (-\theta_0) = \frac{\frac{\sigma_0^2}{\sigma_1^2}\theta_0 - \theta_1}{\frac{\sigma_1^2}{\sigma_0^2} + 1} \tag{6}$$

Fig. 2 – Illustration of the cue combination. *The reference (i.e., posterior) is shifted in the direction of the adaptor cues, leading to a shift of the perceived orientation in the opposite direction.*

Results

Linear additivity for temporally segregated contexts

First, we considered the change in the perceived orientation caused by temporal combinations of spatially isolated Gabor patches. Specifically, we considered two possible presentation times, 650 and 1300 ms, before a target, which we referred to as 1-back and 2-back, respectively. In each time frame, the fixated center of the display presented a Gabor tilted +20° clockwise to vertical, -20° to vertical, or was empty (Fig. 3A). We measured the change in the perceived vertical orientation of the target stimulus due to the adaptor(s)' exposure (tilt aftereffect, TAE), and checked whether the combined influence of both adaptors can be explained as a linear sum of the isolated adaptor influences. The results showed standard TAE magnitudes of around ~1° (Fig. 3D), and importantly, the additivity of the 1- and 2-back adaptation intervals (Fig. 3B). We analyzed this using a 3×3 repeated measures ANOVA of {-20°,+20°,empty}×{-20°,+20°,empty}, which showed the significant influence of the 1- and the 2-back presentations, and importantly, no interaction, indicating additivity (1-back: $F_{(2,16)} = 26.99$, $p < 0.0001$; 2-back: $F_{(2,16)} = 20.25$, $p < 0.0001$, interaction: $F_{(4,32)} = 1.08$, $p = 0.38$, $N = 9$ observers, Fig. 3B). To strengthen this result, we also considered a linear mixed-effects regression analysis, which showed the same findings (1-back: $t_{(77)} = 8.81$, $p < 0.0001$; 2-back: $t_{(77)} = 4.64$, $p < 0.0001$; interaction: $t_{(77)} = -0.05$, $p = 0.95$).

Similarly, using presentations in the near-periphery (±2° of visual angle, Fig. 3A), we considered the 1- and 2-back adaptors, each presented at the same side as the target (retinotopic, "R") or at the contralateral side (non-retinotopic, "N"). We analyzed the data from two experiments, with four adaptor presentations before the target (AAAAT, $N = 5$ observers, $n = 13{,}271$ trials), or with up to four adaptor presentations before the target (AT, AAT, AAAT, or AAAAT, $N = 6$ observers, $n = 15{,}608$ trials). The results showed retinotopic TAEs of about 1°, non-retinotopic TAEs of about 0.25° (Fig. 3D), and importantly, additivity of the 1- and 2-back adaptors (Fig. 3C). Specifically, we applied a 4×4 repeated measures ANOVA of {R-20°,N-20°,N+20°,R+20°}×{R-20°,N-20°,N+20°,R+20°}, which showed significant 1- and 2-back adaptor influences, but non-significant interaction, for "AAAAT" (1-back: $F_{(3,12)} = 28.71$, $p < 0.0001$, 2-back: $F_{(3,12)} = 23.98$, $p < 0.0001$, interaction: $F_{(9,36)} = 0.41$, $p = 0.92$), and similarly for "AT, AAT, AAAT, AAAAT" (1-back: $F_{(3,15)} = 3.38$, $p = 0.05$, 2-back: $F_{(3,15)} = 7.25$, $p = 0.003$, interaction: $F_{(9,45)} = 0.25$, $p = 0.98$). We therefore merged the data of the two experiments (1-back: $F_{(3,30)} = 14.96$, $p < 0.0001$, 2-back: $F_{(3,30)} = 22.81$, $p < 0.0001$, interaction: $F_{(9,90)} = 0.19$, $p = 0.99$;

$N = 11$ total observers from both experiments, Fig. 3C). Therefore, the results showed an additivity of
peripheral adaptation, and moreover, an additivity of retinotopic with a non-retinotopic adaptation. The
3- and 4-back adapters led to a TAE (~0.25° retinotopically); however, the effect was weaker and less
consistent than that observed for the 1- and 2-back adapters; therefore, analyzing the additivity was not
immediately possible (see the Methods).

Overall, across all conditions, the perceived vertical orientation (PV), following a sequence of
temporally segregated adaptors, can accurately be described as the sum of the isolated adaptor
influences (see the Methods, Eqs. (1) and (2)). That is, the combined effect was well explained as the
sum of the component influences, with non-significant deviations from additivity. One possibility of
interpreting these findings is by using the theory that a bias in the perceived orientation reflects an
update of an internal reference frame by using the contextual orientations as cues (Andrews, 1964;
Dakin & Rosenberg, 2018). For example, an adaptor oriented +20° might shift the vertical reference
from 0° to +1°, in which case a physical 0° orientation will be perceived as -1° (relative to the
reference). We modelled this idea as a Bayesian cue combination of independent Gaussian cues (see the
Methods; Oruç, Maloney, & Landy, 2003). Remarkably, model fitting showed that in the relevant
parameter range ($\frac{\sigma_0^2}{\sigma_1^2} \leq 0.06$), the model is practically equivalent to additivity. That is, the cue-
combination predictions (in orange) were superimposed on the additive predictions (in black, see Fig. 3).
Quantitatively, for both additive and cue-combination models, the resulting R^2 values (the prediction
errors in Fig. 3BC) showed $R^2 = 0.97$ in the fixation experiment, and $R^2 = 0.99$ in the periphery
experiment. Therefore, the independent cue-combination model accounts for the observation of the
additivity of temporally segregated contexts. Note that the cue combination account provides a theory,
whereas additivity is an arbitrary rule (see the Discussion).

Fig. 3 – Additivity in temporal combinations of contexts. *(**A**) Stimuli illustrations (actual stimuli were Gabor patches, see Fig. 1A). (**B and C**) The shown data points are the average across observers of the perceived vertical orientation (PV), measured for the "Target", following exposure to different combinations of 1- and 2-back adaptors. The shown lines are the predictions based on additivity (black, Eqs. (1) and (2), using the averages shown in panel D) or the cue combination (orange, Eq. (3)). (**B**) Fixation experiment. (**C**) Periphery experiment, where each of the 1- and 2-back adaptors was either presented at the same side as the target (retinotopic, "R") or at the contra-lateral side (non-retinotopic, "N"). Note that the 1-back, 2-back, and Target locations were each randomized independently (Left/Right). The results showed linear additivity across all conditions. (**D**) Average TAE magnitudes. Note that magnitudes of TAE and TI, as shown in the subsequent figures, represent the change in PV. Error bars are ±1SEM.*

Sub-additivity for overlapping or adjacent contexts

Here, we consider combinations of similar contexts (the same grating or the same Gabor pattern) that are adjacent or overlapping in space. That is, the similar contexts are presented at the same time, in adjacent or overlaying positions, and tested at the same (TI) or at a later (TAE) time.

First, we consider the additivity of an increase in the context area (Fig. 4). We measured the TI due to a surrounding annulus that is close to the central target ("Near"), far from the central target ("Far"), or when both annuli were presented ("Both") (Fig. 4A). Remarkably, the results showed the same TI magnitude of 2° when the Near annulus was presented either with or without the Far annulus, despite a TI of 1° caused by the Far surround alone. Indeed, comparing "Both" to the sum of "Near" and "Far" showed a large and significant difference indicating a strong sub-additivity ($t_{(20)} = 3.77$, $p = 0.001$, $N = 21$ MTurk observers, using a presentation duration of 400 ms).

Using a similar design, we measured the TI from the upper half ("Up"), lower half ("Down"), or full ("Both") surrounding annulus (Fig. 4B). The results showed for both surround halves a TI of between half to two-thirds of the TI of the full annulus, i.e., close to additivity but with a statistically significant trend toward sub-additivity (comparing "Both" to the sum "Up" + "Down" showed, 100 ms presentation: $p = 0.0002$, $t_{(18)} = 4.69$, $N = 19$ MTurk observers; 400 ms presentation: $p = 0.05$, $t_{(25)} = 2.05$, $N = 26$ MTurk observers; Fig. 4B).

A similar sub-additivity was found when considering different quadrants of the surrounding annulus (Fig. 4C). For example, with a 100 ms presentation duration, the results showed a TI of ~1° when one quadrant was presented ("1Q": left, right, up, or down), a TI of ~1.5° when two quadrants were presented ("2Q": left and right, or up and down), and a TI of only ~2.5° when all four quadrants were presented ("4Q"). That is, a strong sub-additivity is evident by the "4Q" measurement showing much less TI than four times the "1Q" measurement (see Fig. 4C) (100 ms: $t_{(9)} = -3.71$, $p = 0.005$, N=10 MTurk observers; 400 ms: $t_{(16)} = -3.22$, $p = 0.005$, $N = 17$). Moreover, a weak sub-additivity is evident by the "2Q" measurement showing slightly less TI than twice the "1Q" measurement (100 ms presentation duration: $p = 0.01$, $t_{(32)} = -2.66$, $N = 33$ MTurk observers; data were pooled from a condition having an onset jitter, see Table 1; no 400 ms condition). Importantly, comparing the left/right to the up/down measurements showed that the left and right quadrants were nearly additive ("1Q" averaged 0.86°, "2Q" averaged 1.62°, and 2·"1Q" vs. "2Q" showed: $t_{(32)} = -0.72$, $p = 0.9$ corrected),

whereas the up and down quadrants measured a strong sub-additivity ("1Q" averaged 0.99°, "2Q" averaged 1.44°, and 2·"1Q" vs. "2Q" showed: $t_{(32)}$ = -3.53, p=0.003; Bonferroni correction for two multiple comparisons). This anisotropy can possibly be explained by the fact that the sub-additivity is caused by low-level, intra-hemisphere interactions (see the Discussion).

Finally, we considered an increase in an adaptor's contrast (Fig. 5). That is, we measured the TAE at different adaptor contrasts, and compared the result to what is expected from linearity of the TAE vs. contrast function. The results showed a lack of linearity (Fig. 5), consistent with previous reports (Blake, Tadin, Sobel, Raissian, & Chong, 2006). For example, an increase of contrast by a factor of almost 10, from 11% to 100%, led to an increase in TAE by a factor of only ~2.5. More generally, the 100% contrast measurement was lower than the multiplication of a lower-contrast measurement by the factor of the contrast change, for Gabor adaptors (compared to 33% contrast: $t_{(8)}$ = 3.42, p = 0.01; compared to 11% contrast: $t_{(8)}$ = 2.20, p = 0.06; two-tailed repeated measures t-test, not corrected for multiple comparisons; N = 9 observers), and for 'biased' image adaptors (compared to 30% contrast: $t_{(9)}$ = -3.50, $p < 0.01$; compared to 10% contrast: $t_{(9)}$ = -3.46, $p < 0.01$; N = 10 observers, using the 'Detection task' double-task design, see the Methods). Therefore, an increase in contrast led to a sub-linear increase in TAE.

Importantly, the same Gabor stimuli were used in the Gabor contrast experiments and in the temporal segregation experiments (Fig. 3BC and Fig. 5, "Gabor"). Moreover, the magnitudes of TAE were comparable. Therefore, the finding of sub-linearity here rules out the possibility that the additivity of the segregated contexts found above is due to measuring a TAE with a small magnitude. (It is also interesting to note that the manipulation of contrast is, in a sense, the opposite of temporal segregation: sufficiently adjacent presentations, within a period of less than ~50 ms can be assumed to be temporally integrated, and thus to be almost interchangeable with a change in the contrast of a single presentation, see, e.g., Watson & Nachmias, 1977.)

Overall, increasing context strength, either by increasing the contrast or by increasing the area leads to a sub-linear increase in the context influence. The finding in the previous section, of linear additivity for temporally segregated contexts, suggests that the sub-linearity observed here is caused by a non-linear encoding of the context (see the Discussion).

Fig. 4 – Sub-linear additivity for an increased area of surround*. The TI is shown (Fig. 1B), averaged across observers. (**A and B***) TI from component contexts (Near/Far or Up/Down), from the combined context ("Both"), and for the prediction based on additivity ("Sum" of the TI of the two components). (**A***) Near/Far (400 ms presentation duration). (**B***) Up/Down. Sub-additivity is evident by Both<Sum. (**C***) TI from one quadrant ("1Q": left/right/up/down), two quadrants ("2Q": left and right, or up and down), and four quadrants ("4Q"), pooled from all "Quadrants" experiments (Table 1). The continuous line indicates the prediction from the "1Q" measurement based on additivity. The results showed, under all conditions, that an increase in the context area led to a sub-linear increase in the influence of context. The "Near" annulus in "Near/Far" (panel A) has a size identical to the full annulus in the "Up/Down" and the "Quadrants" experiments (panels B and C), and the background was always uniform gray. Error bars are ±1SEM.*

Fig. 5 – Sub-linear influence for increased adaptor contrast. *The average TAE across observers measured when using different adaptor contrasts is shown. TAE was caused by exposure to either a "Gabor" patch or an oriented "Image" adaptor, and measured using a near-vertical Gabor target (comparable to Fig. 3 for the "Gabor" case). The results show the sub-linearity of the TAE vs. contrast function. Error bars are ±1SEM.*

Non-linear competition between adjacent, dissimilar contexts

A possible account of the sub-linearity observed above for overlapping or adjacent contexts is that it reflects self-inhibition (for overlapping contexts) or lateral inhibition (for adjacent contexts). Here, we considered experiments using combinations of dissimilar contexts (e.g., one tilted and one not). These experiments showed a form of competition between contexts that cannot be readily explained by self- or lateral inhibition.

First, we measured how the TI, caused by the orientation of a surrounding annulus ("near-surround", tilted ±20° to vertical), is influenced by a far-surround annulus that is either horizontal or vertical. The results showed about a 30% reduction in TI in the presence of vertical compared with horizontal far surround (Fig. 6) (100 ms presentation duration: $N = 9$ MTurk observers, $t_{(8)} = -4.77$, $p = 0.002$, two-tailed repeated-measures t-tests; 400 ms presentation duration: $N = 14$ MTurk observers, $t_{(13)} = -4.62$, $p = 0.0005$). The observed effect was robust, as evident for all observers except one (22 out of 23 showing V<H), which is especially striking considering the inherent variability of the display parameters in the web-based platform.

Moreover, the interaction with the far surround was selective to the vertical far surround orientation ("H/V/E" conditions, Fig. 6). That is, the TI for absent vs. horizontal far surrounds was almost identical as indicated by the non-significance of the null hypothesis ("No Far" vs. "Far=H", 100 ms: $t_{(25)} = -0.3$, $p = 1$ corrected, $N = 26$ MTurk observers; 400 ms with onset jitter: $t_{(14)} = -1.16$, $p = 0.79$ corrected, $N = 15$ MTurk observers; Bonferroni correction for three multiple comparisons), whereas TI in the presence of a vertical far surround was reduced by about 30% relative to both the absent and horizontal far surrounds ("Far=V" vs. "No Far", 100 ms: $t_{(25)} = -3.32$, $p = 0.008$ corrected, 400 ms: $t_{(14)} = -3.53$, $p = 0.01$ corrected; "Far=V" vs. "Far=H", 100 ms: $t_{(25)} = -5.29$, $p < 0.0001$ corrected, 400 ms: $t_{(14)} = -4.09$, $p = 0.003$ corrected). It is reassuring to note the comparable TI magnitudes for identical stimuli under the different conditions: "No Far" condition here, the "Near" condition of Fig. 4A, and the "Both" condition of Fig. 4B (~2° on average for the 400 ms measurement).

Next, to address the possibility that the effect is mediated by an interaction between near and far surrounds (e.g., lateral inhibition, Clifford, 2014; O'Toole & Wenderoth, 1977), we used a near surround tilted at ±45°, with the far surround oriented horizontally or vertically as before. In this experiment, the orientation difference between the near and the far surround is always the same (45° for both vertical and horizontal far surrounds), equating the interaction between the near and the far

surrounds. Importantly, the results again showed a robust reduction in TI in the presence of a vertical

530 compared with a horizontal far surround (Fig. 6, "400 ms N=45°" condition, $N = 12$ MTurk observers of whom 11 show the effect, $t_{(11)} = -4.45$, $p = 0.001$; a presentation duration of 400 ms was used). This suggests that the effect is mediated by an interaction between the far surround and the target, rather than an interaction between the far surround and the near surround. Note that although the far surround affects the orientation of the near surround (i.e., a TI of near surround toward or away from the

535 orientation of the far surround), this effect is probably not too large (~$\pm 2°$); hence, it seems unlikely to lead to a large change in the TI magnitude (i.e., TI from 43° vs. 47° seems unlikely to lead to a ~25% reduction in the effect magnitude). Overall, the results showed that the interaction between the target and the near surround (as measured by TI) interacts with the interaction between the target and the far surround (a three-way interaction: target × near surround × far surround). Put differently, the results

540 showed that competition exists between the different contextual interactions of the target (see the Discussion).

To address the possibility that the reduction in TI is mediated by an improved orientation acuity in the presence of a vertical orientation cue in the far surround (Wei & Stocker, 2017), we compared JNDs between vertical and horizontal far surrounds. JNDs were calculated as the inverse of the slope of the

545 psychometric function (i.e., percent clockwise reports as a function of target orientation, see the Methods). The results showed minor and non-significant modulated in JND in the presence of the vertical compared with the horizontal far surround, and even a nonsignificant deterioration in three of the five conditions ("100 ms": $t_{(8)} = -0.43$, $p = 0.67$; "400 ms": $t_{(13)} = 1.61$, $p = 0.13$; "100 ms H/V/E ": $t_{(25)} = -0.94$, $p = 0.35$, "400 ms H/V/E": $t_{(14)} = -0.19$, $p = 0.84$; "400 ms N=45°": $t_{(11)} = 0.65$, $p = 0.52$).

550 Therefore, the main effect was not explained by improved orientation acuity.

Fig. 6 – Non-linear competition between lateral interactions. *Average across observers of the TI due to a near-surround annulus, when a far-surround annulus is horizontal ("Far=H"), vertical ("Far=V"), or empty ("No Far"). The orientation of the near-surround was ±45° ("400 ms N=45°") or ±20° (all other conditions). The results showed a lower TI for "Far=V" compared with "Far=H" and "No Far". Error bars are ±1SEM.*

Non-linear competition between overlapping, dissimilar contexts

560 Next, we considered combinations of overlapping contexts that are dissimilar (Figs. 7 and 8). These experiments are important, because natural stimuli are composed of dissimilar orientations. Moreover, the findings may prove diagnostic for comparing the different theoretical accounts of bias (see the Discussion).

We first considered the TI caused by a tilted surround ("A"), a mix of a tilted and a vertical surround

565 ("A+V" plaid), and a mix of a tilted and a horizontal surround ("A+H" plaid) (Fig. 7). Importantly, in terms of first-order orientation (i.e., contrast), the mixtures maintain the orientation of "A", with the added vertical or horizontal orientations having no tilt relative to the vertical target. However, unlike a prediction based on the first-order orientation, the results showed that TI depends on the identity of the mixed grating, as indicated by a repeated measures ANOVA (Lab, 400 ms: $F_{(2,18)} = 27.76$, $p < 0.0001$, N

570 $= 10$ observers; MTurk, 400 ms: $F_{(2,12)} = 23.95$, $p = 0.0001$, $N = 7$ MTurk observers; MTurk, 100 ms: $F_{(2,22)} = 22.17$, $p < 0.0001$, $N = 12$ MTurk observers). Specifically, the "A+V" and "A" conditions measured a similar TI (differences showing $p > 0.05$ in each of the three experiments), whereas the "A+H" condition measured much less TI than "A+V", and much less TI than "A" (all post-hoc t-tests showing $p < 0.005$, Bonferonni corrected for three multiple comparisons). Moreover, the "A+H"

575 measurement was typically negative ($p < 0.05$ in each of the three experiments, again corrected for three multiple comparisons). This finding shows the non-additivity of the mixing.

The direction of TI can be explained by the perceived, second-order orientation (Dakin, Williams, & Hess, 1999; Smith, Clifford, & Wenderoth, 2001; Wei, Zhou, & Chen, 2018). As evident in a visual inspection (Fig. 7), the checkerboard pattern is grouped along either the orientation of the elements, or

580 the orthogonal orientation. For "A+V" with the near-surround titled -45°, this results in either a -22.5° or a +67.5° second-order orientation. For "A+H" with the near-surround titled -45°, this results in either a - 67.5° or a +22.5° second-order orientation. (We quantitatively verified this claim by finding the orientations with the most Fourier energy in the "contrast" of the stimuli, where "contrast" was calculated as the squared difference between the stimuli image and the background value.) Importantly,

585 the sign of TI depends on the orientation difference between the surround and the target: at ~5 to ~65° orientation differences, the shift in the perceived orientation is in the opposite direction relative to the surround orientation ("repulsive effect"), whereas orientation differences of ~80° lead to a shift of the perceived orientation in the direction of the surround ("attractive effect", also known as the indirect tilt

illusion) (Clifford, 2014). Therefore, under the "A+V" condition, the first-order orientation content (e.g., repulsive from -45°) leads to the same sign of TI as the second-order orientation (repulsive from -22.5°, or attractive toward +67.5°). However, under the "A+H" condition, the first-order orientation content (e.g., repulsive from -45°) leads to an opposite sign of TI than does the second-order orientation (attractive toward -67.5°, or repulsive from +22.5°). Therefore, under the mixed conditions, the direction of TI was consistent with the second-order (or perceived) orientation. This account suggest that when first- and second-order orientations are not consistent, the influence follows the higher-order orientation.

It is important to emphasize that although both adjacent (Fig. 6) and overlapping (Fig. 7) combinations with vertical and horizontal gratings showed non-linearity, the measured direction of influence was opposite (V>H for overlapping, and H>V for adjacent, see Fig. 6 and Fig. 7, respectively; see the Discussion). This observation is particularly interesting when considered in the context of phase scrambling approaches, showing that the phase (i.e., relative position) of an oriented component determines its influence.

Fig. 7 – Non-linear competition between first- and second-order orientations. Shown is the magnitude of TI, averaged across observers, measured for an oriented surround ("A"), and for mixtures of the oriented surround with vertical or horizontal surrounds ("A+V" and "A+H" plaids, respectively). Note that the vertical and horizontal surrounds do not cause TI because they are not titled relative to the vertical target. The results showed TI following the first-order orientation only in the absence of a higher-order oriented structure, as indicated by the measurement of negative TI under the A+H condition. Error bars are ±1SEM.

Non-linear competition between natural and phase-scrambled images

Finally, we considered combinations of overlapping contexts that are natural vs. reduced. To this end, we used 'biased' image adaptors, which are unmodified natural images that were selected to maximize the orientation content around a particular orientation (Dekel & Sagi, 2015). Observers were exposed to biased images ("Img", with content oriented around +25° or -25° to vertical), phase scrambles of the biased images ("Phs", the same oriented content as "Img", but with scrambled high-order image content), and a mixture of the two image types ("Mixed"). Importantly, the "Img" and "Phs" components of the mix were always taken from opposite orientation groups (-25° with +25°, or vice versa). Therefore, TAE under the mixed condition could either reflect the image component, the phase scramble component, or neither (additive).

The results showed that the TAE under the "Mixed" condition was in a direction consistent with the image content, that is, non-additive (Fig. 8). To quantify this effect, we compared the measurement under the "Mixed" condition to the average of the "Img" and "Phs" conditions, and found a significant difference ($N = 9$ MTurk observers, $t_{(8)} = 2.39$, $p = 0.05$ two-tailed repeated measures t-test, Fig. 8, "Common"). This result was replicated when tested under laboratory conditions instead of using the Mechanical Turk ($N = 10$ observers, $t_{(9)} = 4.24$, $p = 0.002$), when using color images instead of grayscale ($N = 10$ MTurk observers, $t_{(9)} = 3.28$, $p = 0.01$), and when using a briefer image adaptor presentation duration of 100 ms instead of 300 ms, to rule out the influence of eye movements ($N = 11$ MTurk observers, $t_{(10)} = 3.17$, $p = 0.01$).

Interestingly, when a detection task was used instead of the image-content classification task, the results showed additivity, i.e., the "Mixed" measured the average of "Img" and "Phs" (as indicated by the non-significance of the null hypothesis, $N = 19$ MTurk observers, $t_{(18)} = 0.28$, $p = 0.78$). The absence of non-linearity with the detection task suggests that the image classification task is implicated in mediating the competition between the contexts (see the Discussion). Moreover, this finding rules out the possibility that the direction of TAE in the "Mixed" condition reflects a higher local contrast in the "Img" stimuli (since phase scrambling maintains the global but not necessarily the local contrast in the image; see also Stojanoski & Cusack, 2014). (We note that the "Img" and "Phs" components indeed measured a minor and practically negligible difference, with a slightly higher "Img" TAE in most conditions, and a slightly above-zero "Mixed" TAE for the "Detection task", $p = 0.04$, see Fig. 8.) Overall, the results suggest that

selection between the component influences is task dependent. Specifically, when the task required attending to the image component over the phase-scrambled component, the TAE followed the image.

645 ***Fig. 8 – Non-linear competition between a natural image and a phase-scrambled image.*** *TAE, averaged across observers, is shown, with the sign determined by the image content orientation (i.e., opposite the phase content orientation). In the experiments reported here, we measured the TAE due to exposure to biased images ("Img", which are unmodified natural images with content oriented around +25° or -25° to vertical, Dekel & Sagi, 2015), phase scrambles of the biased images ("Phs"), or a*
650 *mixture of the two stimulus types having opposite orientations ("Mixed", a +25° image mixed with a phase scramble of a -25° image, or vice versa). Observers either performed an image classification task (animal vs. no animal), or a detection task (adaptor-present vs. adaptor-absent). The results showed, for the "Mixed" stimuli, a significant tilt in a direction consistent with the image content only in the presence of the classification task (not for the detection task), that is, non-additivity that depends on the*
655 *task. The results also showed no significant change in TAE magnitude due to phase scrambling, as indicated by a similar TAE magnitude for "Img" and "Phs" (note that the sign of TAE is the same; in the plot the sign is flipped to reflect the structure of the experiment under the "Mixed" condition). Error bars are ±1SEM.*

Discussion

We measured bias in the perceived orientation due to different spatiotemporal combinations of contexts. Our investigation focused on meaningful qualitative patterns, which we believe to be general, because it was impractical to systematically test all combinatorial combinations of all context types.

Additively of segregated contexts

We found that for adaptors that were presented at separate times (600 ms ISI), the combined influence was explained as an additive sum of the component effects (Fig. 3, see Eqs. (1) and (2)). Notably, there was no evidence for sub-additivity (as indicated by linearity for orientation repetition) and no evidence for selection (as indicated by linearity for orientation switch and side repetition). These findings are strengthened by a previous report on additivity of the TAE from adaptors and from learned expectations (Pinchuk-Yacobi, Dekel, et al., 2016). Further, Magnussen and Kurtenbach (1980) found additivity of the TAE and the TI. Overall, it seems that contexts that are sufficiently segregated in time interact additively in the measured influence, unlike commonly held beliefs.

An important observation concerning the additivity is that it refers to a specific measurement scale of orientation: degrees. Using a measurement scale that is not linear with degrees in the tested range (~2° change in the perceived orientation) would not show an additivity of measurements. For example, if the orientation changes were represented as the logarithm of the orientation difference, then additivity would be evident for the exponent of the measurements.

Theoretically, we expect additivity in degrees from an optimal calibration of orientation (see the cue combination theory below), and from population vector readout theories (whereby sufficiently small context-dependent changes in the representation are linear with the read-out bias in degrees) (Georgopoulos, Schwartz, & Kettner, 1986; Gilbert & Wiesel, 1990; Schwartz, Sejnowski, & Dayan, 2009). Our findings are in line with the existence of multiple, separate mechanisms of adaptation (Bao & Engel, 2012; Gekas, McDermott, & Mamassian, 2019), and it remains to be seen whether additivity is only evident when aggregating over separate mechanisms, or also within a mechanism (as long as the mechanism is not saturated).

Non-additivity of overlapping or adjacent contexts (TI and TAE)

When the contexts were adjacent or overlapping in space, the combined influence could not be generally

690   explained as an additive sum of separate influences (i.e., a non-linear interaction). Here, we parsed the

absence of additivity using two terms: sub-additivity, when a combined influence is lower than the sum

of the separate influences, and selection, when the combined influence reflects more one of the

components than the others.

Sub-additivity.

695   For increased context strength, we found sub-additivity. That is, an increase of the context area (Fig. 4)

or contrast (Fig. 5), which leads to a sub-linear increase in bias. A sub-linear increase in bias for

increased contrast was observed previously for the TAE (Blake et al., 2006; Harris & Calvert, 1989), but

here, the results can be interpreted in light of the finding of additivity (Fig. 3; using the same

experimental design). Moreover, an increase in adaptor presentation duration (which is arguably similar

700   to a change in contrast, Watson & Nachmias, 1977), is also known to lead to a sub-linear increase in

TAE (Dekel & Sagi, 2015; Sekuler & Littlejohn, 1974). Additivity for increased context area was, to the

best of our knowledge, never directly tested. (An experiment by Song & Rees, 2018 is similar to the

Up/Down experiment in Fig. 4B, but in their design the central target circle was halved along with the

surround, so it is unclear whether the findings reflect the change in the target or in the surround. Still,

705   their results of sub-additivity are consistent with our own.) Overall, the reports of sub-additivity seem

ubiquitous. Moreover, electrophysiological investigations of contextual influence seem consistent with

sub-linearity, though the mapping from neural activations to perception is not necessarily

straightforward (see, e.g., the results reviewed by Carandini & Heeger, 2012).

Interestingly, the results measured a spatial anisotropy of the additivity. Specifically, left and right

710   combinations were nearly additive, whereas up and down combinations measured a strong sub-additivity

(Fig. 4C). The anisotropy in additivity was not explained by anisotropy of TI of isolated components:

the left, right, up, and down quadrants measured nearly identical TI. A possible interpretation of this

finding is that the sub-additivity of combinations reflects low-level, intra-hemisphere interactions. This

claim seems consistent with a suggestion by Song & Rees (2018), based on fMRI analysis, that TI

715   reflects intra-hemisphere integration. Alternatively, the interaction between the components may depend

on the configuration (parallel vs. collinear).

For combinations of dissimilar contexts (Figs. 6-8), we often found a selection between the components, whereby the higher-level or task-relevant context was selected over a lower-level or a task-irrelevant

720   context.

We first considered the case where a "neutral" context is adjacent to a tilted context. By neutral, we refer to vertical or horizontal orientations, which cannot cause TI in the vertical target (due to symmetry). The results showed reduced or unchanged TI. Specifically, the TI from a tilted near-surround context was reduced in the presence of a vertical, but not horizontal, far surround context (Fig. 6). Moreover, the

725   reduction in TI cannot be explained by an interaction between the near and the far surrounds. Specifically, under a condition with a fixed orientation difference between the near and the far surrounds, we still observed a change in TI due to the far surround orientation (Fig. 6, "N=45°" condition). Therefore, the results indicate an interaction between the far surround and the target in mediating the effect. Based on these findings, we suggest that a selection process exists between

730   contexts, whereby inclusion of a context leads to a reduced influence of another context. Contexts that are similar to the target may be given preference for inclusion.

When a neutral and a tilted context were mixed in the same position (i.e., overlapping), the second-order orientations of the resulting plaid pattern determined the sign of the TI (Fig. 7). Specifically, for TI caused by a 45° tilt, a mix with a horizontal orientation inverted the sign of the TI, whereas a mix with

735   the vertical did not significantly change the TI. Given the results with the adjacent contexts above, it is clear that the TI is determined by the arrangement of the contexts' orientations: H<V when overlapping (Fig. 7), and V<H or V=H when adjacent (Fig. 6, "400 ms N=45°" condition). This observation is particularly relevant for phase scrambling approaches, showing that the phase (i.e., relative position) of an oriented component determines its influence. That is, the TI cannot be reduced to basic (first-order)

740   orientation processing circuits. In the current case, the direction of TI could be described by the second-order orientation (see Dakin, Williams, & Hess, 1999; Smith, Clifford, & Wenderoth, 2001; Wei, Zhou, & Chen, 2018). Possibly, in competition between first and second-order orientations, the TI follows the higher order orientation, consistent with the Gestalt approach to visual perception (Herzog, Thunell, & Ögmen, 2016; Jäkel, Singh, Wichmann, & Herzog, 2016). (An alternative account is that both first and

745   second order orientations receive a fixed weight, but that the "direct effect" from the second-order

orientation is stronger than the "indirect effect" from the first-order orientation even when tested with a first-order target (see Smith et al., 2001.))

When two contexts were matched based on the low- but not the high-level orientation content, the context with the higher-level orientations usually caused more bias when in competition (Fig. 8). Moreover, this effect was mediated by the task, indicating that the contexts were correctly balanced in terms of low-level statistics (see the "Detection task" in Fig. 8). Therefore, the spatial arrangement of the oriented content in the image affect the TAE. This result is consistent with modulation of TAE by perceptual grouping (He et al., 2012; Suzuki et al., 2005). Interestingly, the task dependence of the effect, and the lack of an effect in the absence of competition (compare "Img" to "Phs" in Fig. 8) are both consistent with a selection process between the different orientations, with preference in favor of the more task-relevant content (Ismail, Solomon, Hansard, & Mareschal, 2019; Pinchuk-Yacobi, Harris, et al., 2016). It is also interesting to note that visual attention, known to increase TAE magnitudes, seems to lead to a relatively modest and inconsistent effect (~25%) (Spivey & Spirn, 2000). The large effect magnitudes observed here when the different contexts are overlapping, suggests that selection has more of an effect in the presence of strong competition.

Contextual influence is the sum of segregated, separately bound contexts

Taken together, the results in this work showed an additive (linear) influence only when the contexts were segregated. Here, we attempted to answer the question of why some contextual combinations are additive whereas others are not. Above, we identified two patterns of non-linearity: sub-additivity and selection. We argue that the findings of sub-additivity can be explained by compression during early stimulus encoding, which binds visual stimulation using a narrow window of space and time into a higher-order structure. This idea is supported by accounts of contextual influence in terms of mechanisms exploiting statistical regularities in the visual environment (Coen-Cagli et al., 2015; Schwartz et al., 2007; Summerfield & De Lange, 2014). The contextual influence is then determined by selection between the competing encoded contexts (e.g., between high vs. low alternatives). Moreover, the influence (bias in degrees) is linear, given a selected, encoded context. To summarize, sufficiently segregated contexts are bound separately, and will accumulate additively in terms of measured bias, weighted by selection:

$$\text{Influence(contexts)} = \text{Influence(bound}_1, \text{bound}_2, \dots) = \qquad\qquad (7)$$

$$w_1 \cdot \text{Influence(bound}_1) + w_2 \cdot \text{Influence(bound}_2) + \cdots,$$

where w_i are the selection weights (possibly binary). A simple interpretation of these findings is that contextual influence and visual perception, both non-linear transformations, are at least partially shared during visual processing. This can be useful for the system in two ways. First, resolving an orientation from a stimulus is a computationally challenging task, so it seems reasonable to "recycle" the previously resolved orientations. Second, the information about the properties of an object becomes saturated when the object is fully resolved. For example, with a large enough grating, the orientation of the grating can be estimated to a sufficiently high accuracy, so an increase in grating size, contrast, or viewing duration will not provide much new information. However, a different object can be considered as a separate source of information, so its influence is not saturated. Overall, it seems possible that an efficient implementation of orientation context dependence relies on previously resolved orientations.

Bayesian cue combination

A possible realization of the above idea was investigated using a Bayesian framework with independent Gaussian cues that are used to infer an internal reference (Fig. 2). The inferred reference is partially, but not fully, local, as evident in the measurement of non-retinotopic TAEs (Fig. 3). Importantly, using model fitting, we found that the predictions of the cue-combination model were indistinguishable from additivity for the relevant parameter range, accounting for the observation of additivity with segregated contexts (Fig. 3).

Interestingly, when the cues cannot be assumed to be independent, the weight of the correlated cues is reduced (Oruç et al., 2003). This suggests an explanation of the observed sub-additivity of adjacent, and hence, correlated, contexts (Fig. 4). A similar argument applies for the observed sub-additivity of increased adaptor contrast (Fig. 5 and Blake et al., 2006) or duration (Dekel & Sagi, 2015; Sekuler & Littlejohn, 1974) (see the discussion above). In the TAE experiments with temporally segregated contexts, the correlation between cues was presumably negligible, which led to additivity. Note that estimating the relevant correlations using natural statistics is inaccurate, because reference cues probably change more slowly in space than is implied by image correlations (for example, two neighboring image regions can have very different image properties but the same frame of reference).

Another prediction of the theory is that the influence of each cue depends on the ratio between the cue reliability and the unadapted reference reliability (see Eq. (6)) (the unadapted reference encompasses all the unknown unadapted reference cues). Based on this prediction, the TI and TAE are expected to be
805   reduced in the presence of better reference cues. This claim is consistent with the behavioral observation of reduced TI in the presence of vertical compared with horizontal far surround (Fig. 6). (Of note, we did not observe an improvement in JND. Still, orientation discrimination JND depends on both internal and external noise contributions, so a reduction in internal noise may lead to a negligible improvement in JND when the external noise is the performance-limiting factor.)

810   It can also be observed that based on the model, bias is expected to be linear with the orientation difference between the context and the test (Eq. (6)). This prediction is not consistent with behavior: bias is known to reach a maximum for orientation differences of ~20° (e.g., Schwartz et al., 2007). Still, this discrepancy should probably not be considered as a strong argument against the theory, because orientations that are sufficiently far from the reference are obviously not suited to serve as good
815   reference cues (and hence deserve a lower weight). More extensive modeling is outside the scope of this work, but TAE may indeed be linear with small orientation differences (Campbell & Maffei, 1971; Magnussen & Kurtenbach, 1980).

Overall, the theory suggests how different reference cues are combined, which can account for the TAE and the TI. What counts as a reference may vary, including physical frames (e.g., the addition of a
820   luminance frame, as in the appendix of Knapen et al., 2010), predictions (Pinchuk-Yacobi, Dekel, et al., 2016), gravity (Wolfe & Held, 1982), and proprioception. In addition, the proposed theory probably does not account for all cases of orientation bias (for possible components of bias, see Clifford, Wenderoth, & Spehar, 2000). For example, the TAE experiments considered here use brief adaptor presentations (50 ms) and comparatively long inter-stimulus intervals (600 ms), a situation where
825   contrast adaptation can be argued to be negligible. Possibly, other experimental situations could be explained better by contrast adaptation, which may be an additional, separate cause of the TAE.

Acknowledgments

This research was supported by the Basic Research Foundation, administered by the Israel Academy of
830   Science. We thank Drs. Misha Katkov and Noga Pinchuk-Yacobi for their suggestions and comments.

Data and code availability

The data that support the findings of this study, the images used in the behavioral experiments, and a workable example of the code used in the web experiments are all available from the corresponding author upon reasonable request.

References

Alam, M. M., Vilankar, K. P., Field, D. J., & Chandler, D. M. (2014). Local masking in natural images: A database and analysis. *Journal of Vision*, *14*(8), 22–22.

Andrews, D. P. (1964). Error-correcting perceptual mechanisms. *Quarterly Journal of Experimental Psychology*, *16*(2), 104–115.

Bao, M., & Engel, S. A. (2012). Distinct mechanism for long-term contrast adaptation. *Proceedings of the National Academy of Sciences*, *109*(15), 5898–5903.

Blake, R., Tadin, D., Sobel, K. V, Raissian, T. A., & Chong, S. C. (2006). Strength of early visual adaptation depends on visual awareness. *Proceedings of the National Academy of Sciences*, *103*(12), 4783–4788.

Blakemore, C., & Campbell, F. W. (1969). On the existence of neurones in the human visual system selectively sensitive to the orientation and size of retinal images. *The Journal of Physiology*, *203*(1), 237–260.

Blakemore, C., Carpenter, R. H. S., & Georgeson, M. A. (1971). Lateral thinking about lateral inhibition. *Nature*, *234*(5329), 418.

Campbell, F. W., & Maffei, L. (1971). The tilt after-effect: a fresh look. *Vision Research*, *11*(8), 833–840.

Carandini, M., Demb, J. B., Mante, V., Tolhurst, D. J., Dan, Y., Olshausen, B. A., … Rust, N. C. (2005). Do we know what the early visual system does? *The Journal of Neuroscience*, *25*(46), 10577–10597.

Carandini, M., & Heeger, D. J. (2012). Normalization as a canonical neural computation. *Nature Reviews Neuroscience*, *13*(1), 51–62.

Clifford, C. W. G. (2014). The tilt illusion: Phenomenology and functional implications. *Vision Research*, *104*, 3–11.

Clifford, C. W. G., & Rhodes, G. (2005). *Fitting the mind to the world: Adaptation and after-effects in*

high-level vision (Vol. 2). Oxford University Press.

Clifford, C. W. G., Webster, M. A., Stanley, G. B., Stocker, A. A., Kohn, A., Sharpee, T. O., & Schwartz, O. (2007). Visual adaptation: neural, psychological and computational aspects. *Vision Research*, *47*(25), 3125–3131.

Clifford, C. W. G., Wenderoth, P., & Spehar, B. (2000). A functional angle on some after-effects in cortical vision. *Proceedings of the Royal Society of London. Series B: Biological Sciences*, *267*(1454), 1705–1710.

Coen-Cagli, R., Kohn, A., & Schwartz, O. (2015). Flexible gating of contextual influences in natural vision. *Nature Neuroscience*, *18*(11), 1648–1655.

Coltheart, M. (1971). Visual feature-analyzers and aftereffects of tilt and curvature. *Psychological Review*, *78*(2), 114.

Dakin, C. J., & Rosenberg, A. (2018). Gravity estimation and verticality perception. In *Handbook of clinical neurology* (Vol. 159, pp. 43–59). Elsevier.

Dakin, Williams, & Hess. (1999). The interaction of first-and second-order cues to orientation. *Vision Research*, *39*(17), 2867–2884.

Dekel, R., & Sagi, D. (2015). Tilt aftereffect due to adaptation to natural stimuli. *Vision Research*, *117*, 91–99.

Dekel, R., & Sagi, D. (2019). Seeing, fast and slow: the effects of processing time on perceptual bias. *BioRxiv*, 556944. https://doi.org/10.1101/556944

Deng, J., Dong, W., Socher, R., Li, L.-J., Li, K., & Fei-Fei, L. (2009). Imagenet: A large-scale hierarchical image database. In *Computer Vision and Pattern Recognition, 2009. CVPR 2009. IEEE Conference on* (pp. 248–255).

Dong, X., Gao, Y., Lv, L., & Bao, M. (2016). Habituation of visual adaptation. *Scientific Reports*, *6*.

Fründ, I., Haenel, N. V., & Wichmann, F. A. (2011). Inference for psychometric functions in the presence of nonstationary behavior. *Journal of Vision*, *11*(6).

Gekas, N., McDermott, K. C., & Mamassian, P. (2019). Disambiguating serial effects of multiple timescales. *Journal of Vision*, *19*(6), 24.

Georgopoulos, A. P., Schwartz, A. B., & Kettner, R. E. (1986). Neuronal population coding of movement direction. *Science*, *233*(4771), 1416–1419.

Gibson, J. J. (1937). Adaptation, after-effect, and contrast in the perception of tilted lines. II. Simultaneous contrast and the areal restriction of the after-effect. *Journal of Experimental Psychology*, *20*(6), 553.

Gibson, J. J., & Radner, M. (1937). Adaptation, after-effect and contrast in the perception of tilted lines.

I. Quantitative studies. *Journal of Experimental Psychology*, *20*(5), 453.

Gilbert, C. D., & Wiesel, T. N. (1990). The influence of contextual stimuli on the orientation selectivity of cells in primary visual cortex of the cat. *Vision Research*, *30*(11), 1689–1701.

Goddard, E., Clifford, C. W. G., & Solomon, S. G. (2008). Centre-surround effects on perceived orientation in complex images. *Vision Research*, *48*(12), 1374–1382.

Harris, J. P., & Calvert, J. E. (1989). Contrast, spatial frequency and test duration effects on the tilt aftereffect: Implications for underlying mechanisms. *Vision Research*, *29*(1), 129–135.

He, D., Kersten, D., & Fang, F. (2012). Opposite Modulation of High- and Low-Level Visual Aftereffects by Perceptual Grouping. *Current Biology*, *22*(11), 1040–1045.

Herzog, M. H., Thunell, E., & Ögmen, H. (2016). Putting low-level vision into global context: Why vision cannot be reduced to basic circuits. *Vision Research*, *126*, 9–18.

Ismail, A. M. H., Solomon, J. A., Hansard, M., & Mareschal, I. (2016). A tilt after-effect for images of buildings: evidence of selectivity for the orientation of everyday scenes. *Royal Society Open Science*, *3*(11), 160551.

Ismail, Solomon, J. A., Hansard, M., & Mareschal, I. (2019). A perceptual bias for man-made objects in humans. *Proceedings of the Royal Society B: Biological Sciences*.

Jäkel, F., Singh, M., Wichmann, F. A., & Herzog, M. H. (2016). An overview of quantitative approaches in Gestalt perception. *Vision Research*, *126*, 3–8.

Knapen, T., Rolfs, M., Wexler, M., & Cavanagh, P. (2010). The reference frame of the tilt aftereffect. *Journal of Vision*, *10*(1).

Kohn, A. (2007). Visual adaptation: physiology, mechanisms, and functional benefits. *Journal of Neurophysiology*, *97*(5), 3155–3164.

Lamme, V. A. (1995). The neurophysiology of figure-ground segregation in primary visual cortex. *Journal of Neuroscience*, *15*(2), 1605–1615.

Magnussen, S., & Kurtenbach, W. (1980). Linear summation of tilt illusion and tilt aftereffect. *Vision Research*, *20*(1), 39–42.

Mitchell, D. E., & Muir, D. W. (1976). Does the tilt after-effect occur in the oblique meridian? *Vision Research*, *16*(6), 609–613.

Movshon, J. A., & Simoncelli, E. P. (2014). Representation of Naturalistic Image Structure in the Primate Visual Cortex. *Cold Spring Harbor Symposia on Quantitative Biology*, *79*, 115–122.

Nakashima, Y., & Sugita, Y. (2014). Surround-contingent tilt aftereffect. *Journal of Vision*, *14*(14), 5.

O'Toole, B., & Wenderoth, P. (1977). The tilt illusion: Repulsion and attraction effects in the oblique meridian. *Vision Research*, *17*(3), 367–374.

Olshausen, B. A., & Field, D. J. (2005). How close are we to understanding V1? *Neural Computation*, *17*(8), 1665–1699.

Oruç, I., Maloney, L. T., & Landy, M. S. (2003). Weighted linear cue combination with possibly correlated error. *Vision Research*, *43*(23), 2451–2468.

Pinchuk-Yacobi, N., Dekel, R., & Sagi, D. (2016). Expectations and visual aftereffects. *Journal of Vision*, *16*(15), 19.

Pinchuk-Yacobi, N., Harris, & Sagi, D. (2016). Target-selective tilt aftereffect during texture learning. *Vision Research*, *124*, 44–51.

Schwartz, O., Hsu, A., & Dayan, P. (2007). Space and time in visual context. *Nature Reviews. Neuroscience*, *8*(7), 522–535.

Schwartz, O., Sejnowski, T. J., & Dayan, P. (2009). Perceptual organization in the tilt illusion. *Journal of Vision*, *9*(4), 19.

Sekuler, R., & Littlejohn, J. (1974). Tilt aftereffect following very brief exposures.

Smith, S., Clifford, C. W. G., & Wenderoth, P. (2001). Interaction between first-and second-order orientation channels revealed by the tilt illusion: psychophysics and computational modelling. *Vision Research*, *41*(8), 1057–1071.

Snow, M., Coen-Cagli, R., & Schwartz, O. (2017). Adaptation in the visual cortex: a case for probing neuronal populations with natural stimuli. *F1000Research*, *6*.

Solomon, S. G., & Kohn, A. (2014). Moving sensory adaptation beyond suppressive effects in single neurons. *Current Biology : CB*, *24*(20), R1012-22.

Song, C., & Rees, G. (2018). Intra-hemispheric integration underlies perception of tilt illusion. *NeuroImage*, *175*, 80–90.

Spivey, M. J., & Spirn, M. J. (2000). Selective visual attention modulates the direct tilt aftereffect. *Perception & Psychophysics*, *62*(8), 1525–1533.

Stojanoski, B., & Cusack, R. (2014). Time to wave good-bye to phase scrambling: creating controlled scrambled images using diffeomorphic transformations. *Journal of Vision*, *14*(12), 6.

Summerfield, C., & De Lange, F. P. (2014). Expectation in perceptual decision making: neural and computational mechanisms. *Nature Reviews Neuroscience*, *15*(11), 745.

Suzuki, S., Clifford, C., & Rhodes, G. (2005). High-level pattern coding revealed by brief shape aftereffects. *Fitting the Mind to the World: Adaptation and after-Effects in High-Level Vision*, *2*, 135–172.

Watson, A. B., & Nachmias, J. (1977). Patterns of temporal interaction in the detection of gratings. *Vision Research*, *17*(8), 893–902.

Webster, M. A. (2011). Adaptation and visual coding. *Journal of Vision*, *11*(5).

Wei, N., Zhou, T., & Chen, L. (2018). Objective measurement of gestalts: Quantifying grouping effect by tilt aftereffect. *Behavior Research Methods*, *50*(3), 963–971.

Wei, & Stocker. (2017). Lawful relation between perceptual bias and discriminability. *Proceedings of the National Academy of Sciences*, *114*(38), 10244–10249.

Wolfe, J. M., & Held, R. (1982). Gravity and the tilt aftereffect. *Vision Research*.

Yehezkel, O., Sagi, D., Sterkin, A., Belkin, M., & Polat, U. (2010). Learning to adapt: Dynamics of readaptation to geometrical distortions. *Vision Research*, *50*(16), 1550–1558.