Male genital lobe morphology affects the chance to copulate in *Drosophila pachea*

Bénédicte M. Lefèvre[1], Diane Catté[1], Virginie Courtier-Orgogozo[1], Michael Lang[1]

[1]Team "Evolution and Genetics", Institut Jacques Monod, CNRS, UMR7592, Université de Paris, 15 rue Hélène Brion, 75013 Paris

[*]corresponding author: michael.lang@ijm.fr

Abstract

Introduction: Male genitalia are thought to ensure transfer of sperm through direct physical contact

with female during copulation. However, little attention has been given to their pre-copulatory role

with respect to sexual selection and sexual conflict. Males of the fruitfly *Drosophila pachea* have a

pair of asymmetric external genital lobes, which are primary sexual structures and stabilize the

copulatory complex of female and male genitalia.

Results: We tested for a pre-copulatory role of these lobes with a *D. pachea* stock where males

have variable lobe lengths. In 92 mate competition trials with a single female and two males,

females preferentially engaged into a first copulation with males that had a longer left lobe and that

displayed increased courtship vigor. In 53 additional trials with both males having partially

amputated left lobes of different lengths, we observed a weaker and non-significant effect of left

lobe length on copulation success. Courtship durations significantly increased with female age and

when two males courted the female simultaneously, compared to trials with only one courting male.

Conclusion: Left lobe length affects the chance of a male to engage into copulation. The

morphology of this primary sexual trait may affect reproductive success by mediating courtship

signals or by facilitating the establishment of genital contacts at the onset of copulation.

keywords: *Drosophila pachea*, primary sexual trait, mating behaviour, mate-competition

experiments, genitalia

Introduction

Males and females exhibit different reproduction strategies due to higher energy costs of larger female gametes compared to smaller and more abundant male gametes [1]. This implies male-male intrasexual mate-competition for siring the limited female gametes. In turn, females may optimize reproduction by choosing males that confer survival and fecundity benefits to her and to the offspring. This dynamics has been formalized into genetic models [2, 3] involving female preferences for particular male characters and male mate competition, and can contribute to the rapid evolution of female preferences and male sexual attributes. There is also evidence for male mate-choice and female intra-sexual selection [4–6], showing that traditional sex-roles can change between species and in individuals under diverse conditions. In general, sexual selection implies a sexual conflict whenever male and female reproductive fitness strategies differ [7].

Across animals with internal fertilization, genitalia are usually the most rapidly evolving organs [8]. Several hypotheses propose that the evolution of genitalia is based on sexual selection and a consequence of sexual conflict between males and females at different levels of reproduction, including competition of sperm from different males inside the female (sperm competition) [9], female controlled storage and usage of sperm to fertilize eggs (cryptic female choice) [8, 10, 11], or sexually antagonistic co-evolution of aggressive and defensive morphologies and/or behaviors between male and female to control fertilization decisions [12, 13]. Male courtship is well known as a behaviour to attract females before copulation begins, but has also been reported to occur during or even after copulation [14, 15]. It is thought to be a widespread and key aspect of copulation in the cryptic female choice scenario, where males stimulate the female to utilize his own sperm [8, 10]. A variety of insect species were reported where males had evolved elaborated male genital structures that are apparently used for courtship throughout copulation to stimulate the female through tapping or other physical stimuli [14, 15].

Traditionally, sexual traits are categorized as primary or secondary. Genital structures are

considered "primary" when they are directly used for the transfer of gametes during copulation or

when they contribute to the complexing of female and male copulatory organs [8, 16]. Other traits

that differ between sexes and that are linked to reproduction are considered "secondary" sexual

traits. For primary sexual traits, sexual selection is traditionally thought to act during or after

copulation [8, 15], while secondary sexual traits can be involved in pre-copulatory mate-choice via

visual, auditory or chemical long range signalling [17–25].

Primary genitalia could also be possibly involved in pre-copulatory mate choice. Supposing

that a particular male genital trait enhances or favours female fecundity, it might become preferred

by the female during pre-copulatory courtship or at the moment of genitalia coupling [26–28]. This

preference could rely on direct benefits to reduce energy investment or predation risk caused by

unsuccessful copulation attempts, or on indirect male genetic quality to be inherited into the

offspring generation. Several examples of primary genitalia used in pre-copulatory courtship

signalling are known in vertebrates. For example, male and female genital display, swelling, or

coloration changes are associated with sexual activity [29–31] and penile display was also observed

in lizards [32]. Human male penis size was found to influence male attractiveness to women [33]. In

diverse mammalian species, females have also evolved external and visible structures that resemble

male genitals, a phenomenon called andromimicry [31, 34, 35]. These structures are likely used in

visual signaling, so it is possible that the male analogous parts also do. Furthermore, females of

some live-bearing fish species were reported to prefer to mate with males with large gonopodia [36,

37]. These cases involve visual stimuli and large genital organs. Only a few studies examined the

roles of primary genitalia in invertebrates during courtship or at the onset of copulation before

phallus intromission is achieved [26–28, 38, 39]. Longer male external genitalia of the water strider

Aquarius remigis were associated with higher mating success [28]. In addition, male genital spine

morphology of *Drosophila bipectinata* and *Drosophila ananassae* were reported to evolve in

response to direct competition among males for securing mates in a so called 'scramble

competition' scenario [26, 27]. In general, the role of primary sexual traits in establishing

copulation or their role during courtship needs further investigation.

The fruitfly species *Drosophila pachea* is a promising model to study the evolution of

primary sexual traits, especially with respect to the evolution of left-right asymmetry. Male *D.*

pachea have an asymmetric aedeagus (intromission organ) and a pair of asymmetric external lobes

with the left lobe being approximately 1.5 times longer than the right lobe [40–42] (Figure 1). The

genital lobes have likely evolved during the past 3-6 Ma and are not found in closely related species

[43]. *D. pachea* couples also mate in a right-sided copulation position where the male rests on top

of the female abdomen with the antero-posterior midline shifted about 6-8 degrees to the right side

of the female midline [41, 42, 44]. Male and female genitalia form an asymmetric complex during

copulation and the asymmetric lobes stabilize this genital complex [44]. Furthermore, *D. pachea* is

among the *Drosophila* species that produce the longest (giant) sperm [45, 46], and their ejaculates

contain in average just about 40 sperm cells [47]. Thus, a particular right-sided mating position

could potentially be associated with optimal transfer of giant sperm during copulation [42].

The aim of this study was to test if the asymmetric genital lobes of *D. pachea* would have an

effect on pre-copulatory mate-choice, in addition to their role in stabilizing the complex of male and

female genitalia during copulation. Previously, we found that males originating from one of our *D.*

pachea laboratory stocks possess short and rather symmetric lobes [41] (Figure 1), while others

have the typical size asymmetry. The variability in left genital lobe length that we observed within

this fly stock enabled us to test if lobe length might affect pre-copulatory courtship or mate

competition. We selected *D. pachea* males with short left lobes and produced a stock with an

increased variance of left lobe length. Next, we tested sibling males of this "selection stock" in

mate-competition assays for their success to engage first into copulation with a single female. We

also tested whether copulation success in our assay would be affected by male courtship vigour.

Then, we surgically shortened the length of the left lobe in males that had developed long left lobes,

to further test whether left lobe length affects copulation success. Finally, we assessed whether lobe

length affects ejaculate allocation into female storage organs.

Results

Genital lobe lengths differ between _D. pachea_ stocks

In our laboratory stock 15090-1698.01, _D. pachea_ males display a characteristic left-right

size asymmetry of genital lobes with the left lobe being consistently larger than the right lobe

(Figure 1A,C, Figure 2 A, Table S4, Additional datafile 3). Similarly, in stock 15090-1698.02, most

males reveal larger left lobes, but a few individuals are observed with particularly small lobes that

are rather symmetric in length (Figure 1B-D, Figure 2B). We selected for males with short lobe

lengths and crossed them with sibling females for 50 generations (Material and Methods). The

resulting stock was named "selection stock". We observed an increased variance of left lobe length

compared to the source stock (Levene's test: selection stock / 15090-1698.02: DF = 1/147, F =

11.506, P = 0.0008913) (Figure 2C, Additional datafile 3). In contrast, the right lobe length differed

only marginally among the two stocks (Table S4) and the variances of right lobe lengths were not

significantly different among them (Levene's test, DF = 1/147, F = 0.5164, P = 0.4735).

Male genital left lobe length, male courtship and failed copulation attempts have an effect on copulation success

119 We performed a mate competition assay (Material and Methods) with two males and a

120 single female to test if differences in genital lobe length and courtship behavior may influence the

121 chance of a male to copulate first with a single female. This first engagement into copulation will be

122 referred to as "copulation success". We filmed courtship and copulation behavior of two males of

123 the selection stock and a single female of stock 15090-1698.01 or 15090-1698.02. We annotated

111 trials where both males simultaneously courted the female (Material and Methods) (51 and 60 trials with females of stocks 15090-1698.01 or 15090-1698.02, respectively). In particular, we waited for a first copulation to occur, then isolated the non-copulating male and finally the other two specimen once the first copulation had ended. Then, we dissected the genitalia of both males to determine lobe lengths. We examined 54 additional trials with two courting males from the selection stock that had partially amputated left lobes and females of stock 15090-1698.01. This was done to test if other characters might co-vary with left lobe length and could potentially also have an effect on copulation success. We shortened the left lobes of both males in order to be able to neglect possible effects of the caused wound. The relative courtship activity or vigor of the two males in each trial was compared by quantifying periods that each male spent touching the female ovipositor or the ground next to it with the proboscis (mouth-parts), hereafter referred to as "licking behavior" (Material and Methods). We counted the sum duration of these periods (sum licking duration) and the number of licking periods (number of licking sequences). A copulation was considered to take place, once a male had mounted the female abdomen and achieved to settle into an invariant copulation position (Material and Methods).

In the majority of trials (130/165), the first mounting attempt of a male resulted in a stable copulation. We did not observe any male behavior that could indicate an attempt of the non-copulating male to directly usurp the female as observed for *D. ananassae* or *D. bipectinata* [26, 27]. However, in 35 trials, a total of 51 failed mounting attempts were observed (Material and Methods). These were not proportionally higher among trials with lobe amputated males (9 failed mounting attempts in 7 / 54 trials 13%) compared to unmodified males (42 failed mounting attempts in 28 / 111 trials 25%). Thus, the partial lobe amputation probably did not decrease the ability of a male to mount a female. Only 15/51 failed mounting attempts lasted 8 seconds or longer. Among those, the non-mounted male continued to court the female in 10/15 attempts. However, this was also observed in 131/165 successful mounting attempts and these proportions are not

significantly different (χ^2-test, χ^2 : 1.4852, df = 1, P = 0.2230). Altogether, mounting failure was not

found to be significantly associated with the courtship activity of the other male.

Different variables were tested to have en effect on male copulation success by using a

Bradley Terry Model [48] (Material and Methods). In trials with unmodified males, copulation

success was positively associated with left lobe length and the sum licking duration of the males

during courtship (Table 1). The effect of sum licking duration is illustrated when comparing the

ratio of sum licking duration and total courtship duration between the copulating and the non-

copulating male (Figure 3A). This ratio was higher in the copulating male compared to the non-

copulating male. Copulation success was negatively associated with the number of licking

sequences and with the number of failed mounting attempts (Table 1). In particular, we identified a

positive interaction of failed mounting attempts and female age, indicating that the negative effect

of failed mounting attempts is pronounced in young females but not in older females. However, this

difference turned out to be based on three outlier observations when comparing the number of failed

mounting attempts of either male over female age (Additional datafile 1, Figure S5). Therefore,

more observations are required to evaluate this interaction and the effect of failed mounting

attempts on copulation success.

In trials with males that underwent lobe surgery, left lobe length was not significantly

associated with copulation success (Table 2), although a similar positive effect was detected. Thus,

we cannot exclude the possibility that left lobe length co-varies with yet another unknown selected

trait that reveals an advantage for the male to copulate. Similarly, a non-significant negative effect

of the number of licking events on copulation success was observed with those males (Table 2) but

a positive significant effect of sum licking duration.

Courtship duration is increased when two males court a female simultaneously

Regardless of the stock for the female used, courtship durations appeared to be longer when

both males courted the female simultaneously compared to trials where only one male courted

(Mann-Whitney-Wilcoxon test, W = 1906.5, N = 48/165, P = 4.7×10^{-8}) (Figure 3B). We also

observed that courtship duration was strikingly increased in females that were older than 11 days

(after emerging from the pupa) compared to younger females (Additional datafile 1: Figure S4)

(Mann-Whitney test, W = 660, females younger than 12 days / females 12 days or older = 33 / 132,

P = 6.333×10^{-10}). Thus, older females may be less likely to copulate and may also discriminate less

between males based on failed mounting attempts (see above).

Lobe length is not associated with the amount of ejaculate in female sperm storage organs

To test whether lobe length influences the amount of ejaculate deposited into the female

after copulation, we dissected a random subset of females (54 of stock 15090-1698.01 and 24 of

stock 15090-1698.02) in trials with non-modified lobes that revealed a copulation. Neither right

lobe length, left lobe length, sum licking duration or female age (days after emerging from the

pupa) was significantly associated with ejaculate content in the spermathecae (Table 3). However,

male adult age and copulation duration had a negative effect. Given that *D. pachea* males become

fertile about 13 d after emerging from the pupa [49], our results indicate an optimal period of male

ejaculate transfer at the beginning of their reproductive period. Copulation duration had a negative

effect on sperm transfer (Table 3), indicating that failed sperm transfer may result in prolonged

attempts and longer copulation duration.

Discussion

Left lobe length affects the chance of a male to copulate

We observed a tendency of females to copulate first with the male that had the longer left lobe when two males courted simultaneously in our trials. These results suggest that a long left lobe might not only stabilize an asymmetric genital complex during copulation [44], but also increases the chance of a male to engage into copulation. With our data, we cannot exclude that the sensing of the lobe length by the female occurs just before intromission, while male and female genitalia are contacting each other or if it is used even during courtship before genital contacts establish. Other factors, unrelated to lobe length, can also bias copulation. For example, females might prefer to copulate with males of larger body size or that display particular courtship behaviors. In our study, we found no significant effect of tibia length on copulation success, which was used as an approximation for overall body size. However, the mutation(s) associated with lobe length variation in this stock are unknown. It is possible that one or several alleles associated with lobe length still segregate in this stock, even after 50 generations of inbreeding. If such segregating mutations have pleiotropic effects on other male characters (morphological, physiological or behavioural), these traits are expected to co-vary together with lobe length and the effect of lobe length that we detected might actually be due to the other factors. In trials with artificially introduced lobe length variation (through surgery), males were taken from the selection stock but had the expected wild-type lobe asymmetry. Left lobe length difference was then artificially introduced in those males, so that it would not be expected to co-vary with the relative phenotypic expression of the supposed underlying mutation. In such trials, we observed a trend of the left lobe length being associated with copulation success, but this trend was not significant. Thus, we cannot completely rule out the possibility that left lobe length affects copulation success indirectly via a pleiotropic mutation affecting both, lobe length and other traits.

Male intra-sexual competition is potentially reflected by male courtship intensity

Previous analyses in *D. ananassae* and *D. bipectinata* relate male genital spines to pre-copulatory abilities to achieve a copulation due to increased male intrasexual competitiveness in so called "scramble competition" or vigorous copulation attempts with females that are optionally already copulating with anoher male [26, 27]. In *D. pachea*, this behavior was not observed and most mounting events of a male resulted in a stable copulation. Once a copulation started, the non-copulating male was never observed to separate the couple. Instead, it continued to display courtship behaviors such as licking, wing vibration or touching the female side in the majority of cases. In addition, male mounting was only observed upon female wing opening and oviscapt protrusion. This indicates that mounting in *D. pachea* is a complex behavior involving both sex partners. Upon mounting, the chance of a *D. pachea* male to engage into copulation thus may largely depend on male-female communication rather than on male male competition.

Total courtship durations were increased in trials where both males courted the female simultaneously compared to trials with only one male courting. This indicates that females potentially require more time to accept one of the males for copulation in cases where two males court simultaneously. Alternatively, longer courtship durations might indeed rely on male-male competition, causing mutual disturbances in courtship display. We detected a strong association of copulation success with the sum licking duration, which we used as an approximation for overall courtship activity or vigor of each male. We also observed a negative effect of the number of licking sequences on copulation success, indicating that a male has highest chances to copulate if he continuously presents licking behavior at the female abdomen for a prolonged time period. This is achieved by maintaining its position close to the female oviscape in competition with the other male, suggesting that licking behavior in part reports male-intrasexual conflict.

Potential female benefits gained from choosing males with long left lobes

Female mate choice on male sexual traits was suggested to rely at least in part on the way a sexual trait is displayed or moved [50]. This implies that a certain quantity but also accuracy of locomotor activity would matter in courtship and might better reflect overall male quality and "truth in advertising" than a morphological character or ornament alone. Our data suggest that male settling upon mounting the female might be crucial for a successful copulation since we identified a negative effect of failed mounting attempts on copulation success and a positive interaction of failed mounting attempts with female age, possibly indicating that young females are more discriminative compared to older females with respect to male mounting performance. Female mate choice is hypothesized to be based on direct benefits, affecting fecundity and survival of the female, but also on indirect benefits that relate to genetic quality, fecundity and survival of the progeny [50, 51]. Left lobe length and male mounting performance might therefore influence female mate-choice through enhancing the quality of male courtship signals and by enhancing the coupling efficiency of male and female genitalia upon mounting.

Alternatively, left lobe length might potentially associate with direct copulation benefits for male and female by ensuring the efficiency of ejaculate transfer during copulation. Males of *D. pachea* produce giant sperm and previous estimates revealed that a *D. pachea* male transmits only about 44 ± 6 sperm cells per copulation [47], while the maximum female sperm storage capacity in the spermathecae was estimated to be much higher and to be about 304 sperm cells [47]. Wild-caught *D. pachea* females were found to contain sperm from at least 3-4 males based on spermathecae filling levels [47]. As asymmetric lobes stabilize the genital complex [44], they might have a positive effect on the amount of sperm transferred per copulation. However, we have not observed any effect of lobe length on ejaculate presence in the spermathecae after copulation. We note that our quantification method was not precise and that sample size was limited. Accurate direct sperm counts requires radio-labeling methods [47], which was not applicable in our

experimental setup. In future approaches, improved sperm counts should be applied to test for a

correlation between lobe length, female-male copulation complex stability and sperm quantity in

female spermathecae. Based on our data, we note that left lobe length affects the chance of a male

to establish genital contacts that lead to a firm complex of female and male genitalia, congruent

with previous observations in other insects [26–28, 52]. In particular, the male genital spines in *D.*

ananassae and *D. bipectinata* were observed to play a role in pre-copulatory events but had no

measurable influence later on with respect to sperm transfer or fecundity [26, 27, 52].

Interestingly, copulation duration negatively affected ejaculate presence in the female. This

observation suggests that the couple separates once successful sperm transfer is achieved. However,

not all copulations reveal sperm transfer. In occasions of failure the copulation duration may

therefore be prolonged to further attempt to mate. Copulation duration varies largely among

Drosophila species from fractions of a minute to up to about 2 h in *D. acanthoptera* [42, 49].

Copulation duration in *D. pachea* is thus comparatively long with respect to other species of the

genus, pointing to presumably complex copulatory interactions affecting sperm transfer. Future

studies must analyze the internal processes of sperm transfer during *D. pachea* copulation to better

understand this process and its possible limitations.

Indirect paternal effects on the growth and survival of offspring could potentially play a role

in *D. pachea* female choice for males with long left lobes. It has been demonstrated that mate

choice based on particular male sexual characters correlates with offspring survival in diverse

species [18, 20–23, 53]. However, our experiment did not allow us to test for these effects because

females were sacrificed after the mating experiment and no progeny was obtained. We assessed

ejaculate filling levels instead of the amount of female progeny because *D. pachea* females rarely

lay eggs after a single insemination and about four copulations are necessary for a female to start

oviposition [54]. To test for indirect effects, future studies should ideally use progeny males from

wild caught *D. pachea* females and test if subtle lobe length variation in those males would

correlate with offspring survival in single couple crosses. A similar approach was used by Hoikkala et al. (1998) [53], who found that *Drosophila montana* courtship song frequency of wild caught males correlated with the survival rate of the male's progeny.

Age of reproductive activities appear to be non-overlapping in sibling males and females

The amount of ejaculate present in the female spermathecae after a single copulation was negatively affected by the age of the copulating male. Similarly, it was previously found that the number of progeny in single couple crosses (presumably involving multiple copulations) was also dependent on male adult age in *D. pachea* [55]. Males in *D. pachea* need about 10-14 days at 25° C after emergence to become sexually mature [49, 54, 55]. This is related to adult testis growth and production of giant sperm [47]. It was shown that the relatively long time to reach male sexual maturity impacts the proportion of sexually active *D. pachea* adults (operational sex ratio), which is female-biased [49]. Our findings suggest that males potentially have a maximum fertility period at the beginning of their sexual maturity at approximately 13-16 days after emerging from the pupa. The detected reduction of transferred ejaculate in males older than 16 days implies an additional male reproductive limitation.

It was hypothesized that the delay in male sexual maturity might potentially lead to decreased sibling mating in *D. pachea* [54]. In *D. melanogaster*, sibling matings yield fewer progeny compared to crosses with unrelated individuals [55]. In our study, the female was not a sibling of the two males. Courtship durations were shortest in trials with "young" virgin females, between 6 and 11 days. Given that females reach sexual maturity at about 4 days after emerging from the pupa [49, 55], our observation suggests that young females engage more quickly into copulation compared to older females. Overall, female and male maximum reproductive activities appear to be non-overlapping among siblings of opposite sex and similar age. Such an effect can be relevant in the wild if a cactus rot is colonized by a single previously fertilized female.

Conclusion

Longer male left genital lobes potentially enhance the probability of *D. pachea* males to engage into copulation, which is expected to confer a reproductive advantage over males with shorter left lobes. The chance of a male to copulate first with a female was also affected by the relative male courtship activity and potentially by mounting attempt failures, especially for females at the beginning of their reproductive activity. The evolution of left lobe length might not only relate to its previously known function as a stabilizing device during copulation, but also through pre-copulatory mate choice by increasing the chances to copulate and to increase the coupling efficiency of genitalia upon mounting. Lobes do not appear to impact sperm transfer in later copulatory events. In addition, reproductive ages are non-overlapping between the two sexes potentially decreasing the amount of sibling mating in this species.

Material and Methods

Fly stock establishment and maintenance

Two *D. pachea* stocks 15090-1698.01 and 15090-1698.02 were retrieved from the San Diego Drosophila Species Stock Center (now The National Drosophila Species Stock Center, College of Agriculture and Life Science, Cornell University, USA) (Additional datafile 1: Table S1). Flies were maintained in 25 x 95 mm plastic vials containing 10 mL of standard Drosophila medium (60 g/L brewer's yeast, 66.6 g/L cornmeal, 8.6 g/L agar, 5 g/L methyl-4-hydroxybenzoate and 2.5% $^v/_v$ ethanol) and a ~ 10 x 50 mm piece of bench protection sheet (Bench guard). As *D. pachea* requires 7-dehydrocholesterol for proper development [56–58], we mixed the medium of each vial with 40 µL of 5 mg/mL 7-dehydrocholesterol, dissolved in ethanol (standard *D. pachea* food). Flies were kept at 25°C inside incubators (Velp) with a self-made light installation for a 12 h light: 12h dark photo-periodic cycle combined with a 30-min linear illumination change between

light (1080 lumen) and dark (0 lumen). We used males of stock 15090-1698.02 to generate a new

stock with increased proportions of males with short gential lobes. For this, we chose 3 males with

apparently symmetric (aberrant) genital lobes and crossed them with 3-4 sibling virgin females. We

repeated the selection with the progeny for a total of 36 generations. Then, we discarded males with

clearly visible asymmetric (wild-type) lobes from the progeny for another 14 generations to derive

the final stock (selection stock).

Virgin fly selection

Virgin flies at 0-1 d after emerging from the pupa were CO_2 anaesthetised on a CO_2-pad

(INJECT+MATIC Sleeper) under a stereo-microscope Stemi 2000 (Zeiss), separated according to

sex and maintained in groups of 20-30 individuals. Males and females were raised until reaching

sexual maturity, about 14 days for males and 4 days for females at 25° C [49]. This allowed us to

use virgin individuals in each mating experiment. Males were anaesthetised on the CO_2 pad (see

above), sorted according to lobe morphology (asymmetric and symmetric lobes) and isolated into

single vials at least two days before each mating experiment took place.

Mate-competition assay

We used a single virgin female of wild-type stock 15090-1698.01 or 15090-1698.02 and two

sibling males of the selection stock that visually differed in lobe length when inspected with a

binocular microscope (Additional datafile 1: Figure S1). This selection was done to increase the

average pairwise difference of lobe lengths between the two males. Specimens were introduced into

a white, cylindrical mating cell (Additional datafile 1: Figure S2) with a diameter of 20 mm, a depth

of 4 mm and a transparent 1 mm Plexiglas top-cover. Optionally, mating cells were concave with a

diameter of 20 mm and a depth of 4 mm at the center (Additional datafile 2: trials 1-16). Flies were

transferred without CO_2 anaesthesia using a fly aspirator: a 7 mm diameter silicone tube closed at

the tip with cotton and a 1000 µL wide bore (3 mm) micro-tip. Movie recordings were started as

soon as the chamber was immediately put under the camera (see below).

All mating trials were carried out inside a temperature and humidity controlled climate

chamber at 25° C ± 0.1° C and 80% ± 5% or 60% ± 5% (trials 1-15) humidity. We used digital

cameras MIRAZOOM MZ902 (OWL) or DigiMicro Profi (DNT) to record copulation and

courtship behaviour. The MIRAZOOM MZ902 (OWL) camera was mounted on a modified

microscope stand (191348, Conrad) equipped with a 8-cm LED white light illumination ring (EB-

AE-COB-Cover, YM E-Bright) and a platform to hold four individual cylindrical mating cells

(Additional datafile 1: Figure S2). Experiment 16 was filmed with the OWL camera and a concave

shaped mating cell that was put on a flat plastic cover on top of the microscope stand. For trials 1-

15, we used concave shaped mating cells that were put onto the stand of the DigiMicro Profi (DNT)

camera. Data was acquired with the programs Cheese (version 3.18.1)

(https://wiki.gnome.org/Apps/Cheese) or GUVCVIEW (version 0.9.9) GTK UVC (trials 1-15) on

an ubuntu linux operating system in webm or mkv format. Up to four trials were recorded in a

single movie, which was split after recording and converted into mp4 format with Avidemux 2.6

(http://www.avidemux.org/) to obtain a single movie per experiment.

We waited for a copulation to take place between one of the males and the female (see

below). The mating cell was then shortly recovered from the climate chamber in order to remove

the non-copulating male with an aspirator and to transfer it into a 2-mL reaction tube filled with

70% ethanol. This was performed after the copulating male appeared to have settled into a stable

mating posture on the female abdomen, which was controlled by live-imaging. The copulating male

and the female were isolated from the mating cell after copulation had ended and were also isolated

into single 2-mL reaction tubes filled with 70% ethanol. Optionally, the female was kept alive for

12-24 h in a vial containing 5-10 mL grape juice agar (24 gr/L agar, 26 gr/L sucrose, 120 mg/L

Tegosept, 20% $^v/_v$ grape juice, and 1.5% $^v/_v$ ethanol). The presence of eggs on the plate was

systematically checked but never observed. Females were finally sacrificed to prepare

spermathecae.

Left lobe surgery

Epandrial lobe surgery was done on 5-6 day-old *D. pachea* adult males of the selection stock

following Rhebergen et al. (2016). Males were anaesthetised on a CO_2 pad (see above) and then

further immobilized with a small copper wire, which was slightly pressed onto the male abdomen.

The left epandrial lobe was shortened to various lengths with micro dissection scissors (SuperFine

Vannas Scissors, World Precision Instruments). Flies were let to recover for at least 7 days on

standard *D. pachea* food at 25° C in groups. No mortality was observed in males that underwent

lobe surgery, similar to what was observed by Rhebergen et al. (2016). Partially amputated lobes

ranged in length from lacking at least their most distal tip to more reduced stumps whose length

approximated the corresponding right lobe.

Adult dissection and imaging

Adults were dissected in water with forceps (Forceps Dumont #5, Fine Science Tool) inside

a transparent 25 mm round dissection dish under a stereo-microscope (Zeiss Stemi 2000). Male

genitalia were isolated by piercing the abdomen with the forceps between the genital arch and the

A6 abdominal segment and thereby separating the genitalia from the abdomen. We also dissected

the left anterior leg of each dissected male. Female spermathecae were recovered after opening the

ventral abdomen with forceps and removal of the gut and ovaries. The spermathecae were isolated

and separated according to left and right sides and immediately examined using a microscope (see

below). All dissected tissues were stored in 200 µL storage solution (glycerol:acetate:ethanol, 1:1:3)

at 4° C.

For imaging, male genitalia were transferred into a dissection dish filled with pure glycerol

and examined with a VHX2000 (Keyence) microscope equipped with a 100-1000x VH-Z100W

(Keyence) zoom objective at 300-400 fold magnification. Genitalia were oriented to be visible in

posterior view. The left and right lateral spines, as well as the dorsal edge of the genital arch were

aligned to be visible in the same focal plane. In some preparations, the genital arch broke and lobes

were aligned without adjusting the position of the dorsal genital arch. The experiment was

discarded from analysis in cases where lobes could not be aligned. Male lobe lengths were

measured on acquired images as the distance between the base of each lateral spine and the tip of

each lobe (Figure 1c,d) using ImageJ version 1.50d (https://imagej.nih.gov/ij). Male legs were put

on a flat glycerol surface with the inner side of the tibia facing to the camera. Legs were imaged at

200 fold magnification with the VHX 2000 microscope (Keyence) as described above.

We prepared the female paired sperm storage organs (spermathecae) at 12 h - 24 h after

copulation (54 females from stock 15090-1698.01 and 24 females from stock 15090-1698.02,

additional file 2) and examined the presence of ejaculate. *D. pachea* has giant sperm [47], which

makes direct sperm counts difficult. Therefore, we determined an apparent average spermathecae

filling level per female based on visual inspection, similar to Jefferson (1977) [55]. Female

spermathecae were arranged on the bottom of the transparent plastic dissection dish, filled with

water. Images were acquired using transmission light in lateral view with the Keyence VHX2000

microscope at 400-500 fold magnification. Ejaculate was directly visible inside the transparent

spermathecae. Ejaculate filling levels were annotated for each spermatheca separately to match

three different categories: 0, 1/6 or 1/3 of its total volume. Then, the average filling level for both

spermathecae was calculated for each female.

Annotation of courtship and copulation behaviour

Videos were analysed with the OpenShot Video Editor software, version 1.4.3

(https://www.openshot.org) to annotate the relative timing of courtship and copulation in our

experiments (Additional datafile 1: Figure S3). Data was manually entered into spreadsheets. We

annotated the beginning of male courtship as the start of at least three consecutive male courtship

behaviours according to Spieth (1952) [59], such as male touching the female abdomen with the

forelegs, wing vibration, male following the female, and male licking behavior (see above). We estimated licking behavior throughout courtship of each male in trials were both males courted the female simultaneously. This data was used to approximate male courtship vigor. We calculated the sum duration of all licking events per male and trial and the number of licking sequences. However, this behavior could not be quantified in 7 trials because the males changed positions very fast so that they could not be unambiguously distinguished (Additional datafile 2, Additional datafile 4). In all analysed trials, licking behaviour was observed throughout *D. pachea* courtship from shortly after courtship start (Materials and Methods) to the start of copulation (Additional datafile 1: Figure S3). Both males reached at least once the licking position in all trials (Additional datafile 1, Table S3). This indicates that the female had physical contact with the proboscis of both males before copulation started (Additional datafile 1: Figure S3).

The beginning of copulation was defined as the moment when the male mounted the female abdomen. However, we only scored a copulation start if the male also settled into an invariant copulation posture. Upon mounting, the male moves its abdomen tip forth and back along the female oviscape. This behavior lasts for up to about 3 minutes and during this time the male grasps the female wings and abdomen with its legs. This behavior was previously described as settling period and ends with a characteristic right-sided, invariant copulation posture [41, 42, 44]. Any copulation that ended within the first minutes after mounting without settling was considered to be a "failed mounting attempt" (Additional datafile 4). The end of copulation was considered as the moment when male and female genitalia were separated and the male had completely descended with the forelegs from the female abdomen.

Copulation duration (Additional datafile 1, Table S3)) was comparable to previous analyses of *D. pachea* copulation durations [41, 44, 47, 55]. We observed that copulation ended upon isolation of the non-copulating male in 3 trials with females of stock 15090-1698.01 and in 4 trials with females of stock 15090-1698.02, probably indicating premature copulation end due to

460 specimen handling . However, the estimates were included into the dataset and had little impact on

461 average copulation duration when comparing mean and median values.

Mate-competition analysis

Our aim was to compare at least 50 mate-competition trials, where both males courted the

female simultaneously and where copulation was observed. In total, we carried out 98 and 89 trials

with females of stocks 15090-1698.01 and 15090-1698.02, respectively (Additional datafile 1:

Table S2). We performed another 62 trials with males that underwent surgical manipulation of the

left lobe and females of stock 15090-1698.01 (Additional datafile 1: Table S2). In total, we removed

36 trials from the analysis because either copulation was not observed until 1 h after recording

started (21 trials), flies escaped, got injured or died inside the mating cell (4 trials) or the dissections

of male genitalia failed (11 trials, Additional datafile 1: Table S2). We thus examined 152 trials with

unmodified males and 61 trials with males that had undergone genital surgery, (see below,

Additional file 1: Table S2). Among those, we observed that both males courted the female

simultaneously in 111 and 54 trials, respectively. Statistical assessement with the Bradley Terry

Model (see below) further required a complete dataset so that we excluded 20 additional trials with

missing values, 19 trials with unmodified males (left lobe length: 2 trials, right lobe length: 2 trials,

licking behavior and/or failed mounting attempts: 7 trials, tibia length: 8 trials) and one trial with

lobe-modified males (no information on tibia length). In total, we considered 92 trials with

unmodified males and 53 trials with lobe-modified males.

Data analysis was performed in R version 3.6 [60]. Male copulation success (see above) was

evaluated with a Bradley Terry Model [48] using the R package BradleyTerry2 [61]. We

incorporated data (Additional dtafile 2) into a list with three objects: 1) "contest" with the pairings

of males, a trial ID and a factor indicating trials with males that had modified left genital lobes, 2)

trial-specific variables "conditions" containing female age, female stock, courtship duration and a

factor indicating which male initiated courtship, 3) male-specific predictor variables

"male_predictors" with left and right genital lobe lengths, male tibia length, sum licking duration,

number of licking sequences and failed mounting attempts (Additional datafile 5). We developed

the statistical model (Additional datafile 1, Tables S5-11) by evaluating the effect of different

variables on copulation success and by comparing pairs of nested models, using Chi-square tests

with the generic anova() function [61].

We tested trial specific variables in interaction with all male-specific variables, but only

found an interaction of female age with failed mounting attempts in the dataset with unmodified

males (Additional datafile 1, Figure S5, Table S8, Table 1). We also tested for interactions between

male specific predictor variables but did not find any. The best fitting model for the dataset with

unmodified males was: copulation success ~ sum licking duration + number of licking sequences +

left_lobe length + failed mounting attempt x female age; and for the trials with lobe modified

males: copulation success ~ sum licking duration + number of licking sequences + left lobe length.

The male specific variables tibia length (body size approximation) and right lobe length were

excluded from the final model.

Spermathecae filling levels were evaluated with a linear model using the lm() function:

spermatheca filling level ~ left lobe length + right lobe length + sum licking duration + Number of

licking sequences + copulation duration + female age + male age, and significance of each variable

was evaluated with an F-test using anova().

Declarations

Ethics approval and consent to participate

"Not applicable" --- This research focuses on invertebrate insects. There are no ethical

considerations mentioned for these species according to EU Directive 86/609-STE123.

Consent for publication

"Not applicable"

Availability of data and material

The data sets supporting the results of this article are available in the DRYAD repository,

https://datadryad.org/stash/share/jzbHRTO70QV55ejT_BxIcopEoG-wVDjRI_HpQK0ergY.

Competing interests

The authors declare no competing financial interests

Funding

This study was supported by the CNRS and by a grant of the European Research Council

under the European Community's Seventh Framework Program (FP7/2007-2013 Grant Agreement

no. 337579) given to VCO. BL was supported by a pre-doctoral fellowship Sorbonne Paris Cité of

the Université Paris 7 Denis Diderot.

Authors' contributions

ML and BL designed the experiments. BL and DC recorded fly copulation and performed

light microscopy analysis of *Drosophila* male genitalia. BL and ML analysed the data. ML, VCO

and BL wrote the manuscript. All authors have read and approved the manuscript.

Acknowledgements

We thank Therese A. Markow for discussions. We are very grateful to Alexandre Peluffo for

help with the statistical analysis, Claus Lang for technical support and Juliette Royer for dissections

of *D. pachea* male genitalia.

References

1. Bateman AJ. Intra-sexual selection in Drosophila. Heredity. 1948;2:349–68.

2. Fisher RA. The Genetical Theory of Natural Selection. Oxford: Clarendon Press. 272 p. 1930.

3. Lande R. Models of speciation by sexual selection on polygenic traits. Proceedings of the
National Academy of Sciences. 1981;78:3721–5. doi:10.1073/pnas.78.6.3721.

4. Bonduriansky R. The evolution of male mate choice in insects: a synthesis of ideas and evidence.
Biological Reviews. 2001;76:305–39. doi:10.1017/S1464793101005693.

5. Leonard J, Córdoba-Aguilar A. The evolution of primary sexual characters in animals. Oxford
University Press; 2010.

6. Edward DA, Chapman T. The evolution and significance of male mate choice. Trends in Ecology
& Evolution. 2011;26:647–54. doi:https://doi.org/10.1016/j.tree.2011.07.012.

7. Kokko H, Jennions MD. The Relationship between Sexual Selection and Sexual Conflict. Cold
Spring Harbor Perspectives in Biology . 2014;6. doi:10.1101/cshperspect.a017517.

8. Eberhard WG. Sexual selection and animal genitalia. Harvard University Press; 1985.

9. Birkhead TR, Pizzari T. Postcopulatory sexual selection. Nat Rev Genet. 2002;3.
doi:10.1038/nrg774.

10. Thornhill R. Cryptic Female Choice and Its Implications in the Scorpionfly Harpobittacus
nigriceps. The American Naturalist. 1983;122:765–88. doi:10.1086/284170.

11. Eberhard WG. Evolution of genitalia: theories, evidence, and new directions. Genetica.
2010;138:5–18. http://www.springerlink.com/content/h375483656l11773/export-citation/.

12. Arnqvist G. Comparative evidence for the evolution of genitalia by sexual selection. Nature.
1998;393:784–6. doi:10.1038/31689.

13. Chapman T, Arnqvist G, Bangham J, Rowe L. Sexual conflict. Trends Ecol Evol. 2003;18:41–7.
doi:10.1016/S0169-5347(02)00004-6.

14. Eberhard WG. Copulatory Courtship and Cryptic Female Choice in Insects. Biological Reviews.
1991;66:1–31. doi:10.1111/j.1469-185X.1991.tb01133.x.

15. Eberhard WG. Evidence for widespread courtship during copulation in 131 species of insects
and spiders, and implications for cryptic female choice. Evolution. 1994;48:711–33.
doi:10.1111/j.1558-5646.1994.tb01356.x.

16. Ghiselin MT. The distinction between primary and secondary sexual characters. In: Leonard JL,
Córdoba-Aguilar A, editors. The Evolution of Primary Sexual Characters in Animals. New York:
Oxford University Press; 2010. p. 9–14.

17. Andersson M. Female choice selects for extreme tail length in a widowbird. Nature.
1982;299:818–20. doi:10.1038/299818a0.

18. von Schantz T, Göransson G, Andersson G, Fröberg I, Grahn M, Helgée A, et al. Female choice
selects for a viability-based male trait in pheasants. Nature. 1989;337:166–9.
doi:10.1038/337166a0.

19. Hill GE. Plumage coloration is a sexually selected indicator of male quality. Nature.
1991;350:337–9. doi:10.1038/350337a0.

20. Norris K. Heritable variation in a plumage indicator of viability in male great tits Parus major.
Nature. 1993;362:537–9. doi:10.1038/362537a0.

21. Petrie M. Improved growth and survival of offspring of peacocks with more elaborate trains.
Nature. 1994;371:598–9. doi:10.1038/371598a0.

22. Hasselquist D, Bensch S, von Schantz T. Correlation between male song repertoire, extra-pair
paternity and offspring survival in the great reed warbler. Nature. 1996;381:229–32.
doi:10.1038/381229a0.

23. Welch AM, Semlitsch RD, Gerhardt HC. Call Duration as an Indicator of Genetic Quality in
Male Gray Tree Frogs. Science. 1998;280:1928 LP – 1930. doi:10.1126/science.280.5371.1928.

24. Birch M. Pre-courtship use of abdominal brushes by the nocturnal moth, Phlogophora
meticulosa (L.) (Lepidoptera: Noctuidae). Animal Behaviour. 1970;18:310–6.
doi:https://doi.org/10.1016/S0003-3472(70)80043-4.

25. Sreng L. Cockroach mating behaviors, sex pheromones, and abdominal glands (Dictyoptera:
Blaberidae). Journal of Insect Behavior. 1993;6:715–35. doi:10.1007/BF01201672.

26. Polak M, Rashed A. Microscale laser surgery reveals adaptive function of male intromittent
genitalia. Proc R Soc B. 2010;277. doi:10.1098/rspb.2009.1720.

27. Grieshop K, Polak M. The precopulatory function of male genital spines in Drosophila
ananassae [Doleschall] (Diptera: Drosophilidae) revealed by laser surgery. Evolution. 2012;66.
doi:10.1111/j.1558-5646.2012.01638.x.

28. Bertin A, Fairbairn DJ. One tool, many uses: precopulatory sexual selection on genital
morphology in Aquarius remigis. Journal of Evolutionary Biology. 2005;18:949–61.
doi:10.1111/j.1420-9101.2005.00913.x.

29. Ploog DW, MacLean PD. Display of penile erection in squirrel monkey (Sainuri sciureus).
Animal Behaviour. 1963;11:32–9. doi:https://doi.org/10.1016/0003-3472(63)90005-8.

30. Wickler W. Ursprung und biologische Deutung des Genitalprasentierens männlicher Primaten.
Zeitschrift für Tierpsychologie. 1966;23:422–37. doi:10.1111/j.1439-0310.1966.tb01605.x.

31. Miller EH. Genitalic traits of mammals. In: Leonard J, Córdoba-Aguilar A, editors. The
evolution of primary sexual characters in animals. New York: Oxford University Press; 2010. p.
471–93.

32. Bohme W. The Tucano Indians of Colombia and the Iguanid Lizard Plica plica: Ethnological,
Herpetological and Ethological Implications. Biotropica. 1983;15:148–50. doi:10.2307/2387961.

33. Mautz BS, Wong BBM, Peters RA, Jennions MD. Penis size interacts with body shape and
height to influence male attractiveness. Proceedings of the National Academy of Sciences.
2013;110:6925–30. doi:10.1073/pnas.1219361110.

34. Estes RD. The significance of horns and other male secondary sexual characters in female
bovids. Applied Animal Behaviour Science. 1991;29:403–51. doi:https://doi.org/10.1016/0168-
1591(91)90264-X.

35. Muller MN, Wrangham R. Sexual Mimicry in Hyenas. The Quarterly Review of Biology.
2002;77:3–16. doi:10.1086/339199.

36. Kahn AT, Mautz B, Jennions MD. Females prefer to associate with males with longer
intromittent organs in mosquitofish. Biology Letters. 2010;6:55–8. doi:10.1098/rsbl.2009.0637.

37. Langerhans RB, Layman CA, DeWitt TJ. Male genital size reflects a tradeoff between attracting
mates and avoiding predators in two live-bearing fish species. Proceedings of the National Academy
of Sciences. 2005;102:7618–23. doi:10.1073/pnas.0500935102.

38. Moreno-García M, Cordero C. On the function of male genital claspers in Stenomacra
marginella (Heteroptera: Largidae). Journal of Ethology. 2008;26:255–60. doi:10.1007/s10164-007-
0058-8.

39. Michiels NK. Mating Conflicts and Sperm Competition in Simultaneous. In: Birkhead TR,
Møller AP, editors. Sperm competition and sexual selection. New York: Academic Press; 1998. p.
219–254.

40. Pitnick S, Heed W. New Species of Cactus-Breeding Drosophila (Diptera: Drosophilidae) in the
Nannoptera Species Group. Annals of the Entomological Society of America. 1994;87:307–10.
http://www.nature.com/nature/journal/v440/n7085/full/nature04623.html.

41. Lang M, Orgogozo V. Distinct copulation positions in Drosophila pachea males with symmetric
or asymmetric: External genitalia. Contributions to Zoology. 2012;81:87–94.

42. Acurio A, Rhebergen FT, Paulus SE, Courtier-Orgogozo V, Lang M. Repeated evolution of
asymmetric genitalia and right-sided mating behavior in the Drosophila nannoptera species group.
BMC Evolutionary Biology. 2019;19:109. doi:10.1186/s12862-019-1434-z.

43. Lang M, Polihronakis Richmond M, Acurio AE, Markow TA, Orgogozo V. Radiation of the
Drosophila nannoptera species group in Mexico. Journal of Evolutionary Biology. 2014;27:575–84.
doi:10.1111/jeb.12325.

44. Rhebergen FT, Courtier-Orgogozo V, Dumont J, Schilthuizen M, Lang M. Drosophila pachea
asymmetric lobes are part of a grasping device and stabilize one-sided mating. BMC Evolutionary
Biology. 2016;16:176. doi:10.1186/s12862-016-0747-4.

45. Pitnick S, Markow TA, Spicer GS. Delayed male maturity is a cost of producing large sperm in
Drosophila. Proceedings of the National Academy of Sciences. 1995;92:10614–8.
http://www.pnas.org/content/92/23/10614.abstract.

46. Pitnick S, Markow T, Spicer GS. Evolution of Multiple Kinds of Female Sperm-Storage Organs
in Drosophila. Evolution. 1999;53:1804–22. doi:10.2307/2640442.

47. Pitnick S, Markow TA. Male gametic strategies: sperm size, testes size, and the allocation of
ejaculate among successive mates by the sperm-limited fly Drosophila pachea and its relatives. The
American Naturalist. 1994;143:785–819.

48. Bradley RA, Terry ME. Rank Analysis of Incomplete Block Designs: I. The Method of Paired
Comparisons. Biometrika. 1952;39 3/4:324–45. doi:10.2307/2334029.

49. Pitnick S. Operational sex ratios and sperm limitation in populations of Drosophila pachea.
Behavioral Ecology and Sociobiology. 1993;33:383–91.

50. Byers J, Hebets E, Podos J. Female mate choice based upon male motor performance. Animal
Behaviour. 2010;79:771–8. doi:https://doi.org/10.1016/j.anbehav.2010.01.009.

51. Kirkpatrick M, Ryan MJ. The evolution of mating preferences and the paradox of the lek.
Nature. 1991;350:33–8. doi:10.1038/350033a0.

52. Grieshop K, Polak M. Evaluating the post-copulatory sexual selection hypothesis for genital
evolution reveals evidence for pleiotropic harm exerted by the male genital spines of Drosophila
ananassae. J Evol Biol. 2014;27. doi:10.1111/jeb.12524.

53. Hoikkala A, Aspi J, Suvanto L. Male courtship song frequency as an indicator of male genetic
quality in an insect species, Drosophila montana. Proceedings of the Royal Society of London
Series B: Biological Sciences. 1998;265:503–8. doi:10.1098/rspb.1998.0323.

54. Pitnick SS. Sexual selection and sperm production in Drosophila. Thesis (Ph. D.)--Arizona State
University, 1992. ; 1992.

55. Jefferson M. Breeding biology of Drosophila pachea and its relatives. 1977.

56. Heed WB, Kircher HW. Unique Sterol in the Ecology and Nutrition of Drosophila pachea.
Science. 1965;149:758–61. doi:10.1126/science.149.3685.758.

57. Lang M, Murat S, Clark AG, Gouppil G, Blais C, Matzkin LM, et al. Mutations in the neverland
gene turned Drosophila pachea into an obligate specialist species. Science. 2012;337:1658–61.

58. Warren JT, Wismar J, Subrahmanyam B, Gilbert LI. Woc (without children) gene control of
ecdysone biosynthesis in Drosophila melanogaster. Molecular and cellular endocrinology.
2001;181:1–14.

59. Spieth HT. Mating behavior within the genus Drosophila (Diptera). Bull Am Museum Nat Hist.
1952;99.

60. R Core Team. R: A language and environment for statistical computing. R Foundation for
Statistical Computing, Vienna, Austria. 2013. 2014.

61. Turner H, Firth D. Bradley-Terry Models in R: The BradleyTerry2 Package. Journal of
Statistical Software, Articles. 2012;48:1–21. doi:10.18637/jss.v048.i09.

Supporting Information

Additional datafile 1: Supplementary Figures and Tables, **Figure S1:** Camera device for movie recording, **Figure S2:** Pairwise lobe length differences of males used in the competition mating experiment, **Figure S3:** Courtship and copulation duration in competition mating experiments **Figure S4:** Courtship duration increases with female age, **Figure S5:** Failed copulation attempts compared to female age, **Table S1:** *Drosophila pachea* resources, **Table S2:** Competition mating trials used for courtship analysis, **Table S3:** Courtship duration, licking index and copulation duration in trials where both males courted the female, **Table S4:** Genital lobe lengths in *D. pachea* stocks.

Additional datafile 2: Measurements of mating behaviour, genital lobe lengths and spermathecae filling levels of individuals used in the competition mating experiments.

Additional datafile 3: Raw measurements of lobe lengths in different *D. pachea* stocks

Additional datafile 4: Dataset for male licking behavior and male mounting.

Additional datafile 5: R list object "PacData", dataset for Bradley-Terry model fit

Tables

Table 1: Bradley–Terry (BT) model examining the effects of left lobe length, sum licking duration, number of licking sequences and failed mounting attempts on copulation success, in 92 trials with non-lobe modified males. The main effect for "female age" cannot be estimated in the BT model because it is specific to each trial and not to each male. Bolded estimates are significant.

variable	estimate	std. error	z value	P (>|z|)
sum licking duration	1.224514	0.334856	3.657	**0.0003**
number of licking sequences	-0.259216	0.127859	-2.027	**0.0426**
left lobe	0.007471	0.003582	2.086	**0.0370**
failed mounting attempts	-4.493660	1.834694	-2.449	**0.0143**
female age	NA	NA	NA	NA
failed mounting attempts x female age	0.295355	0.119397	2.474	**0.01337**

Table 2: Bradley–Terry (BT) model examining the effects of left lobe length, sum licking duration, and number of licking sequences on copulation success, in 53 trials with males that had surgically modified left genital lobes. Bolded estimates are significant.

variable	estimate	std. error	z value	P (>|z|)
sum licking duration	0.561757	0.256917	2.187	**0.0288**
number of licking sequences	-0.176035	0.090341	-1.949	0.0513
left lobe	0.009199	0.005090	1.807	0.0707

Table 3: Linear model, examining the effects of lobe length, sum licking duration, copulation duration and age on spermathecae filling levels. degrees of freedom: numerator / denominator 38

variable	F value	P
sum licking duration	1.6841	0.2022
number of licking sequences	0.1107	0.7411
left lobe length	0.1139	0.7377
right lobe length	1.4739	0.2322
copulation duration	10.94	**0.0021**
female age	0.5036	0.4822
male age	22.828	**0.00003**

Figure Legends

Figure 1: *Drosophila pachea* male genital lobes. (a) Male of stock 15090-1698.02 in ventral view and with asymmetric lobes. (b) Male of the selection stock with apparently symmetric lobes. The scale bars correspond to 200 μm. (c) Posterior view of a dissected male terminalia of stock 15090-1698.02 with asymmetric lobes. (d) Posterior view of a dissected male terminalia of the selection stock; the scale bars correspond to 100 μm, dots indicate lobe length measurement points.

Figure 2: Genital lobe lengths differ between *D. pachea* stocks. Lengths of the left and right epandrial lobe lobes are presented for (a) stock 15090-1698.01, (b) stock 15090-1698.02, and (c) the selection stock. Sibling males of those used in the mating experiment are shown in panel (c). Each point represents one male. The variance of left lobe length is increased in the selection stock

compared to the source stock 15090-1698.02 (Levene's test: selection stock / 15090-1698.02: DF =

1/147, F = 11.506, P = 0.0009), while the variance of the right lobe length were not significantly

different (Levene's test, DF = 1/147, F = 0.5164, P = 0.4735).

Figure 3: Courtship durations and relative courtship activity. (a) The ratio of sum licking

duration and total courtship duration in each trial shows that the copulating male (cop, grey box-

plots) had a higher average sum licking duration compared to the non-copulating male (non-cop,

white box-plots). Each point represents one male. (b) The total courtship duration was shorter in

trials with only one male displaying courtship signs (one, white box-plots) compared to trials with

both males courting simultaneously (both, grey box-plots) (Mann-Whitney-Wilcoxon test, female

stock 15090-1698.01, non modified males, W = 385, N = 25/51, P = 0.0053 ; females of stock

15090-1698.02, W = 219, N = 16/60, P = 0.0009 ; females stock 15090-1698.01, modified males, W

= 62.5, N = 7/54, P = 0.0044). Horizontal bars indicate pairwise comparisons with P-values ** <

0.01 or *** < 0.001. Each point represents one mating experiment.

Figure 1

Figure 2

Figure 3