Essential Metabolic Routes as a Way to ESKAPE from Antibiotic Resistance

Angélica Luana C. Barra[1], Lívia de Oliveira C. Dantas[1], Luana Galvão Morão[1], Raíssa F.
Gutierrez[1], Igor Polikarpov[1], Carsten Wrenger[2*] and Alessandro S. Nascimento[1*]

[1] São Carlos Institute of Physics, University of Sao Paulo. Av. Trabalhador São-carlense, nº 400, São
Carlos, SP. Brazil.

[2] Department of Parasitology, Institute of Biomedical Sciences, University of Sao Paulo. Av. Prof.
Lineu Prestes, 1374 – Cidade Universitária, São Paulo, SP, Brazil.

* Correspondence:
Corresponding Authors
cwrenger@icb.usp.br (CW) or asnascimento@ifsc.usp.br (ASN)

Keywords: ESKAPE pathogens, thiamine, pyridoxal 5'-phosphate, antibiotic resistance.

Abstract

The antibiotic resistance is a worldwide concern that requires a concerted action from physicians,
patients, governmental agencies and academia to prevent infections and the spread of resistance, to
track resistant bacteria, to improve the use of current antibiotics and to develop new antibiotics.
Despite the efforts spent so far, the current antibiotics in the market are restricted to only five general
targets/pathways highlighting the need for basic research focusing on the discovery and evaluation of
new potential targets. Here we interrogate two biosynthetic pathways as potentially druggable
pathways in bacteria. The biosynthesis pathway for thiamine (vitamin B1), absent in humans, but
found in many bacteria, including organisms in the group of the ESKAPE pathogens (*Enterococcus*
faecium, *Staphylococcus aureus*, *Klebsiella pneumoniae*, *Acinetobacter baumanii*, *Pseudomonas*
aeruginosa and *Enterobacter* species) and the biosynthesis pathway for pyridoxal 5'-phosphate and
its vitamers (vitamin B6), found in *S. aureus*. Using current genomic data, we discuss the
possibilities of inhibition of enzymes in the pathway and review the current state of the art in the
scientific literature.

1   Introduction

Antibiotic resistance is an urgent threat to human health and requires urgent actions from physicians, patients, industries, governmental agencies and the academic community worldwide. According to the last document from the Centers for Disease Control and Prevention (CDC) regarding antibiotic resistance in the United States, from 2013, the number of people with serious infections caused by resistant bacteria reaches two million every year, with at least 23,000 deaths per year directly caused by these infections (1). The situation is similarly warning in Europe, where 25,100 deaths were reported from the European Centre for Disease Prevention and Control in 2007 (2). Globally, 700,000 deaths are estimated every year as a consequence of antibiotic resistance (3). The same CDC document lists four general action lines to address antibiotic resistance: (*i*) preventing infections and the spread of resistance; (*ii*) tracking resistant bacteria; (*iii*) improving the use of current antibiotics; (*iv*) developing of new antibiotics (1).

Since bacteria have a short doubling time and efficient mechanisms for plasmid sharing, the development of antibiotic resistance is a very efficient defense mechanism. So, as previously said by Walsh and Wencewicz, the development of resistance is not a matter of *if*, but rather a matter of *when* (4) and, despite the title of this paper (which is rather provocative), there is no way we can escape from it (5). On the contrary, the development/discovery of new antibiotics will tend to be a continuous goal to be achieved in drug discovery pipelines.

Interestingly, even after what has been named as the 'golden age' of antibiotics development, the drugs currently in the market are restricted to only five molecular targets and/or pathways (4): (*i*) the peptidoglycan/cell wall biosynthesis, site of action of beta-lactam antibiotics, for example; (*ii*) the protein biosynthesis, where the ribosome is an important target; (*iii*) DNA replication and RNA transcription; (*iv*) the folate biosynthesis pathway, and (*v*) the disruption of the bacterial membrane. Given the relevance of the antibiotic therapy and the emergence of antibiotic resistance, the available choices of current drugs are very narrow in the context of the mechanism of action restricted to only five molecular/pathway targets.

Antibiotic resistance, as a threat to human health, should be addressed in multiple, and simultaneous ways. For academia, an interesting way to address antibiotic resistance is the discovery and validation of new targets/pathways that could be specifically targeted by new antibiotic candidates. It is worthy of note that, according to Kelly and Davies (3), no new class of antibiotics was discovered and released for routine treatment since the 1980s, highlighting the outstanding role that academia can have in the preliminary research for discovery and validation of druggable targets/pathways.

About 7% of the *E. coli* genome, 303 genes, were shown to be composed of essential genes (5,6). Under stress conditions caused by a limited medium, additional 119 genes show some condition-dependent essentiality, including several genes related to the metabolism and synthesis of essential molecules such as amino acids, and nucleotides (5) and, obviously, some of these targets/pathways can be an interesting choices for the development of new antibiotic candidates. A promising approach to screen compounds for their activity in metabolic pathways was shown by Zlitni and coworkers (7). The authors searched for antibacterial compounds under poor nutrient media and found three potential compounds in this screening strategy (7). Worth of note, the metabolic profile of existing antibiotics showed that the supplementation of vitamins B5, B6, B1, and B2 did not significantly reverse the antimicrobial effect of any of the 24 inhibitors assayed (7), suggesting that the current antibiotics do not explore these essential pathways.

In this context, the enzymatic routes for the biosynthesis of vitamins are very interesting
pathways to be explored as potential targets for the discovery of new antibiotic candidates. Vitamin
B1, for example, cannot be synthesized by humans, although several microorganisms can synthesize
this vitamin, including pathogens. In the absence of thiamin (vitamin B1), the activity of several
carbohydrate metabolism enzymes is impaired, including pyruvate dehydrogenase, which connects
glycolysis and the citric acid cycle (8). Other thiamin dependent enzymes are transketolase, α-
ketoacid decarboxylase, α-ketoacid dehydrogenase and acetolactate synthase (9). A very similar
scenario was observed for pyridoxal 5'-phosphate (vitamin B6) in the model Gram-positive organism
Bacillus subtilis (10).

The pathways involved in the synthesis of vitamins, in particular, vitamin B1 (thiamin) and
B6 (pyridoxal), seem to be of great relevance, since they are involved in central processes in the
metabolism of carbohydrates and amino acids and the corresponding enzymes are found in most
bacteria, fungi and plants but not in humans (8), favoring the development of specific drugs with
minimal side effects due to the interaction with the host. However, a few questions still stand: (i) how
feasible are the targets involved in the pathways for the biosynthesis of thiamin and pyridoxal for the
microorganisms with a higher emergency in terms of resistance, or ESKAPE: *Enterococcus faecium,*
Staphylococcus aureus, Klebsiella pneumoniae, Acinetobacter baumanii, Pseudomonas aeruginosa
and *Enterobacter* species? (ii) What enzymes are present and what do we know about them?

Here we used the available genomic and proteomic data to address these questions focusing
on the ESKAPE pathogens. Vitamins B1 and B6 were chosen for this analysis, since the biosynthetic
pathways for these enzymes have already been validated as molecular targets for some human
pathogens, such as thiamin for *Plasmodium falciparum* (11), or pyridoxal for *P. falciparum* (12) or
Trypanosoma brucei (13), for example.

95   **2   Methods**

The analyses provided here are the result of the interrogation of the enzymes in the
biosynthesis pathways for thiamin and pyridoxine phosphate or pyridoxal phosphate using the KEGG
database (14,15). For this purpose, the KEGG pathway for thiamin metabolism (map 00730) was
listed for the ESKAPE pathogens using, whenever possible, commercial strains rather than specific
or antibiotic-resistant strains. Briefly, for each pathogen (*Enterococcus faecium*, *Staphylococcus*
aureus, *Klebsiella pneumoniae*, *Acinetobacter baumanii*, *Pseudomonas aeruginosa* and
Enterobacter) a KEGG search for the maps of thiamin and pyridoxal metabolism was done, looking
for the existing genes in commercial strains for each of the ESKAPE organisms. For the sake of
comparison, some additional resistant strains were listed, with no differences in the observed genes
for the pathways under study. A similar search was carried out for the vitamin B6 metabolism (map
00750) in KEGG, interrogating the existing enzymes for the ESKAPE pathogens in comparison with
humans. The analysis was focused in the enzymes in the biosynthesis of B1 and B6 vitamin in an
attempt to identify the most promising targets for future medicinal chemistry studies. Finally, a table
listing the existing genes for each pathogen was compiled and is presented in the following sections.

Using the existing literature, we sought to identify whenever possible the cases of enzymes
with functional redundancy with another enzyme for the same organism. The identified cases were
discussed as less promising targets for medicinal chemistry campaigns. Additionally, the existing

data for enzyme inhibition using natural or synthetic compounds was compiled to provide initial
proof-of-concept clues about the *druggability* of the identified promising targets.

When necessary, the sequence of an enzyme of the pathway was used to search for homologues using
BLAST (16) searches against the PDB (17) or the non-redundant database of proteins using BLAST
default parameters, i.e., minimum expected threshold of 10, a word size of 6 and the BLOSUM62
substitution matrix.

3   Results and Discussion

3.1   Thiamin Biosynthesis

The biosynthesis pathway for thiamin involves two branches, as shown in Figure 1. In
summary, thiamin is synthesized from 4-amino-5-hydroxymethyl-2- methylpyrimidine (HMP,
superior branch in Figure 1) and 5-(2-hydroxyethyl)-4-methylthiazole (THZ, inferior branch). Both
compounds are phosphorylated by ThiD and ThiM, respectively. Finally, the enzyme ThiE, central to
the pathway, is responsible to synthesize thiamine phosphate (TMP) by merging the two branches
(18). Thiamin phosphate can be dephosphorylated to thiamine and then pyrophosphorylated to
thiamin diphosphate (TPP) by TPK. Alternatively, thiamin can be degraded to HMP and THZ by
TenA, a thiaminase II enzyme.

According to the KEGG pathway database, humans lack many of the enzymes of this
pathway, but not all of them. In humans, a TPK (TPK1) enzyme is found, with UNIPROT ID
Q9H3S4. Other than that, the remaining enzymes in the biosynthetic pathway for thiamin are missing
in humans, making them very attractive for the design of chemical probes that could be used as
proof-of-concept compounds.

The analysis of the thiamin pathway for *Enterococcus faecium* (ATCC 8459) in the KEGG
database did not identify genes for ThiE, ThiM and TenA. For this species, only ThiD and TPK were
positively identified. However, a BLAST search for TenA homologous within the non-redundant
database restricted to *Enterococcus faecium* (TAXID 1352) identified a single result for thiaminase II
(GenBank: SAM74984.1), four results for ThiE (SAZ10134.1, WP_086323306.1, WP_010732763.1
and WP_072538983.1) and two results for ThiM (SAZ10238.1, EJY48288.1). The small number of
hits in the BLAST search may suggest some issues in the annotation.

Interestingly, when the *Enterococcus faecalis* thiamin pathway is compared with the pathway
observed for *E. faecium*, many differences are found. The KEGG pathway for *E. faecalis* (ATCC
29212) shows that the entire pathway, as described in Figure 1 is found: ThiD, ThiM, ThiE, TPK and
TenA, in contrast with *E. faecium*. Additionally, there is no change in the repertoire of enzymes in
the pathway for the Vancomycin-resistant strain V583 as compared to the ATCC 29121 strain.

For *Staphylococcus aureus* (NCTC 8325) and *Klebsiella pneumoniae* (subsp. pneumoniae
ATCC 43816 KPPR1), the KEGG pathway indicates that all the enzymes in the pathway are
observed, with no changes to an *S. aureus* resistant strain such as COL (MRSA) as compared to the
NCTC 8325. For *Acinetobacter baumannii* (ATCC 14978), all enzymes are observed but TPK.
Instead, thiamin phosphate may be converted directly to thiamin diphosphate by a thiamin-
monophosphate kinase, and then converted to thiamin triphosphate by an adenylate kinase. In the

case of *Pseudomonas aeruginosa* (NCGM 1900), some enzymes are missing: ThiD, TenA, TPK.
Finally, *Enterobacter sp.* (638) has ThiD, ThiE, ThiM and misses TenA and TPK.

The overall panel of enzymes for the thiamin pathway for the ESKAPE pathogens is
summarized in Table 1.

TenA was shown to play a dual role in thiamine synthesis and salvage (19): beyond its
function to hydrolyze thiamin, TenA also deaminates aminopyrimidine to form 4-amino-5-
hydroxymethyl-2-methylpyrimidine (HMP), with the latter activity being about 100 times faster than
the former (19,20). In some organisms, such as *Bacillus subtilis*, the deaminase activity is somewhat
redundant with the ThiC (ThiA) activity, since both activities provide HMP or HMP phosphate, that
can be further phosphorylated by ThiD (9).

In the other branch of the thiamin biosynthesis pathway, thiazole phosphate can be generated
from thiazole alcohol incorporated from the medium and phosphorylated by ThiM or it can be
synthesized in a series of enzymatically catalyzed reactions using amino acids such as glycine or
tyrosine as substrates (9). In this context, according to Begley and coworkers (9), lesions in the ThiM
gene were observed to be prototrophic. In contrast, selective mutations of ThiD and ThiE led to the
requirement of externally provided thiamin (9).

ThiE is central to the pathway since the enzyme merges the two branches to finally synthesize
thiamin phosphate (Figure 1). It is important to highlight that many of the mutational analysis was
carried out in *E. coli* and, as Table 1 shows, there is significant variance among bacterial species.

Taking together, the analysis of the literature suggests that ThiD and ThiE may represent
interesting targets for the development of chemical probes to further evaluate their function. TPK is
expressed in humans and the inhibition of this enzyme could lead to harmful effects on the human
host. TenA inhibition could be overcome by the somewhat functional redundancy with thiC (thiA)
and ThiM function can be dispensable since thiazole phosphate can be synthesized *de novo*.

ThiD and ThiE are found in most ESKAPE organisms. Exceptions are *P. aeruginosa*, which
lacks ThiD and *E. faecium* that lacks ThiE, according to KEGG data. For ThiE, in particular, it was
reported that 4-amino-2-trifluoromethyl-5-hydromethylpyrimidine (CF$_3$-HMP), an HMP analogue,
can be converted by ThiD to CF$_3$-HMP pyrophosphate, which in turn inhibits ThiE (8,21). This
enzymatic inhibition culminated with the inhibition of *E. coli* growth (8), suggesting that this strategy
might be promising for the development of new inhibitors.

In terms of the structural biology, some crystal structures of ThiD are available by the time of
the writing of this paper, including the enzymes from *Salmonella enterica* (PDB ID 1JXH (22)),
Clostridioides difficile (PDB ID 4JJP), *B. subtilis* (PDB ID 2I5B (23)), *Thermus thermophilus* (PDB
ID 1UB0), *A. baumannii* (PDB ID 4YL5), *Bacteroides thetaiotaomicron* (PDB ID 3MBH) and the
bifunctional enzyme from *Saccharomyces cerevisiae* (PDB ID 3RM5). Typically, ThiDs are folded
as typical ribokinases, with a central 8-stranded β-sheet surrounded by 8 α-helices and structural
studies suggest that some surface loops may have a structural change based on the presence of the
nucleotide (23).

Interestingly, the *S. aureus* pyridoxal kinase enzyme (*Sa*PdxK) has a dual role,
phosphorylating pyridoxal and pyridoxine in the pyridoxal *de novo* biosynthesis pathway as well as
HMP in the thiamin biosynthesis pathway, with a K$_M$ almost 20 times greater for HMP than for

pyridoxal and k_{cat} values 3 times faster for pyridoxal (24). So, SaPdxK has some redundancy with *S.*
aureus ThiD (SaThiD), with less efficiency, though.

*Sa*ThiD was shown to be potentially inhibited by Rugulactone (Ru0), a natural product as
well as by its derivatives Ru1 and Ru2, with IC_{50} values ranging from 14 to 32 μM (25). In the
absence of thiamine in the medium, the MIC observed for Ru1 was four times lower than in the
presence of thiamine for *L. monocytogenes* (25), suggesting that the inhibitory effect of Rugulactone
is due to ThiD inhibition, although Rugulactone also inhibits other kinases.

For ThiE, the crystal structures of a few enzymes are available, including the enzyme from
Pyrococcus furiosus (PDB ID 1XI3), *B. subtilis* (PDB IDs 1G4T, 3O15, 3O16, 1G69, 1G4E, 1G67,
1G4P, 1YAD (26)), *Mycobacterium tuberculosis* (3O63) and for the bifunctional enzyme from
Candida glabrata (PDB IDs 3NL2 and 3NM1 (27)). No crystal structure of an ESKAPE pathogen
ThiE is available to date. From the structural point of view, ThiE is typically folded as an α/β TIM
barrel, with thiamine binding at the top of the barrel, as elegantly shown in the set of crystal
structures determined for the *B. subtilis* enzyme (28).

At this point, an important question has to be addressed: can the blockade of thiamine *de novo*
biosynthesis pathway be overcome by internalization of the vitamin? In principle, several
microorganisms can import thiamine from the environment (29). So, how effective can be the
inhibition of ThiD and ThiE particularly for pathogenic bacteria? This question can be addressed in
several levels. On the first level, the essentiality of ThiD, for example, has been shown some
pathogenic microorganisms (23), such as *Streptococcus pneumoniae* (30), *Haemophilus influenzae*
(31) and *Mycobacterium tuberculosis* (32). This is not the case for other bacteria. For *B. subtilis*, for
example, ThiD gene was shown to be dispensable (23). Whether ThiD and ThiE are essential for the
ESKAPE pathogens is still an open question to be addressed by further investigation. On a second
level, Nodwell and coworkers showed that *Listeria monocytogenes*, which is known to uptake
thiamine from the environment, is still sensible to Rugulactone and its derivatives Ru1 and Ru2 (25).
When grown in a chemical defined media without thiamine, the inhibitory effects of Rugulactone are
potentiated with a 4-fold reduction in the MIC (25). However, the inhibitory effect observed even in
the presence of thiamine highlight that the blockade of the *de novo* biosynthesis pathway is still a
promising strategy. As a third level, the natural compound bacimethrin, isolated from *Bacillus*
megaterium and from *Streptomyces albus* is known to be toxic for bacteria with MIC at low
micromolar range (33,34). The mechanism of action of bacimethrin is based on the formation of 2'-
methoxy-thiamine where bacimethrin is used by ThiD instead of HMP. 2'-methoxy-thiamine
pyrophosphate inhibits *E. coli* growth at concentrations 15 times lower than bacimethrin (35).
Interestingly, some *Salmonella enterica* thiD mutants were shown to be bacimethrin resistant (36).
Together, these data suggest that a very promising approach can be devised by the design of suicide
drugs, i.e., compounds that are recognized by the enzymes in the pathway but lead to final
compounds that can't be used as thiamine substituents. Bacimethrin is a good example of a suicide
inhibitor and, although its effects can be reverted by increased thiamine concentrations, the vitamin
concentration that pathogens are exposed are usually defined within a narrow range.

3.2 Pyridoxal Biosynthesis

Pyridoxal, pyridoxine and pyridoxamine, together with their respective phosphate esters
compose the six vitamers for vitamin B6 and can be interconverted (37). There are two biosynthetic
routes for pyridoxal 5'-phosphate: a deoxyxylulose 5-phosphate-dependent pathway found in some
bacteria and a deoxyxylulose 5-phosphate-independent pathway found in all kingdoms (37,38). This
second and widespread pathway, curiously, depends on only two enzymes, as shown in Figure 2.

In this pathway, the enzyme Pdx2 converts L-glutamine into L-glutamate and an ammonium
molecule. The latter is thought to diffuse to Pdx1 active site, which also uses glyceraldehyde 3-
phosphate and D-ribose 5-phosphate to synthesize pyridoxal 5'-phosphate (37,39). Interestingly, to
properly synthesize pyridoxal 5'-phosphate, a complex involving twelve Pdx1 molecules and twelve
Pdx2 molecules has to be assembled (40–42) and this assembly is further stabilized by the interaction
with L-glutamine (43).

Similar to what is observed to the thiamine biosynthesis pathway, humans lack the genes for
Pdx1 and Pdx2, making the development of specific antibiotic candidates an attracting strategy. In
the same direction, for some pathogenic organisms such as *Helicobacter pylori* (44), *Mycobacterium*
tuberculosis (45), and *Streptococcus pneumoniae* (46), the depletion in vitamin B6 resulted in
reduced virulence, indicating that the biosynthesis of this vitamin may be a good strategy for the
design of new antibiotic candidates.

Very interestingly and in contrast to what was observed for the thiamine pathways, from the
ESKAPE pathogens, *S. aureus* is the only microorganism that has the ribose 5-phosphate dependent
pathway for the biosynthesis of pyridoxal 5'-phosphate, as shown in Table 2. A single enzyme (Pdx1
or Pdx2 alone) is never observed for these organisms (Table 2).

A comparison between the pyridoxal/pyridoxine phosphate pathways for *E. faecium* and *E.*
faecalis shows that both pathogens miss the Pdx enzymes. On the other hand, some other pathogenic
organisms, including organisms with high antibiotic resistance rates, have already been identified as
susceptible to modulations of the pyridoxal phosphate biosynthesis pathway. Some organisms which
have been the focus of basic research include *Plasmodium falciparum* and *Plasmodium vivax,* the
malaria pathogens (47–50).

In terms of the structural biology of the enzymes in the vitamin B6 biosynthesis pathway, the
S. aureus Pdx1 enzyme (UNIPROT ID Q2G0Q1) is a close homologue of *Bacillus subtilis* Pdx1
(81% identity, PDB ID 2NV1 (42)), *Geobacillus stearothermophilus* Pdx1 (78% identity, PDB ID
1ZNN (51)) and *Thermus thermophilus* enzyme (67% identity, PDB ID 2ZBT). *S. aureus* Pdx2, has
Geobacillus kaustophilus (PDB ID 4WXY (40)), *Geobacillus stearothermophilus* (PDB ID 1Q7R)
and *B. subtillis* (PDB IDs 2NV0 and 2NV2 (42)) enzymes as its close homologues, with 60%, 60%
and 58% identity in sequence similarity respectively.

In terms of its structure, the Pdx1 enzyme has a typical $(\beta/\alpha)_8$ barrel fold with a central 8-
stranded β-barrel surrounded by 8 α-helices (51). The most impressing feature in the Pdx1 enzyme
structure is its quaternary arrangement, where six Pdx1 enzymes interact with each other to form a
'*donut-like*' arrangement with about 100 Å in diameter. For Pdx2, a Rossman fold is observed with
an α/β/α sandwich topology (42). Curiously, in the active PLP synthase complex, 12 Pdx1 molecules
(a two-layer *donut*) interact with 12 Pdx2 enzymes. However, Pdx2 molecules do not interact with
each other in this complex (42).

Using a structural homology model, Reeksting and coworkers identified some ribose 5'-
phosphate analogues with interesting *in vitro* inhibitory effects on the enzyme Pdx1 from *P.*
falciparum (12). The compounds were identified in a structure-based computational screening
campaign and were shown to have inhibitory effects *in vivo*, as well as the *in vitro* effect. The authors
also showed that the *in vivo* effects could be at least partially suppressed in a mutant strain
overexpressing Pdx1 and Pdx2, in a clear indication that the effect of the analogues was due their
inhibition of the pyridoxal 5'-phosphate biosynthesis pathway (12).

The inhibition of the Pdx2 activity by the glutamine analogue acivicin was also demonstrated
to be a feasible strategy for the inhibition of the vitamin B6 biosynthesis pathway (52). Raschle and
coworkers showed that when inhibited by acivicin, Pdx2 is incapable of interacting with Pdx1, thus
disrupting the pyridoxal 5'-phosphate synthase activity. Interestingly, acivicin is a covalent inhibitor,
that binds to a cysteine residue. So, it seems that multiple strategies may be available for the design
of new binders, including the inhibition of Pdx2 (covalent and non-covalent), the inhibition of Pdx1
and possibly the inhibition of the assembly of the 24-mer complex with Pdx1/Pdx2. Another
possibility to address the inhibition of PLP biosynthesis might be at the level of the Pdx1
transcription, which was shown to be regulated by PdxR (46). In principle, if PdxR action could be
regulated by an exogenous chemical probe, the PLP synthase activity would be regulated as well.
However, a proof-of-concept experimental validation is still necessary.

Finally, a salvage pathway for vitamin B6 is found in many bacteria, as well as in humans. In
E. coli, it involves two enzymes: PdxK and PdxY. These kinases act phosphorylating the vitamers
pyridoxal, pyridoxine and pyridoxamine into their phosphorylated forms (23,37,53). Interestingly, in
some organisms, including pathogenic microorganisms such as *S. aureus*, the PdxK activity is carried
out by the HMP kinase ThiD, showing some functional convergence between the vitamin B6 and B1
pathways (24). Not surprisingly, this enzyme has also been shown to be validated druggable target
for some pathogens, such as *Trypanosoma brucei* for example (13), although specificity may be an
important issue, since humans also have genes for PdxK known to be inhibited by drugs like
theophylline, for example, with known neurotoxic effects (54).

Again, the same question asked for the thiamine biosynthesis pathway can be asked here: how
can the microbial uptake of vitamin B6 overcome the blockade of the biosynthetic pathway? Some
evidences suggest that the blockade might be effective. For example, in *S. pneumoniae*, the deletion
of Pdx1 resulted in defective growth of the bacteria. The growth was shown to be restored under
increased concentrations of vitamin B6 (46). A similar scenario was also observed for *M.*
tuberculosis (45), suggesting that the uptake might not be enough to meet the pathogen's requirement
of the vitamin.

In conclusion, the biosynthesis pathways for vitamin B1 (thiamine) and B6 (pyridoxal 5'-
phosphate) may be interesting molecular targets for the development of new chemical probes aiming
to inhibit the synthesis of these essential cofactors in pathogenic organisms. Model compounds,
mainly based on substrate analogues showed that inhibition of the vitamin B1 biosynthetic enzymes,
ThiD and ThiE, as well as Pdx1, Pdx2 and PdxK in pyridoxal 5'-phosphate biosynthesis pathway
have shown an initial proof of concept for the model and it is up the scientific community and/or
researcher in industry to explore these target further.

4   Conflict of Interest

The authors declare that the research was conducted in the absence of any commercial or financial
relationships that could be construed as a potential conflict of interest.

5   Author Contributions

CW, IP and ASN conceived the project. LGM and RFG analyzed the thiamine pathway, while ALCB
and LOCD analyzed the pyridoxal pathway. All authors wrote and approved the manuscript.

6   Funding

329 **7   Acknowledgments**

330 The authors thank the funding agencies FAPESP and CNPq. We are also indebted to Maria
331 Auxiliadora M. Santos, Lívia Regina M. Margarido, Josimar Sartori and João Fernando Possatto for
332 their technical support. This manuscript has been released as a Pre-Print at BioRxiv (55).

8 References

1. CDC. Antibiotic resistance threats in the USA. (2013)

2. WHO. Prioritization of Pathogens to Guide Discovery, Research and Development of New Antibiotics for Drug-Resistant Bacterial Infections, Including Tuberculosis. (2017) doi:10.1192/bjp.111.479.1009-a

3. Kelly R, Davies SC. Tackling antimicrobial resistance globally. *Med J Aust* (2017) **207**:371-373.e1. doi:10.5694/mja17.00865

4. Walsh CT, Wencewicz T a. Prospects for new antibiotics: a molecule-centered perspective. *J Antibiot (Tokyo)* (2013) **67**:7–22. doi:10.1038/ja.2013.49

5. Brown ED, Wright GD. Antibacterial drug discovery in the resistance era. *Nature* (2016) **529**:336–343. doi:10.1038/nature17042

6. Baba T, Ara T, Hasegawa M, Takai Y, Okumura Y, Baba M, Datsenko KA, Tomita M, Wanner BL, Mori H. Construction of Escherichia coli K-12 in-frame, single-gene knockout mutants: The Keio collection. *Mol Syst Biol* (2006) doi:10.1038/msb4100050

7. Zlitni S, Ferruccio LF, Brown ED. Metabolic suppression identifies new antibacterial inhibitors under nutrient limitation. *Nat Chem Biol* (2013) **9**:796–804. doi:10.1038/nchembio.1361

8. Du Q, Wang H, Xie J. Thiamin (vitamin B1) biosynthesis and regulation: A rich source of antimicrobial drug targets? *Int J Biol Sci* (2011) **7**:41–52. doi:10.7150/ijbs.7.41

9. Begley TP, Downs DM, Ealick SE, McLafferty FW, Van Loon APGM, Taylor S, Campobasso N, Chiu HJ, Kinsland C, Reddick JJ, et al. Thiamin biosynthesis in prokaryotes. *Arch Microbiol* (1999) **171**:293–300. doi:10.1007/s002030050713

10. Richts B, Rosenberg J, Commichau FM. A survey of pyridoxal 5'-phosphate-dependent proteins in the gram-positive model bacterium Bacillus subtilis. *Front Mol Biosci* (2019) **6**: doi:10.3389/fmolb.2019.00032

11. Chan XWA, Wrenger C, Stahl K, Bergmann B, Winterberg M, Müller IB, Saliba KJ. Chemical and genetic validation of thiamine utilization as an antimalarial drug target. *Nat Commun* (2013) **4**: doi:10.1038/ncomms3060

12. Reeksting SB, Müller IB, Burger PB, Burgos ES, Salmon L, Louw AI, Birkholtz LM, Wrenger C. Exploring inhibition of Pdx1, a component of the PLP synthase complex of the human malaria parasite Plasmodium falciparum. *Biochem J* (2013) **449**:175–187. doi:10.1042/BJ20120925

13. Jones DC, Alphey MS, Wyllie S, Fairlamb AH. Chemical, genetic and structural assessment of pyridoxal kinase as a drug target in the African trypanosome. *Mol Microbiol* (2012) **86**:51–64. doi:10.1111/j.1365-2958.2012.08189.x

14. Kanehisa M. KEGG: Kyoto Encyclopedia of Genes and Genomes. *Nucleic Acids Res* (2000)
28:27–30. doi:10.1093/nar/28.1.27

15. Kanehisa M, Furumichi M, Tanabe M, Sato Y, Morishima K. KEGG: New perspectives on
genomes, pathways, diseases and drugs. *Nucleic Acids Res* (2017) **45**:D353–D361.
doi:10.1093/nar/gkw1092

16. Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K, Madden TL.
BLAST+: Architecture and applications. *BMC Bioinformatics* (2009) **10**:1–9.
doi:10.1186/1471-2105-10-421

17. Berman HM, Battistuz T, Bhat TN, Bluhm WF, Bourne PE, Burkhardt K, Iype L, Jain S,
Fagan P, Marvin J, et al. The Protein Data Bank. *Acta Crystallogr Sect D-Biological
Crystallogr* (2002) **58**:899–907. doi:10.1107/S0907444902003451

18. Backstrom AD, McMordie RAS, Begley TP. Biosynthesis of Thiamin I: The Function of the
thiE Gene Product. *J Am Chem Soc* (1995) **117**:2351–2352. doi:10.1021/ja00113a025

19. Begum A, Drebes J, Kikhney A, Müller IB, Perbandt M, Svergun D, Wrenger C, Betzel C.
Staphylococcus aureus thiaminase II: Oligomerization warrants proteolytic protection against
serine proteases. *Acta Crystallogr Sect D Biol Crystallogr* (2013) **69**:2320–2329.
doi:10.1107/S0907444913021550

20. Jenkins AH, Schyns G, Potot S, Sun G, Begley TP. A new thiamin salvage pathway. *Nat
Chem Biol* (2007) **3**:492–497. doi:10.1038/nchembio.2007.13

21. Reddick JJ, Nicewonger R, Begley TP. Mechanistic Studies on Thiamin Phosphate Synthase:
Evidence for a Dissociative Mechanism †. *Biochemistry* (2001) **40**:10095–10102.
doi:10.1021/bi010267q

22. Cheng G, Bennett EM, Begley TP, Ealick SE. Crystal structure of 4-amino-5-hydroxymethyl-
2-methylpyrimidine phosphate kinase from Salmonella typhimurium at 2.3 Å resolution.
Structure (2002) **10**:225–235. doi:10.1016/S0969-2126(02)00708-6

23. Newman JA, Das SK, Sedelnikova SE, Rice DW. The Crystal Structure of an ADP Complex
of Bacillus subtilis Pyridoxal Kinase Provides Evidence for the Parallel Emergence of Enzyme
Activity During Evolution. *J Mol Biol* (2006) **363**:520–530. doi:10.1016/j.jmb.2006.08.013

24. Nodwell MB, Koch MF, Alte F, Schneider S, Sieber SA. A subfamily of bacterial ribokinases
utilizes a hemithioacetal for pyridoxal phosphate salvage. *J Am Chem Soc* (2014) **136**:4992–
4999. doi:10.1021/ja411785r

25. Nodwell MB, Menz H, Kirsch SF, Sieber SA. Rugulactone and its Analogues Exert
Antibacterial Effects through Multiple Mechanisms Including Inhibition of Thiamine
Biosynthesis. *ChemBioChem* (2012) **13**:1439–1446. doi:10.1002/cbic.201200265

26. Peapus DH, Chiu HJ, Campobasso N, Reddick JJ, Begley TP, Ealick SE. Structural
characterization of the enzyme-substrate, enzyme-intermediate, and enzyme-product
complexes of thiamin phosphate synthase. *Biochemistry* (2001) **40**:10103–10114.
doi:10.1021/bi0104726

27.   Paul D, Chatterjee A, Begley TP, Ealick SE. Domain organization in Candida glabrata THI6, a
409   bifunctional enzyme required for thiamin biosynthesis in Eukaryotes. *Biochemistry* (2010)
410   **49**:9922–9934. doi:10.1021/bi101008u

411   28.   Peapus DH, Chiu H-J, Campobasso N, Reddick JJ, Begley TP, Ealick SE. Structural
412   Characterization of the Enzyme–Substrate, Enzyme–Intermediate, and Enzyme–Product
413   Complexes of Thiamin Phosphate Synthase † , ‡. *Biochemistry* (2001) **40**:10103–10114.
414   doi:10.1021/bi0104726

415   29.   Manzetti S, Zhang J, van der Spoel D. Thiamin Function, Metabolism, Uptake, and Transport.
416   *Biochemistry* (2014) **53**:821–835. doi:10.1021/bi401618y

417   30.   Hava DL, Camilli A. Large-scale identification of serotype 4 Streptococcus pneumoniae
418   virulence factors. *Mol Microbiol* (2002) **45**:1389–1406. doi:10.1046/j.1365-2958.2002.t01-1-
419   03106.x

420   31.   Akerley BJ, Rubin EJ, Novick VL, Amaya K, Judson N, Mekalanos JJ. A genome-scale
421   analysis for identification of genes required for growth or survival of Haemophilus influenzae.
422   *Proc Natl Acad Sci U S A* (2002) **99**:966–971. doi:10.1073/pnas.012602299

423   32.   Sassetti CM, Boyd DH, Rubin EJ. Comprehensive identification of conditionally essential
424   genes in mycobacteria. *Proc Natl Acad Sci U S A* (2001) **98**:12712–12717.
425   doi:10.1073/pnas.231275498

426   33.   Reddick JJ, Saha S, Lee J, Melnick JS, Perkins J, Begley TP. The mechanism of action of
427   bacimethrin, a naturally occurring thiamin antimetabolite. *Bioorg Med Chem Lett* (2001)
428   **11**:2245–2248. doi:10.1016/S0960-894X(01)00373-0

429   34.   Breiding-Mack S, Zeeck A, Zahner H. Metabolic Products Of Microorganisms. 239
430   Bacimethrin Isolated From Streptomyces Albus Identification, Derivatives, Synthesis And
431   Biological Properties. *J Antibiot (Tokyo)* (1987) **40**:1431–1439.
432   doi:10.7164/antibiotics.40.1431

433   35.   Reddick JJ, Saha S, Lee J, Melnick JS, Perkins J, Begley TP. The mechanism of action of
434   bacimethrin, a naturally occurring thiamin antimetabolite. *Bioorg Med Chem Lett* (2001)
435   **11**:2245–2248. doi:10.1016/S0960-894X(01)00373-0

436   36.   Zilles JL, Croal LR, Downs DM. Action of the Thiamine Antagonist Bacimethrin on Thiamine
437   Biosynthesis. *J Bacteriol* (2000) **182**:5606–5610. doi:10.1128/JB.182.19.5606-5610.2000

438   37.   Mukherjee T, Hanes J, Tews I, Ealick SE, Begley TP. Pyridoxal phosphate: Biosynthesis and
439   catabolism. *Biochim Biophys Acta - Proteins Proteomics* (2011) **1814**:1585–1596.
440   doi:10.1016/j.bbapap.2011.06.018

441   38.   Ehrenshaft M, Bilski P, Li M, Chignell CF, Daub ME. A highly conserved sequence is a novel
442   gene involved in de novo vitamin B6 biosynthesis. *Proc Natl Acad Sci U S A* (1999) **96**:9374–
443   9378. doi:10.1073/pnas.96.16.9374

444   39.   Neuwirth M, Strohmeier M, Windeisen V, Wallner S, Deller S, Rippe K, Sinning I,
445   Macheroux P, Tews I. X-ray crystal structure of Saccharomyces cerevisiae Pdx1 provides

insights into the oligomeric nature of PLP synthases. *FEBS Lett* (2009) **583**:2179–2186.
doi:10.1016/j.febslet.2009.06.009

40. Smith AM, Brown WC, Harms E, Smith JL. Crystal structures capture three states in the
catalytic cycle of a pyridoxal phosphate (PLP) Synthase. *J Biol Chem* (2015) **290**:5226–5239.
doi:10.1074/jbc.M114.626382

41. Guédez G, Hipp K, Windeisen V, Derrer B, Gengenbacher M, Böttcher B, Sinning I, Kappes
B, Tews I. Assembly of the eukaryotic PLP-synthase complex from Plasmodium and
activation of the Pdx1 enzyme. *Structure* (2012) **20**:172–184. doi:10.1016/j.str.2011.11.015

42. Strohmeier M, Raschle T, Mazurkiewicz J, Rippe K, Sinning I, Fitzpatrick TB, Tews I.
Structure of a bacterial pyridoxal 5'-phosphate synthase complex. *Proc Natl Acad Sci* (2006)
103:19284–19289. doi:10.1073/pnas.0604950103

43. Neuwirth M, Flicker K, Strohmeier M, Tews I, Macheroux P. Thermodynamic
characterization of the protein-protein interaction in the heteromeric Bacillus subtilis
pyridoxalphosphate synthase. *Biochemistry* (2007) **46**:5131–5139. doi:10.1021/bi602602x

44. Grubman A, Phillips A, Thibonnier M, Kaparakis-Liaskos M, Johnson C, Thiberge JM,
Radcliff FJ, Ecobichon C, Labigne A, de Reuse H, et al. Vitamin B6 is required for full
motility and virulence in Helicobacter pylori. *MBio* (2010) **1**:1–9. doi:10.1128/mBio.00112-10

45. Dick T, Manjunatha U, Kappes B, Gengenbacher M. Vitamin B6 biosynthesis is essential for
survival and virulence of Mycobacterium tuberculosis. *Mol Microbiol* (2010) **78**:980–988.
doi:10.1111/j.1365-2958.2010.07381.x

46. Qaidi S El, Yang J, Zhang JR, Metzger DW, Bai G. The vitamin B6 biosynthesis pathway in
streptococcus pneumoniae is controlled by pyridoxal 5'-phosphate and the transcription factor
PdxR and has an impact on ear infection. *J Bacteriol* (2013) **195**:2187–2196.
doi:10.1128/JB.00041-13

47. Wrenger C, Knöckel J, Walter RD, Müller IB. Vitamin B1 and B6 in the malaria parasite:
Requisite or dispensable? *Brazilian J Med Biol Res* (2008) **41**:82–88. doi:10.1590/S0100-
879X2008005000006

48. Kronenberger T, Lindner J, Meissner KA, Zimbres FM, Coronado MA, Sauer FM, Schettert I,
Wrenger C. Vitamin B6-dependent enzymes in the human malaria parasite plasmodium
falciparum: A druggable target? *Biomed Res Int* (2014) **2014**: doi:10.1155/2014/108516

49. Kronenberger T, Schettert I, Wrenger C. Targeting the vitamin biosynthesis pathways for the
treatment of malaria. *Future Med Chem* (2013) **5**:769–779. doi:10.4155/fmc.13.43

50. Wrenger C, Eschbach ML, Müller IB, Warnecke D, Walter RD. Analysis of the vitamin B6
biosynthesis pathway in the human malaria parasite Plasmodium falciparum. *J Biol Chem*
(2005) **280**:5242–5248. doi:10.1074/jbc.M412475200

51. Zhu J, Burgner JW, Harms E, Belitsky BR, Smith JL. A new arrangement of (β/α)8 barrels in
the synthase subunit of PLP synthase. *J Biol Chem* (2005) **280**:27914–27923.
doi:10.1074/jbc.M503642200

52.   Raschle T, Amrhein N, Fitzpatrick TB. On the two components of pyridoxal 5′-phosphate
485   synthase from Bacillus subtilis. *J Biol Chem* (2005) **280**:32291–32300.
486   doi:10.1074/jbc.M501356200

487   53.   Newman JA, Das SK, Sedelnikova SE, Rice DW. Cloning, purification and preliminary
488   crystallographic analysis of a putative pyridoxal kinase from Bacillus subtilis. *Acta
489   Crystallogr Sect F Struct Biol Cryst Commun* (2006) **62**:1006–1009.
490   doi:10.1107/S1744309106035779

491   54.   Gandhi AK, Desai J V., Ghatge MS, di Salvo ML, Di Biase S, Danso-Danquah R, Musayev
492   FN, Contestabile R, Schirch V, Safo MK. Crystal structures of human pyridoxal kinase in
493   complex with the neurotoxins, ginkgotoxin and theophylline: Insights into pyridoxal kinase
494   inhibition. *PLoS One* (2012) **7**:1–11. doi:10.1371/journal.pone.0040954

495   55.   Barra ALC, Dantas L de OC, Morão LG, Gutierrez RF, Polikarpov I, Wrenger C, Nascimento
496   AS. Essential Metabolic Routes as a Way to ESKAPE from Antibiotic Resistance. *bioRxiv*
497   (2019)817015. doi:10.1101/817015

500   **Table 1. Panel of observed enzymes for the thiamine pathway in ESKAPE pathogens.**

Enzyme	*Enterococcus faecium (ATCC 8459)*	*Staphylococcus aureus (NCTC 8325)*	*Klebsiella pneumoniae (subsp. Pneumoniae ATCC 43816 KPPR1)*	*Acinetobacter baumannii (ATCC 17978)*	*Pseudomonas aeruginosa (NCGM 1900)*	*Enterobacter sp. (638)*
ThiD	✓	✓	✓	✓	✗	✓
THiM	✗	✓	✓	✓	✓	✓
ThiE	✗	✓	✓	✓	✓	✓
TPK	✓	✓	✓	✗	✗	✗
TenA	✗	✓	✓	✓	✗	✗

502 **Table 2. Panel of observed enzymes for the pyridoxal 5'-phosphate pathway in ESKAPE**
503 **pathogens.**

Enzyme	*Enterococcus faecium (ATCC 8459)*	*Staphylococcus aureus (NCTC 8325)*	*Klebsiella pneumoniae (subsp. Pneumoniae ATCC 43816 KPPR1)*	*Acinetobacter baumannii (ATCC 17978)*	*Pseudomonas aeruginosa (NCGM 1900)*	*Enterobacter sp. (638)*
Pdx1	✗	✓	✗	✗	✗	✗
Pdx2	✗	✓	✗	✗	✗	✗