1

Co-expression clustering across flower development identifies modules and candidate genes for diverse floral forms in *Achimenes* (Gesneriaceae)

Wade R Roberts[1,2], Eric H Roalson[1]

[1] School of Biological Sciences, Washington State University, Pullman, WA, United States
[2] Current address: Department of Biological Sciences, University of Arkansas, Fayetteville, AR, United States

Corresponding Author:
Wade Roberts[1]
Department of Biological Sciences, University of Arkansas, SCEN 601, Fayetteville, AR, 72701, United States
Email address: wader@uark.edu

Abstract

Background. Genetic pathways involved with flower color and shape are thought to play an important role in the development of floral forms associated with different pollination syndromes, such as those associated with bee, butterfly, or hummingbird pollination. Because pollination syndromes are complex traits that are orchestrated by multiple genes and pathways, the gene regulatory networks have not been explored. Gene co-expression networks provide a systems level approach to identify important contributors to floral diversification.

Methods. RNA-sequencing was used to assay gene expression across two stages of flower development (an early bud and an intermediate stage) in 10 species of *Achimenes* (Gesneriaceae). Two stage-specific co-expression networks were created from 9503 orthologs and analyzed to identify module hubs and the network periphery. Module association with bee, butterfly, and hummingbird pollination syndromes was tested using phylogenetic mixed models. The relationship between network connectivity and evolutionary rates (d_N/d_S) was tested using linear models.

Results. Networks contained 65 and 62 modules that were largely preserved between developmental stages and contained few stage-specific modules. Over a third of the modules in both networks were associated with floral form and pollination syndrome. Within these modules, several hub nodes were identified that related to the production of anthocyanin and carotenoid pigments and the development of flower shape. Evolutionary rates were decreased in highly connected genes and elevated in peripheral genes.

38   **Discussion.** This study aids in the understanding of the genetic architecture and network
39   properties underlying the development of floral form and provides valuable candidate modules
40   and genes for future studies.
41

Introduction

Flowers display an extraordinary range of forms, shapes, and colors. Understanding the
evolution and development of diverse flowers across angiosperms has long been a major goal in
biology. This variation has often been attributed to pollinator-mediated selection and shifts in
floral form are often observed over time (Stebbins 1970; O'Meara et al. 2016). The evolution of
flowers and the relationship between plant and pollinator is considered one of the key
innovations in angiosperm diversification (Faegri and van der Pijl 1979; Fenster et al. 2004; Van
der Niet and Johnson 2012; Barrett 2013; Sauquet and Magallón 2018). Characters of the floral
phenotype are tightly linked to pollinator efficiency and overall plant reproductive success
(Rosas-Guerrero et al. 2014). Pollinators may positively select specific floral traits across
evolutionary time (Gervasi and Schiestle 2017) and flowers may be under constant and strong
selective pressure. The genetic basis of floral and pollination syndrome divergence has been
examined in model systems extensively, particularly in *Mimulus* (Bradshaw et al. 1998; Yuan et
al. 2016) and *Petunia* (Hoballah et al. 2007; Hermann et al. 2015), and more recently in many
non-model systems (e.g., Wessinger et al. 2014; Alexandre et al. 2015).
Among the most widely used large-scale experimental approaches to investigate genome
function are transcriptome analyses (e.g., Pickrell et al. 2010; Raherison et al. 2015), particularly
in non-model organisms where no reference genome or functional genomics data exists. The
transcriptome is highly dynamic as a consequence of quantitative variation over spatial and
temporal scales. Changes in expression often result from the combinatorial action of genetic
regulatory pathways orchestrating development and responses to environmental stimuli.
Therefore, the transcriptome can be viewed as a link between the genotype and phenotype and
may be acted upon through selection (Romero et al. 2012; Prasad et al. 2013). As changes in
gene expression may underlie many of the phenotypic changes between species (Brawand et al.
2011; Romero et al. 2012; Uebbing et al. 2016), studying the transcriptome may shed light on
important pathways and targets of selection. Phenotypic changes frequently arise through
functional changes in conserved developmental pathways among closely related species. For
example, changes in enzyme function and transcriptional regulation of the anthocyanin pathway
has been implicated frequently across angiosperms in the evolution of red flowers (e.g., Des
Marais and Rausher 2010; Smith and Rausher 2011). It is now recognized that most genes act as
members of biological pathways or of co-regulated modules (Stuart et al. 2003).
Here, we undertake a comparative study of gene expression during flower development
in the genus *Achimenes*. This group is a member of the diverse African violet family
(Gesneriaceae) and distributed throughout Mexico and Central America. *Achimenes* is a young
lineage (*c*. 7-12 Mya; Roalson and Roberts 2016) known for its floral diversity, a feature thought

to be associated with speciation (Ramírez Roa 1987; Roalson et al. 2003). Four pollination
syndromes are found in *Achimenes*, including bee (melittophily), butterfly (psychophily), female
euglossine bee (euglossophily), and hummingbird (ornithophily) (Figure 1). These syndromes
have traditionally been defined on the basis of flower color and flower shape (Ramírez Roa
1987) and recently through pollinator observation (Martén-Rodríguez et al. 2015; Ramírez-
Aguirre et al. 2019). Repeated origins and transitions between these pollination syndromes have
been hypothesized in *Achimenes*, making this small lineage an attractive system to understand
the genetic and ecological factors underlying floral diversification.
In this study, we aimed to characterize and compare patterns of gene co-expression
during flower development across 10 *Achimenes* species. So far, few studies have investigated
the genetic basis of floral form in the Gesneriaceae family (e.g., Alexandre et al. 2015; Roberts
and Roalson 2017). Given that bee, butterfly, and hummingbird pollination syndromes evolved
in parallel across *Achimenes,* we hypothesized that: (1) distinct sets of co-expressed genes would
correlate to each syndrome during flower development, and (2) genes in pathways involved with
flower color and shape may be central in each network during flower development. Using a
comparative transcriptome approach, we clustered a shared set of 9503 orthologs into two
different co-expression networks, each corresponding to two stages of flower development: an
early bud stage and an intermediate stage. This strategy allowed us to compare gene co-
expression patterns temporally. We compared these networks and tested whether gene co-
expression clusters correlated with different floral traits (flower color, flower shape, spurs). We
also tested whether network properties, such as connectivity and expression, influence
evolutionary constraints. Our results allowed us to quantify the extent of shared gene co-
expression among closely related species and to identify candidate genes and pathways that may
be associated with the repeated evolution of floral forms underlying pollinator specificity.
102
103   ## Materials & Methods
104
Plant materials
Ten of the 26 currently recognized species in *Achimenes* were sampled: five from Clade 1
and five from Clade 2 (Figure 1). These species were chosen based on their hypothesized close
relationships and their diversity in floral form (Roalson et al. 2003). On the basis of previous
molecular studies (Roalson and Roberts 2016), two species were chosen as outgroups to
represent related lineages: *Eucodonia verticillata* and *Gesneria cuneifolia* (Figure 1). All
selected species were grown in standard greenhouse conditions at Washington State University,
under 16-hour days, 24-27°C, and 80-85% humidity. Whole flower buds from two stages of
flower development were sampled for each species: an immature bud stage (Bud stage; Figure 1)
and an intermediate stage before anthesis (D stage; Figure 1). Each stage was sampled with two
or three biological replicates, contributing four or six samples for each species. The bud stage
was chosen to represent a baseline level of gene expression when floral phenotypes are largely
similar that could then be compared to the intermediate stage where many important pathways

involved in flower development and pigmentation show increased expression levels (Roberts and
Roalson 2017). All tissues were immediately frozen into liquid nitrogen and stored at -80°C.
120
RNA extraction and RNA-seq library construction
Total RNA was extracted from frozen tissue using the RNEasy Plant Kit (Qiagen) and
ribosomal depleted RNA samples were prepared using the RiboMinus Plant Kit (Thermo Fisher
Scientific). Stranded libraries were prepared using the NEBNext Ultra Directional RNA Library
Kit (New England Biolabs), barcoded, and pooled based on nanomolar concentrations. Library
quality and quantity were assessed using the Qubit (dsDNA HS assay; Thermo Fisher Scientific)
and BioAnalyzer 2100 (Agilent). The library pool was sequenced across 4 lanes of an Illumina
HiSeq2500 for paired-end 101 bp reads at the Genomics Core Lab at Washington State
University, Spokane. All new sequence data were deposited in the NCBI Sequence Read Archive
(BioProject: PRJNA435759).
Single replicate libraries for four of the species in the current study (*A. cettoana*, *A.*
erecta, *A. misera,* and *A. patens*) were previously sequenced for the same two timepoints (Bud
and D stages) (Roberts and Roalson 2017). These samples were combined with the newly
generated data to assembly de novo transcriptomes and perform co-expression clustering. This
data was deposited in the NCBI Sequence Read Archive (BioProject: PRJNA340450).
136
Transcriptome assembly
Per base quality, adapter contamination, and per base sequence content of raw reads was
assessed with FastQC tools (http://bioinformatics.babraham.ac.uk/projects/fastqc). One library
(AT6) returned poor yields and low-quality reads and was excluded from further analyses.
Trimmomatic v.0.36 (Bolger et al. 2014) was then used to remove contaminating adapter
sequences and dynamically trim low-quality bases (ILLUMINACLIP:TruSeq3-PE-2.fa:2:30:10
HEADCROP:13 LEADING:3 TRAILING:3 SLIDINGWINDOW:4:15 MINLEN:50). Trimmed
reads derived from an individual species were combined and de novo assembled into contigs
using Trinity (--SS_lib_type RF; Haas et al. 2013; Grabherr et al. 2011), generating a reference
transcriptome for each species. Sequence redundancy was reduced with CDHIT-EST (-c 0.99 -n
5; Li and Godzik 2006). Open reading frames were then predicted for each transcriptome
assembly with TransDecoder (Haas et al. 2013) using a BLASTp search against the SwissProt
database (www.uniprot.org) to maximize sensitivity. Finally, transcriptome assembly
completeness was assessed using BUSCO (Simão et al. 2015) against a set of single-copy plant
orthologs (embryophyte_odb10 dataset).
152
Orthogroup inference
To help facilitate orthogroup inference, transcriptome assemblies from flowers in
Sinningia eumorpha and *S. magnifica* were downloaded and combined with our assemblies
(https://doi.org/10.5061/dryad.4r5p1; Serrano-Serrano et al. 2017a). *Sinningia* is closely related
to our focal lineage in the Neotropical Gesneriaceae (Roalson and Roberts 2016). To identify

orthogroup sequences across all 14 species, we used a modified version of the approach
described in Yang and Smith (2014). All peptide sequences were clustered using an all-by-all
BLASTp search performed using DIAMOND (--evalue 1e-6; Buchfink et al. 2014) and results
were clustered using MCL v14-137 (-I 1.4; Enright et al. 2002). Homolog clusters were aligned
using MAFFT v7.271 (--genafpair --maxiterate 1000; Katoh and Standley 2013) and alignments
were trimmed using Phyutility v2.7.1 (-clean 0.1; Smith and Dunn 2008). Codon alignments
were then produced for each cluster using PAL2NAL v14 (Suyama et al. 2006) and maximum-
likelihood trees were constructed using FastTree v2.1.8 (Price et al. 2010) using the GTR model.
Trees were then trimmed to exclude branches >10 times longer than its sister or longer than 0.2
substitutions per site. Final homolog group alignments were created from the trimmed trees using
MAFFT and used for final homolog tree inference using RAxML v8.2.9 (Stamatakis 2014) and
the GTRGAMMA model. Orthogroups were inferred from the final homolog trees using the
rooted ingroup (RT) method of Yang and Smith (2014), using *S. eumorpha* and *S. magnifica* as
outgroups. Homolog trees were trimmed to include at least 2 ingroup species, resulting in 83595
orthogroups.
173
Quantifying gene expression
Trimmed reads for each sample were aligned to the corresponding transcriptome
assembly to quantify expression using default parameters in Kallisto v0.43.0 (Bray et al. 2016).
The resulting counts for each contig were matched with the inferred orthogroups derived above,
creating two separate matrices for the Bud and D stage samples. Rows corresponded to
orthogroups and columns corresponded to individual samples. Counts for the Bud and D stage
samples were separately transformed using variance-stabilizing transformation implemented in
the Bioconductor package DESeq2 (Love et al. 2014) and quantile normalized using the
Bioconductor package preprocessCore (Bolstad 2018). Lastly, counts for both stages were
separately corrected for confounding effects, such as batch or species effects, using principal
component regression with the 'sva_network' function in the Bioconductor package sva (Parsana
et al. 2019).
186
Co-expression networks and identification of modules
The Bud and D stage RNAseq data was clustered into gene co-expression networks using
the R package WGCNA (Langfelder and Horvath 2008). Prior to network construction,
orthogroups were filtered to remove any with >50% missing values or <0.3 expression variance.
In total, 9503 orthogroups common to both Bud and D stage data were used to construct separate
networks for each stage. Parameters for the Bud stage network were as follows: power = 18,
networkType = "signed", corType = "bicor", maxPOutliers = 0.05, TOMType = "signed",
deepSplit = 2, mergeCutHeight = 0.25. For the D stage network, the following parameters were
used: power = 16, networkType = "signed", corType = "bicor", maxPOutliers = 0.05, TOMType
= "signed", deepSplit = 2, mergeCutHeight = 0.25.

Module eigengenes and orthogroup connectivity were calculated in each network
separately using the 'moduleEigengenes' and 'intramodularConnectivity' functions in WGCNA,
respectively. The module eigengene is defined as the first principal component of a module and
represents the gene expression profile of a sample within a module (Langfelder and Horvath
2008). Connectivity refers to the sum of connection strengths a node has to other network nodes.
Module hub genes (defined as the most highly connected nodes within a module) were
identified in both networks based on module membership (kME) scores >0.9, calculated using
the 'signedKME' function of WGCNA. kME is defined as the correlation between the
expression levels of a gene and the module eigengene (Horvath and Dong 2008). Additionally,
using connectivity measures, we defined peripheral genes in each network as the lowest 10% of
connected nodes.
We tested whether Bud stage modules were preserved in the D stage and vice versa using
module preservation statistics in WGCNA (Langfelder et al. 2011). Median Rank and Zsummary
statistics were computed using the 'modulePreservation' function of WGCNA, using 200
permutations (Langfelder et al. 2011). Median rank statistics were compared to the "gold"
module, which consists of 1000 randomly selected orthogroups. Modules with a lower median
rank exhibit stronger preservation than modules with a higher median rank. Zsummary scores >
10 indicate strong evidence of preservation and scores < 2 indicate no evidence of preservation.
215
Phylogeny
We previously inferred a phylogeny for the 12 sampled species using 1306 single-copy
orthologs identified from the same transcriptome dataset used here (Roberts and Roalson 2018).
For comparative analyses of module-trait relationships, we randomly sampled 50 single-copy
ortholog gene trees and rescaled branch lengths to be proportional to time (ultrametric) using the
'chronos' function in the R package ape (Paradis et al. 2004). Bootstrap support was 100 for
nearly every branch in the Roberts and Roalson (2018) phylogeny, therefore we chose to use
randomly sampled ortholog trees to account for phylogenetic uncertainty.
224
Module and trait relationship
We correlated external traits (e.g., red flower color) with the module eigengenes using a
phylogenetic Markov Chain Monte Carlo method, implemented in the R packages MCMCglmm
(Hadfield 2010) and mulTree (Guillerme and Healy 2014). These analyses were performed
individually for each module in both the Bud and D stage networks. Three floral traits and
pollination syndrome were coded: primary flower color (purple, red, white, yellow), flower
shape (funnelform, salverform, tubular), corolla spur (absent, present), and syndrome (bee,
butterfly, hummingbird). We fit a multivariate mixed model with a non-informative prior, where
the shape and scale parameters equal 0.001, and residual variance was fixed. Analyses were
performed over the 50 randomly selected ortholog trees to account for phylogenetic uncertainty.
For each tree, two MCMC chains were run for 250k generations and discarding the first 50k as
burn-in. This process was run individually over each module in both the Bud and D stages. We

checked for convergence between model chains using the Gelman-Rubin statistic (Gelman and
Rubin 1992). Potential scale reduction values were all less than 1.1 and effective sample sizes for
all fixed effects were greater than 400. We considered fixed effects to be statistically significant
when the probabilities in the 95% credible region did not include zero.
241
Evolutionary rate analysis
Codon alignments of the filtered orthogroups (n=9503) were produced using PAL2NAL
v14 (Suyama et al. 2006). Subsequent alignments and corresponding gene trees were used as
input to the codeml package in PAML v4.9 (Yang 2007) to estimate d_N/d_S (omega, ω). d_N/d_S for
each orthogroup was estimated using a one-ratio model (model = 0, NSsites = 0), providing a
single estimate for each orthogroup to match other single orthogroup metrics, such as
connectivity. If d_N/d_S values were exceptionally large (= 999) because of zero synonymous
differences, those values were removed from the dataset.
First, we performed linear regression (LM) to test the effect of orthogroup connectivity,
average expression levels, and their interactions (explanatory variables) on the estimated d_N/d_S
values (main effect), LM = d_N/d_S ~ connectivity * expression. This was performed using the lm
function in R with 10000 bootstrap pseudoreplicates. Second, we ran two-sample t-tests with
10000 permutations to test whether hub nodes and periphery nodes in each network had lower
(for hubs) or higher (for periphery) d_N/d_S values than all background nodes. Third, we fit a LM
with 10000 bootstrap pseudoreplicates to test whether modules associated with a pollination
syndrome (bee, butterfly, or hummingbird; see Results below) had increased d_N/d_S values, LM =
d_N/d_S ~ syndrome. For all LM and permutation analyses, d_N/d_S, connectivity, and expression values
were log transformed to reach a normal distribution before being processed. Analyses were
performed separately for each network using the separately estimated connectivity and
expression levels.
262
Functional annotation and GO enrichment analysis
Filtered orthogroups (n=9503) were annotated based on a search against all green plant
proteins in the SwissProt database (release 2018_07; The UniProt Consortium 2017) using
Diamond BLASTp (--evalue 1e-6). Results were used to assign annotations and gene ontology
(GO) terms. GO enrichment analyses were performed using the R package TopGO with the
default "weight01" algorithm with the recommended cutoff of p < 0.05 (Alexa et al. 2006).
Enriched GO term redundancy was removed and summarized using the semantic similarity
measures implemented in the REVIGO web server (Supek et al. 2011).
271

Results

Transcriptome assembly
Twelve transcriptomes were sequenced (72 libraries) and assembled in 12 species after
sampling across two floral developmental stages (Supplemental Table 1). The transcriptomes had

an average assembly length of 244.9 Mb (± 33.9 Mb), an average of 217,510 (± 47k) transcripts,
and an average N50 length of 1988 bp (± 306 bp). Between 22,942 and 33,501 putative genes
were detected with 2.34 (± 0.35) putative isoforms. BUSCO identified an average of 87% of the
complete, conserved single-copy plant orthologs (Supplemental Table 1). De novo assembly
enables functional genomic studies in non-model systems without a reference genome, but has
the potential for misassemblies (Hornett and Wheat 2012) and subsequent downstream biases.
Our focus in our comparative study was on the presence of conserved orthologs, and not on the
presence of taxonomically restricted genes, which are not likely preserved across species. We
subsequently identified 83595 orthologous clusters (orthogroups), each containing at least 2
species.
287
Constructing the co-expression network for 12 species
After removing 74092 orthogroups with too many missing values (>50% missing) or low
expression variance (<0.3 variance) and keeping common orthogroups shared between the Bud
and D stages, 9503 orthogroups were used to construct two co-expression networks using the
weighted gene correlation network analysis (WGCNA) approach. Modules of co-expressed
orthogroups are inferred using the expression profiles of each sample regardless of species.
Clustering identified 65 and 62 co-expression modules in the Bud and D stage networks,
respectively (Figure 2). Module size in the Bud stage ranged from 22 ('mediumorchid') to 592
('turquoise') orthogroups (mean = 146) and ranged from 21 ('antiquewhite4') to 704
('turquoise') orthogroups (mean = 153) in the D stage. Out of the 9503 orthogroups, 7845 (83%)
and 8735 (92%) were assigned to modules in each network.
The remaining unassigned genes from each network that were not placed into a well-
defined module were assigned to the 'grey' module by WGCNA. Despite being unplaced, these
orthogroups may still be functionally relevant within each network. Unassigned genes in the
'grey' module were enriched for many processes, including transcription (Bud, n = 173; D, n =
84) and organ development (D, n = 55), among other functions (Supplemental Tables 2-3).
We tested whether modules in the Bud stage were preserved in the D stage, and vice
versa. Overall, preservation of modules between the Bud and D stage networks was high. Eight
of the 65 Bud stage modules ('brown4', 'coral2', 'floralwhite', 'lavenderblush3', 'lightcyan1',
'lightsteelblue1', 'maroon', 'mediumpurple3') were not strongly preserved in the D stage, while
only 1 of the 62 D stage modules ('darkseagreen4') was not strongly preserved in the Bud stage
(Supplemental Table 4). These patterns suggest that these non-preserved modules may play
unique roles within their network. Among the 8 Bud stage-specific modules were many
orthogroups related to primary cell growth, meristem identity and differentiation, stamen
development, and flowering (Supplemental Table 2). In the single D stage-specific module were
many orthogroups related to methylation, flowering, and carotenoid biosynthesis (Supplemental
Table 3). Despite the relatively high preservation, correspondence between module membership
in the Bud and D stages was moderate (Figure 2). Orthogroup module membership tended to

shift between each network. For example, most members of module 'midnightblue' in the Bud
stage are transferred to modules 'brown' and 'black' in the D stage (Figure 2).
We defined network hubs (the most highly connected nodes within a module) with
module membership (kME) scores > 0.9. Using this definition, all modules in each network
contained at least one hub, with the percentage of hubs ranging from 1% to 24% of orthogroups
in a module (Figures 3 and 4; Supplemental Table 5). Of the 632 and 691 hubs identified in the
Bud and D stages, a significant portion (110) were shared between networks (Fisher's Exact
Test, odds ratio = 3.02, P < 0.001). Functional enrichment of hubs in the Bud stage revealed that
many were involved in the regulation of development and metabolic processes (n = 40), transport
and protein localization (n = 39), biosynthetic processes (n = 19), and DNA methylation (n = 7),
consistent with the notion that hubs usually play important roles in integrating other genes during
network evolution (Ravasz et al. 2002) (Table 1). In the D stage, hubs were enriched for
functions related to translation (n = 33), transport (n = 9), ethylene and abscisic acid signaling
pathways (n = 17), carotenoid and anthocyanin biosynthetic processes (n = 11), and the
regulation of gene expression (n = 20), among others (Table 1).
Nodes in the periphery of each network were loosely defined by considering the lowest
10% connected (connectivity$_{Bud}$ < 2.824, connectivity$_D$ < 4.651). These definitions identified 951
periphery nodes (Supplemental Table 6), of which 301 were peripheral in both networks
(Fisher's Exact Test, odds ratio = 5.67, P < 0.001). These periphery nodes in the Bud stage were
enriched for functions related to reproductive structure development (n = 75), transport and
localization (n = 107), and signaling (n = 389), among others (Table 2). In the D stage, the
periphery was enriched for functions related to the regulation of flower development (n = 64),
the regulation of flowering time (n = 18), the regulation of flavonoid biosynthesis (n = 7), and
cell fate specification (n = 12), among others (Table 2). Greater than expected numbers of nodes
in the D stage were related to regulation (Fisher's exact test, odds ratio = 1.191, p = 0.02752) and
transcription (Fisher's exact test, odds ratio = 1.370, p = 0.001549). In contrast, there were no
differences in the expected number of nodes related to regulation (Fisher's exact test, odds ratio
= 0.955, p = 0.7044) or transcription (Fisher's exact test, odds ratio = 1.047, p = 0.3549) in the
Bud stage.
345
Modules associated with floral traits
We calculated module eigengenes, single values which represent the gene expression
profiles of a sample within a module. The extent of module involvement in various biological
processes can be tested by correlating eigengenes with external traits, such as flower color. We
tested the relationship between the module eigengenes and the four floral traits (pollination
syndrome [bee, butterfly, hummingbird], flower color [purple, red, yellow, white], flower shape
[funnelform, salverform, tubular], and corolla spurs [presence]), while controlling for any
phylogenetic bias in the dataset using a phylogenetic mixed model (Figures 3-4; Supplemental
Figures 3-6). We also examined whether eigengene values (> 0.1 or < -0.1) in single species
contributed to a strong correlation or whether eigengene values (> 0.1 or < -0.1) in multiple

species contributed to a strong correlation (Figures 3-4; Supplemental Figures 3-6). We
considered strong evidence for trait association when a significant correlation from MCMCglmm
was coupled with elevated eigengene expression in multiple species. WGCNA does not use trait
information when inferring modules, so modules that correlated strongly with any floral trait
were inferred without a priori knowledge.

Pollination syndrome—We included 3 species with the bee pollination syndrome (*A.*
candida, *A. misera*, and *E. verticillata*), 4 species with the butterfly pollination syndrome (*A.*
cettoana, *A. grandiflora*, *A. longiflora*, and *A. patens* 'Major'), and 5 species with the
hummingbird pollination syndrome (*A. admirabilis*, *A. antirrhina*, *A. erecta*, *A. pedunculata*, and
G. cuneifolia) (Figure 1). These pollination syndromes are traditionally defined in *Achimenes*
and other gesneriads based on primary flower color, flower shape, and the presence of corolla
spurs (Ramírez Roa 1987). Expression of 22 and 28 modules were positively associated with the
pollination syndromes in the Bud and D stage networks, respectively (Figures 3-4). There were
3, 7, and 13 modules associated with bee, butterfly, and hummingbird pollination syndrome in
the Bud stage network, while 2, 8, and 12 modules were associated with the same syndromes in
the D stage. Modules were never associated with more than one syndrome. Butterfly and
hummingbird syndromes were never correlated to the same modules and were often correlated in
opposite directions (i.e., butterfly was positive correlated and hummingbird was negative
correlated).

Flower color—Primary flower color is closely associated to pollination syndrome in
Achimenes, and we identified 26 and 14 modules associated to any flower color in the Bud and D
stage networks, respectively (Figures 3-4). Red and yellow are associated with hummingbird
pollination, purple with butterfly pollination, and white with bee pollination. In the Bud stage,
there were 14, 11, and 1 module associated with red, purple, or yellow flower color. In the D
stage, no modules were associated with yellow, but 8 and 6 modules were associated with red or
purple flower color. In both networks, no modules were corelated to white flower color. Similar
to syndrome, purple and red flower color were never associated with the same modules and were
always correlated in opposite directions.

Flower shape—Flower shape is also important but not as closely associated to pollination
syndromes as primary flower color in *Achimenes*. Flowers with bee pollination tend to have
funnelform flowers, butterfly pollinated flowers all have salverform flowers, and hummingbird
pollinated flowers have either salverform or tubular flowers. Here, there were 1, 8, and 5
modules whose expression correlated to funnelform, salverform, or tubular flowers in the Bud
stage (Figure 3). Fewer modules were correlated in the D stage network, but expression in 5 and
1 module were associated with salverform and tubular shapes (Figure 4). No modules were
positively associated with more than one flower shape.

Corolla spurs—The presence of corolla spurs is found only on butterfly pollinated
species, two of which were included: *A. grandiflora* and *A. patens*. There was expression in 5
modules associated with spurs in the Bud stage network, while only 1 module was associated in
the D stage network (Figures 3-4).

Evolutionary rates correlate to network location

We ran several analyses to explore how several network variables may affect evolutionary rates of the orthogroups within each network (as measured by d_N/d_s). First, we tested the effects of orthogroup connectivity, expression levels, and their interaction on the d_N/d_s values (Supplemental Table 7). Results indicated a collective effect between connectivity and expression on d_N/d_s in both the Bud stage (LM, $F(3,9548) = 92.58$, p = 0, $\eta^2 = 0.03$) and the D stage networks (LM, $F(3,9548) = 78.36$, p = 0, $\eta^2 = 0.02$). We found that orthogroup connectivity was negatively correlated with d_N/d_s in the Bud stage network (LM, t = -16.31, p = 0) and the D stage network (LM, t = -15.13, p = 0) (Figure 5), while expression was positively correlated with d_N/d_s in the Bud stage (LM, t = 2.12, p = 0.0344) and not in the D stage (LM, t = 1.51, p = 0.1324) (Supplemental Figure 7). The interaction between connectivity and expression also indicated a marginal dependency of both predictors on d_N/d_s in the Bud stage (LM, t = -2.022, p = 0.0433) but not in the D stage network (LM, t = -1.823, p = 0.0683). These patterns have been observed in model species and may be universal to all eukaryotes.

Second, we tested whether the nodes we defined as hubs in each network had lower d_N/d_s values, suggestive of increased evolutionary constraint (Supplemental Table 7). Hubs are considered to functionally significant and may control large portions of the network. We found that the hubs in the Bud stage network had lower d_N/d_s than all background genes (Two-sample t-test, $t(679.31) = 7.0934$, p = 1.65e-12, Cohen's d = 0.383; Permutation t-test. mean diff. = 0.266, p = 0.0001) (Supplemental Figure 8). Similarly, hubs in the D stage network also had lower d_N/d_s than all background genes (Two-sample t-test, $t(755.91) = 4.1918$, p = 1.55e-05, Cohen's d = 0.206; Permutation t-test, mean diff. = 0.143, p = 0.0001) (Supplemental Figure 8).

Third, we tested whether the most peripheral nodes (the lowest 10% connected nodes) had higher d_N/d_s values, which may suggest relaxed evolutionary constraint (Supplemental Table 7). These nodes are loosely connected within the network and their roles may be more apt to fluctuate during evolution and development. These peripheral genes had higher d_N/d_s than all background genes in the Bud stage network (Two-sample t-test, $t(1487.70) = -10.272$, p = 0, Cohen's d = -0.253; Permutation t-test, mean diff. = -0.176, p = 0.0001) and the D stage network (Two-sample, $t(1512.30) = -10.387$, p = 0, Cohen's d = -0.252; Permutation t-test, mean diff. = -0.175, p = 0.0001) (Supplemental Figure 8).

Lastly, we tested whether modules correlated with bee, butterfly, or hummingbird pollination syndromes showed increased evolutionary rates over the non-associated modules. Results indicated a very weak effect of syndrome association on d_N/d_s in the Bud network (LM, $F(3,9548) = 5.186$, p = 0.001406, $\eta^2 = 0.0016$) and the D network (LM, $F(3,9548) = 4.277$, p = 0.005038, $\eta^2 = 0.0013$) (Supplemental Table 7). Only bee syndrome associated modules had marginally lower d_N/d_s values in both networks (Bud stage, LM, t = -3.58, p = 0.000346; D stage, LM, t = -2.508, p = 0.0122). Butterfly associated modules had marginally lower d_N/d_s only in the D stage (LM, t = -2.548, p = 0.0108), while hummingbird associated modules did not have any effect in either network.

Discussion

Here we report on patterns of gene co-expression across two stages of flower
development in 12 species of *Achimenes*, *Eucodonia*, and *Gesneria*. Our comparative and
phylogenetic context allowed us to reveal gene co-expression patterns across a group of closely
related plant species that correlated to diverse pollination syndromes and floral traits, including
flower color, flower shape, and corolla spurs. Our results suggest floral forms that are associated
with bee, butterfly, and hummingbird pollinator groups are correlated with many co-expression
modules. We found that nearly a third of modules in each network (23 and 21 modules in the
Bud and D stages, respectively) had evidence for association with the three pollination
syndromes (Figures 3-4). In our analysis, large fractions of our 9503 orthogroups set could be
partitioned into modules, network hubs and peripheral nodes identified, and candidate
orthogroups were identified for their potential importance in floral phenotypic differentiation.
Lastly, orthogroup position in each network (measured by connectivity) had an association with
evolutionary rate (d_N/d_S) (Figure 5). These analyses provide a first overview of floral
transcriptional architecture across members of the same genus that display diverse floral forms.

Hub nodes contain important candidates for floral form

To date, a large number of studies conducted in diverse angiosperm systems, have found
that relatively few genes are associated with floral form differentiation (e.g., Quattrocchio et al.
1999; Howarth et al. 2011). Despite evidence for extensive gene/genome duplications that could
expand the genetic repertoire (e.g., Tang et al. 2008), many recent studies have suggested that the
generation of novel genes is rare (Patterson et al. 2009; De Smet et al. 2013). Thus, the evolution
and development of diverse floral forms may predominantly occur through the co-option of
existing pathways (Preston et al. 2011). To our knowledge, gene co-expression network analyses
have not been previously used across multiple species in a non-model lineage to understand
flower development and evolution.

Hub nodes (genes) are considered to be conserved, highly connected nodes that are
central to network architecture and may directly orchestrate numerous biological processes
(Ravasz et al. 2002; Hu et al. 2016; Mähler et al. 2017). Our analyses identified over 600 gene
nodes as hubs within each network, the majority of which were only hubs within a specific floral
stage. These hub nodes tended to have lower d_N/d_S estimates, suggesting that they not prone to
rapid evolutionary change (Supplemental Table 7). Unsurprisingly, hub nodes in the Bud stage
were enriched for many processes related to early flower development, including cell growth,
leaf formation, and photosystem assembly (Table 1). Similarly, hub nodes in the D stage were
enriched for genes that may be related the development of distinct floral forms, including
carotenoid biosynthesis, anthocyanin biosynthesis, and cell morphogenesis (Table 1). Within this
context of flower development, we identified a number of hubs that we consider important
candidates for involvement in the production of flower color, the development of floral form, and
the production of corolla spurs. Many of these candidates were considered previously (Roberts

and Roalson 2017) as potentially important for the diversification of floral form in *Achimenes*.
Undoubtedly, this list of hubs contains many genes that may important for other biological traits
that we do not consider here, such as scent, nectar, and pollen production.
Flower color in *Achimenes* is primarily due to the accumulation of anthocyanin and
carotenoid pigments (Roberts and Roalson 2017). We identified two members of the core
anthocyanin pathway as hubs. Homologs of flavonoid 3'-hydroxylase (*F3'H*;
cluster8703_1rr.inclade1.ortho3; module 'tan') and dihydroflavonol 4-reductase (*DFR*;
cluster5082_1rr.inclade1.ortho6; module 'darkslateblue') were identified as hubs in the Bud and
D stages, respectively (Supplemental Table 8). *F3'H* is involved in the shifting metabolic flux of
the pathway between production of red pelargonidins and purple cyanidins while *DFR* acts
downstream of *F3'H* and directs the production of anthocyanins. Shifting expression of *F3'H*
and shifting substrate specificity of *DFR* have been identified in many systems as important
contributors to transitions between red and blue flowers (e.g., Des Marais and Rausher 2010;
Smith and Rausher 2011). We also identified two homologs of carotenoid biosynthetic pathway
genes as hubs in the D stage: homologs of geranylgeranyl pyrophosphate synthase (*GGPPS*;
cluster10945_2rr.inclade1.ortho1; module 'sienna3') and lycopene beta-cyclase (*LCYB*;
cluster13774_1rr.inclade1.ortho1; module 'black'). *GGPPS* is involved in the production of
precursor molecules while *LCYB* directs production of either alpha-carotenes or beta-carotenes,
the latter of which is thought to be the predominant carotenoids found in *Achimenes* flowers
(Roberts and Roalson 2017).
Less is known about the genetic pathways giving rise to the diverse flower shapes in
Achimenes. Among the hubs found in the Bud stage were homologs of many proteins involved in
floral development and cell differentiation, including the floral homeotic protein agamous (*AG*;
cluster7311_1rr.inclade1.ortho3; module 'darkturquoise'), phantastica (*PHAN* or rough sheath 2
[*RS2*]; cluster11220_1rr.inclade1.ortho3; module 'yellow'), and tcp4 (*TCP4*;
cluster12380_1rr.inclade1.ortho2; module 'darkgreen'). *AG* plays a role in maintaining floral
meristem determinancy and is required for normal development of stamens and carpels (Gómez-
Mena et al. 2005). *PHAN* is involved in cell differentiation and adaxial identity by repressing
numerous knox homeobox genes (Waites et al. 1998). *TCP4* is a cell division regulator that was
identified as a candidate for corolla spur development in columbine (Yant et al. 2015) and
showed increased expression during later stages of floral development in *Achimenes patens*
(Roberts and Roalson 2017). Finally, among the hubs in the D stage network was a homolog of
the floral homeotic protein globosa (*GLO*; cluster2750_1rr.inclade1.ortho3; module 'skyblue3'),
which acts in conjunction with deficiens (*DEFA*) to specify petal identity (Tröbner et al. 1990).
510
Peripheral genes are enriched for transcriptional regulators in later stages of development
Peripheral nodes have low connectivity within co-expression networks (Supplemental
Table 7), tend to have higher rates of molecular evolution (e.g., Masalia et al. 2017), and have
been hypothesized to contribute more to evolutionary innovations than network hubs (e.g.,
Ichihashi et al. 2014). Of the 951 nodes we defined as peripheral in both Bud and D stage

networks, roughly a third were consistently peripheral under our definitions (lowest 10%
connected; Supplemental Table 6). After identifying the nodes with homologies to transcriptional
regulators, we found there was no difference in the expected number in the early Bud stage,
while there were greater than expected numbers of transcriptional regulators in the periphery of
the D stage (see Results). No such patterns existed when we tested these transcriptional
regulators as hub genes. While Mähler et al. (2017) found an enrichment for transcription factors
among hub genes in a single species network of *Populus tremula,* our results in a multi-species
network find the opposite pattern. Shifting the expression patterns of various genes and pathways
can lead to phenotypic divergence among closely related species (West-Eberhard 1989). This can
be seen during later stages of flower development when distinct traits, such as red or purple
flower color, can appear in different species due to shifting expression patterns within the
anthocyanin biosynthetic pathway (Smith and Rausher 2011). The connection between the
apparent overabundance of transcriptional regulators in later stages of flower development and
the rapid phenotypic divergence in floral form will need to be explored further.
Some notable peripheral nodes included homologs of divaricata (*DIV*), dichotoma
(*DICH*), and wuschel (*WUS*) (Supplemental Table 6). Both *DIV*
(cluster13684_1rr.inclade1.ortho1, module 'grey) and *DICH* (cluster13245_1rr.inclade1.ortho1;
module 'grey') are involved in the dorsoventral asymmetry of flowers (Almeida et al. 1997; Luo
et al. 1999) and were only peripheral in the Bud stage. Flowers at this developmental stage are
lacking or only just beginning to establish asymmetry and the relegation of both *DIV* and *DICH*
to the periphery may reflect this small role. Meanwhile, the homolog of *WUS*
(cluster9273_1rr.inclade1.ortho1; module 'grey') was only peripheral during the D stage. *WUS* is
an important transcription factor that plays an important role to maintain meristem identity and
participates in the vegetative to reproductive transition (Lohmann et al. 2001; Zuo et al. 2002).
The relegation of *WUS* to the periphery at later developmental stages might be expected given
that it plays such a prominent role during early floral development.
542
Evolutionary rates are correlated to network connectivity
Increased protein evolutionary rates during rapid diversification has been suggested in
other radiations of animals and plants (e.g., Kapralov et al. 2013; Brawand et al. 2014; Pease et
al. 2016). Our study is among the first to examine patterns of network evolution across multiple
closely-related species of plants with multiple transitions in floral form and pollination
syndrome. Hummingbirds are hypothesized to have had an important influence on diversification
in Neotropical Gesneriaceae (Roalson and Roberts 2016; Serrano-Serrano et al. 2017b), while
the role of butterflies has not been explored. Modules correlated with hummingbird or butterfly
pollination syndromes did not exhibit clear increases or decreases in evolutionary rates compared
to other modules (Supplemental Table 7). Detecting selection is subject to many factors, such as
a gene's transcriptional abundance and its importance within the protein interaction network
(Lemos et al. 2005). For instance, we found that higher network connectivity was strongly
associated with lower evolutionary rates across both networks in our study (Figure 5;

Supplemental Table 7). This corroborates the recent results found in other eukaryotic systems
(Morandin et al. 2016; Masalia et al. 2017; Josephs et al. 2017; Mähler et al. 2017) while also
showing the apparent strength of correlation between connectivity and evolutionary rate differs
between studies and systems.
Network effects (such as the type of network analysis employed) and the expression
levels of highly abundant genes are known to affect evolutionary rates, potentially confounding
analysis of selection pressure (Krylov et al. 2003). We found that d_N/d_S in early development may
be dependent on both connectivity and expression levels, while this interactive effect was gone
during later development (Supplemental Table 7). As there are few highly connected genes
(hubs, which are important determinants of the observed network structure), a random mutation
would be less likely to affect such a gene. Changing the amino acid sequence of a hub gene
could have multiple associated effects, some deleterious (Hahn and Kern 2004; Luisi et al.
2015). In comparison, a random mutation would be more likely to affect a gene with lower
connectivity, while also resulting in variants with fewer negative consequences. The hub genes
have more direct interactions and are more likely to be essential and involved in more biological
processes, placing them under greater constraint.
572
Numerous network modules correlated to floral form
One primary use for gene co-expression network analyses is to identify modules of co-
expressed genes whose overall expression (as measured by the eigengene) may correlate with
different traits. These traits can be both qualitative or quantitative, such as social caste (Morandin
et al. 2016), seed oil production (Hu et al. 2016), or anther development (Hollender et al. 2014).
Since our dataset was composed of samples from 12 species, we performed correlation analyses
using a phylogenetic model to test whether module expression correlated to different flower
colors, flower shapes, corolla spurs, and pollination syndromes. Nearly half of our modules in
both networks had evidence for an association with these floral traits (Figures 3-4). There were
no modules whose correlation to a pollination syndrome had increased eigengene expression in
all species sharing that trait (Figures 3-4). Instead many modules with correlation to a trait
showed a phylogenetic pattern with a shared increased eigengene expression among the most
closely related species. For example, module 'darkturquoise' in the Bud stage network had a
correlation to butterfly pollination and increased eigengene expression in *A. cettoana* and *A.
longiflora,* both members of Clade 1 (Figure 1). Our aim was to identify sets of co-expressed
genes that would provide candidates for involvement in the development of these traits.
For instance, many of the candidate hubs we identified (see above) were found in
modules that were correlated to one or multiple traits to which they might contribute, such as
PHAN (module 'yellow', correlated to corolla spurs) and *DFR* (module 'darkslateblue',
correlated to hummingbird pollination and purple flowers). Additionally, modules correlated to
various floral traits were enriched for multiple genes that might be involved in the development
of those traits. Module 'palevioletred3' was correlated to corolla spurs in the D stage network
and was enriched for many genes involved in cell division and cell growth (Figure 4;

Supplemental Table 3). Other times, modules were correlated to a specific trait but the enriched
GO terms did not provide a clear idea of how these co-expressed orthogroups may contribute
(Supplemental Tables 2-3). In our analyses, we considered relatively few traits and only focused
on those that have traditionally been considered the most important for the differentiation of
floral form in *Achimenes*, namely flower color, flower shape, and corolla spurs. Undoubtedly
there are a myriad number of molecular and biochemical pathways that we have not considered
here that underlie the development of these complex floral phenotypes. With the idea that
networks represent complex systems of gene and protein interactions, it may be that these
seemingly simple and generalized floral phenotypes are produced through much more complex
means than many have previously considered.
606
Strengths and limitations of network analyses in non-model organisms
Our analysis covers genes with orthologs found in most species, representing a set of
evolutionarily conserved genes. While no reference genome currently exists for Neotropical
Gesneriaceae, we constructed, annotated, and presented de novo transcriptomes for 10
Achimenes species as well as 2 outgroup species, *E. verticillata* and *G. cuneifolia*. Our approach
enables functional genomic studies in this non-model system, but the potential for misassemblies
may bias our downstream results (Hornett and Wheat 2012). Using a gene tree-based approach to
carefully identify orthogroups, we focused our attention on conserved gene regulatory machinery
and not on taxonomically restricted genes. The latter may be involved in species-specific
functions. Nevertheless, it is worth noting that many recent studies have indicated that transitions
in floral form may be due to the shifting expression of genes found in conserved genetic
pathways (e.g., Wessinger and Rausher 2014). Taxonomically restricted genes are likely
interacting with existing regulatory pathways, and the manner in which they integrate into the
co-expression modules identified here will be a fascinating topic for future research. With
extended sampling within *Achimenes* we could infer species-specific co-expression networks and
perform more detailed cross-species comparisons. We expect that these taxonomically restricted
genes may be poorly connected with the network, which may allow for increased diversification.
We characterized gene expression patterns by sequencing RNA across two timepoints
during flower development. Previous gene expression experiments across three timepoints (early
Bud, intermediate D, and late Pre-anthesis stages) in four *Achimenes* species suggested that the
majority of differential gene expression occurred between the early Bud stage and the
intermediate D stage (Roberts and Roalson 2017). We believe our approach allows us to capture
dynamic transcriptional patterns during flower development for comparison across multiple
species. We found that the co-expression networks for the Bud and D stages were largely
preserved, despite there being a few stage specific modules (Supplemental Table 4). However,
this approach may greatly over-simplify the nature of transcriptional regulation in these species.
Organ- and structure-specific expression studies may have greater power to detect floral form-
specific differences, both in terms of the number of differentially expressed genes, and co-
expression network structure (Hollender et al. 2014; Suzuki et al. 2017; Shahan et al. 2018). As a

result, our data likely represents an underestimate of the true number of modules. Future studies

could focus on the comparative analysis of expression patterns in specific organs, such as the

petals, stigma, and stamens, and additional timepoints, which would provide greater functional

insight into the evolution, development, and diversification of these organs.

640   Network analysis represents a more complex approach than differential expression

analyses because it can capture system-level properties (Langfelder and Horvath 2008). Most

phenotypes involve interactions of proteins from diverse biochemical pathways. Although

inferred modules do not necessarily correspond to entire biochemical pathways, or other

components of cellular organization, the approach performs well in reconstructing the

complexity of protein-protein interaction networks (Allen et al. 2012). In many systems this

approach has succeeded in identifying candidate genes for various biological traits, for example

floral organ development in strawberry (Hollender et al. 2014; Suzuki et al. 2018) or carotenoid

accumulation in carrots (Iorizzo et al. 2016). We were able to identify numerous candidate

orthogroups involved in the development of floral form in our non-model system. While no

functional genomic data exists yet for *Achimenes,* our annotations are based on the manually

curated SwissProt database. Applying this network-based approach to the diversification of floral

form across multiple species in *Achimenes*, we recovered distinct modules of co-expressed genes

that may underlie a range of floral phenotypic traits tied to different pollination syndromes.

Although our data come from whole flowers during development, and do not allow us to test

specific hypotheses regarding whether particular genes have shifted expression location in the

developing flower, they do suggest that there are conserved regulatory modules that may have

been co-opted in convergent floral forms multiple times. It will be interesting to apply network

analyses to study floral diversification in additional lineages and examine how conserved or

divergent the patterns are across angiosperms.

Conclusions

662   Floral forms corresponding to bee, butterfly, and hummingbird pollination have evolved

multiple times across the Neotropical plant genus *Achimenes*. Genome-wide gene expression

estimates were taken from flowers in 12 species across two development stages in order to

construct, analyze, and compare stage-specific gene co-expression networks. We hypothesized

that numerous modules in each network would correlate to these pollination syndromes and that

central genes in the network may be candidates for involvement in the development of important

floral traits, such as flower color and shape. We found that nearly a third of modules were

correlated to the pollination syndromes and many more were correlated to different flower colors

and shapes. Several of the hub genes in the networks were homologs of the anthocyanin and

carotenoid biosynthetic pathways, important for the production of floral pigments that attract

different pollinators. A negative relationship between network connectivity and d_N/d_S corroborates

the findings in model systems that more centrally located nodes (likely hubs) are under increased

evolutionary constraint. We found that the less connected genes (peripheral) are under more

relaxed constraints and contained numerous transcriptional regulators. Our results demonstrate

the utility of applying co-expression network analyses in non-model plant lineages to begin
identifying important modules and pathways that will be useful starting points for additional
analyses into the evolution and development of floral form.
679

Acknowledgements

We thank Joanna Kelley, Amit Dhingra, and Andrew McCubbin for helpful comments that
improved this work. Sequencing was performed at the Genomics Core Lab at Washington State
University, Spokane, and the Genomics Sequencing and Analysis Facility at the University of
Texas, Austin.

References

Alexa A, Rahnenführer J, Lengauer T. 2006. Improved scoring of functional groups from gene
expression data by decorrelating GO graph structure. Bioinformatics 22:1600-1607.
Alexandre H, Vrignaud J, Mangin B, Joly S. 2015. Genetic architecture of pollination syndrome
transition between hummingbird-specialist and generalist species in the genus
Rhytidophyllum (Gesneriaceae). PeerJ 3:e1028.
Allen JD, Xie Y, Chen M, Girard L, Xiao G. 2012. Computing statistical methods for
constructing large scale gene networks. PLOS One 7:e29348.
Almeida J, Rocheta M, Galego L. 1997. Genetic control of flower shape in Antirrhinum majus.
Development 124:1387-1392.
Barrett SC. 2013. The evolution of plant reproductive systems: how often are transitions
irreversible? P Roy Soc B-Biol Sci 280: 20130913.
Bolger AM, Lohse M, Usadel B. 2014. Trimmomatic: a flexible trimmer for Illumina sequence
data. Bioinformatics 30:2114-2120.
Bolstad B. 2018. preprocessCore: A collection of pre-processing functions. R package version
1.46.0, https://github.com/bmbolstad/preprocessCore.
Bradshaw HD, Otto KG, Frewen BE, McKay JK, Schemske DW. 1998. Quantitative trait loci
affecting differences in floral morphology between two species of monkeyflower
(Mimulus). Genetics 149:367-382.
Brawand D, Soumillon M, Necsulea A, Julien P, Csárdi G, Harrigan P, Weier M, Liechti A,
Aximu-Petri A, Kircher M, Albert FW. 2011. The evolution of gene expression levels in
mammalian organs. Nature 478:343.
Brawand D, Wagner CE, Li YI, Malinsky M, Keller I, Fan S, Simakov O, Ng AY, Lim ZW,
Bezault E, Turner-Maier J. 2014. The genomic substrate for adaptive radiation in African
cichlid fish. Nature 513:375.
Bray NL, Pimentel H, Melsted P, Pachter L. 2016. Near-optimal probabilistic RNA-seq
quantification. Nat Biotechnol 34:525.
Buchfink B, Xie C, Huson DH. 2014. Fast and sensitive protein alignment using DIAMOND.
Nat Methods 12:59-60.

Des Marais DL, Rausher MD. 2010. Parallel evolution at multiple levels in the origin of hummingbird pollination flowers in Ipomoea. Evolution 64:2044-2054.

De Smet R, Adams KL, Vandepoele K, Van Montagu MC, Maere S, Van de Peer Y. 2013. Convergent gene loss following gene and genome duplications creates single-copy families in flowering plants. Proc Natl Acad Sci USA 110:2898-2903.

Enright AJ, Van Dongen S, Ouzounis CA. 2002. An efficient algorithm for large-scale detection of protein families. Nucleic Acids Res 30:1575-1584.

Faegri K, van der Pijl L. 1979. *Principles of Pollination Ecology*. Oxford: Pergamon Press.

Fenster CB, Armbruster WS, Wilson P, Dudash MR, Thomson JD. 2004. Pollination syndromes and floral specialization. Annu Rev Ecol Evol S 35:375-403.

Gelman A, Rubin DB. 1992. Inference from iterative simulation using multiple sequences. Stat Sci 7:457-472.

Gervasi DD, Schiestl FP. 2017. Real-time divergent evolution in plants driven by pollinators. Nat Commun 8: 14691.

Gómez-Mena C, De Folter S, Costa MMR, Angenent GC, Sablowski R. 2005. Transcriptional program controlled by the floral homeotic gene AGAMOUS during early organogenesis. Development 132:429-438.

Grabherr MG, Haas BJ, Yassour M, Levin JZ, Thompson DA, Amit I, Adiconis X, Fan L, Raychowdhury R, Zeng Q, Chen Z. 2011. Trinity: reconstructing a full-length transcriptome without a genome from RNA-Seq data. Nat Biotechnol 29:644.

Guillerme T, Healy K. 2014. mulTree: a package for running MCMCglmm analysis on multiple trees. ZENODO. 10.5281/zenodo.12902.

Hahn MW, Kern AD. Comparative genomics of centrality and essentiality in three eukaryotic protein-interaction networks. Mol Biol Evol 22:803-806.

Haas BJ, Papanicolaou A, Yassour M, Grabherr M, Blood PD, Bowden J, Couger MB, Eccles D, Li B, Lieber M, MacManes MD. 2013. De novo transcript sequence reconstruction from RNA-seq using the Trinity platform for reference generation and analysis. Nat Protoc 8:1494.

Hadfield JD. 2010. MCMC methods for multi-response generalized linear mixed models: the MCMCglmm R pckage. J Stat Softw 33:1-22.

Hermann K, Klahre U, Venail J, Brandenburg A, Kuhlemeier C. 2015. The genetics of reproductive organ morphology in two Petunia species with contrasting pollination syndromes. Planta 241:1241-1254.

Hoballah ME, Gübitz T, Stuurman J, Broger L, Barone M, Mandel T, Dell'Olivo A, Arnold M, Kuhlemeier C. 2007. Single gene-mediated shift in pollinator attraction in Petunia. Plant Cell 19: 779-790.

Hollender CA, Kang C, Darwish O, Geretz A, Matthews BF, Slovin J, Alkharouf N, Liu Z. 2014. Floral transcriptomes in woodland strawberry uncover developing receptacle and anther gene networks. Plant Physiol 165:1062-1075.

Hornett EA, Wheat CW. 2012. Quantitative RNA-Seq analysis in non-model species: assessing
transcriptome assemblies as a scaffold and the utility of evolutionary divergent genomic
reference species. BMC Genomics 13:361.
Horvath S, Dong J. 2008. Geometric interpretation of gene co-expression network analysis.
PLOS Comput Biol 4:e1000117.
Howarth DG, Martins T, Chimney E, Donoghue MJ. 2011. Diversification of CYCLOIDEA
expression in the evolution of bilateral flower symmetry in Caprifoliaceae and *Lonicera*
(Dipsacales). Ann Bot-London 107:1521-1532.
Hu G, Hovav R, Grover CE, Faigenboim-Doron A, Kadmon N, Page JT, Udall JA, Wendel JF.
2016. Evolutionary conservation and divergence of gene coexpression networks in
Gossypium (cotton) seeds. Genome Biol Evol 8:3765-3783.
Ichihashi Y, Aguilar-Martínez JA, Farhi M, Chitwood DH, Kumar R, Millon LV, Peng J, Maloof
JN, Sinha NR. 2014. Evolutionary developmental transcriptomics reveals a gene network
module regulating interspecific diversity in plant leaf shape. Proc Natl Acad Sci-USA
111:E2616-E2621.
Iorizzo M, Ellison S, Senalik D, Zeng P, Satapoomin P, Huang J, Bowman M, Iovene M,
Sanseverino W, Cavagnaro P, Yildiz M. 2016. A high-quality carrot genome assembly
provides new insights into carotenoid accumulation and asterid genome evolution. Nat
Genet 48:657.
Josephs EB, Wright SI, Stinchcombe JR, Schoen DJ. 2017. The relationship between selection,
network connectivity, and regulatory variation within a population of *Capsella*
grandiflora. Genome Biol Evol 9:1099-1109.
Katoh K, Standley DM. 2013. MAFFT multiple sequence alignment software version 7:
improvements in performance and usability. Mol Biol Evol 30:772-780.
Kapralov MV, Votintseva AA, Filatov DA. 2013. Molecular adaptation during a rapid adaptive
radiation. Mol Biol Evol 30:1051-1059.
Krylov DM, Wolf YI, Rogozin IB, Koonin EV. 2003. Gene loss, protein sequence divergence,
gene dispensability, expression level, and interactivity are correlated in eukaryotic
evolution. Genome Res 13:2229-2235.
Langfelder P, Horvath S. 2008. WGCNA: an R package for weighted correlation network
analysis. BMC Bioinformatics 9:559.
Langfelder P, Luo R, Oldham MC, Horvath S. 2011. Is my network module preserved and
reproducible? PLOS Comput Biol 7:e1001057.
Lemos B, Bettencourt BR, Meiklejohn CD, Hartl DL. 2005. Evolution of proteins and gene
expression levels are coupled in *Drosophila* and are independently associated with
mRNA abundance, protein length, and number of protein-protein interactions. Mol Biol
Evol 22:1345-1354.
Li W, Godzik A. 2006. Cd-hit: a fast program for clustering and comparing large sets of protein
or nucleotide sequences. Bioinformatics 22:1658-1659.

Lohmann JU, Hong RL, Hobe M, Busch MA, Parcy F, Simon R, Weigel D. 2001. A molecular link between stem cell regulation and floral patterning in Arabidopsis. Cell 105:793-803.

Love MI, Huber W, Anders S. 2014. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. Genome Biol 15:550.

Luisi P, Alvarez-Ponce D, Pybus M, Fares MA, Bertranpetit J, Laayouni H. 2015. Recent positive selection has acted on genes encoding proteins with more interactions within the whole human interactome. Genome Biol Evol 7:1141-1154.

Luo D, Carpenter R, Copsey L, Vincent C, Clark J, Coen E. 1999. Control of organ asymmetry in flowers of Antirrhinum. Cell 99:367-376.

Mähler N, Wang J, Terebieniec BK, Ingvarsson PK, Street NR, Hvidsten TR. 2017. Gene co-expression network connectivity is an important determinant of selective constraint. PLOS Genet 13:e1006402.

Martén-Rodríguez S, Quesada M, Castrao AA, Lopezaraiza-Mikel M, Fenster CB. 2015. A comparison of reproductive strategies between island and mainland Caribbean Gesneriaceae. J Ecol 103:1190-1204.

Masalia RR, Bewick AJ, Burke JM. 2017. Connectivity in gene coexpression networks negatively correlates with rates of molecular evolution in flowering plants. PLOS One 12:e182289.

Morandin C, Tin MM, Abril S, Gómez C, Pontieri L, Schiøtt M, Sundström L, Tsuji K, Pedersen JS, Helanterä H, Mikheyev AS. 2016. Comparative transcriptomics reveals the conserved building blocks involved in parallel evolution of diverse phenotypic traits in ants. Genome Biol 17:43.

O'Meara BC, Smith SD, Armbruster WS, Harder LD, Hardy CR, Hileman LC, Hufford L, Litt A, Magallón S, Smith SA, Stevens PF. 2016. Non-equilibrium dynamics and floral trait interactions shape extant angiosperm diversity. P Roy Soc B-Biol Sci 283:20152304.

Paradis E, Claude J, Strimmer K. 2004. APE: analyses of phylogenetics and evolution in R language. Bioinformatics 20:289-290.

Parsana P, Ruberman C, Jaffe AE, Schatz MC, Battle A, Leek JT. 2019. Addressing confounding artifacts in reconstruction of gene co-expression networks. Genome Biol 20:94.

Patterson AH, Bowers JE, Ruggmann R, Dubchak I, Grimwood J, Gundlach H, Haberer G, Hellsten U, Mitros T, Poliakov A, Schmutz J. 2009. The *Sorghum bicolor* genome and the diversification of grasses. Nature 457:551.

Pease JB, Haak DC, Hahn MW, Moyle LC. 2016. Phylogenomics reveals three sources of adaptive variation during a rapid radiation. PLOS Biol 14:e1002379.

Pickrell JK, Marioni JC, Pai AA, Degner JF, Engelhardt BE, Nkadori E, Veyrieras JB, Stephens M, Gilad Y, Pritchard JK. 2010. Understanding mechanisms underlying human gene expression variation with RNA sequencing. Nature 464:768.

Prasad A, Kumar SS, Dessimoz C, Bleuler S, Laule O, Hruz T, Gruissem W, Zimmerman P. 2013. Global regulatory architecture of human, mouse and rat tissue transcriptomes. BMC Genomics 14:716.

Preston JC, Hileman LC, Cubas P. 2011. Reduce, reuse, and recycle: developmental evolution of
trait diversification. Am J Bot 98:397-403.
Price MN, Dehal PS, Arkin AP. 2010. FastTree 2—approximately maximum-likelihood trees for
large alignments. PLOS One 5:e9490.
Quattrocchio F, Wing J, van der Woude K, Souer E, de Vetten N, Mol J, Koes R. 1999.
Molecular analysis of the *anthocyanin2* gene of petunia and its role in the evolution of
flower color. Plant Cell 11:1433-1444.
Raherison ESM, Giguère I, Caron S, Lamara M, MacKay JJ. 2015. Modular organization of the
white spruce (*Picea glauca*) transcriptome reveals functional organization and
evolutionary signatures. New Phytol 207:172-187.
Ramírez-Aguirre E, Martén-Rodríguez S, Quesada-Avila G, Quesada M, Martínez-Díaz Y,
Oyama K, Espinosa-García FJ. 2019. Reproductive isolation among three sympatric
Achimenes species: pre- and post-pollination components. Am J Bot 106:1021-1031.
Ramírez Roa MA. 1987. Revision de *Achimenes* (Gesneriaceae). PhD dissertation, Universidad
Nacional Autonoma de Mexico, Mexico City, Mexico.
Ravasz E, Somera AL, Mongru DA, Oltvai ZN, Barbási AL. 2002. Hierarchical organization of
modularity in metabolic networks. Science 297:1551-1555.
Roalson EH, Roberts WR. 2016. Distinct processes drive diversification in different clades of
Gesneriaceae. Syst Biol 65:662-684.
Roalson EH, Skog LE, Zimmer EA. 2003. Phylogenetic relationships and the diversification of
floral form in *Achimenes* (Gesneriaceae). Syst Bot 28:593-608.
Roberts WR, Roalson EH, 2017. Comparative transcriptome analyses of flower development in
four species of *Achimenes* (Gesneriaceae). BMC Genomics 18:240.
Roberts WR, Roalson EH. 2018. Phylogenomic analyses reveal extensive gene flow within the
magic flowers (*Achimenes*). Am J Bot 105:726-740.
Romero IG, Ruvinsky I, Gilad Y. 2012. Comparative studies of gene expression and the
evolution of gene regulation. Nat Rev Genet 13:505-516.
Rosas-Guerrero V, Aguilar R, Martén-Rodriguez S, Ashworth L, Lopezaraiza-Mikel M, Bastida
JM, Quesada M. 2014. A quantitative review of pollination syndromes: do floral traits
predict effective pollinators? Ecol Lett 17:388-400.
Sauquet H, Magallón S. 2018. Key questions and challenges in angiosperm macroevolution.
New Phytol 219:1170-1187.
Serrano-Serrano ML, Marcionetti A, Perret M, Salamin N. 2017a. Transcriptome resources for
an endemic Neotropical plant lineage (Gesneriaceae). Appl Plant Sci 5:1600135.
Serrano-Serrano ML, Rolland J, Clark JL, Salamin N, Perret M. 2017b. Hummingbird
pollination and the diversification of angiosperms: an old and successful association in
Gesneriaceae. P Roy Soc B-Biol Sci 284:20162816.
Shahan R, Zawora C, Wight H, Sittmann J, Wang W, Mount SM, Liu Z. 2018. Consensus
coexpression network analysis identifies key regulators of flower and fruit development
in wild strawberry. Plant Physiol 178:202-216.

Simão FA, Waterhouse RM, Ioannidis P, Kriventseva EV, Zbodnov EM. 2015. BUSCO:
assessing genome assembly and annotation completeness with single-copy orthologs.
Bioinformatics 31:3210-3212.

Smith SA, Dunn CW. 2008. Phyutility: a phyloinformatics tool for trees, alignments and
molecular data. Bioinformatics 24:715-716.

Smith SD, Rausher MD. 2011. Gene loss and parallel evolution contribute to species difference
in flower color. Mol Biol Evol 28:2799-2810.

Stamatakis A. 2014. RAxML version 8: a tool for phylogenetic analysis and post-analysis of
large phylogenies. Bioinformatics 30:1312-1313.

Stebbins GL. 1970. Adaptive radiation of reproductive characteristics in angiosperms, I:
pollination mechanisms. Annu Rev Ecol Syst 1:307-326.

Stuart JM, Segal E, Koller D, Kim SK. 2003. A gene co-expression network for global discovery
of conserved genetic modules. Science 302:249-255.

Supek F, Bošnjak M, Škunca N, Šmuc T. 2011. REVIGO summarizes and visualizes long lists of
gene ontology terms. PLOS One 6:e21800.

Suyama M, Torrents D, Bork P. 2006. PAL2NAL: robust conversion of protein sequence
alignments into the corresponding codon alignments. Nucleic Acids Res 34:W609-W612.

Suzuki JY, Amore TD, Calla B, Palmer NA, Scully ED, Sattler SE, Sarath G, Lichty JS, Myers
RY, Keith LM, Matsumoto TK. 2017. Organ-specific transcriptome profiling of
metabolic and pigment biosynthetic pathways in the floral ornamental progenitor species
Anthurium amnicola Dressler. Sci Rep-UK 7:1596.

Tang H, Wang X, Bowers JE, Ming R, Alam M, Paterson AH. 2008. Unraveling ancient
hexaploidy through multiply-aligned angiosperm gene maps. Genome Res 18:1944-1954.

Tröbner W, Ramirez L, Motte P, Hue I, Huijser P, Löning WE, Saedler H, Sommer H, Schwarz-
Sommer Z. 1990. GLOBOSA: a homeotic gene which interacts with DEFICIENS in the
control of Antirrhinum floral organogenesis. EMBO J 11:4693-4704.

Uebbing S, Künster A, Mäkinen H, Backström N, Bolivar P, Burri R, Dutoit L, Mugal CF, Nater
A, Aken B, Flicek P. 2016. Divergence in gene expression within and between two
closely related flycatcher species. Mol Ecol 25:2015-2028.

The UniProt Consortium. 2017. UniProt: the universal protein knowledgebase. Nucleic Acids
Res 45:D158-D169.

Van der Niet T, Johnson SD. 2012. Phylogenetic evidence for pollinator-driven diversification of
angiosperms. Trends Ecol Evol 27: 353-361.

Waites R, Selvadurai HR, Oliver IR, Hudson A. 1998. The PHANTASTICA gene encodes a
MYB transcription factor involved in growth and dorsoventrality of lateral organs in
Antirrhinum. Cell 93:779-789.

Wessinger CA, Rausher MD. 2014. Ecological transition predictably associated with gene
degeneration. Mol Biol Evol 32:347-354.

West-Eberhard MJ. 1989. Phenotypic plasticity and the origins of diversity. Ann Rev Ecol Syst
20:249-278.

Yang Y, Smith SA. 2014. Orthology inference in nonmodel organisms using transcriptomes and
914   low-coverage genomes: improving accuracy and matrix occupancy for phylogenomics.
915   Mol Biol Evol 31:3081-3092.
916   Yang Z. 2007. PAML 4: phylogenetic analysis by maximum likelihood. Mol Biol Evol 24:1586-
917   1591.
918   Yant L, Collani S, Puzey J, Levy C, Kramer EM. 2015. Molecular basis for three-dimensional
919   elaboration of the Aquilegia petal spur. P Roy Soc B-Biol Sci 282:20142778.
920   Yuan YW, Rebocho AB, Sagawa JM, Stanley LE, Bradshaw HD. 2016. Competition between
921   anthocyanin and flavonol biosynthesis produces spatial pattern variation of floral
922   pigments between Mimulus species. Proc Natl Acad Sci USA 113: 2448-2453.
923   Zuo J, Niu Q-W, Frugis G, Chua N-H. 2002. The WUSCHEL gene promotes vegetative-to-
924   embryonic transition in Arabidopsis. Plant J 30:349-359.
925

Tables

Table 1. Gene Ontology enrichment of network hubs.

Table 2. Gene Ontology enrichment of network periphery.

Figures

Figure 1. Phylogenetic relationships in *Achimenes* and floral form. (A) Phylogeny of *Achimenes* adapted from Roberts and Roalson (2018). Flowers of each species are shown alongside the pollination syndrome (bees, butterflies, or hummingbirds). Clades 1 and Clades 2 (sensu Roalson et al. 2003) are indicated on the phylogeny with a circle and box, respectively. (B) Two stages of flower development were sampled for each of the 12 species, an early Bud stage and an intermediate D stage.

Figure 2. Co-expression clustering and the correspondence between the Bud and D stage networks. (A) Hierarchical clustering by WGCNA showing co-expressed modules for the Bud stage and (B) the D stage. Each of the 9503 orthogroups were assigned to 65 and 62 modules, respectively, and the color rows underneath show the module assignment. The 'grey' module is for unassigned orthogroups. (C) Correspondence of Bud stage modules and D stage modules. Each row corresponds to one Bud stage module (labeled by color and module size), and each column corresponds to one D stage module (labeled by color and module size). Numbers in the table indicate gene counts in the intersection of the corresponding modules. Coloring of the table encodes -log(p), with p being the Fisher's exact test p-value for the overlap of the two modules. The stronger the red color, the more significant the overlap.

Figure 3. Summary of Bud stage network module association to floral traits and pollination syndrome. Each row represents an individual module in the Bud stage network and shows the total module size (column All) and the number of hubs (column Hubs). Each column in the central table represents the floral traits (Pollination Syndrome, Flower color, Flower shape, and Spurs) that were tested for an association to module eigengenes. The eigengene expression is indicated for each module in which all three replicates from a species either had > 1.0 or < -1.0 expression. An association was considered 'Strongly positive' when both the eigengene expression was > 1.0 in species that had the trait and MCMCglmm results were significant. An association was considered 'Strongly negative' when both the eigengene expression was < -1.0 in species that had the trait and MCMCglmm results were significant. Associations were considered 'Weakly positive' or 'Weakly negative' when only MCMCglmm results were significant.

Figure 4. Summary of D stage network module association to floral traits and pollination syndrome. Each row represents an individual module in the D stage network and shows the total module size (column All) and the number of hubs (column Hubs). Each column in the central table represents the floral traits (Pollination Syndrome, Flower color, Flower shape, and Spurs) that were tested for an association to module eigengenes. The eigengene expression is indicated for each module in which all three replicates from a species either had > 1.0 or < -1.0 expression.

An association was considered 'Strongly positive' when both the eigengene expression was > 1.0 in species that had the trait and MCMCglmm results were significant. An association was considered 'Strongly negative' when both the eigengene expression was < -1.0 in species that had the trait and MCMCglmm results were significant. Associations were considered 'Weakly positive' or 'Weakly negative' when only MCMCglmm results were significant.

Figure 5. Relationship between orthogroup dN/dS (omega) and orthogroup connectivity. (A) Bud stage network. (B) D stage network. Each point indicates an individual orthogroup. The red line indicates the linear regression line for the relationship between dN/dS and connectivity. The dotted line indicates dN/dS = 1 with orthogroups above this line considered to be under relaxed or positive selection.

Supplemental Files

Suppl. Table 1. Sequencing and transcriptome assembly results.

Suppl. Table 2. Gene Ontology enrichment for the Bud stage modules.

Suppl. Table 3. Gene Ontology enrichment for the D stage modules.

Suppl. Table 4. MedianRank and Zsummary module preservation scores.

Suppl. Table 5. Orthogroups identified as hubs, including module membership and annotation.

Suppl. Table 6. Orthogroups identified as periphery, including module membership and annotation.

Suppl. Table 7. Statistical results of the evolutionary rates analyses.

Suppl. Figure 1. Module eigengenes in the Bud stage network. Plots show the module eigengenes for each module in the Bud stage network. Each bar represents the eigengene expression (y-axis) for each sample in the Bud stage dataset (x-axis). Eigengene expression represents the expression of a sample within a module.

Suppl. Figure 2. Module eigengenes in the D stage network. Plots show the module eigengenes for each module in the D stage network. Each bar represents the eigengene expression (y-axis) for each sample in the D stage dataset (x-axis). Eigengene expression represents the expression of a sample within a module.

Suppl. Figure 3. Effects of floral traits on the eigengene expression in the Bud stage modules summarized over 50 phylogenetic trees. Each plot shows a summary of the important floral predictors (flower color, flower shape, and corolla spur) for each module in the Bud stage network. The posterior distribution of an independent variable with negligible effect on

eigengene expression is centered around zero (dashed line). The distribution of an influential variable is expected to be substantially different from zero. Black dots indicate the mode of the posterior distribution, and boxes indicate the 50 and 95% credible intervals.

Suppl. Figure 4. Effects of pollination syndrome on the eigengene expression in the Bud stage modules summarized over 50 phylogenetic trees. Each plot shows a summary of the important pollination syndrome predictors for each module in the Bud stage network. The posterior distribution of an independent variable with negligible effect on eigengene expression is centered around zero (dashed line). The distribution of an influential variable is expected to be substantially different from zero. Black dots indicate the mode of the posterior distribution, and boxes indicate the 50 and 95% credible intervals.

Suppl. Figure 5. Effects of floral traits on the eigengene expression in the D stage modules summarized over 50 phylogenetic trees. Each plot shows a summary of the important floral predictors (flower color, flower shape, and corolla spur) for each module in the D stage network. The posterior distribution of an independent variable with negligible effect on eigengene expression is centered around zero (dashed line). The distribution of an influential variable is expected to be substantially different from zero. Black dots indicate the mode of the posterior distribution, and boxes indicate the 50 and 95% credible intervals.

Suppl. Figure 6. Effects of pollination syndrome on the eigengene expression in the D stage modules summarized over 50 phylogenetic trees. Each plot shows a summary of the important pollination syndrome predictors for each module in the D stage network. The posterior distribution of an independent variable with negligible effect on eigengene expression is centered around zero (dashed line). The distribution of an influential variable is expected to be substantially different from zero. Black dots indicate the mode of the posterior distribution, and boxes indicate the 50 and 95% credible intervals.

Suppl. Figure 7. Relationship between orthogroup dN/dS (omega) and average expression. (A) Bud stage. (B) D stage. Each point indicates an individual orthogroup. The red line indicates the linear regression line between fit for dN/dS ~ expression. The dotted line indicates dN/dS = 1, where orthogroups with dN/dS > 1 may be under stronger relaxed selection.

Suppl. Figure 8. Distribution of dN/dS (omega) among hubs and periphery nodes. (A) dN/dS for hub and non-hub nodes in the Bud stage network. (B) dN/dS for hub and non-hub nodes in the D stage network. (C) dN/dS for periphery and non-periphery nodes in the Bud stage network. (D) dN/dS for periphery and non-periphery nodes in the D stage network. Violin plots show the distribution of dN/dS values for each category. Horizontal lines in each distribution indicate the median dN/dS value.

Table 1. Gene Ontology enrichment of network hubs.

	GO.ID	Term	Annotated	Significant	Expected	P
A. Bud stage hubs	GO:0031325	positive regulation of cellular metabolic process	224	16	14.52	0.000
	GO:0006607	NLS-bearing protein import into nucleus	7	4	0.45	0.001
	GO:0015991	ATP hydrolysis coupled proton transport	9	4	0.58	0.002
	GO:0046470	phosphatidylcholine metabolic process	9	4	0.58	0.002
	GO:0018298	protein-chromophore linkage	32	7	2.07	0.004
	GO:0070507	regulation of microtubule cytoskeleton organization	6	3	0.39	0.005
	GO:1990778	protein localization to cell periphery	6	3	0.39	0.005
	GO:0051592	response to calcium ion	7	3	0.45	0.008
	GO:0006412	translation	233	24	15.10	0.009
	GO:0042594	response to starvation	60	10	3.89	0.012
	GO:0007035	vacuolar acidification	8	3	0.52	0.012
	GO:0006534	cysteine metabolic process	9	4	0.58	0.012
	GO:0009833	plant-type primary cell wall biogenesis	15	4	0.97	0.013
	GO:0048831	regulation of shoot system development	123	12	7.97	0.017
	GO:0032509	endosome transport via multivesicular body sorting pathway	9	3	0.58	0.017
	GO:0051568	histone H3-K4 methylation	9	3	0.58	0.017
	GO:0000271	polysaccharide biosynthetic process	123	11	7.97	0.023
	GO:0006732	coenzyme metabolic process	116	8	7.52	0.023
	GO:0010338	leaf formation	5	2	0.32	0.037
	GO:0048564	photosystem I assembly	5	2	0.32	0.037
	GO:0017001	antibiotic catabolic process	26	3	1.69	0.037
	GO:0044282	small molecule catabolic process	75	10	4.86	0.038
	GO:0006833	water transport	13	3	0.84	0.048
	GO.ID	**Term**	**Annotated**	**Significant**	**Expected**	**P**
B. D stage hubs	GO:0006412	translation	233	30	16.03	0.001
	GO:0042939	tripeptide transport	8	4	0.55	0.001
	GO:0016117	carotenoid biosynthetic process	20	6	1.38	0.002
	GO:0006744	ubiquinone biosynthetic process	5	3	0.34	0.003
	GO:0032259	methylation	88	15	6.05	0.004
	GO:0045814	negative regulation of gene expression, epigenetic	37	5	2.55	0.005
	GO:0009206	purine ribonucleoside triphosphate biosynthetic process	44	6	3.03	0.013
	GO:0000271	polysaccharide biosynthetic process	123	11	8.46	0.013
	GO:0009873	ethylene-activated signaling pathway	78	12	5.37	0.014
	GO:0015991	ATP hydrolysis coupled proton transport	9	3	0.62	0.020
	GO:0009718	anthocyanin-containing compound biosynthetic process	26	5	1.79	0.030
	GO:0000018	regulation of DNA recombination	5	2	0.34	0.041
	GO:0015976	carbon utilization	5	2	0.34	0.041
	GO:0034314	Arp2/3 complex-mediated actin nucleation	5	2	0.34	0.041
	GO:0080144	amino acid homeostasis	5	2	0.34	0.041
	GO:1901615	organic hydroxy compound metabolic process	121	9	8.32	0.041
	GO:0000902	cell morphogenesis	174	9	11.97	0.042

Table 2. Gene Ontology enrichment of network periphery.

	GO.ID	Term	Annotated	Significant	Expected	P
A. Bud stage periphery	GO:0016579	protein deubiquitination	24	16	8.70	0.002
	GO:0048825	cotyledon development	21	12	7.61	0.005
	GO:0015846	polyamine transport	5	5	1.81	0.006
	GO:0034314	Arp2/3 complex-mediated actin nucleation	5	5	1.81	0.006
	GO:0006020	inositol metabolic process	12	7	4.35	0.011
	GO:0010229	inflorescence development	9	7	3.26	0.014
	GO:0009744	response to sucrose	31	16	11.24	0.021
	GO:1903338	regulation of cell wall organization or biogenesis	11	7	3.99	0.026
	GO:0006809	nitric oxide biosynthetic process	6	5	2.18	0.026
	GO:0009773	photosynthetic electron transport in photosystem I	8	6	2.90	0.030
	GO:0034613	cellular protein localization	183	75	66.34	0.031
	GO:0006004	fucose metabolic process	12	8	4.35	0.032
	GO:0009926	auxin polar transport	42	22	15.23	0.037
	GO:0010090	trichome morphogenesis	38	19	13.78	0.041
	GO:0009414	response to water deprivation	182	69	65.98	0.045
	GO:0009873	ethylene-activated signaling pathway	78	38	28.28	0.046
	GO:0007166	cell surface receptor signaling pathway	93	42	33.71	0.046
	GO:0006163	purine nucleotide metabolic process	96	38	34.80	0.047
	GO:0035195	gene silencing by miRNA	15	7	5.44	0.048
	GO:0016236	macroautophagy	21	9	7.61	0.048
	GO:0055083	monovalent inorganic anion homeostasis	13	5	4.71	0.048
	GO:0051338	regulation of transferase activity	38	13	13.78	0.048
	GO:0006473	protein acetylation	23	7	8.34	0.048
	GO.ID	Term	Annotated	Significant	Expected	weight01
B. D stage periphery	GO:0045490	pectin catabolic process	20	19	12.88	0.002
	GO:0007166	cell surface receptor signaling pathway	93	71	59.90	0.006
	GO:0000398	mRNA splicing, via spliceosome	46	38	29.63	0.007
	GO:0009740	gibberellic acid mediated signaling pathway	31	24	19.97	0.015
	GO:0006310	DNA recombination	54	41	34.78	0.015
	GO:0009909	regulation of flower development	80	57	51.53	0.017
	GO:0009058	biosynthetic process	2121	1360	1366.19	0.019
	GO:0042545	cell wall modification	34	26	21.90	0.021
	GO:0001708	cell fate specification	13	12	8.37	0.027
	GO:0042538	hyperosmotic salinity response	25	21	16.10	0.027
	GO:0002181	cytoplasmic translation	21	18	13.53	0.029
	GO:0006897	endocytosis	40	27	25.76	0.029
	GO:0010197	polar nucleus fusion	20	17	12.88	0.039
	GO:0071702	organic substance transport	492	319	316.91	0.046
	GO:0006949	syncytium formation	7	7	4.51	0.046
	GO:0008295	spermidine biosynthetic process	7	7	4.51	0.046
	GO:0009963	positive regulation of flavonoid biosynthetic process	7	7	4.51	0.046

A

Pollinator group

Bees
(Apidae)

Butterflies
(Lepidoptera)

Hummingbirds
(Trochilidae)

Clade 1

Clade 2

G. cuneifolia

E. verticillata

A. misera

A. admirabilis

A. erecta

A. longiflora

A. cettoana

A. candida

A. antirrhina

A. pedunculata

A. grandiflora

A. patens

0.006

B

Bud stage

D stage

Scale bar equals 1 cm.

Figure 1. Phylogenetic relationships in *Achimenes* and floral form.

Figure 2. Co-expression clustering and the correspondence between the Bud and D stage networks.

Figure 3. Summary of Bud stage network module association to floral traits and pollination syndrome.

Figure 4. Summary of D stage network module association to floral traits and pollination syndrome.

Figure 5. Relationship between orthogroup dN/dS (omega) and orthogroup connectivity.

Supplemental Figure 1. Module eigengenes in the Bud stage network.

Supplemental Figure 2. Module eigengenes in the D stage network.

Supplemental Figure 3. Effects of floral traits on the eigengene expression in the Bud stage modules summarized over 50 phylogenetic trees.

Supplemental Figure 4. Effects of pollination syndrome on the eigengene expression in the Bud stage modules summarized over 50 phylogenetic trees.

Supplemental Figure 5. Effects of floral traits on the eigengene expression in the D stage modules summarized over 50 phylogenetic trees.

Supplemental Figure 6. Effects of pollination syndrome on the eigengene expression in the D stage modules summarized over 50 phylogenetic trees.

Supplemental Figure 7. Relationship between orthogroup dN/dS (omega) and average expression.

Supplemental Figure 8. Distribution of dN/dS (omega) among hubs and periphery nodes.