Co-expression clustering across flower development identifies modules for diverse floral forms in *Achimenes* (Gesneriaceae)

Wade R Roberts[1,2], Eric H Roalson[1]

[1] School of Biological Sciences, Washington State University, Pullman, WA, United States
[2] Current address: Biological Sciences, University of Arkansas, Fayetteville, AR, United States

Corresponding Author:
Wade Roberts[1]
Biological Sciences, University of Arkansas, SCEN 601, Fayetteville, AR, 72701, United States
Email address: wader@uark.edu

Abstract

Background. Genetic pathways involved with flower color and shape are thought to play an important role in the development of flowers associated with different pollination syndromes, such as those associated with bee, butterfly, or hummingbird pollination. Because pollination syndromes are complex traits that are orchestrated by multiple genes and pathways, the gene networks have not been explored. Gene co-expression networks provide a systems level approach to identify important contributors to floral diversification.

Methods. RNA-sequencing was used to assay gene expression across two stages of flower development (an early bud and an intermediate stage) in 10 species of *Achimenes* (Gesneriaceae). Two stage-specific co-expression networks were created from 9503 orthologs and analyzed to identify module hubs and the network periphery. Module association with bee, butterfly, and hummingbird pollination syndromes was tested using phylogenetic mixed models. The relationship between network connectivity and evolutionary rates (d_N/d_S) was tested using linear models.

Results. Networks contained 65 and 62 modules that were largely preserved between developmental stages and contained few stage-specific modules. Over a third of the modules in both networks were associated with flower color, shape, and pollination syndrome. Within these modules, several hub nodes were identified that related to the production of anthocyanin and carotenoid pigments and the development of flower shape. Evolutionary rates were decreased in highly connected genes and elevated in peripheral genes.

Discussion. This study aids in the understanding of the genetic architecture and network properties underlying the development of floral form and provides valuable candidate modules and genes for future studies.

Introduction

Flowers display a diverse range of colors, shapes, and sizes, and understanding the ecological and genetic factors contributing to their diversity across angiosperms has long been a major goal in biology. Often this diversity has been attributed to pollinator-mediated selection (Stebbins 1970; O'Meara et al. 2016; Gervasi and Schiestle 2017) and the relationship between a plant and its mode of pollination is considered to be one of the key innovations contributing to angiosperm diversification (Faegri and van der Pijl 1979; Fenster et al. 2004; van der Niet and Johnson 2012; Barrett 2013; Sauquet and Magallón 2018). Most plants evolved to rely on biotic or abiotic means in order to move pollen and ensure reproductive success (Faegri and van der Pijl 1979; Fenster et al. 2004). These biotic pollinators are often attracted to flowers that contain specific traits, such as red flowers that provide high nectar rewards for bird visitors (Cronk and Ojeda 2008) or the ultraviolet markings on some flowers that attract bee visitors (Papiorek et al 2016). A wide range of floral traits are thought to contribute to successful pollination, such as color (Sletvold et al. 2016), odor (Piechowski et al. 2010), nectar composition (Amorim et al. 2013), and time of flowering (Cortés-Flores et al. 2017). The genetic basis for these and other traits and their role in floral and pollination syndrome divergence has been examined extensively in model systems, particularly in *Mimulus* (Bradshaw et al. 1998; Yuan et al. 2016) and *Petunia* (Hoballah et al. 2007; Hermann et al. 2015), and more recently in many non-model systems (Wessinger et al. 2014; Alexandre et al. 2015).

Among the most widely used large-scale experimental approaches to investigate genome function are transcriptome analyses (Pickrell et al. 2010; Raherison et al. 2015), particularly in non-model organisms where no reference genome or functional genomics data exists. Changes in expression often result from the combinatorial action of genetic regulatory pathways orchestrating development and responses to environmental stimuli. Therefore, the transcriptome can be viewed as a link between the genotype and phenotype and may be acted upon through selection (Romero et al. 2012; Prasad et al. 2013). As changes in gene expression may underlie many of the phenotypic changes between species (Brawand et al. 2011; Romero et al. 2012; Uebbing et al. 2016), studying the transcriptome may shed light on important pathways and targets of selection. Phenotypic changes can frequently arise through functional changes in conserved developmental pathways among closely related species. For example, changes in enzyme function and transcriptional regulation of the anthocyanin pathway has been implicated frequently across angiosperms in the evolution of red flowers (Des Marais and Rausher 2010; Smith and Rausher 2011). It is now recognized that most genes act as members of biological pathways or of co-regulated modules (Hollender et al. 2014; Ma et al. 2018).

Here, we undertake a comparative study of gene expression during flower development in the genus *Achimenes*. This group is a member of the diverse African violet family (Gesneriaceae) and distributed throughout Mexico and Central America. *Achimenes* is a young lineage (*c*. 7-12 Mya; Roalson and Roberts 2016) known for its floral diversity (Figure 1), a

feature thought to be associated with speciation (Ramírez Roa 1987; Roalson et al. 2003). Four
pollination syndromes are found in *Achimenes,* including melittophily (bees), psychophily
(butterflies), euglossophily (female euglossine bees), and ornithophily (hummingbirds) (Figure
2). These syndromes have traditionally been defined on the basis of flower color and flower
shape (Figure 2; Ramírez Roa 1987) and recently through pollinator observations (Martén-
Rodríguez et al. 2015; Ramírez-Aguirre et al. 2019). Currently there have been pollinator
observations made for four *Achimenes* species: *A. antirrhina* (hummingbirds, *Amazilia*
beryllina), *A. flava* (bees, Anthophoridae), *A. obscura* (bees, *Trigona fulviventris*), and *A. patens*
(unidentified butterflies) (Martén-Rodriguez et al. 2015; Ramírez-Aguirre et al. 2019). Based on
these observations and the floral traits of these species (Figure 2), we can hypothesize the likely
pollinators for the other *Achimenes* species (Figure 2; Table 1). Repeated origins and transitions
between these pollination syndromes have been hypothesized in *Achimenes* (Roalson et al.
2003), making this small lineage an attractive system to understand the genetic and ecological
factors underlying floral diversification.
In this study, we aimed to characterize and compare patterns of gene co-expression
during flower development across 10 *Achimenes* species. So far, few studies have investigated
the genetic basis of floral development in the Gesneriaceae family (Alexandre et al. 2015;
Roberts and Roalson 2017). Given that bee, butterfly, and hummingbird pollination syndromes
evolved in parallel across *Achimenes,* we hypothesized that: (1) distinct sets of co-expressed
genes would correlate to each syndrome during flower development, and (2) genes in pathways
involved with flower color and shape may be central in each network during flower
development. Using a comparative transcriptome approach, we clustered a shared set of 9503
orthologs into two different co-expression networks, each corresponding to two stages of flower
development: an early bud stage and an intermediate stage prior to anthesis. This strategy
allowed us to compare gene co-expression patterns temporally. We compared these networks and
tested whether gene co-expression clusters correlated with different floral traits (pollination
syndrome, flower color, flower shape, corolla spurs). We also tested whether network properties,
such as connectivity and expression, influence evolutionary constraints. Our results allowed us to
quantify the extent of shared gene co-expression among closely related species and to identify
pathways and genes that may be associated with the repeated evolution of flower types
underlying pollinator specificity.
110
Materials & Methods
112
Plant materials
Ten of the 26 currently recognized species in *Achimenes* were sampled: five from Clade 1
and five from Clade 2 (Figure 2). These species were chosen based on their hypothesized close
relationships and their diversity in floral form (Figure 1; Roalson et al. 2003). On the basis of
previous molecular studies (Roalson and Roberts 2016), two species were chosen as outgroups to
represent related lineages: *Eucodonia verticillata* and *Gesneria cuneifolia* (Figures 1-2). All

selected species were grown in standard greenhouse conditions at Washington State University,
under 16-hour days, 24-27°C, and 80-85% humidity.
Whole flower buds from two stages of flower development were sampled for each
species: an immature bud stage (Bud stage; Figure 1) and an intermediate stage before anthesis
(D stage; Figure 1). Our definitions of the floral developmental stages in *Achimenes* were
adapted from the stages determined for *Antirrhinum majus*, which uses a morphological and
temporal framework (Vincent and Enrico 2004). The Bud stage represents the smallest
reproductive bud distinguishable from vegetative buds, having lengths between 1-2 mm (Figure
1). The appearance is marked by the emergence of the floral primordia, no accumulation of
anthocyanin pigments, and no elongation or growth of corolla tissue cells. The D stage represents
a halfway point between the Bud and a pre-anthesis flower, where the length is about half the
length of a fully developed flower (Figure 1). Its appearance is marked by the accumulation of
anthocyanins in the corolla tissue, the cells have elongated in the corolla tube, and trichomes
have emerged. Corolla spurs are found only in *A. grandiflora* and *A. patens*, and the spur begins
development during the D stage as an outward growth of the corolla in the opposite direction
from the tube opening (Figure 1).
Each stage was sampled with two or three biological replicates, contributing four or six
samples for each species. The bud stage was chosen to represent a baseline level of gene
expression when floral phenotypes are largely similar between species that could then be
compared to the intermediate stage where floral phenotypes diverge and many important
pathways involved in flower development and pigmentation show increased expression levels
(Roberts and Roalson 2017). All tissues were immediately frozen into liquid nitrogen and stored
at -80°C.
142

RNA extraction and RNA-seq library construction
Total RNA was extracted from frozen tissue using the RNEasy Plant Kit (Qiagen) and
ribosomal depleted RNA samples were prepared using the RiboMinus Plant Kit (Thermo Fisher
Scientific). Stranded libraries were prepared using the NEBNext Ultra Directional RNA Library
Kit (New England Biolabs), barcoded, and pooled based on nanomolar concentrations. Library
quality and quantity were assessed using the Qubit (dsDNA HS assay; Thermo Fisher Scientific)
and BioAnalyzer 2100 (Agilent). The library pool was sequenced across 4 lanes of an Illumina
HiSeq2500 for paired-end 101 bp reads at the Genomics Core Lab at Washington State
University, Spokane. All new sequence data were deposited in the NCBI Sequence Read Archive
(BioProject: PRJNA435759).
Single replicate libraries for four of the species in the current study (*A. cettoana, A.*
erecta, A. misera, and *A. patens*) were previously sequenced for the same two timepoints (Bud
and D stages) (Roberts and Roalson 2017). These samples were combined with the newly
generated data to assembly de novo transcriptomes and perform co-expression clustering. This
data was deposited in the NCBI Sequence Read Archive (BioProject: PRJNA340450).
158

Transcriptome assembly

Per base quality, adapter contamination, and per base sequence content of raw reads was assessed with FastQC tools (http://bioinformatics.babraham.ac.uk/projects/fastqc). One library (AT6) returned poor yields and low-quality reads and was excluded from further analyses. Trimmomatic v.0.36 (Bolger et al. 2014) was then used to remove contaminating adapter sequences and dynamically trim low-quality bases (ILLUMINACLIP:TruSeq3-PE-2.fa:2:30:10 HEADCROP:13 LEADING:3 TRAILING:3 SLIDINGWINDOW:4:15 MINLEN:50). Trimmed reads derived from an individual species were combined and de novo assembled into contigs using Trinity (--SS_lib_type RF; Grabherr et al. 2011; Haas et al. 2013), generating a reference transcriptome for each species. Sequence redundancy was reduced with CDHIT-EST (-c 0.99 -n 5; Li and Godzik 2006). Open reading frames were then predicted for each transcriptome assembly with TransDecoder (Haas et al. 2013) using a BLASTp search against the SwissProt database (www.uniprot.org) to maximize sensitivity. Finally, transcriptome assembly completeness was assessed using BUSCO (Simão et al. 2015) against a set of single-copy plant orthologs (embryophyte_odb10 dataset).

Orthogroup inference

To help facilitate orthogroup inference, transcriptome assemblies from flowers in *Sinningia eumorpha* and *S. magnifica* were downloaded and combined with our assemblies (https://doi.org/10.5061/dryad.4r5p1; Serrano-Serrano et al. 2017a). *Sinningia* is closely related to our focal lineage in the Neotropical Gesneriaceae (Roalson and Roberts 2016). To identify orthogroup sequences across all 14 species, we used a modified version of the approach described in Yang and Smith (2014). All peptide sequences were clustered using an all-by-all BLASTp search performed using DIAMOND (--evalue 1e-6; Buchfink et al. 2014) and results were clustered using MCL v14-137 (-I 1.4; Enright et al. 2002). Homolog clusters were aligned using MAFFT v7.271 (--genafpair --maxiterate 1000; Katoh and Standley 2013) and alignments were trimmed using Phyutility v2.7.1 (-clean 0.1; Smith and Dunn 2008). Codon alignments were then produced for each cluster using PAL2NAL v14 (Suyama et al. 2006) and maximum-likelihood trees were constructed using FastTree v2.1.8 (Price et al. 2010) using the GTR model. Trees were then trimmed to exclude branches >10 times longer than its sister or longer than 0.2 substitutions per site. Final homolog group alignments were created from the trimmed trees using MAFFT and used for final homolog tree inference using RAxML v8.2.9 (Stamatakis 2014) and the GTRGAMMA model. Orthogroups were inferred from the final homolog trees using the rooted ingroup (RT) method of Yang and Smith (2014), using *S. eumorpha* and *S. magnifica* as outgroups. Homolog trees were trimmed to include at least 2 ingroup species, resulting in 83595 orthogroups.

Quantifying gene expression

Trimmed reads for each sample were aligned to the corresponding transcriptome assembly to quantify expression using default parameters in Kallisto v0.43.0 (Bray et al. 2016).

The resulting counts for each contig were matched with the inferred orthogroups derived above,
creating two separate matrices for the Bud and D stage samples. In the resulting gene expression
matrices, rows corresponded to orthogroups and columns corresponded to individual samples
(Supplemental Data). Counts for the Bud and D stage samples were separately transformed using
variance-stabilizing transformation implemented in the Bioconductor package DESeq2 (Love et
al. 2014) and quantile normalized using the Bioconductor package preprocessCore (Bolstad
2018). Lastly, counts for both stages were separately corrected for confounding effects, such as
batch or species effects, using principal component regression with the 'sva_network' function in
the Bioconductor package sva (Parsana et al. 2019).

Co-expression networks and identification of modules
The Bud and D stage RNAseq data was clustered into gene co-expression networks using
the R package WGCNA (Langfelder and Horvath 2008). Prior to network construction,
orthogroups were filtered to remove any with >50% missing values or <0.3 expression variance.
In total, 9503 orthogroups common to both Bud and D stage data were used to construct separate
networks for each stage. Parameters for the Bud stage network were as follows: power = 18,
networkType = "signed", corType = "bicor", maxPOutliers = 0.05, TOMType = "signed",
deepSplit = 2, mergeCutHeight = 0.25. For the D stage network, the following parameters were
used: power = 16, networkType = "signed", corType = "bicor", maxPOutliers = 0.05, TOMType
= "signed", deepSplit = 2, mergeCutHeight = 0.25. Soft-thresholding powers for each network
("power" parameter) were chosen as the lowest value such that the scale free topology model fit
(R^2) was ≥ 0.9 (Supplemental Data). Signed networks consider positively correlated nodes
connected, while unsigned networks consider both positively and negatively correlated nodes
connected (van Dam et al. 2018). We used signed networks here ("networkType" and
"TOMType" parameters) because this type is considered more robust to biological functions
with more specific expression patterns (Mason et al. 2009; Song et al. 2012). Unsigned networks
can capture only strong correlations and many negative regulatory relationships in biological
systems are weak or moderate (Ritchie et al. 2009). Biweight midcorrelation ("corType"
parameter) was used as this method is often more robust than Pearson correlation and more
powerful than Spearman correlation (Song et al. 2012). All other WGCNA parameters were kept
at their default values.
Module eigengenes and orthogroup connectivity were calculated in each network
separately using the 'moduleEigengenes' and 'intramodularConnectivity' functions in WGCNA,
respectively. The module eigengene is defined as the first principal component of a module and
represents the gene expression profile of a sample within a module (Langfelder and Horvath
2008). Connectivity refers to the sum of connection strengths a node has to other network nodes.
Module hub genes (defined as the most highly connected nodes within a module) were
identified in both networks based on module membership (kME) scores > 0.9, calculated using
the 'signedKME' function of WGCNA. kME is defined as the correlation between the
expression levels of a gene and the module eigengene (Horvath and Dong 2008). Additionally,

using connectivity measures, we defined peripheral genes in each network as the lowest 10% of
connected nodes.
We tested whether Bud stage modules were preserved in the D stage and vice versa using
module preservation statistics in WGCNA (Langfelder et al. 2011). Median Rank and Zsummary
statistics were computed using the 'modulePreservation' function of WGCNA, using 200
permutations (Langfelder et al. 2011). Median rank statistics were compared to the "gold"
module, which consists of 1000 randomly selected orthogroups. Modules with a lower median
rank exhibit stronger preservation than modules with a higher median rank. Zsummary scores >
10 indicate strong evidence of preservation and scores < 2 indicate no evidence of preservation.
248
Phylogeny
We previously inferred a phylogeny for the 12 sampled species using 1306 single-copy
orthologs identified from the same transcriptome dataset used here (Roberts and Roalson 2018).
For comparative analyses of module-trait relationships, we randomly sampled 50 single-copy
ortholog gene trees and rescaled branch lengths to be proportional to time (ultrametric) using the
'chronos' function in the R package ape (Paradis et al. 2004). Bootstrap support was 100 for
nearly every branch in the Roberts and Roalson (2018) phylogeny, therefore we chose to use
randomly sampled ortholog trees to account for phylogenetic uncertainty.
257
Module and trait relationship
We correlated external traits (e.g., red flower color) with the module eigengenes using a
phylogenetic Markov Chain Monte Carlo method, implemented in the R packages MCMCglmm
(Hadfield 2010) and mulTree (Guillerme and Healy 2014). These analyses were performed
individually for each module in both the Bud and D stage networks. Three floral traits and
pollination syndrome were coded: primary flower color (purple, red, white, yellow), flower
shape (funnelform, salverform, tubular), corolla spur (absent, present), and syndrome (bee,
butterfly, hummingbird). We fit a multivariate mixed model with a non-informative prior, where
the shape and scale parameters equal 0.001, and residual variance was fixed. Analyses were
performed over the 50 randomly selected ortholog trees to account for phylogenetic uncertainty.
A phylogenetic model was used to account for any interspecific non-dependency in the dataset
by using a random effects structure that incorporated a phylogenetic tree model. Floral traits
were set as categorical response variables and the module eigengenes were set as the predictor
variable. For each tree, two MCMC chains were run for 250k generations and discarding the first
50k as burn-in. This process was run individually over each module in both the Bud and D
stages. We checked for convergence between model chains using the Gelman-Rubin statistic
(Gelman and Rubin 1992). Potential scale reduction values were all less than 1.1 and effective
sample sizes for all fixed effects were greater than 400. We considered fixed effects to be
statistically significant when the probabilities in the 95% credible region did not include zero
(Supplemental Figures 3-6).
278

Evolutionary rate analysis

Codon alignments of the filtered orthogroups (n=9503) were produced using PAL2NAL v14 (Suyama et al. 2006). Subsequent alignments and corresponding gene trees were used as input to the codeml package in PAML v4.9 (Yang 2007) to estimate d_N/d_S (omega, ω). d_N/d_S for each orthogroup was estimated using a one-ratio model (model = 0, NSsites = 0), providing a single estimate for each orthogroup to match other single orthogroup metrics, such as connectivity. If d_N/d_S values were exceptionally large (= 999) because of zero synonymous differences, those values were removed from the dataset.

First, we performed linear regression (LM) to test the effect of orthogroup connectivity, average expression levels, and their interactions (explanatory variables) on the estimated d_N/d_S values (main effect), LM = d_N/d_S ~ connectivity * expression. This was performed using the lm function in R with 10000 bootstrap pseudoreplicates. Second, we ran two-sample t-tests with 10000 permutations to test whether hub nodes and periphery nodes in each network had lower (for hubs) or higher (for periphery) d_N/d_S values than all background nodes. Third, we fit a LM with 10000 bootstrap pseudoreplicates to test whether modules associated with a pollination syndrome (bee, butterfly, or hummingbird; see Results below) had increased d_N/d_S values, LM = d_N/d_S ~ syndrome. For all LM and permutation analyses, d_N/d_S, connectivity, and expression values were log transformed to reach a normal distribution before being processed. Analyses were performed separately for each network using the separately estimated connectivity and expression levels.

Functional annotation and GO enrichment analysis

Filtered orthogroups (n=9503) were annotated based on a search against all green plant proteins in the SwissProt database (release 2018_07; The UniProt Consortium 2017) using Diamond BLASTp (--evalue 1e-6). Results were used to assign annotations and gene ontology (GO) terms. GO enrichment analyses were performed using the R package TopGO with the default "weight01" algorithm with the recommended cutoff of p < 0.05 (Alexa et al. 2006). Enriched GO term redundancy was removed and summarized using the semantic similarity measures implemented in the REVIGO web server (Supek et al. 2011).

Data and code availability

Raw sequencing reads are available from the NCBI Sequence Read Archive under BioProjects: PRJNA435759 and PRJNA340450. Data and code for all analyses can found from https://drive.google.com/open?id=1pvzw1t1pObY_kAEYmxQ484PCTuEWrsrb.

Results

Transcriptome assembly

Twelve transcriptomes were sequenced (72 libraries) and assembled in 12 species after sampling across two floral developmental stages (Supplemental Table 1). The transcriptomes had

an average assembly length of 244.9 Mb (± 33.9 Mb), an average of 217,510 (± 47k) transcripts,

and an average N50 length of 1988 bp (± 306 bp). Between 22,942 and 33,501 putative genes

were detected with 2.34 (± 0.35) putative isoforms. BUSCO identified an average of 87% of the

complete, conserved single-copy plant orthologs (Supplemental Table 1). Our focus in our

comparative study was on the presence of conserved orthologs, and not on the presence of

taxonomically restricted genes, which are not likely preserved across species.

Constructing the co-expression network for 12 species

We identified 83595 orthologous clusters (orthogroups), each containing at least 2

species. After removing 74092 orthogroups with too many missing values (> 50% missing) or

low expression variance (< 0.3 variance) and keeping common orthogroups shared between the

Bud and D stages, 9503 orthogroups were used to construct two co-expression networks using

the weighted gene correlation network analysis (WGCNA) approach. Modules of co-expressed

orthogroups are inferred using the expression profiles of each sample regardless of species.

Clustering identified 65 and 62 co-expression modules in the Bud and D stage networks,

respectively (Figure 3; Supplemental Table 2). Module size in the Bud stage ranged from 22

(module ME63) to 592 (ME1) orthogroups (mean = 146) and ranged from 21 (ME61) to 704

(ME1) orthogroups (mean = 153) in the D stage. Out of the 9503 orthogroups, 7845 (83%) and

8735 (92%) were assigned to modules in each network.

The remaining unassigned genes from each network that were not placed into a well-

defined module were assigned to the ME0 module by WGCNA. Despite being unplaced, these

orthogroups may still be functionally relevant within each network. Unassigned genes in the

ME0 module were enriched for many processes, including transcription (Bud, n = 173; D, n =

84) and organ development (D, n = 55), among other functions (Supplemental Tables 3-4).

We tested whether modules in the Bud stage were preserved in the D stage, and vice

versa. Overall, preservation of modules between the Bud and D stage networks was high. Eight

of the 65 Bud stage modules (ME46, ME62, ME44, ME57, ME42, ME41, ME55, ME40) were

not strongly preserved in the D stage, while only 1 of the 62 D stage modules (ME59) was not

strongly preserved in the Bud stage (Supplemental Table 5). These patterns suggest that these

non-preserved modules may play unique roles within their network. Among the 8 Bud stage-

specific modules were many orthogroups related to primary cell growth, meristem identity and

differentiation, stamen development, and flowering (Supplemental Table 3). In the single D

stage-specific module were many orthogroups related to methylation, flowering, and carotenoid

biosynthesis (Supplemental Table 4). Despite the relatively high preservation, correspondence

between module membership in the Bud and D stages was moderate (Figure 4). Orthogroup

module membership tended to shift between each network. For example, most members of

module ME15 in the Bud stage are transferred to modules ME3 and ME7 in the D stage (Figure

4).

We defined network hubs (the most highly connected nodes within a module) with

module membership (kME) scores > 0.9. Using this definition, all modules in each network

contained at least one hub, with the percentage of hubs ranging from 1% to 24% of orthogroups
in a module (Figures 5-6; Supplemental Table 6). Of the 632 and 691 hubs identified in the Bud
and D stages, a significant portion (110) were shared between networks (Fisher's Exact Test,
odds ratio = 3.02, P < 0.001). Functional enrichment of hubs in the Bud stage revealed that many
were involved in the regulation of development and metabolic processes (n = 40), transport and
protein localization (n = 39), biosynthetic processes (n = 19), and DNA methylation (n = 7),
consistent with the notion that hubs usually play important roles in integrating other genes during
network evolution (Ravasz et al. 2002) (Table 2). In the D stage, hubs were enriched for
functions related to translation (n = 33), transport (n = 9), ethylene and abscisic acid signaling
pathways (n = 17), carotenoid and anthocyanin biosynthetic processes (n = 11), and the
regulation of gene expression (n = 20), among others (Table 2).
Nodes in the periphery of each network were loosely defined by considering them to be
the lowest 10% connected (connectivity$_{Bud}$ < 2.824, connectivity$_D$ < 4.651). These definitions
identified 951 periphery nodes (Supplemental Table 7), of which 301 were peripheral in both
networks (Fisher's Exact Test, odds ratio = 5.67, P < 0.001). These periphery nodes in the Bud
stage were enriched for functions related to reproductive structure development (n = 75),
transport and localization (n = 107), and signaling (n = 389), among others (Table 3). In the D
stage, the periphery was enriched for functions related to the regulation of flower development (n
= 64), the regulation of flowering time (n = 18), the regulation of flavonoid biosynthesis (n = 7),
and cell fate specification (n = 12), among others (Table 3). Greater than expected numbers of
peripheral nodes in the D stage were related to regulation (Fisher's exact test, odds ratio = 1.191,
p = 0.02752) and transcription (Fisher's exact test, odds ratio = 1.370, p = 0.001549). In contrast,
there were no differences in the expected number of peripheral nodes related to regulation
(Fisher's exact test, odds ratio = 0.955, p = 0.7044) or transcription (Fisher's exact test, odds
ratio = 1.047, p = 0.3549) in the Bud stage.
384
Modules associated with floral traits
We calculated module eigengenes, single values which represent the gene expression
profiles of a sample within a module (Supplemental Figures 1-2). The extent of module
involvement in various biological processes can be tested by correlating eigengenes with external
traits, such as flower color. We tested the relationship between the module eigengenes and the
four floral traits (pollination syndrome [bee, butterfly, hummingbird], flower color [purple, red,
yellow, white], flower shape [funnelform, salverform, tubular], and corolla spurs [absence,
presence]), while controlling for any phylogenetic bias in the dataset using a phylogenetic mixed
model (Figures 5-6; Supplemental Figures 3-6). We also examined whether eigengene values (>
0.1 or < -0.1) in single species contributed to a strong correlation or whether eigengene values (>
0.1 or < -0.1) in multiple species contributed to a strong correlation (Figures 5-6; Supplemental
Figures 3-6). We considered there to be strong evidence for trait association when a significant
correlation from MCMCglmm was coupled with elevated eigengene (> 0.1) or decreased
eigengene (< -1.0) expression in multiple species that shared any of the significant traits.

Likewise, weak evidence for trait association was considered when the MCMCglmm results
were significant, but only a single species or no species sharing the significant traits had elevated
(> 0.1) or decreased (< -1.0) eigengene expression. WGCNA does not use trait information when
inferring modules, so modules that correlated strongly with any floral trait were inferred without
a priori knowledge.

Pollination syndrome—We included 3 species with the bee pollination syndrome (*A.*
candida, A. misera, and *E. verticillata*), 4 species with the butterfly pollination syndrome (*A.*
cettoana, A. grandiflora, A. longiflora, and *A. patens* 'Major'), and 5 species with the
hummingbird pollination syndrome (*A. admirabilis, A. antirrhina, A. erecta, A. pedunculata,* and
G. cuneifolia) (Figure 2; Table 1). These pollination syndromes are traditionally defined in
Achimenes and other gesneriads based on primary flower color, flower shape, and the presence
of corolla spurs (Wiehler 1983; Ramírez Roa 1987; Roalson et al. 2003). Expression of 23 (35%)
and 21 (34%) modules were positively associated with the pollination syndromes in the Bud and
D stage networks, consisting of 2967 and 2880 orthogroups, respectively. (Figures 5-6). There
were 3 (4.6%), 7 (10.8%), and 13 (20%) modules associated with bee, butterfly, and
hummingbird pollination syndrome in the Bud stage network, while 1 (1.5%), 8 (12.9%), and 12
(19.4%) modules were associated with the same syndromes in the D stage. Modules were never
associated with more than one syndrome. Butterfly and hummingbird syndromes were never
correlated to the same modules and were often correlated in opposite directions (i.e., butterfly
was positive correlated and hummingbird was negative correlated).

Among the 3 modules that were correlated to the bee pollination syndrome in the Bud
stage were many orthogroups involved in hormone signaling pathways (particularly ethylene and
abscisic acid, n = 14), cell wall and lipid biosynthesis (n = 7), photomorphogenesis (n = 2),
nitrogen compound metabolism (n = 4), and the transport of potassium and monocarboxylic
acids (n = 5) (Table 4). In contrast, the 1 module correlated to the bee pollination syndrome in
the D stage was enriched for different functions from the Bud stage, including the positive
regulation of molecular functions and response to biotic stimuli (n = 7), phospholipid catalysis (n
= 2), xyloglucan biosynthesis (n = 2), mRNA stability and catalysis (n = 5), and chromatin
modification (n = 4) (Table 5).

The 7 modules correlated to the butterfly pollination syndrome in the Bud stage
contained orthogroups involved in RNA splicing (n = 16), fertilization (n = 3), DNA methylation
(n = 2), cell growth and shoot formation (n = 8), the development of organ boundaries (n = 3),
and stress responses (n = 83) (Table 4). Within the 8 modules correlated with butterfly
pollination in the D stage were orthogroups enriched for functions in photorespiration (n = 7),
microtubule organization (n = 5), chromatin silencing (n = 3), red and far-red light responses (n
= 3), modified amino acid biosynthesis and metabolism (n = 10), and mRNA transport (n = 9)
(Table 5).

Thirteen modules were correlated to the hummingbird pollination syndrome in the Bud
stage and the orthogroups were enriched for numerous functions, including many developmental
processes (Table 4). These modules were enriched for floral whorl development (n = 12), RNA

and mRNA modifications (n = 19), cell growth and division (n = 54), organelle assembly (n = 3),
transportation of carbohydrates and potassium (n = 22), and responses to external stimuli (n =
17) (Table 4). Many of the same enrichments were seen in the 12 modules correlated to the
hummingbird pollination syndrome in the D stage. These modules had enrichments for floral
whorl development (n = 14), cell polarity and proliferation (n = 8), adaxial/abaxial specification
(n = 8), organ formation (n = 3), the regulation of growth (n = 26), and endosperm development
(n = 6). Additionally, these modules had enrichment for RNA and mRNA modification (n = 79),
transcription (n = 3), protein-DNA complex formation (n = 7), signaling (n = 25), and UV
response (n = 9) (Table 5).
Flower color—Primary flower color is closely associated to pollination syndrome in
Achimenes (Ramírez Roa 1987; Roalson et al. 2003), and we identified 26 (40%) and 17 (27%)
modules associated to any flower color in the Bud and D stage networks, consisting of 2935 and
1500 orthogroups respectively (Figures 5-6). Red and yellow are associated with hummingbird
pollination, purple with butterfly pollination, and white with bee pollination. In the Bud stage,
there were 14 (21.5%), 11 (16.9%), and 1 (1.5%) module associated with red, purple, or yellow
flower color. In the D stage, no modules were associated with yellow, but 8 (12.9%) and 9
(14.5%) modules were associated with red or purple flower color. In both networks, no modules
were corelated to white flower color. Similar to syndrome, purple and red flower color were
never associated with the same modules and were always correlated in opposite directions.
The 14 modules associated with red flowers in the Bud stage overlapped significantly
with the modules associated with the hummingbird pollination syndrome in the Bud stage and
contained the same functional enrichments (Table 4; Supplemental Table 8). However, in the 8 D
stage modules associated with red flowers, the orthogroups were enriched for functions involved
in flower morphogenesis (n = 8), cell wall modification (n = 3), positive regulation of gene
expression (n = 14), and histone modification (n = 7), among others (Supplemental Table 8).
Within the 11 modules associated with purple flowers in the Bud stage were orthogroups
enriched for transcription (n = 14), meristem initiation (n = 4), DNA methylation (n = 3),
fertilization (n = 5), ethylene signaling (n = 5), response to biotic stimuli (n = 55), and others
(Supplemental Table 8). In contrast, the 9 modules associated with purple flowers in the D stage
were enriched for orthogroups involved in anther and endosperm development (n = 9),
translation (n = 4), the regulation of developmental processes (n = 9), lipid modification (n = 8),
cell division (n = 3), and hormone signaling (gibberellic acid and jasmonic acid, n = 6), among
many other processes (Supplemental Table 8).
A single module (ME36) was associated with yellow flower color, only in the Bud stage,
and contained few orthogroups involved in floral meristem determinacy (n = 1), floral organ
identity (n = 1), gene silencing (n = 1), sulfur compound biosynthesis (n = 2), and vegetative
phase change (n = 1) (Supplemental Table 8).
Flower shape—Flower shape is also important but not as closely associated to pollination
syndromes as primary flower color in *Achimenes* (Ramírez Roa 1987; Roalson et al. 2003).
Flowers with bee pollination tend to have funnelform flowers, butterfly pollinated flowers all

have salverform flowers, and hummingbird pollinated flowers have either salverform or tubular
flowers (Table 2). Overall, there were 14 (22%) and 13 (21%) modules associated with any of
the flower shapes in the Bud and D stages, consisting of 2940 and 2504 orthogroups,
respectively (Figures 5-6). Here, there were 1 (1.5%), 8 (12.3%), and 5 (7.7%) modules whose
expression correlated to funnelform, salverform, or tubular flowers in the Bud stage (Figure 5).
Fewer modules were correlated in the D stage network, but expression in 1 (1.6%), 7 (12.3%),
and 5 (8.1%) modules were associated with funnelform, salverform, and tubular shapes (Figure
4). No modules were positively associated with more than one flower shape.
Only two modules were associated with funnelform flowers, one in the Bud stage and the
other in the D stage, and these were enriched for functions related to ethylene signaling (n = 3),
lipid biosynthesis (n = 3), chromatin modification (n = 4), cell wall organization (n = 2), and
many metabolic processes (Supplemental Table 8). Within the 8 modules associated with
salverform flowers in the Bud stage, there were many orthogroups involved in mRNA processing
and modification (n = 50), ovule development (n = 7), chlorophyll biosynthesis (n = 4), and the
regulation of flower development (n = 20) (Supplemental Table 8). In the D stage, the 7 modules
associated with salverform flowers contained orthogroups involved in chromatin silencing (n =
7), anthocyanin biosynthesis (n = 10), cell cycle and cell division (n = 95), stamen development
(n = 12), auxin biosynthesis (n = 7), and transport (n = 17) (Supplemental Table 8).
Among the 5 modules associated with tubular flowers in the Bud stage were orthogroups
enriched for functions in the positive regulation of transcription and organelle assembly (n = 13),
RNA processing (n = 5), xylem development (n = 4), metabolism (n = 12), and many
biosynthetic processes (Supplemental Table 8). In the 5 modules associated with tubular flowers
in the D stage were enrichments for floral whorl development (n = 6), positive regulation of gene
expression (n = 15), mRNA modification (n = 5), and signaling (n = 12) (Supplemental Table 8).
Corolla spurs—The presence of corolla spurs is found only on butterfly pollinated
species, two of which were included: *A. grandiflora* and *A. patens* (Figure 1; Table 1). There was
expression in 5 (7.7%) modules associated with spurs in the Bud stage network, while only 1
(1.6%) module was associated in the D stage network (Figures 3-4). Within the 5 modules
associated with corolla spurs in the Bud stage were orthogroups related to miRNA processing (n
= 4), mRNA splicing (n = 9), the regulation of signal transduction pathways (n = 16), and pollen
tube growth (n = 6) (Supplemental Table 8). In the single module associated with corolla spurs in
the D stage (ME53) were very few orthogroups involved in microtubule organization (n = 2),
stamen development (n = 1), the positive regulation of cell division (n = 1), flavonoid
biosynthesis (n = 1), and indole acetic acid metabolism (n = 1) (Supplemental Table 8).
Non-associated—There were 13 and 9 modules not associated with any traits in the Bud
and D stage networks, respectively (Figures 5-6). Three of these modules in the Bud stage were
not-preserved in the D stage (modules ME46, ME44, and ME41; Figures 5-6; Supplemental
Table 5). The 9 non-associated modules in the D stage were also not associated with any traits in
the Bud stage (Figures 5-6) and the orthogroups had overlapping enrichments for many essential
processes (Supplemental Table 8). These modules were enriched for orthogroups involved in the

development of the inflorescence, stamens, and floral organs, biosynthesis of vitamins and
nucleotides, photosynthesis and photosynthetic electron transport, metabolism of phosphate
compounds, and cell surface receptor signaling (Supplemental Table 8). Numerous orthogroups
were also involved in the positive regulation of reproduction, the negative regulation of growth,
and the regulation of heterochronic development, histone methylation, and gibberellic acid
mediated signaling (Supplemental Table 8).
525
Evolutionary rates correlate to network location
We ran several analyses to explore how several network variables may affect
evolutionary rates of the orthogroups within each network (as measured by d_N/d_S). First, we tested
the effects of orthogroup connectivity, expression levels, and their interaction on the d_N/d_S values
(Supplemental Table 7). Results indicated a collective effect between connectivity and
expression on d_N/d_S in both the Bud stage (LM, $F(3,9548) = 92.58$, p = 0, $\eta^2 = 0.03$) and the D
stage networks (LM, $F(3,9548) = 78.36$, p = 0, $\eta^2 = 0.02$). We found that orthogroup connectivity
was negatively correlated with d_N/d_S in the Bud stage network (LM, t = -16.31, p = 0) and the D
stage network (LM, t = -15.13, p = 0) (Figure 7), while expression was positively correlated with
d_N/d_S in the Bud stage (LM, t = 2.12, p = 0.0344) and not in the D stage (LM, t = 1.51, p =
0.1324) (Supplemental Figure 7). The interaction between connectivity and expression also
indicated a marginal dependency of both predictors on d_N/d_S in the Bud stage (LM, t = -2.022, p =
0.0433) but not in the D stage network (LM, t = -1.823, p = 0.0683). These patterns have been
observed in model species and may be universal to all eukaryotes.
Second, we tested whether the nodes we defined as hubs in each network had lower d_N/d_S
values, suggestive of increased evolutionary constraint (Supplemental Table 9). Hubs are
considered to functionally significant and may control large portions of the network. We found
that the hubs in the Bud stage network had lower d_N/d_S than all background genes (Two-sample t-
test, $t(679.31) = 7.0934$, p = 1.65e-12, Cohen's d = 0.383; Permutation t-test. mean diff. = 0.266,
p = 0.0001) (Supplemental Figure 8). Similarly, hubs in the D stage network also had lower d_N/d_S
than all background genes (Two-sample t-test, $t(755.91) = 4.1918$, p = 1.55e-05, Cohen's d =
0.206; Permutation t-test, mean diff. = 0.143, p = 0.0001) (Supplemental Figure 8).
Third, we tested whether the most peripheral nodes (the lowest 10% connected nodes)
had higher d_N/d_S values, which may suggest relaxed evolutionary constraint (Supplemental Table
9). These nodes are loosely connected within the network and their roles may be more apt to
fluctuate during evolution and development. These peripheral genes had higher d_N/d_S than all
background genes in the Bud stage network (Two-sample t-test, $t(1487.70) = -10.272$, p = 0,
Cohen's d = -0.253; Permutation t-test, mean diff. = -0.176, p = 0.0001) and the D stage network
(Two-sample, $t(1512.30) = -10.387$, p = 0, Cohen's d = -0.252; Permutation t-test, mean diff. = -
0.175, p = 0.0001) (Supplemental Figure 8).
Lastly, we tested whether modules correlated with bee, butterfly, or hummingbird
pollination syndromes showed increased evolutionary rates over the non-associated modules.
Results indicated a very weak effect of syndrome association on d_N/d_S in the Bud network (LM,

$F(3,9548) = 5.186$, p $= 0.001406$, $\eta^2 = 0.0016$) and the D network (LM, $F(3,9548) = 4.277$, p $=$
0.005038, $\eta^2 = 0.0013$) (Supplemental Table 9). Only bee syndrome associated modules had
marginally lower d_N/d_s values in both networks (Bud stage, LM, t $= -3.58$, p $= 0.000346$; D stage,
LM, t $= -2.508$, p $= 0.0122$). Butterfly associated modules had marginally lower d_N/d_s only in the
D stage (LM, t $= -2.548$, p $= 0.0108$), while hummingbird associated modules did not have any
effect in either network.
565

Discussion

Here we report on patterns of gene co-expression across two stages of flower
development in 12 species of *Achimenes, Eucodonia,* and *Gesneria*. Our comparative and
phylogenetic context allowed us to reveal gene co-expression patterns across a group of closely
related plant species that correlated to diverse pollination syndromes and floral traits, including
flower color, flower shape, and corolla spurs. Our results suggest floral forms that are associated
with bee, butterfly, and hummingbird pollinator groups are correlated with many co-expression
modules. We found that nearly a third of modules in each network (23 and 21 modules in the
Bud and D stages, respectively) had evidence for association with the three pollination
syndromes (Figures 5-6). In our analysis, large fractions of our 9503 orthogroups set could be
partitioned into modules (Figure 3), network hubs and peripheral nodes identified (Supplemental
Tables 6-7), and several orthogroups were identified for their potential importance in floral
phenotypic differentiation. Lastly, orthogroup position in each network (measured by
connectivity) had an association with evolutionary rate (d_N/d_s) (Figure 7). These analyses provide
a first overview of floral transcriptional architecture across members of the same genus that
display diverse floral forms.
582

Hub nodes contain important candidates for floral form

To date, a large number of studies conducted in diverse angiosperm systems, have found
that many genes are associated with flower development (Quattrocchio et al. 1999; Howarth et
al. 2011). Despite evidence for extensive gene/genome duplications that could expand the
genetic repertoire (Tang et al. 2008), many recent studies have suggested that the generation of
novel genes is rare (Patterson et al. 2009; De Smet et al. 2013). Thus, the evolution and
development of diverse floral forms may predominantly occur through the co-option of existing
pathways (Preston et al. 2011). To our knowledge, gene co-expression network analyses have not
been previously used across multiple species in a non-model lineage to understand flower
development and evolution.
Hub nodes (genes) are considered to be conserved, highly connected nodes that are
central to network architecture and may directly orchestrate numerous biological processes
(Ravasz et al. 2002; Hu et al. 2016; Mähler et al. 2017). Our analyses identified over 600 gene
nodes as hubs within each network, the majority of which were only hubs within a specific floral
stage. These hub nodes tended to have lower d_N/d_s estimates, suggesting that they not prone to
rapid evolutionary change (Supplemental Table 9). Unsurprisingly, hub nodes in the Bud stage

were enriched for many processes related to early flower development, including cell growth,
leaf formation, and photosystem assembly (Table 2). Similarly, hub nodes in the D stage were
enriched for genes that may be related the development of distinct floral forms, including
carotenoid biosynthesis, anthocyanin biosynthesis, and cell morphogenesis (Table 2). Within this
context of flower development, we identified a number of hubs that we consider important
candidates for involvement in the production of flower color and the development of floral form.
Many of these candidates were considered previously (Roberts and Roalson 2017) as potentially
important for the diversification of floral form in *Achimenes*. Undoubtedly, this list of hubs
contains many genes that may important for other biological traits that we do not consider here,
such as scent, nectar, and pollen production.
Anthocyanin biosynthesis—Anthocyanins are common floral pigments that produce red,
purple, and blue colors and are produced via the highly conserved flavonoid biosynthetic
pathway (Holton and Cornish 1995). With a few exceptions, all anthocyanins are produced from
three precursor molecules: pelargonidin, cyanidin, and delphinidin (Grotewold 2006). There are
six enzymatic reactions required for the production of anthocyanins and many also serve as
branching points for the production of flavonoids, lignins, flavonols, and other phenolics that
have roles in stress response and physiology (Winkel-Shirley 2002; Tahara 2007). We identified
four homologs of anthocyanin biosynthetic pathway enzymes as network hubs: flavonoid 3'-
hydroxylase (*F3'H*; cluster8703_1rr.inclade1.ortho3; module ME12), dihydroflavonol 4-
reductase (*DFR*; cluster5082_1rr.inclade1.ortho6; module ME48), flavonoid 3',5'-
methyltransferase (*FAOMT*; cluster11783_1rr.inclade1.ortho1.tre; module ME32), and flavonol
3-O-glucosyltransferase (*UFOG*; cluster8241_2rr.inclade1.ortho2.tre; module ME1).
These 4 enzymes are involved in distinct steps during anthocyanin production and
chemical modification in floral tissue. Activity of *F3'H* directs the pathway toward the
production of cyanidins and quercetin (Grotewold 2006). Cyanidin can accumulate in floral
tissue as a purple-hued pigment, while quercetin plays a large role in various stress-related
physiological functions (Pollastri and Tattini 2011), including response to high light and UVB
radiation (Gerhardt et al. 2008) and antioxidant properties (Rice-Evans et al. 1996). Therefore,
F3'H as an important hub in the networks makes sense given its enormous pleiotropic role in
various physiological responses. Just prior to the production of anthocyanins, *DFR* catalyzes the
reaction to produce pelargonidins, cyanidins, and delphinidins from the precursor molecules
dihydrokaempferol, dihydroquercetin, and dihyrdomyricetin, respectively (Grotewold 2006).
Shifts in *DFR* expression, function, and substrate specificity have been implicated in many
evolutionary transitions in flower color (Smith et al. 2013; Wu et al. 2013).
Further modifications to anthocyanins, such as methylation, glycosylation, and acylation,
produce a wide variety of compounds and hues (Wessinger and Rausher 2012). We identified
homologs to both a methyltransferase (*FAOMT*) and glucosyltransferase (*UFOG*) as hubs in the
networks. Methylation of anthocyanins have been widely studied in petunia (Provenzano et al.
2014), grapes (Hugueney et al. 2009) and tree peony (Du et al. 2015), demonstrating that
methylation can impact the hue, either shifting it more toward purple (Sakata et al. 1995) or

sometimes red (Tanaka et al. 2008). Glycosylation stabilizes anthocyanins molecules for
subsequent transfer to acidic vacuoles (Saito and Yamazaki 2002), increasing their solubility,
and influencing their color variability by inter- and intra-molecular stacking (Springob et al.
2003). Finding homologs of *FAOMT* and *UFOG* as network hubs reflect the importance of
anthocyanin modification for different flower color hues among *Achimenes* species.
644   *Carotenoid biosynthesis*—Carotenoids are red, orange, and yellow pigments that are
produced in plastids and are essential for photosynthesis, but also accumulate as secondary
metabolites in flowers to attract pollinators (Kevan and Baker 1983). Floral and fruit carotenoids
can also be used to produce volatile apocarotenoids, which may further enhance plant-animal
interactions during pollination and seed dispersal (Dudareva et al. 2006). They may also serve as
precursors to the production of the hormone abscisic acid and strigalactones that are involved in
many plant developmental processes (Rosas-Saavedra et al. 2016). We identified 4 homologs of
carotenoid biosynthetic pathway enzymes as network hubs: geranylgeranyl pyrophosphate
synthase (*GGPPS*; cluster10945_2rr.inclade1.ortho1.tre; module ME35), phytoene
dehydrogenase (*PDS*; cluster13048_1rr.inclade1.ortho1.tre; module ME4), lycopene beta cyclase
(*LCYB*; cluster13774_1rr.inclade1.ortho1.tre; module ME7), and capsanthin-capsorubin synthase
(*CCS*; cluster13583_1rr.inclade1.ortho2.tre; module ME6).
656   Each of these 4 enzymes catalyzes reactions during different steps of carotenoid
production. *GGPPS* produces the main precursor molecular geranylgeranyl pyrophosphate
(GGPP) to carotenoid biosynthesis that is produced as the end-product of the plastid
methylerythritol 4-phosphate (MEP) pathway. Two GGPP molecules are then condensed into
phytoene and undergoes four successive dehydrogenations and isomerizations to produce
lycopene. *PDS* catalyzes the first two desaturation steps that transform the colorless phytoene
into the red-colored lycopene. *PDS* is a rate-limiting enzyme in carotenoid biosynthesis
(Chamovitz et al. 1993) and down-regulation has been shown to lead to large accumulations of
phytoene (Busch et al. 2002) and decreases in total carotenoid, chlorophyll, and photosynthetic
efficiency (Wang et al. 2009).
666   Lycopene is the main starting compound for a large variety of carotenoids. *LCYB* takes
lycopene as a substrate and subsequently produces beta-carotene. Many plants use tissue-specific
isoforms of *LCYB* genes that are expressed in fruits or flowers that often correlate with the
accumulation of beta-carotene or downstream xanthophylls (Ronen et al. 2000; Ahrazem et al.
2010; Devitt et al. 2010). Beta-carotenes are hypothesized to be the primary carotenoids in
Achimenes flowers (Roberts and Roalson 2017). Another functionally related carotenoid cyclase
enzyme is *CCS* (or neoxanthin synthase, *NSY*) which catalyzes the final step in the carotenoid
pathway and converts the yellow-colored violaxanthin into neoxanthin (Parry and Horgan 1991).
Both violaxanthin and neoxanthin can be further used for producing the plant hormone abscisic
acid (Neuman et al. 2014). Both enzymes are likely functionally important in the developing
flowers for the biosynthesis of beta-carotene and xanthophylls.
677   *Flower development*—Very few studies have so far investigated the molecular
mechanisms involved in the development of flower shape in the Gesneriaceae (Gao et al. 2008;

Zhou et al. 2008; Alexandre et al. 2015). However, extensive work has been done in model
organisms, such as *Arabidopsis thaliana* and *Antirrhinum majus*, which can provide insight into
the potential roles that many transcription factors might play in determining floral organ identity
and shape in *Achimenes*. Among the network hubs, we identified several homologs that are likely
involved in floral organ development, including mixta (*MIXTA*;
cluster8699_1rr.inclade1.ortho3.tre; modules ME5 and ME8), agamous (*AG*;
cluster7311_1rr.inclade1.ortho3.tre; module ME23), globosa (*GLO*;
cluster2750_1rr.inclade1.ortho3.tre; module ME37), and tcp4 (*TCP4*;
cluster12380_1rr.inclade1.ortho2.tre; module ME22).
Petal identity and petal surface morphology are both thought to be important components
for pollinator attraction (Noda et al. 1994; Whitney et al. 2011). *MIXTA* was first characterized
from *Antirrhinum* and controls the development of the conical cell shape in the petal epidermis
(Noda et al. 1994). Consequently, *MIXTA* and *MIXTA*-like genes have been characterized in
other model systems, such as *Arabidopsis* and *Mimulus* for their similar role in epidermal cell
morphogenesis (Baumann et al. 2007) as well as trichome development (Gilding and Marks
2010; Scoville et al. 2011). The homolog of *MIXTA* in our study was a hub node in both
networks, suggesting it might play an important role throughout flower development in
determining petal surface morphology.
Early floral organ determination requires the involvement of several MADS-box
transcription factors, such as *AG* and *GLO*. We identified homologs of *AG* as a hub in the Bud
stage and *GLO* as a hub in the D stage (Supplemental Table 7). *AG* has been shown in
Arabidopsis to be expressed in early flower development in order to terminate meristem activity
and promote the development of stamens and carpels, while working in combination with
APETALA3, PISTILLATA, and *SEPALLATA* (Gómez-Mena et al. 2005). More recently homologs
of *AG* were demonstrated to be involved in nectary development in both *Arabidopsis* and
Petunia (Morel et al. 2018), another floral trait important for plant-pollinator interaction. *GLO*
was first characterized in *Antirrhinum* and *Arabidopsis* and is also involved with stamen identity,
as well as petal identity (Tröbner et al. 1992; Goto and Meyerowitz 1994). Outside of these
model systems, its expression patterns and role in the evolution of complex floral forms have
been examined in numerous plant lineages, from monocots (Bartlett and Specht 2010) to
Solanaceae (Zhang et al. 2014). *AG* and *GLO* will be interesting candidates to begin exploring
further how petals, stamens, carpels, and nectary development in *Achimenes*.
Lastly, the homolog of *TCP4* was identified as a hub during the Bud stage, and may be
important for its potential role in petal development and cell proliferation. Looking into the
genetic factors regulating the elaboration of petal growth will be important for understanding
how corolla spurs develop in *Achimenes* (Figure 1). The role of *TCP4* been characterized in
Arabidopsis for its role in both petal growth, showing restricted expression to the developing
petal tissue (Nag et al. 2009). More recent work using gene regulatory networks in *Arabidopsis*
has shown that petal size is controlled by a *SEPALLATA3*-regulated miR319/*TCP4* module
(Chen et al. 2018). Outside of *Arabidopsis, TCP4* was also implicated for a having a role in the

719	development of the petal spur in *Aquilegia* (Yant et al. 2015). Therefore, considering both the
720	role for *TCP4* in petal and spur development demonstrated in *Arabidopsis* and *Aquilegia* and its
721	identification as a hub gene in our networks, we think this gene requires further interrogation.
722	
723	**Peripheral genes are enriched for transcriptional regulators in later stages of development**
724	Peripheral nodes have low connectivity within co-expression networks (Supplemental
725	Table 7), tend to have higher rates of molecular evolution (Masalia et al. 2017), and have been
726	hypothesized to contribute more to evolutionary innovations than network hubs (Ichihashi et al.
727	2014). Of the 951 nodes we defined as peripheral in both Bud and D stage networks, roughly a
728	third were consistently peripheral under our definitions (lowest 10% connected; Supplemental
729	Table 7). After identifying the nodes with homologies to transcriptional regulators, we found
730	there was no difference in the expected number in the early Bud stage, while there were greater
731	than expected numbers of transcriptional regulators in the periphery of the D stage (see Results).
732	No such patterns existed when we tested these transcriptional regulators as hub genes. While
733	Mähler et al. (2017) found an enrichment for transcription factors among hub genes in a single
734	species network of *Populus tremula*, our results in a multi-species network find the opposite
735	pattern. Shifting the expression patterns of various genes and pathways can lead to phenotypic
736	divergence among closely related species (West-Eberhard 1989). This can be seen during later
737	stages of flower development when distinct traits, such as red or purple flower color, can appear
738	in different species due to shifting expression patterns within the anthocyanin biosynthetic
739	pathway (Smith and Rausher 2011). The connection between the apparent overabundance of
740	transcriptional regulators in later stages of flower development and the rapid phenotypic
741	divergence in floral form will need to be explored further.
742	Several notable peripheral nodes were identified for their potential involvement in
743	important floral processes (Supplemental Table 7). Three homologs were identified in the
744	periphery of the Bud stage with potential involvement in the determination of zygmorphic
745	(bilateral) floral symmetry: two homologs of divaricata (*DIV*;
746	cluster13684_1rr.inclade1.ortho1.tre, cluster8585_1rr.inclade1.ortho1.tre; ME26 and ME0) and
747	one homolog of dichotoma (*DICH*; cluster13245_1rr.inclade1.ortho1.tre; ME0). Bilateral
748	symmetry has been proposed as a mechanism that facilitates pollen transfer by insects (Galliot et
749	al. 2006). Both *DIV* and *DICH* are TCP transcription factors that were first characterized in
750	*Antirrhinum* (Almeida et al. 1997; Luo et al. 1996), and show localized expression to the ventral
751	and dorsal portions of the petal, respectively. *Achimenes* flowers at the Bud stage are lacking or
752	only just beginning to establish floral asymmetry and the relegation of both *DIV* and *DICH* to the
753	network periphery may reflect this small role. The role of these genes has not been studied in the
754	Gesneriaceae, but have been examined more recently in other non-model systems for their role in
755	the evolution of pollination syndromes (Zhang et al. 2010; Preston et al. 2011). Additionally, a
756	homolog of the homeobox wuschel gene (*WUS*; cluster9273_1rr.inclade1.ortho1.tre; module
757	ME0) was identified from the periphery of the D stage network. *WUS* was first characterized in
758	*Arabidopsis* for its role in meristem cell maintenance (Laux et al. 1996) and its expression is

759 localized only to the meristem (Mayer et al. 1998). Once the floral meristem has been initiated,
760 *WUS* starts to be repressed by *AG* (hub gene, described above) (Sun et al. 2009). The relegation
761 of *WUS* to the periphery in our networks at the D stage might then be expected given that it plays
762 such a prominent role during early floral development and the vegetative to reproductive
763 transition.
764
Evolutionary rates are correlated to network connectivity
Increased protein evolutionary rates during rapid diversification has been suggested in
other radiations of animals and plants (Kapralov et al. 2013; Brawand et al. 2014; Pease et al.
2016). Our study is among the first to examine patterns of network evolution across multiple
closely-related species of plants with multiple transitions in flower type and pollination
syndrome. Hummingbirds are hypothesized to have had an important influence on diversification
in Neotropical Gesneriaceae (Roalson and Roberts 2016; Serrano-Serrano et al. 2017b), while
the role of butterflies has not been explored. Modules correlated with hummingbird or butterfly
pollination syndromes did not exhibit clear increases or decreases in evolutionary rates compared
to other modules (Supplemental Table 9). Detecting selection is subject to many factors, such as
a gene's transcriptional abundance and its importance within the protein interaction network
(Lemos et al. 2005). For instance, we found that higher network connectivity was strongly
associated with lower evolutionary rates across both networks in our study (Figure 7;
Supplemental Table 9). This corroborates the recent results found in other eukaryotic systems
(Morandin et al. 2016; Masalia et al. 2017; Josephs et al. 2017; Mähler et al. 2017) while also
showing the apparent strength of correlation between connectivity and evolutionary rate differs
between studies and systems.
Network effects (such as the type of network analysis employed) and the expression
levels of highly abundant genes are known to affect evolutionary rates, potentially confounding
analysis of selection pressure (Krylov et al. 2003). We found that d_N/d_S in early development may
be dependent on both connectivity and expression levels, while this interactive effect was gone
during later development (Supplemental Table 9). As there are few highly connected genes
(hubs, which are important determinants of the observed network structure), a random mutation
would be less likely to affect such a gene. Changing the amino acid sequence of a hub gene
could have multiple associated effects, some deleterious (Hahn and Kern 2004; Luisi et al.
2015). In comparison, a random mutation would be more likely to affect a gene with lower
connectivity, while also resulting in variants with fewer negative consequences. The hub genes
have more direct interactions and are more likely to be essential and involved in more biological
processes, placing them under greater constraint.
794
Numerous network modules correlated to floral form
One primary use for gene co-expression network analyses is to identify modules of co-
expressed genes whose overall expression (as measured by the eigengene) may correlate with
different traits. These traits can be both qualitative or quantitative, such as social caste (Morandin

et al. 2016), seed oil production (Hu et al. 2016), or anther development (Hollender et al. 2014).
While we tested only qualitative traits here, future studies could also correlate eigengene
expression to quantitative variables such as nectary size or sucrose concentration, since these
appear to be another important component of pollination syndrome delimitation (Perret et al.
2001; Katzer et al. 2019; Vandelook et al. 2019). Since our dataset was composed of samples
from 12 species, we performed correlation analyses using a phylogenetic model to test whether
module expression correlated to different flower colors, flower shapes, corolla spurs, and
pollination syndromes. Nearly half of our modules in both networks had evidence for an
association with these floral traits (Figures 5-6). There were no modules whose correlation to a
pollination syndrome had increased eigengene expression in all species sharing that trait (Figures
5-6). Instead many modules with correlation to a trait showed a phylogenetic pattern with a
shared increased eigengene expression among the most closely related species. For example,
module ME23 in the Bud stage network had a correlation to butterfly pollination and increased
eigengene expression in *A. cettoana* and *A. longiflora*, both members of Clade 1 (Figure 2). Our
aim was to identify sets of co-expressed genes that would provide candidates for involvement in
the development of these traits.
For instance, many of the candidate hubs we identified (see above) were found in
modules that were correlated to one or multiple traits to which they might contribute, such as
PHAN (module ME4, correlated to corolla spurs) and *DFR* (module ME48, correlated to
hummingbird pollination and purple flowers) (Supplemental Table 2). Additionally, modules
correlated to various floral traits were enriched for multiple genes that might be involved in the
development of those traits (Supplemental Table 8). Module ME53 was correlated to the corolla
spurs found in *A. grandiflora* and *A. patens* in the D stage network and was enriched for many
genes involved in cell division and cell growth (Figure 6; Supplemental Table 4). Other times,
modules were correlated to a specific trait but the enriched GO terms did not provide a clear idea
of how these co-expressed orthogroups may contribute (Supplemental Tables 3-4). In our
analyses, we considered relatively few traits and only focused on those that have traditionally
been considered the most important for the differentiation of floral form in *Achimenes*, namely
flower color, flower shape, and corolla spurs. Undoubtedly there are a myriad number of
molecular and biochemical pathways that we have not considered here that underlie the
development of these complex floral phenotypes. With the idea that networks represent complex
systems of gene and protein interactions, it may be that these seemingly simple and generalized
floral phenotypes are produced through much more complex means than many have previously
considered.
833
Strengths and limitations of network analyses in non-model organisms
Our analysis covers genes with orthologs found in most species, representing a set of
evolutionarily conserved genes. While no reference genome currently exists for Neotropical
Gesneriaceae, we constructed, annotated, and presented de novo transcriptomes for 10
Achimenes species as well as 2 outgroup species, *E. verticillata* and *G. cuneifolia*. Our approach

enables functional genomic studies in this non-model system, but the potential for misassemblies
may bias our downstream results (Hornett and Wheat 2012). Using a gene tree-based approach to
carefully identify orthogroups, we focused our attention on conserved gene regulatory machinery
and not on taxonomically restricted genes. The latter may be involved in species-specific
functions. Nevertheless, it is worth noting that many recent studies have indicated that transitions
in floral form may be due to the shifting expression of genes found in conserved genetic
pathways (Wessinger and Rausher 2014). Taxonomically restricted genes are likely interacting
with existing regulatory pathways, and the manner in which they integrate into the co-expression
modules identified here will be a fascinating topic for future research. With extended sampling
within *Achimenes* we could infer species-specific co-expression networks and perform more
detailed cross-species comparisons. We expect that these taxonomically restricted genes may be
poorly connected with the network, which may allow for increased diversification.

We characterized gene expression patterns by sequencing RNA across two timepoints
during flower development. Previous gene expression experiments across three timepoints (early
Bud, intermediate D, and late Pre-anthesis stages) in four *Achimenes* species suggested that the
majority of differential gene expression occurred between the early Bud stage and the
intermediate D stage (Roberts and Roalson 2017). We believe our approach allows us to capture
dynamic transcriptional patterns during flower development for comparison across multiple
species. We found that the co-expression networks for the Bud and D stages were largely
preserved, despite there being a few stage specific modules (Supplemental Table 5). However,
this approach may greatly over-simplify the nature of transcriptional regulation in these species.
Organ- and structure-specific expression studies may have greater power to detect floral form-
specific differences, both in terms of the number of differentially expressed genes, and co-
expression network structure (Hollender et al. 2014; Suzuki et al. 2017; Shahan et al. 2018). As a
result, our data likely represents an underestimate of the true number of modules. Future studies
could focus on the comparative analysis of expression patterns in specific organs, such as the
petals, stigma, and stamens, and additional timepoints, which would provide greater functional
insight into the evolution, development, and diversification of these organs.

Network analysis represents a more complex approach than differential expression
analyses because it can capture system-level properties (Langfelder and Horvath 2008). Most
phenotypes involve interactions of proteins from diverse biochemical pathways. Although
inferred modules do not necessarily correspond to entire biochemical pathways, or other
components of cellular organization, the approach performs well in reconstructing the
complexity of protein-protein interaction networks (Allen et al. 2012). In many systems this
approach has succeeded in identifying candidate genes for various biological traits, for example
floral organ development in strawberry (Hollender et al. 2014; Shahan et al. 2018) or carotenoid
accumulation in carrots (Iorizzo et al. 2016). Our results demonstrate the difficulty in identifying
gene clusters that underlie highly correlated floral traits, especially in non-model systems, as
seen by the overlapping correlation of modules associated with flower shape and color (Figures
5-6). It has been shown in many plant systems that floral traits tend to be selected together by

pollinators (Cuartas-Domínguez and Medel 2010; Sletvold et al. 2010), making it challenging to
separate correlated modules from causal modules. Co-expression networks are foundationally
about correlations between genes (van Dam et al. 2018), making them useful in identifying novel
sets of genes that may be functionally relevant for trait of interest. Other approaches, such as
quantitative trait locus (QTL) analyses may provide more promise in identifying distinct genes
that are involved in the divergence of floral traits. QTL studies have already shown promise in
identifying genetic loci that underlie pollination syndrome divergence in *Mimulus* (Yuan et al.
2013), *Penstemon* (Wessinger et al. 2014), and *Rhytidophyllum* (Alexandre et al. 2015).
We were able to identify numerous candidate orthogroups involved in the development of
floral form in our non-model system. While no functional genomic data exists yet for *Achimenes*,
our annotations are based on the manually curated SwissProt database. De novo assemblies
provide important data for non-model organisms, but there can be biases introduced during
assembly and annotation due to potentially missing genes (Hornett and Wheat 2012). We
attempted to limit this bias by focusing only on shared orthogroups found in the majority (at least
6) of our sampled *Achimenes*. Applying this network-based approach to the diversification of
flowers across multiple species in *Achimenes*, we recovered distinct modules of co-expressed
genes that may underlie a range of floral phenotypic traits tied to different pollination
syndromes. Although our data come from whole flowers during development, and do not allow
us to test specific hypotheses regarding whether particular genes have shifted expression location
in the developing flower, they do suggest that there are conserved regulatory modules that may
have been co-opted in convergent flower types multiple times. It will be interesting to apply
network analyses to study floral diversification in additional lineages and examine how
conserved or divergent the patterns are across angiosperms.
902
Conclusions
Floral forms corresponding to bee, butterfly, and hummingbird pollination have evolved
multiple times across the Neotropical plant genus *Achimenes*. Genome-wide gene expression
estimates were taken from flowers in 12 species across two development stages in order to
construct, analyze, and compare stage-specific gene co-expression networks. We hypothesized
that numerous modules in each network would correlate to these pollination syndromes and that
central genes in the network may be candidates for involvement in the development of important
floral traits, such as flower color and shape. We found that nearly a third of modules were
correlated to the pollination syndromes and many more were correlated to different flower colors
and shapes. The outgroup species, *E. verticillata* (bee pollinated) and *G. cuneifolia*
(hummingbird pollinated), displayed correlation patterns similar to the ingroup species that
shared floral traits, suggesting that some co-expression patterns might be shared across
evolutionary distances. Several of the hub genes in the networks were homologs of the
anthocyanin and carotenoid biosynthetic pathways, important for the production of floral
pigments that attract different pollinators. A negative relationship between network connectivity
and d_N/d_S corroborates the findings in model systems that more centrally located nodes (likely

hubs) are under increased evolutionary constraint. We found that the less connected genes (peripheral) are under more relaxed constraints and contained numerous transcriptional regulators. Our results demonstrate the utility of applying co-expression network analyses in non-model plant lineages to begin identifying important modules and pathways that will be useful starting points for additional analyses into the evolution and development of floral form.

Acknowledgements

We thank Joanna Kelley, Amit Dhingra, and Andrew McCubbin for helpful comments that improved this work. Sequencing was performed at the Genomics Core Lab at Washington State University, Spokane, and the Genomics Sequencing and Analysis Facility at the University of Texas, Austin.

References

Ahrazem O, Rubio-Moraga A, López RC, Gómez-Gómez L. 2010. The expression of a chromoplast-specific lycopene beta cyclase gene is involved in high production of saffron's apocarotenoid precursors. J Exp Bot 61:105-119.

Alexa A, Rahnenführer J, Lengauer T. 2006. Improved scoring of functional groups from gene expression data by decorrelating GO graph structure. Bioinformatics 22:1600-1607.

Alexandre H, Vrignaud J, Mangin B, Joly S. 2015. Genetic architecture of pollination syndrome transition between hummingbird-specialist and generalist species in the genus Rhytidophyllum (Gesneriaceae). PeerJ 3:e1028.

Allen JD, Xie Y, Chen M, Girard L, Xiao G. 2012. Computing statistical methods for constructing large scale gene networks. PLOS One 7:e29348.

Almeida J, Rocheta M, Galego L. 1997. Genetic control of flower shape in Antirrhinum majus. Development 124:1387-1392.

Amorim FW, Galetto L, Sazima M. 2013. Beyond the pollination syndrome: nectar ecology and the role of diurnal and nocturnal pollinators in the reproductive success of Inga sessilis. Plant Biol 15:317-327.

Baumann K, Perez-Rodriguez M, Bradley D, Venail J, Bailey P, Jin H, Koes R, Roberts K, Martin C. 2007. Control of cell and petal morphogenesis by R2R3 MYB transcription factors. Development 134:1691-1701.

Barrett SC. 2013. The evolution of plant reproductive systems: how often are transitions irreversible? P Roy Soc B-Biol Sci 280: 20130913.

Bartlett ME, Specht CD. 2010. Evidence for the involvement of GLOBOSA-like gene duplications and expression divergence in the evolution of floral morphology in the Zingiberales. New Phytol 187:521-541.

Bolger AM, Lohse M, Usadel B. 2014. Trimmomatic: a flexible trimmer for Illumina sequence data. Bioinformatics 30:2114-2120.

Bolstad B. 2018. preprocessCore: A collection of pre-processing functions. R package version 1.46.0, https://github.com/bmbolstad/preprocessCore.

Bradshaw HD, Otto KG, Frewen BE, McKay JK, Schemske DW. 1998. Quantitative trait loci
affecting differences in floral morphology between two species of monkeyflower
(Mimulus). Genetics 149:367-382.
Brawand D, Soumillon M, Necsulea A, Julien P, Csárdi G, Harrigan P, Weier M, Liechti A,
Aximu-Petri A, Kircher M, Albert FW. 2011. The evolution of gene expression levels in
mammalian organs. Nature 478:343.
Brawand D, Wagner CE, Li YI, Malinsky M, Keller I, Fan S, Simakov O, Ng AY, Lim ZW,
Bezault E, Turner-Maier J. 2014. The genomic substrate for adaptive radiation in African
cichlid fish. Nature 513:375.
Bray NL, Pimentel H, Melsted P, Pachter L. 2016. Near-optimal probabilistic RNA-seq
quantification. Nat Biotechnol 34:525.
Buchfink B, Xie C, Huson DH. 2014. Fast and sensitive protein alignment using DIAMOND.
Nat Methods 12:59-60.
Busch M, Seuter A, Hain R. 2002. Functional analysis of the early steps of carotenoid
biosynthesis in tobacco. Plant Physiol 128:439-453.
Chamovitz D, Sandmann G, Hirschberg J. 1993. Molecular and biochemical characterization of
herbicide-resistant mutants of cyanobacteria reveals that phytoene desaturation is a rate-
limiting step in carotenoid biosynthesis. J Biol Chem 268:17348-17353.
Chen D, Yan W, Fu LY, Kaufmann K. 2018. Architecture of gene regulatory networks
controlling flower development in Arabidopsis thaliana. Nat Commun 9:1-13.
Cortés-Flores J, Hernández-Esquivel KB, González-Rodríguez A, Ibarra-Manríquez G. 2017.
Flowering phenology, growth forms, and pollination syndromes in tropical and dry forest
species: influence of phylogeny and abiotic factors. Am J Bot 104:39-49.
Cronk Q, Ojeda I. 2008. Bird-pollinated flowers in an evolutionary and molecular context. J Exp
Bot 59:715-727.
Cuartas-Domínguez M, Mendel R. 2010. Pollinator-mediated selection and experimental
manipulation of the flower phenotype in Chloraea bletioides. Funct Ecol 24:1219-1227.
Des Marais DL, Rausher MD. 2010. Parallel evolution at multiple levels in the origin of
hummingbird pollination flowers in Ipomoea. Evolution 64:2044-2054.
De Smet R, Adams KL, Vandepoele K, Van Montagu MC, Maere S, Van de Peer Y. 2013.
Convergent gene loss following gene and genome duplications creates single-copy
families in flowering plants. P Natl Acad Sci USA 110:2898-2903.
Devitt LC, Fanning K, Dietzgen RG, Holton TA. 2010. Isolation and functional characterization
of a lycopene β-cyclase gene that controls fruit colour of papaya (Carica papaya L.). J
Exp Bot 61:33-39.
Du H, Wu J, Ji KX, Zeng QY, Bhuiya MW, Su S, Shu QY, Ren HX, Liu ZA, Wang LS. 2015.
Methylation mediated by an anthocyanin O-methyltransferase is involved in purple
flower coloration in Paeonia. J Exp Bot 66:6563-6577.
Dudareva N, Negre F, Nagegowda DA, Orlova I. 2006. Plant volatiles: recent advances and
future perspectives. Cr Rev Plant Sci 25:417-440.

Enright AJ, Van Dongen S, Ouzounis CA. 2002. An efficient algorithm for large-scale detection
of protein families. Nucleic Acids Res 30:1575-1584.
Faegri K, van der Pijl L. 1979. *Principles of Pollination Ecology*. Oxford: Pergamon Press.
Fenster CB, Armbruster WS, Wilson P, Dudash MR, Thomson JD. 2004. Pollination syndromes
and floral specialization. Annu Rev Ecol Evol S 35:375-403.
Galliot C, Stuurman J, Kuhlemeier C. 2006. The genetic dissection of floral pollination
syndromes. Curr Opin Plant Biol 9:78-82.
Gao Q, Tao JH, Yan D, Wang YZ, Li ZY. 2008. Expression differentiation of CYC-like floral
symmetry genes correlated with their protein sequence divergence in Chirita heterotricha
(Gesneriaceae). Dev Genes Evol 218:341-351.
Gelman A, Rubin DB. 1992. Inference from iterative simulation using multiple sequences. Stat
Sci 7:457-472.
Gerhardt KK, Lampi MA, Greenberg BM. 2008. The effect of far-red light on plant growth and
flavonoid accumulation in Brassica napus in the presence of ultraviolet B radiation.
Photochem Photobiol 84:1445-1454.
Gervasi DD, Schiestl FP. 2017. Real-time divergent evolution in plants driven by pollinators.
Nat Commun 8: 14691.
Gilding EK, Marks MD. 2010. Analysis of purified glabra3-shapeshifter trichomes reveals a role
for NOECK in regulating early trichome morphogenic events. Plant J 64:304-317.
Gómez-Mena C, De Folter S, Costa MMR, Angenent GC, Sablowski R. 2005. Transcriptional
program controlled by the floral homeotic gene AGAMOUS during early organogenesis.
Development 132:429-438.
Goto K, Meyerowitz EM. 1994. Function and regulation of the Arabidopsis floral homeotic gene
PISTILLATA. Gene Dev 8:1548-1560.
Grabherr MG, Haas BJ, Yassour M, Levin JZ, Thompson DA, Amit I, Adiconis X, Fan L,
Raychowdhury R, Zeng Q, Chen Z. 2011. Trinity: reconstructing a full-length
transcriptome without a genome from RNA-Seq data. Nat Biotechnol 29:644.
Grotewold E. 2006. The genetics and biochemistry of floral pigments. Annu Rev Plant Biol
57:761-780.
Guillerme T, Healy K. 2014. mulTree: a package for running MCMCglmm analysis on multiple
trees. ZENODO. 10.5281/zenodo.12902.
Hahn MW, Kern AD. 2004. Comparative genomics of centrality and essentiality in three
eukaryotic protein-interaction networks. Mol Biol Evol 22:803-806.
Haas BJ, Papanicolaou A, Yassour M, Grabherr M, Blood PD, Bowden J, Couger MB, Eccles D,
Li B, Lieber M, MacManes MD. 2013. De novo transcript sequence reconstruction from
RNA-seq using the Trinity platform for reference generation and analysis. Nat Protoc
8:1494.
Hadfield JD. 2010. MCMC methods for multi-response generalized linear mixed models: the
MCMCglmm R pckage. J Stat Softw 33:1-22.

Hermann K, Klahre U, Venail J, Brandenburg A, Kuhlemeier C. 2015. The genetics of
reproductive organ morphology in two Petunia species with contrasting pollination
syndromes. Planta 241:1241-1254.

Hoballah ME, Gübitz T, Stuurman J, Broger L, Barone M, Mandel T, Dell'Olivo A, Arnold M,
Kuhlemeier C. 2007. Single gene-mediated shift in pollinator attraction in Petunia. Plant
Cell 19: 779-790.

Hollender CA, Kang C, Darwish O, Geretz A, Matthews BF, Slovin J, Alkharouf N, Liu Z. 2014.
Floral transcriptomes in woodland strawberry uncover developing receptacle and anther
gene networks. Plant Physiol 165:1062-1075.

Hornett EA, Wheat CW. 2012. Quantitative RNA-Seq analysis in non-model species: assessing
transcriptome assemblies as a scaffold and the utility of evolutionary divergent genomic
reference species. BMC Genomics 13:361.

Horvath S, Dong J. 2008. Geometric interpretation of gene co-expression network analysis.
PLOS Comput Biol 4:e1000117.

Howarth DG, Martins T, Chimney E, Donoghue MJ. 2011. Diversification of CYCLOIDEA
expression in the evolution of bilateral flower symmetry in Caprifoliaceae and *Lonicera*
(Dipsacales). Ann Bot-London 107:1521-1532.

Hu G, Hovav R, Grover CE, Faigenboim-Doron A, Kadmon N, Page JT, Udall JA, Wendel JF.
2016. Evolutionary conservation and divergence of gene coexpression networks in
Gossypium (cotton) seeds. Genome Biol Evol 8:3765-3783.

Hugueney P, Provenzano S, Verriès C, Ferrandino A, Meudec E, Batelli G, Merdinoglu D,
Cheynier V, Schubert A, Ageorges A. 2009. A novel cation-dependent O-
methyltransferase involved in anthocyanin methylation in grapevine. Plant Physiol
150:2057-2070.

Ichihashi Y, Aguilar-Martínez JA, Farhi M, Chitwood DH, Kumar R, Millon LV, Peng J, Maloof
JN, Sinha NR. 2014. Evolutionary developmental transcriptomics reveals a gene network
module regulating interspecific diversity in plant leaf shape. P Natl Acad Sci USA
111:E2616-E2621.

Iorizzo M, Ellison S, Senalik D, Zeng P, Satapoomin P, Huang J, Bowman M, Iovene M,
Sanseverino W, Cavagnaro P, Yildiz M. 2016. A high-quality carrot genome assembly
provides new insights into carotenoid accumulation and asterid genome evolution. Nat
Genet 48:657.

Josephs EB, Wright SI, Stinchcombe JR, Schoen DJ. 2017. The relationship between selection,
network connectivity, and regulatory variation within a population of *Capsella*
grandiflora. Genome Biol Evol 9:1099-1109.

Katoh K, Standley DM. 2013. MAFFT multiple sequence alignment software version 7:
improvements in performance and usability. Mol Biol Evol 30:772-780.

Katzer AM, Wessinger CA, Hileman LC. 2019. Nectary size is a pollination syndrome trait in
Penstemon. New Phytol 223:377-384.

Kapralov MV, Votintseva AA, Filatov DA. 2013. Molecular adaptation during a rapid adaptive
radiation. Mol Biol Evol 30:1051-1059.
Kevan PG, Baker HG. 1983. Insects as flower visitors and pollinators. Annu Rev Entomol
28:407-453.
Krylov DM, Wolf YI, Rogozin IB, Koonin EV. 2003. Gene loss, protein sequence divergence,
gene dispensability, expression level, and interactivity are correlated in eukaryotic
evolution. Genome Res 13:2229-2235.
Langfelder P, Horvath S. 2008. WGCNA: an R package for weighted correlation network
analysis. BMC Bioinformatics 9:559.
Langfelder P, Luo R, Oldham MC, Horvath S. 2011. Is my network module preserved and
reproducible? PLOS Comput Biol 7:e1001057.
Laux T, Mayer KF, Berger J, Jurgens G. 1996. The WUSCHEL gene is required for shoot and
floral meristem integrity in Arabidopsis. Development 122:87-96.
Lemos B, Bettencourt BR, Meiklejohn CD, Hartl DL. 2005. Evolution of proteins and gene
expression levels are coupled in *Drosophila* and are independently associated with
mRNA abundance, protein length, and number of protein-protein interactions. Mol Biol
Evol 22:1345-1354.
Li W, Godzik A. 2006. Cd-hit: a fast program for clustering and comparing large sets of protein
or nucleotide sequences. Bioinformatics 22:1658-1659.
Lohmann JU, Hong RL, Hobe M, Busch MA, Parcy F, Simon R, Weigel D. 2001. A molecular
link between stem cell regulation and floral patterning in Arabidopsis. Cell 105:793-803.
Love MI, Huber W, Anders S. 2014. Moderated estimation of fold change and dispersion for
RNA-seq data with DESeq2. Genome Biol 15:550.
Luisi P, Alvarez-Ponce D, Pybus M, Fares MA, Bertranpetit J, Laayouni H. 2015. Recent
positive selection has acted on genes encoding proteins with more interactions within the
whole human interactome. Genome Biol Evol 7:1141-1154.
Luo D, Carpenter R, Copsey L, Vincent C, Clark J, Coen E. 1996. Control of organ asymmetry
in flowers of Antirrhinum. Cell 99:367-376.
Ma X, Zhao H, Xu W, You Q, Yan H, Gao Z, Su Z. 2018. Co-expression gene network analysis
and functional module identification in bamboo growth and development. Front Genetics
9:574.
Mähler N, Wang J, Terebieniec BK, Ingvarsson PK, Street NR, Hvidsten TR. 2017. Gene co-
expression network connectivity is an important determinant of selective constraint.
PLOS Genet 13:e1006402.
Martén-Rodríguez S, Quesada M, Castrao AA, Lopezaraiza-Mikel M, Fenster CB. 2015. A
comparison of reproductive strategies between island and mainland Caribbean
Gesneriaceae. J Ecol 103:1190-1204.
Masalia RR, Bewick AJ, Burke JM. 2017. Connectivity in gene coexpression networks
negatively correlates with rates of molecular evolution in flowering plants. PLOS One
12:e182289.

Mason MJ, Fan G, Plath K, Zhou Q, Horvath S. 2009. Signed weighted gene co-expression
network analysis of transcriptional regulation in murine embryonic stem cells. BMC
Genomics 10:327.
Mayer KF, Schoof H, Haecker A, Lenhard M, Jürgens G, Laux T. 1998. Role of WUSCHEL in
regulating stem cell fate in the Arabidopsis shoot meristem. Cell 95:805-815.
Morandin C, Tin MM, Abril S, Gómez C, Pontieri L, Schiøtt M, Sundström L, Tsuji K, Pedersen
JS, Helanterä H, Mikheyev AS. 2016. Comparative transcriptomics reveals the conserved
building blocks involved in parallel evolution of diverse phenotypic traits in ants.
Genome Biol 17:43.
Morel P, Heijmans K, Ament K, Chopy M, Trehin C, Chambrier P, Bento SR, Bimbo A,
Vandenbussche M. 2018. The floral C-lineage genes trigger nectary development in
Petunia and Arabidopsis. Plant Cell 30:2020-2037.
Nag A, King S, Jack T. 2009. miR319a targeting of TCP4 is critical for petal growth and
development in Arabidopsis. P Natl Acad Sci USA 106:22534-22539.
Neuman H, Galpaz N, Cunningham Jr, FX, Zamir D, Hirschberg J. 2014. The tomato mutation
nxd1 reveals a gene necessary for neoxanthin biosynthesis and demonstrates that
violaxanthin is a sufficient precursor for abscisic acid biosynthesis. Plant J 78:80-93.
Noda KI, Glover BJ, Linstead P, Martin C. 1994. Flower colour intensity depends on specialized
cell shape controlled by a Myb-related transcription factor. Nature 369:661-664.
O'Meara BC, Smith SD, Armbruster WS, Harder LD, Hardy CR, Hileman LC, Hufford L, Litt
A, Magallón S, Smith SA, Stevens PF. 2016. Non-equilibrium dynamics and floral trait
interactions shape extant angiosperm diversity. P Roy Soc B-Biol Sci 283:20152304.
Papiorek S, Junker RR, Alves-dos-Santos I, Melo GA, Amaral-Neto LP, Sazima M, Wolowski
M, Freitas L, Lunau K. 2016. Bees, birds and yellow flowers: pollinator-dependent
convergent evolution of UV patterns. Plant Biol 18:46-55.
Paradis E, Claude J, Strimmer K. 2004. APE: analyses of phylogenetics and evolution in R
language. Bioinformatics 20:289-290.
Parry AD, Horgan R. 1991. Carotenoids and abscisic acid (ABA) biosynthesis in higher plants.
Physiol Plantarum 82:320-326.
Parsana P, Ruberman C, Jaffe AE, Schatz MC, Battle A, Leek JT. 2019. Addressing confounding
artifacts in reconstruction of gene co-expression networks. Genome Biol 20:94.
Patterson AH, Bowers JE, Ruggmann R, Dubchak I, Grimwood J, Gundlach H, Haberer G,
Hellsten U, Mitros T, Poliakov A, Schmutz J. 2009. The *Sorghum bicolor* genome and
the diversification of grasses. Nature 457:551.
Pease JB, Haak DC, Hahn MW, Moyle LC. 2016. Phylogenomics reveals three sources of
adaptive variation during a rapid radiation. PLOS Biol 14:e1002379.
Perret M, Chautems A, Spichiger R, Peixoto M, Savolainen V. 2001. Nectar sugar composition
in relation to pollination syndromes in Sinningieae (Gesneriaceae). Ann Bot-London
87:267-273.

Piechowski D, Dötterl S, Gottsberger G. 2010. Pollination biology and floral scent chemistry of
the Neotropical chieropterophilous Parkia pendula. Plant Biol 12:172-182.
Pickrell JK, Marioni JC, Pai AA, Degner JF, Engelhardt BE, Nkadori E, Veyrieras JB, Stephens
M, Gilad Y, Pritchard JK. 2010. Understanding mechanisms underlying human gene
expression variation with RNA sequencing. Nature 464:768.
Pollastri S, Tattini M. 2011. Flavonols: old compounds for old roles. Ann Bot-London 108:1225-
1233.
Prasad A, Kumar SS, Dessimoz C, Bleuler S, Laule O, Hruz T, Gruissem W, Zimmerman P.
2013. Global regulatory architecture of human, mouse and rat tissue transcriptomes.
BMC Genomics 14:716.
Preston JC, Hileman LC, Cubas P. 2011. Reduce, reuse, and recycle: developmental evolution of
trait diversification. Am J Bot 98:397-403.
Preston JC, Martinez CC. Hileman LC. 2011. Gradual disintegration of the floral symmetry gene
network is implicated in the evolution of a wind-pollination syndrome. P Natl Acad Sci
USA 108:2343-2348.
Price MN, Dehal PS, Arkin AP. 2010. FastTree 2—approximately maximum-likelihood trees for
large alignments. PLOS One 5:e9490.
Provenzano S, Spelt C, Hosokawa S, Nakamura N, Brugliera F, Demelis L, Geerke DP, Schubert
A, Tanaka Y, Quattrocchio F, Koes R. 2014. Genetic control and evolution of
anthocyanin methylation. Plant Physiol 165:962-977.
Quattrocchio F, Wing J, van der Woude K, Souer E, de Vetten N, Mol J, Koes R. 1999.
Molecular analysis of the *anthocyanin2* gene of petunia and its role in the evolution of
flower color. Plant Cell 11:1433-1444.
Raherison ESM, Giguère I, Caron S, Lamara M, MacKay JJ. 2015. Modular organization of the
white spruce (*Picea glauca*) transcriptome reveals functional organization and
evolutionary signatures. New Phytol 207:172-187.
Ramírez-Aguirre E, Martén-Rodríguez S, Quesada-Avila G, Quesada M, Martínez-Díaz Y,
Oyama K, Espinosa-García FJ. 2019. Reproductive isolation among three sympatric
Achimenes species: pre- and post-pollination components. Am J Bot 106:1021-1031.
Ramírez Roa MA. 1987. Revision de *Achimenes* (Gesneriaceae). PhD dissertation, Universidad
Nacional Autonoma de Mexico, Mexico City, Mexico.
Ravasz E, Somera AL, Mongru DA, Oltvai ZN, Barbási AL. 2002. Hierarchical organization of
modularity in metabolic networks. Science 297:1551-1555.
Rice-Evans CA, Miller N, Papanga G. 1996. Structure-antioxidant relationships of flavonoids
and phenolic acids. Free Radical Bio Med 20:933-956.
Ritchie W, Rajasekhar M, Flamant S, Rasko JE. 2009. Conserved expression patterns predict
microRNA targets. PLOS Comput Biol 5:e1000513.
Roalson EH, Roberts WR. 2016. Distinct processes drive diversification in different clades of
Gesneriaceae. Syst Biol 65:662-684.

Roalson EH, Skog LE, Zimmer EA. 2003. Phylogenetic relationships and the diversification of floral form in *Achimenes* (Gesneriaceae). Syst Bot 28:593-608.

Roberts WR, Roalson EH, 2017. Comparative transcriptome analyses of flower development in four species of *Achimenes* (Gesneriaceae). BMC Genomics 18:240.

Roberts WR, Roalson EH. 2018. Phylogenomic analyses reveal extensive gene flow within the magic flowers (*Achimenes*). Am J Bot 105:726-740.

Romero IG, Ruvinsky I, Gilad Y. 2012. Comparative studies of gene expression and the evolution of gene regulation. Nat Rev Genet 13:505-516.

Ronen G, Carmel-Goren L, Zamir D, Hirschberg J. 2000. An alternative pathway to β-carotene formation in plant chromoplasts discovered by map-based cloning of Beta and old-gold color mutations in tomato. P Natl Acad Sci USA 97:11102-11107.

Rosas-Guerrero V, Aguilar R, Martén-Rodriguez S, Ashworth L, Lopezaraiza-Mikel M, Bastida JM, Quesada M. 2014. A quantitative review of pollination syndromes: do floral traits predict effective pollinators? Ecol Lett 17:388-400.

Rosas-Saavedra C, Stange C. 2016. Biosynthesis of carotenoids in plants: Enzymes and color. In: Stange C, editor. *Carotenoids in nature: biosynthesis, regulation and function*. Vol. 79. Springer. p. 35-69.

Saito K, Yamazaki M. 2002. Biochemistry and molecular biology of the late-stage of biosynthesis of anthocyanin: lessons from Perilla frutescens as a model plant. New Phytol 155:9-23.

Sakata Y, Aoki N, Tsunematsu S, Nishikouri H, Johjima T. 1995. Petal coloration and pigmentation of tree peony bred and selected in Daikon island (Shimane Prefecture). J Jpn Soc Hortic Sci 65:351-357.

Sauquet H, Magallón S. 2018. Key questions and challenges in angiosperm macroevolution. New Phytol 219:1170-1187.

Scoville AG, Barnett LL, Bodbyl-Roels S, Kelly JK, Hileman LC. 2011. Differential regulation of a MYB transcription factor is correlated with transgenerational epigenetic inheritance of trichome density in Mimulus guttatus. New Phytol 191:251-263.

Serrano-Serrano ML, Marcionetti A, Perret M, Salamin N. 2017a. Transcriptome resources for an endemic Neotropical plant lineage (Gesneriaceae). Appl Plant Sci 5:1600135.

Serrano-Serrano ML, Rolland J, Clark JL, Salamin N, Perret M. 2017b. Hummingbird pollination and the diversification of angiosperms: an old and successful association in Gesneriaceae. P Roy Soc B-Biol Sci 284:20162816.

Shahan R, Zawora C, Wight H, Sittmann J, Wang W, Mount SM, Liu Z. 2018. Consensus coexpression network analysis identifies key regulators of flower and fruit development in wild strawberry. Plant Physiol 178:202-216.

Simão FA, Waterhouse RM, Ioannidis P, Kriventseva EV, Zbodnov EM. 2015. BUSCO: assessing genome assembly and annotation completeness with single-copy orthologs. Bioinformatics 31:3210-3212.

Sletvold N, Grindeland JM, Ågren J. 2010. Pollinator-mediated selection on floral display, spur
length and flowering phenology in the deceptive orchid Dactylorhiza lapponica. New
Phytol 188:385-392.
Sletvold N, Trunschke J, Smit M, Verbeek J, Ågren J. 2016. Strong pollinator-mediated selection
for increased flower brightness and contrast in a deceptive orchid. Evolution 70:716-724.
Smith SA, Dunn CW. 2008. Phyutility: a phyloinformatics tool for trees, alignments and
molecular data. Bioinformatics 24:715-716.
Smith SD, Rausher MD. 2011. Gene loss and parallel evolution contribute to species difference
in flower color. Mol Biol Evol 28:2799-2810.
Smith SD, Wang S, Rausher MD. 2013. Functional evolution of an anthocyanin pathway enzyme
during a flower color transition. Mol Biol Evol 30:602-612.
Song L, Langfelder P, Horvath S. 2012. Comparison of co-expression measures: mutual
information, correlation, and model based indices. BMC Bioinformatics 13:328.
Springob K. Nakajima JI, Yamazaki M, Saito K. 2003. Recent advances in the biosynthesis and
accumulation of anthocyanins. Nat Prod Rep 20:288-303.
Stamatakis A. 2014. RAxML version 8: a tool for phylogenetic analysis and post-analysis of
large phylogenies. Bioinformatics 30:1312-1313.
Stebbins GL. 1970. Adaptive radiation of reproductive characteristics in angiosperms, I:
pollination mechanisms. Annu Rev Ecol Syst 1:307-326.
Sun B, Xu Y, Ng KH, Ito T. 2009. A timing mechanism for stem cell maintenance and
differentiation in the Arabidopsis floral meristem. Gene Dev 23:1791-1804.
Supek F, Bošnjak M, Škunca N, Šmuc T. 2011. REVIGO summarizes and visualizes long lists of
gene ontology terms. PLOS One 6:e21800.
Suyama M, Torrents D, Bork P. 2006. PAL2NAL: robust conversion of protein sequence
alignments into the corresponding codon alignments. Nucleic Acids Res 34:W609-W612.
Suzuki JY, Amore TD, Calla B, Palmer NA, Scully ED, Sattler SE, Sarath G, Lichty JS, Myers
RY, Keith LM, Matsumoto TK. 2017. Organ-specific transcriptome profiling of
metabolic and pigment biosynthetic pathways in the floral ornamental progenitor species
Anthurium amnicola Dressler. Sci Rep-UK 7:1596.
Tahara S. 2007. A journey of twenty-five years through the ecological biochemistry of
flavonoids. Biosci Biotechnol Biochem 71:1387-1404.
Tanaka Y, Sasaki N, Ohmiya A. 2008. Biosynthesis of plant pigments: anthocyanins, betalains
and carotenoids. Plant J 54:733-749.
Tang H, Wang X, Bowers JE, Ming R, Alam M, Paterson AH. 2008. Unraveling ancient
hexaploidy through multiply-aligned angiosperm gene maps. Genome Res 18:1944-1954.
Tröbner W, Ramirez L, Motte P, Hue I, Huijser P, Löning WE, Saedler H, Sommer H, Schwarz-
Sommer Z. 1992. GLOBOSA: a homeotic gene which interacts with DEFICIENS in the
control of Antirrhinum floral organogenesis. EMBO J 11:4693-4704.

Uebbing S, Künster A, Mäkinen H, Backström N, Bolivar P, Burri R, Dutoit L, Mugal CF, Nater
A, Aken B, Flicek P. 2016. Divergence in gene expression within and between two
closely related flycatcher species. Mol Ecol 25:2015-2028.

The UniProt Consortium. 2017. UniProt: the universal protein knowledgebase. Nucleic Acids
Res 45:D158-D169.

Van Dam S, Võsa U, van der Graaf A, Franke L, de Magalhaes JP. 2018. Gene co-expression
analysis for functional classification and gene-disease predictions. Brief Bioinform
19:575-592.

Vandelook F, Janssens SB, Gijbels P, Fischer E, Van den Ende W, Honnay O, Abrahamczyk S.
2019. Nectar traits differ between pollination syndromes in Balsaminaceae. Ann Bot-
London 124:269-279.

Van der Niet T, Johnson SD. 2012. Phylogenetic evidence for pollinator-driven diversification of
angiosperms. Trends Ecol Evol 27: 353-361.

Vincent CA, Enrico SC. 2004. A temporal and morphological framework for flower
development in Antirrhinum majus. Can J Botany 82:681-690.

Waites R, Selvadurai HR, Oliver IR, Hudson A. 1998. The PHANTASTICA gene encodes a
MYB transcription factor involved in growth and dorsoventrality of lateral organs in
Antirrhinum. Cell 93:779-789.

Wang M, Wang G, Ji J, Wang J. 2009. The effect of pds gene silencing on chloroplast pigment
composition, thylakoid membrane structure and photosynthesis efficiency in tobacco
plants. Plant Sci 177:222-226.

Wessinger CA, Rausher MD. 2014. Ecological transition predictably associated with gene
degeneration. Mol Biol Evol 32:347-354.

West-Eberhard MJ. 1989. Phenotypic plasticity and the origins of diversity. Ann Rev Ecol Syst
20:249-278.

Whitney HM, Glover BJ, Walker R, Ellis AG. 2011. The contribution of epidermal structure to
flower colour in the South African flora. Curtis's Botanical Magazine 28:349-371.

Wiehler H. 1983. A synopsis of the neotropical Gesneriaceae. Selbyana 6:1-219.

Winkel-Shirley B. 2002. Biosynthesis of flavonoids and effects of stress. Curr Opinion Plant
Biol 5:218-223.

Wu CA, Streisfeld MA, Nutter LI, Cross KA. 2013. The genetic basis of a rare flower color
polymorphism in Mimulus lewisii provides insight into the repeatability of evolution.
PLOS One 8:12.

Yang Y, Smith SA. 2014. Orthology inference in nonmodel organisms using transcriptomes and
low-coverage genomes: improving accuracy and matrix occupancy for phylogenomics.
Mol Biol Evol 31:3081-3092.

Yang Z. 2007. PAML 4: phylogenetic analysis by maximum likelihood. Mol Biol Evol 24:1586-
1591.

Yant L, Collani S, Puzey J, Levy C, Kramer EM. 2015. Molecular basis for three-dimensional
elaboration of the Aquilegia petal spur. P Roy Soc B-Biol Sci 282:20142778.

Yuan YW, Rebocho AB, Sagawa JM, Stanley LE, Bradshaw HD. 2016. Competition between
anthocyanin and flavonol biosynthesis produces spatial pattern variation of floral
pigments between Mimulus species. P Natl Acad Sci USA 113: 2448-2453.
Yuan YW, Sagawa JM, Young RC, Christensen BJ, Bradshaw HD. 2013. Genetic dissection of a
major anthocyanin QTL contributing to pollinator-mediated reproductive isolation
between sister species of Mimulus. Genetics 194:255-263.
Zhang JS, Li Z, Zhao J, Zhang S, Quan H, Zhao H, He C. 2014. Deciphering the Physalis
floridana double-layered-lantern1 mutant provides insights into the functional divergence
of the GLOBOSA duplicates within the Solanaceae. Plant Physiol 164:748-764.
Zhang W, Kramer EM, Davis CC. 2010. Floral symmetry genes and the origin and maintenance
of zygomorphy in a plant-pollinator mutualism. P Natl Acad Sci USA 107:6388-6393.
Zhou XR, Wang YZ, Smith JF, Chen R. 2008. Altered expression patterns of TCP and MYB
genes relating to the floral developmental transition from initial zygomorphy to
actinomorphy in Bournea (Gesneriaceae). New Phytol 178:532-543.
Zuo J, Niu Q-W, Frugis G, Chua N-H. 2002. The WUSCHEL gene promotes vegetative-to-
embryonic transition in Arabidopsis. Plant J 30:349-359.
1328

Tables

Table 1. Definition of pollination syndromes in *Achimenes*, *Eucodonia,* and *Gesneria*.

Table 2. Gene Ontology enrichment for the network hubs.

Table 3. Gene Ontology enrichment for the network periphery.

Table 4. Gene Ontology enrichment in Bud stage modules correlated to pollination syndrome.

Table 5. Gene Ontology enrichment in D stage modules correlated to pollination syndrome.

Figures

Figure 1. *Achimenes* flowers and the sampled developmental stages. Flowers of the twelve sampled species: (A) *A. admirabilis,* (B) *A. antirrhina,* (C) *A. candida*, (D) *A. cettoana*, (E) *A. erecta*, (F) *A. grandiflora*, (G) *A. longiflora*, (H) *A. misera*, (I) *A. patens*, (J) *A. pedunculate*, (K) *E. verticillate*, and (L) *G. cuneifolia*. Sampled Bud and D stage flowers for (M-N) *A. cettoana*, (O-P) *A. erecta*, (Q-R) *A. misera*, and (S-T) *A. patens*. Corolla spurs found in (U) *A. grandiflora* and (V) *A. patens* are indicated with arrows. Scale bars equal 1 cm. Abbreviations: A, *Achimenes*; E, *Eucodonia*; G, *Gesneria*.

Figure 2. Phylogenetic relationships and floral traits in *Achimenes*. Phylogeny of *Achimenes* adapted from Roberts and Roalson (2018). Clade 1 and Clade 2 (sensu Roalson et al. 2003) are indicated on the phylogeny with a circle and box, respectively. Flower traits for each species are shown at the tips of the phylogeny, including flower color, flower shape, presence of corolla spurs, pollination syndrome, and pollinator observations.

Figure 3. Co-expression clustering of the Bud and D stages. Hierarchical clustering by WGCNA showing co-expressed modules for (A) the Bud Stage and (B) the D stage. Each of the 9503 orthogroups were assigned to 65 and 62 modules, respectively, and the color rows underneath show the module assignment.

Figure 4. Correspondence of the Bud and D stage networks. Each row corresponds to one Bud stage module (labeled by module name and module size), and each column corresponds to one D stage module (labeled by module name and module size). Numbers in the table indicate gene counts in the intersection of the corresponding modules. Coloring of the table encodes -log(p), with p being the Fisher's exact test p-value for the overlap of the two modules. The stronger the red color, the more significant the overlap.

Figure 5. Summary of Bud stage network module association to floral traits and pollination syndrome. Each row represents an individual module in the Bud stage network and shows the total size (column All), the number of hubs (column Hubs), and whether the module is preserved in the D stage. Each column in the central table represents the floral traits (Pollination Syndrome,

Flower Color, Flower Shape, and Spurs) that were tested for an association to module eigengenes using MCMCglmm. The eigengene expression is indicated for each module in which all three replicates from an individual species had > 1.0 or < -1.0 expression. An association was considered strong when both MCMCglmm results were significant and the eigengene expression was > 1.0 or < -1.0 in multiple species that shared the significant trait. An association was considered weak when MCMCglmm results were significant but the eigengene was > 1.0 or < -1.0 in a single species or no species that shared the significant trait. Eigengene expression for species that share traits to those found significant by MCMCglmm are in bold.

Figure 6. Summary of D stage network module association to floral traits and pollination syndrome. Each row represents an individual module in the D stage network and shows the total size (column All), the number of hubs (column Hubs), and whether the module is preserved in the Bud stage. Each column in the central table represents the floral traits (Pollination Syndrome, Flower Color, Flower Shape, and Spurs) that were tested for an association to module eigengenes using MCMCglmm. The eigengene expression is indicated for each module in which all three replicates from an individual species had > 1.0 or < -1.0 expression. An association was considered strong when both MCMCglmm results were significant and the eigengene expression was > 1.0 or < -1.0 in multiple species that shared the significant trait. An association was considered weak when MCMCglmm results were significant but the eigengene was > 1.0 or < -1.0 in a single species or no species that shared the significant trait. Eigengene expression for species that share traits to those found significant by MCMCglmm are in bold.

Figure 7. Relationship between orthogroup dN/dS (omega) and orthogroup connectivity. (A) Bud stage network. (B) D stage network. Each point represents an individual orthogroup. The red line indicates the linear regression line for the relationship between dN/dS and connectivity. The dotted line indicates dN/dS = 1 with orthogroups above this line considered to be under relaxed or positive selection.

Table 1. Definition of pollination syndromes in *Achimenes, Eucodonia,* and *Gesneria*.

Pollination syndrome	Primary flower color	Flower shape	Presence of corolla spurs	Species
Melittophily (bees)	White or Purple with yellow or purple spotted throat	Funnelform	No	*Achimenes candida*
				Achimenes misera
				Eucodonia verticillata
Ornithophily (hummingbirds)	Red and yellow	Tubular, Salverform	No	*Achimenes admirabilis*
				Achimenes antirrhina
				Achimenes erecta
				Achimenes pedunculata
				Gesneria cuneifolia
Psychophily (butterflies)	Purple	Salverform	Yes	*Achimenes cettoana*
				Achimenes grandiflora
				Achimenes longiflora
				Achimenes patens

Table 2. Gene Ontology enrichment for the network hubs.

	GO ID	Term	Annotated	Significant	Expected	P
A. Bud stage Hubs	GO:0031325	positive regulation of cellular metabolic process	224	16	14.52	0.000
	GO:0006607	NLS-bearing protein import into nucleus	7	4	0.45	0.001
	GO:0015991	ATP hydrolysis coupled proton transport	9	4	0.58	0.002
	GO:0046470	phosphatidylcholine metabolic process	9	4	0.58	0.002
	GO:0018298	protein-chromophore linkage	32	7	2.07	0.004
	GO:0070507	regulation of microtubule cytoskeleton organization	6	3	0.39	0.005
	GO:1990778	protein localization to cell periphery	6	3	0.39	0.005
	GO:0051592	response to calcium ion	7	3	0.45	0.008
	GO:0006412	translation	233	24	15.10	0.009
	GO:0042594	response to starvation	60	10	3.89	0.012
	GO:0007035	vacuolar acidification	8	3	0.52	0.012
	GO:0006534	cysteine metabolic process	9	4	0.58	0.012
	GO:0009833	plant-type primary cell wall biogenesis	15	4	0.97	0.013
	GO:0048831	regulation of shoot system development	123	12	7.97	0.017
	GO:0032509	endosome transport via multivesicular body sorting pathway	9	3	0.58	0.017
	GO:0051568	histone H3-K4 methylation	9	3	0.58	0.017
	GO:0000271	polysaccharide biosynthetic process	123	11	7.97	0.023
	GO:0006732	coenzyme metabolic process	116	8	7.52	0.023
	GO:0010338	leaf formation	5	2	0.32	0.037
	GO:0048564	photosystem I assembly	5	2	0.32	0.037
	GO:0017001	antibiotic catabolic process	26	3	1.69	0.037
	GO:0044282	small molecule catabolic process	75	10	4.86	0.038
	GO:0006833	water transport	13	3	0.84	0.048
	GO ID	Term	Annotated	Significant	Expected	P
B. D stage Hubs	GO:0006412	translation	233	30	16.03	0.001
	GO:0042939	tripeptide transport	8	4	0.55	0.001
	GO:0016117	carotenoid biosynthetic process	20	6	1.38	0.002
	GO:0006744	ubiquinone biosynthetic process	5	3	0.34	0.003
	GO:0032259	methylation	88	15	6.05	0.004
	GO:0045814	negative regulation of gene expression, epigenetic	37	5	2.55	0.005
	GO:0009206	purine ribonucleoside triphosphate biosynthetic process	44	6	3.03	0.013
	GO:0000271	polysaccharide biosynthetic process	123	11	8.46	0.013
	GO:0009873	ethylene-activated signaling pathway	78	12	5.37	0.014
	GO:0015991	ATP hydrolysis coupled proton transport	9	3	0.62	0.020
	GO:0009718	anthocyanin-containing compound biosynthetic process	26	5	1.79	0.030
	GO:0000018	regulation of DNA recombination	5	2	0.34	0.041
	GO:0015976	carbon utilization	5	2	0.34	0.041
	GO:0034314	Arp2/3 complex-mediated actin nucleation	5	2	0.34	0.041
	GO:0080144	amino acid homeostasis	5	2	0.34	0.041
	GO:1901615	organic hydroxy compound metabolic process	121	9	8.32	0.041
	GO:0000902	cell morphogenesis	174	9	11.97	0.042

Table 3. Gene Ontology enrichment for the network periphery.

	GO.ID	Term	Annotated	Significant	Expected	P
A. Bud stage periphery	GO:0016579	protein deubiquitination	24	16	8.70	0.002
	GO:0048825	cotyledon development	21	12	7.61	0.005
	GO:0015846	polyamine transport	5	5	1.81	0.006
	GO:0034314	Arp2/3 complex-mediated actin nucleation	5	5	1.81	0.006
	GO:0006020	inositol metabolic process	12	7	4.35	0.011
	GO:0010229	inflorescence development	9	7	3.26	0.014
	GO:0009744	response to sucrose	31	16	11.24	0.021
	GO:1903338	regulation of cell wall organization or biogenesis	11	7	3.99	0.026
	GO:0006809	nitric oxide biosynthetic process	6	5	2.18	0.026
	GO:0009773	photosynthetic electron transport in photosystem I	8	6	2.90	0.030
	GO:0034613	cellular protein localization	183	75	66.34	0.031
	GO:0006004	fucose metabolic process	12	8	4.35	0.032
	GO:0009926	auxin polar transport	42	22	15.23	0.037
	GO:0010090	trichome morphogenesis	38	19	13.78	0.041
	GO:0009414	response to water deprivation	182	69	65.98	0.045
	GO:0009873	ethylene-activated signaling pathway	78	38	28.28	0.046
	GO:0007166	cell surface receptor signaling pathway	93	42	33.71	0.046
	GO:0006163	purine nucleotide metabolic process	96	38	34.80	0.047
	GO:0035195	gene silencing by miRNA	15	7	5.44	0.048
	GO:0016236	macroautophagy	21	9	7.61	0.048
	GO:0055083	monovalent inorganic anion homeostasis	13	5	4.71	0.048
	GO:0051338	regulation of transferase activity	38	13	13.78	0.048
	GO:0006473	protein acetylation	23	7	8.34	0.048
	GO.ID	Term	Annotated	Significant	Expected	P
B. D stage periphery	GO:0045490	pectin catabolic process	20	19	12.88	0.002
	GO:0007166	cell surface receptor signaling pathway	93	71	59.90	0.006
	GO:0000398	mRNA splicing, via spliceosome	46	38	29.63	0.007
	GO:0009740	gibberellic acid mediated signaling pathway	31	24	19.97	0.015
	GO:0006310	DNA recombination	54	41	34.78	0.015
	GO:0009909	regulation of flower development	80	57	51.53	0.017
	GO:0009058	biosynthetic process	2121	1360	1366.19	0.019
	GO:0042545	cell wall modification	34	26	21.90	0.021
	GO:0001708	cell fate specification	13	12	8.37	0.027
	GO:0042538	hyperosmotic salinity response	25	21	16.10	0.027
	GO:0002181	cytoplasmic translation	21	18	13.53	0.029
	GO:0006897	endocytosis	40	27	25.76	0.029
	GO:0010197	polar nucleus fusion	20	17	12.88	0.039
	GO:0071702	organic substance transport	492	319	316.91	0.046
	GO:0006949	syncytium formation	7	7	4.51	0.046
	GO:0008295	spermidine biosynthetic process	7	7	4.51	0.046
	GO:0009963	positive regulation of flavonoid biosynthetic process	7	7	4.51	0.046

Table 4. Gene Ontology enrichment in Bud stage modules correlated to pollination syndrome.

A. Bee pollination

GO.ID	Term	Annotated	Significant	Expected	P
GO:0046686	response to cadmium ion	99	16	5.69	0.000
GO:0010104	regulation of ethylene-activated signaling pathway	12	3	0.69	0.003
GO:0010189	vitamin E biosynthetic process	8	3	0.46	0.008
GO:0070085	glycosylation	72	7	4.13	0.009
GO:0009226	nucleotide-sugar biosynthetic process	21	5	1.21	0.011
GO:0042136	neurotransmitter biosynthetic process	9	4	0.52	0.018
GO:0070592	cell wall polysaccharide biosynthetic process	32	4	1.84	0.018
GO:0051173	positive regulation of nitrogen compound metabolic process	192	4	11.03	0.019
GO:0046777	protein autophosphorylation	71	9	4.08	0.019
GO:0009651	response to salt stress	215	19	12.35	0.022
GO:0009738	abscisic acid-activated signaling pathway	135	11	7.75	0.028
GO:0015718	monocarboxylic acid transport	12	3	0.69	0.028
GO:0046889	positive regulation of lipid biosynthetic process	12	3	0.69	0.029
GO:0006809	nitric oxide biosynthetic process	5	2	0.29	0.029
GO:0006974	cellular response to DNA damage stimulus	99	9	5.69	0.035
GO:0010099	regulation of photomorphogenesis	6	2	0.34	0.042
GO:0071805	potassium ion transmembrane transport	6	2	0.34	0.042

B. Butterfly pollination

GO.ID	Term	Annotated	Significant	Expected	P
GO:0009415	response to water	155	13	11.47	0.016
GO:0008380	RNA splicing	122	16	9.03	0.016
GO:0010223	secondary shoot formation	14	4	1.04	0.016
GO:0009624	response to nematode	15	4	1.11	0.021
GO:0009695	jasmonic acid biosynthetic process	9	3	0.67	0.024
GO:0090691	formation of plant organ boundary	9	3	0.67	0.024
GO:0006950	response to stress	1173	83	86.79	0.028
GO:0006353	DNA-templated transcription, termination	17	4	1.26	0.032
GO:0009567	double fertilization forming a zygote and endosperm	11	3	0.81	0.042
GO:0009835	fruit ripening	11	3	0.81	0.042
GO:0009607	response to biotic stimulus	374	35	27.67	0.045

GO:0010216	maintenance of DNA methylation	5	2	0.37	0.047
GO:0010363	regulation of plant-type hypersensitive response	5	2	0.37	0.047
GO:0071577	zinc ion transmembrane transport	5	2	0.37	0.047
GO:0009825	multidimensional cell growth	19	4	1.41	0.047

C. Hummingbird pollination

GO.ID	Term	Annotated	Significant	Expected	P
GO:0009451	RNA modification	353	89	62.34	0.000
GO:1900618	regulation of shoot system morphogenesis	9	5	1.59	0.001
GO:0009826	unidimensional cell growth	95	24	16.78	0.002
GO:0016556	mRNA modification	35	11	6.18	0.002
GO:0006012	galactose metabolic process	5	4	0.88	0.004
GO:0031425	chloroplast RNA processing	18	8	3.18	0.008
GO:0008643	carbohydrate transport	44	15	7.77	0.010
GO:1905428	regulation of plant organ formation	6	4	1.06	0.011
GO:0006888	ER to Golgi vesicle-mediated transport	19	8	3.36	0.011
GO:0044272	sulfur compound biosynthetic process	58	12	10.24	0.011
GO:0045944	positive regulation of transcription by RNA polymerase II	37	13	6.53	0.013
GO:0002239	response to oomycetes	18	6	3.18	0.019
GO:0006813	potassium ion transport	16	7	2.83	0.020
GO:0043650	dicarboxylic acid biosynthetic process	10	5	1.77	0.020
GO:0051053	negative regulation of DNA metabolic process	7	4	1.24	0.022
GO:0051301	cell division	147	30	25.96	0.027
GO:0000079	regulation of cyclin-dependent protein serine/threonine kinase activity	11	5	1.94	0.031
GO:0046131	pyrimidine ribonucleoside metabolic process	12	3	2.12	0.031
GO:0032509	endosome transport via multivesicular body sorting pathway	8	4	1.41	0.037
GO:0071496	cellular response to external stimulus	71	17	12.54	0.041
GO:1902117	positive regulation of organelle assembly	5	3	0.88	0.041
GO:0048438	floral whorl development	69	12	12.19	0.042
GO:0009651	response to salt stress	215	49	37.97	0.043

Table 5. Gene Ontology enrichment in D stage modules correlated to pollination syndrome.

A. Bee pollination					
GO.ID	Term	Annotated	Significant	Expected	P
GO:0061014	positive regulation of mRNA catabolic process	6	3	0.26	0.002
GO:0010189	vitamin E biosynthetic process	8	3	0.35	0.004
GO:0044093	positive regulation of molecular function	46	4	2.01	0.006
GO:0046686	response to cadmium ion	99	10	4.33	0.011
GO:0006974	cellular response to DNA damage stimulus	99	8	4.33	0.017
GO:0002833	positive regulation of response to biotic stimulus	19	3	0.83	0.017
GO:0009395	phospholipid catabolic process	5	2	0.22	0.017
GO:0009969	xyloglucan biosynthetic process	5	2	0.22	0.017
GO:0006544	glycine metabolic process	7	2	0.31	0.035
GO:0043488	regulation of mRNA stability	7	2	0.31	0.035
GO:0016569	covalent chromatin modification	58	4	2.54	0.044
GO:0043902	positive regulation of multi-organism process	22	3	0.96	0.045
GO:0009631	cold acclimation	8	2	0.35	0.045

B. Butterfly pollination					
GO.ID	Term	Annotated	Significant	Expected	P
GO:0009853	photorespiration	19	7	2.00	0.002
GO:0043622	cortical microtubule organization	13	5	1.37	0.008
GO:0031349	positive regulation of defense response	51	8	5.37	0.010
GO:0006575	cellular modified amino acid metabolic process	35	7	3.69	0.018
GO:0070919	production of siRNA involved in chromatin silencing by small RNA	6	3	0.63	0.018
GO:0042398	cellular modified amino acid biosynthetic process	7	3	0.74	0.029
GO:0009808	lignin metabolic process	22	5	2.32	0.031
GO:2000030	regulation of response to red or far red light	9	3	0.95	0.031
GO:0019220	regulation of phosphate metabolic process	39	6	4.11	0.031
GO:0010197	polar nucleus fusion	13	4	1.37	0.040
GO:0051028	mRNA transport	44	9	4.63	0.041
GO:0006833	water transport	13	4	1.37	0.043
GO:0043902	positive regulation of multi-organism process	22	4	2.32	0.044

C. Hummingbird pollination					
GO.ID	Term	Annotated	Significant	Expected	P

GO ID	Description				
GO:0009960	endosperm development	10	6	1.53	0.002
GO:0007163	establishment or maintenance of cell polarity	5	4	0.77	0.002
GO:0000272	polysaccharide catabolic process	50	14	7.65	0.005
GO:0010224	response to UV-B	24	9	3.67	0.007
GO:0009089	lysine biosynthetic process via diaminopimelate	7	4	1.07	0.013
GO:0009451	RNA modification	353	70	54.01	0.015
GO:0007166	cell surface receptor signaling pathway	86	21	13.16	0.018
GO:0008285	negative regulation of cell proliferation	8	4	1.22	0.023
GO:0042939	tripeptide transport	8	4	1.22	0.023
GO:0009955	adaxial/abaxial pattern specification	17	8	2.60	0.023
GO:0051656	establishment of organelle localization	36	9	5.51	0.023
GO:0006213	pyrimidine nucleoside metabolic process	14	3	2.14	0.023
GO:0033014	tetrapyrrole biosynthetic process	39	6	5.97	0.024
GO:0009267	cellular response to starvation	47	14	7.19	0.025
GO:0006635	fatty acid beta-oxidation	16	6	2.45	0.026
GO:0016556	mRNA modification	35	9	5.36	0.026
GO:0048438	floral whorl development	69	14	10.56	0.028
GO:0010103	stomatal complex morphogenesis	14	5	2.14	0.028
GO:0006383	transcription by RNA polymerase III	5	3	0.77	0.028
GO:0040008	regulation of growth	126	26	19.28	0.029
GO:0010019	chloroplast-nucleus signaling pathway	9	4	1.38	0.036
GO:0010506	regulation of autophagy	9	4	1.38	0.036
GO:0043547	positive regulation of GTPase activity	9	4	1.38	0.036
GO:0018298	protein-chromophore linkage	27	8	4.13	0.044
GO:0031425	chloroplast RNA processing	18	6	2.75	0.045
GO:0051187	cofactor catabolic process	34	9	5.20	0.050
GO:0002238	response to molecule of fungal origin	6	3	0.92	0.050
GO:0032958	inositol phosphate biosynthetic process	6	3	0.92	0.050
GO:0042219	cellular modified amino acid catabolic process	6	3	0.92	0.050
GO:1905428	regulation of plant organ formation	6	3	0.92	0.050
GO:0065004	protein-DNA complex assembly	45	8	6.89	0.050

Scale bars equals 1 cm.

Figure 1. *Achimenes* flowers and the sampled developmental stages.

Figure 2. Phylogenetic relationships and floral traits in *Achimenes*.

A

Cluster Dendrogram

B

Cluster Dendrogram

Figure 3. Co-expression clustering of the Bud and D stages.

Correspondence of Bud and D stage modules

Figure 4. Correspondence of the Bud and D stage networks.

Figure 5. Summary of Bud stage network module association to floral traits and pollination syndrome.

Figure 6. Summary of D stage network module association to floral traits and pollination syndrome.

Figure 7. Relationship between orthogroup dN/dS (omega) and orthogroup connectivity.

Supplemental Figure 1. Module eigengenes in the Bud stage network.

Supplemental Figure 2. Module eigengenes in the D stage network.

Supplemental Figure 3. Effects of floral traits on the eigengene expression in the Bud stage modules summarized over 50 phylogenetic trees.

Supplemental Figure 4. Effects of pollination syndrome on the eigengene expression in the Bud stage modules summarized over 50 phylogenetic trees.

Supplemental Figure 5. Effects of floral traits on the eigengene expression in the D stage modules summarized over 50 phylogenetic trees.

Supplemental Figure 6. Effects of pollination syndrome on the eigengene expression in the D stage modules summarized over 50 phylogenetic trees.

Supplemental Figure 7. Relationship between orthogroup dN/dS (omega) and average expression.

Supplemental Figure 8. Distribution of dN/dS (omega) among hubs and periphery nodes.