1

Title: External and Internal Signal-to-noise Ratios Alter Timing and Location of Cortical Activities During Speech-in-noise Perception

Abbreviated title: Neural correlates of speech-in-noise performance

Subong Kim,[1, #] Adam T. Schwalje,[2, #] Andrew S. Liu,[2] Phillip E. Gander,[3] Bob McMurray, [1,2,4] Timothy D. Griffiths,[5] and Inyong Choi[1, 2, *]

[1] Department of Communication Sciences and Disorders, University of Iowa, Iowa City, IA 52242, USA

[2] Department of Otolaryngology – Head and Neck Surgery, University of Iowa Hospitals and Clinics, Iowa City, IA 52242, USA

[3] Department of Neurosurgery, University of Iowa Hospitals and Clinics, Iowa City, IA 52242, USA

[4] Department of Psychological and Brain Sciences, University of Iowa, Iowa City, IA 52242, USA

[5] Institute of Neuroscience, Newcastle University, Newcastle upon Tyne, NE1 7RU, UK

[#] SK and ATS contributed equally to this work.

[*] Corresponding author at: Department of Communication Sciences and Disorders, University of Iowa, 250 Hawkins Dr., Iowa City, IA 52242, USA. Email address: inyong-choi@uiowa.edu

Number of pages: 34

Number of figures: 6

Number of words: 238 (abstract), 1185 (introduction), and 1862 words (discussion)

Author contributions:

S.K. and A.T.S. contributed equally to this work. I.C. designed the experiments. S.K. ran the experiments. S.K., A.T.S., A.S.L., P.E.G., B.M., T.D.G., and I.C. analyzed and interpreted data.

S.K., A.T.S., and I.C. prepared the manuscript, with revisions and suggestions from A.S.L,

P.E.G., B.M., and T.D.G..

Acknowledgments

This work was supported by Hearing Health Foundation Emerging Research Grant and

departmental start-up awarded to Inyong Choi, and NIH T32 (5T32DC000040-24) and NIDCD

P50 (DC000242 31A1) awarded to Bruce Gantz. The authors declare no competing financial

interests.

Abstract

Understanding speech in noise (SiN) is a complex task that recruits multiple cortical

subsystems. There is a variance in individuals' ability to understand SiN that cannot be

explained by simple hearing profiles, which suggests that central factors may underlie the

variance in SiN ability. Characterizing central functions that exhibit individual differences during

a SiN task and finding their relative contributions to predicting SiN performance can reveal key

neural mechanisms of SiN understanding. Here, we elucidated a few cortical functions involved

during a SiN task and their hierarchical relationship using both within- and across-subject

approaches. Through our within-subject analysis of source-localized electroencephalography,

we demonstrated how acoustic signal-to-noise ratio (SNR) alters neural activities along the

auditory-motor pathway, or dorsal stream, of speech perception. In quieter noise, left

supramarginal gyrus (SMG, BA40) exhibited dominant activity at an early timing (~300 ms after

word onset). In contrast, in louder noise, left inferior frontal gyrus (IFG, BA44) showed dominant

activity at a later timing (~700 ms). Further, through an individual differences approach, we

showed that listeners show different neural sensitivity to the background noise and target

speech, reflected in the amplitude ratio of cortical responses to speech and noise, named as an

"internal SNR." We found the "softer noise" pattern of activity in listeners with better internal

SNR, who also performed better. This result implies that how well a listener "unmask" target

speech from noise determines the subsequent speech analysis and SiN performance.

Significance

This study elucidated crucial cortical mechanisms underlying speech-in-noise perception using

both within- and across-subject design approaches. We found that cortical auditory evoked

responses to speech involved early activation in the temporo-parietal cortex in an easy condition

while a hard condition cortical activity involved late activation in the frontal cortex. Importantly,

the across-subject analysis showed that pre-speech time cortical activity predicts post-speech

time activity, in such a way that good performers with better neural suppression of background

noise show cortical activity similar to the pattern observed in the easier condition regardless of

given acoustic noise level. This suggests a critical role of pre-lexical sensory gain control

processes affecting performance and cognitive load during speech-in-noise perception.

Introduction

Understanding speech in noise (SiN) is essential for communication in social settings.

Yet young normal-hearing listeners are remarkably good at this: even in challenging SiN

conditions where the speech and noise have the same intensity (i.e., 0 dB signal-to-noise ratio:

SNR) and overlapped frequency components, they often recognize nearly 90% of sentences

correctly (Ohlenforst et al., 2017). This suggests a remarkable capacity of the auditory system to

cope with noise. However, the ability to understand SiN degrades severely with increased

background noise level (Ohlenforst et al., 2017), hearing loss (Harris and Swenson, 1990),

and/or aging (Nabelek, 1988). Moreover, recent studies show that normal hearing listeners

show large individual differences in SiN performance (Liberman et al., 2016). By linking this

variable ability for SiN perception to cortical activity, we may be able to understand the neural

mechanisms by which humans accomplish this ability, and this may shape our understanding of

how best to remediate hearing loss.

Different neuro-cognitive mechanisms might give rise to better or worse SiN

performance. First, listeners may vary in the mechanisms for representing sound in the

ascending pathway to the auditory cortex (AC) and in auditory scene analysis (Bregman, 1999),

both of which are required to separate signal from noise. Auditory scene analysis processes can

inhibit neural responses to task-irrelevant sensory inputs even before the target sound is heard,

based on expectations of differences between the target and masker such as differences in their

spectra (Lee et al., 2013), location (Frey et al., 2014; Goldberg et al., 2014), and timing (Lange,

2009). A successful auditory scene analysis during a SiN task will unmask the target speech

from maskers, which will enhance effective signal-to-noise ratio (SNR) in the neural pathway.

Second, listeners might vary in neural mechanisms for representing speech in the temporo-

parietal-frontal language network that might compensate for a noisy signal. Current models of

the neural processing of speech suggest two distinct cortical networks (i.e., dorsal and ventral

stream) for speech processing (Scott and Johnsrude, 2003; Hickok and Poeppel, 2007; Myers

et al., 2009; Gow, 2012). Such a division of labor is highlighted by work showing that speech in

noise engages the dorsal stream more strongly than in quiet (Hickok and Poeppel, 2007;

Liebenthal et al., 2013; Bidelman and Howell, 2016; Du et al., 2016). However, we have limited

knowledge of the functional roles of these two cortical pathways, their timing, and their

hierarchical relation.

Signal separation and compensatory mechanisms can be distinguished by using

functional neuroimaging in an individual differences approach. That is, we can compare the

degree to which accuracy in a SiN task is correlated to either pattern of activity in auditory and

related (signal analysis), or with frontal areas involving articulation and decision making

(compensation)[1]. A few have studies asked how individual differences in cortical pathways

correlate with SiN performance. These suggest activity in frontal areas (e.g., inferior frontal

gyrus: IFG) predict SiNs performance (Wong et al., 2009; Bidelman and Howell, 2016; Du et al.,

2016). They largely did not find an influence of lower-level areas; however, as we discuss,

methodological limits may have prevented this. Importantly, no work has examined both dorsal

and ventral processes simultaneously to determine if signal separation or compensatory speech

pathways capture unique variance in performance. Our central hypothesis is that the quality of

early signal analysis that occurs before auditory-motor transformations, rather than variation in

later compensatory processes dependent on the dorsal pathway, uniquely predicts speech

perception accuracy.

A secondary test of the importance of each pathway is the relative timing of activity in

these pathways during speech processing. Functionally, work using eye-movements in the

---

[1] A third possible mechanism—differences in auditory attention—likely spans both networks. Auditory attention likely originates in a fronto-parietal network involving the inferior frontal gyrus (IFG), the superior parietal sulcus, and the intraparietal sulcus (Teki et al., 2011; Hausfeld et al., 2018), but affects early-stage auditory activity (Choi et al., 2013; Choi et al., 2014; Bressler et al., 2017). Under an attentional account, if the most variance amongst listeners is due to differences in deploying attention, we should see SiN performance primarily correlated with frontal activity; whereas if most variance is due to using attention to clean up the signal it should primarily be associated with the auditory and related cortex.

visual world paradigm has extensively characterized the time course of word recognition in both

quiet (Allopenna et al., 1998; Dahan and Gareth Gaskell, 2007; Magnuson et al., 2007) and

under challenging conditions such as noise or signal degradation (Huettig and Altmann, 2005;

Ben-David et al., 2011; McQueen and Huettig, 2012; Brouwer and Bradlow, 2016; McMurray et

al., 2017). In general, this work suggests that in quiet, listeners activate a range of lexical

candidates immediately at the onset of the auditory stimulus. This *lexical competition* resolves

rapidly by around 250 ms after the uniqueness point of the word. A moderate amount of

degradation or noise typically imposes about a 75-100 ms delay on this recognition process

(Ben-David et al., 2011; Farris-Trimble et al., 2014), where severe degradation can delay lexical

access by up to 250 ms (Farris-Trimble et al., 2014; McMurray et al., 2017). This is particularly

relevant for evaluating the causal role of downstream compensatory processes – if such

processes are later than 400-500 ms, this may be too late to reflect lexical competition.

However, the timing of cortical activity within each pathway is largely unknown, as most of the

work on speech in noise perception has been conducted with functional magnetic resonance

imaging (fMRI) (Wong et al., 2008; Wong et al., 2009; Du et al., 2014, 2016) which has the poor

temporal resolution.

Assuming that left supramarginal gyrus (SMG: the anterior part of the inferior parietal

lobule) is an early stage in the dorsal pathway works as an interface between

auditory/phonological representations in the superior temporal gyrus and motor ones in the

frontal lobe (Binder et al., 2004; Gow, 2012), we expect to see early SMG activity in the less

adverse listening situation. In contrast, we hypothesize that noisier listening conditions will

evoke late activities in the frontal area, reflecting downstream compensatory processes.

The present study tests above hypotheses using both a within-subject design and

individual differences approaches. First, we identify how both primary auditory pathways and

frontal compensatory processes differ in noise. This within-subject design examined the effect of

acoustic SNR on the a) timing and b) location of cortical activity during word-in-noise

recognition. For this, we used two distinct high-density electroencephalographic (EEG)

analyses: 1) Hypothesis-driven source estimation that examined time courses of evoked

responses within two regions-of interest (ROIs) – SMG and IFG; and 2) sensor-space

microstate analysis as a data-driven approach that cross-validates the ROI-based analysis.

Next, we assessed the relative and unique contributions of both primary and compensatory

cortical processes in predicting SiN performance. We use the same EEG data to quantify an

individual's speech unmasking ability by computing the ratio of cortical evoked responses to

noise- and target speech-onset, or "internal SNR." We also quantify post-speech-time neural

activity in the dorsal speech-motor pathway. These were then used in a regression analysis to

determine the relative contribution of each to SiN performance.

150   **Materials and Methods**

151   *Participants*

152   All study procedures were reviewed and approved by the local Institutional Review

Board. Thirty subjects between 19 and 31 years of age (mean = 22.4 years, SD = 2.8 years;

median = 22 years; 9 (30%) male) were recruited from a population of students at a large

Midwestern university. All subjects were native speakers of American English, with normal

hearing thresholds no worse than 20 dB HL at any frequency, tested in octaves from 250 to

8000 Hz. Four subjects (1 male) were excluded from the analysis due to signal contamination

across several EEG channels.

Task design and procedures

161   We simultaneously measured behavioral performance and cortical neural activity a short

162   (15 minute) experimental sessions.

163   Each trial (**Figure 1**) began with the presentation of a fixation cross ('+') on the screen.

Listeners were asked to fix their gaze on this throughout the trial to minimize eye-movement

artifacts. Next, they heard the cue phrase "check the word." This enabled listeners to predict the

timing of next acoustic event (the noise onset). After fixed-duration (700 ms) silence that

followed the cue phrase, multi-talker babble noise started and continued for 2 seconds. One

second after the noise onset, the target word was heard 100 ms after the composite auditory

stimulus (noise + word) offset, four written choices appeared on the screen. The response

options differed either in the initial or the final consonant (e.g., 'than,' 'van,' 'ban,' and 'pan,' for

target word 'ban'; 'hit,' 'hip,' 'hiss,' 'hitch' for target word 'hiss'). Subjects pressed a button on a

keypad to indicate their choice. No feedback was given to the subject at the end of a trial. The

next trial began 1 second after the button press. This trial structure was intended to minimize

visual, pre-motor, and motor artifacts during the time of interest surrounding the auditory stimuli.

The timing and intervals of auditory stimuli (i.e., cue phrase, noise, and target) were intended to

derive well-distinct cortical evoked responses to the onsets of background noise and target

word.

Figure 1. Trial and stimulus structure. Every trial starts with the cue phrase "check the word." A target word starts 1 second after the noise onset. Four choices are given after the word ends; subjects select the correct answer with a keypad. No feedback is given. The noise level is manipulated to create high (+3 dB) and low (-3 dB) SNR conditions. Subjects complete 50 trials for each condition.

Since we are particularly interested in SMG/IFG regions that are involved in

phonological/lexical processing (Hickok and Poeppel, 2007), we elected to use natural

monosyllabic words, rather than simpler non-sense speech tokens used by prior studies

(Parbery-Clark et al., 2009; Bidelman and Howell, 2016). This engaged lexical processing,

placed high demands on cortical processing, and maximized ecological validity (Gagne et al.,

2017). Thus, target words consisted of 100 hundred CVC words from the California Consonant

Test (CCT) (Owens and Schubert, 1977), spoken by a male speaker with a General American

accent.

The RMS level of noise was either 68 and 62 dB SPL, and target word was always

presented at 65 dB SPL. This led to either +3 or -3dB SNR (referred to as "high SNR" and "low

SNR," respectively). Fifty words were presented at each SNR (±3 dB). -3 dB SNR was chosen

to emulate a highly effortful listening condition yielding ~70% accuracy from pilot experiments.

+3 dB SNR condition emulates an easy listening condition.

The task was implemented using the Psychtoolbox 3 package (Brainard, 1997; Pelli,

1997) for Matlab (R2016b, The Mathworks). Participants were tested a sound-treated,

electrically shielded booth with a single loudspeaker (model #LOFT40, JBL) positioned at a 0°

azimuth angle at a distance of 1.2 m. A computer monitor was located 0.5m in front of the

subject at eye level. The auditory stimuli were presented at the same levels for all subjects.

EEG acquisition and preprocessing

Scalp electrical activity (EEG) was recorded during the SiN task using the BioSemi

ActiveTwo system at a 2048 Hz sampling rate. Sixty-four active electrodes were placed

according to the international 10-20 configuration. Trigger signals were sent from Matlab

(R2016b, The Mathworks) to the ActiView acquisition software (BioSemi). The recorded EEG

data from each channel were bandpass filtered from 1 to 50 Hz using a 2048-point FIR filter.

Epochs were extracted from -500 ms to 3 s relative to stimulus onset. After baseline correction

using the average voltage between -200 and 0 ms, epochs were down-sampled to 256 Hz.

Since we were interested in the speech-evoked responses from frontal brain regions, we

opted for a non-modifying approach to eye blink rejection: Trials that were contaminated by an

eye blink artifact were rejected based on the voltage value of the Fp1 electrode (bandpass

filtered between 1 and 20 Hz). Rejection thresholds for eye blink artifacts were chosen

individually for each subject, and separately for the noise and target word periods. After

rejecting bad trials, grand averages for each electrode were calculated for the two conditions.

For analysis of speech-evoked responses, we repeated baseline correction using the average

signal in the 300 ms preceding the word onset.

Source analysis

The source-space analysis was based on minimum norm estimation (Gramfort et al.,

2013; Gramfort et al., 2014) as a form of multiple sparse priors (Friston et al., 2008). After co-

registration of average electrode positions to the reconstructed average head model MRI, the

forward solution (a linear operator that transforms source-space signals to sensor space) was

computed using a single-compartment boundary-element model (Hämäläinen, 1989). The

cortical current distribution was estimated assuming that the orientation of the source is

perpendicular to the cortical mesh. Cross-channel EEG-noise covariance, computed for each

subject, was used to calculate the inverse operators. A noise-normalization procedure was used

to obtain dynamic statistical parametric maps (dSPMs) as z-scores (Dale et al., 2000). The

inverse solution estimated the source-space time courses of event-related activity at each of

10,242 cortical voxels per hemisphere. In the present study, two predetermined ROIs will be

included: (1) bilateral SMG, and (2) bilateral pars opercularis and pars triangularis of IFG. The

SMG is an early stage in the dorsal pathway and, among its many roles, works as an interface

between auditory/phonological representations in the superior temporal gyrus and motor ones in

frontal lobe including precentral/postcentral and IFG (Binder et al., 2004; Gow, 2012). To ensure

the fidelity of source localization at our ROIs, we applied the same analysis to HG, which

expected to be active in auditory tasks, before running ROI-based source analysis explained

below (**supplement Figure 1**).

Statistical analyses

    ROI-based source analysis. For ROI-based source analysis, one-sample *t*-tests were used to compare the distributions of mean source event-related potential (ERP) to zero (baseline voltage) in each SNR condition. The t-value envelope was then computed (as a form of smoothing). This was done by applying a bandpass filter, then calculating the absolute value of the Hilbert transform. The bandpass filter was set to one of two center frequencies, depending on the specific ROIs. Because the neural oscillations evoked by early ERP components such as N1-P2 have peak latencies of about 100 ms at their half cycle (approximately 5 Hz), and late ERP components such as N2-P3 have latencies of about 200 ms at their half cycle (approximately 2.5 Hz), the bandpass filter was set to between either 3 to 7 or 1 to 5 Hz to highlight these components. The *t*-value envelope calculated using the bandpass filter between 3 to 7 Hz was used to investigate HG and SMG ROIs, and earlier times of interest, while the envelope using the 1 – 5 Hz filter was used for IFG ROI and later times of interest. For each SNR condition, the whole brain *t*-value envelope time courses were obtained.

    Since we did not have individual structural MRI head models, it was not ideal to take the summed activity (mean or median) for all the voxels within ROIs. This is because individual difference in functional and anatomical structure of the brain may result in spatial blurring since current densities across adjacent voxels can overlap each other. Instead, representative voxels were identified for each ROI, for each SNR condition. We used a combination of previously-described methods to select voxels of interest that were used in fMRI studies (Tong et al., 2016). The cross-correlation coefficients for *t*-value envelopes between all voxels in an ROI were calculated across time (up to 800 ms after the target word onset), and then the mean coefficient was calculated for each voxel. The most representative voxel was defined as having the maximum value mean coefficient, while also being above threshold at two or more

continuous timepoints based on voxel's p value, as determined using one-sample *t*-tests (Tong

et al., 2016).

Once the most representative voxel was chosen for each SNR condition, a leave-one-

out procedure (i.e., Jackknife approach) was used to compare the population means between

the two SNR conditions, at those voxels, using paired t-tests. Prior to computing *p*-values, t-

statistics were adjusted for jackknifing (Luck, 2014). The positive false discovery rate (pFDR)

was estimated from those p-values after downsampling the time sequence according to Nyquist

theorem, and used to find timepoints that showed significant difference between SNR conditions

(Storey, 2002). Finally, the whole cortical surface source space was evaluated at those

timepoints.

Microstate analyses. Microstate analysis was conducted to cross-validate our ROI-

based analyses that assess effect of external SNR on ERPs at the group level. Microstates

describe transient, quasi-stable topographic orientations which provide information about the

timing of cognitive processes (Koenig et al., 2014). Microstates have been used to characterize

both resting state and event-related EEG activity (Ott et al., 2011; Schiller et al., 2016). The

microstate analysis was conducted on grand mean data (averaged across all subjects),

separately in each SNR condition and were implemented in the RAGU program (Koenig et al.,

2011; Koenig et al., 2014).

To conduct this analysis, we first identified four microstate cluster maps based on spatial

clustering of ERP topographies. To this, a k-means algorithm was used for microstate

identification with ten random re-initializations. As a cross-validation procedure, the 26 subjects

were randomly split 50 times into a training set and a test set, each comprising 13 subjects.

Next, the grand mean voltages at each timepoint of each SNR condition were assigned to the

best fit cluster map (Koenig et al., 2011; Koenig et al., 2014). To do this, each timepoint of the

ERP was assigned to the specific microstate cluster map that had the highest correlation

coefficients with the topography of the ERP, across all electrodes, at that timepoint.

To assess how well the grand mean ERPs were explained by the different microstate

clusters, we calculated the relative area of global field power (GFP) for each cluster, after

assigning the timepoints to microstates. The GFP at time t is equal to the standard deviation of

the signal at all N electrodes, which is defined as

$$(1)\;\; \text{GFP}(t) = \sqrt{\frac{\sum_{i=1}^{N}(v_i(t) - \bar{v}(t))^2}{N}},$$

where $v_i(t)$ is the voltage at electrode i, and $\bar{v}(t)$ is the average voltage across all electrodes at

time t.

After identification of the microstate explaining the most variance for grand average ERP

in each SNR condition, the timepoints of maximum GFP for that microstate were used to create

whole brain maps showing cortical source activity. The t-value envelope calculated by one-

sample t-tests in each SNR condition was used to investigate the source activity at early

timepoints (about up to 400 ms after the word onset) using the bandpass filter between 3 to 7

Hz, and the source activity at later timepoints (about after 400 ms) using the 1 – 5 Hz filter.

Regression approaches. To conduct multiple linear regressions, we used a jackknifing

approach (Stahl and Gibbons, 2004; Luck, 2014). In this approach the relevant neural factors

were computed for all subjects but one. This was repeated leaving out each subject in turn, and

the resulting data submitted to a linear regression with SiN performance (accuracy) as the

dependent variable, and SMG/IFG activation, and internal SNR as the predictor variables. Test

statistics were then adjusted to account for the fact that each data-point represents N-1

participants.

Correlation/regression analyses used these techniques to simultaneously examine

bottom-up and compensatory related SiN performance to three factors. Our first, analysis

examined raw cortical evoked activity to the target and noise. To best represent bilateral

auditory cortical activity, we used sensor-space ERP envelopes using the 3 – 7 Hz bandpass

filter from channel Cz at around 200 ms after the noise onset and at about 200 ms after the

target word onset, based on the timing determined by microstate analysis. Then, "internal SNR"

was defined as the ratio of target word-evoked ERP envelope peaks to noise-evoked ERP

envelope peaks magnitude described above, in dB scale, obtained from channel Cz (Equation

3). The internal SNR is different for each subject, and is separate from the fixed external, or

acoustic, SNR (here, ±3 dB).

317
$$(2) \ \ \mathrm{Internal} \ SNR = 20log_{10} \frac{\text{Word evoked potential}}{\text{Noise evoked potential}}$$

318　　　　　Second, to examine dorsal regions, we used cortical regions that were previously

319　　identified in the ROI-based source analysis described above. The peak magnitudes of the *t*-

320　　value envelopes were obtained for early SMG activation in the high SNR condition and late IFG

321　　activation in the low SNR condition.

323　　　　　　　　　　　　　　　　　　　**Results**

SiN performance

There was a large variance in performance among participants. This was observed in

both the high SNR condition (mean accuracy = 80.64%, SD = 7.81%, median = 83.01%; mean

reaction time = 1.53 s, SD = 0.32 s, median = 1.55 s) and the low SNR condition (mean

accuracy = 68.21%, SD = 8.92%, median = 70.37%; mean reaction time = 1.70 s, SD = 0.36 s,

median = 1.69 s). Both accuracy (*t*(25) = 6.99*, p* < 0.001, paired t-test) and reaction time (*t*(25) =

-3.81*, p* < 0.001, paired t-test) differed significantly between the two SNR conditions (**Figure**

2A). Reaction time and accuracy were correlated in the high SNR condition (**Figure 2B**,

Pearson correlation *r* = -0.50, *p* = 0.009), but not in the low SNR condition (**Figure 2C**, *r* = -0.19,

p = 0.34). As a whole, these results validate that the SNR manipulation was sufficient to create

more challenging speech perception conditions.

Figure 2. Behavioral results. **A.** Summary of behavioral performance for the two conditions (+3 and -3 dB SNR). Boxes denote the 25^{th} – 75^{th} percentile range; the horizontal bars in the center denote the median; the ranges are indicated by vertical dashed lines. Solid lines connect points for the same subject in different conditions. **B.** Average accuracy as a function of reaction time in +3 dB SNR condition. **C.** Average accuracy and reaction time in -3 dB SNR condition.

The effect of SNR on cortical activity

To assess the cortical activity, we converted sensor-space EEG signals to whole brain

source-activity. This allowed us to localize the effects of noise on targeted ROIs at specific

times. Within left hemisphere SMG, the high SNR condition showed significantly greater activity

than the low SNR condition from 200 to 330 ms (FDR adjusted *p* < 0.05) (**Figure 3B left**).

Within left hemisphere IFG, the low SNR condition showed significantly greater activity than the

high SNR condition from 740 to 830 ms (FDR adjusted *p* < 0.05) (**Figure 3B right**), throughout

triangular and opercula regions.

A visual representation of this finding can be seen in **Figure 3C**. Here, *t*-values reveal

significant differences between high and low SNR conditions. The timepoints were chosen for

display where the greatest number of voxels have significant differences. We observed

increased amplitude of the left hemisphere SMG activation in the high SNR condition at around

250 ms (*t*-value envelope). This may indicate efficient lexical processing in a relatively favorable

listening condition. However, in the low SNR condition, the peak amplitude of the left

hemisphere IFG activation (*t*-value envelope) seen around 700 ms after word onset. Given most

target words were around 500 ms in duration, SMG was primarily activated during the

presentation of the target word, while IFG was activated after its offset.

Figure 3. Region-of-interest (ROI) based source analysis. **A.** Cortical labels for two ROIs in left and right hemispheres: supramarginal gyrus (SMG), and the pars opercularis and triangularis of the inferior frontal gyrus (IFG), respectively. **B.** The time course of the *t*-value envelope, with the standard error of the mean (±1 SEM), obtained at representative voxels in each SNR condition (red color: +3 dB SNR, blue color: -3 dB SNR). An asterisk shows the timing of significant difference between +3 and -3 dB SNR conditions (paired *t*-test, FDR adjusted *p* < 0.05). **C.** Whole brain maps showing statistical contrasts (*t*-values obtained from paired *t*-tests between the two SNR conditions) of source activation at each voxel, only displaying those with *p*-value less than 0.05, at the timepoint that shows significant differences over the broadest area in the ROIs within the time range described above.

To more precisely timelock these neural events to the stimulus, the webMAUS (Kisler et

al., 2017) was used to identify the starting location of second and third phonemes in each of the

100 stimuli. A histogram of these acoustic time points is shown in **Figure 4A**. **Figure 4B** shows

the timing of SMG and IFG activity relative to the distribution of phoneme onsets in the target

word stimuli. This confirms that the early SMG activation occurs within the timecourse of target

words before the final phoneme is presented; the late IFG activation occurs after all words are

presented.

Figure 4. Timings of significant cortical activity relative to distributions of phonological events. **A.** Top and second panel show a histogram of the onsets of second and final phoneme of each stimulus. The third panel shows superimposed temporal envelopes extracted from waveforms of the 100 words. **B.** The whole brain maps at the bottom are from **Figure 3C** that shows statistical contrasts of source activation at the timepoints that show significant differences between the two SNR conditions. Purple curves on the cortical maps represent the conceptual illustration of ascending information flow through the dorsal pathway.

External (acoustic) SNR effects on timepoints and regions of interest based on

microstate analysis

To further assess temporal dynamics of neural activity during SiN perception in a data-

driven way, and to cross-validate our ROI-based analyses, we performed microstate analysis

(Lehmann, 1989b, a; Wackermann et al., 1993). Microstate analysis clusters time series of ERP

data into multiple different brief brain states, which may indicate different stages of the

information processing (Lehmann, 1989b). Four microstate maps were identified. (**Figure 5C**).

The grand mean ERP at each timepoint was assigned to one of the microstate clusters, and

GFP was calculated at those timepoints (**Figure 5B, C**).

Calculation of the relative area of GFP revealed that microstates 1 and 2 explained the

largest variance in sensor-space ERPs over time in low SNR (area = 37%) and high SNR (area

= 37%) condition, respectively. The timing, suggested by maximum GFP and the maximum

deflection of ERPs at frontal-central electrodes among the timepoints assigned to microstate 1,

was 0.668 seconds after word onset for the low SNR condition. The timing for microstate 2 was

0.320 seconds after word onset for the high SNR condition.

We next conducted a whole-brain source analysis at the timepoint assigned to

microstate 1 using one-sample t-tests against 0. This revealed increased activity in left

hemisphere IFG activation for the low SNR condition (**Figure 5D**). The same analysis at the

timepoint assigned to microstate 2 showed increased activity in the left hemisphere SMG for the

high SNR condition (**Figure 5D**). These results from microstate analysis are consistent with the

results of ROI-based analysis (illustrated in **Figure 3**) regarding timings and regions of neural

activation.

Figure 5. Microstate analysis. **A.** Evoked responses over time after the word onset for +3 and -3 dB SNR condition. Each color represents ERPs from a different channel of interest. **B.** Global field power (GFP) is calculated at each timepoint that is assigned to one of the microstate clusters. **C.** Four microstate cluster maps. Dark blue, light blue, red, and dark pink colors represent microstates 1, 2, 3, and 4, respectively. The relative area of GFP is calculated and reveals the highest value for the microstate 1 and 2 for -3 and +3 dB SNR condition, respectively. **D.** Whole brain maps obtained at the times assigned to microstates 1 and 2, that show maximum GFP and the maximum peak of ERPs at the frontal-central electrodes.

Individual differences in internal SNR predict SiN performance

For visualization purposes, we identified good and poor performers by conducting a

median split based on their performance in the task. **Figure 6A** shows GFP of the grand mean

ERPs for good and poor performance and topographies obtained at two timepoints identified by

microstate maps in the low SNR condition: 1) 220 ms after the noise onset, corresponding to the

auditory P2 to the noise; and 2) 240 ms after target word onset, corresponding to the AC-driven

N1 response. The word-evoked N1 was chosen as the first clearly AC-driven response to the

word onset as evidenced by its assignment to microstate 4 in the previous analysis, while the

timing of the noise-evoked P2 was suggested by microstate 3. Despite the same noise level for

these two groups of subjects, good performers exhibited less AC response to the background

noise, and greater AC response to the target word at central channels including Cz. This

validates that each component of the internal SNR measure seems to contribute separately to

SiN performance.

Figure 6. Individual differences in speech-in-noise processing. **A.** Global field power of the grand mean evoked potentials after the noise onset and after the target word onset, separately in the low SNR condition. Scalp topographies were examined at the timepoints, suggested by microstate analysis from **Figure 5**, and compared between good and poor performers, as determined by the median split. **B.** A series of scatter plots showing Pearson correlation coefficients among internal SNR, early SMG, late IFG activation, and behavioral accuracy. **C.** A scatter plot showing the regression coefficients from a linear regression model where behavioral accuracy is the dependent variable while internal SNR, early SMG, late IFG activation are the predictor variables. *The linear model significantly predicts behavioral accuracy while internal SNR is the only significant predictor.

To address our primary research question, we evaluated the simultaneous contribution

of primary auditory pathways and frontal compensatory processes. For this, we computed

internal SNR as the amplitude ratio of the noise and target related cortical evoked responses

(e.g., **Figure 6A**), expecting to quantify an individual's speech unmasking ability. We also

computed the mean activity in SMG and IFG separately averaged over 150 to 350 ms (the early

component), and 500 to 800 ms (late).

Figure 6B shows correlations among internal SNR, SMG/IFG source activation, and SiN

performance (accuracy) in the low SNR condition, that are obtained from 26 leave-one-out

grand averages using jackknifing approach (Luck, 2014; Stahl & Gibbons, 2004). Greater

internal SNR predicted better performance ($R = 0.45$, $p = 0.022$). Stronger early SMG activation

also predicted better performance ($R = 0.39$, $p = 0.047$). However, greater late IFG activity

predicted poorer performance ($R = -0.42$, $p = 0.033$). This suggests that both internal SNR and

SMG activation are positively related to SiN performance, while IFG activation has negative

relation to the performance.

To test their joint contributions, we conducted a linear regression in which internal SNR,

SMG, IFG activation were simultaneously related to SiN performance (**Figure 6C**). As a whole,

these factors accounted for a large proportion of the variance ($R = 0.65$, $p = 0.00033$). However,

among these three factors, internal SNR was the only significant predictor ($t = 2.25$, $df = 22$, $p =$

0.035), whereas SMG ($t = 1.80$, $df = 22$, $p > 0.05$) and IFG activation ($t = -2.03$, $df = 22$, $p >$

0.05) did not significantly contribute in the prediction SiN performance. This suggests that

internal SNR (representing the contribution of lower level signal analysis) uniquely predicts

variation in accuracy.

Discussion

We found SNR effects on timing and location of cortical activity for a speech-in-noise

recognition task. In a relatively easy SNR condition (in which subjects achieved ~80%

accuracy), left SMG showed a relatively early evoked response (~250 ms after target word

onset). In contrast, a challenging SNR (~68% accuracy) elicited the late response in left IFG

(~700 ms after target word onset). Within the same "external" SNR condition, individual

differences in such SMG and IFG responses predicted SiN performance; good performers

showed stronger early SMG response while poor performers showed stronger late IFG activity.

Individual differences in the ratio of noise- to target word-evoked cortical responses—the

"internal SNR,"—also predicted SiN performance; subjects with lower internal SNR exhibited

poorer accuracy. Importantly, both SMG and IFG responses did not contribute to the prediction

of SiN performance when internal SNR was added to the linear regression model. These results

from correlational analyses could be explained by auditory scene analysis mechanisms for

target unmasking and by temporal cortical processes for speech perception. A poorer ability to

unmasking target speech from background noise may lead to increased frontal lobe processing

that is employed when lower-level auditory pathways are unable to secure favorable speech

quality due to background noise.

Internal SNR: A measure of pre-speech processing for speech unmasking

Pre-speech time cortical activity for speech unmasking should be localized to the

primary and secondary AC, appearing as enhanced neural representation of the target sound

(the speech) and suppressed neural representation of ignored stimuli (the noise). This is

consistent with work suggesting that the auditory N1 (the numerator of the measure) can be

localized to AC and the planum temporale (Schneider et al., 2002).

Such responses could reflect auditory selective attention, which shows a similar pattern

in previous studies (Hillyard et al., 1973; Hillyard et al., 1998; Mesgarani and Chang, 2012). In

the present study, good performers showed significantly weaker noise-evoked AC response,

compared with poor performers, approximately 200 ms after the noise onset (**Figure 6A**).

Decreased AC response to background noise in good performers is compatible with the

presence of a sensory gain control mechanism (Hillyard et al., 1998). The variation in the

sensory gain control may originate from multiple factors. It may reflect the acuity of encoding

spectro-temporal acoustic cues from speech and noise or grouping of such acoustic cues for

auditory object formation. It may also reflect endogenous mechanisms for active suppression of

background sounds along with neural enhancement of foreground sounds (Shinn-Cunningham

and Best, 2008).

Our goal was not to disentangle the sources of variation in sensory gain control but

rather to quantify the effectiveness of sensory gain control by internal SNR and test how it

predicts later speech processes and behavioral accuracy. In this regard, we found a significant

correlation between accuracy and the relative amplitude word- and noise-evoked potentials.

This demonstrates that individuals have a differential ability to suppress the noise effectively via

early auditory processes (indexed by internal SNR). The first-order correlations between IFG

and SMG activation and behavioral performance were compatible with the effect of SNR on

ERPs: Good performers had stronger SMG activation (as in the high SNR condition), while poor

performers had stronger IFG activation (as the low SNR condition). However, the amplitude of

SMG and IFG response did not uniquely contribute to accuracy when internal SNR was added

to the model. This indicates that changes in SMG or IFG activity are the outcome of pre-speech

sensory gain control processing, rather than an independent causal factor predicting speech

perception performance.

This result conflicts with some findings from earlier studies but also clarifies their

findings. For example, Wong et al. (2009) did not find a relationship between SiN performance

and AC or auditory related cortex. However, this study used fMRI, which may have missed the

contribution of much shorter-lived bottom-up processes (Parbery-Clark et al., 2009). Like us,

they did find correlations with activity in the IFG (and also the precentral gyrus a second dorsal

route site). However, this study included both younger and older adults (who showed

differences in cortical networks). Thus, some of these correlations may have been driven by age

differences. Also, it is unclear whether these differences in cortical activity are necessary for

successful SiN understanding (or at least helpful), as such differences could also reflect

processes like increased effort (consistent with the view that IFG may be a domain-general

control process) (Fedorenko et al., 2013), error monitoring, or even just increased uncertainty.

Such processes may be engaged by noise without necessarily playing a causal role in

improving perception.

Similarly, Bidelman and Howell (2016) related both AC and dorsal route activity to

performance. They found no contribution of AC, but a correlation between speech performance

and early (~115 ms) activity in the IFG. However, their measure of AC activity represented the

response to both the speech and noise, not the ability of AC to pull speech from the noise.

Moreover, speech perception accuracy was assessed in an offline task. As a result, the cortical

activity measures did not reflect cortical processes leading up to accurate (or inaccurate

response); thus, these correlations may reflect broader individual differences, rather than a

causal chain leading to accurate SiN processing.

SNR effect on timing and location of cortical activity

Previous studies have suggested that spoken-word recognition occurs via a process of

dynamic lexical competition as speech unfolds over time. For many words, this competition

begins to resolve (e.g., the target separates from competitors) around 300 ms after word onset

(Huettig and Altmann, 2005; Farris-Trimble and McMurray, 2013). In significantly challenging

conditions (high noise) however, lexical processing can be delayed about 250 ms until most of

the word has been heard (Farris-Trimble et al., 2014; McMurray et al., 2017), which may

minimize competition. Based on the timing predicted by these studies, we expected early SMG

processing in the high SNR condition, and late IFG processing in the low SNR condition.

Our secondary analysis, a data-driven approach based on spatiotemporal clustering

analysis of ERPs (microstate analysis), supports the conclusion from the ROI-based analysis.

As microstates 1 and 2 explained the greatest amount of the signal's variance in the low and the

high SNR condition, respectively, we focused on the highest GFP peak timepoints, within

corresponding microstates for each SNR condition. Whole brain maps obtained from those

timepoints were supportive of the ROI analysis: in the high SNR condition, SMG was strongly

activated in the left hemisphere, while left IFG and bilateral Heschl's gyrus (HG) were activated

in the low SNR condition.

Increased SMG activity between the second and the third phonemes (see **Figure 4**) in

the high SNR condition may indicate a neural substrate of immediate lexical access (Farris-

Trimble et al., 2014; McMurray et al., 2017), consistent with Gow (2012). This immediacy was

observed when speech sounds were relatively clean (high SNR), and it does not appear in

previous EEG studies using non-word synthesized phonemes (Bidelman and Dexter, 2015;

Bidelman and Howell, 2016). However, we note that Bidelman and Howell used a single non-

word stimulus (a vowel-consonant-vowel) that would not be expected to engage lexical

processing. Bidelman and Howell's results also demonstrated an early activity (~115 ms) in IFG

with a clearly intelligible VCV phoneme. This was not observed in our study. However, because

we used naturally spoken CVC words, we can limit the interpretation of the late IFG activity to

the decision-making process in which listeners are trying to clean up the results of lexical

competition in SMG.

The idea that greater IFG activity is linked with poorer SiN recognition performance

seems to be inconsistent with some fMRI studies that showed the positive correlation between

the IFG activity and speech recognition performance (Zekveld et al., 2006; Wong et al., 2009;

Vaden et al., 2015; Du et al., 2016). This may stem from the difference between fMRI and EEG

in temporal resolution and in sensitivity to either neural metabolic activity or the equivalent

current dipoles (Bridwell et al., 2013). In the present study, we exhibited event-related potentials

at a specific latency of ~700 ms after target word onset, i.e., ~200 ms after target word "offset."

Previous fMRI studies might have demonstrated IFG activity at different latencies or

accumulated BOLD signal that is not time-locked to a specific sensory event. Alternatively, this

difference between fMRI and the current EEG results might be due to an error in the estimation

of our source location. However, the fact that both of our approaches – an ROI-based approach

and a data-driven approach without a space constraint – resulted in exhibiting the same IFG

activity with a similar latency might make our results more reliable.

So, what is IFG contributing to the process? It is unknown whether these processes are

due to active compensation for the noise or increased effort (both of which may help) or are a

simply marker of increased response uncertainty. Our correlational analyses do not suggest a

causal role for frontal activity in predicting an individual's accuracy. The first order correlation

was negative – more frontal activity was linked to *less* overall accuracy. More importantly, this

correlation was not significant when internal SNR was added to the model suggesting IFG

activity does not offer a unique contribution to accuracy. Rather it may simply reflect the clarity

of the signal offered by the earlier auditory processes that deal with noise. That is, if we view

IFG as primarily serving a decision-making role in this task when dorsal route areas do not

output clear representations of the signal, IFG must work harder to resolve on a decision. It may

be then that activity in frontal areas is not causally necessary for good SiN performance, but

rather reflects the additional response uncertainty created by noisy listening situations. This

challenges accounts like Du et al. (2016) that argue for a causal role of dorsal route processing

in SiN understanding.

Methodological advances and justifications for source time course analysis

548   Our approach to identifying a single voxel within an ROI deserves a particular

discussion. Identification of the representative voxel of an ROI is a problem common to EEG

source analysis, fMRI, and other functional brain imaging studies. Many relevant neuroimaging

analysis approaches have been described, including univariate, multivariate, and machine

learning; however, most of these are intended for the identification of regions of interest or

functional connections from a whole brain map. Drawbacks of this type of whole-brain analysis

include the need for strict multiple comparisons correction and, therefore, decreased statistical

power. Using strong a priori hypotheses to generate regions of interest allowed us to circumvent

these issues, but still requires identification of representative voxels within our regions of

interest. Favored approaches generally require identification of peak activity within an ROI

(Tong et al., 2016). However, to avoid the assumption that choosing peak activity implies, we

opted instead to choose the voxel that has the maximum average correlation to every other

voxel within the ROI. In the present study, we chose not to constrain the location of the voxel of

interest within an ROI for each condition. Because our anatomic resolution is unlikely to be at

the voxel level, we elected to choose a different representative voxel for each condition,

unconstrained by the location of the representative voxel from other conditions.

565   **Conclusion**

We found that clean, intelligible speech elicits early processing at SMG, while sensory

degradation results in late processing at IFG for less intelligible speech. Better speech

unmasking in good performers modulated the ratio of cortical evoked responses to the

background noise and target sound, which effectively changed SNR internally, resulting in

facilitated lexical/phonological processing through SMG. These findings may collectively form a

neural substrate of individual differences in speech-in-noise understanding ability. Crucially,

however, only neural representation of SNR uniquely predicted variation in performance,

suggesting that individual differences in SiN comprehension are largely a matter of primary

processes that extract the signal from noise rather than later compensatory ones.

References

Allopenna PD, Magnuson JS, Tanenhaus MK (1998) Tracking the Time Course of Spoken Word Recognition Using Eye Movements: Evidence for Continuous Mapping Models. Journal of memory and language 38:419-439.

Ben-David BM, Chambers CG, Daneman M, Pichora-Fuller MK, Reingold EM, Schneider BA (2011) Effects of aging and noise on real-time spoken word recognition: evidence from eye movements. Journal of speech, language, and hearing research : JSLHR 54:243-262.

Bidelman GM, Dexter L (2015) Bilinguals at the "cocktail party": dissociable neural activity in auditory-linguistic brain regions reveals neurobiological basis for nonnative listeners' speech-in-noise recognition deficits. Brain Lang 143:32-41.

Bidelman GM, Howell M (2016) Functional changes in inter- and intra-hemispheric cortical processing underlying degraded speech perception. Neuroimage 124:581-590.

Binder JR, Liebenthal E, Possing ET, Medler DA, Ward BD (2004) Neural correlates of sensory and decision processes in auditory object identification. Nat Neurosci 7:295-301.

Brainard DH (1997) The Psychophysics Toolbox. Spat Vis 10:433-436.

Bregman AS (1999) Auditory scene analysis : the perceptual organization of sound. Cambridge, Mass.: MIT Press.

Bridwell DA, Wu L, Eichele T, Calhoun VD (2013) The spatiospectral characterization of brain networks: fusing concurrent EEG spectra and fMRI maps. Neuroimage 69:101-111.

Brouwer S, Bradlow AR (2016) The Temporal Dynamics of Spoken Word Recognition in Adverse Listening Conditions. Journal of psycholinguistic research 45:1151-1160.

Dahan D, Gareth Gaskell M (2007) The temporal dynamics of ambiguity resolution: Evidence from spoken-word recognition. Journal of memory and language 57:483-501.

Dale AM, Liu AK, Fischl BR, Buckner RL, Belliveau JW, Lewine JD, Halgren E (2000) Dynamic statistical parametric mapping: combining fMRI and MEG for high-resolution imaging of cortical activity. Neuron 26:55-67.

Du Y, Buchsbaum BR, Grady CL, Alain C (2014) Noise differentially impacts phoneme representations in the auditory and speech motor systems. Proc Natl Acad Sci U S A 111:7126-7131.

Du Y, Buchsbaum BR, Grady CL, Alain C (2016) Increased activity in frontal motor cortex compensates impaired speech perception in older adults. Nat Commun 7:12241.

Farris-Trimble A, McMurray B (2013) Test-retest reliability of eye tracking in the visual world paradigm for the study of real-time spoken word recognition. Journal of Speech Language and Hearing Research 56:1328-1345.

Farris-Trimble A, McMurray B, Cigrand N, Tomblin JB (2014) The process of spoken word recognition in the face of signal degradation. Journal of Experimental Psychology: Human Perception and Performance Journal of Experimental Psychology: Human Perception and Performance 40:308-327.

Fedorenko E, Duncan J, Kanwisher N (2013) Broad domain generality in focal regions of frontal and parietal cortex. Proceedings of the National Academy of Sciences Proceedings of the National Academy of Sciences 110:16616-16621.

Frey JN, Mainy N, Lachaux JP, Muller N, Bertrand O, Weisz N (2014) Selective Modulation of Auditory Cortical Alpha Activity in an Audiovisual Spatial Attention Task. JOURNAL OF NEUROSCIENCE 34:6634.

Friston K, Harrison L, Daunizeau J, Kiebel S, Phillips C, Trujillo-Barreto N, Henson R, Flandin G, Mattout J (2008) Multiple sparse priors for the M/EEG inverse problem. Neuroimage 39:1104-1120.

Gagne JP, Besser J, Lemke U (2017) Behavioral Assessment of Listening Effort Using a Dual-Task Paradigm. Trends Hear 21:1-25.

Goldberg HR, Choi I, Varghese LA, Bharadwaj H, Shinn-Cunningham BG (2014) Auditory
attention in a dynamic scene: Behavioral and electrophysiological correlates. The
Journal of the Acoustical Society of America The Journal of the Acoustical Society of
America 135:2415.
Gow DW, Jr. (2012) The cortical organization of lexical knowledge: a dual lexicon model of
spoken language processing. Brain Lang 121:273-288.
Gramfort A, Luessi M, Larson E, Engemann DA, Strohmeier D, Brodbeck C, Parkkonen L,
Hamalainen MS (2014) MNE software for processing MEG and EEG data. Neuroimage
86:446-460.
Gramfort A, Luessi M, Larson E, Engemann DA, Strohmeier D, Brodbeck C, Goj R, Jas M,
Brooks T, Parkkonen L, Hamalainen M (2013) MEG and EEG data analysis with MNE-
Python. Front Neurosci 7:267.
Hämäläinen MS, Sarvas, J. (1989) Realistic conductivity geometry model of the human head for
interpretation of neuromagnetic data. IEEE Trans Biomed Eng 36:165-171.
Harris RW, Swenson DW (1990) Effects of reverberation and noise on speech recognition by
adults with various amounts of sensorineural hearing impairment. Audiology : official
organ of the International Society of Audiology 29:314-321.
Hickok G, Poeppel D (2007) The cortical organization of speech processing. Nat Rev Neurosci
8:393-402.
Hillyard SA, Vogel EK, Luck SJ (1998) Sensory gain control (amplification) as a mechanism of
selective attention: electrophysiological and neuroimaging evidence. Philosophical
transactions of the Royal Society of London Series B, Biological sciences 353:1257-
1270.
Hillyard SA, Hink RF, Schwent VL, Picton TW (1973) Electrical signs of selective attention in the
human brain. Science 182:177-180.
Huettig F, Altmann GT (2005) Word meaning and the control of eye fixation: semantic
competitor effects and the visual world paradigm. Cognition 96:B23-32.
Kisler T, Reichel UD, Sciel F (2017) Multilingual processing of speech via web services.
Computer Speech & Language 45:326-347.
Koenig T, Kottlow M, Stein M, Melie-Garcia L (2011) Ragu: a free tool for the analysis of EEG
and MEG event-related scalp field data using global randomization statistics. Comput
Intell Neurosci 2011:1-14.
Koenig T, Stein M, Grieder M, Kottlow M (2014) A tutorial on data-driven methods for
statistically assessing ERP topographies. Brain Topogr 27:72-83.
Lange K (2009) Brain correlates of early auditory processing are attenuated by expectations for
time and pitch. Brain and cognition 69:127.
Lee AKC, Rajaram S, Xia J, Bharadwaj H, Larson E, Hämäläinen MS, Shinn-Cunningham BG
(2013) Auditory Selective Attention Reveals Preparatory Activity in Different Cortical
Regions for Selection Based on Source Location and Source Pitch. Front Neurosci
Frontiers in Neuroscience 6.
Lehmann D (1989a) Microstates of the Brain in EEG and ERP Mapping Studies.72-83.
Lehmann D (1989b) From Mapping to the Analysis and Interpretation of EEG/EP Maps.53-75.
Liberman MC, Epstein MJ, Cleveland SS, Wang H, Maison SF (2016) Toward a differential
diagnosis of hidden hearing loss in humans. PLoS One 11:1-16.
Liebenthal E, Sabri M, Beardsley SA, Mangalathu-Arumana J, Desai A (2013) Neural dynamics
of phonological processing in the dorsal auditory stream. J Neurosci 33:15414-15424.
Luck SJ (2014) An introduction to the event-related potential technique, Second edition. Edition.
Cambridge, Massachusetts: The MIT Press.
Magnuson JS, Dixon JA, Tanenhaus MK, Aslin RN (2007) The dynamics of lexical competition
during spoken word recognition. Cognitive science 31:133-156.

McMurray B, Farris-Trimble A, Rigler H (2017) Waiting for lexical access: Cochlear implants or
severely degraded input lead listeners to process speech less incrementally. Cognition
169:147-164.
McQueen JM, Huettig F (2012) Changing only the probability that spoken words will be distorted
changes how they are recognized. The Journal of the Acoustical Society of America The
Journal of the Acoustical Society of America 131:509-517.
Mesgarani N, Chang EF (2012) Selective cortical representation of attended speaker in multi-
talker speech perception. Nature 485:233-236.
Myers EB, Blumstein SE, Walsh E, Eliassen J (2009) Inferior Frontal Regions Underlie the
Perception of Phonetic Category Invariance. PSCI Psychological Science 20:895-903.
Nabelek AK (1988) Identification of vowels in quiet, noise, and reverberation: relationships with
age and hearing loss. The Journal of the Acoustical Society of America 84:476-484.
Ohlenforst B, Zekveld AA, Lunner T, Wendt D, Naylor G, Wang Y, Versfeld NJ, Kramer SE
(2017) Impact of stimulus-related factors and hearing impairment on listening effort as
indicated by pupil dilation. Hear Res 351:68-79.
Ott CG, Langer N, Oechslin MS, Meyer M, Jancke L (2011) Processing of voiced and unvoiced
acoustic stimuli in musicians. Front Psychol 2:195.
Owens E, Schubert ED (1977) Development of the California Consonant Test. J Speech Lang
Hear Res 20:463-474.
Parbery-Clark A, Skoe E, Kraus N (2009) Musical experience limits the degradative effects of
background noise on the neural processing of sound. J Neurosci 29:14100-14107.
Pelli DG (1997) The VideoToolbox software for visual psychophysics: transforming numbers into
movies. Spat Vis 10:437-442.
Schiller B, Gianotti LR, Baumgartner T, Nash K, Koenig T, Knoch D (2016) Clocking the social
mind by identifying mental processes in the IAT with electrical neuroimaging. Proc Natl
Acad Sci U S A 113:2786-2791.
Schneider P, Scherg M, Dosch HG, Specht HJ, Gutschalk A, Rupp A (2002) Morphology of
Heschl's gyrus reflects enhanced activation in the auditory cortex of musicians. Nature
neuroscience 5:688-694.
Scott SK, Johnsrude IS (2003) The neuroanatomical and functional organization of speech
perception. Trends Neurosci 26:100-107.
Shinn-Cunningham BG, Best V (2008) Selective attention in normal and impaired hearing.
Trends Amplif 12:283-299.
Stahl J, Gibbons H (2004) The application of jackknife-based onset detection of lateralized
readiness potential in correlative approaches. Psychophysiology 41:845-860.
Storey JD (2002) A direct approach to false discovery rates. JOURNAL- ROYAL STATISTICAL
SOCIETY SERIES B STATISTICAL METHODOLOGY 64:479-498.
Tong Y, Chen Q, Nichols TE, Rasetti R, Callicott JH, Berman KF, Weinberger DR, Mattay VS
(2016) Seeking optimal region-of-interest (ROI) single-value summary measures for
fMRI studies in imaging genetics. PLoS One 11:1-20.
Vaden KI, Jr., Kuchinsky SE, Ahlstrom JB, Dubno JR, Eckert MA (2015) Cortical activity
predicts which older adults recognize speech in noise and when. J Neurosci 35:3929-
3937.
Wackermann J, Lehmann D, Michel CM, Strik WK (1993) Adaptive segmentation of
spontaneous EEG map series into spatially defined microstates. INTPSY</cja:jid>
International Journal of Psychophysiology 14:269-283.
Wong PC, Uppunda AK, Parrish TB, Dhar S (2008) Cortical mechanisms of speech perception
in noise. Journal of speech, language, and hearing research : JSLHR 51:1026-1041.
Wong PC, Jin JX, Gunasekera GM, Abel R, Lee ER, Dhar S (2009) Aging and cortical
mechanisms of speech perception in noise. Neuropsychologia 47:693-703.

Zekveld AA, Heslenfeld DJ, Festen JM, Schoonhoven R (2006) Top-down and bottom-up
processes in speech comprehension. Neuroimage 32:1826-1836.
728

Figure Legends

Figure 1. Trial and stimulus structure. Every trial starts with the cue phrase "check the word." A target word starts 1 second after the noise onset. Four choices are given after the word ends; subjects select the correct answer with a keypad. No feedback is given. The noise level is manipulated to create high (+3 dB) and low (-3 dB) SNR conditions. Subjects complete 50 trials for each condition.

Figure 2. Behavioral results. **A**. Summary of behavioral performance for the two conditions (+3 and -3 dB SNR). Boxes denote the 25th – 75th percentile range; the horizontal bars in the center denote the median; the ranges are indicated by vertical dashed lines. Solid lines connect points for the same subject in different conditions. **B**. Average accuracy as a function of reaction time in +3 dB SNR condition. **C**. Average accuracy and reaction time in -3 dB SNR condition.

Figure 3. Region-of-interest (ROI) based source analysis. **A**. Cortical labels for two ROIs in left and right hemispheres: supramarginal gyrus (SMG), and the pars opercularis and triangularis of the inferior frontal gyrus (IFG), respectively. **B**. The time course of the t-value envelope, with the standard error of the mean (±1 SEM), obtained at representative voxels in each SNR condition (red color: +3 dB SNR, blue color: -3 dB SNR). An asterisk shows the timing of significant difference between +3 and -3 dB SNR conditions (paired t-test, FDR adjusted p < 0.05). **C**. Whole brain maps showing statistical contrasts (t-values obtained from paired t-tests between the two SNR conditions) of source activation at each voxel, only displaying those with p-value less than 0.05, at the timepoint that shows significant differences over the broadest area in the ROIs within the time range described above.

Figure 4. Timings of significant cortical activity relative to distributions of phonological events. **A**. Top and second panel show a histogram of the onsets of second and final phoneme of each stimulus. The third panel shows superimposed temporal envelopes extracted from waveforms of the 100 words. **B**. The whole brain maps at the bottom are from **Figure 3C** that shows statistical contrasts of source activation at the timepoints that show significant differences between the two SNR conditions. Purple curves on the cortical maps represent the conceptual illustration of ascending information flow through the dorsal pathway.

Figure 5. Microstate analysis. **A**. Evoked responses over time after the word onset for +3 and -3 dB SNR condition. Each color represents ERPs from a different channel of interest. **B**. Global field power (GFP) is calculated at each timepoint that is assigned to one of the microstate clusters. **C**. Four microstate cluster maps. Dark blue, light blue, red, and dark pink colors represent microstates 1, 2, 3, and 4, respectively. The relative area of GFP is calculated and reveals the highest value for the microstate 1 and 2 for -3 and +3 dB SNR condition, respectively. **D**. Whole brain maps obtained at the times assigned to microstates 1 and 2, that show maximum GFP and the maximum peak of ERPs at the frontal-central electrodes.

Figure 6. Individual differences in speech-in-noise processing. **A**. Global field power of the grand mean evoked potentials after the noise onset and after the target word onset, separately in the low SNR condition. Scalp topographies were examined at the timepoints, suggested by microstate analysis from **Figure 5**, and compared between good and poor performers, as

determined by the median split. **B**. A series of scatter plots showing Pearson correlation
coefficients among internal SNR, early SMG, late IFG activation, and behavioral accuracy. **C**. A
scatter plot showing the regression coefficients from a linear regression model where behavioral
accuracy is the dependent variable while internal SNR, early SMG, late IFG activation are the
predictor variables. *The linear model significantly predicts behavioral accuracy while internal
SNR is the only significant predictor.

Supplement Figure 1. Region-of-interest (ROI) based source analysis. **A.** Cortical labels for Heschl's gyrus in left and right hemispheres. **B.** The time course of the t-value envelope, with the standard error of the mean (±1 SEM), obtained at representative voxels in each SNR condition (red color: +3 dB SNR, blue color: -3 dB SNR). An asterisk shows the timing of significant difference between +3 and -3 dB SNR conditions (paired t-test, FDR adjusted $p < 0.05$). **C.** Whole brain maps showing statistical contrasts (t-values obtained from paired t-tests between the two SNR conditions) of source activation at each voxel, only displaying those with p-value less than 0.05.