A *Francisella tularensis* L,D-carboxypeptidase plays important roles in cell morphology,

envelope integrity, and virulence.

Briana Zellner[1], Dominique Mengin-Lecreulx[2], Brenden Tully[1], William T. Gunning 3rd[3],

Robert Booth[3], Jason F. Huntley[1*]

[1] Department of Medical Microbiology and Immunology, University of Toledo, Toledo, OH,

U.S.A.; [2] Institute for Integrative Biology of the Cell (I2BC), CEA, CNRS, Univ. Paris-Sud,

Université Paris-Saclay, 91198 Gif-sur-Yvette, France; [3]Department of Pathology, University of

Toledo, Toledo, OH, U.S.A.

Running Head: *F. tularensis* L,D-carboxypeptidase

To whom correspondence should be addressed: Jason F. Huntley, Department of Medical

Microbiology and Immunology, University of Toledo College of Medicine and Life Sciences,

3000 Arlington Ave., MS1021, HEB 200D, Toledo, OH, U.S.A., 43614. Tel.: 419-383-5456;

Email: Jason.Huntley@UToledo.edu

Summary

Francisella tularensis is a Gram-negative, intracellular bacterium that causes the zoonotic disease tularemia. Intracellular pathogens, including *F. tularensis*, have evolved mechanisms to survive in the harsh environment of macrophages and neutrophils, where they are exposed to cell membrane-damaging molecules. The bacterial cell wall, primarily composed of peptidoglycan (PG), maintains cell morphology, structure, and membrane integrity. Intracellular Gram-negative bacteria protect themselves from macrophage and neutrophil killing by recycling and repairing damaged PG – a process that involves over 50 different PG synthesis and recycling enzymes. Here, we identified a PG recycling enzyme, L,D-carboxypeptidase A (LdcA), of *F. tularensis* that is responsible for converting PG tetrapeptide stems to tripeptide stems. Unlike *E. coli* LdcA and most other orthologs, *F. tularensis* LdcA does not localize to the cytoplasm and also exhibits L,D-endopeptidase activity, converting PG pentapeptide stems to tripeptide stems. Loss of *F. tularensis* LdcA led to altered cell morphology and membrane integrity, as well as attenuation in a mouse pulmonary infection model and in primary and immortalized macrophages. Finally, an *F. tularensis ldcA* mutant protected mice against virulent Type A *F. tularensis* SchuS4 pulmonary challenge.

Keywords: tularemia, peptidoglycan, L,D-carboxypeptidase, virulence, *Francisella tularensis*

Introduction

The Gram-negative bacterial cell wall plays an important role in maintaining cell shape, protecting against external insults, and preventing cell lysis amid fluctuations in internal turgor pressure (Dhar, 2018). The cell wall is composed of peptidoglycan (PG), a network of alternating *N*-acetylglucosamine (GlcNAc) and *N*-acetylmuramic acid (MurNAc) glycan chains that are crosslinked through peptide stems, and lies just outside of the cytoplasmic membrane of most bacteria (Johnson, 2013; Mengin-Lecreulx & Lemaitre, 2005). In *Escherichia coli*, PG has been shown to be covalently attached to the outer membrane (OM) by Braun's lipoprotein and noncovalently attached by Pal and other lipoproteins (Bouveret, 1999; Braun, 1975; Braun, 1969; Leduc, 1992). A loss of membrane integrity can occur if interactions between PG and attached lipoproteins are disturbed, thus maintenance of correct PG architecture is extremely important (Braun & Hantke, 2019; Braun & Rehn, 1969).

Synthesis of Gram-negative PG precursors, studied mainly in *E. coli*, occurs in the bacterial cytoplasm and requires a series of enzymes to build a pentapeptide PG monomer before transporting this molecule into the periplasm. Periplasmic PG cross-linking most often occurs between the fourth residue (D-Ala) of newly-formed pentapeptide stems to the third residue (*meso*-A_2pm) of existing tripeptide stems (4-3 cross links), resulting in the release of the pentapeptide terminal D-Ala and forming tetrapeptide stems (Glauner, 1988; Pazos & Peters, 2019). PG is not a static structure, rather, PG degradation/remodeling is necessary to incorporate new PG and expand the cell wall during bacterial growth and replication, to insert flagella or secretion systems, and to septate during bacterial division (Scheurwater, 2008). Up to 60% of PG is recycled per generation, helping to repair damaged PG and providing energy during periods of stress or starvation (Dhar, 2018; Holtje, 1998; Park, 2008). Using *E. coli* as a model, more than 50 different PG synthesis and hydrolysis/recycling enzymes have been identified, most of which appear to be

cytoplasmic. While deletion studies in *E. coli* have indicated that some PG synthesis enzymes are

essential, few PG hydrolases have been found to be essential and, in fact, substantial redundancy

in hydrolase activity appears to exist. Indeed, alterations in cell morphology, membrane integrity,

and ability to replicate/divide sometimes are observed only after multiple PG metabolism genes

have been deleted (Dhar, 2018; van Heijenoort, 2011; Vollmer & Bertsche, 2008). Given the

increasing threat of antimicrobial-resistant organisms and link between PG homeostasis and

bacterial virulence, more studies are needed to understand how a diverse range of Gram-negative

bacteria synthesize and recycle PG (Juan, 2018). As proof that PG studies can reveal new

information about bacterial pathogenesis and new treatment options, an *Acinetobacter baumannii*

penicillin-binding protein (PBP) mutant was reported to be more sensitive to complement-

mediated killing than wild-type bacteria (Russo, 2009) and a *Helicobacter pylori* PG hydrolase

(AmiA) mutant was unable to colonize mouse stomachs (Chaput, 2016).

PG recycling, mainly characterized in *E. coli,* begins with periplasmic lytic

transglycosylase (LT) cleavage of the β-1,4-glycosidic linkage between MurNAc and GlcNAc,

forming a GlcNAc-1,6-anhydro-MurNAc product, which allows for insertion of new muropeptides

and recycling of the GlcNAc-1,6-anhydro-MurNAc peptide (Scheurwater, 2008). Low molecular

mass penicillin binding proteins (LMM PBPs) can function as endopeptidases, cleaving the cross-

links between adjacent tetrapeptide stems, and/or as D,D-carboxypeptidases, removing the

terminal D-Ala of pentapeptides during transpeptidation (cross-linking) reactions, forming the

tetrapeptide (Dhar, 2018). Inner membrane permeases, such as AmpG, then transfer GlcNAc-1,6-

anhydro-MurNAc disaccharides, with or without attached peptides, to the cytoplasm where they

can be disassembled. Cytoplasmic NagZ and AmpD further degrade disaccharides by cleaving

the bond between GlcNAc and 1,6-anhydro-MurNAc, and separating the peptide chain from 1,6-

anhydro-MurNAc, respectively.   Additional cytoplasmic hydrolases, such as L,D-

carboxypeptidases (Ldc), act on free peptide chains to cleave the terminal D-Ala from tetrapeptides, resulting in tripeptides (Dhar, 2018). PG is unusual in that it is both highly dynamic (*e.g.,* allowing for bacterial division and molecular transport across the periplasm), yet tightly regulated to prevent membrane collapse and bacterial death. As such, PG recycling enzymes have been speculated to be important virulence determinants in Gram-negative bacteria (Juan, 2018). Indeed, *E.coli ldc* mutants lyse in stationary phase (Templin, 1999) and are more susceptible to β-lactam antibiotics (Ursinus, 1992), *Helicobacter* and *Campylobacter ldc* mutants have altered cell morphology and defects in motility and biofilm formation (Frirdich, 2012; 2014; Sycuro, 2013), and *Neisseria gonorrhoeae ldc* mutants are unable to stimulate NOD1-dependent responses in the host (Lenz, 2017). However, very little is known about the importance of PG recycling enzymes in the pathogenesis of intracellular pathogens such as *Francisella tularensis*.

       F. tularensis, the causative agent of tularemia, is a Gram-negative, intracellular, coccobacillus that can infect and cause lethal disease in many species, including humans (Dennis, 2001; Keim, 2007). There are three subspecies of *F. tularensis*, subsp. *tularensis* (Type A), subsp. *holarctica* (Type B), and subsp. *mediasiatica*, although only subsp. *tularensis* and subsp. *holarctica* are virulent for humans (Kingry, 2014). *F. tularensis* poses a severe threat to public health and has been classified as an NIH Category A Priority Pathogen and a CDC Tier 1 Select Agent due to its low infectious dose (<10 CFU), ease of aerosolization, and high morbidity and mortality rates (up to 60%) (Ellis, 2002; Sjostedt, 2007). Like other intracellular pathogens, *F. tularensis* has evolved different mechanisms to infect, survive, and replicate within host cells, including macrophages and neutrophils (Ray, 2009). However, this lifestyle exposes the bacteria to reactive oxygen species (ROS), reactive nitrogen species (RNS), antimicrobial peptides, and other cell membrane- and cell wall-damaging molecules (Jones, 2012). Our group previously demonstrated that the *F. tularensis* disulfide bond formation protein A (DsbA) ortholog repairs

damaged outer membrane proteins and known virulence factors. We additionally showed that *F.*

tularensis DsbA, unlike periplasmic DsbA in *E. coli* and most other Gram-negative bacteria, is

outer membrane-bound and is a multifunctional protein with both oxidoreductase and isomerase

activities. Finally, using a molecular trapping approach, we identified over 50 *F. tularensis* DsbA

substrates, many of which we speculate are involved in virulence (Ren, 2014).

Here, we determined the function of one of those *F. tularensis* DsbA substrates – a

previously unstudied hypothetical protein containing a putative LdcA domain – and assessed its

role in bacterial virulence. Deletion of *F. tularensis* LdcA resulted in bacteria with altered cell

morphology, increased sensitivity to β-lactam antibiotics, yet increased resistance to several

stressors (*e.g.,* H_2O_2, NaCl, low pH). Next, we demonstrated that *F. tularensis* LdcA exhibits L,D-

carboxypeptidase and L,D-endopeptidase activities on pentapeptide and tetrapeptide residues of

PG. Finally, we established that *F. tularensis* LdcA is required for virulence, as mutants were

unable to replicate in macrophages or cause disease in mice.

Results

FTL1678 contains a putative L,D-carboxypeptidase domain

Previous studies by our group and others have shown that *F. tularensis* DsbA mutants are

attenuated in mice (Qin, 2009; Ren, 2014). However, additional work by our group, demonstrating

that DsbA possesses both oxidoreductase and isomerase activities to repair damaged envelope and

cell membrane proteins, indicated that other envelope proteins likely are responsible for *F.*

tularensis virulence (Ren, 2014). To identify new *F. tularensis* virulence factors, we used a

molecular trapping approach and identified over 50 *F. tularensis* DsbA substrates (Ren, 2014).

One of those DsbA substrates, FTL1678, is annotated in the *F. tularensis* genome as a conserved

membrane hypothetical protein. Here, a conserved domain search revealed that a large portion of

FTL1678 contains a putative Ldc domain, part of the peptidase_S66 superfamily (Figure S1). Ldc

proteins have been studied in a number of Gram-negative bacteria, including *E. coli* (Metz, 1986a;

b; Templin, 1999; Ursinus, 1992), *Pseudomonas aeruginosa* (Korza, 2005), *N. gonorrhoeae* (Lenz,

2017), and *Campylobacter jejuni* (Frirdich, 2014).   To further explore this conserved domain,

amino acid sequences of FTL1678 (*F. tularensis* subsp. *holarctica* [Type B] LVS) and FTT0101

(homolog of FTL1678 in *F. tularensis* subsp. *tularensis* [Type A] SchuS4) were aligned with LdcA

orthologs from *E. coli*, *P. aeruginosa*, *N. gonorrhoeae*, and *C. jejuni* (named Pgp2). Despite low

percentages of amino acid identities among the LdcA orthologs (6.3% to 30.3%; Figure 1), there

was a higher degree of amino acid similarity among LdcA orthologs (13.0% [*E. coli* and *C. jejuni*]

to 44.7% [*E. coli* and *N. gonorrhoeae*]; Figure 1). Notably, the LdcA Ser-Glu-His catalytic triad,

previously shown to be required for *P. aeruginosa* LdcA activity (Korza, 2005), was absent from

C. jejuni Pgp2 but was present in all LdcA homologs, including FTL1678 and FTT0101 (Figure

1).

E. coli and *P. aeruginosa* Ldcs have been localized to the bacterial cytoplasm (Korza, 2005;

Templin, 1999). However, *C. jejuni* Pgp2 is unusual in that it contains a signal peptide and has

been speculated to be periplasmic (Frirdich, 2014). In addition, *N. gonorrhoeae* LdcA was found

to be periplasmic and outer membrane-associated (Lenz, 2017). As noted above, we previously

demonstrated that FTL1678 is a DsbA substrate (Ren, 2014), indicating that FTL1678 is located

in the *F. tularensis* envelope (*i.e.,* in the inner membrane [IM], periplasm, or outer membrane

[OM]). Bioinformatic analyses of FTL1678 indicated that it is a periplasmic protein due to the

presence of a signal peptide but absence of OM or lipoprotein signatures (Table S1).   To

experimentally confirm FTL1678 localization, we generated an *F. tularensis* strain with 6×

histidine-tagged FTL1678, then performed spheroplasting, osmotic lysis, and sucrose density

gradient centrifugation to separate IM and OM fractions and probe for protein subcellular

localization. Immunoblotting of whole-cell lysates (WCL), OM fractions, and IM fractions

demonstrated that the OM control protein, FopA (Huntley, 2007), only was present in WCL and

OM fractions (but not IM fractions; Figure 2) and the IM control protein, SecY (Huntley, 2007),

only was present in WCL and IM fractions (but not OM fractions; Figure 2). By comparison,

FTL1678 only was detected in WCL and OM fractions, demonstrating OM-association (Figure 2).

As noted above, because *N. nonorrhoeae* LdcA was found to fractionate to both the OM and

soluble fractions (indicating periplasmic localization) (Lenz, 2017), we next examined the

localization of *F. tularensis* periplasmic proteins in our fractions to better understand FTL1678

localization. TolB is a well-known periplasmic protein in Gram-negative bacteria and binds to PG

with the peptidoglycan associated lipoprotein, Pal (Clavel, 1998; Walburger, 2002). We

previously demonstrated that the *F. tularensis* Pal homolog is OM-localized (Huntley, 2007),

similar to its OM-localization in other Gram-negative bacteria. Here, the *F. tularensis* TolB

homolog was detected in OM fractions, but not IM fractions (Figure S2), demonstrating that some

F. tularensis periplasmic proteins fractionate with OMs. Additional fractionation and periplasmic

protein localization (Jones et al, 2016) experiments were performed but specific localization of

FTL1678 to the periplasm could not be confirmed (data not shown). Regardless, data that

FTL1678 is a DsbA substrate (*i.e.,* FTL1678 is an envelope protein)(Ren, 2014), FTL1678

contains a signal peptidase I cleavage site (Table S1), FTL1678 does not contain membrane protein

signatures (Table S1), and FTL1678 is OM-associated (Figure 2), provides strong evidence that

FTL1678 is not a cytoplasmic protein, unlike *E. coli* LdcA. Instead, our data indicate that, similar

to *N. nonorrhoeae* LdcA, FTL1678 may be a periplasmic protein.

FTL1678 exhibits L,D-carboxypeptidase and L,D-endopeptidase activities

To confirm the predicted Ldc activity of FTL1678 and FTT0101, recombinant FTL1678

and FTT0101 were expressed and affinity purified from *E. coli*. As a control, lysates from *E. coli*

containing the empty vector (pPROEX HTb) also were affinity-purified . Recombinant FTL1678,

lysate from the vector control, or buffer alone were incubated with various PG precursors and PG

intermediates (Table 1) to determine substrate specificity and specific activity. The vector control

and buffer alone did not demonstrate activity against any of the PG substrates (data not shown).

When FTL1678 was incubated with various PG substrates, the highest specific activity was

detected against the tetrapeptide substrates GlcNAc-anhydroMurNAc-L-Ala-γ-D-Glu-*meso*-

A$_2$pm-D-Ala (tracheal cytotoxin; TCT; 21.5 nmol/min/mg of protein) and GlcNAc-MurNAc-L-

Ala-γ-D-Glu-*meso*-A$_2$pm-D-Ala (reducing PG monomer; 15.6 nmol/min/mg of protein),

confirming that FTL1678 exhibits L,D-carboxypeptidase activity (Table 1). Interestingly,

FTL1678 activity against free tetrapeptide, L-Ala-γ-D-Glu-*meso*-A$_2$pm-D-Ala, was approx. 6-fold

lower (3.4 nmol/min/mg of protein) than TCT and 5-fold lower than the reducing PG monomer

(Table 1), indicating that GlcNAc and MurNAc may be important for tetrapeptide recognition or

FTL1678 binding. Importantly, FTL1678 exhibited specific activity against pentapeptide

substrates MurNAc-L-Ala-γ-D-Glu-*meso*-A$_2$pm-D-Ala-D-Ala (9.8 nmol/min/mg of protein;

Table 1) and UDP-MurNAc-L-Ala-γ-D-Glu-*meso*-A$_2$pm-D-Ala-D-Ala (5.9 nmol/min/mg of

protein; Table 1), indicating that FTL1678 also functions as an L,D-endopeptidase (cleavage of

the pentapeptide between *meso*-A$_2$pm and D-Ala) by FTL1678.

To further investigate the endopeptidase activity of FTL1678, FTL1678 was incubated

with various TCT monomer and dimer substrates, containing different peptide lengths and cross-

link locations (Table S2). Although FTL1678 exhibited the highest specific activity against TCT

monomers containing tetrapeptide stems with either an Alanine or a Glycine in the fourth position

(20 nmol/min/mg and 18 nmol/min/mg of protein, respectively), FTL1678 also was able to cleave

all four variations of the TCT dimer (two cross-linked TCT monomers), demonstrating

endopeptidase activity (Table S2). Of the four different TCT dimers tested, FTL1678 was most

active on TCT dimers connected by a 4-3 cross linkage (6.6 nmol/min/mg of protein), followed by

an approx. 24-fold reduction in activity on 3-3 cross links connecting two tripeptides (0.28

nmol/min/mg of protein), a tripeptide and tetrapeptide with a Glycine at the fourth position (0.21

nmol/min/mg of protein), and a tripeptide and tetrapeptide with an Alanine at the fourth position

(0.16 nmol/min/mg of protein) (Table S2). These data suggest that, while FTL1678 exhibits

endopeptidase activity on 4-3 and 3-3 cross-links, cleavage of 3-3 cross links is likely not its main

physiological function.

Finally, FTL1678 had negligible activity against L-lysine-containing substrates (0.7 to 1.3

nmol/min/mg of protein; Table 1), where L-lysine replaced *meso*-A$_2$pm at the third amino acid

position, indicating the importance of *meso*-A$_2$pm. Assays were repeated with recombinant

FTT0101 (SchuS4 homolog) and, due to 99.4% amino acid identify with FTL1678, FTT0101

demonstrated similar tetrapeptide cleavage activity (*i.e.,* LdcA activity) as FTL1678 (Table 1).

FTT0101 also was not active on a peptidoglycan polymer and had either no or negligible activity

on PG monomers that were amidated at the *meso*-A$_2$pm or D-Glu residues (Table 1). Additionally,

FTT0101 was not inhibited by 5 mM EDTA and did not require the presence of cations (Mg^{2+}) for

tetrapeptide cleavage (data not shown).

FTL1678 controls bacterial morphology

LdcA has been shown to be important for maintenance of bacterial morphology and

structural integrity (Frirdich, 2014; Sycuro, 2013).   In addition, mutations/deletions or

combinations of mutations/deletions in PG-modifying proteins can result in abnormal bacterial

morphology, emphasizing the importance of PG modification and recycling (Guinane, 2006;

Heidrich, 2001; Juan, 2018; Nelson, 2000; Priyadarshini, 2007; Sycuro, 2010). To assess if

FTL1678 plays a similar role in *F. tularensis*, we generated an isogenic deletion of *FTL1678*,

referred to hereafter as Δ*FTL1678*, in *F. tularensis* LVS. When examined by transmission electron

microscopy (TEM), wild-type (WT) bacterial width ranged from 350 to 800 nm (Figures 3A and

3D), whereas Δ*FTL1678* bacteria were more uniform in cell width, averaging approx. 350 nm

(Figures 3B and 3D). WT bacteria were observed to be coccobacilli with loosely-associated OM

(Figure 3A), while Δ*FTL1678* bacteria were found to be more coccoid in appearance, the OM was

tightly-associated, and three prominent structures were present around the periphery of each

bacterium, likely the OM, PG, and IM (Figure 3B). Additionally, Δ*FTL1678* bacteria appeared

more electron dense and had a significantly-thicker OM than WT bacteria (Figures 3A, 3B, and

3C).

251   Previous studies have shown that deletion of genes for PG-modifying proteins (*e.g.,* murein

hydrolases) can result in abnormal growth characteristics, including lysis during stationary phase

(Templin, 1999) and an inability to separate daughter cells at the septa during cell division,

resulting in abnormal bacterial chains (Chaput, 2016; Denome, 1999; Heidrich, 2001; 2002; Juan,

2018; Priyadarshini, 2007; 2006). Although *N*-acetylmuramyl-L-alanine amidases have been

shown to be predominantly involved in the cleavage of bacterial septa, deletion of lytic

transglycosylases and some endopeptidases, in combination with amidase deletions, also have

resulted in abnormal bacterial chains (Heidrich, 2001; 2002). To examine any potential replication

defects of Δ*FTL1678*, we compared both OD_{600} values (Figure S3A) and CFUs over time (Figure

S3B) of WT and Δ*FTL1678* in supplemented Mueller Hinton Broth (sMHB; standard growth

medium for *F. tularensis*; (Huntley, 2007)), finding that Δ*FTL1678* did not have any inherent

growth defects. When examining both WT and Δ*FTL1678* by TEM for any septation defects or

abnormal bacterial chains, approximately 10% of Δ*FTL1678* bacteria grew in chains of three to

four bacteria (Figure S4A and S4B), whereas no WT bacteria exhibited this septation defect (data

not shown).   Taken together, our findings that Δ*FTL1678* is 1.5- to 2-times smaller than WT

(Figure 3D), Δ*FTL1678* is more coccoid in shape (Fig 3B), Δ*FTL1678* has a pronounced three-

layered envelope, including a prominent electron dense layer between the IM and OM (likely PG;

Figure 3B), and Δ*FTL1678* has a partial septum defect (Figure S4), further support the role of

FTL1678 as a PG-modifying enzyme that is important for bacterial elongation and division.

Deletion of FTL1678 affects antibiotic, detergent, and stressor sensitivity

Given the above noted morphological differences in the Δ*FTL1678* envelope (*e.g.,* thicker

OM; tightly-associated OM; three prominent envelope structures) using TEM*,* we assessed

envelope (*i.e.,* IM, PG, and OM) integrity by growing both WT and Δ*FTL1678* bacteria in the

presence of various antibiotics, detergents, and dyes, and measuring zones of inhibition after 48 h

of growth (Table 2).   Δ*FTL1678* was found to be more susceptible than WT to ampicillin,

vancomycin, lysozyme, and SDS (Table 2), indicating potential changes to PG (ampicillin

sensitivity), OM integrity (vancomycin and lysozyme sensitivity), or efflux pumps (SDS

sensitivity).   Conversely, Δ*FTL1678* was found to be more resistant than WT to gentamicin,

tetracycline, chloramphenicol, ciprofloxacin, ethidium bromide, and Triton X-100 (Table 2).

Given that the majority of these latter reagents must enter the cytoplasm to exert their inhibitory

effects (*i.e.,* gentamicin, tetracycline, and chloramphenicol inhibit protein synthesis; ciprofloxacin

and ethidium bromide interfere with DNA replication), these results suggest that Δ*FTL1678*

bacteria exclude these inhibitory molecules from entering the cytoplasm.

To better understand potential differences in the Δ*FTL1678* envelope, WT and Δ*FTL1678*

were grown in either sMHB at 37°C or in sMHB with various stress conditions. In sMHB at 37°C,

Δ*FTL1678* did not exhibit a growth defect but, instead, appeared to grow to a higher optical density

(OD_{600}) than WT (Figure 4A). However, as noted above, despite higher OD_{600} measurements for

Δ*FTL1678* at several time points, bacterial numbers were not significantly different between WT

and Δ*FTL1678* (Figure S3). Although speculative, the disassociation between Δ*FTL1678* optical

densities and bacterial numbers may be due to the observed TEM morphological differences of

Δ*FTL1678* (Figure 3 and Figure S4). Compared with growth in sMHB at 37°C, no substantial

differences in the growth rates of WT and Δ*FTL1678* were observed at either 40°C or in the

presence of 60 μM $CuCl_2$ (Figure 4B and 4C). However, Δ*FTL1678* grew considerably better than

WT in the presence of 5 mM H_2O_2, 5% NaCl, and pH 5.5 (Figure 4D, 4E, and 4F, respectively),

providing further evidence of modifications to the Δ*FTL1678* envelope. Although many

mechanisms can account for increased resistance to H_2O_2, high NaCl, and low pH, the combined

results from these antibiotic, detergent, and stressor sensitivity/resistance assays suggest that

Δ*FTL1678* bacteria may have decreased OM integrity, altered PG, altered activity of efflux pumps,

and a less permeable IM. Indeed, decreased permeability of the IM and altered activity of IM

efflux pumps may help explain the more electron-dense staining of Δ*FTL1678*, compared to WT

(Figure 3).

Δ*FTL1678 is fully-attenuated in a mouse pulmonary infection model*

To examine if FTL1678 plays a role in *F. tularensis* virulence, mice were intranasally

infected with 10^4 CFU of either WT or Δ*FTL1678* and monitored daily for signs of disease.

Whereas all WT-infected mice died by day 9 post-infection (median time-to-death day 6),

Δ*FTL1678* was completely attenuated (100% survival through day 21 post-infection),

demonstrating that FTL1678 is required for *F. tularensis* virulence (Figure 5A). To confirm that

the observed attenuation was solely due to the deletion of *FTL1678*, and not to polar effects, we

complemented $\Delta FTL1678$ with a 6× His-tagged FTL1678 *in trans*, which fully-restored virulence

to WT levels (all mice died by day 7; median time-to-death day 6; Figure 5A).

To more carefully assess $\Delta FTL1678$ attenuation *in vivo*, we intranasally-infected mice with

10^4 CFU of either WT LVS or $\Delta FTL1678$, and enumerated bacterial CFUs from lungs, livers,

spleens, and blood on days 2 and 5 post-infection to examine bacterial replication and

dissemination to these organs/tissues over time. On day 2 post-infection, WT LVS replicated to

$>10^7$ CFU/mg lung and had disseminated to livers, spleens (approx. 10^3 CFU/mg), and blood (10^3

CFU/ml; Figure 5B). In contrast, $\Delta FTL1678$ had an initial (day 2) colonization defect in the lungs

(4-logs less than WT) and was unable to disseminate to livers, spleens, or blood (Figure 5B). By

day 5 post-infection, the attenuation of $\Delta FTL1678$ was even more apparent, with WT LVS

replicating to extremely high numbers (approx. 10^8 CFU/mg) in lungs, livers, and spleens,

compared with $\Delta FTL1678$, which replicated approx. 1-log in lungs (between day 2 and 5), but was

4-log attenuated in lungs and was not detectable in livers or spleens (Figure 5B). Although

$\Delta FTL1678$ was detected in the blood on day 5, it was 2-logs less than WT LVS (Figure 5B).

The Type A *F. tularensis* strain SchuS4 originally was isolated from a human tularemia

patient and requires BSL3 containment. Given its relevance to human disease, we next generated

an isogenic deletion mutant of the *FTL1678* homolog, *FTT0101*, in SchuS4. When mice were

intranasally-infected with either WT SchuS4 or $\Delta FTT0101$, all mice died by day 7 post-infection,

indicating that *FTT0101* is not required for SchuS4 virulence (Figure S5). Whereas $\Delta FTT0101$-

infected mice exhibited a slightly delayed time-to-death (median time-to-death day 6; Figure S5),

compared with WT SchuS4-infected mice (median time-to-death day 5; Figure S5), this may be

due to differences in the infectious dose administered to mice in this experiment (80 CFU SchuS4;

12 CFU $\Delta FTT0101$). However, it also remains possible that the extreme virulence of SchuS4

(intranasal LD_{50} <10 CFU in our hands) complicates assessments of mutant attenuation *in vivo*.

To further assess Δ*FTT0101*, the susceptibilities of Δ*FTT0101* and WT SchuS4 to various

antibiotics, detergents, and dyes were compared, with no significant differences observed (Table

S3). At this time, we cannot fully explain why the *F. tularensis* Type B LdcA mutant Δ*FTL1678*

is fully-attenuated in mice (Figure 5) and displays altered sensitivity/resistance to antibiotics,

detergents, and dyes (Table 2), while the *F. tularensis* Type A LdcA mutant Δ*FTT0101* retains

virulence in mice (Figure S5) and does not demonstrate altered sensitivity/resistance to antibiotics,

detergents, and dyes (compared to WT SchuS4; Table S3). However, this finding is not

unexpected given genomic studies indicating that, despite >97% nucleotide identity between Type

A and Type B *F. tularensis*, there are over 100 genomic rearrangements between Type A and Type

B and each subspecies encodes over 100 unique genes that likely influence known differences in

Type A and Type B virulence (Petrosino et al, 2006).

Putative FTL1678 catalytic triad and Ldc activity are required for F. tularensis virulence

As noted above and highlighted in Figure 1, *P. aeruginosa* LdcA contains a Ser-Glu-His

catalytic triad which is essential for function and is characteristic of Ldc in the Peptidase_S66

family (Korza, 2005). The Ser-Glu-His catalytic triad also has been confirmed in Ldc from *E. coli*

(Meyer, 2018), *Novosphingobium aromaticivorans* (Das, 2013), and *N. gonorrhoeae* (Lenz, 2017).

Given the relatively conserved spacing of Ser134-Glu239-His308 residues in FTL1678 (Figure 1),

we tested if these residues were required for *F. tularensis* virulence (similar to Figure 5A virulence

assessments for Δ*FTL1678* and the FTL1678 complemented strain). Site-directed mutagenesis

was performed to independently generate FTL1678 complementation constructs containing either

S134A, E239A, or H308A mutations. Next, Δ*FTL1678* was complemented *in-trans* with each of

these FTL1678 catalytic triad point mutants, and mice were intranasally infected with either WT,

Δ*FTL1678*, Δ*FTL1678 trans*-complemented with *FTL1678*, or Δ*FTL1678 trans*-complemented

with one of the FTL1678 catalytic triad point mutants (referred to hereafter as S134A, E239A, and

H308A). Confirming our previous findings, Δ*FTL1678* was completely attenuated (100% survival

through day 21), while complementation of Δ*FTL1678* with either FTL1678 (all mice dead by day

8), S134A (all mice dead by day 7), E239A (all mice dead by day 8), or H308A (all mice dead by

day 8), fully-restored virulence to WT LVS levels (all mice dead by day 10; Figure S6). These

results indicated that although a putative Ser-Glu-His catalytic triad is present in FTL1678,

mutation of individual residues does not affect *F. tularensis* virulence *in vivo*.

Given the complexity of *in vivo* animal infections and the possibility that single mutations

in the putative Ser134-Glu239-His308 catalytic triad may not be sufficient to abolish enzyme

function or alter bacterial virulence, we next assessed if mutations of two or three residues in the

putative catalytic triad affected *in vitro* enzyme activity. Recombinant FTL1678 mutant proteins

were generated and purified, each containing either two amino acid mutations (S134A/E239A,

S134A/H308A, and E239A/H308A) or three amino acid mutations (S134A/E239A/H308A).

Similar to what is described above, enzymatic assays were performed, using the TCT monomer as

a substrate. While no specific activity to TCT monomer was detected for the empty vector control

or buffer alone, WT FTL1678 was active against the TCT monomer (11.0 nmols/min/mg). By

comparison, no activity was detected for any of the double or triple mutant proteins (Table S4),

suggesting that two or more of the catalytic triad residues are required for FTL1678 Ldc activity.

The Ldc ortholog alignment (Figure 1) highlighted that, of the six Ldc orthologs examined

here, only *C. jejuni* Pgp2 lacked the Ser-Glu-His catalytic triad. However, *C. jejuni* Pgp2 has been

shown to exhibit LdcA activity (Frirdich, 2014), indicating that a Ser-Glu-His catalytic triad is not

required for LdcA function. To test if an LdcA ortholog, without the Ser-Glu-His catalytic triad,

could restore virulence in the Δ*FTL1678* mutant, we complemented Δ*FTL1678* with a 6× His-

tagged Pgp2 from *C. jejuni* and infected mice with either WT LVS, Δ*FTL1678*, Δ*FTL1678 trans-*

complemented with FTL1678, or Δ*FTL1678 trans*-complemented with *C. jejuni* Pgp2. While

Δ*FTL1678* was fully attenuated (100% survival through day 21), the Pgp2 *trans*-complement was

fully-virulent (median time-to-death 6 days; all mice dead by day 7), nearly identical to WT LVS

(median time-to-death 7 days; all mice dead by day 7) and the FTL1678 *trans*-complement

(median time-to-death 6 days; all mice dead by day7; Figure 5C). This *in vivo* data provides further

evidence that FTL1678 is an Ldc and Ldc activity is required for *F. tularensis* virulence.

To provide additional evidence that FTL1678 exhibits Ldc activity and that *C. jejuni* Pgp2

functionally complements Δ*FTL1678*, the FTL1678 *trans*-complement and Pgp2 *trans*-

complement were examined by TEM, revealing that both complemented strains had similar

morphology as WT LVS (Figure S7A and S7B; Figure 3A). Additionally, OM thickness and cell

width were measured for WT, Δ*FTL1678*, and both complemented strains, demonstrating that both

complemented strains had OM thicknesses and cell widths similar to WT, and both complemented

strains were significantly different from Δ*FTL1678* (Figure S7C and S7D). Finally, when grown

in the presence of various stressors (*e.g.,* 5 mM H_2O_2, 5% NaCl, pH 5.5, antibiotics, SDS,

lysozyme, or ethidium bromide), both complemented strains exhibited similar phenotypes as WT

LVS (Figure S8A-D and Table S5). Taken together, these studies indicated that at least two

residues of the putative Ser134-Glu239-His308 catalytic triad are required for FTL1678 LdcA

enzymatic activity. These studies also demonstrated that an LdcA ortholog can functionally

complement Δ*FTL1678,* further supporting that FTL1678 is an LdcA ortholog.

FTL1678 is required for F. tularensis replication in macrophages

F. tularensis is an intracellular pathogen and macrophages appear to be one of the major

targets for *F. tularensis* infection and replication (De Pascalis, 2018; Hall, 2008; Steiner, 2017).

To investigate potential replication defects of Δ*FTL1678* in macrophages, J774A.1 macrophages

or murine bone marrow-derived macrophages (mBMDM) were infected with either WT LVS or

Δ*FTL1678* (MOI 100:1) and bacterial numbers were enumerated at 0 h (entry), 6 h, and 24 h post-

infection. At entry (0 h), approx. 2.5-logs more Δ*FTL1678* were present in both macrophage lines,

compared with WT LVS (Figure 6A). This likely was due to the above noted gentamicin resistance

of Δ*FTL1678* (Table 2). Attempts to normalize entry numbers for both WT LVS and Δ*FTL1678*,

using different antibiotics or combinations of antibiotics, were not successful. Despite higher

numbers of Δ*FTL1678* in both macrophages at entry (0 h) and 6 h, Δ*FTL1678* was unable to

replicate in either macrophage, and decreased approx. 1-log from 6 h to 24 h (Figure 6A). By

comparison, WT LVS numbers increased 1.5- to 2-logs from 6 h to 24 h (Figure 6A). To normalize

WT LVS and Δ*FTL1678* bacterial numbers and replication rates in both macrophages, fold change

in bacterial numbers was calculated from 6 h to 24 h: WT was found to increase 2.5- and 1.5-logs

in J774A.1 and mBMDMs, respectively, whereas Δ*FTL1678* was found to decrease 0.5- to 1-log,

respectively (Figure 6B). Taken together, these *in vitro* results (Figures 6A and 6B) confirm the

observed *in vivo* attenuation of Δ*FTL1678* (Figures 5A and 5B).

ΔFTL1678 protects mice against Type A F. tularensis infection

No FDA-approved vaccine currently is available to prevent tularemia. In addition, *F.*

tularensis is designated as an NIH Category A priority pathogen and CDC Tier 1 Select Agent,

highlighting the extreme virulence of this bacterium and the need for a safe and effective vaccine

to prevent tularemia. Given our above findings that 10^5 CFU of Δ*FTL1678* did not cause disease

or death in mice (Figures 5A and 5C), we next examined whether high doses (10^7 or 10^9 CFU) of

Δ*FTL1678* were attenuated or if Δ*FTL1678* immunization could protect mice from fully-virulent

Type A *F. tularensis* SchuS4 challenge. First, all mice intranasally immunized with either 10^5,

10^7, or 10^9 CFU of Δ*FTL1678* survived through day 28 post-infection, with no signs of clinical

disease (Figure 7A). Next, on day 29, all mice were boosted with 10^9 CFU of $\Delta FTL1678$ and no

mice demonstrated any signs of disease through day 50 (Figure 7A). Finally, on day 51, mice

were intranasally-challenged with 120 CFU ($6\times$ the LD$_{50}$) of SchuS4 and the health status of each

immunization group was monitored for 26 days post-challenge. In a dose-dependent manner, the

10^9 prime-10^9 boost regimen conferred 80% protection, the 10^7 prime-10^9 boost regimen conferred

40% protection, and the 10^5 prime-10^9 boost regimen conferred 20% protection (Figure 7B).

These data demonstrate that $\Delta FTL1678$ is highly attenuated (up to 10^9 CFU) and that $\Delta FTL1678$

may be able to be used as a live, attenuated vaccine.

ΔFTL1678 does not cause tissue damage

The *in vitro* (Figure 6) and *in vivo* (Figures 5A, 5B, 7A) attenuation of $\Delta FTL1678,$ as well

as protection against SchuS4 pulmonary challenge (Figure 7B), indicated that $\Delta FTL1678$ could be

used as a live, attenuated vaccine. While live, attenuated vaccines have been extremely effective

at preventing a number of diseases, they can pose safety challenges (Minor, 2015; Roberts, 2018).

To assess whether $\Delta FTL1678$ immunization induced any pathology in immunized mice, lungs,

livers, and spleens from uninfected, WT LVS-, or $\Delta FTL1678$-infected mice were assessed for

pathologic changes on day 5 post-infection/immunization. Day 5 is when mice exhibit severe

signs of disease and is one day before the majority of WT-infected mice begin succumbing to

disease (Figures 5 and 7). WT LVS-infected lungs demonstrated alveolar wall thickening, large

areas of inflammation, and severe neutrophil infiltration (Figure 8A). By comparison, little

inflammation was observed in $\Delta FTL1678$-infected lungs, although some red blood cell congestion

was present, indicating a limited, acute immune response that was quickly resolved (Figure 8A).

Whereas WT LVS-infected livers were characterized by diffuse inflammation with focal areas of

necrosis, $\Delta FTL1678$-infected livers were virtually indistinguishable from uninfected livers, with

no observable pathology (Figure 8A). Finally, although the architecture of WT LVS-infected

spleens lacked distinct areas of white pulp or red pulp, indicative of a severe infection, Δ*FTL1678*-

infected spleens were observed to contain distinct areas of red pulp and white pulp, with some red

blood cell congestion – indicating a limited, acute immune response that was quickly resolved

(Figure 8A). All tissues were blindly scored using a pathology severity index (scale from 0 to 4,

with 4 indicating severe pathology), confirming that Δ*FTL1678*-infected tissues were virtually

indistinguishable from uninfected tissues (pathology scores of 1 for lungs, 0 for liver, and 1.5 for

spleens) and WT LVS-infected tissues had significantly higher pathology scores (pathology scores

>3.5 for all tissues; Figure 8B).

Discussion

466   Bacterial PG is a complex, mesh-like structure, composed of a glycan backbone,

crosslinked to varying degrees, by peptide chains (de Pedro, 2015). It is well known that this

structure plays an important role in maintaining Gram-negative bacterial cell morphology,

membrane integrity, regulating changes in osmotic pressure, and providing a platform for

attachment of the OM (den Blaauwen, 2008; Silhavy et al, 2010). Although a majority of PG

studies have focused on how the thick layer of PG in Gram-positive bacteria contributes to

virulence and antibiotic resistance, more recent studies have highlighted that Gram-negative PG

also is intimately linked with pathogenicity (Juan, 2018).

474   PG recycling is an essential function of Gram-negative bacteria during cell growth and

division to produce new cell wall components. In fact, Gram-negative bacteria recycle up to 60%

of their PG with every generation, suggesting that both PG synthesis and PG recycling are dynamic

(Dhar, 2018; Typas, 2011). A number of proteins are involved in these processes and, while they

are well-characterized in *E. coli*, very little is known about these pathways in intracellular

pathogens such as *Burkholderia pseudomallei*, *Legionella pneumophila*, or *F. tularensis* (Jenkins,

2019; Kijek, 2019; Spidlova, 2018; van Heijenoort, 2011).

E. coli LdcA, a cytoplasmic protein, was the first L,D-carboxypeptidase to be identified

and was shown to be important for PG recycling and survival during stationary phase (Templin,

1999; Ursinus, 1992). More recently, Ldc orthologs have been identified in *P. aeruginosa* (Korza,

2005), *C. jejuni* (Frirdich, 2014), *N. gonorrhoeae* (Lenz et al, 2017), and *N. aromaticivorans* (Das,

2013). In this study, we identified an *F. tularensis* Ldc ortholog, FTL1678, which we propose

naming LdcA based on its confirmed L,D, carboxypeptidase activity (Table 1) and role in

maintaining bacterial morphology. Unlike well-characterized cytoplasmic LdcA orthologs from

E. coli and *P. aeruginosa*, we demonstrated that *F. tularensis* LdcA was localized to OM fractions

and, given co-localization with PG-associated proteins Pal and TolB, is most likely located on the

inner leaflet of the OM or in the periplasm (associated with PG). At this time, we can only

speculate on the OM-association or periplasmic localization of *F. tularensis* LdcA, but in the

context of PG repair and recycling, periplasmic LdcA certainly offers a fitness advantage. In

addition, this is not the first report of a periplasmic LdcA, as *C. jejuni* Pgp2 is predicted to be

periplasmic and *N. gonorrhoeae* LdcA previously was reported to be periplasmic (Frirdich, 2014;

Lenz, 2017).

Our results demonstrated, for the first time, that *F. tularensis* LdcA directly acts on the

TCT tetrapeptide and the reducing PG monomer. More importantly, we demonstrated that *F.*

tularensis LdcA directly cleaves PG pentapeptides to tripeptides, without a prior cleavage event

by a D,D-carboxypeptidase/penicillin binding protein (PBP), such as DacD, and that FTL1678

cleaves TCT dimers with 4-3 and 3-3 cross links, highlighting that *F. tularensis* LdcA is a multi-

functional enzyme that performs both L,D-carboxypeptidase and L,D- and D,D-endopeptidase

activities (Table 1 and Table S2). . It should be noted that although *N. gonorrhoeae* LdcA has

been reported to have L,D-endopeptidase activity, this activity is specific to tetrapeptide cross links

(3-3 cross links) (Lenz, 2017). Interestingly, only two previous studies have examined putative *F.*

tularensis PG modifying enzymes and both studies primarily focused on the role of an *F. tularensis*

DacD ortholog in virulence, with no PG activity assays to confirm function (Kijek, 2019; Spidlova,

2018). In our PG cleavage analysis, *F. tularensis* LdcA demonstrated the highest specific activity

on disaccharide-tetrapeptide PG substrates (GlcNAc-anhydroMurNAc-L-Ala-γ-D-Glu-*meso*-

A_2pm-D-Ala [TCT] and GlcNAc-MurNAc-L-Ala-γ-D-Glu-*meso*-A_2pm-D-Ala [reducing PG

monomer]), followed by cleavage of pentapeptide PG substrates (MurNAc-L-Ala-γ-D-Glu-*meso*-

A_2pm-D-Ala-D-Ala and UDP-MurNAc-L-Ala-γ-D-Glu-*meso*-A_2pm-D-Ala-D-Ala). Despite high

specific activity of *F. tularensis* LdcA on tetrapeptide attached to the disaccharide, *F. tularensis*

LdcA demonstrated approximately 6-times lower specific activity on free tetrapeptide (no sugars)

(Table 1). In contrast, *E. coli* LdcA has been shown to have the highest specific activity on free

tetrapeptide, monosaccharide-tetrapeptide (MurNAc-L-Ala-γ-D-Glu-*meso*-A_2pm-D-Ala), and

monosaccharide tetrapeptide linked to a glycan lipid carrier (UDP-MurNAc-tetrapeptide)

(Templin, 1999), but is unable to cleave dimeric muropeptides. Additionally, *F. tularensis* LdcA

was active against TCT dimers (two crosslinked TCT monomers), cleaving the 4-3 cross linkage

between D-Ala and A_2pm and the 3-3 cross linkage between A_2pm and A_2pm, but was not active

on PG polymers. This TCT dimer cleavage indicates that, in addition to L,D carboxypeptidase

activity, *F. tularensis* LdcA possesses L,D- and D,D-endopeptidase activity on peptide cross-links.

Previous studies have shown that Ldc orthologs are important for bacterial morphology

and membrane integrity. Deletion of the *ldc* orthologs *csd6* from *H. pylori* (Sycuro, 2013) and

pgp2 from *C. jejuni* (Frirdich, 2014) resulted in loss of helical morphology. Here, we demonstrated

that FTL1678 is essential for maintaining both the size (width) and the coccobacillus morphology

of *F. tularensis,* as Δ*FTL1678* were significantly-smaller than WT and exhibited a more-rounded,

cocci shape than WT (Figure 3). Further evidence for the role of *F. tularensis* LdcA in modifying

and recycling PG, which impacts bacterialmorphology, is provided by TEM images of Δ*FTL1678*

bacteria that had prominent three-layered structures at their periphery, including a thick middle

layer (presumably PG), compared to WT (Figure 3). In Δ*FTL1678* bacteria, it is possible that loss

of LdcA activity may have reduced PG recycling or may have affected the breakdown of existing

PG (important for cell division), resulting in a buildup of pentapeptides or tetrapeptides that are

highly-crosslinked. Our lab and others have repeatedly attempted to isolate and analyze *F.*

tularensis PG but theses attempts have not been unsuccessful (data not shown; Martin Pavelka, U.

Rochester, personal communication). As such, we can only speculate on the true nature of the

thick PG and OM layers in Δ*FTL1678* (Figure 3).

Because Δ*FTL1678* bacteria were found to have a thicker OM (Figure 3C), a prominent

middle layer in their envelope (presumably PG; Figure 3B), and altered cell morphology (Figure

3B), we investigated differences in WT and Δ*FTL1678* susceptibility to various antibiotics,

detergents, and stressors. Vancomycin and lysozyme, usually not effective against Gram-

negatives due to their inability to penetrate the OM, inhibited Δ*FTL1678* growth (Table 2),

indicating increased permeability of the Δ*FTL1678* OM. Vancomycin, in particular, may have

been effective on Δ*FTL1678* because its mechanism of action includes binding to the two terminal

D-Ala-D-Ala residues of PG pentapeptide chains and preventing cross-linking of monomers.

Δ*FTL1678* may have increased amounts of pentapeptides present in its PG, providing more targets

for vancomycin action. Similarly, ampicillin inhibits bacterial transpeptidases, which blocks

cross-linking of peptide side chains of PG strands. Taken together, the enhanced susceptibility of

Δ*FTL1678* bacteria to vancomycin and ampicillin supports the role of *F. tularensis* LdcA as a PG-

modifying enzyme. Conversely, Δ*FTL1678* was more resistant to antibiotics and molecules that

must cross the IM to exert their toxic effects, including gentamicin, tetracycline, chloramphenicol,

ciprofloxacin, and ethidium bromide (Table 2), suggesting that Δ*FTL1678* bacteria have a less

permeable IM. Mechanisms to explain why Δ*FTL1678* bacteria were more resistant to other

stressors, including Triton X-100, H_2O_2, high salt, and low pH, are less clear, but could include

increased expression/activity of chaperone proteins, efflux pumps, antioxidant/scavenger proteins,

and membrane stabilizing proteins (Knodler et al, 2003; Lund et al, 2014; Mishra & Imlay, 2012).

The extreme virulence of Type A *F. tularensis* and its designation as a Tier 1 Select Agent

highlight why studies to identify *F. tularensis* virulence factors and the development of new

vaccines is important. In this study, we identified the role of a previously unstudied protein,

FTL1678, in PG recycling, PG integrity, and bacterial morphology. In addition, we found that

FTL1678 was required for *F. tularensis* LVS virulence and demonstrated that Δ*FTL1678*

conferred 80% protection against fully-virulent, Type A *F. tularensis* SchuS4 pulmonary

challenge. Further studies are needed to determine specific immune responses induced by

Δ*FTL1678* immunization, as well as to identify the most effective immunization regimen (*e.g.,*

number of immunizations and time between immunizations).

Finally, given our findings that PG maintenance and recycling are important for *F.*

tularensis virulence, and that future studies may reveal additional PG-associated enzymes, we used

bioinformatic approaches to predict other proteins involved in *F. tularensis* PG synthesis and

recycling (Figure 9). While at least seven PG synthesis and recycling genes/proteins orthologs

could not be identified, 22 putative PG synthesis and recycling proteins were identified in *F.*

tularensis (Figure 9). Of these, only DacD (FTL1060/FTT1029) has been studied in *F. tularensis*.

Given our observed attenuation of Δ*FTL1678*, future studies to better understand PG synthesis and

recycling pathways may offer more opportunities to better understand the virulence of *F. tularensis*

and other intracellular pathogens. Characterization of other proteins involved in PG pathways may

provide clues as to why *F. tularensis* LdcA is OM-associated or periplasmic and encodes both

L,D-carboxypeptidase and L,D-endopeptidase activities.

Experimental Procedures

Bacterial strains and culture conditions

F. tularensis Type A strain SchuS4 and *F. tularensis* Type B strain LVS were obtained from BEI

Resources and cultured as previously described (Ren, 2014; Wu, 2016). All experiments with

SchuS4 were performed under BSL3 containment conditions at the University of Toledo Health

Science Campus BSL3 laboratory. Routine *F. tularensis* cultures were grown overnight at 37°C

with 5% CO_2 on supplemented Mueller-Hinton agar (sMHA): Mueller-Hinton broth powder

(Becton Dickinson) was mixed with 1.6% (wt/vol) Bacto Agar (Becton Dickinson), autoclaved,

and further supplemented with 2.5% (vol/vol) bovine calf serum (Hyclone), 2% (vol/vol)

IsoVitaleX (Becton Dickinson), 0.1% (wt/vol) glucose, and 0.025% (wt/ vol) iron pyrophosphate.

For mouse infections, *F. tularensis* was first grown on sMHA then transferred to Brain Heart

Infusion agar (BHI; Becton Dickinson). Chocolate agar for mutant strain generation was prepared

by mixing Mueller Hinton broth powder with 1.6% (wt/vol) agar, 1% (wt/vol) tryptone, and 0.5%

(wt/vol) sodium chloride, autoclaved, and further supplemented with 1% (wt/vol) hemoglobin and

1% (vol/vol) IsoVitaleX. For macrophage infections, *F. tularensis* was first grown on sMHA then

transferred to modified chocolate agar: Mueller-Hinton broth powder was mixed with 1.6%

(wt/vol) Bacto Agar, 1% hemoglobin (wt/vol), and 1% (vol/vol) IsoVitaleX. All growth curves

were performed in sMHB: Mueller-Hinton broth powder was mixed with 182 mg/L calcium

chloride dihydrate, and 210 mg L^{-1} magnesium chloride hexahydrate, 0.1% (wt/vol) glucose,

0.025% (wt/vol) iron pyrophosphate, and 2% (vol/vol) IsoVitaleX. All bacterial strains and

plasmids are listed in Table S6. *E. coli* S17-1 and *E. coli* NEB10-β were grown in Luria Bertani

(LB) broth or on LB agar at 37°C, supplemented as needed with antibiotics.

Sequence Alignments and Bioinformatic Predictions

Amino acid alignments of *F. tularensis* subsp. *holarctica* FTL_1678, *F. tularensis* subsp.

tularensis FTT_0101, *E. coli* LdcA (BAA36050.1), *P. aeruginosa* LdcA (Q9HTZ1), *N.*

gonorrhoeae (YP_208343.1), and *C. jejuni* Pgp2 (WP_002856863) were performed using Clustal

Omega   (https://www.ebi.ac.uk/Tools/msa/clustalo/)   and   MView

(https://www.ebi.ac.uk/Tools/msa/mview/). Pairwise sequence alignments were performed and

amino acid identities among Ldc homologues were calculated by EMBOSS Needle

(https://www.ebi.ac.uk/Tools/psa/emboss_needle/). The Prokaryotic Genome Analysis Tool

(PGAT)   (http://tools.uwgenomics.org/pgat/),   BlastP,   and   BlastX   analyses

(http://blast.ncbi.nlm.nih.gov) were used to identify *F. tularensis* homologues. Bacterial protein

sub-localization was predicted by PSORTb version 3.0.2 (https://www.psort.org/psortb/). Protein

signal   sequence   prediction   was   performed   by   LipoP   version   1.0

(http://www.cbs.dtu.dk/services/LipoP/)   and   SignalP   version   4.1

(http://www.cbs.dtu.dk/services/SignalP-4.1/).

Generation of F. tularensis gene deletion mutants

F. tularensis isogenic deletion mutants were generated by homologous recombination as

previously described (Wu, 2015). Briefly, 500 bp regions upstream and downstream from the gene

of interest (*FTL1678* or *FTT0101*) were PCR-amplified from *F. tularensis* genomic DNA using

the following primers: FTL1678_A and FTL1678_B; FTL1678_C and FTL1678_D; FTT0101_A

and FTT0101_B; FTT0101_C and FTT0101_D (Table S7). A FLP recombination target (FRT)-

flanked Pfn-kanamycin resistance cassette, FRT-Pfn-kan-FRT, was PCR amplified from pLG66a

(Gallagher, 2008) and splicing overlap extension PCR (SOE PCR) was used to join the upstream

(A-B) and downstream (C-D) regions with FRT-Pfn-kan-FRT, which replaced the gene of interest.

The resulting insert and a suicide plasmid, pTP163 (Robertson, 2013), were digested with ApaI

(New England Biolabs), and ligated using T4 DNA ligase (New England Biolabs). Gene deletion

constructs were transformed into NEB10-β *E. coli* (New England Biolabs), sequence-verified,

transformed into *E. coli* S17-1, and conjugation was performed with *F. tularensis* LVS on sMHA

plates. Conjugants were recovered on chocolate agar supplemented with 200 mg L^{-1} hygromycin

and 100 mg L^{-1} polymyxin B. Individual mutants were selected by sequential plating on sMHA

supplemented with 10 mg L^{-1} kanamycin (sMHA-kan10), sMHA-kan10 with 8% (wt/vol) sucrose,

and final replica plating onto sMHA containing either 200 mg L^{-1} hygromycin (sMHA-hyg200) or

sMHA-kan10. Hyg-sensitive and kan-resistant colonies were sequence verified (referred to

hereafter as either Δ*FTL1678* or Δ*FTT0101*).

FTL1678 complementation in trans

Complementation *in trans* was performed as previously described, with some modifications (Wu,

2016). *FTL1678* was PCR-amplified from *F. tularensis* LVS using primers

5'FTL1678_NEBuilder and 3'FTL1678_NEBuilder (Table S7), pQE-60 (Qiagen) was double-

digested with NcoI and BglII (New England Biolabs), and the NEBuilder HiFi DNA Assembly

Cloning kit was used to ligate the *FTL1678* amplicon and digested pQE-60. The construct was

transformed into NEB 10-β *E. coli* and transformants were selected on LB agar supplemented with

100 mg L^{-1} ampicillin (LB-amp). Plasmids were purified from individual clones using the Qiagen

QIAprep Spin Miniprep kit (Qiagen), diagnostic PCR was performed to confirm insert presence

and correct size, and DNA sequencing was performed to verify insert integrity. The resulting

construct, *FTL1678* with a C-terminal 6x histidine tag, was PCR-amplified using primers

5'FTL1678_pFNLTP6 and 3'FTL1678_pFNLTP6 (Table S7), the amplicon and pFNLTP6-gro-

GFP (Maier, 2004) were double-digested with XhoI and BamHI (New England Biolabs), and

ligated using T4 DNA Ligase. The construct, pFNLTP6-gro-*FTL1678*-6xHis, was transformed

into NEB10-β *E. coli*, transformants were selected on LB plates supplemented with 50 mg L^{-1}

kanamycin (LB-kan), and DNA sequencing was performed to verify FTL1678-6xHis integrity.

Next, the kan resistance gene was removed from Δ*FTL1678* by suspending the strain in 0.5 M

sucrose (in 1 mM EDTA, pH 7.5), washing three times, and electroporating the shuttle plasmid

pTP405 (Robertson, 2013), which encodes the Flp recombinase to remove FRT-Pfn-kan-FRT from

the genome. Bacteria were grown overnight on sMHA-hyg200, hyg-resistant transformants were

passaged three times on sMHA, then transformants were replica plated onto sMHA-hyg200 and

sMHA-kan10 to confirm sensitivity to both antibiotics (kan-cured Δ*FTL1678*). pFNLTP6-gro-

FTL1678-6xHis was transformed into kan-cured Δ*FTL1678* by electroporation, transformants

were selected on sMHA-kan10, and expression of FTL1678 was confirmed by immunoblot

analysis (referred to hereafter as Δ*FTL1678 trans*-complement).

C. jejuni Pgp2 complementation in trans

Complementation of *C. jejuni pgp2* (CJJ81176_0915) into Δ*FTL1678* was performed as described

above, with several modifications. The *pgp2* gene, with the FTL1678 signal sequence (amino acid

residues 1-29) in place of the native Pgp2 signal sequence (amino acid residues 1-18), was

synthesized and inserted in pQE-60 by GenScript USA. pQE-60-*pgp2* was transformed into

NEB10-β *E. coli* and selection was performed on LB-amp. Pgp2-6xHis was amplified from pQE-

60 using primers 5'FTL1678_pFNLTP6 and 3'FTL1678_pFNLTP6 (Table S7), the amplicon was

ligated into similarly digested pFNLTP6, pFNLTP6-gro-*pgp2*-6xHis was transformed into

NEB10-β *E. coli*, and transformants were selected on LB-kan. Plasmids were purified from kan-

resistant transformants, sequence verified, then electroporated into kan-cured Δ*FTL1678*. Pgp2

expression was confirmed by immunoblot analysis.

Mouse Infections

All animal studies were approved by the University of Toledo Institutional Animal Care and Use

Committee (IACUC). Mouse infections were performed as previously described (Huntley, 2008),

with some modifications.   Briefly, *F. tularensis* strains were grown on sMHA overnight,

transferred to BHI agar for an additional 20-24 h, suspended in sterile PBS, and diluted to the

desired concentration (20 to 10^9 CFU/20 µl) based on previous OD_{600} measurements and bacterial

enumeration studies.   Groups of 4-8 female C3H/HeN mice (6-8 weeks old; Charles River

Laboratories) were anesthetized with a ketamine-xylazine sedative and intranasally (i.n.) infected

with 20 µl of prepared bacterial suspensions. Bacterial innocula were serially-diluted and plated

in quadruplet on sMHA to confirm CFUs. For survival studies, mice were monitored daily, for

signs of disease, with health status scores (scale of 1-5, with 1 indicating healthy and 5 indicating

mouse found dead) being recorded for each mouse. Moribund mice were humanely euthanized to

minimize suffering. To quantitate bacterial tissue burdens, groups of 4 mice were euthanized on

days 2 and 5 post-infection, blood was collected by cardiac puncture and plated onto sMHA, lungs,

livers, and spleens were aseptically harvested, homogenized, 25 µl of PBS/mg of tissue was added

to each tissue, serially-diluted, and dilutions were plated onto sMHA. Following 72 h of

incubation, the number of colonies per plate were counted and CFU/mg (tissues) or CFU/ml

(blood) were calculated based on tissue weight and dilution factor.   For immunization and

challenge studies, groups of 4-10 mice were i.n. immunized with either 100-300 CFU LVS or 10^4-

10^9 CFU ΔFTL1678, boosted 3-4 weeks later with either 10^3 CFU LVS or 10^9 CFU ΔFTL1678,

transported to the ABSL3 facility 3-weeks later, and i.n. challenged with 20-120 CFU of *F.*

tularensis SchuS4. Mice were monitored daily for signs of disease with health status scores being

recorded for each mouse.

Membrane Integrity Testing

Sensitivity of LVS, Δ*FTL1678*, FTL1678 *trans*-complement, and the Pgp2 *trans*-complement to

various antibiotics, detergents, dyes, and cell wall stressors was determined by disk diffusion

assays or in liquid cultures, as previously described (Wu, 2016), with some modifications.

Bacterial strains were grown on either sMHA or sMHA-kan10 (ΔFTL1678 and complement

strains), scraped and resuspended in sterile PBS, adjusted to an OD_{600} of 0.2 (approx. $9x10^7$

CFU/ml), diluted 1:1 in PBS, and 100 µl was plated onto sMHA plates using cotton tipped

applicators (Puritan). Sterile paper disks (Whatman; 0.8 mm thick, 6.5 mm in diameter) were

placed in the center of each plate and antibiotics, detergents, or dyes were added to the disks at the

concentrations listed in Table 2. Antibiotics tested were: gentamicin (Gibco), tetracycline (Fisher

Scientific), chloramphenicol (Acros Organics), ciprofloxacin (Oxoid), ampicillin (Fisher

Scientific), vancomycin (Acros Organics), bacitracin (Oxoid), bacitracin (Oxoid), ciprofloxacin

(Oxoid), and polymyxin b (MP Biomedicals). Detergents tested were: sodium dodecyl sulfate

(SDS; anionic; Fisher Scientific), Triton X-100 (nonionic; Acros Organics), cetyltrimethyl

ammonium bromide (CTAB; cationic; MP Biomedicals), 3-cholamidopropyl dimethylammonio

1-propanesulfonate (CHAPS; zwitterionic; Thermo Scientific). In addition, sensitivity to ethidium

bromide (Thermo Scientific) and lysozyme (Thermo Fisher) also was tested. After 48 h, diameters

of zones of inhibition around the disks were measured. Experiments were performed in triplicate.

For liquid cultures, bacteria were suspended in sMHB, adjusted to OD_{600} 0.4, and 5 ml of each

bacterial suspensions was inoculated into 100 ml of either sMHB or sMHB with 5 mM hydrogen

peroxide (H_2O_2), 5% sodium chloride (NaCl), or pH 5.5 (pH of sMHB is 6.5). Cultures were

grown at 37˚C with rotation at 180 rpm for 24 h with OD_{600} readings recorded every 4 hours.

Electron Microscopy

Electron microscopy was used to visualize differences in bacterial envelope structure and cell

shape, as previously described (Wu, 2016), with some modifications. LVS, Δ*FTL1678*, FTL1678

trans-complement, and the Pgp2 *trans*-complement were grown overnight in sMHB, approx.

$1x10^9$ CFU of each bacterial strain was pelleted by centrifugation at $7000 \times g$ at 4˚C, washed three

times in PBS, fixed in 3% (vol/vol) glutaraldehyde (Electron Microscopy Sciences [EMS]) for

approx. 24 hours, washed twice in sodium cacodylate buffer (pH 7.4; EMS) for 10 min, suspended

in 1% (wt/vol) osmium tetroxide (EMS) in s-collidine buffer (pH 7.4; EMS) for 45 min at room

temperature (r/t) to stain and fix the samples, washed two times with sodium cacodylate buffer for

10 min each, and tertiary fixation was performed using an aqueous saturated solution of uranyl

acetate (pH 3.3; EMS) for 45 min at r/t. Samples were then dehydrated at room temperature using

a series of ethanol washes: two washes with 30% ethanol for 10 min each; two washes with 50%

ethanol for 10 min each; two washes with 95% ethanol for 10 min each; two washes with 100%

ethanol for 10 min each; and two washes with 100% acetone for 10 min each. Samples were then

infiltrated with 50% acetone and 50% embedding media (Hard Plus Resin 812, EMS) for 8 h to

overnight at r/t. Samples were embedded in 100% embedding media (EMS) and allowed to

polymerize for 8 h to overnight at 85°C, then sectioned at 85–90 nm, and visualized using a Tecnai

G2 Spirit transmission electron microscope (FEI) at 80 kV and Radius 1.3 (Olympus) camera

software at the University of Toledo Electron Microscopy Facility. Experiments were performed

twice to confirm reproducibility, with two bacterial preparations fixed, stained, embedded,

sectioned, and visualized per experiment.

Spheroplasting and Sucrose Density Gradient Centrifugation

Spheroplasting, osmotic lysis, and sucrose density gradient centrifugation was performed as previously described (Huntley, 2007) to determine subcellular localization of FTL1678. Briefly, the histidine-tagged FTL1678 *trans*-complement was grown in sMHB to an OD_{600} of 0.3-0.4, pelleted at 7500 × g for 30 min at 10˚C, supernatants were removed, pellets were resuspended in 0.75 M sucrose (in 5 mM Tris, pH 7.5) with gentle mixing, 10 mM EDTA (in 5 mM Tris, pH 7.8) was slowly added over 10 min, and the suspension was incubated for 30 min at r/t. After incubation, lysozyme was slowly added to a final concentration of 200 μg ml^{-1}, incubated for 30 min at r/t, bacteria were osmotically lysed by dilution into 4.5 × volume of molecular-grade water (Corning) over 11 min with gentle mixing, and incubated for 30 min at r/t. Lysates were centrifuged at 7,500 × g for 30 min at 10°C to remove intact cells and cellular debris. Supernatants were collected and centrifuged at 182,500 × g for 2 h at 4°C in a F37L 8 × 100 Fiberlite Ultracentrifuge rotor. Following centrifugation, supernatants were removed, membrane pellets were gently resuspended in 6 ml of resuspension buffer (25% [wt/wt] sucrose, 5 mM Tris, 30 mM $MgCl_2$, 1 tablet of Pierce Protease Inhibitor Mini Tablets, EDTA-Free [Thermo Scientific], 5 U Benzonase [Novagen]), suspensions were incubated with gentle mixing for 30 min at room temperature to degrade DNA, and a DC protein assay (Bio-Rad) was performed to determine total protein yield. Linear sucrose gradients were prepared by layering 1.8 ml each of sucrose solutions (wt/wt; prepared in 5 mM EDTA, pH 7.5) into 14- by 95-mm ultracentrifuge tubes (Beckman) in the following order: 55%, 50%, 45%, 40%, 35%, and 30%. Membrane suspensions were layered on top of each sucrose gradient, with less than 1.5 mg of protein per gradient. Sucrose gradients were centrifuged in an SW40 swinging bucket rotor (Beckman) at 256,000 × g for 17 h at 4°C. After centrifugation, 500 μl fractions were collected from each gradient by puncturing the bottom

of each tube and allowing fractions to drip into microcentrifuge tubes. The refractive index of

each fraction was determined using a refractometer (Fisher Scientific) and correlated with a

specific density in g ml^{-1} (Price, 1982) to identify outer membrane (OM; 1.17-1.20 g/ml) and inner

membrane (IM; 1.13-1.14 g/ml) fractions. Sucrose gradient fractions were examined by

immunoblotting as described below.

Immunoblotting

Whole cell lysates of FTL1678 *trans*-complement were prepared by suspending bacteria (pelleted

at 7000 × *g*) in molecular biology grade water, diluting with SDS-PAGE loading buffer, and

boiling for 10 min. Whole cell lysates, OM fractions, IM fractions, and molecular mass standards

(Precision Plus protein all blue prestained protein standards; BioRad Laboratories) were separated

on a 12.5% polyacrylamide gel, transferred to nitrocellulose, and blots were incubated overnight

in blot block (0.1% (vol/vol) Tween 20 and 2% (wt/vol) bovine serum albumin in PBS) at 4°C.

Immunoblotting was performed using rat polyclonal antiserum specific for either *F. tularensis* OM

protein FopA, *F. tularensis* IM protein SecY (Huntley, 2007) or the Penta-His HRP conjugate

antibody (Qiagen).

Infections of Mouse Bone Marrow Derived Macrophages (mBMDMs) and J774A.1 cells

Macrophage culture (37˚C with 5% CO$_2$ unless otherwise indicated) and infections were

performed as previously described (Wu, 2016), with some modifications. Bone marrow

macrophages were harvested from female C3H/HeN mice. Mice were euthanized by CO$_2$

asphyxiation and cervical dislocation. Femurs and tibias of both hind legs were aseptically-

harvested, marrow was flushed from each bone with RPMI-1640 (Hyclone) containing 10% heat-

inactivated fetal bovine serum ([HI-FBS], Atlanta Biologicals) and 30% supernatants from day 7

L929 cultures (ATCC). Bone marrow was disrupted by repeated passage through a 23-gauge

needle and cultured for 4 days. Next, cell media was removed and replaced with RPMI containing

10% HI-FBS and 30% supernatant from day 14 L929 cultures, and cells were cultured for 2 days.

Approx. 24 h before infection, media was removed, cells were harvested by scraping and

centrifugation at $400 \times g$ for 10 min at 10°C, cells were enumerated using a hemocytometer, and

diluted to $1x10^5$ cells in RPMI containing 10% HI-FBS. J774A.1 cells (ATCC) were cultured in

Dulbecco's Modified Eagle Medium ([DMEM], Gibco) containing 10% HI-FBS. Approx. 24 h

before infection, cells were harvested as described above, seeded into individual wells of 24-well

plates (Corning) at a concentration of $1x10^5$ cells/well, and incubated overnight. mBMDMs and

J774A.1 cells were infected with a multiplicity of infection (MOI) of 100 bacteria to 1 cell (100:1).

Following infection, cells were centrifuged at $1,000 \times g$ for 10 min at 4°C, incubated at 37°C with

5% CO_2 for 1 h, washed $1 \times$ with RPMI (or DMEM), media containing 100 µg ml^{-1} gentamicin

was added to kill extracellular bacteria, cells were incubated at 37°C with 5% CO_2 for 1 h, washed

1x with RPMI (or DMEM), lysed with 1% saponin for 4 min, serially diluted in PBS, plated onto

sMHA plates, and bacteria were enumerated (entry) after 48h. Alternatively, after gentamicin

treatment and washing, RPMI (or DMEM) containing 10% HI-FBS was added to cells and they

were incubated for 6 or 24 h, lysed, serially-diluted, and plated to determine bacterial numbers.

Expression and Purification of Recombinant FTL1678 and FTT0101

F. tularensis LVS and SchuS4 genomic DNA were extracted using phenol/chloroform/isoamyl

alcohol (Fisher Bioreagents). *FTL1678* and *FTT0101*, without signal sequences (amino acid

residues 1-29), were PCR-amplified from LVS and SchuS4 genomic DNA, respectively, using

High Fidelity Platinum Taq Polymerase (Life Technologies), and primers 5'FTL1678_BamHI and

3'FTL1678_XhoI and 5'FTT0101_BamHI and 3'FTT0101_XhoI, respectively (Table S7).

Amplicons and pPROEX HTb were double-digested with BamHI and XhoI, ligated using T4 DNA

ligase, and transformed into NEB 10-β *E. coli*. Plasmids were purified using the Qiagen QIAprep

Spin Miniprep kit and diagnostic PCR was performed to confirm presence and correct size of the

insert. DNA sequencing was performed to confirm insert integrity and plasmid constructs were

transformed into Rosetta DE3 *E. coli* (Millipore) for protein expression. Recombinant proteins

were expressed and purified as previously described (Ren, 2014) with some modifications.

Bacteria were grown in LB-amp to an OD_{600} of 0.4, protein expression was induced for 2 h by the

addition of isopropyl β-D-thiogalactopyranoside (IPTG) to a final concentration of 100 mM,

bacteria were pelleted by centrifugation, and frozen overnight at −80°C to aid in lysis. Cell pellets

were suspended in 10 mM Tris, 500 mM NaCl, and 10 mM imidazole, pH 8.0, sonicated on ice

for 10 min with 30 sec intervals, insoluble material was removed by centrifugation at $8,000 \times g$,

and supernatants were collected for affinity purification over pre-equilibrated Ni-nitrilotriacetic

acid (Ni-NTA) agarose (Qiagen) columns. Eluted recombinant proteins were concentrated in

Amicon Ultra-4 centrifugal filter units with 30-kDa cutoff (Millipore), concentrations were

determined using the DC BCA protein assay (BioRad), and purity was assessed by SDS-PAGE

and Imperial protein staining (Thermo Scientific). An empty vector construct also was expressed

and purified as a control in enzymatic assays.

Enzymatic Assays for FTL1678/FTT0101 Activity

FTL1678 and FTT0101 recombinant protein activity toward various PG-related compounds were

tested in 50 μl reaction mixtures containing 50 mM Tris-HCl, pH 8.0, 0.1 mM substrate, and

partially purified enzyme stock (10 μl in 1 M NaCl, 10 mM Tris, pH 8.0). Mixtures were incubated

for 30 min to 2 h at 37°C and reactions were stopped by freezing. Substrate and reaction products

were separated by HPLC on an ODS-Hypersil 3 μm particle-size C18 column (250 by 4.6 mm;

Thermo Scientific). Elutions were performed with 50 mM sodium phosphate buffer, pH 4.5, with

or without application of a linear gradient of methanol (from 0 to 50% in 50 min), at a flow rate of

0.5 ml/min. Peaks were detected by measuring the absorbance at 207 nm or at 262 nm for UDP-

containing nucleotide precursors. Identification of compounds was based on their retention times,

compared to authentic standards, as well as on their amino acid and amino sugar composition,

determined with a Hitachi model L8800 analyzer (Sciencetec) after hydrolysis of samples in 6 M

HCl for 16 h at 95°C. Enzyme activity was calculated by integration of peaks corresponding to

substrate and product. Amounts of alanine released by the L,D-carboxypeptidase activity also

were determined using an amino acid analyzer. Depending on the substrate used, the amount of

partially purified protein varied from 0.9 to 5 µg per assay and incubation times varied from 30

min to 4 h. To ensure linearity, substrate consumption was < 20% in all cases. Values represent

the means for three independent experiments; the standard deviation was < 10% in all cases.

Specific activities were calculated from the amounts of D-Ala (tetrapeptide substrates) or D-Ala-

D-Ala (pentapeptide substrates) released during the reaction. FTL1678 double and triple active

site mutants were tested in 50 µl reaction mixtures containing 50 mM Tris-HCl, pH 8.0, 0.05 mM

substrate, and partially purified enzyme stock (5-10 µl in 1 M NaCl, 10 mM Tris, pH 8.0).

Reactions were incubated for 18 hours then analyzed by HPLC as described above. Peaks were

detected by measuring absorbance at 207 nm.

Peptidoglycan Precursors and Muropeptides

UDP-MurNAc-pentapeptide precursors containing either *meso*-diaminopimelic acid (A_2pm) or L-

lysine were prepared by enzymatic synthesis using purified Mur ligases, and UDP-MurNAc-

tetrapeptides were generated by treatment of the UDP-MurNAc-pentapeptide precursors with

purified *E. coli* PBP5 DD-carboxypeptidase as previously described (Herve, 2007). MurNAc-

peptides were obtained by mild acid hydrolysis of UDP-MurNAc-peptides (0.1 M HCl, 100°C, 15

min) and were not reduced and thus purified as a mixture of the two α and β anomers (Blanot,

1983). Free peptides were prepared by cleavage of MurNAc-peptides with *E. coli* AmiD *N*-

acetylmuramoyl-L-alanine amidase (Pennartz, 2009). The *E. coli* peptidoglycan polymer was

purified from a Δ*lpp* mutant strain that does not express the Lpp lipoprotein (Leulier, 2003).

GlcNAc-1,6-anhydro-MurNAc-L-Ala-γ-D-Glu-*meso*-A$_2$pm-D-Ala (TCT) and its dimer (two

cross-linked TCT monomers) were produced by digestion of peptidoglycan with *E. coli* SltY lytic

transglycosylase and the non-anhydro forms of these monomer and dimer were generated by

digestion of the polymer with mutanolysin (Stenbak, 2004). All these compounds were HPLC-

purified and their composition was controlled by amino acid and amino sugar content analysis

and/or by MALDI-TOF mass spectrometry.

Statistics

GraphPad Prism6 was used in various statistical analyses, including: differences in antibiotic,

detergent, dye, or lysozyme susceptibility were calculated by one-way ANOVA with multiple

comparisons and the Holm-Sidak post-hoc test; differences in EM measurements were determined

by unpaired t-tests; differences in median time-to-death and percent survival following *F.*

tularensis infection of mice were calculated using the log-rank Mantel-Cox test; differences in

pathology scores of *F. tularensis*-infected tissues were calculated by two-way ANOVA with

multiple comparisons and a Tukey post-hoc test. Differences in lung, liver, spleen, and blood

bacterial burdens from infected mice were calculated by one-way ANOVA with multiple

comparisons using R software.

Acknowledgments

We would like to acknowledge Drs. Joe Dillard and Ryan Schaub (University of Wisconsin-

Madison) for all their help and expertise with peptidoglycan isolation/purification. We would also

like to thank Dr. Erin Gaynor for her generous contribution of *C. jejuni* Pgp2-pGEM plasmid

construct. This work was supported by grant R01 AI093351 from the National Institute of Allergy

and Infectious Disease of the National Institutes of Health (NIAID-NIH) to J.F.H.

Data Availability Statement

Data that support the findings of this study are available in the supplementary material of this

article.

References

Blanot, D., Kretsovali A, Abo-Ghalia M, Mengin-Lecreulx D, van Heijenoort J (1983) Synthesis

of analogues of precursors of bacterial peptidoglycan, in Malon, B. a. P. (ed), *Peptides*.

Berlin, Germany: Walter de Gruyter, 311-314.

Bouveret, E., Benedetti H, Rigal A, Loret E, Lazdunski C (1999) In vitro characterization of

peptidoglycan-associated lipoprotein (PAL)-peptidoglycan and PAL-TolB interactions. *J

Bacteriol*, 181(20), 6306-11.

Braun, V. (1975) Covalent lipoprotein from the outer membrane of *Escherichia coli*. *Biochim

Biophys Acta*, 415(3), 335-77.

Braun, V. & Hantke, K. (2019) Lipoproteins: Structure, Function, Biosynthesis. *Subcell

Biochem*, 92, 39-77.

Braun, V., Rehn K (1969) Chemical characterization, spatial distribution and function of a

lipoprotein (murein-lipoprotein) of the *E. coli* cell wall. The specific effect of trypsin on

the membrane structure. *Eur J Biochem*, 10(3), 426-38.

Braun, V. & Rehn, K. (1969) Chemical characterization, spatial distribution and function of a

lipoprotein (murein-lipoprotein) of the *E. coli* cell wall. The specific effect of trypsin on

the membrane structure. *Eur J Biochem*, 10(3), 426-38.

Chaput, C., Ecobichon C, Pouradier N, Rousselle JC, Namane A, Boneca IG (2016) Role of the

N-Acetylmuramoyl-l-Alanyl Amidase, AmiA, of *Helicobacter pylori* in Peptidoglycan

Metabolism, Daughter Cell Separation, and Virulence. *Microb Drug Resist*, 22(6), 477-

86.

Clavel, T., Germon P, Vianney A, Portalier R, Lazzaroni JC (1998) TolB protein of *Escherichia*

coli K-12 interacts with the outer membrane peptidoglycan-associated proteins Pal, Lpp

and OmpA. *Mol Microbiol*, 29(1), 359-67.

Das, D., Herve M, Elsliger MA, Kadam RU, Grant JC, Chiu HJ, Knuth MW, Klock HE, Miller

MD, Godzik A, Lesley SA, Deacon AM, Mengin-Lecreulx D, Wilson IA (2013)

Structure and function of a novel LD-carboxypeptidase a involved in peptidoglycan

recycling. *J Bacteriol*, 195(24), 5555-66.

De Pascalis, R., Hahn A, Brook HM, Ryden P, Donart N, Mittereder L, Frey B, Wu TH, Elkins

KL (2018) A panel of correlates predicts vaccine-induced protection of rats against

respiratory challenge with virulent *Francisella tularensis*. *PLoS One*, 13(5), e0198140.

de Pedro, M., Cava F (2015) Structural constraints and dynamics of bacterial cell wall

architecture. *Front Microbiol*, 6, 449.

den Blaauwen, T., de Pedro MA, Nguyen-Disteche M, Ayala JA (2008) Morphogenesis of rod-

shaped sacculi. *FEMS Microbiol Rev*, 32(2), 321-44.

Dennis, D., Inglesby TV, Henderson DA, Bartlett JG, Ascher MS, Eitzen E, Fine AD,

Friedlander AM, Hauer J, Layton M, Lillibridge SR, McDade JE, Osterholm MT,

O'Toole T, Parker G, Perl TM, Russell PK, and Tonat K (2001) Tularemia as a biological

weapon: medical and public health management. *JAMA*, 285(21), 2763-73.

Denome, S., Elf PK, Henderson TA, Nelson DE, Young KD (1999) *Escherichia coli* mutants

lacking all possible combinations of eight penicillin binding proteins: viability,

characteristics, and implications for peptidoglycan synthesis. *J Bacteriol*, 181(13), 3981-

93.

Dhar, S., Kumari H, Balasubramanian D, Mathee K. (2018) Cell-wall recycling and synthesis in

Escherichia coli and *Pseudomonas aeruginosa* - their role in the development of

resistance. *J Med Microbiol*, 67(1), 1-21.

Ellis, J., Oyston PC, Green M, Titball RW (2002) Tularemia. *Clin Microbiol Rev*, 15(4), 631-46.

Frirdich, E., Biboy J, Adams C, Lee J, Ellermeier J, Gielda LD, Dirita VJ, Girardin SE, Vollmer

W, Gaynor EC (2012) Peptidoglycan-modifying enzyme Pgp1 is required for helical cell

shape and pathogenicity traits in *Campylobacter jejuni. PLoS Pathog*, 8(3), e1002602.

Frirdich, E., Vermeulen J, Biboy J, Soares F, Taveirne ME, Johnson JG, DiRita VJ, Girardin SE,

Vollmer W, Gaynor EC (2014) Peptidoglycan LD-carboxypeptidase Pgp2 influences

Campylobacter jejuni helical cell shape and pathogenic properties and provides the

substrate for the DL-carboxypeptidase Pgp1. *J Biol Chem*, 289(12), 8007-18.

Gallagher, L., McKevitt M, Ramage ER, Manoil C (2008) Genetic dissection of the *Francisella*

novicida restriction barrier. *J Bacteriol*, 190(23), 7830-7.

Glauner, B., Holtje JV, Schwarz U (1988) The composition of the murein of *Escherichia coli. J*

Biol Chem, 263(21), 10088-95.

957 Guinane, C., Cotter PD, Ross RP, Hill C (2006) Contribution of penicillin-binding protein

958   homologs to antibiotic resistance, cell morphology, and virulence of *Listeria*

959   *monocytogenes* EGDe. *Antimicrob Agents Chemother*, 50(8), 2824-8.

960 Hall, J., Woolard MD, Gunn BM, Craven RR, Taft-Benz S, Frelinger JA, Kawula TH (2008)

961   Infected-host-cell repertoire and cellular response in the lung following inhalation of

962   *Francisella tularensis* Schu S4, LVS, or U112. *Infect Immun*, 76(12), 5843-52.

963 Heidrich, C., Templin MF, Ursinus A, Merdanovic M, Berger J, Schwarz H, de Pedro MA,

964   Holtje JV (2001) Involvement of N-acetylmuramyl-L-alanine amidases in cell separation

965   and antibiotic-induced autolysis of *Escherichia coli*. *Mol Microbiol*, 41(1), 167-78.

966 Heidrich, C., Ursinus A, Berger J, Schwarz H, Holtje JV (2002) Effects of multiple deletions of

967   murein hydrolases on viability, septum cleavage, and sensitivity to large toxic molecules

968   in *Escherichia coli*. *J Bacteriol*, 184(22), 6093-9.

969 Herve, M., Boniface A, Gobec S, Blanot D, Mengin-Lecreulx D (2007) Biochemical

970   characterization and physiological properties of *Escherichia coli* UDP-N-

971   acetylmuramate:L-alanyl-gamma-D-glutamyl-meso-diaminopimelate ligase. *J Bacteriol*,

972   189(11), 3987-95.

973 Holtje, J. (1998) Growth of the stress-bearing and shape-maintaining murein sacculus of

974   *Escherichia coli*. *Microbiol Mol Biol Rev*, 62(1), 181-203.

975 Huntley, J., Conley PG, Hagman KE, Norgard MV (2007) Characterization of *Francisella*

976   *tularensis* outer membrane proteins. *J Bacteriol*, 189(2), 561-74.

977 Huntley, J., Conley PG, Rasko DA, Hagman KE, Apicella MA, Norgard MV (2008) Native

978   outer membrane proteins protect mice against pulmonary challenge with virulent type A

979   *Francisella tularensis*. *Infect Immun*, 76(8), 3664-71.

Jenkins, C., Wallis R, Allcock N, Barnes KB, Richards MI, Auty JM, Galyov EE, Harding SV,
Mukamolova GV (2019) The lytic transglycosylase, LtgG, controls cell morphology and
virulence in *Burkholderia pseudomallei. Sci Rep*, 9(1), 11060.

Johnson, J., Fisher JF, Mobashery S (2013) Bacterial cell-wall recycling. *Ann N Y Acad Sci*,
1277, 54-75.

Jones, A. S., Austerberry, J. I., Dajani, R., Warwicker, J., Curtis, R., Derrick, J. P. & Robinson,
C. (2016) Proofreading of substrate structure by the Twin-Arginine Translocase is highly
dependent on substrate conformational flexibility but surprisingly tolerant of surface
charge and hydrophobicity changes. *Biochim Biophys Acta*, 1863(12), 3116-3124.

Jones, C., Napier BA, Sampson TR, Llewellyn AC, Schroeder MR, Weiss DS (2012) Subversion
of host recognition and defense systems by *Francisella* spp. *Microbiol Mol Biol Rev*,
76(2), 383-404.

Juan, C., Torrens G, Barcelo IM, Oliver A (2018) Interplay between Peptidoglycan Biology and
Virulence in Gram-Negative Pathogens. *Microbiol Mol Biol Rev*, 82(4), e00033-18.

Keim, P., Johansson A, Wagner DM (2007) Molecular epidemiology, evolution, and ecology of
Francisella. Ann N Y Acad Sci, 1105, 30-66.

Kijek, T., Mou S, Bachert BA, Kuehl KA, Williams JA, Daye SP, Worsham PL, Bozue JA
(2019) The D-alanyl-d-alanine carboxypeptidase enzyme is essential for virulence in the
Schu S4 strain of *Francisella tularensis* and a dacD mutant is able to provide protection
against a pneumonic challenge. *Microb Pathog*, 137, 103742.

Kingry, L., Petersen JM (2014) Comparative review of *Francisella tularensis* and *Francisella*
novicida. Front Cell Infect Microbiol, 4, 35.

Knodler, L. A., Vallance, B. A., Hensel, M., Jackel, D., Finlay, B. B. & Steele-Mortimer, O. (2003) *Salmonella* type III effectors PipB and PipB2 are targeted to detergent-resistant microdomains on internal host cell membranes. *Mol Microbiol*, 49(3), 685-704.

Korza, H., Bochtler M (2005) *Pseudomonas aeruginosa* LD-carboxypeptidase, a serine peptidase with a Ser-His-Glu triad and a nucleophilic elbow. *J Biol Chem*, 280(49), 40802-12.

Leduc, M., Ishidate K, Shakibai N, Rothfield L (1992) Interactions of *Escherichia coli* membrane lipoproteins with the murein sacculus. *J Bacteriol*, 174(24), 7982-8.

Lenz, J., Hackett KT, Dillard JP (2017) A Single Dual-Function Enzyme Controls the Production of Inflammatory NOD Agonist Peptidoglycan Fragments by *Neisseria gonorrhoeae*. *MBio*, 8(5), e01464-17.

Lenz, J. D., Hackett, K. T. & Dillard, J. P. (2017) A Single Dual-Function Enzyme Controls the Production of Inflammatory NOD Agonist Peptidoglycan Fragments by *Neisseria gonorrhoeae*. *MBio*, 8(5).

Leulier, F., Parquet C, Pili-Floury S, Ryu JH, Caroff M, Lee WJ, Mengin-Lecreulx D, Lemaitre B (2003) The *Drosophila* immune system detects bacteria through specific peptidoglycan recognition. *Nat Immunol*, 4(5), 478-84.

Lund, P., Tramonti, A. & De Biase, D. (2014) Coping with low pH: molecular strategies in neutralophilic bacteria. *FEMS Microbiol Rev*, 38(6), 1091-125.

Maier, T., Havig A, Casey M, Nano FE, Frank DW, Zahrt TC (2004) Construction and characterization of a highly efficient *Francisella* shuttle plasmid. *Appl Environ Microbiol*, 70(12), 7511-9.

Mengin-Lecreulx, D. & Lemaitre, B. (2005) Structure and metabolism of peptidoglycan and molecular requirements allowing its detection by the *Drosophila* innate immune system. *J Endotoxin Res*, 11(2), 105-11.

Metz, R., Henning S, Hammes WP (1986a) LD-carboxypeptidase activity in *Escherichia coli*. I.

The LD-carboxypeptidase activity in ether treated cells. *Arch Microbiol*, 144(2), 175-80.

Metz, R., Henning S, Hammes WP (1986b) LD-carboxypeptidase activity in *Escherichia coli*. II.

Isolation, purification and characterization of the enzyme from *E. coli* K 12. *Arch*

Microbiol, 144(2), 181-6.

Meyer, K., Addy C, Akashi S, Roper DI, Tame JRH (2018) The crystal structure and oligomeric

form of *Escherichia coli* L,D-carboxypeptidase A. *Biochem Biophys Res Commun*,

499(3), 594-599.

Minor, P. (2015) Live attenuated vaccines: Historical successes and current challenges. *Virology*,

479-480, 379-92.

Mishra, S. & Imlay, J. (2012) Why do bacteria use so many enzymes to scavenge hydrogen

peroxide? *Arch Biochem Biophys*, 525(2), 145-60.

Nelson, D., Young KD (2000) Penicillin binding protein 5 affects cell diameter, contour, and

morphology of *Escherichia coli*. *J Bacteriol*, 182(6), 1714-21.

Park, J., Uehara T (2008) How bacteria consume their own exoskeletons (turnover and recycling

of cell wall peptidoglycan). *Microbiol Mol Biol Rev*, 72(2), 211-27.

Pazos, M. & Peters, K. (2019) Peptidoglycan. *Subcell Biochem*, 92, 127-168.

Pennartz, A., Genereux C, Parquet C, Mengin-Lecreulx D, Joris B (2009) Substrate-induced

inactivation of the *Escherichia coli* AmiD N-acetylmuramoyl-L-alanine amidase

highlights a new strategy to inhibit this class of enzyme. *Antimicrob Agents Chemother*,

53(7), 2991-7.

Petrosino, J. F., Xiang, Q., Karpathy, S. E., Jiang, H., Yerrapragada, S., Liu, Y., Gioia, J.,

Hemphill, L., Gonzalez, A., Raghavan, T. M., Uzman, A., Fox, G. E., Highlander, S.,

Reichard, M., Morton, R. J., Clinkenbeard, K. D. & Weinstock, G. M. (2006)

Chromosome rearrangement and diversification of *Francisella tularensis* revealed by the type B (OSU18) genome sequence. *J Bacteriol*, 188(19), 6977-85.

Price, C. (1982) Centrifugation in density gradients. *Academic Press*, 335-343.

Priyadarshini, R., de Pedro MA, Young KD (2007) Role of peptidoglycan amidases in the development and morphology of the division septum in *Escherichia coli*. *J Bacteriol*, 189(14), 5334-47.

Priyadarshini, R., Popham DL, Young KD (2006) Daughter cell separation by penicillin-binding proteins and peptidoglycan amidases in *Escherichia coli*. *J Bacteriol*, 188(15), 5345-55.

Qin, A., Scott DW, Thompson JA, Mann BJ (2009) Identification of an essential *Francisella tularensis* subsp. *tularensis* virulence factor. *Infect Immun*, 77(1), 152-61.

Ray, K., Marteyn B, Sansonetti PJ, Tang CM (2009) Life on the inside: the intracellular lifestyle of cytosolic bacteria. *Nat Rev Microbiol*, 7(5), 333-40.

Ren, G., Champion MM, Huntley JF (2014) Identification of disulfide bond isomerase substrates reveals bacterial virulence factors. *Mol Microbiol*, 94(4), 926-44.

Roberts, L., Powell DA, Frelinger JA (2018) Adaptive Immunity to *Francisella tularensis* and Considerations for Vaccine Development. *Front Cell Infect Microbiol*, 8, 115.

Robertson, G., Child R, Ingle C, Celli J, Norgard MV (2013) IglE is an outer membrane-associated lipoprotein essential for intracellular survival and murine virulence of type A *Francisella tularensis*. *Infect Immun*, 81(11), 4026-40.

Russo, T., MacDonald U, Beanan JM, Olson R, MacDonald IJ, Sauberan SL, Luke NR, Schultz LW, Umland TC (2009) Penicillin-binding protein 7/8 contributes to the survival of *Acinetobacter baumannii* in vitro and in vivo. *J Infect Dis*, 199(4), 513-21.

Scheurwater, E., Reid CW, Clarke AJ (2008) Lytic transglycosylases: bacterial space-making autolysins. *Int J Biochem Cell Biol*, 40(4), 586-91.

Silhavy, T. J., Kahne, D. & Walker, S. (2010) The bacterial cell envelope. *Cold Spring Harb*

Perspect Biol, 2(5), a000414.

Sjostedt, A. (2007) Tularemia: history, epidemiology, pathogen physiology, and clinical

manifestations. *Ann N Y Acad Sci*, 1105, 1-29.

Spidlova, P., Stojkova P, Dankova V, Senitkova I, Santic M, Pinkas D, Philimonenko V, Stulik J

(2018) *Francisella tularensis* D-Ala D-Ala Carboxypeptidase DacD Is Involved in

Intracellular Replication and It Is Necessary for Bacterial Cell Wall Integrity. *Front Cell*

Infect Microbiol, 8, 111.

Steiner, D., Furuya Y, Jordan MB, Metzger DW (2017) Protective Role for Macrophages in

Respiratory *Francisella tularensis* Infection. *Infect Immun*, 85(6).

Stenbak, C., Ryu JH, Leulier F, Pili-Floury S, Parquet C, Herve M, Chaput C, Boneca IG, Lee

WJ, Lemaitre B, Mengin-Lecreulx D (2004) Peptidoglycan molecular requirements

allowing detection by the *Drosophila* immune deficiency pathway. *J Immunol*, 173(12),

7339-48.

Sycuro, L., Pincus Z, Gutierrez KD, Biboy J, Stern CA, Vollmer W, Salama NR (2010)

Peptidoglycan crosslinking relaxation promotes *Helicobacter pylori*'s helical shape and

stomach colonization. *Cell*, 141(5), 822-33.

Sycuro, L., Rule CS, Petersen TW, Wyckoff TJ, Sessler T, Nagarkar DB, Khalid F, Pincus Z,

Biboy J, Vollmer W, Salama NR (2013) Flow cytometry-based enrichment for cell shape

mutants identifies multiple genes that influence *Helicobacter pylori* morphology. *Mol*

Microbiol, 90(4), 869-83.

Templin, M., Ursinus A, Holtje JV (1999) A defect in cell wall recycling triggers autolysis

during the stationary growth phase of *Escherichia coli*. *EMBO J*, 18(15), 4108-17.

Typas, A., Banzhaf M, Gross CA, Vollmer W (2011) From the regulation of peptidoglycan

synthesis to bacterial growth and morphology. *Nat Rev Microbiol*, 10(2), 123-36.

Ursinus, A., Steinhaus H, Holtje JV (1992) Purification of a nocardicin A-sensitive LD-

carboxypeptidase from *Escherichia coli* by affinity chromatography. *J Bacteriol*, 174(2),

441-6.

van Heijenoort, J. (2011) Peptidoglycan hydrolases of *Escherichia coli*. *Microbiol Mol Biol Rev*,

75(4), 636-63.

Vollmer, W. & Bertsche, U. (2008) Murein (peptidoglycan) structure, architecture and

biosynthesis in *Escherichia coli*. *Biochim Biophys Acta*, 1778(9), 1714-34.

Walburger, A., Lazdunski C, Corda Y (2002) The Tol/Pal system function requires an interaction

between the C-terminal domain of TolA and the N-terminal domain of TolB. *Mol*

Microbiol, 44(3), 695-708.

Wu, X., Ren G, Gunning WT 3rd, Weaver DA, Kalinoski AL, Khuder SA, Huntley JF (2016)

FmvB: A *Francisella tularensis* Magnesium-Responsive Outer Membrane Protein that

Plays a Role in Virulence. *PLoS One*, 11(8), e0160977.

Wu, X., Ren G, Huntley JF (2015) Generating Isogenic Deletions (Knockouts) in *Francisella*

tularensis, a Highly-infectious and Fastidious Gram-negative Bacterium. *Bio Protoc*,

5(12), e1500.

1115

1116 **Table 1.** Specific Activity and Substrate Specificity of FTL1678 and FTT0101 Enzymes

Substrate	Specific Activity (nmol/min/mg of protein) [a]	
	FTL1678	FTT0101
A$_2$pm-containing substrates		
GlcNAc-anhydroMurNAc-L-Ala-γ-D-Glu-*meso*-A$_2$pm-D-Ala (TCT)	21.5	29.0
GlcNAc-MurNAc-L-Ala-γ-D-Glu-*meso*- A$_2$pm-D-Ala (PG monomer)	15.6	24.2
TCT dimer	7.7	11.8
MurNAc-L-Ala-γ-D-Glu-*meso*-A2pm-D-Ala	5.8	7.8
UDP-MurNAc-L-Ala-γ-D-Glu-*meso*-A$_2$pm-D-Ala	4.6	6.3
L-Ala-γ-D-Glu-*meso*-A$_2$pm-D-Ala (free tetrapeptide)	3.4	8.0
MurNAc-L-Ala-γ-D-Glu-*meso*-A$_2$pm-D-Ala-D-Ala (pentapeptide)	9.8	6.2
UDP-MurNAc-L-Ala-γ-D-Glu-*meso*-A$_2$pm-D-Ala-D-Ala (pentapeptide)	5.9	6.4
L-Lysine-containing substrates		
MurNAc-L-Ala-γ-D-Glu-L-Lys-D-Ala	1.3	2.1
L-Ala-γ-D-Glu-L-Lys-D-Ala	0.7	2.1
UDP-MurNAc-L-Ala-γ-D-Glu-L-Lys-D-Ala-D-Ala	1.2	1.7
Other		
GlcNAc-MurNAc-L-Ala-γ-D-Glu-*meso*-A$_2$pm(NH$_2$)-D-Ala		Very low (<0.5)
GlcNAc-MurNAc-L-Ala-γ-D-Glu(NH$_2$)-*meso*-A$_2$pm-D-Ala		Not detected [b]
GlcNAc-MurNAc-L-Ala-γ-D-Glu(NH$_2$)-*meso*-A$_2$pm(NH$_2$)-D-Ala		Not detected [b]
Peptidoglycan polymer		Not detected [b]

1117

1118 [a] Standard enzyme assay conditions are described in the Materials and Methods.

1119 [b] Not detected indicates that no release of alanine was detected.

1120   **Table 2.** Sensitivity of WT and Δ*FTL1678* to Antibiotics, Detergents, and Dyes

Compound	Concentration (µg/disk)	Average Zone of Inhibition, mm (mean ± SD)[a]	
		WT	**Δ*FTL1678***
Gentamicin	4	2.50 ± 0.10	0.93 ± 0.15 (R)
Tetracycline	5	1.83 ± 0.06	0.80 ± 0.00 (R)
Chloramphenicol	5	3.37 ± 0.12	1.00 ± 0.00 (R)
Ciprofloxacin	5	4.73 ± 0.23	1.90 ± 0.10 (R)
Ampicillin	200	2.57 ± 0.12	3.53 ± 0.15 (S)
Vancomycin	20	1.27 ± 0.21	2.60 ± 0.10 (S)
Bacitracin	182	1.10 ± 0.10	1.20 ± 0.10
Polymyxin B	100	0.70 ± 0.00	0.70 ± 0.00
Lysozyme	1000	0.70 ± 0.00	1.67 ± 0.06 (S)
Ethidium Bromide	5	2.73 ± 0.15	0.80 ± 0.00 (R)
Triton-X100	750	3.40 ± 0.27	2.63 ± 0.06 (R)
SDS	1000	1.20 ± 0.10	1.80 ± 0.00 (S)
CTAB	50	0.73 ± 0.06	0.80 ± 0.10
CHAPS	50	0.70 ± 0.00	0.70 ± 0.00

1122

[a] (R) indicates Δ*FTL1678* is significantly more resistant than WT by one-way ANOVA ($P<0.05$)

[a] (S) indicates Δ*FTL1678* is significantly more sensitive than WT by one-way ANOVA ($P<0.05$)

Figure 1. Amino acid alignment of bacterial L,D-carboxypeptidases. Clustal Omega amino

acid alignment of *E. coli* LdcA (BAA36050.1), *P. aeruginosa* LdcA (Q9HTZ1), *N. gonorrhoeae*

LdcA (YP_208343.1), *F. tularensis* Type B FTL1678, and *F. tularensis* Type A FTT0101, and *C.*

jejuni Pgp2 (WP_002856863). Percent identities (pid), compared to *E. coli* LdcA, are indicated.

Black shading indicates similar residues. Red shading indicates the catalytic triad.

1130

Figure 2. FTL1678 is OM-associated. Spheroplasting, osmotic lysis, and sucrose density

gradient centrifugation were performed to separate inner membranes (IM) and outer membranes

(OM) from *F. tularensis* ΔFTL1678 *trans*-complemented with a 6× histidine-tagged FTL1678.

Whole-cell lysates (WCL), OM fractions, and IM fractions were separated by SDS-PAGE,

transferred to nitrocellulose, and immunoblotting was performed using antisera specific for the

OM control protein FopA (αFopA), IM control protein SecY (αSecY), or histidine-tagged

FTL1678.

1138

1139 **Figure 3. FTL1678 controls bacterial morphology.** Electron micrograph images of: (A) Wild-

1140 type LVS or (B) Δ*FTL1678* grown in sMHB to OD_{600} of 0.4. Scale bars represent 100 nm; (C)

1141 Outer membrane thickness measurements (nm) were measured for WT and Δ*FTL1678*, n=50; (D)

1142 Cell width measurements (nm) for WT and Δ*FTL1678*, n=175. **** indicates $P<0.0001$.

1143

Figure 4. Deletion of *FTL1678* increases resistance to stressors. WT and Δ*FTL1678* were

grown in 75 ml sMHB at: (A) 37˚C, (B) sMHB at 40˚C, (C) sMHB + 60 µM $CuCl_2$, (D) sMHB +

5 mM H_2O_2, (E) sMHB + 5% NaCl, or (F) sMHB at pH 5.5. Bacteria were grown for 24 hours

and OD_{600} measurements were recorded every 4 hours.

Figure 5. ΔFTL1678 is fully-attenuated in a mouse pulmonary infection model. (A) Groups

of 5 C3H/HeN mice were intranasally-infected with 10^5 CFU of either wild-type WT, Δ*FTL1678*,

or Δ*FTL1678 trans*-complemented with *FTL1678* [FTL1678 compl]. Animal health was

monitored daily through day 21 post-infection. **** indicates *P*<0.0001; (B) Lungs, livers,

spleens, and blood were aseptically harvested from mice infected with 10^4 CFU of either WT or

Δ*FTL1678* on days 2 and 5 post-infection and plated to enumerate bacterial numbers. * indicates

P<0.01; (C) Groups of 5 C3H/HeN mice were intranasally-infected with 10^5 CFU of either LVS,

Δ*FTL1678*, FTL1678 *trans*-complement [FTL1678 compl], or *C. jejuni* Pgp2 *trans*-complement

[Pgp2 compl]. Animal health was monitored through day 21 post-infection. ** indicates *P*<0.01.

1157

1158 **Figure 6. FTL1678 is required for *F. tularensis* replication in macrophages.** (A) J774A.1

1159 macrophages or mouse bone marrow-derived macrophages (mBMDMs) were infected with WT

1160 or Δ*FTL1678* at an MOI of 100:1 and bacterial numbers were enumerated at entry (0 h), 6 h, and

1161 24 h post-infection. (B) Fold change in bacterial numbers from 6 to 24 h post-infection was

1162 calculated. * indicates *P*< 0.01.

1163

Figure 7. Δ*FTL1678* protects against fully-virulent Type A *F. tularensis* SchuS4. (A) Groups

of 5 C3H/HeN mice were intranasally infected with either 10^5 CFU WT or 10^5, 10^7, or 10^9 CFU

Δ*FTL1678*. On day 29 post-infection, mice were boosted with 10^9 CFU Δ*FTL1678* and animal

health was monitored daily through day 50 post-infection. *** *P*<0.001; (B) Mice from A were

intranasally-challenged with 120 CFU of wild-type SchuS4 [BSL3; 6× LD50]. Animal health was

monitored daily through day 26 post-infection. * indicates *P*<0.001.

1170

1171   **Figure 8. Δ*FTL1678* does not induce tissue damage.** (A) Hematoxylin and eosin (H&E)-

1172   stained lungs, livers, and spleens were examined from either uninfected, *F. tularensis* WT LVS-,

1173   or Δ*FTL1678*-infected mice at 10× objective. (B) Tissues were graded on a scale of 0 to 4, with 4

1174   being the most severe. * indicates *P*<0.05.

1175

Figure 9. Model of *F. tularensis* peptidoglycan synthesis and recycling pathways.

Bioinformatic analyses were used to predict proteins that may be involved in peptidoglycan

synthesis and recycling in *F. tularensis*. *F. tularensis* LVS gene locus tags are indicated, with *E.*

coli or Gram-negative ortholog protein names. OM, outer membrane. IM, inner membrane.

GlcNAc, N-Acetylglucosamine. MurNAc, N-Acetylmuramic acid. Pal, OM-localized

peptidoglycan-associated lipoprotein. TolB, periplasmic protein that interacts with Pal and

peptidoglycan. HMM PBP, high molecular weight penicillin binding protein. LMM PBP, low

molecular weight penicillin binding protein.

1184

Supporting Information

Table S1. Bioinformatic analyses of FTL1678 localization

Table S2. Endopeptidase activity of FTL1678

Table S3. Sensitivity of WT *F. tularensis* SchuS4 and Δ*FTT0101* to antibiotics, detergents, and dyes

Table S4. Specific activity of FTL1678, FTL1678 double mutants, and FTL1678 triple mutants to the TCT monomer

Table S5. Sensitivity of WT LVS, ΔFTL1678, FTL1678 trans-complement, and Pgp2 trans-complement to antibiotics, detergents, and dyes

Table S6. Bacterial strains and plasmids used in this study

Table S7. Primers used in this study

Figure S1. FTL1678 contains a putative L,D-carboxypeptidase domain. NCBI Conserved Domain search results for *F. tularensis* FTL1678.

Figure S2. F. tularensis TolB is OM-localized. Spheroplasting, osmotic lysis, and sucrose density gradient centrifugation were performed to separate inner membranes (IM) and outer

1209   membranes (OM) from *F. tularensis* Δ*FTL1678 trans*-complemented with a 6x histidine-tagged

1210   FTL1678. Whole-cell lysates (WCL), OM fractions, and IM fractions were separated by SDS-

1211   PAGE, transferred to nitrocellulose, and immunoblotting was performed using antisera specific

1212   for the periplasmic protein TolB (αTolB).

1213

1214   **Figure S3. ΔFTL1678 does not have a growth defect.** WT and ΔFTL1678 were grown in

1215   sMHB for 24 hours at 37˚C. Samples were taken every 4 hours for (A) OD600 measurements

1216   and (B) CFU enumeration following serial-dilution and plating.

1217

Figure S4. ΔFTL1678 has septation defects. Transmission electron micrograph images of

ΔFTL1678 showing aberrant septal formation and reduced ability to separate cells. Images taken

at: (A) 49,000×, scale bar represents 200 nm; and (B) 6,800×, scale bar represents 2 μm. In

(A), red arrows point to formed septa that have not separated in ΔFTL1678 and white arrows

point to new septa that are forming in ΔFTL1678. Experiments were performed twice to confirm

reproducibility, with two bacterial preparations fixed, stained, embedded, sectioned, and

visualized per experiment. Representative images shown.

1225

1226   **Figure S5. FTT0101 is not required for *F. tularensis* Type A strain SchuS4 virulence.**

1227   C3H/HeN mice were intranasally infected with either 80 CFU SchuS4 (n=3 mice) or 12 CFU

1228   Δ*FTT0101* (n=5 mice).

1229

Figure S6. LdcA catalytic triad is not essential for *F. tularensis* virulence. Groups of 5

C3H/HeN mice were intranasally-infected with 10^5 CFU of either *F. tularensis* WT LVS,

Δ*FTL1678*, Δ*FTL1678 trans*-complemented with FTL1678 (FTL1678 compl), Δ*FTL1678 trans*-

complemented with S134A (S134A), Δ*FTL1678 trans*-complemented with E239A (E239A), or

Δ*FTL1678 trans*-complemented with H308A (H308A). Animal health was monitored daily

through day 21 post-infection. ** *P*<0.01.

1236

Figure S7. FTL1678 *trans*-complement and *C. jejuni* Pgp2 *trans*-complement restore *F.*

***tularensis* phenotype.** Transmission electron micrograph images of: (A) Δ*FTL1678 trans*-

complemented with *FTL1678* [FTL1678 compl] or (B) Δ*FTL1678 trans*-complemented with *C.*

jejuni pgp2 [Pgp2 compl]. Bacteria were grown in sMHB to OD_{600} of 0.4. Scale bars represent

100 nm. (C) Outer membrane thickness and (D) cell width of FTL1678 compl and Pgp2 compl

were compared to WT LVS and D*FTL1678.* **** indicates *P*<0.0001.

1243

Figure S8. FTL1678 *trans*-complement and *C. jejuni* Pgp2 *trans*-complement exhibit

similar phenotypes to stressors as WT *F. tularensis*. WT LVS [WT], Δ*FTL1678,* Δ*FTL1678*

trans-complemented with *FTL1678* [FTL1678 compl], or Δ*FTL1678 trans*-complemented with

C. jejuni pgp2 [Pgp2 compl] were grown in either: (A) sMHB; (B) sMHB with 5 mM H_2O_2; (C)

sMHB with 5% NaCl; or (D) sMHB at pH 5.5. Cultures were incubated for 24 hours and OD_{600}

measurements were recorded every 4 hours.

```
              pid   1                                                                           80
Ec  LdcA 100.0%  ----------------------------------------------------------MSLFHLIAPSGYCIKQHAALRG
Pa  LdcA  23.0%  -----------------------------------------------MTSRPSSDQTWQPIDGRVALIAPASAIATDVLEAT
Ng  LdcA  30.3%  MTEPTSRRRFLKTCTAAGAGLLQACGTSATSVPPLPSSHSVVKARTVPLQTPRRQSSDGNLLRVVASSGFAEDTNRVNTA
FTL1678   17.6%  --------------------------MLLKNNYLVSIILVVLIMIVTKSFACAATDYNKVALINVS-TQYYPNDIKQA
FTT0101   17.6%  --------------------------MLLKNNYLVSIILVVLIMIVTKSFACAATDYNKVALINVS-TQYYPNDIKQA
Cj  Pgp2   6.3%  ------------------------------MLKRLALLITLSSIMLHASDLVKIYLNQGLDAVGVAIEKELTQKD

              pid  81                                                                          160
Ec  LdcA 100.0%  IQRLTDAGHQVNNVEVIARRCERFAGTETERLEDLNSLAR-LTTPNTIVLAVRGGYG----ASRLLADIDWQALVARQQH
Pa  LdcA  23.0%  LRQLEVHGVDYHLGRHVEARYRYLAGTVEQRLEDLHNAFD--MPDITAVWCLRGGYG----CGQLLPGLDWGRLQAASPR
Ng  LdcA  30.3%  LTRLYNAGFTVTNQQAGSRRFQRFAGTDAQRAADFQEVASGRVATPKVLMGLRGGYG----AARILPHIDFASLGARM
FTL1678   17.6%  EKALKDTGYNTT-YKYLDIYPSDFGYSNPDSIRAKILLDALLDKNIDIIWFLKGGG----AFNLLPYL-YDHINELKKA
FTT0101   17.6%  EKALKDTGYNTT-YKYLDIYPSDFGYSNPDSIRAKILLDALLDKNIDIIWFLKGGG----AFNLLPYL-YDHINELKKA
Cj  Pgp2   6.3%  FWLSEIGDKNISLGYYDDNVAIVLTNKTDKILRVYSYEDGKIRKDFEQKEIITGLMGDKKIEGDLKTPVGFYELGRKFNP

              pid 161                                                                          240
Ec  LdcA 100.0%  DPLLICGHSDFTAIQCGLLAHGNVITFSGPMLVANFGADELNAFTEHHFWIALRNETFTIEWQ-----------GEGPTC
Pa  LdcA  23.0%  P---LIGFSDISVLLSAFFRHGLPAIHGPVATGLGLSPLSAPREQQERIASLASVSRLLAGIDHELP----VQHLGGHKQ
Ng  LdcA  30.3%  HGTLFFGFSDVCAVQLALLAKGNMMSFAGPMAYSDFGKPAPGAFTMDAFIKGATONRLTVDVPY----------IQRADV
FTL1678   17.6%  KPKILVGFSDVTAIHFFVNNVLGWKSLHGVVAAYNKNAYSSQKIEKIRINDLERIPNITEIINNGISYDKLMPMNKMAYN
FTT0101   17.6%  KPKILVGFSDVTAIHFFVNNVLGWKSLHGVVAAYNKNAYSSQKIGKIRINDLERIPNITEIINNGISYDKLMPMNKMAYN
Cj  Pgp2   6.3%  GDPYYGPFAFATTYPNLLDKVQGKTGGGIWIHGYPLDGSRLDEFKTRGCIALFNNN----------------LEKFAQV

              pid 241                                                                          320
Ec  LdcA 100.0%  RAEGTLWGGNLAMLISLIGTP-WMPKIENGILVLEDINEHPFRVERMLLQLYHAGILPRQKAIILGSFSGSTP-NDYDAG
Pa  LdcA  23.0%  RVEGALIGGNLTALACMAGTLGGLHAPAGSILVLEDVGEPYYRLERSLWQLLESIDARQLGAICLGSFTDCPR---KEVA
Ng  LdcA  30.3%  ETEGTLWGGNLSVLASLAGTP-YMPDIDGGILFLEDVGEQPYRIERMLNTLYLSGILGKQRAIVFGDFRMEKIRDLYDSS
FTL1678   17.6%  GTDGSIVGGNMTLIYSYFSTV-YQQDISTKILFLEDTGISFRQLDRSLHQLLYLPENKKPEAIIFGQFYPLDP--TDDQR
FTT0101   17.6%  GIDGSIVGGNMTLIYSYFSTV-YQQDISTKILFLEDTGISFRQLDRSLHQLLYLPENKKPEAIIFGQFYPLDP--TDDQR
Cj  Pgp2   6.3%  VQDKKVFVMTEEKEKIRAKKDQIASLLADLFTWKLAWTNSDTNTYLSFYDEQEFKRFDKMKFEQFASMKKSIFSRKEDKK

              pid 321                                                                          400
Ec  LdcA 100.0%  YNLESVYAFLRSRLSIPLITGLDFGHEQRTVTLPLGAHA-ILNNTREGTQLTISGHPVLKM--------------------
Pa  LdcA  23.0%  HSLERIFGEYAAAIEVPLYHHLPSGHGAQNRAWPYGKTAVLEGNRLRW-----------------------
Ng  LdcA  30.3%  YDFSAVAKHISRTAKIPVLTGFPFGHIADKITFPLGAHTRIRMNGNGGYSVAFEGYPTLDASALTLDTLLPPPDLPIFPE
FTL1678   17.6%  LIYKTVIKKFAKTFNRPVYYFPFIGHGQYNKPLLLGVTSNIKCSKETIFCTLKQK------------------------
FTT0101   17.6%  LIYKTVIKKFAKTFNRPVYYFPFIGHGQYNKPLLLGVTSNIKCSKETIFCTLKQK------------------------
Cj  Pgp2   6.3%  IKFSDINISPYPNLENETMYRISFYEDYYTKNYQFRGDKILYVKIDSKGKMKILAEQ-----------------------

              pid 401          408
Ec  LdcA 100.0%  --------
Pa  LdcA  23.0%  --------
Ng  LdcA  30.3%  SGVADISE
FTL1678   17.6%  --------
FTT0101   17.6%  --------
Cj  Pgp2   6.3%  --------
```

A.

B.

C.

D.

A.

Percent survival vs Days Post-Infection

10⁹ CFU Boost

***

- LVS
- Δ*FTL1678*, 10⁹ CFU
- Δ*FTL1678*, 10⁷ CFU
- Δ*FTL1678*, 10⁵ CFU

B.

Percent survival vs Days Post-Infection

*
*
*