Article, discoveries

Evolution of multiple CDK1 paralogs towards specializations in late cell cycle events

in a lineage with fast developing planktonic embryos.

Xiaofei Ma[1], Jan Inge Øvrebø[1,2#], Eric M Thompson[1,2 *]

[1]Sars International Centre, and [2]Dept. of Biological Sciences, University of Bergen, Bergen, Norway

[#]Present address: Huntsman Cancer Institute, University of Utah, USA

*Correspondence to: Eric.Thompson@uib.no, ORCID 0000-0002-3756-9036

Running Title: Evolution of CDK1 paralog specializations

Abbreviations: CDK, Cyclin Dependent Kinase; DoS, Determinant of Specificity; RNAi, RNA

interference; NEBD, Nuclear Envelope Breakdown;

Key words: PSTAIRE helix; Cyclin B; Tunicate; Coenocyst; Oogenesis; Embryogenesis; Meiosis;

X-linkage; Gene duplication

Abstract

The CDK1 kinase is required for most eukaryotic cell cycles. The active site is generated by essential,

structural elements, among which the PSTAIRE motif in the critical αC-helix, is universally

conserved in metazoans. The CDK2 kinase, sharing the PSTAIRE, arose early in metazoan evolution

and permitted subdivision of tasks along the S-M-phase axis. The marine planktonic chordate,

Oikopleura dioica, is the only metazoan known to possess more than a single CDK1 ortholog, and

all of its 5 paralogs show sequence divergences in the PSTAIRE. Through assessing CDK1 gene

duplications in the rapidly evolving oikopleurid lineage, we show that the CDK1 activation loop

substrate binding platform, ATP entrance site, hinge region, and main Cyclin binding interface, have

all diversified under positive selection. Three of the 5 CDK1 paralogs, a, b and d, are required for

embryonic divisions and knockdown phenotypes indicate further subdivision of functions along the

S-M-phase axis. In parallel to the CDK1 gene amplification there has also been amplification in the

Cyclin B complement. Among these, the CDK1d:Cyclin Ba pairing is required for completion of

oogenic meiosis and early embryonic divisions and shows evidence of coevolution of an exclusive

interaction. In an intriguing twist on the general rule that it is Cyclin B oscillations on a background

of stable CDK1 levels that regulate M-phase MPF activity, it is the CDK1d protein levels that

oscillate, rather than Cyclin Ba levels, to drive rapid, early embryonic cell cycles. Strikingly, the

modified PSTAIRE of odCDK1d shows convergence over great evolutionary distance with plant

CDKB, and in both *O. dioica*, and plants, these variants exhibit increased specialization to M-phase.

Introduction

The complexity of the eukaryotic cell and its larger genome necessitates greater control over the organization of events during cellular division than for prokaryotic life. The eukaryotic protein kinase, CDK1, whose activity depends on unstable Cyclin regulatory subunits, has key roles in orderly triggering S and M phases from yeast to mammals (Santamaria et al. 2007), by sequential, regulatory phosphorylation of hundreds of substrate proteins. CDK activity levels progressively surpass response thresholds, to trigger serial events in S and M phases (Swaffer et al. 2016). Quantitative changes in CDK activity function as a coarse organizing mechanism for cell cycle transitions, with lower CDK activity resulting in higher affinity S-phase substrate phosphorylation, to carry out DNA replication, and higher activity driving mitotic entry via the phosphorylation of lower affinity mitotic substrates. CDK1 protein levels are constant during the cell cycle (Arooz et al. 2000), and are activated at each phase of the cell cycle by binding of stage-specific cyclins. The intrinsic activity of Cyclin–CDK1 complexes increases in correlation with the appearance of particular cyclins in the cell cycle (Koivomagi et al. 2001; Ord and Loog 2019). G1/S cyclins generate CDK complexes with the lowest activity, which manifests in the lowest affinities (k_m) and the lowest catalytic activities (k_{cat}) toward substrates, whereas S-, G2-, and M-phase Cyclins form CDK complexes with higher affinities and catalytic activities towards substrates. The lower CDK activities generated by S- versus M-phase Cyclins also means that S-phase Cyclins do not achieve sufficient CDK1 activity levels to initiate mitosis, helping to ensure resolution of S- and M-phases. Compared to single-celled yeast, metazoans have also evolved CDK2, with lower intrinsic activity than CDK1 in the regulation of S-phase (Ord et al. 2019; Holmes and Solomon 1996). Although metazoan CDK1 has become more specialized to M-phase functions it still retains indispensable, ancestral, late S-phase functions (Hochegger et al. 2007; Katsuno et al. 2009; Farrell et al. 2012; Seller and O'Farell 2018; Szmyd et al. 2019).

In addition to modulating CDK activity, the Cyclins are also implicated in specifying CDK substrate targets. Cyclins bind different substrate docking motifs: RxF for S Cyclins and LxF for budding yeast M Cyclins, and these Cyclin docking pockets increase specificity to target subsets to further fine-tune the CDK threshold ladder (Koivomagi et al. 2011; Ord et al. 2019). Most CDK substrates contain multiple target sites clustered in disordered regions (Holt et al. 2009), and phosphorylation cascades along these sites are shaped by precisely oriented docking interactions mediated by Cks1, the phospho-adaptor subunit of Cdk1 (Koivomagi et al. 2011). Distances between phosphorylation sites and docking sites are critical for both Cyclin and Cks docking. The composition of phosphorylation sites with respect to serine versus threonine, and surrounding residues, combined with positioning of Cyclin and Cks docking motifs, creates a unique barcode on each substrate. The Cyclin−CDK−Cks1 complexes read the barcodes and assign execution of CDK-triggered switches to specified time points during the cell cycle (Ord et al. 2019).

Canonical cell cycle regulation can be modified during development and growth. This is true of rapid mitotic cell cycles observed during early embryogenesis in a wide range of phyla and in the promotion of growth through endocycles in a number of organisms (Edgar and Orr-Weaver 2011). Externally developing, nutrient-rich, non-feeding, embryos are at risk of predation, and rapid development is a means of mitigating this risk (Strathmann et al. 2002; Staver and Strathmann 2002). Many marine planktonic larvae also continue fast development to first swimming in order to enhance survival (Staver and Strathmann 2002). The pelagic, tunicate, *Oikopleura dioica*, exhibits very rapid embryonic and larval development, undergoing metamorphosis

to a feeding juvenile within 12 h at 15 °C (Bouquet et al. 2009). Very rapid growth and extensive modulation

of reproductive output over several orders of magnitude are then achieved through deployment of a variety of

endoreduplicative cell cycle variants (Ganot and Thompson 2002; Ganot et al. 2007). These adaptations allow

the organism to rapidly adjust population levels in response to algal blooms, during an extremely short chordate

life cycle (6 days at 15 °C). Accompanying this life history strategy is considerable modification of the

organism's molecular cell cycle regulatory complement (Campsteijin et al. 2012), including gene duplications

to generate multiple CDK1 and Cyclin B paralogs.

All metazoans examined to date possess a single CDK1 ortholog that exhibits a highly conserved PSTAIRE

motif in the critical αC-helix structural element located within the N-lobe of CDK1 and CDK2. This element

is an important interface in docking interactions with respective Cyclin partners and in contributing to

formation of the kinase active site (Wood et al. 2019). *O. dioica*, aligns with its metazoan relatives in possessing

a CDK2 with a conserved PSTAIRE motif, but deviates distinctively from other metazoans in that it has

evolved 5 CDK1 paralogs, all of which show deviations in the critical PSTAIRE motif. Here, we have

examined how these paralogs have arisen in the appendicularian lineage and to what extent they exhibit

functional specializations along the CDK2-CDK1, S- to M-phase, cell cycle, regulatory axis. We show that the

CDK1 substrate binding platform, activation loop, ATP entrance site, hinge region, and the main Cyclin

binding interface, have all diversified under positive selection. In meiosis and early embryogenesis, CDK1

paralogs were sequentially activated across the cell cycle. Combinatorial knockdowns revealed collaboration

of CDK2, CDK1a, CDK1b and CDK1d along the S- to M-phase axis in oogenic meiosis and early embryonic

divisions. Interestingly, the Cyclin Ba and CDK1d paralogs are located in proximity on the recently evolved

O. dioica X chromosome, were co-expressed specifically during oogenic meiosis and early embryogenesis,

and were required for these processes. Modifications of critical residues in the Cyclin B:CDK1 interface and

the salt bridge, suggest that CDK1d and Cyclin Ba may have coevolved an exclusive interaction that does not

interfere with the more canonical CDK1a/b/c:CycBc interactions in regulating M-phases during early

developmental events. It is generally universal in metazoan mitotic cycles that CDK1 protein levels remain

relatively constant throughout the cell cycle whereas Cyclin B levels oscillate and are up-regulated during M-

phase. In contrast, we reveal a novel regulatory oscillation of levels of the CDK1d paralog, as opposed to its

Cyclin Ba interacting partner, that drive early, very rapid embryonic cell cycles in this planktonic chordate.

Results

Episodic positive selection of CDK1 orthologs after duplication

Metazoans possess a single CDK1, with the exception of *Oikopleura dioica*, where 5 CDK1 paralogs have

been identified (Campstejin et al. 2012). To trace the origins of this amplification we analyzed genomes and

transcriptomes of five appendicularian species in the *Oikopleura* and *Fritillaria* genera (Naville et al. 2019).

CDK1 is amplified in all five species. *CDK1a, b and c* were uniformly present, whereas *CDK1d* and *e* appear

to have arisen from *CDK1c* specifically in *O. dioica*, the only known dioecious appendicularian (fig. 1A).

Interspecies clustering of *CDK1* paralogs and interspecies conservation of their intron-exon structures

(supplementary fig. S1) suggest that *CDK1* amplification arose through early DNA-based duplication events

in the *Oikopleura* lineage. Based on phylogenetic analyses, *CDK1a* duplication gave rise to *CDK1b* that was

subsequently duplicated to generate *CDK1c,* which then gave rise to the *O. dioica* specific, *odCDK1d*. This

duplication gradient pattern, and young paralogs-biased duplication, have not previously been reported in
metazoans. *O. albicans CDK1a*, *b* and *c* locate on the same chromosome. *O. dioica CDK1a* and *b* locate on the
same autosome whereas *CDK1c* is present on a different autosome that also contains *CDK1e* (fig. 1C). *CDK1d*
is found on the X chromosome where, at a distance of 0.1 MB (Japan population), the *CycBa* locus is also
located. Following the classification system for paralog subtypes (Sonnhammer and Koonin 2002), *Oikopleura*
genes in the *CDK1a* clade are orthologs, and *odCDK1a* and *b* are outparalogs to each other when comparing
O. dioica with *O. albicans*, since CDK1 duplication happened before *O. dioica* and *O. albicans* speciation. *O.*
dioica CDK1c, d and *e* are inparalogs to each other when comparing *O. dioica* with *O. albicans,* since CDK1c
duplication happened after *O. dioica* and *O. albicans* speciation.

Both the new Nei-Gojobori method (Zhang et al. 1998) and Codon-based models (Yang 2007) were applied
to analysis of *Oikopleura* CDK1s. After the 3 rounds of gene duplication, the lineage subtending *CDK1a* (fig.
1A) experienced positive selection (bn/bs = 2.48, P=0.0386, dN/dS(ω) = 41.01, p = 0.081, P = 0.0481), as did
CDK1b (bn/bs = 4.72, P = 0.0305, dN/dS(ω) = 28.14, p = 0.046, P = 0.003) and *CDK1c* (bn/bs = infinite, P =
0.033, dN/dS(ω) = 999, p = 0.066, P=0.002) (fig. 1A, supplementary table S1). To better understand positively
selected sites identified by Codon-based models, we aligned CDK1 sequences from 15 metazoan phyla, as well
as Cdc28 from budding yeast. Sites conserved in metazoan sequences, but substituted in *Oikopleura* CDK1
orthologs, were mapped to human CDK1 and CDK2 crystal structures (supplementary figs. S2 and 3).
Determinants of kinase specificities (DoS) identified by machine learning (Creixell et al. 2015) were aligned
and the DoS that are conserved in metazoan CDK1 orthologs but substituted in *Oikopleura* CDK1 sequences
were mapped to human CDK1-Cyclin B1 crystal structures. During kinase evolution, the critical catalytic
residues, regulatory hydrophobic spine and catalytic hydrophobic spine have been conserved. The
diversification of the substrate binding platform-activation segment, and indirect ATP binding sites, drive
diversification of substrate specificity and substrate phosphorylation efficiency. *Oikopleura* CDK1s have the
same active site as human *CDK1* and yeast cdc28, including the Mg^{2+}-binding loop (DFG), important residues
in the catalytic loop, the catalytic hydrophobic spine, the regulatory hydrophobic spine and the salt bridge
(Wood et al. 2019) (supplementary figs. S2 and 3). The bn/bs values below 1, observed among CDK1
sequences (fig. 1A), suggest that these orthologs are functional kinases under purifying selection. Conservation
in the DFG+1 residues (Leu) in the 17 *Oikopleura* CDK1 sequences, suggests that they share with human
CDK1, a preference for serine over threonine as a phosphor-acceptor residue in substrates (Chen et al. 2014).
However, the ATP binding sites, substrate binding sites and Cyclin binding interfaces have diversified in
Oikopleura CDK1 paralogs.

Substitutions in ATP binding sites

In the protein kinase family, the ATP binding site is very conserved, but adjacent areas not occupied by ATP
are more variable, particularly in the hinge region (E81-K89, human CDK1 numbering). These sites modulate
ATP affinity. This region may determine the different catalytic efficiencies of CDK1 and CDK2, as
substitutions in the hinge of CDK2, that mimic amino acids in this region of CDK1 (substitute N84, Q85 in
CDK2 to 84S, 85M in CDK1), increase both ATP affinity (k_m ATP) and turn over (k_{cat}) (Echalier et al. 2012).
In the opposite direction, a K89D/E mutation in the CDK2 sequence, reduces k_{cat}. *Oikopleura* CDK1 paralogs

have diversified in the hinge region. CDK1c paralogs exhibit an S84F/Y substitution whereas this site is conserved in CDK1a paralogs compared to the metazoan consensus (supplementary fig. S2). The M85 site is a DoS, and a M85C substitution occurred in CDK1a of *O. vanhoeffeni*, *O. albicans* and *O. dioica*, after they split from *O. longicauda*. This substitution has been positively selected (supplementary table S1), possibly driving diversification of ATP affinity among CDK1 paralogs following their duplication. *Oikopleura* CDK1a and b orthologs have substitutions at K89, a residue that is conserved in CDK1 and CDK2 from yeast to metazoans (supplementary fig. S2). This raises the possibility that CDK1a and b may have lower ATP affinity and substrate phosphorylation efficiency compared to CDK1c, d and e paralogs. Indeed, as shown later, we found that CDK1d peaks in M phase whereas CDK1a was present earlier in interphase. The CDK1 L83 residue is also conserved form yeast to human, with the amide of L83 forming a hydrogen bond with the heteroaromatic adenine ring of ATP (Bao et al. 2011). This residue is conserved in CDK1a and c, but has been modified to L83M in the hinge region of all *Oikopleura* CDK1b orthologs. Thus, CDK1 paralogs have different signatures in their hinge regions, suggesting diversification in their ATP affinities and catalytic efficiencies.

The conserved DoS, P61, is located in the N-lobe, in the loop connecting the PSTAIRE helix and β4 of human CDK1 (supplementary fig. S2). Positive selection of a P61E substitution in all *Oikopleura* CDK1a orthologs, suggests that positive selection occurred after the initial CDK1 duplication but prior to speciation. This asymmetric adaptive evolution of an evolutionary conserved DoS in one of the duplicates, and conservation of this substitution within CDK1a orthologs, could be best explained by escape from adaptive conflicts after duplication (Des Marais and Rausher 2008).

Substitution in the activation segment

The activation segment forms an integral part of the substrate binding groove, and diversity in this loop contributes to the wide range of substrate specificities and affinities of protein kinases (Nolen et al. 2004). The structure of CDK1 suggests a mechanism through which activation loop flexibility, embedded in a more flexible CDK1 fold, allows CDK1 to accommodate a more diverse substrate set than CDK2 (Kriwacki et al. 1996; Brown et al. 2015). These plastic properties may also contribute to its ability to partner non-cognate cyclins in the absence of other CDKs to drive a complete metazoan cell cycle[1]. In *Oikopleura* CDK1 paralogs, sites in the activation segment that bind to substrate, P0 (L149, T166), P+1 (V164, V165) and P+3 (T161), are the same as in yeast and human CDK1 (supplementary fig. S2). However, several conserved sites in the activation segment have been dramatically modified in *Oikopleura CDK1* paralogs, particularly in *O. dioica CDK1c, d* and *e* inparalogs. G154 is located at the tip of the human CDK1 activation segment where it is important in forming the β turn in the short β hairpin structure when CDK1 binds to Cyclin B1 (Brown et al. 2015) (supplementary fig. S3). Human CDK2 cannot form this hairpin structure (Jeffrey et al. 1995), and this is a major difference in the activation segment between CDK1 and CDK2. The G154 site is conserved in yeast and metazoans, but has diversified in *Oikopleura* CDK1 paralogs. A G154N substitution occurred in *Oikopleura* CDK1a, G154S in *Oikopleura* CDK1b, and G154K in *O. dioica* CDK1c, d and e (supplementary fig. S2). This site is also identified as having experienced positive selection (supplementary table S1), suggesting adaptive diversification of the activation segment by disrupting the β-turn though substitution with large side chain residues. Immediately downstream of this substitution, *O. dioica* CDK1c, d and e all have an I157F substitution, whereas CDK1c paralogs from other *Oikopleura* species have an I157M substitution. *O.*

dioica CDK1c, d and e also exhibit substitutions at theV159 and Y160 sites, two residues immediately upstream

of the important, regulatory phospho T161 site. The V159 site is also a DoS, conserved from yeast to humans.

Finally, in human CDK1, L167 (conserved from yeast to human) is adjacent to the substrate P-2 residue (Brown

et al. 1999; Bao et al. 2011) (supplementary fig. S4). The L167M substitution found in all CDK1b and c

orthologs, presents a larger side chain, which may affect affinity towards P-2 residues.

Substitutions in the PSTAIRE helix

The CDK1 PSTAIRE helix is invariant in metazoans and yeast. Mutation screens in yeast *cdc28* revealed that

the exact PSTAIRE helix sequence (PSTAIREISLLKE) is required for cell division, and any mutations in this

domain are lethal (MacNeill et al. 1991; MacNeill and Nurse 1993; Levine et al. 1999; Ahn et al. 2001; Child

et al. 2010). The A48V substitution in cdc2 in *S. pombe* is dominant negative, as the mutant cdc2 could bind

Cyclin but could not be activated (Fleig et al. 1992), whereas an A48T substitution in *S. cerevisiae cdc28*

inhibits CLB2 association (Ahn et al. 2001). Surprisingly, the PSTAIRE helix is modified in all *Oikopleura*

CDK1 paralogs (fig. 1A) whereas *Oikopleura* CDK2s retain the canonical PSTAIRE sequence. All of the 17

identified *Oikopleura* CDK1 sequences share the A48S substitution (supplementary figs. S1 and S2). The S48

residues in *Oikopleura* CDK1 sequences are encoded by two disjoint codon sets, TCN and AGY, which cannot

be interconverted by a single nucleotide mutation. Accordingly, switches between serine codons from the two

sets can occur either directly, by simultaneous double (tandem) mutation (TC > AG), or indirectly, via two

consecutive single-nucleotide substitutions (TC > AC > AG or TC > TG > AG). This is consistent with strong

positive selection (Averof et al. 2000; Rogozin et al. 2016) at these sites in CDK1 sequences within the entire

Oikopleura genus. Interestingly, coincident with the A48S substitution, codons for S48 in all CDK1b and c

paralogs, except olCDK1c, have phase 2 intron insertions (supplementary fig. S1). This intron insertion is

specific to *Oikopleura*, and is not found in any other metazoan CDK1.

O. dioica CDK1b and d (and *O. longicauda* CDK1b) have an additional S46P substitution in the PSTAIRE

helix whereas *O. dioica* CDK1c and e have an S46A substitution at this residue. The double proline predicts

that the odCDK1b and d PSTAIRE helices would be shorter and exhibit an altered orientation compared to

human CDK1. The *O. dioica* CDK1a, b and d versions of the PSTAIRE helix complemented yeast *cdc28*

temperature sensitive mutants, whereas *O. dioica* CDK1c and e, both possessing the S46A substitution, did not

(fig. 1D). *O. dioica* CDK1c, d and e inparalogs also share double S53C and L54T/S substitutions in the

PSTAIRE helix (supplementary fig. S2). Human CDK1 S53 forms hydrogen bonds with Cyclin B1 L296 and

G297 amide groups (Brown et al. 2015) (located at the C-terminal loop connected to the α5 helix)

(supplementary fig. S3). Substituting serine at this site with cysteine may reduce CDK1c inparalog interaction

interfaces with Cyclin B.

In summary, critical elements for catalysis are conserved in *Oikopleura* CDK1 paralogs, but they diverge

in binding sites for ATP and substrates, suggesting qualitative differences in catalysis. Therefore, we set out to

determine expression profiles and evaluate knockdowns of *O. dioica* CDK1 paralogs to characterize their

individual functions in meiosis and early embryonic cell cycles.

Transcription of *CDK1d* and *cycBa* is restricted to oogenesis

CDK1a, b, c and *cycBc* are generally present in the *Oikopleura* genus (fig. 1A and B). In *O. dioica*, they were

broadly expressed, in a sexually unbiased manner, in embryos, juveniles, and mature animals (Fig. 2). In

contrast, *CDK1d* and *e*, and *cycBa* are thus far, specific to *O. dioica*, and were transcribed during oogenesis.

Their transcripts persisted during early embryogenesis (figs. 2 and 6B). *CycBa* was not expressed in juveniles

or testes. Interestingly, *O. dioica CDK1d* has different transcription start sites (TSS) in testes and ovaries. The

ovarian *CDK1d* TSS occurs before the ATG start codon located in the first exon. In testis, a sharp TSS located

within the first intron yields nonsense transcripts (fig. 1E). Western blots confirmed that the CDK1d protein

was present in ovaries but absent in testes (fig. 2B). *CDK1d* mRNAs began to degrade after fertilization and

dropped to very low levels at 2.5 h post-fertilization (fig. 6B). Given these differential expression patterns, we

then undertook to knockdown the five CDK1 paralogs and 2 Cyclin Bs in *O. dioica* individually, or

simultaneously, to explore functions of the expanded CDK1-Cyclin B complement.

CDK1a, b and d collaborate in embryonic divisions and a conserved PSTAIRE motif is not required

The five odCDK1 paralogs and odCDK2 were each targeted for knockdown by RNAi. After injections of

dsRNA in gonads of day 4 or 5 animals, oocytes were collected and processed to verify specific knockdown

by RT-qPCR, and western blots. Successfully knocked down oocytes were exposed to wild type sperm and

embryonic divisions were examined. Previous reports of CDK1a RNAi at day 4 demonstrated that this impaired

vitellogenesis and meiosis resumption[46,47]. CDK1a RNAi at day 5 avoided these defects and injected animals

produced normal size oocytes. However, embryonic divisions were delayed and they failed to hatch (fig. 3A).

When CDK1b was knocked down, embryonic cell cycles were also delayed and embryos arrested at

gastrulation, corresponding to 2.5 h post-fertilization of normal embryos (fig. 3B). Knockdown of CDK1d

yielded 3 phenotypes: a complete failure to emit polar bodies, arrest after polar body emission, or abnormal

divisions after polar body emission (fig. 3C-G). These phenotypes reflect roles of odCDK1d in nuclear

envelope breakdown (NEBD) (Ovrebo et al. 2015) and spindle assembly in meiosis I, for meiosis completion

(Feng and Thompson 2018), and subsequent mitotic divisions. No significant differences were observed in the

proportions of normal hatching embryos between wild type, and any of the *CDK1c, e,* or *c+e* RNAi knockdown

embryos (supplementary fig. S5). CDK1c and e have very similar PSTAIRE helices (PATSI/VRE, fig. 1A),

they failed to rescue yeast *cdc28* mutants, and they were also dispensable for embryonic divisions.

CDK2 is dispensable for early mouse early embryogenesis and mitotic cycles in cultured vertebrate cells

(Santamaria et al. 2007). Consistent with these observations, *odCDK2* RNAi embryos divided normally and

hatched with correct timing (fig. 4A). CDK2 is the only CDK in *Oikopleura* that has a canonical PSTAIRE

helix. This indicates that the exact sequence in the conserved metazoan CDK1, PSTAIRE helix, is not required

for *Oikopleura* embryonic divisions. After emission of polar bodies, *CDK1a + CDK2* RNAi zygotes arrested

prior to pronuclear fusion (fig. 4B). The synergistic defects during the cell cycle caused by *CDK1a+CDK2*

RNAi suggests CDK1 and CDK2 have overlapping roles in interphase. Since *CDK2* RNAi embryos divided

normally, defects arising from *CDK1a+CDK2* RNAi suggest *CDK1a* could complement CDK2's roles in

interphase during embryonic divisions. Immunostaining of tadpoles developed from oocytes injected with

CDK1a-GFP capped mRNA, showed CDK1a located to the nucleus during interphase (supplementary fig. S6).

CDK1a+b RNAi embryos arrested at the 4th or 5th division at prophase with super-condensed chromosomes

closely juxtaposed to the nuclear lamina (fig. 4D). The synergistic defects during the cell cycle cause by

CDK1a+b RNAi suggests collaborative roles of CDK1a and CDK1b during prophase, the point at which
CDK1b localized to chromatin (supplementary fig. S6).

Coevolution of *Oikopleura dioica* CDK1d and Cyclin Ba

Vertebrate Cyclin B1 is required for meiosis and mitosis. Most invertebrates, except *C. elegans* (van der Voet
et al. 2009), have a single ortholog of vertebrate Cyclin B1/B2 (fig. 1B), and this ortholog is required for
meiosis and mitosis (Okano-Uchida et al. 1998). Oscillation of Cyclin B levels orchestrate CDK1 kinase
activity during M-phase. As for other invertebrates, all 4 hermaphroditic *Oikopleura* species have a single
Cyclin B (Cyclin Bc), whereas dioecious *O. dioica* has Cyclins Ba and Bb (almost identical to Ba) in addition
to Cyclin Bc. The *cycBa* gene is located on the X chromosome, and is specifically expressed in ovaries.
Knockdown of Cyclin Bc, showed no significant differences in Cyclin Ba levels, H1 kinase activity in oocytes,
or embryonic development compared to wild type (fig. 5A), indicating that Cyclin Bc is dispensable for meiosis
and embryonic divisions. In contrast, *cycBa* RNAi oocytes were defective for embryonic division (fig. 5B).
Therefore, in *O. dioica* oogenic meiosis and early embryonic cleavage cycles, Cyclin Ba was required, whereas
Cyclin Bc was dispensable.

Among the oikopleurids, CDK1d and Cyclin Ba specifically appear in *O. dioica* and share several features:
X chromosome location, ovary-specific expression, similar degradation dynamics after fertilization (fig. 6B)
and similar RNAi phenotypes in meiosis and mitosis. Therefore, we wished to determine if they form an active
kinase complex. Cyclin Ba immunoprecipitates possessed H1 kinase activity, whereas lysates from infertile
cycBa RNAi oocytes lose H1 kinase activity (fig. 5C). This led us to test whether *in intro* constituted odCDK1d-
Cyclin Ba complexes possesses kinase activity. GST-odCDK1d recombinants expressed in insect sf9 cells
were insoluble. Co-expression with his-Cyclin Ba improved GST-odCDK1d solubility. Purified GST-
odCDK1d: his-Cyclin Ba complexes possessed H1 kinase activity (fig. 5E). Surprisingly, the conserved Cyclin
B Y170 residue which is critical for CDK1 binding (Goda et al. 2001; Bentley et al. 2007; Levasseur et al.
2013; Levasseur et al. 2019) is divergent in odCyclin Ba (replaced by His), whereas it is conserved in odCyclin
Bc (supplementary fig. S7). A Y170A substitution is known to completely abolish CDK1:Cyclin B interaction
in mouse, human, *Xenopus laevis* and *Ciona intestinalis* (Goda et al. 2001; Bentley et al. 2007; Levasseur et
al. 2013; Levasseur et al. 2019). The CDK1d:Cyclin Ba complex also has modified residues at the position of
the salt bridge (Goda et al. 2001) (supplementary fig. S7), which is in spatial proximity to Y170. This suggests
that CDK1d and Cyclin Ba may have coevolved an exclusive interaction that does not interfere with the more
canonical CDK1a/b/c:CycBc interactions.

Oscillation of CDK1 levels, instead of Cyclin B levels, orchestrates early embryonic divisions

In marine invertebrates and in *Xenopus* embryos, mitosis is driven by accumulation of Cyclin B to a threshold
required to activate CDK1, and mitotic degradation of Cyclin B inactivates CDK1, allowing mitotic exit
(Murray and Kirschner 1989). Cyclin Ba was stocked in *O. dioica* oocytes and was progressively degraded
during early embryogenesis, dropping to low levels by 2.5 hpf (fig. 6A). CDK1d also reached low levels during
this same period (fig. 6B). We then performed higher resolution sampling of Cyclin Ba levels at 2 min intervals
during the first embryonic cell cycle. Morphologies of the zygotes were recorded at each time point and samples

were divided into 2 aliquots and assayed for Cyclin Ba protein levels and H1 kinase activity. As shown in fig. 6F, consistent with cell cycle progression, H1 kinase activity oscillated during the cell cycle, peaking 20 min post-fertilization, when the nuclear envelope had disappeared and spindles were visible. H1 kinase levels dropped at 23 min post fertilization, when cleavage furrows formed. H1 kinase activity dropped to a basal level at 28 min post fertilization, abscission occurred, and two nuclei were visible. Surprisingly, CDK1d levels oscillated during the first cell cycle (fig. 6 C-G), in contrast to CDK1b and c levels which remained constant. CDK1d levels and H1 kinase activity peaked together at 20 min post fertilization, when NEBD occurred and the spindle assembled. CDK1d levels then peaked again, coincident with a rise in H1 kinase activity during the second cell cycle (fig. 6Fand G). To further support this unusual and unexpected oscillation in CDK1d levels, we labeled oocytes with ^{35}S-methionine, and sampled zygotes after *in vitro* fertilization. Autoradiography of SDS-PAGE detected a 37 KD band at the expected CDK1d MW, which appeared 20 min post-fertilization, corresponding to M phase (fig. 6H). Contrary to the oscillation of CDK1d levels, Cyclin Ba levels did not oscillate during the first 2 embryonic cycles (fig. 6H and G), instead, dropping at later embryonic stages. This latter observation is similar to *Drosophila* preblastoderm cycles, where neither bulk levels of Cyclin B, nor the bulk activity of Cdk1 kinase oscillate, with progressive depletion of maternally stocked Cyclin B only leading to oscillations in Cyclin B levels at later developmental stages (Edgar et al. 1994). A key, and novel difference in *O. dioica* is that CDK1d levels did oscillate during these first embryonic cycles.

Discussion

Oikopleura dioica exhibits high intrinsic rates of natural population increase (*r*) (0.68 d–1 to 1.07 d–1), well above the normal range for a metazoan of its size (Troedsson et al. 2002). The parameter, *r,* is increased by decreasing generation time*,* and/or by increasing egg number. In *O. dioica,* generation time is highly heritable and genetically correlated with *r*, whereas fecundity is much less heritable and not genetically correlated with *r* (Lobon et al. 2011). This emphasizes the importance of generation time in setting *O. dioica* fitness and is evidence of a life-history trait that has been strongly constrained by evolution (Deibel and Lowen 2012). The short duration of embryogenesis can be a direct target for selection to increase *r*.

We propose a working model for *O. dioica* cell cycle regulation under the framework of multiple CDK1 paralogs during oogenesis and very rapid, early embryogenesis (fig. 7). Individual, or combined, knockdowns of CDK1c and e, did not generate any detectable phenotypes during either oogenic meiosis or early embryonic divisions. Of note, neither of these CDK1 paralogs was able to complement yeast *cdc28* mutants. Similar to observations in mouse embryos and cultured vertebrate cells, CDK2, the only CDK with a canonical PTSAIRE helix in *Oikopleura,* was dispensable for completion of meiosis and early embryonic divisions, as *CDK2* RNAi zygotes divided and hatched normally (fig. 4A). On the other hand, normal early embryonic divisions required 3 CDK1 paralogs (figs. 4, 5, 7 and supplementary fig. S5), none of which contain a canonical PSTAIRE helix (fig. 1A). Depletion of CDK1a or b alone, delayed embryonic divisions (fig. 3A and B), whereas depletion of CDK1d abolished meiotic NEBD (Ovrebo et al. 2015), disrupted meiosis completion and caused abnormal embryonic cell cycles (fig. 3C-G). Knockdown of CDK1d also depleted oocytes of histone H1kinase activity (Feng and Thompson 2018), as did knockdown of its binding partner, Cyclin Ba (fig. 5C). Double knockdowns of CDK2 and CDK1a arrested zygotes at the pronuclear stage (fig. 4B) whereas double knockdowns of CDK1a and b generated multinucleate embryos with irregular cell division patterns that failed to hatch (fig. 4C and D).

CDK1d had critical functions in completion of meiotic M-phases and early embryonic divisions, during which
it showed an unusual oscillation of protein levels that peaked in M-phase and decreased during interphase.
CDK1d protein dynamics correlated well with the peak of H1 kinase activity and spindle formation during M
phase (fig. 6C-H).

The 3 CDK1 paralogs are sequentially activated in a tiling pattern when embryos traverse late S- to M-
phase, suggesting a further subdivision of CDK1 tasks and substrate targeting, beyond the CDK1-CDK2
subdivision that occurred much earlier in metazoan evolution. An additional, interesting feature of the *O. dioica*
CDK1 paralog complement, is that it is CDK1d protein levels rather than its partner, Cyclin Ba, the only
essential Cyclin B for *O. dioica* oogenic meiosis and early embryonic cell cycles, which oscillates during early
embryonic cell cycles (fig. 6F and G). This contrasts the generally universal oscillation of Cyclin B levels on
a background of stable CDK1 protein levels characteristic of metazoan cell cycle regulation.

During the maternal to zygotic transition, initial, rapid cell cycles, followed by progressive slowing, are a
widespread occurrence in a number of species, indicating a consistent benefit to this strategy throughout the
evolution of animals with externally deposited eggs (Farrell and O'Farrel 2014; Yuan et al. 2016). This strategy
converges on modulating the activity of the cell cycle driver, CDK1: weak inhibitory phosphorylation of CDK1
drives rapid cycles without gap phases, and when inhibitory phosphorylation is introduced, catalyzed by Wee
and Myt1 kinases, the cell cycle slows and gap phases are inserted (Tsai et al. 2014). *O. dioica* and, the model
arthropod, *Drosophila,* are characterized by a number of similarities with respect to oogenesis and early
embryogenesis. Both utilize a syncytial strategy during oogenesis (albeit with significant differences in details),
where polyploid nurse nuclei generate transcripts that code for materials stocked in the oocytes, which house
largely transcriptionally quiescent meiotic nuclei. Both generate excess Cyclin B protein storage in oocytes,
with progressive Cyclin B destruction during succeeding embryonic cycles. They diverge, however, in
adaptations of rapid mitotic cycles to Cyclin B preloading. *Drosophila* syncytial embryos adapt Cyclin B
preloading to mitosis without cytokinesis, compatible with little oscillation of CDK1 kinase levels due to
localized Cyclin B destruction (Edgar et al. 1994). In contrast, completion of cytokinesis during the rapid
mitoses of *O. dioica* embryonic cycles appears to require oscillation of CDK1 kinase levels (fig. 6F and G).
CDK1d oscillations may be an adaptation to the coenocystic ovary structure and Cyclin Ba preloading. Another
possible benefit to storage of Cyclin Ba, rather than CDK1d, could be that progressive destruction of Cyclin
Ba by the anaphase promoting complex during embryogenesis would deplete Cyclin Ba, facilitating switching
of key cell cycle regulators after hatching. In this regard, regulators, such as CDC25, Wee1, and Emi2, known
to be critical in controlling important embryonic cell cycle transitions in other organisms (Hormanseder et al.
2013; Farrel and O'Farrell 2014; Tsai et al. 2014; Yuan et al. 2016), are all duplicated in the *O. dioica* genome.

Coenocystic oogenesis is a common feature of the Oikopleuridae (Ganot et al. 2006), allowing synchronous
oocyte growth and maturation as an adaption to a semelparous, opportunistic life style. The oikopleurid lineage
also shares the presence of CDK1a, b and c paralogs. This raises questions as to the relationship between the
specific gain of the CDK1d-Cyclin Ba complex in *O. dioica*, their linkage on the recently evolved X
chromosome, and their ovary-specific expression. This pattern could be explained by resolution of sexual
antagonism by gene duplication (Connallon and Clark 2011) and higher linkage disequilibrium on the X
chromosome (Schaffner 2004). For genes that function both in hermaphrodite ovary and testes, the coenocystic
ovary structure evokes conflict between different transcriptional regulation of those genes in polyploid nurse

nuclei in the ovary, versus diploid germline progenitor cells in the testis. CDK1c does not show sexually biased
gene expression (fig. 2A) (Campsteijin et al. 2012). CDK1c duplication giving rise to CDK1d and e in *O. dioica*
may have mitigated sexual antagonistic conflict by allowing CDK1d (sexually biased expression to ovaries) to
evolve essential ovarian functions. The X chromosome location of CDK1d might facilitate CDK1d ovary-
specific expression since duplicates beneficial to females preferentially accumulate on the X chromosome
where they are exposed to selection twice as often in females versus males (Connallon and Clark 2011). Indeed,
when compared to zygotic promoters, maternal promoters in *O. dioica* are located on the X-chromosome more
frequently than expected, ($\chi2 = 43.34$, df = 1, p = 4.61×10^{-11}), revealing a female-bias of X-linked genes in
O. dioica (Danks et al. 2018). X-linkage of *CDK1d* and *Cyclin Ba* loci would then efficiently sustain linkage
disequilibrium, reinforced through selection on the recently formed X chromosome (Bachtrog et al. 2009).
Since there will be no recombination in the heterogametic male, this would contribute to an increase in
sweeping to fixation in the population (Bachtrog et al. 2009; Kayserili et al. 2012).

In *O. dioica*, the 3 CDK1 paralogs (a, b and d), which are required for correction execution of early
embryonic cell cycles, exhibit sequence diversification on their activation segments, ATP binding sites and
Cyclin B interaction interfaces. This raises the possibility of different k_{cat}/k_m catalytic outputs or
phosphorylation site specificities during these rapid cell cycles. With respect to the K89 residue in the hinge
region, both CDK1a and b, which participate in events preceding CDK1d, share a K89E/D substitution. This
substitution would likely reduce CDK1 k_{cat}. CDK1a, b and d also have non-overlapping activation segments,
suggesting CDK1 paralogs may have different affinities toward the same substrate phosphorylation site and/or
different phosphorylation site preferences. The sequential activation of CDK1a, b and d may provide a more
stepwise, as opposed to progressive, increase of kinase activity, which may enhance and sharpen transitions
towards early, middle and late substrates across early embryonic cell cycles. In metazoan cell cycles, the
division of labor between CDK1 and CDK2 is not absolute. CDK1 retains essential roles in late S phase that
CDK2 does not efficiently complement (Widrow et al. 1998; Katsuno et al. 2009; Nakanishi et al. 2010; Farrell
et al. 2012). Inactivation of Cdk1 in temperature-sensitive Cdk1 mutant cell lines (FT210) resulted in a
prolonged S phase accompanied by ineffective firing of late replicon clusters and reduced the density of active
origins during late S phase (Katsuno et al. 2009). In *Drosophila* pre-mild blastula transition S phases, it is Cdk1
activity, rather than Cdk2 activity that regulates late replication (Farrell et al. 2012; Seller and O'Farrell 2018).
The requirement for CDK1 in late S phase, coexists with its need to be repressed during early S phase
(Larochelle et al, 2007) and its control in late S phase by Chk1 (Lemmens et al. 2018), to prevent premature
triggering of M phase (Szmyd et al. 2019). In *O. dioica*, the conflict of CDK1's roles in the metazoan cell cycle
of regulating both late S and M phase, might be relaxed by CDK1 duplication and subsequent partitioning of
ancestral CDK1's roles in S and M phase to specific paralogs. Our results indicate that CDK1a collaborates
with CDK2 to drive S phase and suggest that this paralog may exhibit reduced risk of triggering M phase,
through lower intrinsic kinase activity, arising from substitution in hinge region (supplementary fig. S2). On
the other hand, the CDK1d paralog has become a specialist of M phase events without additional constraints
imposed by regulation of late DNA replication. This may reflect adaptation to very rapid embryonic cell
divisions in this planktonic organism.

A conformation change of the very highly conserved PSTAIRE helix is central to CDK1
inactivation/activation transitions (Huang et al. 2012). It is therefore surprising that this sequence was modified

in all *O. dioica* CDK1 paralogs. *O. dioica* CDK2, the only CDK in this species with a conserved PSTAIRE,

was dispensable for embryonic divisions, recalling the situation in plants, where the CDK1 ortholog, CDKA,

which has the canonical PSTAIRE, is dispensable for the cell cycle, whereas CDKB, deviating in the

PSTAIRE, has essential functions in the absence of CDKA (Nowack et al. 2012). Strikingly, the modified

PSTAIRE of the *O. dioica* CDK1d paralog (PPTSLRE) shows convergence over great evolutionary distance

with plant CDKB sequences (P[P/S]T[A/T]LRE), and in both *O. dioica*, and plants, these variants exhibit

increased specialization to M-phase (Joubes et al. 2000; Nowack et al. 2012). The suite of naturally evolved *O.*

dioica, CDK1:Cyclin B complexes, with critical residue substitutions in their activation segments, ATP binding

sites, CDK1-Cyclin B interaction interfaces, and Cyclin B-substrate interaction elements, provide a fertile

foundation for structural studies of how these changes elicit enhanced kinase selection for subsets of G2-M,

CDK1 cell cycle substrates.

Materials and methods

Animal culture and collection

O. dioica were maintained in culture at 15°C (Bouquet et al. 2009). Day 4–6 animals were placed in filtered

seawater, removed from their houses and anesthetized in cold ethyl 3-aminobenzoate methanesulfonate salt

(MS-222, 0.125 mg/ml; Sigma) before collection.

In vitro **fertilization (IVF) and collection of synchronously dividing zygotes**

Sperm solutions were prepared by transferring 5 mature males to a petri dish containing 10 ml artificial

seawater, on ice, and sperm quality was assessed using a Nikon Eclipse E400 microscope. Mature females

were transferred to artificial sea water in 24-well plates coated with 0.9% agarose, artificial sea water was

changed every hour. Oocytes were fertilized with 400 μl sperm solution and fertilized oocytes were washed in

8 ml artificial sea water within 50 seconds. Synchronicity of fertilization was verified by following polar body

exclusion and early cleavages.

Phylogenetic Analyses

A search of larvacean CDK1, CDK2 and Cyclin B proteins was performed using TBLASTN against genomes

and transcriptomes of *Oikopleura albicans, O. longicauda, O. vanhoeffeni* and *Fritillaria borealis*. *O. dioica*

CDK1, CDK2 and Cyclin Bc were used as queries with E-value cutoff set to 1e-005. *Ciona intestinalis* and

Branchiostoma floridae proteins were obtained from the JGI Genome Portal

(https://genome.jgi.doe.gov/mycocosm/home). For identified targets, amino acid sequences of the N-terminal

cyclin box or the kinase domain of CDK1/2 were aligned (clustalX). Alignments were manually edited (Bio-

edit), and secondary structure analysis (psipred) was used to guide alignments. Phylogenetic relationships were

inferred using maximum likelihood (ML) using MEGA7 (Kumar et al. 2016), with amino acid sequence

alignment based on Jones-Taylor-Thornton (Jones et al 1992). Estimates of the gamma-shape parameter and

proportion of invariant sites were then used to obtain the 500 bootstrap replicates.

Test of Selection

Branch-sites tests of selection are phylogeny-based tests that require the *a priori* identification of a foreground

branch that is hypothesized to be evolving at a different rate than the background branches in the rest of the

tree. Codons are alignment with PRANK, gaps in alignment were deleted. We first set a series of nested

models specifying particular branches as foreground branches: the lineage subtending the *CDK1a* clade,

the lineage subtending the *CDK1b* clade, the lineage subtending the *CDK1c* clade. Using a likelihood ratio

test, we then tested the hypothesis, that a model which included one class of sites with $\omega > 1$, indicating positive

selection, was superior to a model where $\omega = 1$. We estimated codon frequencies on the basis of observed

nucleotide frequencies at each of the three codon positions, as a procedure to reduce false positives due to

potential nucleotide biases. We also used the empirical Bayes method as implemented in PAML 4 (Yang 2007)

to identify the positively selected sites along the 3 branches. The synonymous (silent) and non-synonymous

(non-silent) nucleotide substitution rates of each branch were also estimated by the modified Nei–Gojobori

method (Zhang et al. 1998), based on tree topology inferred by maximum likelihood (ML) using MEGA

(Kumar et al. 2016).

Homology Modeling

Active and inactive hsCDK1 was predicted with I-TASSER (Yang et al. 2015), using the hsCDK1-cyclin B

crystal structure (Brown et al. 2015). 3D models were visualized using Pymol (DeLano 2002). Based on

alignment of odCDK1 paralogs and hsCDK1, nonconservative variations between odCDK1s and hsCDK1

were colored in the models.

Yeast Complementation

We used a W303a-background yeast wild-type strain SCU893 (*MATa ura3 ade2-1 trp1-1 leu2-3112 his3-11*

bar1::hisG) and its derivative SCU112 (*cdc28-4*) (gifts from Jorrit Enserink, Univ. of Oslo). To complement

a *cdc28* mutation of *S. cerevisiae*, the open reading frames of the *O. dioica CDK1* paralogs and *CDK2* cDNAs

were cloned in pCR™4-TOPO® TA Vector (Thermofisher). cDNA fragments were isolated by digestion with

*EcoR*I and *Not*I and ligated into the yeast pYES2 expression vector (Invitrogen) downstream of the inducible

GAL1 promoter. The resulting CDK1 paralog plasmids were individually transformed by the lithium acetate

method (Ito et al. 1984) into the *S. cerevisiae cdc28-4* strain. As a control, *cdc28-4* was transformed with the

empty pYES2 vector. The transformants were grown for 4 days at 25°C, and the resulting colonies were

cultivated overnight at 25°C in CM-ura medium containing glucose as a carbon source. The cells were pelleted,

washed, resuspended in CM-ura medium containing 2% glycerol, and grown further at 25°C. After 4 h, the

cells were pelleted and resuspended in CM-ura medium containing either glucose (2%), or galactose (2%)–

raffinose (1%) as carbon source, and grown at 25°C, or the restrictive 36°C, respectively.

Antibodies

Rabbit polyclonal affinity–purified anti-Cyclin Ba/b (acetyl-NRDLNIQESGPVKAVVNAC-amide; acetyl-

CLEFLRRFSRVAEETIDPKEY-amide), anti-CDK1b (acety1-CPDFTKPSTLNVNTSLNEMDM-amide),

anti-CDK1c (acetyl-MRKRNIDRPPTQDLNNYC–amide), ant-Cdk1d (acety1-MIKAKSGTQDLSNYKRC-

amide), and anti-lamin1 (acetyl-QSPISLPPLSGSTC-amide) were from 21st Century Biochemicals (Marlboro,

MA). Other antibodies: anti-PSTAIR (Abcam, ab10345), anti-ATPB (Abcam, ab37922), anti-histone 3

(Abcam, ab10799). Secondary antibodies for immunofluorescence: goat anti-rabbit IgG (H+L) Alexa Fluor

568 (Thermofisher, A-11011); and for western blots: goat anti-rabbit IgG (H+L) superclonal™ secondary

antibody, HRP (Thermofisher, A27033), Peroxidase-AffiniPure Goat Anti-Mouse IgG (H+L) (Jackson ImmunoResearch, 115-035-003).

Western blots

Equal numbers of animals or embryos were collected in clean seawater, centrifuged, washed in PBS, and snap frozen. Animal lysis was performed in $1\times$ Laemmli (50 mM Tris [pH 6.8], 2% sodium dodecyl sulfate [SDS], 10% glycerol, 250 mM β-mercaptoethanol, 0.001% bromophenol blue) by heating to 80°C for 15 min followed by brief vortexing. Insoluble material was removed by centrifugation (5 min at $13,000 \times g$). Equal volumes of lysate were applied to 10% SDS-polyacrylamide gel electrophoresis (PAGE) and transferred onto nitrocellulose membranes (Whatman) in transfer buffer (250 mM Tris, 190 mM glycine, 0.5% SDS, 20% methanol). Membranes were rinsed in Tris-buffered saline (TBS) and blocked for 1 h in 5% skim milk/TBST (TBS, 0.1% Tween 20) or 5% BSA/TBST prior to incubation with antibodies diluted in 5% skim milk/TBST (1:1000), O/N, at 4°C, under gentle agitation. Membranes were washed in TBS, TBST (twice), and TBS followed by incubation with secondary antibody in 5% skim milk/TBST for 1 h at room temperature. Membranes were washed as above, followed by a wash in PBST (PBS, 0.1% Tween 20) and processed using ECL (Pierce).

Immunofluorescence

Embryos were fixed in 4% PFA, 0.1% triton X-100, 0.1 M MOPS (pH 7.5), and 0.5 M NaCl at 4°C O/N. Fixed animals were washed once with PBSTE (PBS, 0.1% tween-20, 1mM EDTA), thrice with PBSTEG (PBSTE, 0.1 M glycine), twice with PBSTE, and then blocked with 3% BSA/PBSTE at 4°C O/N. Primary antibodies were either injected into embryos at 1:50 dilution or incubated with embryos at 1:200 in 3% BSA/PBSTE at 4°C for at least 1 week. Samples were washed six times in PBSTE, post-fixed O/N in 4% PFA/PBSTE, and washed with PBSTE ($1\times$), PBSTEG ($3\times$), and PBSTE ($2\times$). Following secondary antibody incubation (1:500; 4–6 days), samples were washed six times in PBSTE, DNA was counterstained with 1 μM To-Pro-3 iodide (Molecular Probes), and specimens were mounted in Vectashield (Vector Laboratories). Confocal analysis was performed at 20°C using a Leica TCS laser scanning confocal microscope using Leica (LAS AF v2.3) software.

Quantitative Reverse Transcription–PCR

Total RNA was isolated from snap-frozen animals. For qRT-PCR, 1 μg of purified total RNA was subjected to RT using M-MLV reverse transcriptase (Invitrogen). The quantitative polymerase chain reaction (qPCR) (CFX-96; Bio-Rad) was performed using cDNA templates synthesized from an equivalent of 5 ng total RNA, 10 μl of qPCR 2 Master Mix (Bio Rad), and 500 nM primers in a total volume of 20 μl. After initial denaturation for 5 min at 95 °C, 45 cycles of 95 °C for 15 s, 60 °C for 20 s, and 72 °C for 20 s were conducted, with a final extension for 5 min at 72 °C. RT-negative controls were run to 45 cycles. RPL23 and EF-1β mRNA levels served as normalization controls in all qRT-PCRs. For RNAi experiments, target mRNA levels were normalized to control zygotes. Primer pair specificities were confirmed by melting curve evaluation, cloning, and sequencing of amplicons. In these experiments, mean values normalized to EF-1β control transcripts are presented.

Microinjection

Gonad microinjections were performed as previously (Ovrebo et al. 2015). Injection solutions were prepared by mixing 300 ng/µl double stranded RNA (dsRNA) in RNase free PBS. Anesthetized day 4 animals in filtered seawater were transferred to Petri dishes coated with 2% agarose. Gonads were injected on a Nikon Eclipse TE2000-S inverted microscope equipped with Narishige micromanipulators and transjector 5246 micro injector (Eppendorf). Holding pipettes were crafted from borosilicate capillaries with an outer diameter of 1 mm and inner diameter of 0.5 mm (Sutter instruments). Injection needles (quartz, Sutter instruments) were pulled with outer diameter of 1 mm and inner diameter of 0.7 mm on a P-1000 Micropipette puller (Sutter instruments) and coated with Dichlorodimethulsilane (Fluka). Injected day 4/5 animals were transferred to watch glasses containing filtered seawater for recovery and then transferred to 6 L beakers for subsequent culture. Animals were cultured until spawning for in vitro fertilization (IVF) and immunofluorescence. Before oocyte injection, oocytes were transferred to 24-well plate coated with 0.9% agarose containing fresh artificial seawater (Red Sea) for 1 h. Samples of 50 oocytes were used for qRT-PCR. Remaining oocytes were fertilized and the statistical significance of any developmental defects arising from various treatments was assessed by unpaired Student's t-test for equal variances, or, unpaired, Welch's t-test for unequal variances. Effect size was calculated as Cohen's $d = 2t/SQRT\ (df)$.

In vitro **CDK1 kinase activities**

CDK1 kinase activities were determined as previously (Feng and Thompson 2018). Equal numbers of oocytes were snap frozen and thawed twice. Cyclin Ba immunoprecipitates, purified his-Cyclin Ba-GST OdCdk1d complexes were aliquoted, kept at 80°C and thawed before use. Samples were mixed with biotinylated MV peptide and 1 mM ATP incubated at 30°C for 30 min. After stopping the phosphorylation reaction, ELISA was performed using anti-phospho MV peptide antibody (MESACUP) and the CDK1 Kinase Assay Kit (MBL). Finally, POD-conjugated streptavidin was used to detect phosphorylated MV peptide and color development measured at 492 nm. Differences in CDK1 kinase activities from different knockdown treatments were assessed by the Student's T-test. H1 kinase assays were performed as previously (Levasseur and McDougall 2000). After thawing snap frozen oocyte samples on ice, 2 µl of 6× kinase reaction mixture (pH 7.2) containing histone H1 (Sigma type III from calf thymus), 0.6 mM ATP, 0.5 mCi/ml [32P]ATP, and 60 µM cAMP-dependent protein kinase inhibitor was added to 10 µl of oocyte sample. The reaction was started synchronously by transfer to a 30°C heat block for 30 min. The reaction was stopped synchronously by adding 5× Laemmli buffer. The samples were then heated to 80°C for 10 minutes and resolved on 15% polyacrylamide gels. The gels were placed in a Phosphorimager (Fujix; 1500 Bas Reader) and the incorporation of [32P] was quantified.

Metabolic labeling of zygotes

Unfertilized oocytes obtained from individual females were transferred into one well of 24-well plate coated with 0.9% agarose. After addition of a mixture of 35S-labeled amino acids (Pro-mix, Amersham) to a final concentration of 330 µCi/ml and incubation for 60 min at room temperature, oocytes were fertilized and harvested at different times post fertilization. The samples were briefly centrifuged to eliminate excess of labeled mixture and 5x Laemmli buffer was added to the pellets. Samples were then heated to 80°C for 10 min and resolved on 10% polyacrylamide gels. Gels were stained with 0.5% Coomassie blue R250 in 45%

588   methanol, 45% water, 10% acetic acid, and destained in 20% methanol, 75% water, 5% acetic acid; and dried

589   on Whatman 3MM paper. Dried gels were autoradiographed using a Phosphorimager (Fujix; 1500 Bas Reader).

Capped mRNA and dsRNA synthesis

For indestructible Cyclin Ba injection, site directed mutagenesis (New England Biolabs) was used to delete the

destruction box (NTEREQLQALN) using a cyclin Ba/b entry clone as template with subsequent recombination

with pSPE3-Venus-RfA. The resulting plasmid was used as a template for PCR amplification. Amplicons were

purified by phenol-chloroform extraction followed by ethanol precipitation. Capped mRNAs were synthesized

from these templates using mMessage mMachine (Ambion) followed by poly(A) tailing (Ambion) according

to the manufacturer's protocol and purified by Lithium chloride precipitation. DNA templates for dsRNA

synthesis were prepared by PCR using gene-specific primers with T7 overhang and purified by phenol-

chloroform extraction. Sense and antisense RNA were synthesized in a single reaction following the

recommended protocol using T7 RiboMAX^{TM} Express RNAi system (Promega).

Recombinant protein expression in insect cells

The coding regions of Cyclin Ba/b and OdCdk1d were cloned into the pCR™8/GW/TOPO entry plasmid and

then recombined into pDEST™10 (N-terminal His$_6$-TEV fusion tag) or pDEST™20 (N-terminal GST fusion

tag), respectively, according to the manufacturer's (Thermofisher Scientific) instructions. The EMBL Protein

Expression and Purification Core Facility performed protein expression and purification. The

pDEST10_Cyclin Ba/b and pDEST20_OdCdk1d plasmid DNA were used for transposition into *E. coli*

DH10EMBacY cells. Bacmid DNA was isolated from a 3 ml overnight culture of positive (white) colonies

according to the MultiBac protocol (Nie et al. 2013). The bacmid DNA was mixed with X-tremeGENE

transfection reagent and Sf-900 III SFM medium and the mixture was added to Sf21 insect cells. After 3 days,

the V$_0$ virus was harvested and used to infect a larger Sf21 culture for V$_1$ virus generation. For co-expression

of His$_6$-tagged Cyclin Ba/b and GST-tagged OdCdk1d, 500 ml Sf21 insect cells was co-infected with 2 ml

Cyclin Ba/b V$_1$ virus and 2 ml OdCdk1d V$_1$ virus and cultured for 72 h at 27°C. Cells were harvested by

centrifugation and resuspended in running buffer (50 mM Tris pH 8.0, 250 mM NaCl and 20 mM imidazole)

supplemented with benzonase and cOmplete EDTA-free protease inhibitors. Cells were lysed using a Dounce

homogenizer and the lysate was cleared by centrifugation (100000 x *g*, 30 min, 4°C). The cleared lysate was

incubated with 200 µl Ni-NTA (Qiagen) beads for 2 h at 4°C. After incubation, the mixture was transferred to

an empty PD10 column, the beads were washed with running buffer and then eluted in 200 µl fractions with

running buffer supplemented with 300 mM imidazole. All fractions were analysed by SDS-PAGE and the

fractions containing the Cyclin Ba/b–OdCdk1d complex were concentrated, flash-frozen in liquid nitrogen and

stored at -80°C until further use.

Acknowledgments

622 We thank Haiyang Feng, Sars Centre, for helpful discussion and A. Aasjord and K. N. Nøkling for providing
623 animals from the *Oikopleura* culture facility. We thank the Chourrout S1 group at the Sars Centre for early
624 access to sequence data from hermaphroditic appendicularian species.

Disclosure Statement

626 No potential conflict of interest was reported by the authors.

Author Contributions

628 X.M., and E.M.T. conceived and designed the experiments. X.M. performed experiments and X.M. and E.M.T.
629 analyzed the data. J.I.Ø performed CDK1-GFP cmRNA injection and immunostaining. X.M. and E.M.T. wrote
630 the manuscript. All authors approved the manuscript.

632 **References**
633
Ahn SH, Tobe BT, Gerald JNF, Anderson SL, Acurio A, Kron SJ. 2001. Enhanced cell polarity in mutants of the budding
yeast cyclin-dependent kinase Cdc28p. *Mol Biol Cell* 12(11):3589-3600.
Arooz T, Yam CH, Siu WY, Lau A, Li KKW, Poon RYC. 2000. On the concentrations of cyclins and cyclin-dependent
kinases in extracts of cultured human cells. *Biochemistry* 39(31):9494-9501.
Averof M, Rokas A, Wolfe KH, Sharp PM. 2000. Evidence for a high frequency of simultaneous double-nucleotide
substitutions. *Science* 287(5456):1283-1286.
Bachtrog D, Jensen JD, Zhang Z. 2009. Accelerated adaptive evolution on a newly formed X chromosome. *Plos Biol.*
7(4):e1000082
Bao ZQ, Jacobsen DM, Young MA. 2011. Briefly bound to activate: Transient binding of a second catalytic magnesium
activates the structure and dynamics of cdk2 kinase for catalysis. *Structure* 19(5):675-690.
Bentley AM, Normand G, Hoyt J, King RW. 2007. Distinct sequence elements of cyclin B1 promote localization to
chromatin, centrosomes, and kinetochores during mitosis. *Mol Biol Cell* 18(12):4847-4858.
Bouquet JM, Spriet E, Troedsson C, Ottera H, Chourrout D, Thompson EM. 2009. Culture optimization for the emergent
zooplanktonic model organism *Oikopleura dioica*. *J Plankton Res.* 31(4):359-370.
Brown NR, Korolchuk S, Martin MP, Stanley WA, Moukhametzianov R, Noble MEM, Endicott JA. 2015. Cdk1
structures reveal conserved and unique features of the essential cell cycle Cdk. *Nat Commun.* 6:e6769
Brown NR, Noble MEM, Endicott JA, Johnson LN. 1999. The structural basis for specificity of substrate and recruitment
peptides for cyclin-dependent kinases. *Nat Cell Biol.* 1(7):438-443.
Campsteijn C, Ovrebo JI, Karlsen BO, Thompson EM. 2012. Expansion of cyclin D and Cdk1 paralogs in *Oikopleura*
dioica, a chordate employing diverse cell cycle variants. *Mol Biol Evol.* 29(2):487-502.
Chen C, Ha BH, Thevenin AF, Lou HJ, Zhang R, Yip KY, Peterson JR, Gerstein M, Kim PM, Filippakopoulos P et al.
2014. Identification of a major determinant for serine-threonine kinase phosphoacceptor specificity. *Mol Cell*
53(1):140-147.
Child ES, Hendrychova T, McCague K, Futreal A, Otyepka M, Mann DJ. 2010. A cancer-derived mutation in the pstaire
helix of cyclin-dependent kinase 2 alters the stability of cyclin binding. *Biochim Biophys Acta.* 1803(7):858-
864.

Connallon T, Clark AG. 2011. The resolution of sexual antagonism by gene duplication. *Genetics* 187(3):919-937.

Creixell P, Palmeri A, Miller CJ, Lou HJ, Santini CC, Nielsen M, Turk BE, Linding R. 2015. Unmasking determinants of
specificity in the human kinome. *Cell* 163(1):187-201.

Danks GB, Navratilova P, Lenhard B, Thompson EM. 2018. Distinct core promoter codes drive transcription initiation at
key developmental transitions in a marine chordate. *BMC Genomics* 19:164

Danks GB, Raasholm M, Campsteijn C, Long AM, Manak JR, Lenhard B, Thompson EM. 2015. Trans-splicing and
operons in metazoans: Translational control in maternally regulated development and recovery from growth
arrest. *Mol Biol Evol.* 32(3):585-599.

Deibel D, Lowen B. 2012. A review of the life cycles and life-history adaptations of pelagic tunicates to environmental
conditions. *ICES J Mar Sci.* 69(3):358-369.

DeLano WL. 2002. The PyMOL Molecular Graphics System. Delano Scientific, San Carlos.

Des Marais DL, Rausher MD. 2008. Escape from adaptive conflict after duplication in an anthocyanin pathway gene.
Nature 454(7205):762-785.

Echalier A, Cot E, Camasses A, Hodimont E, Hoh F, Jay P, Sheinerman F, Krasinska L, Fisher D. 2012. An integrated
chemical biology approach provides insight into Cdk2 functional redundancy and inhibitor sensitivity. *Chem
Biol.* 19(8):1028-1040.

Edgar BA, Orr-Weaver TL. 2001. Endoreplication cell cycles: More for less. *Cell* 105(3):297-306.

Edgar BA, Sprenger F, Duronio RJ, Leopold P, Ofarrell PH. 1994. Distinct molecular mechanisms regulate cell-cycle
timing at successive stages of *Drosophila* embryogenesis. *Gene Dev.* 8(4):440-452.

Farrell JA, O'Farrell PH. 2014. From egg to gastrula: How the cell cycle is remodeled during the *Drosophila* mid-blastula
transition. *Annu Rev Genet.* 48:269-294.

Farrell JA, Shermoen AW, Yuan K, O'Farrell PH. 2012. Embryonic onset of late replication requires Cdc25 down-
regulation. *Gene Dev.* 26(7):714-725.

Feng HY, Thompson EM. 2018. Specialization of Cdk1 and cyclin B paralog functions in a coenocystic mode of oogenic
meiosis. *Cell Cycle* 17(12):1425-1444.

Fleig UN, Gould KL, Nurse P. 1992. A dominant negative allele of p34(cdc2) shows altered phosphoamino acid content
and sequesters p56(cdc13) cyclin. *Mol Cell Biol.* 12(5):2295-2301.

Ganot P, Bouquet JM, Kallesoe T, Thompson EM. 2007. The *Oikopleura* coenocyst, a unique chordate germ cell
permitting rapid, extensive modulation of oocyte production. *Dev Biol.* 302(2):591-600.

Ganot P, Bouquet JM, Thompson EM. 2006. Comparative organization of follicle, accessory cells and spawning anlagen
in dynamic semelparous clutch manipulators, the urochordate Oikopleuridae. *Biol Cell* 98(7):389-401.

Ganot P, Thompson EM. 2002. Patterning through differential endoreduplication in epithelial organogenesis of the
chordate, *Oikopleura dioica*. *Dev Biol.* 252(1):59-71.

Goda T, Funakoshi M, Suhara H, Nishimoto T, Kobayashi H. 2001. The N-terminal helix of *Xenopus* cyclins A and B
contributes to binding specificity of the cyclin-Cdk complex. *J Biol Chem.* 276(18):15415-15422.

Hochegger H, Deisuphong D, Sonoda E, Sciberi A, Rajendra E, Kirk J, Hunt T, Takeda S. 2007. An essential role for
Cdk1 in s phase control is revealed via chemical genetics in vertebrate cells. *J Cell Biol.* 178(2):257-268.

Holmes JK, Solomon MJ. 1996. A predictive scale for evaluating cyclin-dependent kinase substrates - a comparison of
p34(cdc2) and p33(cdk2). *J Biol Chem.* 271(41):25240-25246.

Holt LJ, Tuch BB, Villen J, Johnson AD, Gygi SP, Morgan DO. 2009. Global analysis of Cdk1 substrate phosphorylation
sites provides insights into evolution. *Science* 325(5948):1682-1686.

Hormanseder E, Tischer T, Mayer TU. 2013. Modulation of cell cycle control during oocyte-to-embryo transitions.
EMBO J. 32(16):2191-2203.

Huang H, Zhao RJ, Dickson BM, Skeel RD, Post CB. 2012. Alpha C helix as a switch in the conformational transition of
Src/Cdk-like kinase domains. *J Phys Chem B* 116(15):4465-4475.

Ito H, Murata K, Kimura A. 1984. Transformation of intact yeast-cells treated with alkali cations or thiol compounds. *Agr*
Biol Chem. 48(2):341-347.

Jeffrey PD, Ruso AA, Polyak K, Gibbs E, Hurwitz J, Massague J, Pavletich NP. 1995. Mechanism of Cdk activation
revealed by the structure of a cyclina-Cdk2 complex. *Nature.* 376(6538):313-320.

Jones DT, Taylor WR, Thornton JM. 1992. The rapid generation of mutation data matrices from protein sequences.
Comput Appl Biosci. 8(3):275-282.

Joubes J, Chevalier C, Dudits D, Heberle-Bors E, Inze D, Umeda M, Renaudin JP. 2000. Cdk-related protein kinases in
plants. *Plant Mol Biol.* 43(5-6):607-620.

Katsuno Y, Suzuki A, Sugimura K, Okumura K, Zineldeen DH, Shimada M, Niida H, Mizuno T, Hanaoka F, Nakanishi
M. 2009. Cyclin A-Cdk1 regulates the origin firing program in mammalian cells. *Proc Natl Acad Sci U S A.*
106(9):3184-3189.

Kayserili MA, Gerrard DT, Tomancak P, Kalinka AT. 2012. An excess of gene expression divergence on the X
chromosome in *Drosophila* embryos: Implications for the faster-X hypothesis. *Plos Genet.* 8(12):1003200

Koivomagi M, Valk E, Venta R, Iofik A, Lepiku M, Balog ERM, Rubin SM, Morgan DO, Loog M. 2011a. Cascades of
multisite phosphorylation control Sic1 destruction at the onset of S phase. *Nature* 480(7375):128-U301.

Koivomagi M, Valk E, Venta R, Iofik A, Lepiku M, Morgan DO, Loog M. 2011b. Dynamics of Cdk1 substrate
specificity during the cell cycle. *Mol Cell* 42(5):610-623.

Kriwacki RW, Hengst L, Tennant L, Reed SI, Wright PE. 1996. Structural studies of p21(waf1/cip1/sdi1) in the free and
cdk2-bound state: Conformational disorder mediates binding diversity. *Proc Natl Acad Sci U S A.*
93(21):11504-11509.

Kumar S, Stecher G, Tamura K. 2016. MEGA7: Molecular evolutionary genetics analysis version 7.0 for bigger datasets.
Mol Biol Evol. 33(7):1870-1874.

Larochelle S, Merrick KA, Terret ME, Wohlbold L, Barboza NM, Zhang C, Shokat KM, Jallepalli PV, Fisher RP. 2007.
Requirements for Cdk7 in the assembly of Cdk1/cyclin B and activation of Cdk2 revealed by chemical genetics
in human cells. *Mol Cell* 25(6):839-850.

Lemmens B, Hegarat N, Akopyan K, Sala-Gaston J, Bartek J, Hochegger H, Lindqvist A. 2018. DNA replication
determines timing of mitosis by restricting CDK1 and PLK1 activation. *Mol Cell* 71(1):117-128.

Levasseur M, Dumollard R, Chambon JP, Hebras C, Sinclair M, Whitaker M, McDougall A. 2013. Release from meiotic
arrest in ascidian eggs requires the activity of two phosphatases but not camkii. *Development* 140(22):4583-
4593.

Levasseur M, McDougall A. 2000. Sperm-induced calcium oscillations at fertilisation in ascidians are controlled by
cyclin B1-dependent kinase activity. *Development* 127(3):631-641.

Levasseur MD, Thomas C, Davies OR, Higgins JMG, Madgwick S. 2019. Aneuploidy in oocytes is prevented by
sustained Cdk1 activity through degron masking in cyclin B1. *Dev Cell* 48(5):672-684.

Levine K, Kiang L, Jacobson MD, Fisher RP, Cross FR. 1999. Directed evolution to bypass cyclin requirements for the
Cdc28p cyclin-dependent kinase. *Mol Cell* 4(3):353-363.

Lobon CM, Acuna JL, Lopez-Alvarez M, Capitanio FL. 2011. Heritability of morphological and life history traits in a
pelagic tunicate. *Mar Ecol Prog Ser.* 422:145-154.

Macneill SA, Nurse P. 1993. Mutational analysis of the fission yeast p34cdc2 protein-kinase gene. *Mol Gen Genet.*
236(2-3):415-426.

MacNeill SA, Warbrick E, Fantes PA. 1991. Controlling cell cycle progress in the fission yeast *Schizosaccharomyces*
pombe. *Curr Opin Genet Dev.* 1(3):307-312.

Murray AW, Kirschner MW. 1989. Cyclin synthesis drives the early embryonic-cell cycle. *Nature.* 339(6222):275-280.

Nakanishi M, Katsuno Y, Niida H, Murakami H, Shimada M. 2010. Chk1-cyclin A/Cdk1 axis regulates origin firing
programs in mammals. *Chromosome Res.* 18(1):103-113.

Naville M, Henriet S, Warren I, Sumic S, Reeve M, Volff JN, Chourrout D. 2019. Massive changes of genome size
driven by expansions of non-autonomous transposable elements. *Curr Biol.* 29(7):1161-1168.

Nolen B, Taylor S, Ghosh G. 2004. Regulation of protein kinases: Controlling activity through activation segment
conformation. *Mol Cell* 15(5):661-675.

Nie Y, Bellon-Echeverria I, Trowitzsch S, Bieniossek C, Berger I. 2014. Multiprotein complex production in insect cells
by using polyproteins. *Methods Mol Biol.* 1091: 131-141.

Nowack MK, Harashima H, Dissmeyer N, Zhao XA, Bouyer D, Weimer AK, De Winter F, Yang F, Schnittger A. 2012.
Genetic framework of cyclin-dependent kinase function in arabidopsis. *Dev Cell* 22(5):1030-1040.

Okano-Uchida T, Sekiai T, Lee KS, Okumura E, Tachibana K, Kishimoto T. 1998. In vivo regulation of cyclin A/Cdc2
and cyclin B/Cdc2 through meiotic and early cleavage cycles in starfish. *Dev Biol.* 197(1):39-53.

Ord M, Loog M. 2019. How the cell cycle clock ticks. Mol Biol Cell. 30(2):169-172.

Ord M, Moll K, Agerova A, Kivi R, Faustova I, Venta R, Valk E, Loog M. 2019a. Multisite phosphorylation code of
Cdk. *Nat Struct Mol Biol.* 26(7):649-658.

Ord M, Venta R, Moll K, Valk E, Loog M. 2019b. Cyclin-specific docking mechanisms reveal the complexity of M-Cdk
function in the cell cycle. *Mol Cell.* 75(1):76-89.

Ovrebo JI, Campsteijn C, Kourtesis I, Hausen H, Raasholm M, Thompson EM. 2015. Functional specialization of
chordate Cdk1 paralogs during oogenic meiosis. *Cell Cycle* 14(6):880-893.

Rogozin IB, Belinky F, Pavlenko V, Shabalina SA, Kristensen DM, Koonin EV. 2016. Evolutionary switches between
two serine codon sets are driven by selection. *Proc Natl Acad Sci U S A.* 113(46):13109-13113.

Santamaria D, Barriere C, Cerqueira A, Hunt S, Tardy C, Newton K, Caceres JF, Dubus P, Malumbres M, Barbacid M.
2007. Cdk1 is sufficient to drive the mammalian cell cycle. *Nature* 448(7155):811-U818.

Schaffner SF. 2004. The x chromosome in population genetics. Nat Rev Genet. 5(1):43-51.

Seller CA, O'Farrell PH. 2018. Rif1 prolongs the embryonic s phase at the *Drosophila* mid-blastula transition. *Plos Biol.*
16(5).

Sonnhammer ELL, Koonin EV. 2002. Orthology, paralogy and proposed classification for paralog subtypes. *Trends*
Genet. 18(12):619-620.

Staver JM, Strathmann RR. 2002. Evolution of fast development of planktonic embryos to early swimming. *Biol Bull.*
203(1):58-69.

Strathmann RR, Staver JM, Hoffman JR. 2002. Risk and the evolution of cell-cycle durations of embryos. *Evolution*
56(4):708-720.

Swaffer MP, Jones AW, Flynn HR, Snijders AP, Nurse P. 2016. Cdk substrate phosphorylation and ordering the cell
cycle. *Cell* 167(7):1750-1761.

Szmyd R, Niska-Blakie J, Diril MK, Nunes PR, Tzelepis K, Lacroix A, van Hul N, Deng LW, Matos J, Dreesen O et al.
2019. Premature activation of Cdk1 leads to mitotic events in s phase and embryonic lethality. *Oncogene*
38(7):998-1018.

Troedsson C, Bouquet JM, Aksnes DL, Thompson EM. 2002. Resource allocation between somatic growth and
reproductive output in the pelagic chordate *Oikopleura dioica* allows opportunistic response to nutritional
variation. *Mar Ecol Prog Ser.* 243(11):83-91.

Tsai TYC, Theriot JA, Ferrell JE. 2014. Changes in oscillatory dynamics in the cell cycle of early *Xenopus laevis*
embryos. *Plos Biol.* 12(2):e1001788.

van der Voet M, Lorson MA, Srinivasan DG, Bennett KL, van den Heuvel S. 2009. *C. elegans* mitotic cyclins have
distinct as well as overlapping functions in chromosome segregation. *Cell Cycle* 8(24):4091-4102.

Widrow RJ, Hansen RS, Kawame H, Gartler SM, Laird CD. 1998. Very late DNA replication in the human cell cycle.
Proc Natl Acad Sci U S A. 95(19):11246-11250.

Wood DJ, Korolchuk S, Tatum NJ, Wang LZ, Endicott JA, Noble MEM, Martin MP. 2019. Differences in the
conformational energy landscape of CDK1 and CDK2 suggest a mechanism for achieving selective CDK
inhibition. *Cell Chem Biol.* 26(1):121-130.

Yang JY, Yan RX, Roy A, Xu D, Poisson J, Zhang Y. 2015. The I-TASSER Suite: Protein structure and function
prediction. *Nat Methods* 12(1):7-8.

Yang ZH. 2007. Paml 4: Phylogenetic analysis by maximum likelihood. *Mol Biol Evol*. 24(8):1586-1591.

Yuan K, Seller CA, Shermoen AW, O'Farrell PH. 2016. Timing the *Drosophila* mid-blastula transition: A cell cycle-
centerec view. *Trends Genet*. 32(8):496-507.

Zhang JZ, Rosenberg HF, Nei M. 1998. Positive darwinian selection after gene duplication in primate ribonuclease
genes. *Proc Natl Acad Sci U S A*. 95(7):3708-3713.

808
809
F_{IG}. 1 Amplification of CDK1 paralogs in the *Oikopleura* genus. (*A*) Left panel: Maximum likelihood inference analysis
of selected chordate CDK1 proteins using the kinase domain of CDK1s with bootstrap values indicated at nodes. bn/bs
values in all branches were calculated using a modified Nei-Gojobori method (Zhang et al. 1998). Blue branches have
significant bn/bs values over 1 as determined by one-sided Z tests: lineage subtending CDK1a, P = 0.0386;   lineage
subtending CDK1b, P = 0.0305, lineage subtending CDK1c, P = 0.0334. Modifications in the highly conserved PSTAIRE
domain of each CDK1 are shown. The codons resulting in the alanine to serine substitution within the Oikopleuridae (dark
grey box) are also indicated. Codons for serine with phase 2 intron insertions (splitting codons between the second and
third nucleotides) are denoted by "2". (*B*) Maximum likelihood inference analysis of selected chordate Cyclin B proteins.
The Oikopleuridae paralogs are indicated by a grey box. (*C*) CDK1 paralog localizations on chromosomes among
appendicularians. Arrows indicate possible chronological duplication steps and chromosomal translocations consistent with
the phylogenetic analyses. (*D*) Complementation studies using budding yeast cdc28-4ts mutants substituted with *O. dioica*
CDK1 paralogs. CDK1a, b and d paralogs were able to complement whereas c and e did not. (*E*) Alternative *CDK1d*
transcription start sites (TSS) in testes versus ovaries. CDK1d maternal transcripts have a broad TSS before the ATG start
codon. Testes *CDK1d* transcripts originated from a single TSS immediately before the AG splice acceptor site of fist intron,
42 nt downstream of the TCTAGA male-specific promoter element, which is also located within the first intron. Male
CDK1d transcripts lack an ATG start codon, and yield nonsense transcripts in all three possible reading frames. Vertical
axis scale for CAGE data (Danks et al. 2018) is in reads per million. P = P value.

835 **F**ɪɢ. **2. CDK1 and Cyclin B paralog expression profiles.** (*A*) *CDK1-Cyclin B* paralog expression

836 levels during gonad maturation from Day 3 to spawning. Mean ± s.d. based on three independent

837 developmental time courses. Expression data were normalized to transcription of the EF1b and Rpl23

838 housekeeping genes. Levels are normalized to maximum levels. (*B*) Western blots of CDK1-Cyclin

839 B paralog levels during development, gonad maturation and embryogenesis.

Fɪɢ. 3. *CDK1a*, *b*, and *d* knockdowns generate abnormal embryonic phenotypes. (*A*) *CDK1a* RNAi delays embryonic divisions and tadpoles fail to form. Left panel: efficient (P = 0.0058, n1 = 3, n2 = 3, one-sided, unpaired Welch's t-test) knockdown of *CDK1a* in oocytes spawned from ovaries injected with dsRNA against *odCDK1a* at D5. Mid panel: When exposed to wild-type sperm, *CDK1a* deficient oocytes generated embryos that divided more slowly than wild-type and failed to hatch at 4 hpf (P = 0.0002, n1 = 3, n2 = 3, one-sided, unpaired Welch's t-test). Right panel: images of delayed development in CDK1a deficient embryos. When wild-type embryos had developed to the late tailbud stage, CDK1a deficient embryos had only begun to gastrulate. At 7 hpf, tadpoles were observed in wild-type whereas CDK1a deficient embryos failed to reach the tailbud stage. The legend in (*A*) also applies to the respective panel in (*B*). (*B*) Upper left panel: efficient (P = 0.00006, n1 = 3, n2 =4, one-sided, unpaired Student t-test) knockdown *CDK1b* in oocytes spawned from ovaries injected with dsRNA against *odCDK1b* at D5. Upper mid panel: When exposed to wild-type sperm, *CDK1b* deficient oocytes generated embryos that failed to hatch at 4 hpf (P = 0.0003, n1 = n2 = 3, one-sided, unpaired Welch's t-test). Upper right panel: Western blot showing absence of CDK1b protein in *CDK1b* RNAi embryos. Bottom left panel: CDK1b deficient embryos arrested before gastrulation. Lower right panel: distribution of developmental stages in *CDK1b* deficient and wild type embryos assessed at 1 hpf. (*C-E*), Knockdown of *CDK1d* generated 3 different phenotypes at 1 hpf. Left panels reveal efficient knockdown ((*C*) P = 0.0005, one-sided, unpaired Student's t-test; (*D*) P = 0.0039, n1 = 3, n2 = 3, one-sided, unpaired Student's t-test; (*E*) P = 0.000008, n1 = 3, n2 = 3, one-sided, unpaired Student's t-test), in oocytes spawned from ovaries that had been injected with dsRNA against *odCDK1d* at D5. Right panels revealed abnormal division ((E) P = 0.0021, n1 = 3, n2 = 3, one-sided

unpaired Student's t-test) in all cases with a subset of embryos exhibiting more severe phenotypes of polar body extrusion but no cell division ((*D*), P = 0.0034, n1 = 3, n2 = 3, one-sided, unpaired Welch's t-test), or a complete lack of polar body extrusion and absence of any division ((*C*), P = 0.0019, n1 = 3, n2 = 3, one-sided, unpaired Welch's t-test). The legend in (*E*) also applies to the respective panels in (*C*) and (*D*). (*F*) Knockdown of CDK1d did not have any effect on the levels of Cyclin Ba, the Cyclin component of MPF in *O. dioica* oocytes (Feng and Thompson 2018). (*G*) Image comparisons of wild-type to *CDK1d* knockdown oocyte/embryo phenotypes 1 h post exposure to wide type sperm. Left to right: WT cleavage stage embryos, infertile oocytes with no polar body extrusion, polar bodies extruded, but no division, and abnormally dividing embryos. Where indicated, data are mean ± s.d.; ***P<0.001, **P<0.01, *P<0.05; P, P-values. Scale bars: 50 µm.

FIG. **4. CDK2 is dispensable for *Oikoplerua dioica* embryogenesis. Synergistic defects of *CDK2* + *CDK1a* RNAi and**
***CDK2* + *CDK1b* RNAi in embryonic divisions.** (*A*) CDK2 is dispensable for embryonic divisions. Right panel: efficient
knockdown of *CDK2* in oocytes (P = 0.0081, n1 = 2, n2 = 3, one-sided, unpaired Welch's t-test) and tadpoles (P = 0.0258,
n1 = 3, n2 = 3, one-sided, unpaired Welch's t-test), spawned from ovaries injected with dsRNA against *odCDK2* at D4.
Mid panel: Western blot showing absence of CDK2 protein in *CDK2* RNAi oocytes and tadpoles. Right panel: no
significant differences in developmental outcome between *CDK2* deficient and wild-type embryos (P = 0.1247, n1 = 3, n2
= 3, two-sided, unpaired Student's t-test). (*B*) Synergistic effect of *CDK2* + *CDK1a* RNAi on embryonic divisions. Left
Panel: efficient knockdown of *CDK1a* (P = 0.0180, n1 = 3, n2 = 3, one-sided, unpaired Welch's t-test) and *CDK2* (P =
0.0119, n1 = 3, n2 = 3; one-sided, unpaired Welch's t-test) in oocytes from ovaries that had been injected with dsRNA
against *odCDK2* and *CDK1a* at D5. Mid panel: Western blot showing absence of CDK2 protein in CDK1a + *CDK2* RNAi
in oocytes. Right panel: at 45 min pf, *CDK1a* + *CDK2* deficient zygotes were arrested prior to pronuclear fusion. Scale
bars: 50 µm. (*C*) Synergistic effect of *CDK1a* + *b* RNAi on embryonic division. Upper left panel: efficient knockdown of
CDK1a (P = 0.00007, n1 = 3, n2 = 3, one-sided, unpaired Student's t test) and *CDK1b* (P = 0.00001, n1 = 3, n2 = 3, one-
sided, unpaired Student's t-test) in oocytes from ovaries that had been injected with dsRNA against *CDK1a* + *b* at D5.
Lower left: Western blot showing absence of CDK1b protein in *CDK1a* + *b* RNAi oocytes. Right panel: *CDK1a* + *b*
deficient zygotes finished 3 rounds of slowed divisions before arresting at the 4th-5th embryonic division. Scale bars: 50
µm. Where indicated, data are mean ± s.d. ***P<0.001, **P<0.01, *P<0.05. (*D*) Confocal images of Lamin 1 and DNA
staining of wild-type versus *CDK1a* + *b* RNAi embryos at 80 min post-fertilization. *CDK1a* + *b* deficient zygotes arrest at
prophase with super-condensed chromosomes at the nuclear periphery (inset) after the third division. Scale bars: 20 µm.
928

932
933
934

FIG. 5. Cyclin Ba, is required for *O. dioica* embryonic cell cycles, whereas Cyclin Bc is dispensable. (*A*) Cyclin Bc is dispensable for embryonic divisions. Left panel: efficient knockdown *cycBc* in oocytes (P = 0.006, n1 = 3, n2 = 3, one-sided, unpaired Welch's t-test) and tadpoles (P = 0.0155, n1 = 2, n2 = 3, one-sided, unpaired Welch's t-test) spawned from ovaries that had been injected with dsRNA against *cycBc* at D5. Mid panel: no significant difference in mean H1 kinase activity between Cyclin Bc deficient and wild-type oocytes (metaphase I arrested). (P = 0.2971, n1 = 3, n2 = 3, two-sided, unpaired Student's t test). Upper right panel: Western blot showing that knockdown of *cycBc* had no effect on protein levels of Cyclin Ba. Lower right panel: images show Cyclin Bc deficient and wild-type embryos had identical phenotypes at the tailbud stage, 4 hpf. (*B*) Efficient knockdown of *cycBa* in oocytes (left panel, P = 0.0121, n1 = 3, n2 = 3; one-sided, unpaired Welch's t-test; Mid-right panel, P = 0.0123, n1 = 3, n2 = 3, one-sided, unpaired Welch's t-test) spawned from ovaries that had been injected with dsRNA against *cycBa* at D5. Cyclin Ba deficient oocytes exposed to wild-type sperm generated two embryonic phenotypes: infertile (P = 0.00003, n1 = 3, n2 = 3, one-sided, unpaired Welch's t-test) and abnormal divisions (P = 0.0143, n1 = 3, n2 = 3, one-sided, unpaired Welch's t-test). Abnormal division phenotypes, compared to wild-type, are illustrated in the bottom panels at 1 hpf. Where indicated, data are mean± s.d. ***P<0.001, **P<0.01, *P< 0.05. (*C-E*) CycBa-CDK1d complexes have histone H1 kinase activity. (*C*) H1 kinase activity assay was present in wild type oocytes but absent in *cycBa* RNAi knockdown oocytes (Feng and Thompson 20118). (*D*) Cyclin Ba immunoprecipitates from oocytes had H1 kinase activity whereas IgG oocyte immunoprecipitates did not. (*E*) H1 kinase activity levels of *O. dioica* Cyclin Ba pulldowns compared to IgG pulldowns and *in vitro* expressed complexes of Human (Hs) CDK1:Cyclin B1 and *O. dioica* (Od) CDK1d:Cyclin Ba. Scale bars: 50 µm.

958
F_{IG}. 6. CDK1d levels oscillate during early embryonic cell cycles. (*A*) Cyclin Ba levels do not oscillate but instead, decline steadily
during early embryogenesis. Upper panel: Western blot of Cyclin Ba during the first 4 h of development with ATPase beta as a loading
control. Lower panel: quantification of Cyclin Ba levels, normalized to levels in unfertilized oocytes. Data are mean ± s.d. for n = 3. (*B*)
Upper panel: Western blot of CDK1d levels during the first 4 h of development with histone H3 as a loading control. Lower panel: *CDK1d*
mRNA levels were quantified during early embryonic development by qRT-PCR, normalized to levels in unfertilized oocytes. (*C-E*)
CDK1d levels peaked in M phase of first two mitotic embryonic cell cycles. (*C*) Western blot of CDK1b, c, and d levels during the first 3
hpf. Arrow indicates the CDK1d band. I, interphase; M, M-phase. (*D*) Sampling of CDK1d levels at 1-minute internals around M phase
of the first embryonic cell cycle. Arrow indicates the oscillation of the CDK1d band. (*E*) Sampling of CDK1d levels around M-phase of
the second division cycle. Arrow indicates the oscillation of the CDK1d band. (*F*) Western blot analyses of Cyclin Ba levels and histone
H1 kinase activity during the first two embryonic cell cycles. Cyclin Ba levels remained elevated, whereas H1 kinase activity oscillated,
with peaks in the first two M-phases. (*G*) Quantification of Cyclin Ba, CDK1d and H1 kinase activity during the first two embryonic cell
cycles. (*H*) Oocytes were incubated in artificial sea water with 35S methionine for 1 h before fertilization. Embryos were collected at 3
min intervals during the course of first embryonic cell cycle. The autoradiograph shows *de novo* synthesized protein during the course of
this cycle. Arrow indicates the CDK1d band.

F_{IG}**. 7. Sequential activation of CDK1a, b and d, and switching from oscillations of CDK1d levels to oscillation of Cyclin Ba levels, drive rapid early embryonic cell cycles in *O. dioica*.** Knockdowns revealed that sequential activation of 3 *O. dioica* CDK1 paralogs, a, b and d, are required for proper execution of embryonic divisions leading to embryo hatching. Cyclin Ba, the partner of CDK1d, is maternally stocked and gradually degrades over the first embryonic cycles, whereas unusually, it is CDK1d protein levels that initially oscillate to regulate rapid M-phases. CDK1a and b levels remain constant during this period. During the maternal to zygotic transition, the cell cycle slows and divisions become less synchronous. In the coenocystic ovary, polyploid nurse nuclei are afforded the time to produce maternally stocked transcripts, many arising from genes containing multiple introns of variable length, with the genes often organized in polycistronic operons (Danks et al. 2015). During the period of rapid embryonic cell divisions, many zygotically activated genes are monocistronic, and contain short introns, or are intronless.