Title

Insights into the origin of metazoan multicellularity from predatory unicellular relatives of

animals

Authors

Denis V. Tikhonenkov[1,2*], Elisabeth Hehenberger[3], Anton S. Esaulov[4], Olga I. Belyakova[4], Yuri

A. Mazei[5], Alexander P. Mylnikov[1,†], Patrick J. Keeling[2]

Affiliations

[1]Papanin Institute for Biology of Inland Waters, Russian Academy of Sciences, Borok, 152742,

Russia

[2]Department of Botany, University of British Columbia, Vancouver, V6T 1Z4, British

Columbia, Canada.

[3]GEOMAR - Helmholtz Centre for Ocean Research Kiel, Division of Experimental Ecology,

Duesternbrookerweg 20, 24105 Kiel, Germany

[4]Penza State University, Penza, 440026, Russia

[5]Moscow State University, Moscow, 119991, Russia

†Deceased 30 May 2019

*Correspondence to: tikho-denis@yandex.ru, pkeeling@mail.ubc.ca

Abstract

The diversity and biology of unicellular relatives of animals has strongly informed our

understanding of the transition from single-celled organisms to the multicellular Metazoa. Here

we analyse the cellular structures and complex life cycles of the novel unicellular holozoans

Pigoraptor and *Syssomonas* (Opisthokonta). Both lineages are characterized by complex life

cycles with a variety of cell types, the formation of multicellular aggregations and syncytium-

like structures, and an unusual diet for single-celled opisthokonts (partial cell fusion and joint

sucking of a large eukaryotic prey), all of which provide new insights into the origin of

multicellularity in Metazoa. The ability to feed on large eukaryotic prey could have been a

powerful trigger in the formation and development both aggregative (e.g., joint feeding, which

also implies signalling) and clonal (e.g., hypertrophic growth followed by palintomy)

multicellular stages that played important roles in the emergence of multicellular animals.

Introduction

The origin of animals (Metazoa) from their unicellular ancestors is one of the most

important evolutionary transitions in the history of life. Questions about the mechanisms of this

transformation arose about 200 years ago, but is still far from being resolved today. Most investigations on the origin of Metazoa have focused on determining the nature of the shared, multicellular ancestor of all contemporary animals (Moroz et al., 2014; Srivastava et al., 2008; 2010). However, even the branching order of early, non-bilaterian lineages of animals on phylogenetic trees is still debated: some consider either sponges (Porifera) (Feuda et al., 2017; Philippe et al., 2009; Simion et al., 2017;) or Ctenophora (Dunn et al., 2008; Ryan et al., 2013; Whelan et al., 2015) or Placozoa (Schierwater et al., 2009; Signorovitch et al., 2007;) to be the first branch of extant metazoans. While molecular clock-based studies and paleontological evidence indicate that multicellular animals arose more than 600 million years ago (Maloof et al., 2010; Sharpe et al., 2015), we know less about how animals arose. To establish the sequence of events in the origin of animals from unicellular ancestors, we also need to investigate their closest relatives, the unicellular opisthokont protists. Information on the diversity and biology of the unicellular relatives of animals, their placement within the phylogenetic tree of opisthokonts, and the identification of molecular and morphological traits thought to be specific for animals within their unicellular sisters, have all strongly informed our understanding of the transition from single-celled organisms to the multicellular Metazoa (King et al., 2008; Suga et al., 2013; Suga, Ruiz-Trillo, 2013; Torruella et al., 2015).

Until recently, only three unicellular lineages, the choanoflagellates, filastereans, ichthyosporeans, as well as *Corallochytrium limacisporum*, a mysterious marine osmotrophic protist described in association with corals, have been described as collectively being sisters to animals. Together with animals they form the Holozoa within the Opisthokonta (Aleshin et al., 2007; Lang et al., 2002; Torruella et al., 2015). These unicellular organisms have extremely variable morphology and biology. Choanoflagellates represent a species-rich group of filter-feeding, bacterivorous, colony-forming protists, which possess a single flagellum surrounded by a collar of tentacles (microvilli). They are subdivided into two main groups – the predominantly marine Acanthoecida and the freshwater and marine Craspedida (Carr et al., 2017). Filastereans are amoeboid protists producing pseudopodia. Until recently, they were represented by only two species: the endosymbiont of a freshwater snail, *Capsaspora owczarzaki*, and the free-living marine heterotroph, *Ministeria vibrans* (Hertel et al., 2002; Shalchian-Tabrizi et al., 2008), which was recently shown to also possess a single, real flagellum (Mylnikov et al., 2019). Ichthyosporeans are parasites or endocommensals of vertebrates and invertebrates characterized by a complex life cycle, reproduction through multinucleated coenocytes colonies and flagellated and amoeboid dispersal stages (Suga, Ruiz-Trillo, 2013). *Corallochytrium* is a unicellular coccoid organism, which produces rough, raised colonies and amoeboid limax-like spores

(Raghukumar, 1987.). Additionally, molecular data predict a cryptic flagellated stage for
Corallochytrium (Torruella et al., 2015).

A large number of hypotheses about the origin of multicellular animals have been
proposed. The most developed model for the origin of metazoan multicellularity is based on a
common ancestor with choanoflagellates (James-Clark, 1866; Ivanov, 1967; King et al., 2008,
Mikhailov et al., 2009; Nielsen, 1987, Ratcliff et al., 2012). This idea was initially based on the
observed similarity between choanoflagellates and specialized choanocyte cells in sponges.
Molecular investigations also supported the idea by consistently indicating that choanoflagellates
are the closest sister group to Metazoa. However, molecular phylogeny itself does not reveal the
nature of ancestral states, it only provides a scaffolding on which they might be inferred from
other data. The evolutionary positions of the other unicellular holozoans (filastereans,
ichthyosporeans, and *Corallochytrium*) are less clear and sometimes controversial (e.g. Cavalier-
Smith, Chao, 2003; del Campo, Ruiz-Trillo, 2013; Hehenberger et al., 2017; Medina et al., 2003;
Ruiz-Trillo et al., 2008; Shalchian-Tabrizi et al., 2008; Torruella et al., 2012, 2015;).

As noted above, many molecular traits that were thought to be "animal-specific" are now
known to be present in unicellular holozoans, while conversely the loss of other traits have been
shown to correlate with the origin of the animals. But gene content alone is not sufficient to
provide a comprehensive understanding of the cell biology, life cycle and regulation capabilities
of the unicellular ancestor, it requires also analysis of the biology of the extant unicellular
relatives of animals (Sebé-Pedrós et al., 2017).

Recently we described a phylogenomic and transciptome analyses of three novel
unicellular holozoans (Hehenberger et al., 2017), which are very similar in morphology and life
style but not closely related. *Pigoraptor vietnamica* and *Pigoraptor chileana* are distantly related
to filastereans, and *Syssomonas multiformis* forms a new phylogenetic clade, "Pluriformea", with
Corallochytrium. Both new genera of unicellular holozoans form the shortest and most slowly
evolving branches on the tree, which improved support for many nodes in the phylogeny of
unicellular holozoans. Also, comparison of gene content of the new taxa with the known
unicellular holozoans revealed several new and interesting distribution patterns for genes related
to multicellularity and adhesion (Hehenberger et al., 2017).

Here we report the detailed morphological and ultrastructural analyses of these new
species, as well as describing their life cycle in culture, which are important implications for
understanding the origin of animals as are the genetic analyses. All three species are shown to be
predatory flagellates that feed on large eukaryotic prey, which is very unusual for unicellular
Holozoa. They also appear to exhibit complex life histories with several distinct stages,
including interesting multicellular structures that might offer important clues as to precursors of

multicellularity. On the basis of these findings we discuss the current hypotheses about the origin
of multicellular animals from their unicellular ancestors.
106
107   **Results and discussion**
108
109   Detailed morphological descriptions of the cells and their aggregates are presented below.
110
111   *Syssomonas multiformis* Tikhonenkov, Hehenberger, Mylnikov et Keeling 2017
112
Morphology and life cycle
The organism is characterized by a large variety of life forms including flagellates,
amoeboflagellates, amoeboid non-flagellar cells, and spherical cysts. The most common stage in
the life cycle, a swimming flagellate cell, resembles a typical opisthokont cell, reminiscent of
sperm cells of most animals and zoospores of the chytrid fungi. Cells are round-to-oval and
propel themselves with a single, long posterior flagellum (Fig. 1A-C, X). The flagellum is
smooth and emerges from the middle-lateral point of the cell, turns back and always directs
backwardduring swimming. The cell rotates during swimming (Video 1). Flagellar beating can
be very fast, which can create the appearance of two flagella. Motile flagellates can suddenly
stop and change the direction of movement. The flagellated cells measure 7-14 µm in diameter.
The flagellum length is 10-24, rarely 35 µm. Cyst diameter is 5 µm (Fig. 1D, Y).
Solitary cells of *Syssomonas* can temporary attach to the substrate by the anterior part of
the cell body. They produce water flow by rapid flagellum beating posteriorly and in that state
resemble cells of choanoflagellates or choanocytes from sponges (Fig. 1E, Video 2).Floating
flagellated cells can also move to the bottom and transform to amoeboflagellates (Fig. 1J, Video
3) by producing both wide lobopodia and thin short filopodia. Flagellar beating becomes slower
and then stops. Amoeboflagellates crawl along the surface using their anterior lobopodia and can
take up clusters of bacteria. The organism can lose the flagellum via three different modes: the
flagellum may be abruptly discarded from its proximal part of the cell; a stretched flagellum may
be retracted into the cell; the flagellum may convolve under the cell-body and then retract into
the cell as a spiral (Video 4). As a result *Syssomonas* turns into an amoeba (Fig. 1K,L, Video 4).
Amoeboid cells produce thin, relatively short filopodia and sometimes have two contractile
vacuoles. Amoeboid cells are weakly motile. The transformation of amoeboflagellates and
amoebae back to flagellates was also observed.
Amoeboid cells can also retract their filopodia, become roundish and transform into a cyst
(Fig. 1D, Video 5). Palintomic divisions may occur inside the cyst and up to 16 (2, 4, 8, or 16)

flagellated cells are released as a result (Fig. 1M, Video 6). Division into two cell structures was also observed in culture (Video 7), but it is hard to tell whether a simple binary longitudinal division of a *Syssomonas* cell with retracted flagellum has taken place, or the final stage of a division inside the cyst has been observed.

Fig.1. External morphology and life forms of *Syssomonas multiformis.* A-C – swimming flagellated cells; D – cyst; E – attached flagellated cell; F-H – sucking of eukaryotic prey; I – simultaneous joint feeding of three cells of *Syssomonas* on one prey cell with attaction of other speciment to the feeding spot; J – amoeboflagellate; K,L – amoeboid cell; M – palintomic cell-division inside the cyst; N – cell with inside vesicles; O – cyst with vesicles; P,Q – cells of *Syssomonas* engulfed starch granules (bright field (P) and fluorescent microscopy, DAPI staining); R – cells of *Syssomonas* with engulfed starch granules hiding into the starch druse, S-U, W – cell aggregations of *Syssomonas* near the bottom of Petri dish; V – floating aggregation of flagellated cells; X – general view of flagellated cell (SEM), Y – cyst (SEM). Scale bars: A-P, S-W – 10 µm, R – 100 µm, X – 3 µm, Y – 2 µm.

Floating, flagellated cells containing vesicular structures were observed (Fig. 1N, Video 8), however the process of formation and the purpose of these vesicles is unknown. After some time

such cells lose their flagellum and transform into vesicular cysts with a thick cover (Fig. 1O).
Division inside vesicular cysts was not observed within 10 days of observation. Such structures
could represent resting cysts or dying cells containing autophagic vacuoles (the partial
destruction of one such cyst was observed after 4 days of observation, see Video 8).

The organism is a predator; it takes up other flagellates (e.g. *Parabodo caudatus* and
Spumella sp.) which can be smaller, about the same size, or larger than *Syssomonas*. But in
contrast to many other eukaryotrophic protists, *Syssomonas* does not possess any extrusive
organelles for prey hunting. After initial contact, *Syssomonas* attaches to the prey cell and sucks
out their cytoplasm (without ingesting the cell membrane) (Fig. 1F-H, Video 9). The organism
feeds better on inactive, slow moving or dead cells and can also capture intact prey cells and
cysts by means unobserved. After attaching to the prey, many other *Syssomonas* cells become
attracted to the same prey cell (likely by chemical signaling) and try to attach to it. Joint feeding
was observed: several cells of *Syssomonas* can suck out the cytoplasm of the same prey cell
together (Fig. 1I, Video 9).

In culture, *Syssomonas* can take up starch granules from rice grains, the granules can be the
same size as the cells (Fig. 1P,Q). In the presence of *Syssomonas* cells, rice grains in Petri dishes
crumble into small fragments and separate granules of starch (Fig. S1). Cells of *Syssomonas* with
engulfed starch granules can hide within the starch druses and lose the flagellum (Fig. 1R).
Numerous cysts integrated into the starch matrix were often observed in culture.

A   **B**
Fig. S1. Rice grain destruction in Petri dish with Pratt medium and presence of the cells of
Parabodo caudatus (prey) only (A) and *Syssomonas multiformis* (B) after 9 days of
incubation.
180

The organism can also feed on clusters of bacteria (Video 10) using short pseudopodia.
After feeding, *Syssomonas* cells become 2-3 times bigger and a large food vacuole is formed at
the posterior end of the cell-body (Fig. 1C). In the absence of eukaryotic prey (cultivation on

bacteria and/or rice grain/starch only), *Syssomonas* either dies or forms resting cysts. Bacteria
alone are not sufficient food for *Syssomonas*.

Solitary cells of *Syssomonas* can partially merge and form temporary cell aggregations.
They are usually shapeless, observed near the bottom, and consist of about 3-10 flagellated or
non-flagellated cells (Fig. 1S-U, Video 11). Another type of aggregation is formed by only
flagellated cells with outwards-directed flagella that can float in the water column and resemble
the rosette-like colonies of choanoflagellates (Fig. 1V, Video 12). Both types of aggregations
break up easily and it seems that the membranes of such aggregated cells are not fused.

However, in rich culture, solitary cells of *Syssomonas* can sometimes merge completely at
the bottom of the Petri dish and form syncytium-like (or pseudoplasmodium) structures (it seems
that nuclei do not merge after cell fusion). The budding of young flagellated daughter cells from
such syncytia was observed (Video 13).Such syncytial structures with budding daughter cells
have not been observed in other eukaryotes, to our knowledge, but multinucleated structures
arising as a result of multiple cell aggregations or fusions of uninuclear cells are also known in
Dictyostelia (Eumycetozoa) and *Copromyxa* (Tubulinea) in the Amoebozoa (sister group of
Opisthokonta), as well as in other protists, such as Acrasidae in the Excavata, *Sorogena* in the
Alveolata, *Sorodiplophrys* in the Stramenopiles, and *Guttulinopsis* in the Rhizaria (Brown et al.,
2012). Within the opisthokonts, aggregation of amoeboid cells is only known in the sorocarpic
species *Fonticula alba* (Holomycota) (Brown et al., 2009). We should also note that a syncytium
is not an unusual cell structure in many fungi and animals; e.g. most of the cytoplasm of glass
sponges (Hexactinellida), the teguments of flatwormsas well as the skeletal muscles and the
placenta of mammals (Gobert et al., 2003; Leyset al., 2006) have a syncytial structure.

In *Syssomonas*, the processes of cells merging attracts (again, likely by chemical
signalling) many other cells of *Syssomonas,* which actively swim near aggregates or syncytium-
like structures and try to attach to them. Some of the these cells succeed to merge and the
aggregates grow.

All aggregations and syncytial-like structures do not seem to form by cell division, but
rather by a merger of the population of cells in the culture (although all cells in the clonal culture
are offspring of a single cell of *Syssomonas*).

All of the above-described life forms and cellular changes do not represent well-defined
phases of the life cycle of *Syssomonas*, but rather embody temporary transitions of cells in
culture which are reversible.

Syssomonas grows at room temperature (22°C) and can survive at temperatures from +5 to
36 °C. At high temperature (30-35 °C) the prey cells (bodonids) in culture become immobile and
roundish; *Syssomonas* actively feeds on such easily accessible cells, multiplies and produces

high biomass. In the absence of live eukaryotic prey, increasing the incubation temperature does
not lead to an increase in cell numbers. The cells grow at pH values from 6 to 11. Agitation of
culture does not lead to the formation of cell aggregates as was observed in the filasterean
Capsaspora (Sebé-Pedrós et al., 2013b).

Cell ultrastructure

The cell is naked and surrounded by the plasmalemma. The naked flagellum ends in a
short, narrowed tip – the acroneme (Fig. 1X, 2D). A single spiral or other additional elements
(e.g. a central filament typical for choanoflagellates) in the transition zone of the flagellum were
not observed (Fig. 2B, 2C). The flagellar axoneme has an ordinary structure (9+2) in section (not
shown). The flagellum can be retracted into the cell (Fig. 2E). A cone-shaped rise at the cell
surface around the flagellum base was observed (Fig. 2B, 2C). The flagellar transition zone
contains a transversal plate which is located above the cell surface (Fig. 2B).

Two basal bodies, one flagellar and one non-flagellar (Fig. 2A–C), lie approximately at a
45-90 degrees angle to each other (Fig. 2B, 2C). The flagellar root system consists of several
elements. Arc-like dense material, representing satellites of the kinetosome, is connected with
the flagellar basal body and initiates microtubules which run into the cell (Fig. 2F). Radial fibrils
originate from the flagellar basal body (Fig. 3A-C, 3G) and resemble transitional fibres. At least
two fibrils connect to the basal bodies (Fig. 2B). It can be seen from serial sections that
microtubules originate near both basal bodies (Fig. 3 A–F). Dense (osmiophilic) spots are
situated near the basal bodies and some of them initiate bundles of microtubules (Fig. 3 I,J,L).
Microtubules originating from both basal bodies singly or in the form of a fan probably run into
the cell (Fig. 2 B, Fig. 3F–K). One group of contiguous microtubules begins from the dense spot
(Fig. 3L) and goes superficially close to the plasmalemma (Fig. 3 E,F,L).

The nucleus is 2.6 µm in diameter, has a central nucleolus and is situated closer to the
posterior part of the cell (Fig. 2 A, D, Fig. 4 H). The Golgi apparatus is of usual structure and is
positioned close to the nucleus (Fig. 2B, Fig. 4A). The cell contains several mitochondria with
lamellar cristae (Fig. 4 B–D). Unusual reticulate or tubular crystal-like structures of unknown
nature were observed inside the mitochondria (Fig. 4C, D). A contractile vacuole is situated at
the periphery of the cell and is usually surrounded by small vacuoles (Fig. 4E).

A large food vacuole is usually located posteriorly or close to the cell center and contains
either remnants of eukaryotic prey, e.g. cells (paraxial flagellar rods are seen) or cysts (fibrous
cyst envelope is seen) of *Parabodo caudatus*, or starch granules (Fig. 2A, Fig. 4F–H).
Exocytosis occurs on the posterior cell end (Fig. 4I).

Thin filopodia are located on some parts of cell surface (Figs. 4D, 4J).

Storage compounds are represented by roundish (presumably glycolipid) granules 0.8 μm
in diameter (Fig. 4 A,D,J).

A flagellum or flagellar axoneme, or two kinetosomes, as well as an eccentric nucleus,
mitochondria with lamellate cristae and dense matrix, and a food vacuole with remnants of the
prey cells are all visible inside cysts containing dense cytoplasm (Fig. 4K, L).

Extrusive organelles for prey hunting were not observed in any cell type.

260
Fig. 2. General view and flagellum structure of *Syssomonas multiformis* (TEM). A –
general view of the cell section, B,C – arrangement of flagellum and basal bodies, D – twisting
of the flagellum around the cell, E – retracted flagellum axoneme inside the cell, F – basal body
of the flagellum and nearest structures.
ac – acroneme, ad – arc-like dense structure, an – flagellum axoneme, ds – dense spot, fb –
fibril, fbb – flagellar basal body, fl – flagellum, fv – food vacuole, Ga – Golgi apparatus, mct –
microtubule, mt – mitochondrion, n – nucleus, nfbb – non-flagellar basal body, nu – nucleolus,
tp – transversal plate.
Scale bars: A – 2, B – 0.5, C – 0.5, D – 2, E – J – 0.5 μm.
270

271
272   Fig. 3. Arrangement of kinetosomes of *Syssomonas multiformis*. A – F – serial sections of
273 the kinetosomal area, G–L – structures nearby the kinetosomes.
274   ds – dense spot, fb – fibril, fbb – flagellar basal body, mct – microtubule, nfbb – non-
275 flagellar basal body, rf – radial fibrils, smmt – submembrane microtubules.
276   Scale bars: A–J, L – 0.5, K – 1 μm.
277

Fig. 4. Cell structures and organelles of *Syssomonas multiformis*. A – Golgi apparatus, B – D – mitochondria, E – contractile vacuole, F – food vacuole with remnants of eukaryotic prey (*Parabodo*), flagella and paraxial rods are seen, G – food vacuole containing cyst of *Parabodo*, H – food vacuole containing starch granule, I – Exocytosis of food vacuole, J – reserve substance and filopodia, K – L – cysts.

ci – crystalloid inclusion, ct – cyst of the prey cell, cv – contractile vacuole, en – cyst envelope, fl – flagellum, fp – filopodia, fv – food vacuole, ga – Golgi apparatus, mt – mitochondrion, n – nucleus, nu – nucleolus, rs – reserve substance, sg – starch granule.

Scale bars: A – 2, B – 1, C – 0.5, D – 2, E – 2, F – 2, G – 2, H – 2, I – 2, J – 2, K – 2, L - 1 μm.

Pigoraptor vietnamica Tikhonenkov, Hehenberger, Mylnikov et Keeling 2017

Morphology and life cycle

The uniflagellated, elongated-oval cells are 5-12 µm long (Fig. 5 A,B,H,I). The flagellum

length is 9-14 µm. Saturated cells with a large food vacuole become roundish. The body plan,

movement, feeding, and growth conditions of *Pigoraptor* are identical to *Syssomonas*

multiformis, except for the feeding on starch granules, which was not observed in *Pigoraptor*.

The main stage of the life cycle is a swimming flagellated cell, which can form thin long,

sometimes branching filopodia that can attach to the substrate (Fig. 5D). Wide lobopodia were

also observed on some cells (Fig. 5E). Non-flagellate crawling amoebas were not observed.

Pigoraptor cells can retract the flagellum and become roundish. After several hours, such

spherical cells either divide into two daughter cells or turn into cysts (Fig. 5F), which stay intact

for a long period. Binary division was observed also inside the cyst (Fig. 5G), resulting in two

daughter cells that produce flagella and disperse.

306
Fig. 5. External morphology and life forms of *Pigoraptor vietnamica.* A, B, H, I – general
view of the cell (DIC and СЭМ); C – aggregation of flagellated cells; D – amoeboflagellate with
filopodia; E – cell with lobodopia; F – cyst; G – binary division.
ac – acroneme, cv – contractile vacuole, gr – groove.
Scale bars: A-G – 10; H – 4, I – 2 µm.

Cells of *Pigoraptor vietnamica* also form easily disintegrating aggregations (Fig. 5C) and

feed jointly (Video 14). The adjacent cells can partially merge during feeding. These

processesalso seem to attract many other cells of *Pigoraptor*.

Cell ultrastructure

A single, naked flagellum with an acroneme originates from a small lateral groove and
directs backward (Fig. 5 H,I). The cell is naked and surrounded by the plasmalemma. Two basal
bodies, flagellar and non-flagellar, are located near the nucleus, lie approximately at a 90 degrees
angle to each other and are not connected by visible fibrils (Fig. 6 A,B; Fig. 7 A,B; Fig. 8 A–F).
The flagellum axoneme has an ordinary structure (9+2) in section (Fig. 6 D,E).A thin central
filament, which connects the central pair of microtubules to the transversal plate, was observed
(Fig. 6 C,F).The flagellum can be retracted into the cell (Fig. 9C).The flagellar root system is
reduced. Radial fibrils arise from the flagellar basal body (Fig. 7C). Microtubules pass near the
flagellar basal body (Fig. 7B, Fig. 8 E,F). Serial sections show that the non-flagellar basal body
does not initiate the formation of microtubules (Fig. 8 A,B)

The roundish nucleus is about 1.5 µm in diameter, contains a prominent nucleolus (Fig. 6
A,B, Fig. 7A, Fig. 9 A, D), and is situated closer to the posterior end of the cell. Chromatin
granules (clumps) are scattered within the nucleoplasm. The Golgi apparatus is adjacent to the
nucleus (Fig. 9B). Cells contain several mitochondria that possess lamellar cristae (Fig. 9 A,C).
Rare thin filopodia have been observed on the cell surface (Fig. 9 D,E). Cells usually contain one
large food vacuole (Fig. 9F), which contains remnants of eukaryotic prey and bacteria.
Exocytosis takes place on the anterior cell end (Fig. 9G). Storage compounds are represented by
roundish (presumably glycolipid) granules 0.3-0.4 µm in diameter (Fig. 6A, 7A, 9 C,H). Cells
contain symbiotic bacteria, which are able to divide in the host cytoplasm (Fig. 9 H,I). A single
contractile vacuole is situated close to the cell surface (not shown on cell sections but visualized
by TEM, Fig. 5H).

339
Fig. 6. General view and flagellum structure of *Pigoraptor vietnamica*, TEM. A –
longitudinal cell section. B–C – longitudinal section of flagellum, D–F – transverse flagellum
sections in transitional area.
cf – central filament, fbb – flagellar basal body, fl – flagellum, n – nucleus, rs – reserve
substance.
Scale bars: A – 1, B – 0.5, C – 0.5, D – F – 0.2 – μm.
346

347
Fig. 7. Nucleus and arrangement of basal bodies of *Pigoraptor vietnamica*. A – part of the
cell containing nucleus and flagellar basal body, B – non-flagellar basal body, C – flagellar basal
body and surrounding structures.
fbb – flagellar basal body, mct – microtubule, n – nucleus, nu – nucleolus, rf – radial
fibrils, rs – reserve substance.
Scale bars: A – 0.5, B – 0.5, C – 0.2 μm.
354

356
357 Fig. 8. Arrangement of two basal bodies of *Pigoraptor vietnamica* relative to one another.
358 A – F – serial sections of basal bodies.
359   fbb – flagellar basal body, mct – microtubule, n – nucleus, nfbb – non-flagellar basal body.
360   Scale bars: A – F – 0.5 μm.
361

362
Fig. 9. Sections of nucleus and other call structures of *Pigoraptor vietnamica.* A – nucleus,
B – Golgi apparatus, C – mitochondrion, D – E – nucleus and filopodia, F – food vacuole, G –
exocytosis, H – reserve substance, I – dividing symbiotic bacteria.

an – flagellar axoneme, bc – bacterium, ec – ectoproct, fb – filopodium, fv – food vacuole,
ga – Golgi apparatus, mt – mitochondrion, n – nucleus, nu – nucleolus, rs – reserve substance, sb
– symbiotic bacteria.
Scale bars: A – 1, B – 0.5, C – 0.5, D – 0.5, E – 0.2, F – 1, G – 1, H – 1, I – 0.5 μm.
370

371　*Pigoraptor chileana* Tikhonenkov, Hehenberger, Mylnikov et Keeling 2017

Morphology and life cycle

The uniflagellated, roundish cells measure 6-14 μm in diameter. The flagellum emerges
from a shallow groove and is 8-16 μm in length (Fig. 10 A,B,H,I). The flagellum ends with the
acroneme. This species is identical to *Pigoraptor vietnamica* in body plan, movement, feeding,
growth conditions*,* joint feeding and aggregation behaviours (Fig. 10 C, Video 15, 16),
encystation (Fig. 10 D), binary division (Fig. 10 E), but additionally characterized by the absence
of symbiotic bacteria and much reduced capability to produce filopodia and lobopodia (Fig. 10
F,G), which are extremely rare in *Pigoraptor chileana*.

381
382　　　Fig. 10. External morphology and life forms of *Pigoraptor chileana.* A, B, H, I – general
383　view of flagellated cell (DIC and SEM); C – cell aggregation; D – cyst; E – binary division; F,G
384　– cell with short lobopodia and filipodia.
385　　　ac – flagella arconeme, gr – groove, fl – flagellum.
386　　　Scale bars: A-G – 10, H – 9, I – 4 μm.
387

Cell ultrastructure

The cell is naked and surrounded by the plasmalemma. The nucleus is positioned close to
the posterior cell end (Fig. 11 A). The flagellum is naked and the flagellar axoneme has an
ordinary structure (9+2) in section (Fig. 11 B–D). The flagellum can be retracted into the cell
which is visible in some sections (Fig. 11 A, C). Flagellar and non-flagellar basal bodies are
located near the nucleus (Fig. 11 A) and lie approximately at a 60-90 degrees angle to each other
(Fig. 12 A–F).The flagellar basal body contains a wheel-shaped structure in the proximal part
(Fig. 11 E,F). Single microtubules and microtubule bundles are situated near this basal body
(Fig. 11 E–H). Some microtubules arise from dense spots close to the basal body (Fig. 11 H).

Rare, thin, sometimes branching filopodia may contain superficially microtubule-like

profiles (Fig. 12 G–I). The roundish nucleus is about 1.5 μm in diameter and has a central

nucleolus (Fig. 11A). Chromatin granules are scattered within the nucleoplasm. The Golgi

apparatus was not observed. Mitochondria contain lamellar cristae and empty space inside (Fig.

13 A, B). Cells usually contain one large food vacuole (Fig. 11A, Fig. 13 C), which contains

remnants of eukaryotic prey and bacteria. Storage compounds are represented by roundish

(presumably glycolipid) granules 0.2-0.4 μm in diameter (Fig. 13 C). The single ultrathin section

of the dividing cell (possible open orthomitosis) was obtained in metaphase stage (Fig. 13 D).

406
Fig. 11. General view, flagellum and flagella root system of *Pigoraptor chileana.* A –
general view of the cell section. B – D – flagellum, E– H – flagellar basal body and surrounding
structures.
an – axoneme, ds – dense spot, fbb – flagellar basal body, fv – food vacuole, mct –
microtubule, n – nucleus, nfbb – non-flagellar basal body, nu – nucleolus.
Scale bars: A – 0.5, B – 0.2, C – 0.2, D – 0.5, E – 0.5, F – 0.5, G – 0.5, H – 0.5 μm.

413
414

417   Fig. 12. Arrangement of basal bodies and structure of filopodia of *Pigoraptor chileana.* A
418 – F – serial sections of basal bodies, G – I – filipodia.
419   fbb – flagellar basal body, fl – flagellum, fp – filopodium, mct – microtubule, n – nucleus,
420 nfbb – non-flagellar basal body.
421   Scale bars: A – F – 0.2, G – I – 0.5 μm.
422

424
425   Fig. 13. Mitochondria, food vacuole and nucleus division of *Pigoraptor chileana.* A, B –
426 mitochondria, C – food vacuole and exocytose, D – nucleus division in metaphase stage.
427   chr – chromosomes, fv – food vacuole, mct – microtubule, mt – mitochondrion, rs –
428 reserve substance.
429   Scale bars: A – 0.5, B – 0.5, C – 0.5, D – 1 μm.
430

Key features of novel unicellular opisthokonts and origin of multicellularity in Metazoa

Our understanding of the origin and early evolution of animals has transformed as a result of the study of their most closely related sister groups of unicellular organisms: choanoflagellates, filastereans, and ichthyosporeans (King et al., 2008; Suga et al., 2013; Suga, Ruiz-Trillo, 2013). Our discovery of previously unknown unicellular Holozoa from freshwater bottom sediments in Vietnam and Chile provides new material for analysis.

Eukaryotrophy

A distinctive feature of all three new species is their feeding on eukaryotic prey of similar or larger size, which is unusual (if not unique) for unicellular Holozoa. While they consume entire prey cells or only the cytoplasmic contents of eukaryotic cells, which resembles the phagocytotic uptake of the contents of *Schistosoma mansoni* sporocysts by the filasterean

Capsaspora owczarzaki in laboratory conditions (Owczarzak et al., 1980), they can also feed on

clusters of bacteria, resembling the phagocytic uptake of bacteria by choanoflagellates (Dayel,

King, 2014). It is particularly noteworthy that the organisms we discovered, do not possess

extrusive organelles for paralyzing and immobilizing the prey, which is typical for

eukaryovorous protists. Our observations show that, prior to absorption, they somehow adhere to

the surface of the prey cell. Studies on the choanoflagellate *Monosiga brevicollis* have shown

that cadherins, that function as cell-cell adhesion proteins in animals, are located on the

microvilli of the feeding collar and colocalize with the actin cytoskeleton (Abedin, King, 2008).

M. brevicollis is non-colonial, thus suggesting that cadherins participate in prey capture, not

colony formation. In addition, studies of the colonial choanoflagellate *Salpingoeca rosetta* did

not indicate a role of the cadherins in colony formation, further supporting the notion that

cadherins do not play a role in cell-cell adhesion between choanoflagellates, and perhaps also did

not in the unicellular ancestor of animals (Sebé-Pedrós et al., 2017). In the case of the unicellular

predators *Syssomonas* and *Pigoraptor*, adherence to a large and actively moving prey seems to

be crucial for feeding and important for survival. Interestingly, the *Syssomonas* transcriptome

does not include cadherin genes, but it does express C-type lectins (carbohydrate-binding

proteins performing various functions in animals, including intercellular interaction, immune

response and apoptosis). A reverse pattern of gene distribution is seen in *Pigoraptor*, where

cadherin domain-containing transcripts were found but no C-type lectins (Hehenberger et al.,

2017).

The presence of eukaryotrophy as a type of feeding within both filastereans and

Pluriformea suggests that predation could be or have once been widespread among unicellular

relatives of animals, and perhaps that the ancestor of Metazoa was able to feed on prey much

larger than bacteria. The "joint feeding" we observed many times in cultures of *Syssomonas* or

Pigoraptor, including the behaviour where cells are attracted to the large prey by other predators

already feeding on it, is probably mediated by chemical signaling of the initially attached

predator cell. The newly arriving cells also adhere to the plasmalemma of the prey, partially

merging with each other and sucking out the contents of the large prey cell together. The

merging of predator cells during feeding is quite unusual, and may represent a new factor to

consider in the emergence of aggregated multicellularity. In addition, putative chemical

signaling to attract other cells of its species is observed during the formation of syncytial

structures in these species. In this context, alpha and beta-integrins and other components of the

so-called integrin adhesome, which are responsible for interaction with the extracellular matrix

and the transmission of various intercellular signals, were found in the transcriptomes of all three

studied species (Hehenberger et al., 2017).

Starch breakdown by Syssomonas

An interesting phenomenon was observed in clonal cultures of *Syssomonas*, where the

predator can completely engulf starch granules of the same size as the cell, also mediating the

rapid destruction of rice grains into smaller fragments and individual starch crystals (Fig. S1). It

is possible that *Syssomonas* secretes hydrolytic enzymes that provide near-membrane

extracellular digestion. Near-membrane extracellular digestion in animals is of great importance

for the breakdown of various biopolymers and organic molecules (for example, in the intestinal

epithelium of mammals in the zone of limbus strigillatus in the glycocalix layer).

This ability of *Syssomonas* to feed on starch is likely promoted by the expression of

numerous enzymes that are putatively involved in starch breakdown (several α-amylases and α-

glucosidases, a glycogen debranching enzyme and a glycogen phosphorylase) (Table 1). For

example, *Syssomonas* has five distinct putative alpha-amylases, two of them not found in any

other Holozoa present in our database (Table 1). Similarly, one of the four α-glucosidases in *S.*

multiformis seems to be specific to this lineage within the Holozoa. While α-amylases and

α-glucosidases are able to hydrolyze α-1,4-linked glycosidic linkages, mobilization of the starch

molecule at the α-1,6 glycosidic bonds at branch points requires the activity of debranching

enzymes. A possible candidate for the catalysis of this reaction is a conserved glycogen

debranching enzyme in *S. multiformis*, orthologous to the human *AGL* gene (Table 1).

Additionally, we identified a transcript for a glycogen phosphorylase (orthologous to the human

PYGB, *PYGL* and *PYGM* genes), an enzyme involved in the degradation of large branched

glycan polymers.

Our observations also show that, in the presence of starch in culture, *Syssomonas* can

form resting stages of unidentified genesis, which tend to adhere to each other and to starch

grains.

Table 1. Transcripts for putative starch-degrading enzymes in *S. multiformis* and the
presence/absence of corresponding orthologs in other unicellular holozoan lineages/Metazoa.
Accession numbers for *S. multiformis* are from the corresponding TransDecoder output files as
described in Hehenberger et al., 2017. Pfam domains used to identify candidates in *S.*
multiformis and/or annotated direct human orthologous genes are indicated in brackets in column
1. NA, no direct orthologs recovered in our dataset; +, direct orthologs present in phylogeny.

	S. multiformis	Ichthyosporea	Filasterea	Choanoflagellates	Metazoa
α-amylases (PF00128)					
1.	Colp12_sorted@267, Colp12_sorted@268, Colp12_sorted@269	NA	NA	NA	NA
2.	Colp12_sorted@19603	NA	NA	NA	NA
3.	Colp12_sorted@18987, Colp12_sorted@16467, Colp12_sorted@17178	NA	+	+	+
4.= paralog to 3.	Colp12_sorted@8715	NA	+	+	+
5. (*SLC3A1*)	Colp12_sorted@10150	+	NA	NA	+
α-glucosidases (PF01055)					
1.	Colp12_sorted@9290, Colp12_sorted@22295	NA	NA	NA	NA
2.	Colp12_sorted@1339, Colp12_sorted@1341	+	+	NA	+
3.= paralog to 2.	Colp12_sorted@9196, Colp12_sorted@14409, Colp12_sorted@18957, Colp12_sorted@21457, Colp12_sorted@31517	+	+	NA	+
4. (*GANAB/GANC*)	Colp12_sorted@13767	+	+	+	+
glycogen phosphorylase (*PYGB/PYGL/PYGM*)					
1.	Colp12_sorted@1564	+	+	+	+
glycogen debranching enzyme (*AGL*)					
1.	Colp12_sorted@14615	+	+	+	+

515 ***Structural features***

516   *Syssomonas* and *Pigoraptor* both display a broad morphological plasticity: all three

517 species have a flagellar stage, form pseudopodia and cysts, and can form aggregations of several

518 cells. *Syssomonas multiformis* also has an amoeboid non-flagellar stage. The dominating life

519 form of all three species in culture is the uniflagellar swimming cell. Interestingly, amoeboid and

pseudopodial life forms were detected in cultures only after two years of cultivation and observation, suggesting they may be extremely rare in nature. Overall, the morphological differences between cells of the same type of the two genera, *Syssomonas* and *Pigoraptor*, are few and subtle. Given these genera are distantly related within the tree of Holozoa, it is interesting to speculate that they may be the result of morphostasis and by extension retain features resembling those of an ancestral state of holozoan lineages.

It has been established that single cells of *Syssomonas* and *Pigoraptor* can temporarily attach themselves to the substrate and, by beating their flagellum, can create water currents to putatively attract food particles, similar to choanoflagellates and sponge choanocytes (Fig. 1 E, Video 2). Choanocytes and choanoflagellates possess, in addition to the flagellum, a collar consisting of cytoplasmic outgrowths reinforced with actin filaments (microvilli) that serve to capture bacterial prey. The thin filopodia that are observed on the cell surface of all three *Syssomonas* and *Pigoraptor* species may thus be homologous to collar microvilli. But this will require further evidence in form of homologous proteins in these structures or evidence of their function in *Syssomonas* and *Pigoraptor*. While the filopodia of *Syssomonas* have no obvious structural contents, the outgrowths of *Pigoraptor* sometimes contain microtubular-like profiles. Cross sections of these structures were not obtained, but they may represent parallel microfilaments such as recently found in the filopodial arms of *Ministeria vibrans* (Mylnikov et al., 2019). The organisation of the *Ministeria* filopodial arms in turn resembles the microvilli of choanoflagellates, which have stable bundles of microfilaments at their base. It has been proposed previously that the ancestor of Filozoa (Filasterea+Choanoflagellida+Metazoa) probably had already developed filose tentacles, which have aggregated into a collar in the common ancestor of choanoflagellates/sponges (Shalchian-Tabrizi et al., 2008), and that microvilli were present in the common ancestor of Filozoa (Mylnikov et al., 2019).

A single, posterior flagellum is the defining characteristic of opisthokonts (Cavalier-Smith, Chao, 2003). However, the flagellum has not yet been found in all known Opisthokonta lineages. Torruella et al. (2015) have found several proteins corresponding to key components of the flagellum in *Corallochytrium* and the filose amoeba *Ministeria vibrans,* which have been considered to lack flagella. The authors have shown that the stalk used by *Ministeria* to attach to the substrate is a modified flagellum. Recently, morphological observations on another strain of *Ministeria vibrans* (strain L27; Mylnikov et al., 2019) revealed that this strain lacks the stalk for substrate attachment, but possesses a typical flagellum that projects forward and beats at attached to the substrate cells (see Fig. 2h and Video S1 in Mylnikov et al., 2019). The authors concluded that the filasterean ancestor possessed a flagellum, which was subsequently lost in *Capsaspora owczarzaki*. In the case of *Corallochytrium limacisporum*, it was suggested that it has a cryptic

flagellate stage in its life cycle (Torruella et al., 2015), as has been proposed for other eukaryotes (*Aureococcus* and *Ostreococcus*, for instance) based on their genome sequences (Wickstead, Gull, 2012). Therefore, the flagellate stage could have been the one morphological trait uniting *Corallochytrium* and *Syssomonas* within "Pluriformea". Interestingly, the ancestor of ichthyosporeans probably also had a flagellum, which is preserved in the Dermocystida (at the stage of zoospores), but was again lost in the Ichthyophonida.

The central filament of the flagellum, which connects the central pair of microtubules with the transversal plate in *Syssomonas* and *Pigoraptor*, is also noteworthy, since this character was previously known only in the choanoflagellates and was considered a unique feature for this lineage. The cone-shaped elevation of the surface membrane around the base of the flagellum in *Syssomonas* was also thought typical for choanoflagellates.

Origins of multicellularity

As mentioned above, numerous theories about the origin of Metazoa exist. One of the first and widely accepted evolutionary theories on the origin of animals is the Gastrea theory of Ernst Haeckel (Haeckel, 1874). Based on the blastula and gastrula stages that various animals undergo during their embryonic development, Haeckel suggested that the first step in the evolution of multicellularity in animals was the formation of a hollow ball, the walls of which consisted of identical flagellated cells, which he called Blastea. This stage is followed by gastrulation, where the ball invaginated, leading to the primary cellular differentiation into ecto- and endoderm. In combination with modern theories, such as the choanoblastea theory, which highlights the similarity between Haeckel's Blastea and the choanoflagellate colony (Nielsen, 2008), this model is still the most commonly used explanation for the origin of multicellular animals. An important assumption of the Gastrea theory is that cell differentiation took place only *after* multicellularity arose, suggesting that animals originated from a single cell type. This in turn generated the hypothesis that multicellular animals originated from a choanoflagellate-like colony-forming ancestor (Sebé-Pedrós et al., 2017), supported by the possible homology between sponge choanocytes and choanoflagellates (Adamska, 2016), and is consistent with the idea behind the Haeckel-Muller Biogenetic Law, that ontogenesis recapitulates phylogenesis (Hashimshony et al., 2015).

However, there are, for example, some basic differences between sponges and choanoflagellates in how their collar and flagella interact, so, though choanocytes and choanoflagellates are superficially similar, homology should not be automatically assumed (see Mah et al., 2014 for details). More importantly, recent ultrastructural studies on the structure of kinetids (flagellar apparatus) of various sponge choanocytes and choanoflagellates show that

they are fundamentally different in many respects (see Pozdnyakov, Karpov, 2016; Pozdnyakov et al., 2017, 2018 for details). These differences are significant, as the kinetid (consisting of the flagella itself, the transition zone, and the kinetosomes with attached microtubular or fibrillar roots) represents one of the very few ultrastructural systems in eukaryotic cells considered a conservative indicator of phylogenetic relationship (Lynn, Small, 1981; Moestrup, 2000; Yubuki and Leander, 2013). Choanocyte kinetids contain more elements that can be considered plesiomorphic for opisthokonts than do choanoflagellate kinetids. For example, significant differences in the spatial arrangement of the non-flagellar basal body, in the structure of the transitional zone and the flagellar root system were found when comparing the flagellate apparatus of the freshwater sponge *Ephydatia fluviatilis* (order Haplosclerida) and the choanoflagellates, but in all of these features the *Ephydatia* choanocyte kinetid is similar to the kinetid of zoospores of chytrids, which are more distantly-related Holomycota (Karpov, Efremova, 1994). Flagellated cells of some ichtyosporeans also possess ultrastructural features in common with flagellated fungi from Holomycota. Therefore, the idea that sponges and by extension all Metazoa descend directly from a single-celled organism similar to choanoflagellates is not supported by the results of the kinetid ultrastructure study (Pozdnyakov et al., 2017).

A more detailed understanding of the unicellular relatives of animals has, however, raised an alternative to the Gastrea theory. Specifically, the presence of diverse life forms in complex life cycles and the prevalence of cellular aggregations that bear little similarity to the blastula all suggest that cell *differentiation might have preceded* the origin of the blastula. Moreover, unicellular relatives of animals were shown to contain a variety of genes homologous to those involved in cell adhesion, differentiation and development, and signal transduction in Metazoa. Some of them were considered to be unique to animals (e.g. transcription factors T-box and Rel/NF-kappa B, Crumbs protein, integrin beta) as they were absent in choanoflagellates (the closest relatives of Metazoa), but later they were found to be present in other unicellular Holozoa (Mikhailov et al., 2009; Sebé-Pedrós et al., 2013a; Shalchian-Tabrizi et al., 2008). To date, it is well known that homologues of most genes controlling the development of animals, their cell differentiation, cell-cell and cell-matrix adhesion are present in various lineages of unicellular organisms (King et al., 2003; Sebé-Pedrós et al., 2016; Suga et al., 2013; Williams et al., 2014), which suggests that genetic programs of cellular differentiation and adhesion arose relatively early in the evolution of opisthokonts and before the emergence of multicellularity (King et al., 2003; Ruiz-Trillo et al., 2007; Shalchian-Tabrizi et al., 2008; Mikhailov et al., 2009; Brunet, King, 2018).

One specific idea positing that cell differentiation preceded the formation of colonies is
the "synzoospore hypothesis" (Sachwatkin, 1956; Zakhvatkin, 1949; and see Mikhailov et al.
2009 for details). In brief, three types of cell cycle alternate in the ontogenesis of multicellular
animals: monotomy (alternate phases of cell growth and division of somatic cells), hypertrophic
growth (in female sex cells) and palintomy (the egg undergoes a series of consecutive divisions).
Zakhvatkin noted that some protists alternate between different types of life cycle, and suggested
that the unicellular ancestor of Metazoa already had differentiated cells as a result of such a
complex life cycle. The life cycle complexity, in turn, results from the fact that monotomic cells
are usually sedentary, or at least less mobile, and can change their phenotype (from flagellated to
amoeboid, etc.) depending on the environment; the process of palintomy is necessary for the
formation of morphologically identical dispersal cells (spores or zoospores). These dispersal
cells remain attached to each other, forming a primary flagellated larva — the synzoospore or
blastula (cited from Mikhailov et al., 2009).

The synzoospore hypothesis is consistent with recent observations of complex life cycles
in unicellular opisthokonts possessing cellular differentiation, the presence of sedentary trophic
phases, and a tendency to aggregation; as seen in choanoflagellates (Cavalier-Smith, 2017; Dayel
et al., 2011; Dayel, King, 2014; Leadbeater, 1983; Maldonado, 2004), filastereans (Sebé-Pedrós
et al., 2013b), *Corallochytrium* (Raghukumar, 1987), ichthyosporeans (Arkush et al., 2003;
Ruiz-Trillo et al., 2007; Suga, Ruiz-Trillo, 2013), chytridiomycetes (Money, 2016), and
nucleariid amoebae (Smirnov, 2000). According to this theory, multicellularity in animals arose
through the temporal integration of various types of cells, which were already present in
different parts of the life cycle. The hypothetical ancestor of animals in this model would thus
already have genetic programs for cell differentiation (including cadherins, integrins, tyrosine
kinases).

Developing the synzoospore hypothesis further, Mikhailov et al. (2009) proposed an
evolutionary mechanism of "*transition from temporal to spatial cell differentiation*" to explain
the emergence of multicellular animals. In this model, the ancestor of Metazoa was a sedentary
colonial protist filter-feeder with colonies formed by cells of different types, which arose
because filtration efficiency is significantly enhanced through the cooperation of cells of
different types. Dispersal cells produced by the sedentary stage, the zoospores, remained
attached together in early metazoans (which increased survivability) as a synzoospore to form a
primary larva, the blastula. Development of a whole colony from such a multicellular larva
occurred through the *differentiation of genetically identical* zoospore cells. This was critical for
the maintenance of long-term cell adhesion and thus emergence of true multicellularity, as
opposite to temporary colonies and aggregations composed of genetically heterogeneous cells.

The authors suggest that the dispersal stages of the sedentary trophic body — primary blastula-
like larvae – acquired adaptations to the predatory lifestyle, which triggered the development of
primary intestine, muscular and nervous systems (Mikhailov et al., 2009).

The origin of multicellularity can in this view be seen as a *transition from temporal to*
spatiotemporal cell differentiation (Sebé-Pedrós et al., 2017). In a unicellular ancestor of
Metazoa that was a sexually reproducing bacteriotroph with many differentiated, temporally-
separated cells, the transitions between different cell states would be regulated by expression of
transcription factor families in response to environmental conditions such as availability of
nutrients or preferred bacterial food. These temporally-regulated cell types then became spatially
integrated, existing simultaneously but in different parts of a now multicellular conglomerate
with different cell types carrying out different functions. Further diversification could then be
accompanied by the evolution of additional mechanisms for complex gene regulation networks
involving signaling pathways, expansion of transcription factors, and the evolution of new
genomic regulatory mechanisms to control spatial differentiation of existing genetic modules
specific to a particular cell type. At that point, the life cycle of the protozoan ancestor of animals
probably included one or more clonal and/or aggregative multicellular stages (Sebé-Pedrós et al.,
2017).

To distinguish between the models for the origin of animal multicellularity, genomics
alone is not sufficient, and data on morphology, life cycle, and structural features of basal
holozoans is also needed. From the current analysis, all three novel species of unicellular
Holozoa have life histories that are consistent with major elements of the synzoospore model
(see Fig. 3A in Mikhailov et al., 2009 and Fig. 5a,b in Sebé-Pedrós et al., 2017). Specifically,
these organisms have complex life histories characterized by a variety of forms: flagellates,
amoebae, amoeboflagellates, cysts. All three species have the tendency to form aggregations.
Syssomonas possesses both clonal and aggregative multicellular stages, as predicted for the
ancestor of animals. Moreover, the formation of aggregations can be associated with feeding on
large eukaryotic prey, but also by cysts, which can adhere to each other and to starch grains in
culture and divide multiply. Both these probably require cellular signaling. Eating a large
eukaryotic prey, sometimes exceeding the size of a predator, also leads to hypertrophic cell
growth (described as proliferative stage in Sebé-Pedrós et al., 2017) with a subsequent phase of
palintomic division (in *Syssomonas*). All these characters are predicted by the synzoospore
model. Interestingly, many of these are apparently triggered by the behaviour of feeding on large
eukaryotic prey, highlighting this as is an interesting and potentially powerful trigger in general
for the formation and development of aggregates (e.g., joint feeding) and clonal multicellularity

(e.g., hypertrophic growth followed by palintomy), perhaps playing a role in the origin of

multicellularity in ancestors of Metazoa.

Concluding remarks

As we acquire more information about the biology of known unicellular relatives of

animals, and even more importantly, describe diverse new species of unicellular Holozoa, more

reliable models for the evolutionary histories of specific characteristics that contributed to the

emergence of multicellularity in animals are possible. *Syssomonas* and *Pigoraptor* are

characterized by complex life cycles, the formation of multicellular aggregations, and an unusual

diet for single-celled opisthokonts (partial cell fusion and joint sucking of large eukaryotic prey),

all of these features providing new insights into the origin of multicellularity in Metazoa.

Genomic and transcriptome analysis of unicellular relatives of animals have shown that

genes encoding proteins for cellular signaling and adhesion, as well as genes for embryonic

development of multicellular organisms, arose before the emergence of multicellular animals

(King et al., 2003; Ruiz-Trillo et al., 2007; Shalchian-Tabrizi et al., 2008; Hehenberger et al.,

2017). While these genes almost certainly have slightly different functions in protists than in

animals, they nevertheless probably relate to the ability to recognize the cells of their own

species, prey, or organic molecules and contribute to the formation of multicellular aggregations,

thus increasing the organism's ability to adapt to environmental change. As we learn more about

the natural history and behaviour of these organisms, the importance of these processes becomes

even more clear. The ancestor of Metazoa probably formed cells of various types that could

aggregate and had molecular mechanisms of cell differentiation and adhesion related to those

processes. Therefore, cellular differentiation likely arose before the emergence of

multicellularity.

The feeding modes of the ancestral metazoan may also have been more complex than

previously thought, including not only bacterial prey, but also larger eukaryotic cells and organic

structures. Indeed, the ability to feed on large eukaryotic prey could have been a powerful trigger

in the formation and development both aggregative and clonal multicellular stages that played

important roles in the emergence of multicellularity in animals. Lastly, we wish to point out that

other new and deep lineages of opisthokonts undoubtedly exist that have not yet been described,

and each of these will play an important role in the development of hypotheses on the origin of

multicellular animals in future.

Materials and Methods

Novel unicellular opisthokont predators were found in freshwater biotopes in Vietnam and Chile. *Syssomonas multiformis* (clone Colp-12) was obtained from the sample of freshwater pool (11°23'08.0"N, 107°21'44.9"E; T = 39°C; pH = 7.18; DO (ppm) = 0.64; conductivity (µS/cm) = 281; TDS (ppm) = 140), Tà Lài, Cát Tiên National Park, Dong Nai Province, S.R. Vietnam on April 29, 2013. *Pigoraptor vietnamica* (clone Opistho-1) was obtained from freshwater Lake Dak Minh, silty sand on the littoral (12°54′50″N, 107°48′26″E; T = 27 °C; pH=7.03; DO (ppm) = 7.43; conductivity (µS/cm) = 109; TDS (ppm) = 54), Dak Lak Province, S.R. Vietnam on March 26 2015. *Pigoraptor chileana* (clone Opistho-2) was obtained from the bottom sediments of freshwater temporary water body (submerged meadow, 54°02'29.7"S, 68°55'18.3"W; T = 16.5°C; pH = 6.62; conductivity (µS/cm) = 141; TDS (ppm) = 72) near the Lake Lago Blanca, Tierra del Fuego, Chile on November 4, 2015.

The samples were examined on the third, sixth and ninth days of incubation in accordance with methods described previously (Tikhonenkov et al., 2008). Following isolation by glass micropipette, freshwater clones Colp-12, Opistho-1, and Opistho-2 were propagated on the bodonid *Parabodo caudatus* (strain BAS-1, IBIW RAS) grown in Pratt's medium or spring water (Aqua Minerale, PepsiCo, Moscow Region, Russia or PC Natural Spring Water, President's Choice, Toronto, Canada) by using the bacterium *Pseudomonas fluorescens* as food (Tikhonenkov et al., 2014). The clone Colp-12 was perished after five years of cultivation. The clones Opistho-1 and Opistho-2 are stored in the "Live culture collection of free-living amoebae, heterotrophic flagellates and heliozoans" at the Institute for Biology of Inland Waters, Russian Academy of Science.

Light microscopy observations were made by using the Zeiss Axio Scope A.1 equipped with a DIC contrast water immersion objective (63x). The images were taken with the AVT HORN MC-1009/S analog video camera and directly digitized by using the Behold TV 409 FM tuner. Cells with engulfed starch granules were inspected by epifluorescence microscopy after DAPI staining using the Zeiss Axioplan 2 Imaging microscope.

For transmission electron microscopy (TEM), cells were centrifuged, fixed at 1 °C for 15-60 min in a cocktail of 0.6% glutaraldehyde and 2% OsO_4 (final concentration) prepared using a 0.1 M cacodylate buffer (pH 7.2). Fixed cells were dehydrated in alcohol and acetone series (30, 50, 70, 96, and 100%, 20 minutes in each step). Afterward, the cells were embedded in a mixture of Araldite and Epon (Luft, 1961). Ultrathin sections were prepared with an LKB ultramicrotome (Sweden) and observed by using the JEM 1011 transmission electron microscope (JEOL, Japan).

For scanning electron microscopy (SEM), cells from exponential growth phase were fixed

as for TEM but only for 10 min at 22 °C and gently drawn onto a polycarbonate filter (diameter

24 mm, pores 0.8 µm). Following the filtration, the specimens were taken through a graded

ethanol dehydration and acetone, and finally put into a chamber of a critical point device for

drying. Then dry filters with fixed specimens were mounted on aluminum stubs, coated with

gold-palladium, and observed with a JSM-6510LV scanning electron microscope (JEOL, Japan).

Analysis of enzymes involved in starch breakdown was based on transcriptomic data

obtained as described earlier (Hehenberger et al., 2017). To identify candidates putatively

involved in starch breakdown, we used the results of a previous hmmscan analysis of *S.*

multiformis (Hehenberger et al., 2017) to search for Pfam domains present in known starch-

degrading enzymes/enzyme families, such as α-amylases (PF00128), glycoside hydrolase

families containing α-glucosidases (PF02056, PF01055, PF03200, PF10566), α-glucan water

dikinase 1 (*GWD1*, PF01326), phosphoglucan phosphatase (*DSP4*, PF00782 and PF16561),

disproportionating enzymes (PF02446) and pullulanases (PF17967). Additionally, we submitted

the *S. multiformis* sorted transcriptome to the KEGG Automatic Annotation Server (KAAS)

(Kanehisa et al., 2014) for functional annotation and investigated the output for transcripts

involved in starch metabolism. All candidates were investigated using phylogenetic

reconstruction. Briefly, they were used as queries in a BLASTp search (e-value threshold 1e-5)

against a comprehensive custom database containing representatives of all major eukaryotic

groups and RefSeq data from all bacterial phyla at NCBI (https://www.ncbi.nlm.nih.gov/, last

accessed December 2017) (Altschul et al., 1990). The database was subjected to CD-HIT with a

similarity threshold of 85% to reduce redundant sequences and paralogs (Li and Godzik, 2006).

Results from blast searches were parsed for hits with a minimum query coverage of 50% and e-

values of less than 1e-5. The number of bacterial hits was restrained to 20 hits per phylum (for

FCB group, most classes of Proteobacteria, PVC group, Spirochaetes, Actinobacteria,

Cyanobacteria (unranked) and Firmicutes) or 10 per phylum (remaining bacterial phyla) as

defined by NCBI taxonomy. Parsed hits were aligned with MAFFT v. 7.212, using the–auto

option, poorly aligned regions were eliminated using trimAl v.1.2 with a gap threshold of 80%

(Katoh and Standley, 2013; Capella-Gutiérrez et al., 2009). Maximum likelihood tree

reconstructions were then performed with FastTree v. 2.1.7 using the default options (Price et al.,

2010). Phylogenies with overlapping taxa were consolidated by combining the parsed hits of the

corresponding queries, removing duplicates and repeating the alignment, trimming and tree

reconstruction steps as described above.

Acknowledgements

We thank Kristina I. Prokina for sample collection in Chile, Dr. Hoan Q. Tran and Tran
Duc Dien for assistance with sample collection and trip management in Vietnam, Jürgen F.H.
Strassert for help with DAPI staining, and Vladimir V. Aleshin and Kirill V. Mikhailov for
fruitful discussion on different aspects of origin of Metazoa. Field work in Vietnam is part of the
project "Ecolan 3.2" of the Russian-Vietnam Tropical Centre.

Funding: This work was supported by the Russian Science Foundation (grant no. 18-14-
00239).

Competing interests: None declared.

References

Abedin M, King N. 2008. The premetazoan ancestry of cadherins. *Science* **319**:946–948.
doi:10.1126/science.1151084

Adamska M. 2016. Sponges as models to study emergence of complex animals. *Curr Opin*
Genet Dev **39**:21–28. doi:10.1016/j.gde.2016.05.026

Aleshin VV, Konstantinova AV, Mikhailov KV, Nikitin MA, Petrov NB. 2007. Do we need
many genes for phylogenetic inference? *Biochemistry Mosc* **72**:1313–1323.
doi:10.1134/S000629790712005X

Altschul, S.F., Gish, W., Miller, W., Myers, E.W., and Lipman, D.J. 1990. Basic local alignment
search tool. *J. Mol. Biol.* **215**, 403–410.

Arkush KD, Mendoza L, Adkison MA, Hedrick RP. 2003. Observations on the life stages of
Sphaerothecum destruens n. g., n. sp., a mesomycetozoean fish pathogen formally referred
to as the rosette agent. *J Eukaryot Microbiol* **50**:430–438. doi:10.1111/j.1550-
7408.2003.tb00269.x

Brown MW, Spiegel FW, Silberman JD. 2009. Phylogeny of the "forgotten" cellular slime mold,
Fonticula alba, reveals a key evolutionary branch within Opisthokonta. *Mol Biol Evol*
26:2699–2709. doi:10.1093/molbev/msp185

Brown MW, Kolisko M, Silberman JD, Roger AJ. 2012. Aggregative multicellularity evolved
independently in the eukaryotic supergroup Rhizaria. *Curr Biol* **22**:1123–1127.
doi:10.1016/j.cub.2012.04.021

Brunet T, King N. 2017. The origin of animal multicellularity and cell differentiation. *Dev Cell*
43:124–140. doi:10.1016/j.devcel.2017.09.016

Capella-Gutiérrez, S., Silla-Martínez, J.M., and Gabaldón, T. 2009. trimAl: a tool for automated
alignment trimming in large-scale phylogenetic analyses. *Bioinformatics* **25**, 1972–1973.

Cavalier-Smith T. 2017. Origin of animal multicellularity: precursors, causes, consequences –
the choanoflagellate/sponge transition, neurogenesis and the Cambrian explosion. *Philos*
Trans R Soc Lond B Biol Sci **372**:20150476. doi:10.1098/rstb.2015.0476

Cavalier-Smith T, Chao EE. 2003. Phylogeny of Choanozoa, Apusozoa, and other Protozoa and
early eukaryote megaevolution. *J Mol Evol* **56**:540–563. doi:10.1007/s00239-002-2424-z

Carr M, Richter DJ, Fozouni P, Smith YJ, Jeuck A, Leadbeater BSC, Nitsche F. 2017. A six-
gene phylogeny provides new insights into choanoflagellate evolution. *Mol Phylogenet*
Evol **107**:166–178. doi:10.1016/j.ympev.2016.10.011

Dayel MJ, King N, 2014. Prey capture and phagocytosis in the choanoflagellate *Salpingoeca*
rosetta. *PLoS ONE* **9**:e95577. doi:10.1371/journal.pone.0095577

Dayel MJ, Agelado RA, Fairclought SR, Levin TC, Nichols SA, McDonald K, King N. 2011.
Cell differentiation and morphogenesis in the colony-forming choanoflagellate
Salpingoeca rosetta. *Dev Biol* **357**:73–82. doi:10.1016/j.ydbio.2011.06.003

del Campo J, Ruiz-Trillo I. 2013. Environmental survey meta-analysis reveals hidden diversity
among unicellular opisthokonts. *Mol Biol Evol* **30**:802–805. doi:10.1093/molbev/mst006

Dunn CW, Hejnol A, Matus DQ, Pang K, Browne WE, Smith SA, Seaver E, Rouse GW, Obst
M, Edgecombe GD, Sørensen MV, Haddock SH, Schmidt-Rhaesa A, Okusu A, Kristensen
RM, Wheeler WC, Martindale MQ, Giribet G. 2008. Broad phylogenomic sampling
improves resolution of the animal tree of life. *Nature* **452**:745–749.
doi:10.1038/nature06614

Feuda R, Dohrmann M, Pett W, Philippe H, Rota-Stabelli O, Lartillot N, Wörheide G, Pisani D.
2017. Improved modeling of compositional heterogeneity supports sponges as sister to all
other animals. *Curr Biol* **27:**3864–3870.e4. doi:10.1016/j.cub.2017.11.008

Gobert GN, Stenzel DJ, McManus DP, Jones MK. 2003. The ultrastructural architecture of the
adult *Schistosoma japonicum* tegument. *Int J Parasitol* **33**:1561–1575. doi:10.1016/S0020-
7519(03)00255-8

Haeckel E. 1874. Die Gastraea-Theorie, die phylogenetische Klassifikation des Thierreichs und
die Homologie der Keimblatter. *Jenaische Z Naturwiss* **8**:1–55. (in German).

Hashimshony T, Feder M, Levin M, Hall BK, Yanai I. 2015. Spatiotemporal transcriptomics
reveals the evolutionary history of the endoderm germ layer. *Nature* **519**:219–222.
doi:10.1038/nature13996

Hehenberger E, Tikhonenkov DV, Kolisko M, del Campo J, Esaulov AS, MylnikovAP, Keeling
PJ. 2017. Novel freshwater predators reshape holozoan phylogeny and reveal the presence
of a two-component signalling system in the ancestor of animals. *Curr Biol* **27**:2043–2050.
doi:10.1016/j.cub.2017.06.006

Hertel LA, Bayne CJ, Loker ES. 2002. The symbiont *Capsaspora owczarzaki*, nov. gen. nov. sp.,
isolated from three strains of the pulmonate snail *Biomphalaria glabrata* is related to
members of the Mesomycetozoea. *Int J Parasitol* **32**:1183–1191. doi:10.1016/s0020-
7519(02)00066-8.

Ivanov AV. 1968. Origin of multicellular animals. Phylogenetic essays. Leningrad: Nauka. 287
p. (in Russian).

James-Clark H. 1866. Note on the *Infusoria flagellata* and the *Spongiae ciliatae*. *Am. J. Sci*
1:113–114.

Kanehisa M, Goto S, Sato Y, Kawashima M, Furumichi M, Tanabe M. 2014. Data, information,
knowledge and principle: back to metabolism in KEGG. *Nucleic Acids Res*. **42**:D199–
D205.

Karpov SA, Efremova SM. 1994. Ultrathin structure of the flagellar apparatus in the choanocyte
of the sponge *Ephydatia fluviatilis*. *Tsitologia* **36**:403–408 (in Russian)

Katoh, K., and Standley, D.M. 2013. MAFFT multiple sequence alignment software version 7:
improvements in performance and usability. *Mol. Biol. Evol.* **30**, 772–780.

King N, Hittinger CT, Carroll SB. 2003. Evolution of key cell signaling and adhesion protein
families predates animal origins. *Science* **301:**361–363. doi:10.1126/science.1083853

King N, Westbrook MJ, Young SL, Kuo A, Abedin M, Chapman J, Fairclough S, Hellsten U,
Isogai Y, Letunic I, Marr M, Pincus D, Putnam N, Rokas A, Wright KJ, Zuzow R, Dirks
W, Good M, Goodstein D, Lemons D, Li W, Lyons JB, Morris A, Nichols S, Richter DJ,
Salamov A, Sequencing JG, Bork P, Lim WA, Manning G, Miller WT, McGinnis W,
Shapiro H, Tjian R, Grigoriev IV, Rokhsar D. 2008. The genome of the choanoflagellate
Monosiga brevicollis and the origin of metazoans. *Nature* **451**:783–788.
doi:10.1038/nature06617

Lang BF, O'Kelly C, Nerad T, Gray MW, Burger G. 2002. The closest unicellular relatives of
animals. *Curr Biol* **12**:1773–1778. doi:10.1016/S0960-9822(02)01187-9

Leadbeater BSC. 1983. Life-history and ultrastructure of a new marine species of
Proterospongia (Choanoflagellida). *J Mar Biol Assoc UK* **63**:135–160.
doi:10.1017/S0025315400049857

Leys SP, Cheung E, Boury-Esnault N. 2006. Embryogenesis in the glass sponge *Oopsacas
minuta*: formation of syncytia by fusion of blastomeres. *Integr Comp Biol* **46**:104–117.
doi:10.1093/icb/icj016

Li, W., and Godzik, A. 2006. Cd-hit: a fast program for clustering and comparing large sets of
protein or nucleotide sequences. *Bioinformatics* **22**, 1658–1659.

Lynn DH, Small EB. 1981. Protist kinetids: structural conservatism, kinetid structure, and
ancestral states. *Biosystems* **14**:377–385. doi:10.1016/0303-2647(81)90044-7

Mah JL, Christensen-Dalsgaard KK, Leys S.P. 2014. Choanoflagellate and choanocyte collar-
flagellar systems and the assumption of homology. *Evolution and Development* **16**:25–37.
doi:10.1111/ede.12060.

Maldonado M. 2004. Choanoflagellates, choanocytes, and animal multicellularity. *Invertebr Biol*
123:1–22. doi:10.1111/j.1744-7410.2004.tb00138.x

Maloof AC, Rose CV, Beach R, Samuels BM, Calmet CC, Erwin DH, Poirier GR, Yao N,
Simons FJ. 2010. Possible animal-body fossils in pre-Marinoan limestones from South
Australia. *Nat Geosci* **3**:653–659. doi:10.1038/ngeo934

Medina M, Collins AG, Taylor JW, Valentine JW, Lipps JH, Amaral-Zettler L, Sogin ML. 2003.
Phylogeny of Opisthokonta and the evolution of multicellularity and complexity in Fungi
and Metazoa. *J Astrobiol* **2**:203–211. doi:10.1017/S1473550403001551

Mikhailov KV, Konstantinova AV, Nikitin MA, Troshin PV, Rusin LY,Lyubetsky VA, Panchin
YV, Mylnikov AP, Moroz LL, Kumar S, Aleoshin VV. 2009. The origin of Metazoa: a
transition from temporal to spatial cell differentiation. *Bio Essays* **31**:758–768.
doi:10.1002/bies.200800214

Moestrup Ø. 2000. The flagellate cytoskeleton: Introduction of a general terminology for
microtubular roots in protists. In: Leadbeater BS, Green JC, editors. The Flagellates: Unity,
Diversity and Evolution. London: Taylor & Francis. pp 69–94.

Money NP. 2016. Fungal Diversity. In: The Fungi (Third Edition). Eds: Watkinson SC, Boddy
L, Money NP. Elsevier. pp. 1–36.

Moroz LL,Kocot KM, Citarella MR, Dosung S, Norekian TP, Povolotskaya IS, Grigorenko AP,
Dailey C., Berezikov E, Buckley KM, Ptitsyn A, Reshetov D, Mukherjee K, Moroz TP,
Bobkova E, Yu F, Kapitonov VV, Jurka J, Bobkov YV, Swore JJ, Girardo DO, Fodor A,
Gusev F, Sanford R, Bruders R, Kittler E, Mills CE, Rast JP,Derelle R, Solovyev VV,
Kondrashov FA, Swalla BJ, Sweedler JV, Rogaev EI, Halanych KM, Kohn AB. 2014. The
ctenophore genome and the evolutionary origins of neural systems. *Nature* **510**: 109–
114.doi: 10.1038/nature13400.

Mylnikov AP, Tikhonenkov DV, Karpov SA, Wylezich C. 2019. Microscopical studies on
picoplanctonic *Ministeria vibrans* Tong, 1997 (Filasterea) highlights the cytoskeletal
structure of the common ancestor of Filasterea, Metazoa and Choanoflagellata. *Protist*.
doi.org/10.1016/j.protis.2019.07.001

Nielsen C. 1987. Structure and function of metazoan ciliary bands and their phylogenetic
significance. *Acta Zoolog* **68**:205–262. doi:10.1111/j.1463-6395.1987.tb00892.x

Nielsen C. 2008. Six major steps in animal evolution: are we derived sponge larvae? *Evol Dev* **10**:241–257. doi:10.1111/j.1525-142X.2008.00231.x

Owczarzak A, Stibbs HH, Bayne CJ. 1980. The destruction of *Schistosoma mansoni* mother sporocysts in vitro by amoebae isolated from *Biomphalaria glabrata*: an ultrastructural study. *J Invertebr Pathol* **35**:26–33. doi:10.1016/0022-2011(80)90079-8

Philippe H, Derelle R, Lopez P, Pick K, Borchiellini C, Boury-Esnault N, Vacelet J, Renard E, Houliston E, Quéinnec E, Da Silva C, Wincker P, Le Guyader H, Leys S, Jackson DJ, Schreiber F, Erpenbeck D, Morgenstern B, Wörheide G, Manuel M. 2009. Phylogenomics revives traditional views on deep animal relationships. *Curr Biol* **19**:706–712. doi:10.1016/j.cub.2009.02.052

Pozdnyakov IR, Karpov SA. 2016. Kinetid Structure in Choanocytes of Sponges (Heteroscleromorpha): Toward the Ancestral Kinetid of Demospongiae. *J Morph* **277**:925–934. doi:10.1002/jmor.20546

Pozdnyakov IR, Sokolova AM, Ereskovsky AV, Karpov SA. 2017. Kinetid structure of choanoflagellates and choanocytes of sponges does not support their close relationship. *Protistology* **11**:248–264. doi:10.21685/1680-0826-2017-11-4-6

Pozdnyakov IR, Sokolova AM, Ereskovsky AV, Karpov SA. 2018. Kinetid structure in sponge choanocytes of Spongillida and Scopalinida in the light of evolutionary relationships within Demospongiae. *Zool J Linnean Soc* **184**:255–272 doi:10.1093/zoolinnean/zlx109

Price, M.N., Dehal, P.S., and Arkin, A.P. 2010. FastTree 2–approximately maximum-likelihood trees for large alignments. *PLoS ONE* **5**, e9490.

Raghukumar S. 1987. Occurrence of the Thraustochytrid, *Corallochytrium limacisporum* gen. et sp. nov. in the coral reef lagoons of the Lakshadweep Islands in the Arabian Sea. *Bot Mar* **30**:83–89. doi:10.1515/botm.1987.30.1.83

Ratcliff WC, Denison RF, Borrello M, Travisano M. 2012. Experimental evolution of multicellularity. *Proc Nat Acad Sci USA* **109:**1595–1600. doi:10.1073/pnas.1115323109

Ruiz-Trillo I, Burger G, Holland PW, King N, Lang BF, Roger AJ, Gray MW. 2007. The origins of multicellularity: a multi-taxon genome initiative. *Trends Genet* **23**:113–118. doi:10.1016/j.tig.2007.01.005

Ruiz-Trillo I, Roger AJ, Burger G, Gray MW, Lang BF. 2008. A phylogenomic investigation into the origin of metazoa. *Mol Biol Evol* **25:**664–672. doi:10.1093/molbev/msn006

Ryan JF, Pang K, Schnitzler CE, Nguyen AD, Moreland RT, Simmons DK, Koch BJ, Francis WR, Havlak P, NISC Comparative Sequencing Program, Smith SA, Putnam NH, Haddock SH, Dunn CW, Wolfsberg TG, Mullikin JC, Martindale MQ, Baxevanis AD. 2013. The

genome of the ctenophore *Mnemiopsis leidyi* and its implications for cell type evolution.
Science **342:**1242592. doi:10.1126/science.1242592

Sachwatkin AA. 1956. Vergleichende embryologie der niederen wirbellosen: ursprung und
gestaltungswege der individuellen entwicklung der vielzeller. Berlin: VEB Deutscher
Verlag der Wissenschaften. S. 401.

Schierwater B, Eitel M, Jakob W, Osigus H-J, Hadrys H, Dellaporta SL, Kolokotronis S-O, De
Salle R. 2009. Concatenated analysis sheds light on early metazoan evolution and fuels a
modern "Urmetazoon" hypothesis. *PLoS Biol* **7:**e1000020.
doi:10.1371/journal.pbio.1000020

Sebé-Pedrós A, Ariza-Cosano A, Weirauch MT, Leininger S, Yang A, Torruella G, Adamski M,
Adamska M, Hughes TR, Gómez-Skarmeta JL, Ruiz-Trillo I. 2013a. Early evolution of the
T-box transcription factor family. *Proc Natl Acad Sci USA* **110:**16050–16055.
doi:10.1073/pnas.1309748110

Sebé-Pedrós A, Irimia M, Del Campo J, Parra-Acero H, Russ C, Nusbaum C, Blencowe BJ,
Ruiz-Trillo I. 2013b. Regulated aggregative multicellularity in a close unicellular relative
of metazoan. *eLife* **2:**e01287. doi:10.7554/eLife.01287

Sebé-Pedrós A, Peña MI, Capella-Gutiérrez S, Antó M, Gabaldón T, Ruiz-Trillo I, Sabido E.
2016. High-throughput proteomics reveals the unicellular roots of animal
phosphosignaling and cell differentiation. *Dev Cell* **39:**186–197.
doi:10.1016/j.devcel.2016.09.019

Sebé-Pedrós A, Degnan BM, Ruiz-Trillo I. 2017. The origin of Metazoa: a unicellular
perspective. *Nat Rev Genet* **18:**498–512. doi:10.1038/nrg.2017.21

Shalchian-Tabrizi K, Minge MA, Espelund M, Orr R, Ruden T, Jakobsen KS, Cavalier-Smith T.
2008. Multigene phylogeny of choanozoa and the origin of animals. *PLoS ONE* **3:**e2098.
doi:10.1371/journal.pone.0002098

Sharpe SC, Eme L, Brown MW, Roger AJ. 2015. Timing the origins of multicellular eukaryotes
through phylogenomics and relaxed molecular clock analyses. Evolutionary transitions to
multicellular life. Advances in Marine Genomics. Ruiz-Trillo I, Nedelcu AM (Eds.).
Netherlands: Springer. P. 3–29.

Signorovitch AY, Buss LW, Dellaporta SL. 2007. Comparative genomics of large mitochondria
in Placozoans. *PLoS Genet* **3:**e13. doi:10.1371/journal.pgen.0030013

Simion P, Philippe H, Baurain D, Jager M, Richter DJ, Di Franco A, Roure B, Satoh N,
Quéinnec É, Ereskovsky A, Lapébie P, Corre E, Delsuc F, King N, Wörheide G, Manuel
M. 2017. A large and consistent phylogenomic dataset supports sponges as the sister group
to all other animals. *Curr Biol* **27:**958–967. doi:10.1016/j.cub.2017.02.031

Smirnov AV. 2000. Subclass Aconchulina. In: Alimov AF, editor. Protista. Pt. 1. Handbook on zoology. St-Petersburg. Nauka. pp. 485–490.

Srivastava M, Begovic E, Chapman J, Putnam NH, Hellsten U, Kawashima T, Kuo A, Mitros T, Salamov A, Carpenter ML, Signorovitch AY, Moreno MA, Kamm K, Grimwood J, Schmutz J, Shapiro H, Grigoriev IV, Buss LW, Schierwater B, Dellaporta SL, Rokhsar DS. 2008.The *Trichoplax* genome and the nature of placozoans. *Nature* **454**: 955–960.doi: 10.1038/nature07191

Srivastava M, Simakov O, Chapman J, Fahey B, Gauthier ME, Mitros T, Richards GS, Conaco C, Dacre M, Hellsten U, Larroux C, Putnam NH, Stanke M, Adamska M, Darling A, Degnan SM, Oakley TH, Plachetzki DC, Zhai Y, Adamski M, Calcino A, Cummins SF, Goodstein DM, Harris C, Jackson DJ, Leys SP, Shu S, Woodcroft BJ, Vervoort M, Kosik KS, Manning G, Degnan BM, Rokhsar DS. 2010.The *Amphimedon queenslandica* genome and the evolution of animal complexity. *Nature* **466**: 720–726. doi: 10.1038/nature09201

Suga H, Ruiz-Trillo I. 2013. Development of ichthyosporeans sheds light on the origin of metazoan multicellularity. *Dev Biol* **377:**284–292. doi:10.1016/j.ydbio.2013.01.009

Suga H, Chen Z, de Mendoza A, Sebé-Pedrós A, Brown MW, Kramer E, Carr M, Kerner P, Vervoort M, Sánchez-Pons N, Torruella G, Derelle R, Manning G, Lang BF, Russ C, Haas BJ, Roger AJ, Nusbaum C, Ruiz-Trillo I. 2013. The *Capsaspora* genome reveals a complex unicellular prehistory of animals. *Nat Commun* **4:**2325. doi:10.1038/ncomms3325

Tikhonenkov DV, Mazei YA, Embulaeva EA. 2008. Degradational Succession of a Community of Heterotrophic Flagellates in Microcosm Experiments. *Zh Obshch Biol* 69:57–64

Tikhonenkov DV, Janouškovec J, Mylnikov AP, Mikhailov KV, Simdyanov TG, Aleoshin VV, Keeling PJ. 2014. Description of *Colponema vietnamica* sp.n. and *Acavomonas peruviana* n. gen. n. sp., two new alveolate phyla (Colponemidia nom. nov. and Acavomonidia nom. nov.) and their contributions to reconstructing the ancestral state of alveolates and eukaryotes. *PLoS ONE* **9**:e95467. doi:10.1371/journal.pone.0095467

Torruella G, Derelle R, Paps J, Lang BF, Roger AJ, Shalchian-Tabrizi K,Ruiz-Trillo I. 2012. Phylogenetic relationships within the Opisthokonta based on phylogenomic analyses of conserved single-copy protein domains. *Mol Biol Evol* **29:**531–544. doi:10.1093/molbev/msr185

Torruella G, de Mendoza A, Grau-Bové X, Antó M, Chaplin MA, del Campo J, Eme L, Pérez-Cordón G, Whipps CM, Nichols KM, Paley R, Roger AJ, Sitjà-Bobadilla A, Donachie S, Ruiz-Trillo I. 2015. Phylogenomics reveals convergent evolution of lifestyles in close relatives of animals and fungi. *Curr Biol* **25:**2404–2410. doi:10.1016/j.cub.2015.07.053

Whelan NV, Kocot KM, Moroz LL, Halanych KM. 2015. Error, signal, and the placement of
Ctenophora sister to all other animals. *Proc Natl Acad Sci USA* **112:**5773–5778.
doi:10.1073/pnas.1503453112

Wickstead B, Gull K. 2012. Evolutionary biology of dyneins. In: King S, editor. Dyneins.
Elsevier. pp. 89–121.

Williams F, Tew HA, Paul CE, Adams JC. 2014. The predicted secretomes of *Monosiga*
brevicollis and *Capsaspora owczarzaki*, close unicellular relatives of metazoans, reveal
new insights into the evolution of the metazoan extracellular matrix. *Matrix Biol* **37:**60–68.
doi:10.1016/j.matbio.2014.02.002

Yubuki N, Leander BS. 2013. Evolution of microtubule organizing centers across the tree of
eukaryotes. *Plant J* **75**:230–244. doi:10.1111/tpj.12145

Zakhvatkin AA. 1949. The comparative embryology of the low invertebrates. Sources and
method of the origin of metazoan development. Moscow, Soviet Science. p. 395. (in
Russian).

Legends for videos

Video 1. Swimming of *Syssomonas multiformis* cell with rotation.

Video 2. Attached cell of *Syssomonas multiformis* and rapid flagellum beating.

Video 3. Amoeboflagellate stage of *Syssomonas multiformis.* Cells of eukaryotic prey
Parabodo caudatus are also visible.

Video 4. Loss of flagellum in *Syssomonas multiformis* and transition to amoeba.

Video 5. Transformation of amoeba into a cyst in *Syssomonas multiformis*.

Video 6. Palintomic divisions inside the cyst of *Syssomonas multiformis*.

Video 7. Division into two cell structures in *Syssomonas multiformis.*

Video 8. Cell and cyst of *Syssomonas multiformis* with vesicular structures inside.

Video 9. Feeding of *Syssomonas multiformis* on eukaryotic prey.

Video 10. Feeding of *Syssomonas multiformis* on bacteria.

Video 11. Temporary cell aggregations of *Syssomonas multiformis.*

Video 12. Floating rosette-like aggregation of *Syssomonas multiformis.*

Video 13. Syncytium-like structures and budding of young flagellated daughter cells in
Syssomonas multiformis.

Video 14. Joint feeding of *Pigoraptor vietnamica* on died cell of *Parabodo caudatus.*

Video 15. Joint feeding of *Pigoraptor chileana* on died cell of *Parabodo caudatus.*

Video 16. Temporary cell aggregation of *Pigoraptor chileana.*

Videos are available at

https://drive.google.com/drive/folders/1ewNvwSQCAVG81eYjJPdhR41aBpUKR05L