Sexually antagonistic coevolution between the sex chromosomes of *Drosophila melanogaster*

Katrine K. Lund-Hansen[1,2*+], Colin Olito[1+], Edward H. Morrow[2], & Jessica K. Abbott[1]

[1] Department of Biology, Section for Evolutionary Ecology, Lund University, Lund 223 62, Sweden.
[2] School of Life Sciences, University of Sussex, Brighton BN1 9QG, United Kingdom
[*] Corresponding author e-mail: Katrine.Lund-Hansen@biol.lu.se
[+] Equal contributions

Manuscript elements: Table 1, Figure 1, Figure 2, Figure 3, Figure 4; Online Supplementary Material: Appendix A – Empirical methods: Figure S1, Table S1, Table S2, Table S3, Table S4; Appendix B – Development of the recursions.

Running Head: Coevolution between the sex chromosomes

Keywords: Coevolution; Compensatory evolution; Interlocus sexual conflict; Sex chromosome evolution; Sperm Competition; Egg-to-adult survival.

Abstract

Antagonistic interactions between the sexes are important drivers of evolutionary divergence. Interlocus sexual conflict is generally theorised as a conflict between alleles at two interacting loci whose identity and genomic location are arbitrary. Here we build on previous theory and suggest that when these two loci are located on the X and Y (or Z and W) chromosomes, it can lead to cycles of antagonistic coevolution between the sex chromosomes and the two sexes. To test this hypothesis, we performed experimental crosses using *Drosophila melanogaster* where we reciprocally exchanged the sex chromosomes between five wild-type populations in a round-robin design. Disrupting putatively coevolved sex chromosome pairs resulted in increased male reproductive success in 16 out of 20 experimental populations (10 of which were significant), but also resulted in lower offspring egg-to-adult viability that affected both male and female fitness. However, after 25 generations of experimental evolution these sexually antagonistic fitness effects appeared to have been resolved. To help formalise our hypothesis, we developed population genetic models of antagonistic coevolution using fitness expressions based on our empirical results. Our models support the conclusion that antagonistic coevolution between the sex chromosomes is plausible under the fitness effects observed in our experiments. Together, our results lend both empirical and theoretical support to the idea that a cycle of antagonistic coevolution can occur between sex chromosomes, and illustrates how this process may drive genetic and phenotypic divergence between populations.

Introduction

Sex chromosomes have a number of unique features distinguishing them from the rest of the nuclear genome and giving them particular importance in evolutionary biology. The sex chromosomes differ both from the autosomes, and from one another, in their mode of inheritance, the selection experienced in the two sexes, and gene content (1). Moreover, these processes are interdependent and strongly influence the genetic variation harboured on each chromosome. The Y (or W) chromosome is inherited only through the heterogametic sex (XY/ZW), whereas the X (or Z) chromosome is inherited from both the hetero- and homogametic (XX/ZZ) sex (1). The Y chromosome will therefore only experience selection in males, but the X chromosome, which spends two-thirds of its time in females, will regularly experience selection in females (1). However, the X chromosome is hemizygous in males and will therefore also always be exposed to selection when present in males (i.e., X-linked genes are unsheltered by dominance effects in males) (2). Though the X chromosome is shared by the sexes, the X chromosome in *Drosophila melanogaster* has been shown to harbour more female-biased than male-biased genetic variation (3, 4), whereas the Y chromosome has been shown to impact male fitness traits (5–7).

These unique properties of the sex chromosomes give them an important role in two major fields within evolutionary biology: speciation (8) and sexual conflict (1, 9). Here we aimed to investigate whether interactions between sex chromosomes contribute to between-population divergence at the intra-specific level. To do so we performed experimental crosses to 'swap' either an X or a Y chromosome between five populations of *D. melanogaster*, allowing us to isolate the effects of novel interactions between the sex chromosomes on male reproductive fitness. As we outline below, we expected to see different patterns of male reproductive success depending on whether population divergence is caused by the accumulation of sex-linked sterility mutations (or other incompatibilities contributing to reproductive isolation between populations), or is instead a result of sexually antagonistic coevolution between the sex chromosomes.

If interactions between the sex chromosomes contribute to divergence between these populations primarily through accumulation of X-linked incompatibilities, we would expect to see a decrease in male fitness in the experimental populations as

these incompatibilities are exposed to selection in males. The X chromosome has a

large effect on hybrid male sterility (10), and the accumulation of incipient 'hybrid

incompatibilities' begin in allopatric populations within a species prior to speciation

(11). However, while previous studies have found evidence of early prezygotic

isolation between sub-Saharan and cosmopolitan populations of *D. melanogaster* (12,

13), there is limited evidence for postzygotic isolation (14). On the other hand, sexual

conflict between the sex chromosomes could, in principle, provide an alternative

mechanism for them to contribute to evolutionary divergence between populations

(15), as Y-linked regulation of gene expression has been indicated to do (16).

Sexually antagonistic coevolution is based on the Red-Queen process among

species playing out between the sexes (17). Simply put, sexually antagonistic

coevolution describes an antagonistic interaction between the sexes in which males

increase their reproductive success through an adaptation that is simultaneously

detrimental to females, which creates selection for counter-adaptation in females to

regain their lost fitness; the process may be repeated in multiple cycles over

evolutionary time until a resolution is reached or a palliative adaptation ends the

conflict (18). Sexually antagonistic coevolution is a form of interlocus sexual conflict

(15) if the traits involved are encoded by different genes in males and females (19).

As first proposed by Rice & Holland in 1997, when the loci in question are located on

the sex chromosomes, this could lead to cycles of sexual antagonistic coevolution

between the sex chromosomes (15). If a Y-linked male-beneficial mutation arises that

increases male reproductive success but decreases female fitness, it should spread

through a population because it is subject to selection in males only. We might then

expect selection for any compensatory mutation that might arise on the X

chromosome or autosomes to increase female fitness again. Such mutations will likely

differ between allopatric populations, and we would therefore expect to find an

increase in male reproductive fitness when a Y chromosome with male beneficial

mutations is paired with an X chromosome without the corresponding compensatory

mutation(s). According to the sexually antagonistic coevolutionary model, we also

predict a decrease in female fitness when mating with males harbouring a Y

chromosome in a novel X chromosome background. Another corollary of the

antagonistic coevolutionary model is that the effect of disrupting coevolved sex

chromosomes on male and female fitness should decay over subsequent generations

as new compensatory mutations arise, or novel combinations of segregating alleles

achieve a similar compensatory effect. We tested this prediction by examining male

and female fitness in our experimental populations again after 25 generation of

experimental evolution. Crucially, the signature of antagonistic coevolution between

the sex chromosomes in terms of male and female fitness differs from that for the

accumulation of sex-linked incompatibilities, allowing us to identify whether one or

other process is driving evolutionary divergence between populations.

To complement our empirical results and help formalize our hypothesis of

sexually antagonistic coevolution between the sex chromosomes, we developed

population genetic models describing the evolution of two interacting loci located in

different genomic regions (i.e., unlinked): a Y-linked locus influencing both adult

male fertilization success (i.e., sperm competition) and subsequent offspring survival,

and a compensatory locus affecting only offspring survival located on either an

autosome or the X chromosome.

Results and Discussion

Empirical Results

To empirically test for evidence of population-specific interactions between the sex

chromosomes in *D. melanogaster*, we crossed five outbred laboratory-adapted wild-

type populations derived from four continents and three climatic zones in a round

robin cross design (Fig. 1). We created 20 novel populations where either the X

chromosome (*novel X* treatment) or the Y chromosome (*novel Y* treatment) from one

wild-type population (*wt*) was incorporated into another population (*SI Appendix A*,

Fig. S1). We found a significant effect of treatment on male relative reproductive

fitness (*SI Appendix A*, Table S1), with males from the *novel X* or *novel Y* treatments

having a significantly higher relative fitness than *wt* males (Fig. 2*A*, *SI Appendix A*,

Table S2). To exclude interactions between the novel sex chromosome and the

autosomes, we also created a *novel XY* treatment where a pair of sex chromosomes

from one *wt* population were introduced into another population (*SI Appendix A*, Fig.

S2). We found no significant effect of treatment on relative fitness in the assay with

novel XY treatments (Fig. 2*B*, *SI Appendix A*, Table S1). It therefore seems that

introducing an individual novel sex chromosome caused the fitness differences rather

than the autosomal background. Mitochondria-nuclear interactions have previously

been shown to affect male fertility in *D. melanogaster* (20), and both *novel X* and

novel Y chromosomes were expressed with novel mitochondria. However, the *novel*

XY chromosomes also were expressed with novel mitochondria and we found no

effect on male fitness in these populations, suggesting that the magnitude of any mito-

nuclear interactions was minor in these populations compared to the effect of

interactions between the sex chromosomes. To investigate the interaction between the

sex chromosomes in more detail, we calculated the difference in fitness between the

novel populations and their *wt* counterparts (ΔFitness = $\omega_{\text{novel population}}$ - $\omega_{\text{wild-type}}$). We

used bootstrap methods to model the data, and found that 10 of the 20 novel

populations were significantly different from zero in a positive direction (Fig. 2*C*, *SI*

Appendix A, Table S3). 16 out of the 20 point estimates were also positive, which is

significantly different from what would be expected under random deviations (*P* =

0.01). Taken together, our results demonstrate that exchanging a sex chromosome

between populations of *D. melanogaster* had a positive overall effect on male fitness.

This was consistent with the expected signature of antagonistic coevolution between

the sex chromosomes on male fitness, but not with that of accumulated sex-linked

incompatibilities influencing male sterility.

To tease apart which fitness components were driving the overall pattern of

increased relative fitness in the *novel* populations, we looked at a number of

phenotypic traits that are correlated to male fitness. Interestingly, we found that the

increase in fitness was related to different traits in the two *novel* sex chromosome

treatments. For *novel X* males, we found that the increase in fitness was correlated

with an increase in size, as *novel X* males were significantly larger than both *wt* and

novel Y males (*SI Appendix A,* Fig. S3*A*, Table S1 & S2). Body size is an important

factor in male fitness (21), and larger males have been shown to have a mating

advantage (22). For *novel Y* males we found that the increase in fitness was correlated

with an increased ability to displace other males' sperm (*SI Appendix A*, Fig. S3*B*,

Table S1), as *novel Y* males were significantly better at displacing sperm compared to

wt males (*SI Appendix A,* Table S2). It therefore seemed that the *novel X* males gained

a mating advantage from an increase in size, whereas the *novel Y* males gained the

advantage through being better at sperm displacement. Thus, the results suggest that

different phenotypic and genetic mechanisms may be responsible for the increased

male fitness observed in the two *novel* sex chromosome treatments.

If the coevolution between the sex chromosomes is antagonistic in nature, the
increase in male fitness should come at a cost to female fitness. We found a
significant effect of treatment on total offspring number (*SI Appendix A*, Fig. S4*A*,
Table S1), with *novel X* males siring a lower number of live offspring compared to *wt*
males (Table S2). However, we did not find any difference in number of eggs laid by
females mated with males from the different treatments (*SI Appendix A*, Fig. S4*B*,
Table S1). The discrepancy between number of eggs and live offspring suggest a
trade-off between number of offspring and offspring quality, which was confirmed by
a significant difference in egg-to-adult viability (SI Appendix A, Fig. S4*C*, Table S1).
Novel X males sired significantly lower numbers of viable offspring than the other
two treatments (*SI appendix A*, Table S2). So mating with *novel X* males decrease a
female's fitness by reducing the overall number of live offspring she produces. A
decrease in offspring survival could be caused by meiotic drives seen in a departure
from 50:50 sex ratio (23). We did not find a significant difference in sex ratio between
the different treatments (*SI Appendix A*, Fig. S4*D*, Table S1). There are also no known
sex ratio meiotic drivers in *D. melanogaster*, so the decrease in offspring survival was
most likely caused by the father's genotype. We were unable to establish which
harmful effects *novel Y* males had on female fitness through the assays performed in
this experiment. But we were able to exclude increased male harassment (e.g. due to
larger size), males inducing high fecundity, and reducing offspring viability. A
possible cause to investigate in the future is the reduction of female receptivity after
mating with males, which could lead to a reduction in female lifetime reproductive
fitness.

If the sex chromosomes coevolve antagonistically any shift towards increased
fitness in one sex at the cost of the other sex could lead to counter-adaption, which we
expected to see as a reduction in male fitness over time. In principle, this subsequent
reduction in male fitness could be due to new X-linked compensatory mutations;
however, given the time-scale of the experiment it is more likely that any
compensatory evolution by females must utilize standing genetic variation in the
stock populations, with novel allelic combinations achieving a compensatory effect.
After 25 generations of experimental evolution (*SI Appendix A,* Fig. S5), we repeated
both the male reproductive fitness assay and two of the significant phenotype assays
to test if the interactions between the novel pairs of sex chromosomes had changed.
As predicted, we no longer found a significant treatment effect on male fitness (Fig.

3*A, SI Appendix A,* Table S4). Indeed, we found that ΔFitness (ΔFitness = ω_{novel}

$_{\text{population}}$ - $\omega_{\text{wild-type}}$) between novel populations and their wild-type counterparts had

diminished in magnitude and were no longer significantly different from 0 (except for

one population, I–O_X) (Fig. 3*B, SI Appendix A,* Table S5). The reduction in fitness

observed after just 25 generations indicates a strong selection pressure for

compensatory evolution, which is supported by the simulation results. We also no

longer found an effect of the treatments on sperm competition (*SI Appendix A,* Fig.

S6*A,* Table S4) or offspring egg-adult survival (*SI Appendix A,* Fig. S6*B,* Table S4).

There was a significant effect of the treatments on sex ratio (*SI Appendix A,* Fig. S6*C,*

Table S4) as *novel Y* males produced significantly more male offspring than *wt* males

(*SI Appendix A,* Table S6). However, this significant difference was not due to a

change in sex ratio in the *novel Y* populations; rather it was due to a change in the sex

ratio in the *wt* populations.

Taken together, our empirical results are consistent with predictions for

antagonistic coevolution between the sex chromosomes. To further validate that we

found evidence of antagonistic coevolution, we built a theoretical model based on our

empirical evidence.

Theoretical Models

Consider two alternative genetic systems involving two unlinked loci: a Y-linked

locus (**Y**, with alleles *Y,* and *y*), and a compensatory locus located on either an

autosome (**A**, with alleles *A,* and *a*) or the X chromosome (**X**, with alleles *X,* and *x*)

(the Autosomal and X-linked models respectively), in a large population with discrete

generations. In both models, a mutant *y* chromosome increases male fertilization

success by a rate of $1 + s_m$ relative to the wild type (*Y*), but also reduces viability of

offspring resulting from matings with females carrying the wild-type allele at the

compensatory locus, *A* (or *X*). At the same time, females carrying the mutant *a* (or *x*)

compensatory allele may incur a 'cost of compensation' in terms of offspring viability

when mating with wild-type (*Y*) males (see Table **1** and Materials and Methods for a

summary and description of fitness expressions). These fitness expressions create a

scenario of antagonistic coevolution between a potentially male-beneficial Y-linked

mutation, and the compensatory locus. Reminiscent of standard theories of

compensatory evolution (e.g., (24, 25)), each of the mutant alleles (*y* and either *a* or *x*)

reduces offspring survival in isolation. However, unlike previous theory, the two loci

are located on different chromosomes that do not recombine with one another; if a

mutant y chromosome spreads to fixation, compensation at the population level

requires that the compensatory mutation also fixes (see also recent models of

"Father's Curse" for similar scenarios, but without compensatory evolution or

parental effects on offspring viability (26); a full description of the models is

presented in the Materials and Methods and in *SI Appendix B*). Our theoretical

analyses focus on (i) the evolutionary invasion of rare mutants at each locus

individually; and (ii) single bouts of coevolution, beginning with invasion of a single-

copy mutant y chromosome in a population initially fixed for the wild-type Y

chromosome and for the wild-type A (or X) allele at the compensatory locus, and

completing when both mutant alleles y and a (or x) become fixed (see (31) for a

similar approach in the context of mito-nuclear coevolution).

Evolutionary invasion analyses of the two models reveal three key results.

Intuitively, invasion of a rare mutant y chromosome into a population initially fixed

for the wild-type alleles at both loci requires that the increase in fertilization success

for mutant males is greater than the accompanying reduction in offspring survival.

Neglecting second order terms, a mutant y chromosome can spread when $\delta = s_m - s_o >$

0. If compensatory evolution is slow relative to the evolution of the Y-linked locus

(e.g., the compensatory mutation rate is much smaller than the mutation rate to a

mutant y), the mutant y chromosome is most likely to fix before a new compensatory

mutation occurs. In this case, the genomic location of the compensatory locus does

not influence the invasion conditions for a compensatory mutation. All males carry

the mutant y, and so compensatory mutations will spread if there is any selection

against the wild-type allele at the compensatory locus (i.e., $0 < h_o, s_o < 1$ for both

models). Differences between the models emerge when new compensatory mutations

can arise while the mutant y chromosome is still segregating in the population. If there

is any cost of compensation for females (i.e., $0 < s_c$), compensatory mutations will

experience purifying selection while q_y is small because most males carry the wild-

type Y chromosome. As q_y increases, more matings involve mutant y males and

females homozygous for the wild-type allele at the compensatory locus. When q_y

reaches a threshold frequency, $\tilde{q}_y^A$ ($\tilde{q}_y^X$ for the X-linked model), compensatory

mutations will be favoured by selection (derivations provided in *SI Appendix B*). As

illustrated in Fig. 4*A* for the case of additive compensatory fitness effects ($h = h_o = h_c$

$= 1/2$), $\tilde{q}_y^X$ is always less than or equal to $\tilde{q}_y^A$, and the difference between the two

thresholds is greatest when $0 < s_o$, $s_c \ll s_m$. Overall, these results suggest that the

genomic location of compensatory mutations (Autosomal or X-linked) will be most

important when (i) the mutant *y* chromosome is strongly beneficial for males and

there is little or no cost of compensation for females (i.e., when $0 < s_o$, $s_c \ll s_m$); and

(ii) when compensatory evolution is not limited by mutational variation (i.e., when

mutant *y* chromosomes and compensatory mutations are likely to co-segregate in the

population). To complement our analytic results, we performed Wright-Fisher

simulations to estimate two other important properties of coevolutionary cycles

predicted by our models: (i) the probability of invasion of single-copy autosomal (Π_A)

and X-linked (Π_X) compensatory mutations into populations of *N* individuals that are

initially fixed for the mutant *y* chromosome; and (ii) the total time to complete a

single bout of coevolution between the Y-linked and compensatory loci (T_A and T_X,

respectively) under recurrent mutation, selection, and drift. Although the invasion

conditions for mutant compensatory mutations are the same for both models when the

mutant *y* chromosome is initially fixed, the probability of eventual fixation for a

single-copy mutation is larger for an X-linked compensatory locus (Π_X) than an

autosomal one (Π_A) except for the special case of complete dominance (i.e., when h_o

$= 1$) (Fig. 4*B*). The average time to complete a co-evolutionary cycle is also smaller

for an X-linked than an autosomal compensatory locus, provided the compensatory

mutation rate (μ_a or μ_x) is not dramatically lower than the male-beneficial mutation

rate (μ_y) (Fig. 4*C*). When compensatory evolution is strongly limited by mutational

variation, compensatory evolution can be faster on the autosomes than the X. This is

due to an implicit trade-off between the probability of invasion, which is higher for

the X than the autosomes (Fig. 4*B*), and the size of the mutational target, which is

lower for the X because males only carry one copy.

Overall, our theoretical results support the conclusion that antagonistic

coevolution is (i) plausible under the fitness effects observed in our empirical

experiments; and (ii) more likely to involve both sex chromosomes than the Y and an

autosome, especially when compensatory dynamics are not slow relative to those of

male-beneficial Y-linked mutations. An interesting corollary to these results is that

antagonistic coevolution should drive more rapid among-population divergence on the
X chromosome than the autosomes.
308
Conclusion
In conclusion, we empirically examined the role of the sex chromosomes in
evolutionary divergence among allopatric populations of *D. melanogaster*. We found
evidence of intragenomic conflict between the sex chromosomes that was independent
of any interaction with the autosomes or mitochondria. Disrupting coevolved sex
chromosomes resulted in an overall increase in reproductive fitness for *novel* males,
through increased body size (for *novel X* males) and sperm displacement (for *novel Y*
males), but also decreased offspring egg-to-adult viability. The accompanying
reduction in offspring viability created indirect selection on females to mitigate the
loss in fitness resulting from mating with a *novel* male. After 25 generations of
experimental evolution *novel* males no longer enjoyed higher reproductive fitness and
there were no differences in offspring survival, indicating that the sexually
antagonistic effects of disrupting coevolved sex chromosomes had been resolved,
probably by standing genetic variation for compensatory alleles in the experimental
populations. Overall, our empirical results were consistent with a sexually
antagonistic coevolutionary model of sex chromosome evolution in allopatric
populations. Analysis of our theoretical models supports the plausibility of
antagonistic coevolution between the sex chromosomes under the fitness effects
observed in the experiments, and predicts that such coadaptation is more likely to
involve sex chromosomes than autosomes provided that compensatory evolution is
not limited by mutational variation. These new insights into the interactions between
the sex chromosomes can help further our understanding of early speciation events.
Antagonistic coevolutionary cycles between the sex chromosomes will most likely
follow different trajectories in different populations due to random mutations and the
interaction between the environment and sexual conflict (27), resulting in genetic and
phenotypic divergence between allopatric populations. Thus, over evolutionary time,
antagonistic coevolutionary cycles could lead to hybrid incompatibility, and thereby
contribute to speciation.
337

Materials and Methods

Empirical methods

***Drosophila* husbandry and fly stocks.** We used five outbred laboratory-adapted wild-type populations: (1) *Dahomey*, Africa, tropical (22); (2) *Innisfail*, Oceania, tropical; (3) LH_M, North America, mediterranean; (4) *Odder*, Europa, temperate (29); (5) *Tasmania*, Oceania, temperate. For at least two generations before the cross, we maintained all wild-type populations under the standard LH_M culturing protocol (25°C, 12-12 light-dark cycle, 60% relative humidity, cornmeal-molasses-yeast medium, *SI Appendix A* (28)). For the fitness assays we used an outbred LH_M population homozygous for the visible brown eye (*bw*) genetic marker (LH_M-*bw*).

Male reproductive fitness. We estimated male reproductive fitness as the proportion of live offspring sired by males of the target treatment. Five adult target males were placed in a vial for two days with ten competitor LH_M-*bw* males and 15 virgin LH_M-*bw* females. The females were transferred into a single test tube to oviposit for 18 hours after which the females were discarded, and the test tubes left under standard LH_M conditions for 12 days. After which we counted the adult offspring and scored them for eye-colour to assess paternity. The *bw* genetic marker is recessive to the wild-type red eye-colour allele, so all red-eyed offspring can be assigned to red-eyed target males. We calculated relative fitness of the target males by dividing the fitness for each replicate by the maximum fitness across all replicates. We measured male fitness for three different assays: at generation 0 (n = 2 blocks x 7 experimental replicates x 25 populations; fitness estimates for 70 individuals per population), at generation 25 (n = 2 blocks x 3 experimental replicates x 2 replicate populations x 14 populations; fitness estimates for 60 individuals per population) and for the sex chromosome-autosome interactions (n = 10 experimental replicates x 10 populations; fitness estimates for 50 individuals per population).

Construction of the *novel* populations, the evolution experiment, male thorax size, sperm competition assay, male effect on female fecundity, and offspring egg-to-adult viability assay are described in *SI Appendix A*.

Statistical procedures. All the statistical analyses were conducted in R version 3.4.4

(30). We fit linear models with treatment as a fixed factor to test each of the

dependent variables: thorax size and male effect on female fecundity, and tested for

significant effects using ANOVA. We performed post-hoc Tukey HSD comparisons

for all analyses with a significant treatment effect. We used linear mixed models

(**lme4** (31)) with treatment as a fixed factor and experimental block as a random

factor to test the dependent variables: male reproductive fitness, egg-to-adult

offspring viability, sex ratio, sperm competition, and total offspring number, and

again tested for significant effects using ANOVA. We performed post-hoc Tukey

HSD (**multcomp** (32)) comparisons for all analyses with a significant treatment

effect. We calculated bootstrap 95% confidence intervals around ΔFitness (ΔFitness =

$\omega_{novel\ population}$ - $\omega_{wild-type}$) randomly resampling 13 out of 14 data points and

recalculating ΔFitness 10,000 times. We used the Exact Binomial Test to test if there

were more positive ΔFitness estimates than expected by chance. For statistical

analysis after the 25 generations of experimental evolution we fit linear mixed models

with treatment nested within replicate populations to test the dependent variables:

male reproductive fitness, egg-to-adult offspring survival, sperm competition and sex

ratio. For assays done in experimental blocks, we added experimental block as a

random factor, tested for significant effects using ANOVA, and performed post-hoc

Tukey HSD tests for analyses with significant treatment effect. We calculated

bootstrap 95% confidence intervals around treatment means for ΔFitness by randomly

resampling 11 out of 12 data points and recalculating ΔFitness 10,000 times.

Theoretical models

We developed two population genetic models, identified by the location of the

compensatory locus: the Autosomal and X-linked models respectively. We assumed

discrete generations, and a life-cycle that proceeds as follows: (i) birth, (ii) selection

on offspring survival, (iii) meiosis and mutation, (iv) selection on male mating

success. Both loci in each model are biallelic. The Y-linked locus, **Y**, has wild-type Y

and mutant y alleles with frequencies q_Y and $q_y = 1 - q_Y$. The compensatory locus is

denoted **A** (with alleles A and a and frequencies q_A and $q_a = 1 - q_A$) for the autosomal

model, and **X** (with alleles X and x and frequencies q_X and $q_x = 1 - q_X$) for the X-

linked model (wild-type alleles are indicated by capital letters, mutant alleles in

lowercase). Following standard population genetics theory, **Y** is effectively haploid

with strictly paternal inheritance, while the autosomal or X-linked compensatory loci

(**A** or **X**) are diploid with bi-parental inheritance.

We based the fitness expressions used in our models loosely on our

experimental results. The mutant y chromosome increases male mating success by a

factor of $1 + s_m$ relative to the ancestral Y chromosome. Offspring survival depends on

the paternal genotype at **Y** and the maternal genotype at the compensatory locus (**A** or

X), as sires effects offspring viability (33) and any such negative effects should create

selection for females to compensate by maternal effects. Offspring sired by mutant y

males may experience reduced survival, depending on their mother's genotype at the

compensatory locus: matings involving parental genotypes $[y : AA]$, $[y : Aa]$, and $[y :$

$aa]$ result in offspring relative fitness of $1 - s_o$, $1 - h_o s_o$, and 1 respectively. To make

the models more general, and analytically tractable, we allowed females carrying the

mutant a allele to incur a 'cost of compensation' when mating with wild-type (Y)

males: matings involving parental genotypes $[Y : AA]$, $[Y : Aa]$, and $[Y : aa]$ result in

relative offspring fitnesses of 1, $1 - h_c s_c$, and $1 - s_c$ respectively (Table **1**). Similar to

standard theories of compensatory evolution (e.g., (24, 25)) each of the mutant alleles

(y and either a or x) is deleterious in isolation. In our models, however, compensation

requires that both parents have the appropriate mutant genotype at the other locus.

We model evolutionary invasion and single co-evolutionary cycles between

the male-beneficial Y-linked locus and the compensatory locus (see (34) for a similar

approach in the context of mito-nuclear coevolution). A bout of coevolution begins

with the invasion of a single-copy mutant y chromosome in a population initially

fixed for the wild-type A (or X) allele at the compensatory locus. The mutant y

evolves under net positive selection if the increase in male mating success outweighs

the reduction in offspring survival (i.e., if $\delta = (s_m - s_o) > 0$) until it becomes fixed (q_y

$= 1$) or is lost from the population ($q_y = 0$). The compensatory locus (**A** or **X**) evolves

under recurrent mutation and selection, with the population initially fixed for the wild

type A allele ($q_a = 1$ for the autosomal model) or X ($q_x = 1$ for the X-linked model).

For simplicity we assume one-way mutation from $A \rightarrow a$ at a rate u_a, and $X \rightarrow x$ at a

rate u_x per meiosis. If females experience a 'cost of compensation' (i.e., when $s_c > 0$),

the mutant compensatory locus will evolve under purifying selection until the mutant

y chromosome reaches a threshold frequency, denoted $\tilde{q}_y^A$ and $\tilde{q}_y^X$, at which y becomes

selectively favoured. A co-evolutionary cycle completes when the mutant allele
becomes fixed at both loci (i.e., $q_y = q_a = 1$ or $q_y = q_x = 1$).

Our analytic results all assume large population sizes (negligible drift), and an
equal sex ratio. To identify the conditions under which rare mutant alleles at each
locus can spread during key points of an coevolutionary cycle, we performed a linear
stability analysis for each model under three different scenarios: (i) invasion of mutant
a y chromosome into populations initially fixed for the wild-type allele at both loci
(initial frequencies of $q_y = 0$, and $q_i = 0$; where $i \in \{a, x\}$); (ii) invasion of a rare
mutant compensatory allele into a population fixed for y ($q_y = 1$, and $q_i = 0$); and (iii)
invasion of a mutant compensatory allele into a population with an arbitrary initial
frequency of y ($q_y = q_y$, $q_i = 0$). Mutant alleles can invade (i.e., the initial equilibrium
is unstable) when the leading eigenvalue of the Jacobian of the system of recursions is
greater than one ($\lambda_L > 1$; 43). When the initial frequency of y is arbitrary, solving the
expression $\lambda_L > 1$ for q_y yields the threshold frequency at which compensatory
mutations become selectively favoured, ($\tilde{q}_y^A$ or $\tilde{q}_y^X$). A full derivation of all models
and analytic results is provided in *SI Appendix B*.

We complemented our analytic results with stochastic Wright-Fisher
simulations for the Autosomal and X-linked models with population size N and an
equal sex ratio. We estimated two important properties of co-evolutionary cycles from
the simulations: (i) the probability of invasion of single copy autosomal (Π_A) and X-
linked (Π_X) compensatory mutations into populations initially fixed for the mutant y
chromosome; and (ii) the total time required to complete a single bout of coevolution
between the Y-linked and compensatory loci (T_A and T_X for the autosomal and X-
linked models respectively) under recurrent mutation, selection, and drift. Additional
details and R code for the simulations can be found in *SI Appendix B*, and online at
https://github.com/colin-olito/sexChromCoAdapt.

References

1. J. E. Mank, Small but mighty: the evolutionary dynamics of W and Y sex chromosomes. *Chromosome Res.* **20**, 21–33 (2012).

2. B. Vicoso, B. Charlesworth, Evolution on the X chromosome: unusual patterns and processes. *Nat. Rev. Genet.* **7**, 645–653 (2006).

3. M. Parisi, *et al.*, Paucity of Genes on the Drosophila X Chromosome Showing Male-Biased Expression. *Science* **299**, 697–700 (2003).

4. J. M. Ranz, C. I. Castillo-Davis, C. D. Meiklejohn, D. L. Hartl, Sex-Dependent Gene Expression and Evolution of the Drosophila Transcriptome. *Science* **300**, 1742–1745 (2003).

5. A. K. Chippindale, W. R. Rice, Y chromosome polymorphism is a strong determinant of male fitness in Drosophila melanogaster. *Proc. Natl. Acad. Sci.* **98**, 5677–5682 (2001).

6. C. Rohmer, J. R. David, B. Moreteau, D. Joly, Heat induced male sterility in Drosophila melanogaster: adaptive genetic variations among geographic populations and role of the Y chromosome. *J. Exp. Biol.* **207**, 2735–2743 (2004).

7. R. M. Griffin, D. Le Gall, H. Schielzeth, U. Friberg, Within-population Y-linked genetic variation for lifespan in Drosophila melanogaster. *J. Evol. Biol.* **28**, 1940–1947 (2015).

8. B. A. Payseur, D. C. Presgraves, D. A. Filatov, Introduction: Sex chromosomes and speciation. *Mol. Ecol.* **27**, 3745–3748 (2018).

9. W. R. Rice, Sex Chromosomes and the Evolution of Sexual Dimorphism. *Evolution* **38**, 735–742 (1984).

10. J. A. Coyne, H. A. Orr, "Two rules of speciation" in *Speciation and Its Consequences*, D. Otte, J. A. Endler, Eds. (Sinauer Associates, Inc., 1989), pp. 180–207.

11. H. A. Orr, The population genetics of speciation: the evolution of hybrid incompatibilities. *Genetics* **139**, 1805–1813 (1995).

12. C. I. Wu, *et al.*, Sexual isolation in Drosophila melanogaster: a possible case of incipient speciation. *Proc. Natl. Acad. Sci.* **92**, 2519–2523 (1995).

13. P. Capy, *et al.*, Sexual isolation of genetically differentiated sympatric populations of Drosophila melanogaster in Brazzaville, Congo: the first step towards speciation? *Heredity* **84**, 468 (2000).

14. J. A. Alipaz, C. Wu, T. L. Karr, Gametic incompatibilities between races of Drosophila melanogaster. *Proc. R. Soc. Lond. B Biol. Sci.* **268**, 789–795 (2001).

15. W. R. Rice, B. Holland, The enemies within: intergenomic conflict, interlocus
contest evolution (ICE), and the intraspecific Red Queen. *Behav. Ecol.*
Sociobiol. **41**, 1–10 (1997).

16. B. Lemos, L. O. Araripe, D. L. Hartl, Polymorphic Y chromosomes harbor cryptic
variation with manifold functional consequences. *science* **319**, 91–93 (2008).

17. G. Arnqvist, L. Rowe, *Sexual Conflict* (Princeton University Press, 2005).

18. L. Van Valen, The red queen. *Am. Nat.* **111**, 809–810 (1977).

19. G. A. Parker, "Sexual selection and sexual conflict" in *Sexual Selection and*
Reproductive Competition in Insects, (Academic Press, 1979), pp. 123–166.

20. M. F. Camus, J. B. W. Wolf, E. H. Morrow, D. K. Dowling, Single Nucleotides in
the mtDNA Sequence Modify Mitochondrial Molecular Function and Are
Associated with Sex-Specific Effects on Fertility and Aging. *Curr. Biol.* **25**,
2717–2722 (2015).

21. S. Pitnick, Male size influences mate fecundity and remating interval in
Drosophila melanogaster. *Anim. Behav.* **41**, 735–745 (1991).

22. L. Partridge, M. Farquhar, Lifetime mating success of male fruitflies (Drosophila
melanogaster) is related to their size. *Anim. Behav.* **31**, 871–877 (1983).

23. D. C. Presgraves, Sex chromosomes and speciation in Drosophila. *Trends Genet.*
24, 336–343 (2008).

24. M. Kimura, The role of compensatory neutral mutations in molecular evolution. *J.*
Genet. **64**, 7 (1985).

25. D. M. Weinreich, L. Chao, Rapid Evolutionary Escape by Large Populations from
Local Fitness Peaks Is Likely in Nature. *Evolution* **59**, 1175–1182 (2005).

26. J. A. Ågren, M. Munasinghe, A. G. Clark, Sexual conflict through mother's curse
and father's curse. *Theor. Popul. Biol.* (2019)
https:/doi.org/10.1016/j.tpb.2018.12.007 (September 10, 2019).

27. D. Arbuthnott, E. M. Dutton, A. F. Agrawal, H. D. Rundle, The ecology of sexual
conflict: ecologically dependent parallel evolution of male harm and female
resistance in Drosophila melanogaster. *Ecol. Lett.* **17**, 221–228 (2014).

28. W. R. Rice, *et al.*, Inter-locus antagonistic coevolution as an engine of speciation:
Assessment with hemiclonal analysis. *Proc. Natl. Acad. Sci.* **102**, 6527–6534
(2005).

29. P. Jensen, *et al.*, Inbreeding effects on standard metabolic rate investigated at
cold, benign and hot temperatures in Drosophila melanogaster. *J. Insect Physiol.*
62, 11–20 (2014).

30. R Core Team, *R: A language and environment for statistical computing. R*
Foundation for Statistical Computing, Vienna, Austria. URL https://www.R-
project.org/ (2018).

31. D. Bates, M. Maechler, B. Bolker, S. Walker, Fitting Linear Mixed-Effects
Models Using lme4 | Bates | Journal of Statistical Software. *J. Stat. Softw.* **67**, 1–
48 (2015).

32. T. Hothorn, *et al.*, *multcomp: Simultaneous Inference in General Parametric*
Models (2017) (October 27, 2018).

33. F. García-González, L. W. Simmons, The evolution of polyandry: intrinsic sire
effects contribute to embryo viability. *J. Evol. Biol.* **18**, 1097–1103 (2005).

34. T. Connallon, F. M. Camus, E. H. Morrow, D. K. Dowling, Coadaptation of
mitochondrial and nuclear genes, and the cost of mother's curse. *Proc. R. Soc. B*
Biol. Sci. **285**, 20172257 (2018).

35. S. P. Otto, T. Day, *A Biologist's Guide to Mathematical Modeling in Ecology and*
Evolution (Princeton University Press, 2007).

547
548

549 Table 1: Fitness expressions for the Y-linked locus (**Y**) influencing male siring
550 success, and the viability of offspring resulting from all possible combinations of
551 parental genotypes at **Y** and the compensatory locus **A** (or **X**).

Male Fitness

Male **Y** genotype	Mating success
Y	1
y	$1 + s_m$

Offspring viability

	Mother's Genotype		
Father's **Y** genotype	AA, XX	Aa, Xx	aa, xx
Y	1	$1 - h_c s_c$	$1 - s_c$
y	$1 - s_o$	$1 - h_o s_o$	1

Note: Subscripts for the offspring fitness expressions indicate ...

553
Fig. 1: Schematic of the round robin experimental cross design. Each of the five wild-type populations is color-
coded: LH_M (L, red), *Innisfail* (I, blue), *Odder* (O, green), *Dahomey* (D, orange), and *Tasmania* (T, purple). We
crossed each wild-type population (large boxes) with two others from either a different continent or climate. This
procedure generated four novel genotypes from each cross, so that every combination of the novel sex
chromosome pairs experienced a different autosomal genetic background. In total, 10 *Novel X* and 10 *Novel Y*
genotypes were generated (small boxes). Coloured bars represent the two major chromosomes (II and III) and
chromosome X, with chromosome Y depicted by a half arrow. *Novel* genotypes are annotated as 'genetic
background – origin of sex chromosome', with the novel sex chromosome indicated by a X or Y subscript.

Fig. 2: Male reproductive fitness assays at generation 0. (A) Relative reproductive fitness for *wt* (circles), *novel X* (triangle), and *novel Y* (diamonds) males. We detected statistically significant ($P = 2.45e^{-04}$) increases in reproductive fitness for both *novel X* and *novel Y* males relative compared to the wild-type populations (Tukey HSD; $P < 0.05$). Mean (±SE) of fitted values from the linear model. The raw data is plotted as grey points. (B) Relative reproductive fitness for *wt* (circles) and *novel XY* (squares) treatment groups. There was no significant difference between *wt* and *novel XY*. Mean (±SE) of fitted values from the linear model. The raw data is plotted as grey points. (C) Change in relative fitness between the *wt* and the *novel* populations. ΔFitness = $\omega_{novel\ population}$ - $\omega_{wild-type}$ with bars indicating bootstrap 95% confidence (see **Materials and Methods**). 10 out of 20 novel populations (*Novel X*: triangle, *novel Y*: diamond) had significantly higher fitness than their *wt* counterpart (indicate by asterisks), and 16 populations had changed in a positive direction, which was more than expected by chance ($P = 0.01$). Each of the five *wild-type*s are indicated underneath their four novel populations. Capital letters indicate which population the novel sex chromosome originates from (D: *Dahomey*, I: *Innisfail*, L: LH$_M$, O: *Odder*, and T: *Tasmania*) and the novel sex chromosome is denoted by X or Y subscripts.

Fig. 3: Reproductive fitness assays at generation 25. We selected 8 of the 10 significant populations for the evolution experiment. (A) Relative reproductive fitness for *wt* (black circle), *novel X* (triangle)**,** and *novel Y* (diamonds) males. *Novel* treatment males no longer had a higher fitness relative to *wt* males after 25 generations of experimental evolution. Mean (±SE) of fitted values from the linear model. The raw data is plotted as grey points. (B) Change in relative fitness between *wt* and *novel X* (triangles) and *novel Y* (diamonds) populations (ΔFitness = $\omega_{\text{novel population}}$ - $\omega_{\text{wild-type}}$), with bootstrap 95% confidence intervals (see **Materials and Methods**). The asterisk indicates whether ΔFitness was significantly greater than zero (Tukey HSD; $P < 0.05$). Each of the *wt* populations are indicated underneath their four *novel* populations. Capital letters indicate which population the novel sex chromosome originates from (D: *Dahomey*, I: *Innisfail*, L: LH_M, O: *Odder*) and the novel sex chromosome is denoted by X or Y subscripts.

Fig. 4: Summary of theoretical results. (A) Threshold frequencies of the mutant y chromosome at which a new compensatory mutation will experience positive selection for the Autosomal ($\tilde{q}_y^A$, solid lines) and X-linked ($\tilde{q}_y^X$, dashed lines) models respectively). As the mutant y chromosome sweeps to fixation, X-linked compensatory mutations will become selectively favoured earlier than autosomal ones. Results are shown for the case of additive fitness ($h_o = h_c = 1/2$), $s_m = 0.1$, and $s_c = 0.005$. (B) The relative probability of invasion for new Autosomal and X-linked compensatory mutations into populations initially fixed for the mutant y chromosome is always greater than or equal to 1, suggesting that when compensatory evolution is limited by mutational variation, the probability of invasion will be higher for X-linked compensatory mutations except under complete dominance. For simplicity we assumed equal dominance for compensatory fitness effects (i.e., $h = h_o = h_c$). Results are shown for three different values of δ (recall that $\delta = s_m - s_o$), where $N = 1,000$, $s_m = 0.1$, and $s_c = 0.005$. Each point indicates the mean of 10^6 replicate Wright-Fisher simulations for $\delta = 0.005$ and $\delta = 0.05$, but 5.0×10^6 for $\delta = 0.095$. (C) The relative time to complete a co-evolutionary cycle for the Autosomal vs. X-linked models plotted as a function of the relative mutation rates at the compensatory and Y-linked loci. To explore the consequences of slow vs. fast compensatory dynamics, μ_Y was held constant at 10^{-3}, while μ_i (where $i \in \{A, X\}$) was incremented from 10^{-7} to μ_Y. At low values of μ_i/μ_Y, the rate of compensatory evolution is limited by mutational variation, but at intermediate to high values, the mutant y chromosome and compensatory mutations are more likely to segregate simultaneously. Results are shown for two different 'costs of compensation' (s_c) and three different dominance scenarios (we again assume equal dominance, $h_o = h_c$), and population size $N = 1,000$.