A hierarchical framework for segmenting movement paths

Wayne M. Getz[1,2]

[1]Dept. ESPM, UC Berkeley, CA 94720-3114, USA

[2]School of Mathematical Sciences, University of KwaZulu-Natal
Private Bag X54001, Durban 4000, South Africa

Abstract

Comparative applications of animal movement path analyses are hampered by the lack of a comprehensive framework for linking structures and processes conceived at different spatio-temporal scales. Although many analyses begin by generating step-length (SL) and turning-angle (TA) distributions from relocation time-series data—some of which are linked to ecological, landscape, and environmental covariates—the frequency at which these data are collected may vary from sub-seconds to several hours, or longer. The kinds of questions that may be asked of these data, however, are very much scale-dependent. It thus behooves us to clarify how the scale at which SL and TA data are collected and relate to one another, as well as the kinds of ecological questions that can be asked. Difficulties arise because the information contained in SL and TA time series is not semantically aligned with the physiological, ecological, and sociological factors that influence the structure of movement paths. I address these difficulties by classifying movement types at salient temporal scales using two different kinds of vocabularies. The first is the language derived from behavioral and ecological concepts. The second is the language derived from mathematically formulated stochastic walks. The primary tools for analyzing these walks are fitting time-series and stochastic-process models to SL and TA statistics (means, variances, correlations, individual-state and local environmental covariates), while paying attention to movement patterns that emerge at various spatial scales. The purpose of this paper is to lay out a more coherent, hierarchical, scale-dependent, appropriate-complexity framework for conceptualizing path segments at different spatio-temporal scales and propose a method for extracting a simulation model, referred to as M^3, from these data when at a relatively high frequencies (ideally minute-by-minute). Additionally, this framework is designed to bridge biological and mathematical movement ecology concepts; thereby stimulating the development of conceptually-rooted methods that facilitates the formulation of our M^3 model for simulating theoretical and analyzing empirical data, which can then be used to test hypothesis regarding mechanisms driving animal movement and make predications of animal movement responses to management and global change.

Keywords: Fundamental movement elements (FuMEs), canonical activity modes (CAMs), diel activity routines (DARs), life-history movement phases (LiMPS), multi-CAM metaFuME Markov (M^3) models, biased correlated random walk models

Introduction   32

The movement of organisms over landscapes can be usefully analyzed from various viewpoints.   33
Four are suggested by the conceptual framework for movement ecology proposed by Nathan et al.   34
(2008), depending on which questions constitute the focal interest—why move? how is movement   35
executed? where to move? and how does the immediate environment, as well as the internal state   36
of individuals, impact movement? Several more viewpoints arise when we focus on the structure   37
of the movement path itself and how this structure may fill space at different spatio-temporal   38
and, hence, ecological relevant scales (Levin 1992)—from the sequencing of various high-resolution   39
movement elements to the emergence of seasonal home ranges and habitat utilization distributions   40
(Worton 1987, Morellet et al. 2013, Torney et al. 2018, Wittemyer et al. 2019). The current lack of a   41
comprehensive framework for linking structures and processes conceived at different spatio-temporal   42
scales, however, hampers the synthesis of these viewpoints and of a comparative meta-analyses of   43
similar questions addressed across disparate species and spatio-temporal scales. With this in mind,   44
a more coherent, hierarchical, scale-dependent, appropriate-complexity framework than currently   45
exists is laid out in this paper for conceptualizing path segments at different spatio-temporal scales   46
and used to develop a model that captures the multimodal, correlated, random walk aspects present   47
in relatively high frequency (i.e., minute-by-minute) relocation data. Of considerable importance   48
is that this framework underscores the need to collect high resolution relocation data (minute or   49
subminute sampling intervals) whenever possible.   50

Nathan et al. (2008) argued that movement arises from the interactions of several processes and   51
motivations that can be understood through an analysis of external factors or stimuli, the internal   52
state of an organism (e.g., see Morelle et al. 2015), and the organism's mechanical and navigational   53
capacities. These interactions then result in a movement process that may be characterized by a   54
time series of relocation points that define movement paths (Siniff & Jessen 1969, Edelhoff et al.   55
2016). From these relocation time series, sequences of step lengths (SL) and turning angles (TA)   56
can be extracted (Kareiva & Shigesada 1983, Turchin 1998) and organized into time-stamped pairs   57
of values. Other data, such as accelerations (Nathan et al. 2012), audio (Sapir et al. 2010, Hurme   58
et al. 2019, Northrup et al. 2019) and magnetometer (Williams et al. 2017, Chakravarty et al. 2019)   59
data, as well as physiological measurements (e.g., temperature, blood glucose levels; Wilmers et al.   60
2015), may be recorded and used in the context of relocation point covariates. Additional covariates   61
may be environmental measurements associated with resistance to movement caused by landscape   62
topology and vegetation structure (Shepard et al. 2013, Getz & Saltz 2008, Spiegel et al. 2015), as   63
well as local environmental features inferred from remotely sensed electromagnetic spectral data   64
(e.g., visual and infra-red satellite imagery or lidar; Pettorelli et al. 2011, 2014, Bork & Su 2007).   65

In this paper, the question is addressed of how we may reconcile the fact that SL and TA   66
distributions are not directly interpretable in terms of behavioral and ecological processes (e.g.,   67
resting, stalking, hiding, gathering); but, depending, on the spatio-temporal scale of the reloca-   68
tion data, phenomenological interpretations are made using scale-appropriate activity designators   69
(Johnson et al. 2002, Owen-Smith & Goodall 2019, Giotto et al. 2015). A deeper understanding   70
of the connections between path SL and TA time series and the designation of the so-called behav-   71
ioral state of an individual at each of its relocation points requires that we find a way to reconcile   72
narratives constructed using two different kinds of vocabularies. The first is the language of bi-   73
ological designators arising from behavioral and ecological concepts. Our primary tools here are   74
the application of path segmentation methods (Nams 2014, Edelhoff et al. 2016, Seidel et al. 2018)   75

that include hidden Markov models (HMMs; Michelot et al. 2016, Franke et al. 2004, Langrock   76
et al. 2012, Zucchini et al. 2016) and behavioral change-point analyses (BCPA; Gurarie et al. 2009,   77
2016, Owen-Smith & Martin 2015). The second is the language of mathematical stochastic walks   78
(Kareiva & Shigesada 1983, Bartumeus et al. 2005, Patterson et al. 2008) that treats movement   79
relocation data as a time series of values and applies existing time-series methods (Chatfield 2016)   80
to finding the parameter values that best fit selected time-series models to movement paths, with   81
the possible inclusion of individual and environmental covariates (Jonsen et al. 2003, Preisler et al.   82
2004, Johnson et al. 2008, Fleming et al. 2014, Calabrese et al. 2016).   83

Moving between stochastic-walk formulations and biological designators in describing segments   84
of movement paths is akin to translating between phonologic (e.g., Latin, cyrillic) and logographic   85
(e.g., hieroglyphics, Chinese) scripts: much context-dependent interpretation needs to take place.   86
But, in fact, the situation is much more dire, since we do not have access to every character—i.e., to   87
the fundamental movement element (FuMEs, see glossary; Getz & Saltz 2008) of a movement path   88
sequence. Rather, we only have access to some kind of "average" value across several characters or   89
potential point-source values. The primary goal of this paper is to flesh out the temporal structure   90
of biological designators of movement-path segments, provide a more coherent insights into the   91
relationship between biological designators and stochastic-process formulations at different tempo-   92
ral scales, and provide a way for extracting useful stochastic process models from a hierarchically   93
framed understanding of move path structure.   94

A Pragmatic View of Path Segmentation   95

The view that large segments of individual movement paths can be parsed into strings of fundamen-   96
tal movement elements (FuMEs—see glossary; Fig. 1, Table 1) with characteristic frequencies and   97
patterns of occurrence is highly idealized. Identifying FuMES is generally a biomechanical prob-   98
lem that is beyond the scope and capabilities of analyses based on relocation data alone. Other   99
kinds of data, such as audio recordings, have been used to identify FuMEs: e.g., wingflaps of small   100
birds (Sapir et al. 2010, Northrup et al. 2019) or footfalls of foraging ungulates (Northrup et al.   101
2019). If the frequency of our relocation data, however, spans strings of around 10-100 FuMEs,   102
occurring either repetitively or in correlated groups of a couple of FuME types, then we may be   103
able to reliably identify different strings of predominantly one type of FuME, or stereotypical mixed   104
sequences of particular FuMES, which we call "metaFuMEs." With this definition, for relatively   105
high frequency relocation data, all the different metaFuMEs, in a set of metaFuME types, have   106
the same fixed length equal to the time between consecutive relocation points and each type has   107
its own characteristic bivariate SL and TA distribution, and auto- and cross-correlation time-series   108
coefficients (Chatfield 2016). Such sets of metaFuMEs can then be used as a basis for representing   109
different types of identifiable activities (e.g., foraging or focused travelling to a selected location),   110
referred to as canonical activity modes (CAMs—see glossary; Getz & Saltz 2008).   111

Segmenting relocation data at a metaFuME level only makes sense if the data have been sampled   112
at frequencies that lie somewhere between the time it takes to execute several typical FuMEs (i.e.,   113
on the order ten seconds) and the time it takes to perform a short-period CAM that is relevant   114
to movement ecologists (i.e., on the order of a few minutes, except for sharp burst of movement   115
related to predation or defense). Fortunately, current technologies facilitate the collection of data   116
at frequencies of 0.01 to 1 Hz in birds (Harel, Horvitz & Nathan 2016, Harel & Nathan 2018)   117
and mammals (de Weerd et al. 2015, McGavin et al. 2018) of even relatively small size. Beyond   118

GPS methods for collecting high frequency data are other methods such as reverse GPS (e.g., the   119
Atlas system; Weiser et al. 2016, Toledo et al. 2016); but reverse GPS is generally limited in spatial   120
extent to some tens or hundreds of square kilometers. Also, high-frequency movement data has been   121
collected using video equipment for small (e.g., ants) to moderately-sized (e.g., mice) organisms in a   122
laboratory setting (Spink et al. 2001, Kane et al. 2004, Delcourt et al. 2013). From a practical point   123
of view, though, particularly since the size of relocation data sets can be rather large when collected   124
at frequencies of around 0.01 to 1 Hz, it may be useful to carry out a metaFuME identification   125
process on selected segments of the full path W (such as CAMs), and then use a suitable method   126
to reconcile disparate identification efforts to obtain a consensus set of metaFuME types that can   127
be applied to the full movement path.   128

Assuming that a set of metaFuMEs has been identified, as discussed in more detail in succeeding   129
sections, then it may be possible to stably identify different kinds of short duration CAMs with   130
periods varying from around 10 metaFuMEs (i.e., a couple of minutes) to a few hours (Fig. 1,   131
Table 1). Following this, longer duration CAMs may be more conveniently parsed in terms of   132
several shorter duration CAMs rather than in terms of metaFuME types. To emphasize this, both   133
short and long duration CAMs are identified in the hierarchical scheme laid out in Table 1. Due to   134
the importance of circadian rhythms as physiological and behavioral drivers (Takahashi et al. 2008,   135
Yerushalmi & Green 2009, Hardin & Panda 2013), the diel cycle provides an empirically obvious   136
segmentation window for the identification of various types of diel activity routines (DARs; set of   137
green boxes in Fig. 1; also see Table 1). Several types of DARs may then be identified in terms of   138
differences in the type and frequency of their constituent CAMs. For example, different types of   139
DARs have been identified in terms of the portion of their diel cycle that various groups of lemurs   140
are active (Donati et al. 2016) and of the variation in the diel travel rates of turtles (Jonsen et al.   141
2006).   142

DARs, in turn, can be strung together to create life-history movement phases (LiMPs; sets of   143
red pentagons in Fig. 1) that, depending on the longevity of the species, may be periodic (e.g.,   144
annual migration; for a review see migration Milner-Gulland et al. 2011) or episodic (e.g., dispersal;   145
for a review see Bowler & Benton 2005). Finally, a sequences of LiMPs sequentially strung together   146
from the birth-to-death of an individual constitute its full life time track (LiTT; Nathan et al. 2008)   147
(indicated by the grey pentagons in Fig. 1; also see Table 1). Beyond LiMPs, LiTTs from several   148
individuals can be used to map out space-use by populations rather than just the movement path   149
of any single individual (Mueller & Fagan 2008).   150

This pragmatic view of metaFuME, CAM, DAR and LiMP segmentation of lifetime tracks, as   151
illustrated in Fig. 1, provides a hierarchical classification system of appropriate complexity (Getz   152
et al. 2018, Larsen et al. 2016) for the parsing of movement paths based on two different fixed   153
time-segment scales: metaFuMES (sampling interval at the order of HZ or 0.1 HZ) and DARs   154
(24 h). The metaFuMEs provide a basis for constructing CAMs as the biologically meaningful   155
segments of DARs; and DARs are used to support the construction of seasonally or life-history   156
relevant LiMPs. Given this ecological relevance of canonical activity modes and lifetime tracks,   157
the classification system presented here also provides a semantic framework for the generation   158
of quantitative narratives (Getz, Salter & Tallam 2019) needed to explain the role that movement   159
plays in the ecology and evolution of organisms. In the context of evolution, for example, we expect   160
natural selection to hone both the mechanics of performing FuMEs and the way these FuMEs are   161
organized into metaFuMEs, CAMs, DARs, and LiMPs to create lifetime tracks that maximize the   162
expected lifetime fitness of individuals. In particular, it is of considerable interest to assess how the   163

Figure 1: **Scale-dependent path segmentation categories of lifetime track segments.** The time scale applies to typical medium and large terrestrial animals and needs to be adjusted for aerial and marine species, and small animals, including invertebrates. Round (blue), oval (lilac), triangular (yellow to orange), square (green), pentagonal (red), and hexagonal (grey) icons are respectively used to represent N_1 FuME types, N_2 metaFuME types, N_3 CAM types, N_4 DAR types, N_5 LiMP types, and an open-ended number of n LiTT types (one for the life time of each movement group type to which an individual can belong). Horizontal whiskers on icons represent variations in the length of examples of the same type within a category (except for metaFuMEs which are the length of time between successive relocation points and DARs which are fixed to the biological relevant diel cycle) and vertical whiskers indicate some building-block variation within the same type (except for FuMEs which are much more stereotyped, and hence much less variable within types than other segmentation categories). Color shades within categories represent different types within those categories. Colorless strings of icons indicate how each category can be considered as a string of the next lower category elements (i.e., shapes and general color denote hierarchical level, while different shades of the same color indicate within level variation). Because FuMEs are hard to identify using relocation data only, a hierarchical segmentation of lifetime tracks will typically be supported by a metaFuME baseset rather than by a set of FuMEs themselves.

frequency and sequencing of different FuMEs and CAMs in the production of DARs and LiMPs [164] affect individual fitness (Hein et al. 2016, Cattarino et al. 2016). [165]

Beyond the relevance of diel cycles to movement behavior (Wittemyer et al. 2008), are lunar [166] (Polansky et al. 2010) and seasonal cycles (Marra et al. 2015); and even weekly cycles if the influence [167] of humans has some impact on the movement of animals in urban and suburban areas. The critical [168] nature of the diel scale in driving movement behavior is reflected in our existence of the fixed- [169] period DAR category in framework presented in Fig. 1. In contrast, the lunar cycle may be more [170] relevant to some organisms than others (Polansky et al. 2010). Thus beyond DARs only seasonally [171] or life-history relevant LiMPs segments are identified with seasonal segments only being relevant [172] to organisms with life-spans long enough to experience several seasonal cycles (Marra et al. 2015, [173] Allen et al. 2018). For some organisms, particular LiMPs, such as dispersal, occur only once; while [174] migration behavior may vary from year to year, influenced by inter-annual variations in climatic [175] conditions—perhaps linked to multiyear marine (Mysterud et al. 2001, Grémillet & Boulinier 2009) [176] or sunspot (Myers 2018) cycles. Further, periodically driven movement patterns may also change [177] with age (e.g., exploration when young) and life-history stage (e.g., elephants in musth). Thus, [178] beyond DARs, other cyclic patterns become either less obvious or more species-specific. [179]

Table 1: Hierarchical organization of a typical vertebrate life time track (LiTT). (Spatio-temporal scaling applies best to typical medium to large vertebrates, but not as well to small vertebrates, birds, or invertebrates.)

Scale/ Segment	Time	Space	Categories	Data	Some approaches and methods
	Constraint level: type of analysis				
FuMEs[*]	0.1–few secs	0–1000 cms	stride, trot, run twist, jump, flap	accelerometer video	machine learning[*†], Newtonian mechanics
	highly constrained: biomechanical analyses				
meta- **FuMEs**[*]	resolution of relocation data[*‡]	average step length	bivariate SL & TA[¶] distributions	SL & TA means, variances & correlations[††]	cluster analysis ensemble filtering
	base level: movement ecology analyses				
short duration **CAMs**[†]	0.1–10 mins	a few meters to a few kms	resting, browsing, riding thermals	secs to mins, metaFuMEs & covariate data	path segmentation[**]
	highly flexible: sub-hourly/hourly analysis				
long duration **CAMs**[†]	10-100 mins	a few meters to many kms	feeding traveling	mins to hours, short CAMs & covariate data	
	highly flexible: sub-diel analysis				
DARs[‡]	fixed 24-hr	diel range	central-place foraging ranging	24 hrs of data CAM sequences	periodograms (Fourier, wavelets)
	time constrained: diel cycle analysis				
LiMPs[§]	several days to a few months	home ranges and their shifts over time	philopatry, dispersal, migration, nomadic, recursion	DAR sequences	clustering methods, ideal-free/despotic distribution analyses
	strongly constrained: lunar and seasonal cycle analyses				
partial or whole **LiTTs**[#]	paths encompassing several LiMPs	habitat types	territorial ranging, migratory nomadic	LiMP sequences	GIS toolbox, deep-learning[*†], pattern anal.
	moderately constrained: question-at-hand analysis				

[*]Fundamental Movement Elements; [†]Canonical Activity Modes; [‡]Diel Activity Routines

[§]Life-history Movement Phases; [#]Lifetime Tracks; [¶]SL=step-length & TA=turning-angle

[*†]Machine/deep learning methods can be applied at any scale but may be particularly useful here

[*‡]Around 5 secs to 1 min

[**]Includes Hidden Markov Models (HMMs) and Behavioral Change-Point Analyses (BCPA)

[††]See Section on **Stochastic Walk Statisics** in main text

The Data Compatibility Problem   180

Parsing a movement path into its elements is somewhat like picking out words from a voice record- 181
ing. The human brain does this very well, as do modern digital machines using deep-learning 182
methods (Hinton et al. 2012, Zhang et al. 2018). In an extremely crude sense, FuMEs are com- 183
parable to syllables, metaFuMEs to words, short duration CAMs to sentences and long duration 184

CAMs to paragraphs. DARs may be thought of as single pages in a book-of-life, where each page can be identified as belonging to one of a rather limited number of types (e.g., a typical winter versus summer day or a day during a migratory versus non-migratory LiMP). Thus, the stories being told are rather boring, page-by-page repetitions, with important cyclic variations, as well as other variations due to environmental influences.

Despite the current, technology driven, exponential growth in the quantity and quality of movement data (including relocation, accelerometer, physiological, and environmental measurements; Williams et al. 2019), collecting the kind of multi-body-placement accelerometer and relocation data needed to identify the start and finish of individual FuMEs (i.e., with a spatial accuracy of centimeters rather than meters or even decimeters) may still be beyond the budgets of most movement ecology studies. Other technologies, such as audio (Sapir et al. 2010, Hurme et al. 2019, Northrup et al. 2019), magnetometer (Williams et al. 2017, Chakravarty et al. 2019), and video (Spink et al. 2001, Kane et al. 2004, Delcourt et al. 2013) may well be more cost effect in terms identifying the start and finish times of individual FuMES. Thus, for some time to come, relocation data collected at fixed frequencies slower than around 1 Hz are going to be significantly misaligned in time (at least to within half a second) with the start and finish of each of the FuMEs that make up a path segment of interest.

At this point, it is worth noting that accelerometer data alone has been used to parse out relatively short time scale (order of 10 secs) behavioral elements along movement paths using machine learning methods (Nathan et al. 2012, Thessen 2016, Wang et al. 2015). These behavioral elements, which almost certainly include several FuME steps, in reality are partial elements of more extensive short-duration CAMs that typically last tens of seconds to several minutes. Accelerometer data, for example, has been used to categorize standing, running, preening, eating, and active and passive flight CAMs in vultures with 80-90% accuracy (Nathan et al. 2012). Another analysis on baboons succeeded in using accelerometer data to categorize foraging, lying, sitting, running, standing, walking, groom (actor), groom (receiver) with high accuracy ($> 90\%$), except for grooming, which was considerably harder to identify, particularly when the focal individual was the receiver (Fehlmann et al. 2017). In pumas, accelerometer has been used to identify low versus high acceleration movement phases, resting, eating and grooming (Wang et al. 2015). In these examples, sitting/resting, grooming, preening and feeding CAMs are identifiable, but none of the basic FuMEs executed during the generation of these CAMs are individually identified. It may be possible for some FuMEs, however, such as the time taken to complete a FuME when walking versus running, to be identifiable directly from accelerometer data (as we find in modern digital watches that are able to monitor the number of steps we take while moving).

In summary, location data on its own (i.e., without accompanying subsecond accelerometer, acoustic, magnetometer or video data) is fundamentally incompatible with the identification of FuMEs because FuMEs are characterized by the movement of body parts while location data apply to a body as a whole. For this reason, it is worth reemphasizing the following.

1. Given that the identification of individual FuMEs from relocation data is not generally possible, we are left with the rather challenging task of extracting a set of metaFuME elements, where each metaFuME type is characterized by its own SL-TA distributional pair and correlations with and between consecutive and simultaneous values.

2. The complete set of metaFuME types, once identified, can then be used as a basis for constructing an appropriate complexity (Getz et al. 2018, Larsen et al. 2016), hierarchical frame-

work of CAM, DAR and LiMP segments, where the latter is used to classify different types   229
of lifetime tracks (Table 1), beginning with CAM sequences constructed using a multi-CAM   230
metaFuME Markov (M^3) modeling approach described in more detail below.   231

Current Path Analysis Methods   232

Before considering how to find a set of metaFuME distribution set that provides an appropriately   233
defined "best fit" to an empirical walk W represented in 2-D, by the relocation time-series   234

$$W = \big\{ \big(t; x(t), y(t)\big) \big| t = 0, ..., t_f \big\} \tag{1}$$

let us briefly review the current state of path analysis that is relevant for analysing relocation data   235
sampled at scales typically appropriate for identifying CAMs and higher-level patterns (Fig. 2).   236
Note that the temporal resolution of the data will determine whether it is possible to identify short   237
duration CAMs (e.g., vigilance behavior during grazing; Fortin et al. 2004), long duration CAMs   238
(e.g., heading to water; Polansky et al. 2015), or only movement patterns at diel scales and beyond   239
(Spiegel & O'Farrell 2019, Owen-Smith 2013, Giotto et al. 2015).   240

At any time scale, movement time series may be analysed using purely statistical methods   241
applied to the SL and TA time series derived from W to generate various running statistics. These   242
statistics may include the means, variance, autocorrelations of SL and TA time series data and   243
cross correlations between the two (Chatfield 2016, McCulloch & Cain 1989, Bergman et al. 2000,   244
Byers 2001). Running versions of these statistics can then be used to identify points in time   245
where abrupt changes in their values occur, using methods referred to as behavioral change-point   246
analyses (BCPAs; Owen-Smith & Martin 2015, Chen & Gupta 2011, Matteson & James 2014,   247
Killick & Eckley 2014, Gurarie et al. 2009, 2016; Fig. 2) or, more generally, path segmentation   248
methods (PS; Nams 2014, Edelhoff et al. 2016, Seidel et al. 2018).   249

Beyond purely statistical characterizations of the SL and TA time series is the notion that some   250
sort of state (typically behavioral; but possibly physiological, disease, etc.) can be associated with   251
each location point in the walk W (Patterson et al. 2008). A very simple example is the stop/move   252
representation of cattle grazing paths (Zhao & Jurdak 2016). The most widely used methods for   253
identifying subsets of points that reflect some underlying state or canonical activity mode (CAM)   254
involve hidden Markov models (HMM) (Michelot et al. 2016, Franke et al. 2004, Langrock et al.   255
2012, Zucchini et al. 2016) (Fig. 2), although an independent mixture modeling approach has been   256
shown to be equally viable, if not better (Goodall et al. 2017). These methods are particularly   257
germane when covariate data are available at each location, such as accelerometer and temperature   258
data from GPS collars, remotely sensed physiological information (Cross et al. 2016), vegetation   259
structure (Tsalyuk et al. 2019), or physical landscape features that, for example, facilitate or retard   260
movement (Zeller et al. 2012). The task of an HMM is to find a best estimate of which state of a   261
current set of "hidden" states H_i belonging to a set of n possible states $\mathcal{H} = \{H_1, ..., H_n\}$ should   262
be assigned to the individual when at the location $(x(t), y(t))$.   263

HMMs can be applied at any temporal scale, including the metaFuME scale, depending only on   264
the relocation and covariate data. Other methods, such as step-selection function (SSF) methods,   265
which address the question of where am I likely to move next as a function of landscape covariates,   266
can be applied at any scale (Fortin et al. 2005, Panzacchi et al. 2016, Thurfjell et al. 2014, Avgar   267
et al. 2016). SSFs, however, address different kinds of questions at different scales: they have   268

Figure 2: **Current scale-dependent analytical methods for analysing movement paths.** The temporal range is only suggestive and best applied to medium and large vertebrates. Also the image placements are not precise and some methods, such as machine learning (Thessen 2016), can be applied to data at any scale, but here are associated with the scale at which they are likely to be most useful. In addition, deep learning (useful for identifying different types of long-term patterns) is actually a subset of machine learning (where other deep-learning techniques, such as random forests and support vector machines have been applied to accelerometer data; Nathan et al. 2012, Fehlmann et al. 2017). Stochastic walks include correlated and biased random walks (Johnson et al. 2008). Space-time residence analyses represents a family methods that include first-passage-time (FTP; Fauchald & Tveraa 2003, McKenzie et al. 2009) and related approaches (Torres et al. 2017), while recurrence analyses cover a plethora of methods used to identify recursive movement patterns (Berger-Tal & Bar-David 2015, Bar-David et al. 2009). The melange of images is extracted from publications in the literature (Nathan et al. 2012, Panzacchi et al. 2016, Morales et al. 2004, DeCesare et al. 2012, Wittemyer et al. 2008, Lyons et al. 2013, Abrahms et al. 2017, Fleming & Calabrese 2017, Pohle et al. 2017, Torres et al. 2017, Gurarie et al. 2017), as well as created for this publication.

more of a behavioral interpretation at a 1-minute scale while more of an ecological (e.g., resource   269
use) interpretation at multi-hour scales. On the other hand, habitat or resource-selection function   270
(HSF or RSF) methods (Boyce et al. 2002, McGarigal et al. 2016, DeCesare et al. 2012) are, by   271
their very nature, low resolution models that typically apply at a seasonal level and assume non-   272
autocorrelated location data. When RSFs are evaluated at a lifetime level over many individuals   273
in a population, they are called niche models—but such models no longer rely on movement data,   274
only on point location data (Townsend Peterson et al. 2007).   275

Other general time-series or signal-analysis methods that have been co-opted by movement   276
ecologists include wavelet analysis (Wittemyer et al. 2008, Polansky et al. 2010)—which provides   277
interpretations at the resolution of one to several orders of magnitude greater than the frequency   278
at which data are collected (e.g., daily movement to monthly movement patterns for data collected   279
at the frequency of 1-hour for at least a year)—and machine learning approaches (Thessen 2016,   280
Valletta et al. 2017) that have been applied both to location (Bennison et al. 2018) and accelerometer   281
data (Nathan et al. 2012) (Fig. 2). These methods, however, do not have an extensive history of   282
application in movement ecology, but are likely to become more ubiquitous as the field of movement   283

ecology matures. ²⁸⁴

Stochastic Walk Statistics ²⁸⁵

As discussed earlier, identifying a set of FuMEs is a problem that requires biomechanical (Delp & ²⁸⁶
Loan 2000), audio (Sapir et al. 2010), or other kinds of covariate data (Chakravarty et al. 2019) ²⁸⁷
than relocation data on its own. With relocation data alone, a hierarchical analysis of a path can ²⁸⁸
only be underpinned by elements that are derived statistically from the relocation time series W ²⁸⁹
(Eq.1). ²⁹⁰
From this time series we can extract a 1-D time series of step lengths and another 1-D time ²⁹¹
series of turning angles as follows: ²⁹²

- Generate the step-length (SL) time series ²⁹³

$$S = \{(t, s(t)) | \ t = 1, ..., t_f\} \tag{2}$$

  using the derived values ²⁹⁴

$$s(t) = \sqrt{(x(t) - x(t-1))^2 + (y(t) - y(t-1))^2} \tag{3}$$

- Generate the turning-angle (TA) time series ²⁹⁵

$$A = \{(t, a(t)) | \ t = 2, ..., t_f\} \tag{4}$$

  using the derived values ²⁹⁶

$$a(t) = \arctan\left(\frac{y(t) - y(t-1)}{x(t) - x(t-1)}\right) - \arctan\left(\frac{y(t-1) - y(t-2)}{x(t-1) - x(t-2)}\right) \tag{5}$$

From these two time series we can obtain their means and variances respectively denoted by (μ_ℓ, σ_ℓ), ²⁹⁷
$\ell = s$, a, and then use them compute the following three variation related time series: ²⁹⁸

- Generate two normalized (by the appropriate variances) "running-term autocorrelation" time ²⁹⁹
  series ³⁰⁰
$$V^{\ell\ell} = \{(t, v^{\ell\ell}(t)) | \ t = t_b, t_b + 1, ..., t_f\} \quad \text{for } \ell = s \ (t_b = 2), \ a \ (t_b = 3), \tag{6}$$

  where ³⁰¹
$$v^{\ell\ell}(t) = \frac{(\ell(t) - \mu_\ell)(\ell(t-1) - \mu_\ell)}{\sigma_\ell} \tag{7}$$

- Generate a normalized "running-term cross-correlation" time series ³⁰²

$$V^{as} = \{(t, v^{as}(t)) | \ t = 2, ..., t_f\} \tag{8}$$

  where ³⁰³
$$v^{as}(t) = \frac{(a(t) - \mu_a)(s(t) - \mu_s)}{\sqrt{\sigma_a \sigma_s}} \tag{9}$$

Figure 3: **Parsing the walk W into a set of metaFuMEs.** Associated with each point t is a set of running values $\big(s(t), a(t), v^{ss}(t), v^{aa}(t), v^{as}(t), \text{other possible covariates}\big)$. Cluster analysis may be used to generate metaFuME categories. For purposes of illustration, we depict three types of metaFuMEs with associated lists of time steps—i.e., metaFuME sequence index sets—that can then be used to define the ensemble of values associated with each metaFuME type: blue $= T_1$, red $= T_2$, and green $= T_3$. Note that we expect transition points between strings of metaFuMEs of one type (broken line segments), such as $t-6$, $t-1$ and $t+4$, could well be outliers in the clustering process. Outliers may be assigned to the cluster to which they are closest in some suitably defined sense (as indicated by the colors of the broken lines).

We are now challenged with the task of using the five time series above (step length, turning angle, two auto correlation and one cross correlation)—and, perhaps, other covariate data when available (e.g., acoustic, accelerometer, local environmental)—to parse W (Eq. 1) into several, say Λ, different single-step metaFuME ensembles (Fig. 3). The most obvious approach is to use clustering methods, but this should be accompanied by some sort of range standardization procedure for each variable to improve the performance of the clustering algorithm (Van Moorter et al. 2010). Each of these Λ ensembles, belonging to the same metaFuME type, can now be regarded as a set of drawings from the same bivariate distribution $D_\lambda(s, a)$, $\lambda = 1, ..., \Lambda$. If the relocation sampling period, except for speed bursts, encompasses around 10-50 FuMEs (i.e., more than around 5 seconds but less than about a minute—e.g, fast walking in humans crudely corresponds to 2 steps per second, though most movement is slower than this); then, in the proposed hierarchical segmentation framework, we are in the metaFuME segment zone (Fig. 1, Table 1). If the relocation sampling period is on the order of tens of minutes to several hours, then each relocation likely encompasses hundreds to tens of thousands of FuMEs. In this case, set of states are more appropriately identified as short or long duration CAMs than as metaFuMEs.

Once a set of metaFuMEs has been identified from a particular ensemble of segments of the same CAM or DAR type, along with the accompanying metaFuME sequence index sets (Fig. 3), we are still faced with the task of reconciling various metaFuME sets obtained from different CAM or DAR ensembles into a comprehensive metaFuME set. Such a set will then underlie all LiMPs of a particular type, but not necessarily across a complete lifetime track, because we might expect metaFuME sets for, say, juveniles versus adults to be different for same species. Many different types of cluster analyses (McGarigal et al. 2016) can be tried to identify metaFuMEs. Also, multicriteria optimization methods using, for example, genetic algorithms or approximate Bayesian computations (Coello 2003, Odu & Charles-Owaba 2013, Marin et al. 2012) could be used to reconcile metaFuME sets obtained from ensembles of different types of segments. Such investigations, however, are well beyond the scope of this "concepts" paper, but are undertaken elsewhere (Getz, Vissat & Salter 2019).

Once the time series W and, hence, the corresponding SL and TA time series have been parsed into ensembles and organized into a set of Λ metaFuMEs, the associated metaFuME sequence index sets T_λ, $\lambda = 1, ..., \Lambda$, (Fig. 3) can be identified. The distributions of all the joint pairs $(s(t), a(t))$, where $t \in \mathbf{T}_\lambda$, $\lambda = 1, ..., \Lambda$, can then be fitted to an appropriate metaFuME bivariate distributions $D_\lambda(s, a)$, $\lambda = 1, ..., \Lambda$ and the metaFuME sequence index set ensemble $\mathcal{T}_\Lambda = \{T_\lambda | \lambda = 1, ..., \Lambda\}$ can be used to estimate transition rates among metaFuME sequences. In essence, we have estimated a set of bivariate metaFuME distributions

$$\mathcal{D}_\Lambda = \{D_1(s, a), ..., D_\Lambda(s, a)\}. \tag{10}$$

and a matrix of values $(\mathbf{P})_{\lambda\kappa} = p_{\lambda\kappa}$ (with λ, $\kappa = 1, ..., \Lambda$) that represent the probability of sampling values from the distribution $D_\lambda(s, a)$ when the previous drawing was from the distribution $D_\kappa(s, a)$.

We can now use the set of distributions $\mathcal{D}_\Lambda$ and transition matrix $\mathbf{P}$, essentially as a Markov metaFuME movement model, to construct a CAM segment of a walk $W(t_f)$. We do this by first generating a set of T drawings (the hat notation represents drawings)

$$T \text{ drawings} := \quad \{(\hat{s}_t, \hat{a}_t) \text{ sampled from } D_\kappa(s, a) \text{ with probability } p_{\lambda\kappa} | t = 0, ..., t_f - 1\} \tag{11}$$

and then generating a segment of $W(t_f)$ from these drawings using the following equations (Patterson et al. 2008), where $\theta(t)$ is the absolute angle of heading at time t

$$\begin{aligned}
x(t+1) &= x(t) + \hat{s}_t \cos(\theta(t) + \hat{a}_t) \\
y(t+1) &= y(t) + \hat{s}_t \sin(\theta(t) + \hat{a}_t) \\
\theta(t+1) &= \theta(t) + \hat{a}_t
\end{aligned} \tag{12}$$

Thus, in short, we use the extracted identified metaFuME-distribution-set and Markov transition-matrix pair $(\mathcal{D}_\Lambda, \mathbf{P})$ to generate a CAM segment of a walk W that is one instantiation of an ensemble of CAM segments generated using Eqs. 11 and 12.

To generate different sets of CAM sequences and string these together to form particular DARs requires that we have additional distributional descriptions for the length of different CAM sequences found in particular DARs, and matrix probabilities for transitioning among CAMs within such DARs. The outlines of model to accomplish this task, referred to as a multi-CAM metaFuME Markov (M^3) model is described in the next section. In terms of a single CAM metaFuME Markov model embodied in Eqs. 11 and 12, the appropriate number of metaFuME distribution pairs Λ in the set $\mathcal{D}_\Lambda$ (Eq. 10) is, *a priori*, unknown. A first guess at this number might be the number of modes in step length time series derived for different types of CAM segments, if such modes are evident. The problem of finding the best mixture of unimodal pairs $(D_\lambda^{\text{SL}}, D_\lambda^{\text{TA}})$ of distributions that best fit such derived empirical data is a challenging estimation problem that can be approached in several ways, including ensemble Kalman filtering (Dovera & Della Rossa 2011) under the assumption that the underlying distributions are log-concave (Walther et al. 2009). A prudent approach may be to proceed by first looking for the two best-fitting pairs of bivariate distributions:

$$\mathcal{D}_2 = \{D_1(s, a), D_2(s, a)\} \tag{13}$$

and then moving on to three, four, and so on, until no improvement in fit is obtained at the next value for Λ in an information theoretic sense (Symonds & Moussalli 2011). We may then use existing segmentation methods (e.g., BCPA) to identify possible CAMs that may emerge from the simulated

data and compare them to the CAMS identified in the original data If the fit is satisfactory, we may then use our multi-CAM metaFuME Markov (M^3) model (Getz, Vissat & Salter 2019) to predict how individuals may respond to management or global change in terms of local and intermediate scale movements. We note, though, that the M^3 modeling approach described in the next section contains no global directional information. Thus, additional constructs will be needed to obtain movement motivated by headings to specific distant target sites. The performance of M^3-type models and their extensions are investigated elsewhere (Getz, Vissat & Salter 2019).

Construction of M^3 Model

The basis of the M^3 model is to take ensembles of CAM segments of the same type and identify a set of metaFuMEs that can be used to simulate the local structure of such CAMs. If this is done for all the kinds of CAMs that constitute a particular DAR type, then the model can be used to string several types of CAMs together to conform to observed sequences and frequencies of CAMs characteristic of type of DAR under consideration. This approach to constructing M^3 models has been explored and evaluated elsewhere (Getz, Vissat & Salter 2019). Such M^3 models, however, do not account for movement towards specific distant targets. To include such phenomena requires that M^3 models be appropriately extended to include directional biases induced by an attraction to distant geographic locations or a repulsion related to the existence of boundaries to movement (e.g., landscape topography or water).

Extraction of an M^3 model from relocation data requires that the frequency of the data be available at the metaFuME scale, which from Table 1 for medium to large terrestrial vertebrates is around 0.01 to 0.1 Hz. Although details of the approach can be found in (Getz, Vissat & Salter 2019), for the sake of completeness, a brief summary of the approach is enumerated here.

Summary of approach

1. Path relocation data, W^{data} (Eq. 1) is parsed into ensembles of DARs of various types, using an appropriate method (e.g., based on net-square daily displacement or other suitable daily measures; see Bunnefeld et al. 2011, Bischof et al. 2012, Abrahms et al. 2017, Owen-Smith & Martin 2015, Owen-Smith et al. 2010).

2. CAM segments are identified from DAR segment ensembles of the same type (Fig. 4), using appropriate methods such as BCPA (Nams 2014, Edelhoff et al. 2016, Seidel et al. 2018) or HMM (Zucchini et al. 2016). In the latter case, the HMM will need to be performed on a subset of the data subsampled at a scale suitable for identifying CAMs of interest (e.g., if CAMs of interest are assumed to last around 15 minutes or longer then data should be subsampled using a consecutive point intervals of, say, around 5 to 10 minutes).

3. All the data from the ensemble of path segments identified as belonging to the same CAM type within the same DAR type should be strung together into a set of points. If we assume that Φ such CAM-DAR compound types are identified, use W^{ensemble_ϕ}, $\phi = 1, ..., \Phi$ to denote these sets of points.

4. Cluster analyses or other type of categorization procedure should be performed, using the values $\left(s(t), a(t), v^{ss}, v^{aa}, v^{as}\right)$ (Eqs. 2-9) for all times t for which there are points in the

sets W^{ensemble_ϕ} to identify sets of metaFuME types specific to each of the sets W^{ensemble_ϕ}, $\phi = 1, ..., \Phi$ (Fig. 3).

5. For each ensemble type ϕ, distributions $\left(D_\lambda^\phi \right)$, $\lambda_\phi = 1, ..., \Lambda_\phi$, should be fitted to each of the Λ_ϕ metaFuME types identified in the previous step, with some effort to reconcile metaFuME types identified across different ensembles W^{ensemble_ϕ}, $\phi = 1, ..., \Phi$

6. Distributions D_λ^ϕ, $\lambda = 1, ..., \Lambda_\phi$ can be fitted to each of the metaFuME types identified in ensemble ϕ, $\phi = 1, ..., \Phi$, and an associated Markov transmission matrix $\left(\mathbf{P}^\phi \right)$ estimated to obtain the characterizing pairs $\left(\mathcal{D}_{\Lambda_\phi}^\phi, \mathbf{P}^\phi \right)$, which can then be used in a Markov metaFuME movement model to generate CAMs of type ϕ, for $\phi = 1, ..., \Phi$.

7. The metaFuME Markov models derived in the previous step for each type of CAM can be combined into a multi-CAM metaFuME Markov (M^3) model that produces sequences of CAMs with CAM lengths and CAM-type transition statistics that are extracted from the DAR ensemble data.

8. The performance of the M^3 model can be evaluated by comparing the CAMs obtained directly from W^{data}, using existing BCPA and HMM methods, with CAMs produced by the M^3 model simulations.

Conclusion

Without an underlying framework to organize information, deep understanding within any scientific field is impossible. Clearly this has been true of the physical sciences with theories of gravity and the structure of matter being central to our current understanding of the physical universe. This is also true of the life sciences where, for example, the Linnaean taxonomic system, with some modifications, has reigned supreme over the past 250 years (Schuh 2003). This reign still holds today, even within the microbial sciences (Rosselló-Móra & Whitman 2019, Moore et al. 2010) where Mayr's biological species concept fails in several critical ways (Gevers et al. 2005).

In the context of movement ecology, the classification of movement types at different spatio-temporal scales is of considerable interest, although approaches to date have been somewhat informal. At the lifetime track level, basic life-history types regarding, *interalia*, dispersal and migration behavior may be identified. In the movement ecology literature, we see many studies interested in diel activity routine (DARs) (Rahimi & Owen-Smith 2007, Owen-Smith 2013, Owen-Smith & Goodall 2014), lifetime phase (LiMPs) (Fahr et al. 2015, Marra et al. 2015) and over-all life time tracks (LiTTs) (Abrahms et al. 2017). Contrasting DAR types may include distinctions among nomadic, central-place foraging, or territorial behavior, where a single lifetime track may have movement phases dominated by one or other daily activity routine, as in male springbok in Etosha National Park, Namibia, exhibiting territorial behavior during the wet season and nomadic behavior during the dry season, with daily excursions to the same water-hole around midday during the wet season (Lyons et al. 2013). Thus daily activity routines are also very likely to be influenced by seasonal factors, as in pandas that change diel levels of activity in response to seasonal drivers (Zhang et al. 2017).

Figure 4: **The segmentation structure of movement paths at different scales. A.** A multi-season life time track (LiTT) W (Eq. 1) can be segmented into different types of life-history movement phases (LiMPs), each of which can then be segmented into diel activity routines (DARs) and again segmented into canonical activity modes (CAMs). Each CAM is constructed from different proportions of metaFuMEs (aggregates of correlated fundamental movement elements), where each metaFuME is derived from a bivariate step-length/turning-angle distribution derived from time series with specific first-order correlative statistics. Short-duration CAMs (e.g., CAM 3) can easily be missed and averaged into longer duration CAMs (e.g., CAMs 1 and 2), if DAR segmentation analyses is not undertaken at a sufficiently fine scale. **B.** An illustrative example of a simple two-CAM *return-to-home-with-a-water-stop* DAR where the two types of metaFuMEs that generate the *traveling* CAM are shown under magnification.

Extensive effort has also been made to parse diel and longer segments into various types of sub-diel activity modes (Owen-Smith et al. 2010, Donati et al. 2016) that, if stably identifiable across different segments, can be organized into a set of canonical activity modes (CAMs; Getz & Saltz 2008). At finer time scales, however, beyond using accelerometer, acoustic and magnetometer

data to identify behavioral states (Williams et al. 2017, Hurme et al. 2019, Nathan et al. 2012, Fehlmann et al. 2017, Chakravarty et al. 2019, Sapir et al. 2010, Zucchini et al. 2016), very little work has been undertaken to identify sets of fundamental movement elements (FuMEs; same as FMEs in Getz & Saltz 2008) from which CAMs are constructed. As pointed out in this paper, using relocation data alone, we cannot endeavour to identify an underlying set of FuMEs. The best we can hope for, provided subminute relocation data is available, is the identification of a set of metaFuMEs consisting of repeated or correlated strings of FuMES, and characterized by particular step-length and turning-angle distributions.

We should not underestimate the analytical and computational challenge required to extract a comprehensive and stable (across many segments at various scales) set of metaFuME elements. The best methods to do this still remain to be developed and how well this may be accomplished remains to be seen; although satisfying progress has already been made (Getz, Vissat & Salter 2019). Future progress may require many of the ideas presented here to be refined or modified. There is not denying, however, that formalized, widely accepted, hierarchical-path-segmentation framework will greatly facilitate efforts to address outstanding questions in movement ecology, particularly those involving comparative analyses within species (Wittemyer et al. 2019), as well as across species. Within species variation may allow us to assess differences along geographic clines with application to the behavioral adaptation of species under landscape and climate change (Seebacher & Post 2015). It may help us assess fitness in the context of feeding strategies, social behavior (Harel et al. 2017, Harel, Duriez, Spiegel, Fluhr, Horvitz, Getz, Bouten, Sarrazin, Hatzofe & Nathan 2016), areas attractive to populations (Giotto et al. 2015), or mapping out landscapes in terms of their overall resistance to movement (Zeller et al. 2012). It may also be diagnostic of changes in movement behavior when individuals are stressed, ill, have genetic defects, or females are pregnant (Spink et al. 2001, Owen-Smith 2013); or it may be predictive in terms of pathogen transmission and the spread of disease (Cross et al. 2007, Tracey et al. 2014, Dougherty et al. 2018, Zidon et al. 2017). Differences in various species' movement profiles at various scales may be critical when it comes to taking a multispecies approach to assessing the impacts of ecosystem management on species conservation (Brodie et al. 2015, Runge et al. 2014).

Additionally, a formalized, hierarchical-path-segmentation framework of appropriate complexity (Getz et al. 2018, Larsen et al. 2016) can be used, for example, to address the dozen plus questions that were recently posed regarding the movement ecology of marine megafauna (Hays et al. 2016)— but are equally applicable to all animal species. These questions include: 1) "Are there simple rules underlying seemingly complex movement patterns and hence common drivers for movement across species?" and 2.) "How will climate change impact animal movements?" An ability to address both of these question, as well as those raised above and others besides, in a comparative way—with a level of consistency and coherency that only a universally accepted classification framework can provide—is needed with great urgency, as the field of movement ecology matures. The framework presented here is a starting point to developing a more comprehensive one that can be fleshed out in the future, supported by analytical and computational methods, some of which still need to be developed or refined from existing methods. The framework proposed here, however, provides a guide to constructing M^3 models for simulating CAM and DAR segments in empirical relocation data (Vissat et al. 2019), with extensions yet to be developed, will ultimately provide a model for simulating walks that conform to empirical data at several local (metaFuME, CAM) and global scales (biased CAM components in DARs and beyond). It will take models of this complexity to address the two questions posed here, as well as those posed elsewhere in the context of theoretical

and applied movement ecology, the latter in particular addressing questions related to conservation   490
biology, resource management, and global change assessment.   491

Glossary   492

An expansion of the various acronyms used in the text are provided here in alphabetical order along   493
with a very brief definition/explanation of their meaning.   494

BCPA: Behavioral Change Point Analysis. Behavioral change-point analysis refers to a group   495
of methods used to determine how the statistics of a biological variable y (e.g., its mean,   496
variance, autocorrelation, or rate of change in slope or curvature), dependent on a second   497
variable x, switches at threshold points with changes in x (e.g., space, time, or the abscissa   498
in a stress-response relationship) (Morales et al. 2004, Andersen et al. 2009, Chen & Gupta   499
2011, Jonsen et al. 2007). Note, more generally, x could be vector-valued.   500

CAM: Canonical Activity Mode. This is a stably classifiable subdiel behavioral mode (pattern of   501
movement) such as a foraging bout, resting period, or purposeful heading (i.e., traveling) to   502
a distant target location.   503

DAR: Diel Activity Routine. This is a stably classifiable 24-hour sequence of CAMs that occur   504
at characteristic frequencies and times of the day.   505

FuME: Fundamental Movement Elements (also FME; Getz & Saltz 2008). This is a relatively   506
rapid, highly repeatable, stereotypic set of body movements that forms the basis of the   507
locomotory capacity of an individual (e.g., a walk step, a running step, a wing flap, a jump,   508
etc.)   509

HMM: Hidden Markov Model. A time series of observable values that depend in a probabilistic   510
sense on the values of an associated, but unobservable Markov chain process (a set of states   511
where the transition from one state to another depends only on the value of the current state)   512
(Zucchini et al. 2016)   513

LiMP: Life-history Movement Phase. This is a path segment that typically reflects a life-history   514
relevant movement behavior such as dispersal (episodic), migration (periodic), or other peri-   515
odic behaviors at a greater-than-diel scale.   516

LiTT: Life Time Track. This is the total movement path of an individual from its birth to its   517
death.   518

M^3: Multi-CAM metaFuME Markov (model). A model based on the metaFuME set $\mathcal{D}_\Lambda$ of CAM-   519
specific distributions (Eq. 10) and CAM-specific Markov transition matrices P_λ, $\lambda = 1, ..., \Lambda$,   520
derived from the set $\mathcal{T}_\Lambda$ of metaFuME sequence index set ensembles (see Fig. 3). This model   521
contains only local movement information and cannot be used to simulate movement patterns   522
motivated by headings to distant targets.   523

MetaFuME: A correlated stereotypical or characteristic sequence of FuMEs of fixed duration   524
equal to the time between consecutive relocation points (only applicable to relatively high   525
frequency data: ideally, metaFuMEs should contain no more than several tens of FuMES).   526

PS: Path Segmentation. This is the process of breaking up a movement path into reliably stable metaFuMEs, CAMs, DARs, and LiMPs using a suite of methods that include BCPA and HMM approaches (Nams 2014, Edelhoff et al. 2016, Seidel et al. 2018).

SL: Step Length. The distance between consecutive relocation points, as generated from Eq. 1 using Eq. 3.

TA: Turning Angle. The change in the angle of heading across three consecutive relocation points, as generated from Eq. 1 using Eq. 5.

Acknowledgements

I thank Briana Abrahms, Shirli Bar-David, Paul Cross, Victoria Goodall, Nir Horvitz, Leif Loe, Ran Nathan, Norman Owen-Smith, Leo Polansky, David Saltz, Dana Seidel, Orr Spiegel, Krti Tallam, Miriam Tsalyuk, George Wittemyer for comments on earlier versions of this paper. All inaccuracies and misconceptions are my own.

References

Abrahms, B., Seidel, D. P., Dougherty, E., Hazen, E. L., Bograd, S. J., Wilson, A. M., McNutt, J. W., Costa, D. P., Blake, S., Brashares, J. S. et al. (2017), 'Suite of simple metrics reveals common movement syndromes across vertebrate taxa', *Movement ecology* **5**(1), 12.

Allen, R. M., Metaxas, A. & Snelgrove, P. V. (2018), 'Applying movement ecology to marine animals with complex life cycles', *Annual Review of Marine Science* **10**, 19–42.

Andersen, T., Carstensen, J., Hernandez-Garcia, E. & Duarte, C. M. (2009), 'Ecological thresholds and regime shifts: approaches to identification', *Trends in Ecology & Evolution* **24**(1), 49–57.

Avgar, T., Potts, J. R., Lewis, M. A. & Boyce, M. S. (2016), 'Integrated step selection analysis: bridging the gap between resource selection and animal movement', *Methods in Ecology and Evolution* **7**(5), 619–630.

Bar-David, S., Bar-David, I., Cross, P. C., Ryan, S. J., Knechtel, C. U. & Getz, W. M. (2009), 'Methods for assessing movement path recursion with application to African buffalo in South Africa', *Ecology* **90**(9), 2467–2479.

Bartumeus, F., da Luz, M. G. E., Viswanathan, G. M. & Catalan, J. (2005), 'Animal search strategies: a quantitative random-walk analysis', *Ecology* **86**(11), 3078–3087.

Bennison, A., Bearhop, S., Bodey, T. W., Votier, S. C., Grecian, W. J., Wakefield, E. D., Hamer, K. C. & Jessopp, M. (2018), 'Search and foraging behaviors from movement data: a comparison of methods', *Ecology and evolution* **8**(1), 13–24.

Berger-Tal, O. & Bar-David, S. (2015), 'Recursive movement patterns: review and synthesis across species', *Ecosphere* **6**(9), 1–12.

Bergman, C. M., Schaefer, J. A. & Luttich, S. (2000), 'Caribou movement as a correlated random walk', *Oecologia* **123**(3), 364–374.

Bischof, R., Loe, L. E., Meisingset, E. L., Zimmermann, B., Van Moorter, B. & Mysterud, A. (2012), 'A migratory northern ungulate in the pursuit of spring: jumping or surfing the green wave?', *The American Naturalist* **180**(4), 407–424.

Bork, E. W. & Su, J. G. (2007), 'Integrating lidar data and multispectral imagery for enhanced classification of rangeland vegetation: A meta analysis', *Remote Sensing of Environment* **111**(1), 11–24.

Bowler, D. E. & Benton, T. G. (2005), 'Causes and consequences of animal dispersal strategies: relating individual behaviour to spatial dynamics', *Biological Reviews* **80**(2), 205–225.

Boyce, M. S., Vernier, P. R., Nielsen, S. E. & Schmiegelow, F. K. (2002), 'Evaluating resource selection functions', *Ecological modelling* **157**(2-3), 281–300.

Brodie, J. F., Giordano, A. J., Dickson, B., Hebblewhite, M., Bernard, H., Mohd-Azlan, J., Anderson, J. & Ambu, L. (2015), 'Evaluating multispecies landscape connectivity in a threatened tropical mammal community', *Conservation Biology* **29**(1), 122–132.

Bunnefeld, N., Börger, L., van Moorter, B., Rolandsen, C. M., Dettki, H., Solberg, E. J. & Ericsson, G. (2011), 'A model-driven approach to quantify migration patterns: individual, regional and yearly differences', *Journal of Animal Ecology* **80**(2), 466–476.

Byers, J. A. (2001), 'Correlated random walk equations of animal dispersal resolved by simulation', *Ecology* **82**(6), 1680–1690.

Calabrese, J. M., Fleming, C. H. & Gurarie, E. (2016), 'ctmm: an r package for analyzing animal relocation data as a continuous-time stochastic process', *Methods in Ecology and Evolution* **7**(9), 1124–1132.

Cattarino, L., McAlpine, C. A. & Rhodes, J. R. (2016), 'Spatial scale and movement behaviour traits control the impacts of habitat fragmentation on individual fitness', *Journal of Animal Ecology* **85**(1), 168–177.

Chakravarty, P., Maalberg, M., Cozzi, G., Ozgul, A. & Aminian, K. (2019), 'Behavioural compass: animal behaviour recognition using magnetometers', *Movement Ecology* **7**(1), 28.

Chatfield, C. (2016), *The analysis of time series: an introduction*, Chapman and Hall/CRC.

Chen, J. & Gupta, A. K. (2011), *Parametric statistical change point analysis: with applications to genetics, medicine, and finance*, Springer Science & Business Media.

Coello, C. A. C. (2003), Evolutionary multi-objective optimization: A critical review, *in* 'Evolutionary optimization', Springer, pp. 117–146.

Cross, P. C., Almberg, E. S., Haase, C. G., Hudson, P. J., Maloney, S. K., Metz, M. C., Munn, A. J., Nugent, P., Putzeys, O., Stahler, D. R. et al. (2016), 'Energetic costs of mange in wolves estimated from infrared thermography', *Ecology* **97**(8), 1938–1948.

Cross, P. C., Edwards, W. H., Scurlock, B. M., Maichak, E. J. & Rogerson, J. D. (2007), 'Effects of management and climate on elk brucellosis in the greater yellowstone ecosystem', *Ecological Applications* **17**(4), 957–964.

de Weerd, N., van Langevelde, F., van Oeveren, H., Nolet, B. A., Kölzsch, A., Prins, H. H. & de Boer, W. F. (2015), 'Deriving animal behaviour from high-frequency gps: tracking cows in open and forested habitat', *Plos one* **10**(6), e0129030.

DeCesare, N. J., Hebblewhite, M., Schmiegelow, F., Hervieux, D., McDermid, G. J., Neufeld, L., Bradley, M., Whittington, J., Smith, K. G., Morgantini, L. E. et al. (2012), 'Transcending scale dependence in identifying habitat with resource selection functions', *Ecological Applications* **22**(4), 1068–1083.

Delcourt, J., Denoël, M., Ylieff, M. & Poncin, P. (2013), 'Video multitracking of fish behaviour: a synthesis and future perspectives', *Fish and Fisheries* **14**(2), 186–204.

Delp, S. L. & Loan, J. P. (2000), 'A computational framework for simulating and analyzing human and animal movement', *Computing in Science & Engineering* **2**(5), 46–55.

Donati, G., Campera, M., Balestri, M., Serra, V., Barresi, M., Schwitzer, C., Curtis, D. J. & Santini, L. (2016), 'Ecological and anthropogenic correlates of activity patterns in eulemur', *International Journal of Primatology* **37**(1), 29–46.

Dougherty, E. R., Seidel, D. P., Carlson, C. J., Spiegel, O. & Getz, W. M. (2018), 'Going through the motions: incorporating movement analyses into disease research', *Ecology letters* **21**(4), 588–604.

Dovera, L. & Della Rossa, E. (2011), 'Multimodal ensemble kalman filtering using gaussian mixture models', *Computational Geosciences* **15**(2), 307–323.

Edelhoff, H., Signer, J. & Balkenhol, N. (2016), 'Path segmentation for beginners: an overview of current methods for detecting changes in animal movement patterns', *Movement ecology* **4**(1), 21.

Fahr, J., Abedi-Lartey, M., Esch, T., Machwitz, M., Suu-Ire, R., Wikelski, M. & Dechmann, D. K. (2015), 'Pronounced seasonal changes in the movement ecology of a highly gregarious central-place forager, the african straw-coloured fruit bat (eidolon helvum)', *PloS one* **10**(10), e0138985.

Fauchald, P. & Tveraa, T. (2003), 'Using first-passage time in the analysis of area-restricted search and habitat selection', *Ecology* **84**(2), 282–288.

Fehlmann, G., ORiain, M. J., Hopkins, P. W., OSullivan, J., Holton, M. D., Shepard, E. L. & King, A. J. (2017), 'Identification of behaviours from accelerometer data in a wild social primate', *Animal Biotelemetry* **5**(1), 6.

Fleming, C. H. & Calabrese, J. M. (2017), 'A new kernel density estimator for accurate home-range and species-range area estimation', *Methods in Ecology and Evolution* **8**(5), 571–579.

Fleming, C. H., Calabrese, J. M., Mueller, T., Olson, K. A., Leimgruber, P. & Fagan, W. F. (2014), 'From fine-scale foraging to home ranges: a semivariance approach to identifying movement modes across spatiotemporal scales', *The American Naturalist* **183**(5), E154–E167.

Fortin, D., Beyer, H. L., Boyce, M. S., Smith, D. W., Duchesne, T. & Mao, J. S. (2005), 'Wolves influence elk movements: behavior shapes a trophic cascade in yellowstone national park', *Ecology* **86**(5), 1320–1330.

Fortin, D., Boyce, M. S., Merrill, E. H. & Fryxell, J. M. (2004), 'Foraging costs of vigilance in large mammalian herbivores', *Oikos* **107**(1), 172–180.

Franke, A., Caelli, T. & Hudson, R. J. (2004), 'Analysis of movements and behavior of caribou (rangifer tarandus) using hidden markov models', *Ecological Modelling* **173**(2-3), 259–270.

Getz, W. M., Marshall, C. R., Carlson, C. J., Giuggioli, L., Ryan, S. J., Romañach, S. S., Boettiger, C., Chamberlain, S. D., Larsen, L., DOdorico, P. et al. (2018), 'Making ecological models adequate', *Ecology letters* **21**(2), 153–166.

Getz, W. M., Salter, R. & Tallam, K. (2019), 'A quantitative narrative on movement, disease and patch exploitation in nesting agent groups', *bioRxiv* .
URL: *https://www.biorxiv.org/content/early/2019/10/03/791400*

Getz, W. M. & Saltz, D. (2008), 'A framework for generating and analyzing movement paths on ecological landscapes', *Proceedings of the National Academy of Sciences* **105**(49), 19066–19071.

Getz, W., Vissat, L. L. & Salter, R. (2019), 'Simulation and analysis of animal movement paths using numerus model builder', *bioRxiv* .
   URL: *https://www.biorxiv.org/content/early/2019/12/15/2019.12.15.876987*

Gevers, D., Cohan, F. M., Lawrence, J. G., Spratt, B. G., Coenye, T., Feil, E. J., Stackebrandt, E., Van de Peer, Y., Vandamme, P., Thompson, F. L. et al. (2005), 'Re-evaluating prokaryotic species', *Nature Reviews Microbiology* **3**(9), 733.

Giotto, N., Gerard, J.-F., Ziv, A., Bouskila, A. & Bar-David, S. (2015), 'Space-use patterns of the asiatic wild ass (*equus hemionus*): complementary insights from displacement, recursion movement and habitat selection analyses', *PloS one* **10**(12), e0143279.

Goodall, V., Fatti, L. & Owen-Smith, N. (2017), 'Animal movement modelling: independent or dependent models?', *South African Statistical Journal* **51**(2), 295–315.

Grémillet, D. & Boulinier, T. (2009), 'Spatial ecology and conservation of seabirds facing global climate change: a review', *Marine Ecology Progress Series* **391**, 121–137.

Gurarie, E., Andrews, R. D. & Laidre, K. L. (2009), 'A novel method for identifying behavioural changes in animal movement data', *Ecology letters* **12**(5), 395–408.

Gurarie, E., Bracis, C., Delgado, M., Meckley, T. D., Kojola, I. & Wagner, C. M. (2016), 'What is the animal doing? tools for exploring behavioural structure in animal movements', *Journal of Animal Ecology* **85**(1), 69–84.

Gurarie, E., Fleming, C. H., Fagan, W. F., Laidre, K. L., Hernández-Pliego, J. & Ovaskainen, O. (2017), 'Correlated velocity models as a fundamental unit of animal movement: synthesis and applications', *Movement ecology* **5**(1), 13.

Hardin, P. E. & Panda, S. (2013), 'Circadian timekeeping and output mechanisms in animals', *Current opinion in neurobiology* **23**(5), 724–731.

Harel, R., Duriez, O., Spiegel, O., Fluhr, J., Horvitz, N., Getz, W. M., Bouten, W., Sarrazin, F., Hatzofe, O. & Nathan, R. (2016), 'Decision-making by a soaring bird: time, energy and risk considerations at different spatio-temporal scales', *Philosophical Transactions of the Royal Society B: Biological Sciences* **371**(1704), 20150397.

Harel, R., Horvitz, N. & Nathan, R. (2016), 'Adult vultures outperform juveniles in challenging thermal soaring conditions', *Scientific reports* **6**, 27865.

Harel, R. & Nathan, R. (2018), 'The characteristic time-scale of perceived information for decision-making: Departure from thermal columns in soaring birds', *Functional ecology* **32**(8), 2065–2072.

Harel, R., Spiegel, O., Getz, W. M. & Nathan, R. (2017), 'Social foraging and individual consistency in following behaviour: testing the information centre hypothesis in free-ranging vultures', *Proceedings of the Royal Society B: Biological Sciences* **284**(1852), 20162654.

Hays, G. C., Ferreira, L. C., Sequeira, A. M., Meekan, M. G., Duarte, C. M., Bailey, H., Bailleul, F., Bowen, W. D., Caley, M. J., Costa, D. P. et al. (2016), 'Key questions in marine megafauna movement ecology', *Trends in ecology & evolution* **31**(6), 463–475.

Hein, A. M., Carrara, F., Brumley, D. R., Stocker, R. & Levin, S. A. (2016), 'Natural search algorithms as a bridge between organisms, evolution, and ecology', *Proceedings of the National Academy of Sciences* **113**(34), 9413–9420.

Hinton, G., Deng, L., Yu, D., Dahl, G. E., Mohamed, A.-r., Jaitly, N., Senior, A., Vanhoucke, V., Nguyen, P., Sainath, T. N. et al. (2012), 'Deep neural networks for acoustic modeling in speech recognition: The shared views of four research groups', *IEEE Signal processing magazine* **29**(6), 82–97.

Hurme, E., Gurarie, E., Greif, S., Flores-Martínez, J. J., Wilkinson, G. S., Yovel, Y. et al. (2019), 'Acoustic evaluation of behavioral states predicted from gps tracking: a case study of a marine fishing bat', *Movement ecology* **7**(1), 21.

Johnson, C. J., Parker, K. L., Heard, D. C. & Gillingham, M. P. (2002), 'A multiscale behavioral approach to understanding the movements of woodland caribou', *Ecological Applications* **12**(6), 1840–1860.

Johnson, D. S., London, J. M., Lea, M.-A. & Durban, J. W. (2008), 'Continuous-time correlated random walk model for animal telemetry data', *Ecology* **89**(5), 1208–1215.

Jonsen, I. D., Myers, R. A. & Flemming, J. M. (2003), 'Meta-analysis of animal movement using state-space models', *Ecology* **84**(11), 3055–3063.

Jonsen, I. D., Myers, R. A. & James, M. C. (2006), 'Robust hierarchical state–space models reveal diel variation in travel rates of migrating leatherback turtles', *Journal of Animal Ecology* **75**(5), 1046–1057.

Jonsen, I. D., Myers, R. A. & James, M. C. (2007), 'Identifying leatherback turtle foraging behaviour from satellite telemetry using a switching state-space model', *Marine Ecology Progress Series* **337**, 255–264.

Kane, A. S., Salierno, J. D., Gipson, G. T., Molteno, T. C. & Hunter, C. (2004), 'A video-based movement analysis system to quantify behavioral stress responses of fish', *Water Research* **38**(18), 3993–4001.

Kareiva, P. & Shigesada, N. (1983), 'Analyzing insect movement as a correlated random walk', *Oecologia* **56**(2-3), 234–238.

Killick, R. & Eckley, I. (2014), 'changepoint: An r package for changepoint analysis', *Journal of statistical software* **58**(3), 1–19.

Langrock, R., King, R., Matthiopoulos, J., Thomas, L., Fortin, D. & Morales, J. M. (2012), 'Flexible and practical modeling of animal telemetry data: hidden markov models and extensions', *Ecology* **93**(11), 2336–2342.

Larsen, L. G., Eppinga, M. B., Passalacqua, P., Getz, W. M., Rose, K. A. & Liang, M. (2016), 'Appropriate complexity landscape modeling', *Earth-science reviews* **160**, 111–130.

Levin, S. A. (1992), 'The problem of pattern and scale in ecology: the robert h. macarthur award lecture', *Ecology* **73**(6), 1943–1967.

Lyons, A. J., Turner, W. C. & Getz, W. M. (2013), 'Home range plus: a space-time characterization of movement over real landscapes', *Movement Ecology* **1**(1), 2.

Marin, J.-M., Pudlo, P., Robert, C. P. & Ryder, R. J. (2012), 'Approximate bayesian computational methods', *Statistics and Computing* **22**(6), 1167–1180.

Marra, P. P., Cohen, E. B., Loss, S. R., Rutter, J. E. & Tonra, C. M. (2015), 'A call for full annual cycle research in animal ecology', *Biology letters* **11**(8), 20150552.

Matteson, D. S. & James, N. A. (2014), 'A nonparametric approach for multiple change point analysis of multivariate data', *Journal of the American Statistical Association* **109**(505), 334–345.

McCulloch, C. & Cain, M. (1989), 'Analyzing discrete movement data as a correlated random walk', *Ecology* **70**(2), 383–388.

McGarigal, K., Wan, H. Y., Zeller, K. A., Timm, B. C. & Cushman, S. A. (2016), 'Multi-scale habitat selection modeling: a review and outlook', *Landscape ecology* **31**(6), 1161–1175.

McGavin, S. L., Bishop-Hurley, G. J., Charmley, E., Greenwood, P. L. & Callaghan, M. J. (2018), 'Effect of gps sample interval and paddock size on estimates of distance travelled by grazing cattle in rangeland, australia', *The Rangeland Journal* **40**(1), 55–64.

McKenzie, H. W., Lewis, M. A. & Merrill, E. H. (2009), 'First passage time analysis of animal movement and insights into the functional response', *Bulletin of mathematical biology* **71**(1), 107–129.

Michelot, T., Langrock, R. & Patterson, T. A. (2016), 'movehmm: An r package for the statistical modelling of animal movement data using hidden markov models', *Methods in Ecology and Evolution* **7**(11), 1308–1315.

Milner-Gulland, E., Fryxell, J. M. & Sinclair, A. R. (2011), *Animal migration: a synthesis*, Oxford University Press.

Moore, E. R., Mihaylova, S. A., Vandamme, P., Krichevsky, M. I. & Dijkshoorn, L. (2010), 'Microbial systematics and taxonomy: relevance for a microbial commons', *Research in microbiology* **161**(6), 430–438.

Morales, J. M., Haydon, D. T., Frair, J., Holsinger, K. E. & Fryxell, J. M. (2004), 'Extracting more out of relocation data: building movement models as mixtures of random walks', *Ecology* **85**(9), 2436–2445.

Morelle, K., Podgórski, T., Prévot, C., Keuling, O., Lehaire, F. & Lejeune, P. (2015), 'Towards understanding wild boar *sus scrofa* movement: a synthetic movement ecology approach', *Mammal Review* **45**(1), 15–29.

Morellet, N., Bonenfant, C., Börger, L., Ossi, F., Cagnacci, F., Heurich, M., Kjellander, P., Linnell, J. D., Nicoloso, S., Sustr, P. et al. (2013), 'Seasonality, weather and climate affect home range size in roe deer across a wide latitudinal gradient within e urope', *Journal of Animal Ecology* **82**(6), 1326–1339.

Mueller, T. & Fagan, W. F. (2008), 'Search and navigation in dynamic environments–from individual behaviors to population distributions', *Oikos* **117**(5), 654–664.

Myers, J. H. (2018), 'Population cycles: generalities, exceptions and remaining mysteries', *Proceedings of the Royal Society B: Biological Sciences* **285**(1875), 20172841.

Mysterud, A., Stenseth, N. C., Yoccoz, N. G., Langvatn, R. & Steinheim, G. (2001), 'Nonlinear effects of large-scale climatic variability on wild and domestic herbivores', *Nature* **410**(6832), 1096.

Nams, V. O. (2014), 'Combining animal movements and behavioural data to detect behavioural states', *Ecology letters* **17**(10), 1228–1237.

Nathan, R., Getz, W. M., Revilla, E., Holyoak, M., Kadmon, R., Saltz, D. & Smouse, P. E. (2008), 'A movement ecology paradigm for unifying organismal movement research', *Proceedings of the National Academy of Sciences* **105**(49), 19052–19059.

Nathan, R., Spiegel, O., Fortmann-Roe, S., Harel, R., Wikelski, M. & Getz, W. M. (2012), 'Using triaxial acceleration data to identify behavioral modes of free-ranging animals: general concepts and tools illustrated for griffon vultures', *Journal of Experimental Biology* **215**(6), 986–996.

Northrup, J. M., Avrin, A., Anderson, C. R., Brown, E. & Wittemyer, G. (2019), 'On-animal acoustic monitoring provides insight to ungulate foraging behavior', *Journal of Mammalogy* .

Odu, G. & Charles-Owaba, O. (2013), 'Review of multi-criteria optimization methods–theory and applications', *IOSR Journal of Engineering (IOSRJEN)* **3**(10), 1–14.

Owen-Smith, N. (2013), 'Daily movement responses by african savanna ungulates as an indicator of seasonal and annual food stress', *Wildlife Research* **40**(3), 232–240.

Owen-Smith, N., Fryxell, J. & Merrill, E. (2010), 'Foraging theory upscaled: the behavioural ecology of herbivore movement', *Philosophical Transactions of the Royal Society B: Biological Sciences* **365**(1550), 2267–2278.

Owen-Smith, N. & Goodall, V. (2014), 'Coping with savanna seasonality: comparative daily activity patterns of a frican ungulates as revealed by gps telemetry', *Journal of Zoology* **293**(3), 181–191.

Owen-Smith, N. & Goodall, V. (2019), 'Movement informatics of large mammalian herbivores across hierarchical scales: inferring biology from statistics', *Under review* .

Owen-Smith, N. & Martin, J. (2015), 'Identifying space use at foraging arena scale within the home ranges of large herbivores', *PloS one* **10**(6), e0128821.

Panzacchi, M., Van Moorter, B., Strand, O., Saerens, M., Kivimäki, I., St. Clair, C. C., Herfindal, I. & Boitani, L. (2016), 'Predicting the continuum between corridors and barriers to animal movements using step selection functions and randomized shortest paths', *Journal of Animal Ecology* **85**(1), 32–42.

Patterson, T. A., Thomas, L., Wilcox, C., Ovaskainen, O. & Matthiopoulos, J. (2008), 'State–space models of individual animal movement', *Trends in ecology & evolution* **23**(2), 87–94.

Pettorelli, N., Laurance, W. F., O'Brien, T. G., Wegmann, M., Nagendra, H. & Turner, W. (2014), 'Satellite remote sensing for applied ecologists: opportunities and challenges', *Journal of Applied Ecology* **51**(4), 839–848.

Pettorelli, N., Ryan, S., Mueller, T., Bunnefeld, N., Jedrzejewska, B., Lima, M. & Kausrud, K. (2011), 'The normalized difference vegetation index (ndvi): unforeseen successes in animal ecology', *Climate Research* **46**(1), 15–27.

Pohle, J., Langrock, R., van Beest, F. M. & Schmidt, N. M. (2017), 'Selecting the number of states in hidden markov models: pragmatic solutions illustrated using animal movement', *Journal of Agricultural, Biological and Environmental Statistics* **22**(3), 270–293.

Polansky, L., Kilian, W. & Wittemyer, G. (2015), 'Elucidating the significance of spatial memory on movement decisions by african savannah elephants using state–space models', *Proceedings of the Royal Society B: Biological Sciences* **282**(1805), 20143042.

Polansky, L., Wittemyer, G., Cross, P. C., Tambling, C. J. & Getz, W. M. (2010), 'From moonlight to movement and synchronized randomness: Fourier and wavelet analyses of animal location time series data', *Ecology* **91**(5), 1506–1518.

Preisler, H. K., Ager, A. A., Johnson, B. K. & Kie, J. G. (2004), 'Modeling animal movements using stochastic differential equations', *Environmetrics* **15**(7), 643–657.

Rahimi, S. & Owen-Smith, N. (2007), 'Movement patterns of sable antelope in the kruger national park from gps/gsm collars: a preliminary assessment', *African Journal of Wildlife Research* **37**(2), 143–152.

Rosselló-Móra, R. & Whitman, W. B. (2019), 'Dialogue on the nomenclature and classification of prokaryotes', *Systematic and applied microbiology* **42**(1), 5–14.

Runge, C. A., Martin, T. G., Possingham, H. P., Willis, S. G. & Fuller, R. A. (2014), 'Conserving mobile species', *Frontiers in Ecology and the Environment* **12**(7), 395–402.

Sapir, N., Wikelski, M., McCue, M. D., Pinshow, B. & Nathan, R. (2010), 'Flight modes in migrating european bee-eaters: heart rate may indicate low metabolic rate during soaring and gliding', *PLoS One* **5**(11), e13956.

Schuh, R. T. (2003), 'The linnaean system and its 250-year persistence', *The Botanical Review* **69**(1), 59.

Seebacher, F. & Post, E. (2015), 'Climate change impacts on animal migration', *Climate Change Responses* **2**(1), 5.

Seidel, D. P., Dougherty, E., Carlson, C. & Getz, W. M. (2018), 'Ecological metrics and methods for gps movement data', *International Journal of Geographical Information Science* **32**(11), 2272–2293.

Shepard, E. L., Wilson, R. P., Rees, W. G., Grundy, E., Lambertucci, S. A. & Vosper, S. B. (2013), 'Energy landscapes shape animal movement ecology', *The American Naturalist* **182**(3), 298–312.

Siniff, D. B. & Jessen, C. (1969), A simulation model of animal movement patterns, *in* 'Advances in ecological research', Vol. 6, Elsevier, pp. 185–219.

Spiegel, O., Harel, R., Centeno-Cuadros, A., Hatzofe, O., Getz, W. M. & Nathan, R. (2015), 'Moving beyond curve fitting: using complementary data to assess alternative explanations for long movements of three vulture species', *The American Naturalist* **185**(2), E44–E54.

Spiegel, O. & O'Farrell, S. (2019), 'Spatial orientation and time: Methods', *Encyclopedia of Animal Behavior* pp. 518–528.

Spink, A., Tegelenbosch, R., Buma, M. & Noldus, L. (2001), 'The ethovision video tracking systema tool for behavioral phenotyping of transgenic mice', *Physiology & behavior* **73**(5), 731–744.

Symonds, M. R. & Moussalli, A. (2011), 'A brief guide to model selection, multimodel inference and model averaging in behavioural ecology using akaikes information criterion', *Behavioral Ecology and Sociobiology* **65**(1), 13–21.

Takahashi, J. S., Hong, H.-K., Ko, C. H. & McDearmon, E. L. (2008), 'The genetics of mammalian circadian order and disorder: implications for physiology and disease', *Nature reviews genetics* **9**(10), 764.

Thessen, A. (2016), 'Adoption of machine learning techniques in ecology and earth science', *One Ecosystem* **1**, e8621.

Thurfjell, H., Ciuti, S. & Boyce, M. S. (2014), 'Applications of step-selection functions in ecology and conservation', *Movement ecology* **2**(1), 4.

Toledo, S., Kishon, O., Orchan, Y., Shohat, A. & Nathan, R. (2016), Lessons and experiences from the design, implementation, and deployment of a wildlife tracking system, *in* 'Software Science, Technology and Engineering (SWSTE), 2016 IEEE International Conference on', IEEE, pp. 51–60.

Torney, C. J., Hopcraft, J. G. C., Morrison, T. A., Couzin, I. D. & Levin, S. A. (2018), 'From single steps to mass migration: the problem of scale in the movement ecology of the serengeti wildebeest', *Philosophical Transactions of the Royal Society B: Biological Sciences* **373**(1746), 20170012.

Torres, L. G., Orben, R. A., Tolkova, I. & Thompson, D. R. (2017), 'Classification of animal movement behavior through residence in space and time', *PloS one* **12**(1), e0168513.

Townsend Peterson, A., Papeş, M. & Eaton, M. (2007), 'Transferability and model evaluation in ecological niche modeling: a comparison of garp and maxent', *Ecography* **30**(4), 550–560.

Tracey, J. A., Bevins, S. N., VandeWoude, S. & Crooks, K. R. (2014), 'An agent-based movement model to assess the impact of landscape fragmentation on disease transmission', *Ecosphere* **5**(9), 1–24.

Tsalyuk, M., Kilian, W., Reineking, B. & Getz, W. M. (2019), 'Temporal variation in resource selection of african elephants follows long-term variability in resource availability', *Ecological Monographs* **89**(2), e01348.

Turchin, P. (1998), *Quantitative analysis of movement*, Sinauer assoc. Sunderland (mass.).

Valletta, J. J., Torney, C., Kings, M., Thornton, A. & Madden, J. (2017), 'Applications of machine learning in animal behaviour studies', *Animal Behaviour* **124**, 203–220.

Van Moorter, B., Visscher, D. R., Jerde, C. L., Frair, J. L. & Merrill, E. H. (2010), 'Identifying movement states from location data using cluster analysis', *The Journal of Wildlife Management* **74**(3), 588–594.

Vissat, L. L., Salter, R. & Getz, W. M. (2019), 'Simulation and analysis of animal movement paths using numerus model builder', *bioRxiv* .
    URL: *https://www.biorxiv.org/content/early/2019/xxxx*

Walther, G. et al. (2009), 'Inference and modeling with log-concave distributions', *Statistical Science* **24**(3), 319–327.

Wang, Y., Nickel, B., Rutishauser, M., Bryce, C. M., Williams, T. M., Elkaim, G. & Wilmers, C. C. (2015), 'Movement, resting, and attack behaviors of wild pumas are revealed by tri-axial accelerometer measurements', *Movement Ecology* **3**(1), 2.

Weiser, A. W., Orchan, Y., Nathan, R., Charter, M., Weiss, A. J. & Toledo, S. (2016), Characterizing the accuracy of a self-synchronized reverse-gps wildlife localization system, *in* 'Information Processing in Sensor Networks (IPSN), 2016 15th ACM/IEEE International Conference on', IEEE, pp. 1–12.

Williams, H. J., Holton, M. D., Shepard, E. L., Largey, N., Norman, B., Ryan, P. G., Duriez, O., Scantlebury, M., Quintana, F., Magowan, E. A. et al. (2017), 'Identification of animal movement patterns using tri-axial magnetometry', *Movement ecology* **5**(1), 6.

Williams, H. J., Taylor, L. A., Benhamou, S., Bijleveld, A. I., Clay, T. A., de Grissac, S., Demšar, U., English, H. M., Franconi, N., Gómez-Laich, A. et al. (2019), 'Optimising the use of bio-loggers for movement ecology research', *Journal of Animal Ecology* .

Wilmers, C. C., Nickel, B., Bryce, C. M., Smith, J. A., Wheat, R. E. & Yovovich, V. (2015), 'The golden age of bio-logging: how animal-borne sensors are advancing the frontiers of ecology', *Ecology* **96**(7), 1741–1753.

Wittemyer, G., Northrup, J. M. & Bastille-Rousseau, G. (2019), 'Behavioural valuation of landscapes using movement data', *Philosophical Transactions of the Royal Society B* **374**(1781), 20180046.

Wittemyer, G., Polansky, L., Douglas-Hamilton, I. & Getz, W. M. (2008), 'Disentangling the effects of forage, social rank, and risk on movement autocorrelation of elephants using fourier and wavelet analyses', *Proceedings of the National Academy of Sciences* pp. pnas–0801744105.

Worton, B. (1987), 'A review of models of home range for animal movement', *Ecological modelling* **38**(3-4), 277–298.

Yerushalmi, S. & Green, R. M. (2009), 'Evidence for the adaptive significance of circadian rhythms', *Ecology letters* **12**(9), 970–981.

Zeller, K. A., McGarigal, K. & Whiteley, A. R. (2012), 'Estimating landscape resistance to movement: a review', *Landscape ecology* **27**(6), 777–797.

Zhang, J., Hull, V., Ouyang, Z., He, L., Connor, T., Yang, H., Huang, J., Zhou, S., Zhang, Z., Zhou, C. et al. (2017), 'Modeling activity patterns of wildlife using time-series analysis', *Ecology and evolution* **7**(8), 2575–2584.

Zhang, Z., Geiger, J., Pohjalainen, J., Mousa, A. E.-D., Jin, W. & Schuller, B. (2018), 'Deep learning for environmentally robust speech recognition: An overview of recent developments', *ACM Transactions on Intelligent Systems and Technology (TIST)* **9**(5), 49.

Zhao, K. & Jurdak, R. (2016), 'Understanding the spatiotemporal pattern of grazing cattle movement', *Scientific reports* **6**, 31967.

Zidon, R., Garti, S., Getz, W. M. & Saltz, D. (2017), 'Zebra migration strategies and anthrax in etosha national park, namibia', *Ecosphere* **8**(8), e01925.

Zucchini, W., MacDonald, I. L. & Langrock, R. (2016), *Hidden Markov models for time series: an introduction using R*, Chapman and Hall/CRC.