A hierarchical framework for segmenting movement paths

Wayne M. Getz[1,2]

[1]Dept. ESPM, UC Berkeley, CA 94720-3114, USA

[2]School of Mathematical Sciences, University of KwaZulu-Natal
Private Bag X54001, Durban 4000, South Africa

Abstract

Comparative applications of animal movement path analyses are hampered by the lack of a comprehensive framework for linking structures and processes conceived at different spatio-temporal scales. Although many analyses begin by generating step-length (SL) and turning-angle (TA) distributions from relocation time-series data—some of which are linked to ecological, landscape, and environmental covariates—the frequency at which these data are collected may vary from sub-seconds to several hours, or longer. The kinds of questions that may be asked of these data, however, are very much scale-dependent. It thus behooves us to clarify how the scale at which SL and TA data are collected related to one another and to the kinds of ecological questions that can be asked. Difficulties arise because the information contained in SL and TA time series is not semantically consilient with the physiological, ecological, and sociological factors that influence the structure of movement paths. I address these difficulties by classifying movement types at salient temporal scales using two different kinds of vocabularies. The first is the language derived from behavioral and ecological concepts. The second is the language derived from mathematically formulated stochastic walks. The primary tools for analyzing these walks are fitting time-series and stochastic-process models to SL and TA statistics (means, variances, correlations, individual-state and local environmental covariates), while paying attention to movement patterns that emerge at various spatial scales. The purpose of this paper is to lay out a more coherent, hierarchical, scale-dependent, appropriate-complexity framework for conceptualizing path segments at different spatio-temporal scales. Additionally, this framework is designed to bridge biological and mathematical movement ecology concepts; thereby, hopefully, stimulating the development of conceptually-rooted methods for simulating hypothetical and analyzing empirical data that can be used to test hypothesis regarding mechanisms driving animal movement.

Keywords: Fundamental movement elements, canonical activity modes, diel activity routines, life-history movement phases, life-time tracks

Introduction   28

The movement of organisms over landscapes can be usefully analyzed from various viewpoints.   29
Four are suggested by the conceptual framework for movement ecology proposed by Nathan et al.   30
(2008), depending on which questions constitute the focal interest—why move? how is movement   31
executed? where to move? and how does the immediate environment, as well as the internal state   32
of individuals, impact movement? Several more viewpoints arise when we focus on the structure   33
of the movement path itself and how this structure may fill space at different spatio-temporal   34
and, hence, ecological relevant scales (Levin 1992)—from the sequencing of various high-resolution   35
movement elements to the emergence of seasonal home ranges and habitat utilization distributions   36
(Worton 1987, Morellet et al. 2013, Torney et al. 2018, Wittemyer et al. 2019). The current lack of a   37
comprehensive framework for linking structures and processes conceived at different spatio-temporal   38
scales, however, hampers the conciliation of these viewpoints and of a comparative meta-analyses of   39
similar questions addressed across disparate species and spatio-temporal scales. With this in mind,   40
a more coherent, hierarchical, scale-dependent, appropriate-complexity framework than currently   41
exists is laid out in this paper for conceptualizing path segments at different spatio-temporal scales.   42
Of considerable importance is that this framework underscores the need for the collection of high   43
resolution relocation data (minute or subminute sampling intervals) whenever possible.   44

Nathan et al. (2008) argued that movement arises from the interactions of several processes and   45
motivations that can be understood through an analysis of external factors or stimuli, the internal   46
state of an organism (e.g. see Morelle et al. 2015), and the organism's mechanical and navigational   47
capacities. These interactions then result in a movement process that may be characterized by a   48
time series of relocation points that define movement paths (Siniff & Jessen 1969, Edelhoff et al.   49
2016). From these relocation time series, sequences of step lengths (SL) and turning angles (TA)   50
can be extracted (Kareiva & Shigesada 1983, Turchin 1998). Other data, such as accelerations   51
(Nathan et al. 2012), audio (Sapir et al. 2010, Hurme et al. 2019, Northrup et al. 2019) and   52
magneto (Williams et al. 2017, Chakravarty et al. 2019) data, as well as physiological measurements   53
(e.g., temperature, blood glucose levels) (Wilmers et al. 2015), may be recorded and used in the   54
context of relocation point covariates. Additional covariates may be environmental, including   55
measurements of local resistance to movement caused by landscape topology, vegetation features,   56
and local atmospheric conditions (Shepard et al. 2013, Getz & Saltz 2008, Spiegel et al. 2015),   57
as well as local ecosystem features that are often inferred using remotely sensed electromagnetic   58
spectral data (e.g., visual and infra-red satellite imagery or lidar) (Pettorelli et al. 2011, 2014, Bork   59
& Su 2007).   60

In this paper, the question is addressed of how we may reconcile the fact that SL and TA   61
distributions are not directly interpretable in terms of behavioral and ecological processes (e.g.,   62
resting, stalking, hiding, gathering); but, depending, on the spatio-temporal scale of the reloca-   63
tion data, phenomenological interpretations are made using scale-appropriate activity designators   64
(Johnson et al. 2002, Owen-Smith & Goodall 2019, Giotto et al. 2015). A deeper understanding of   65
the connections between path SL and TA time series and the designation of the so-called behav-   66
ioral state of an individual at each of its relocation points requires that we find a way to reconcile   67
narratives constructed using two different kinds of vocabularies. The first is the language of bi-   68
ological designators arising from behavioral and ecological concepts. Our primary tools here are   69
the application of path segmentation methods (Nams 2014, Edelhoff et al. 2016, Seidel et al. 2018)   70
that include hidden Markov models (HMMs; Michelot et al. 2016, Franke et al. 2004, Langrock   71

et al. 2012, Zucchini et al. 2016) and behavioral change-point analyses (BCPA; Gurarie et al. 2009, 2016, Owen-Smith & Martin 2015). The second is the language of mathematical stochastic walks (Kareiva & Shigesada 1983, Bartumeus et al. 2005, Patterson et al. 2008) that treats movement relocation data as a time series of values and applies existing time-series methods (Chatfield 2016) to finding the parameter values that best fit selected time-series models to movement paths, with the possible inclusion of individual and environmental covariates (Jonsen et al. 2003, Preisler et al. 2004, Johnson et al. 2008, Fleming et al. 2014, Calabrese et al. 2016).

Moving between stochastic-walk descriptions and biological designators in describing segments of movement paths is akin to translating between phonologic (e.g., Latin, cyrillic) and logographic (e.g., hieroglyphics, Chinese) scripts: much context-dependent interpretation needs to take place. But, in fact, the situation is much more dire, since we do not have access to every character—i.e., fundamental movement element (FuMEs, see glossary; Getz & Saltz 2008)—in the path sequence, but only some kind of "average" value across several characters or potential point-source values. The primary goal of this paper is to flesh out the temporal structure of biological designators of movement-path segments and provide more coherent insights into the relationship between biological designators and stochastic-process models at different temporal scales.

Biological Designators of Path Segments

The view that large segments of individual movement paths can be parsed into strings of fundamental movement elements (FuMEs—see glossary; Fig. 1, Table 1) with characteristic frequencies and patterns of occurrence is highly idealized. Identifying FuMES is generally a biomechanical problem that is beyond the scope and capabilities of analyses based on relocation data alone, although audio recordings, for example, have been used to identify the wingflaps of small birds (Sapir et al. 2010, Northrup et al. 2019). If the frequency of our relocation data, however, spans strings of around 10-100 FuMEs, occurring either repetitively or in correlated groups of a couple of FuME types, then we may be able to reliably identify different multiFuME or "metaFuME" aggregates. All metaFuMEs, in a set of several metaFuME types, has a fixed length equal to the time between consecutive relocation points and each metaFuME type has its own characteristic bivariate SL and TA distribution and time-series auto- and cross-correlations (Chatfield 2016). Such metaFuME sets can then be used as a basis for representing different types of identifiable activities (e.g., foraging or focused travelling to a selected location) as canonical activity modes (CAMs—see glossary; Getz & Saltz 2008).

Segmenting relocation data at a metaFuME level only makes sense if the data have been sampled at frequencies that lie somewhere between the time it takes to execute several typical FuMEs (i.e., on the order ten seconds) and the time it takes to perform a short-period CAM that is relevant to movement ecologists (i.e., on the order of a few minutes). Fortunately, new technologies are leading the collection of data at frequencies of around 0.01 to 0.1 Hz, and even 1 Hz in large birds (Harel, Horvitz & Nathan 2016, Harel & Nathan 2018). For example, relocation data at these frequencies is now being collected for small mammals and birds and, indeed for larger mammals as well (de Weerd et al. 2015, McGavin et al. 2018). Beyond GPS methods for collecting high frequency data are other methods such as reverse GPS (e.g., the Atlas system Weiser et al. 2016, Toledo et al. 2016); but the latter is limited at this time in spatial extent to some tens or hundreds of square kilometers. Also, high frequency movement data has been collected using video equipment for small (e.g., ants) to moderately-sized (e.g., mice) organisms in a laboratory setting (Spink et al, 2001, Kane et al.

2004, Delcourt et al. 2013).   115

From a practical point of view, particularly since the size of relocation data sets can be rather   116
large when collected at frequencies of around 0.01 to 0.1 Hz, it may be useful to carry out a   117
metaFuME identification process on selected segments of the full path W (such as CAMs), and   118
then use a suitable method to reconcile disparate identification efforts to obtain a consensus set   119
of metaFuME types that can be applied to the full movement path. Assuming that a set of   120
metaFuMEs has been identified, as discussed in more detail in succeeding sections, then it may   121
be possible to stably identify different kinds of short duration CAMs with periods varying from   122
around 10 metaFuMEs (i.e., a couple of minutes) to a few hours (Fig. 1, Table 1). Following   123
this, longer duration CAMs may be more conveniently parsed in terms of several shorter duration   124
CAMs rather than in terms of metaFuME types. To emphasize this, I point to the existence of   125
both short duration and long duration CAMs in the hierarchical scheme laid out in Table 1. Due to   126
the importance of circadian rhythms as physiological and behavioral drivers (Takahashi et al. 2008,   127
Yerushalmi & Green 2009, Hardin & Panda 2013), the diel cycle provides an empirically obvious   128
segmentation window for the identification of various types of diel activity routines (DARs: set of   129
green boxes in Fig. 1; also see Table 1). Several types of DARs may then be identified in terms of   130
differences in the type and frequency of their constituent CAMs. For example, different types of   131
DARs have been identified in terms of the portion of their diel cycle that various groups of lemurs   132
are active (Donati et al. 2016) and of the variation in the diel travel rates of turtles (Jonsen et al.   133
2006).   134

DARs, in turn, can be strung together to create life-history movement phases (LiMPs; sets of   135
red pentagons in Fig. 1) that, depending on the longevity of the species, may be periodic (e.g.,   136
annual migration; for a review see migration Milner-Gulland et al. 2011) or episodic (e.g., dispersal;   137
for a review see Bowler & Benton 2005). Finally, a sequences of LiMPs sequentially strung together   138
from the birth-to-death of an individual constitute its full lifetime track (LiTT; Nathan et al. 2008)   139
(indicated by the grey pentagons in Fig. 1; also see Table 1). Beyond LiMPs, LiTTs from several   140
individuals can be used to map out space-use by populations rather than just the movement path   141
of any single individual (Mueller & Fagan 2008).   142

This view of metaFuME, CAM, DAR and LiMP segmentation of LiTTs—in the context of   143
available data and the questions we would like to ask—provides a hierarchical classification system   144
of appropriate complexity (Getz et al. 2018, Larsen et al. 2016) for the parsing of movement paths.   145
It also provides a semantic framework for the quantitative narratives (Getz et al. 2019) needed to   146
explain the role that movement plays in the ecology and evolution of organisms. In the context   147
of evolution, for example, we expect natural selection to hone both the mechanics of performing   148
FuMEs and the way these FuMEs are organized into metaFuMEs, CAMs, DARs, and LiMPs   149
to create LiTTs that maximize the expected lifetime fitness of individuals. In particular, it is   150
of considerable interest how the frequency and sequencing of different FuMEs and CAMs in the   151
production of DARs and LiMPs affect individual fitness (Hein et al. 2016, Cattarino et al. 2016).   152

Beyond the relevance of diel cycles to movement behavior (Wittemyer et al. 2008), are lunar   153
(Polansky et al. 2010) and seasonal cycles (Marra et al. 2015). The critical nature of the diel scale   154
in driving movement behavior is reflected in our creation of the fixed-period DAR category (Fig. 1   155
for identifying key structures within LiMPs. In contrast, the lunar cycle may be more relevant to   156
some organisms than others (Polansky et al. 2010), while only for relatively long-lived organisms   157
will their LiTTs cover sufficiently many seasonal cycles to support a statistical analyses of repeated   158
LiMP motifs (Marra et al. 2015, Allen et al. 2018). For some organisms, particular LiMPs, such   159

Figure 1: **Scale-dependent path segmentation categories of lifetime track segments.** The time scale applies to typical medium and large terrestrial animals and needs to be adjusted for aerial and marine species, and small animals, including invertebrates. Round (blue), oval (lilac), triangular (yellow to orange), square (green), pentagonal (red), and hexagonal (grey) icons are respectively used to represent N_1 FuME types, N_2 metaFuME types, N_3 CAM types, N_4 DAR types, N_5 LiMP types, and an open-ended number of n LiTT types (one for the life time of each movement group type to which an individual can belong). Horizontal whiskers on icons represent variations in the length of examples of the same type within a category (except for metaFuMEs which are the length of time between successive relocation points and DARs which are nominally fixed to 24 hours) and vertical whiskers indicate some building-block variation within the same type (except for FuMEs which are much more stereotyped, and hence much less variable within types than other segmentation categories and metaFuMEs which are characterized by SL and TA distributions). Color shades within categories represent different types within those categories. Colorless strings of icons indicate how each category can be considered as a string of the next lower category elements (i.e., shapes and general color denote hierarchical level, while different shades of the same color indicate within level variation). Because FuMEs are hard to identify using relocation data only, a hierarchical segmentation of LiTTs will typically be supported by a metaFuME baseset rather than by a set of FuMEs themselves.

as dispersal, occur only once; while migration behavior may vary from year to year, influenced by interannual variations in climatic conditions—perhaps linked to multiyear marine (Mysterud et al. 2001, Grémillet & Boulinier 2009) or sunspot (Myers 2018) cycles. Further, periodically driven movement patterns may also change with age (e.g., exploration when young) and life history stage (e.g., elephants in musth). Thus, beyond DARs, other cyclic patterns become less obvious or more species-specific.

The Data Compatibility Problem

Deconstructing a movement path into its elements is somewhat like picking out words from a voice recording. The human brain does this very well, as do modern digital machines using deep-learning methods (Hinton et al. 2012, Zhang et al. 2018). In a very crude sense, FuMEs are comparable to words, and CAMs to sentences or even paragraphs. However, the number of FuMEs (on the order of 10^1-10^2) is likely four orders of magnitude smaller than the number of words in everyday speech, and the story told is a rather boring repetition of some tens of sentences repeated with moderate variation influenced by an individual's environment. DARs may be thought of as single pages, where each page can be identified as belonging to one of several types (e.g., a typical winter versus summer day, a day during a migratory versus non-migratory life-history movement phase—i.e,

Table 1: Hierarchical organization of a typical vertebrate lifetime track (LiTT). (Spatio-temporal scaling applies best to typical medium to large vertebrates, but not as well to small vertebrates, birds, or invertebrates.)

Scale/ Segment	Time	Space Constraint level: agency	Categories	Data	Some approaches and methods
true **FuMEs**[*]	0.1–10 secs	0.1–1000 cms	stride, trot, run twist, jump, flap	accelerometer video	machine learning, Newtonian mechanics
	highly constrained: biomechanical analyses				
meta- **FuMEs**[*]	resolution of relocation data sampling: ≥ 5 secs, ≤ 1 min	bivariate SL & TA[¶] distributions		SL & TA statistics[††]	ensemble filtering on means, variances & correlations
	base level: movement ecology analyses				
short duration **CAMs**[†]	0.1–10 mins	1–100 meters	resting, browsing, riding thermals	secs to mins, metaFuMEs & covariate data	path segmentation[**]
	highly flexible: sub-hourly/hourly focus				
long duration **CAMs**[†]	10-100 mins	0.1-10 kilometers	feeding traveling,	mins to hours, short CAMs & covariate data	
	highly flexible: sub-diel focus				
DARs[‡]	fixed 24-hr	diel range	central-place foraging ranging	mins to hours, CAM sequences	periodograms (Fourier, wavelets)
	time constrained: diel cycle				
LiMPs[§]	several days to a few months	home ranges and their shifts over time	philopatry, dispersal, migration, nomadic, recursion	DAR sequences	clustering methods, ideal-free/despotic distribution analyses
	strongly constrained: lunar and seasonal cycles				
partial or whole **LiTTs**[#]	paths encompassing several LiMPs	habitat types	territorial, migratory, central place foragers, nomads	LiMP sequences	GIS toolbox, deep-learning, pattern anal.
	moderately constrained: question-at-hand				

[*]Fundamental Movement Elements; [†]Canonical Activity Modes; [‡]Diel Activity Routines;
[§]Life-history Movement Phases; [#]Life Time Tracks; [¶]SL=step-length & TA=turning-angle;
[**]Includes Hidden Markov Models (HMMs) and Behavioral Change-Point Analyses (BCPA)
[††]See Section on **Stochastic Walks and MetaFuMEs** in main text

LiMP). ⁠176

Despite the current, technology driven, exponential growth in the quantity and quality of move- ⁠177 ment data (including relocation, accelerometer, physiological, and environmental measurements; ⁠178 Williams et al. 2019), we are still far from collecting the kind of multi-body-placement accelerom- ⁠179 eter and relocation data needed to identify the start and finish of individual FuMEs (i.e., with a ⁠180 spatial accuracy of centimeters rather than meters or even decimeters), although relocation using ⁠181 audio (Sapir et al. 2010, Hurme et al. 2019, Northrup et al. 2019) and magneto (Williams et al. 2017, ⁠182 Chakravarty et al. 2019) data show considerable potential in this respect. In short, how different ⁠183 parts of the body move in relationship to one another may well be required for the identification of ⁠184

certain FuMES. Thus, for some time to come, relocation data collected at fixed frequencies slower   185
than around 1 Hz are going to be temporally misalignment with the approximate (at least to within   186
half a second) of the start and finish of each of the FuMEs that make up a path segment of interest.   187

At this point, it is worth noting that accelerometer data alone has been used to parse out   188
relatively short time scale (order of 10 secs) behavioral elements along movement paths using   189
machine learning methods (Nathan et al. 2012, Thessen 2016). These behavioral elements, which   190
almost certainly include several FuME steps, in reality are partial elements of more extensive   191
short-duration CAMs that typically last tens of seconds to several minutes. Accelerometer data,   192
for example, was able to categorize standing, running, preening, eating, and active and passive flight   193
CAMs in vultures with 80-90% accuracy (Nathan et al. 2012). Another on baboons was able to   194
categorize foraging, lying, sitting, running, standing walking, groom (actor), groom (receiver) with   195
high accuracy ($> 90\%$), except for grooming, which was considerably harder to identify, particularly   196
when the focal individual was the receiver (Fehlmann et al. 2017). In these two examples, sitting,   197
grooming, preening and feeding CAMs are identifiable, but none of the basic FuMEs that are   198
executed during the generation of these CAMs are individually identified. It may be possible for   199
some FuMEs, however, such as the time taken to complete a FuME when walking versus running,   200
to be identifiable directly from accelerometer data (as we find in modern digital watches that are   201
able to monitor the number of steps we take while moving) or audio (Sapir et al. 2010, Hurme   202
et al. 2019, Northrup et al. 2019), magneto (Williams et al. 2017, Chakravarty et al. 2019) or video   203
(Spink et al. 2001, Kane et al. 2004, Delcourt et al. 2013) data.   204

In summary, location data on its own (i.e., without accompanying subsecond accelerometer,   205
accoustic, magneto or video data) is fundamentally incompatible with the identification of FuMEs   206
for two reasons: 207

- Consecutive point locations alone cannot determine what an individual is doing. What is   208
  needed in addition is, at least, the relative position of several different body parts of the   209
  individual to its center of gravity and, more usefully, the acceleration of these parts, or other   210
  kinds of data such as audio or video recordings.   211

- Regular sampling is likely to occur during the execution of FuMEs, rather than at the start   212
  or finish of a FuME; and, further, different FuMEs generally take different amounts of time   213
  to execute.   214

In light of these reasons, it is worth reemphasizing the following. 1.) Given that the identification   215
of individual FuMEs from relocation data is not generally possible, we are left with that the   216
rather challenging task of extracting a set of metaFuME elements, where each metaFuME type is   217
characterized by its own SL-TA distributional pair and correlations with and between consecutive   218
and simultaneous values; 2.). The complete set of metaFuME types, once identified, can then be   219
used as a basis for constructing an appropriate complexity (Getz et al. 2018, Larsen et al. 2016),   220
hierarchical framework of CAM, DAR and LiMP segments, where the latter is used to classify   221
different types of LiTTs (Table 1).   222

Current Path Analysis Methods   223

Before considering how to find a set of metaFuME distribution set that provides an appropriately   224
defined best fit to an empirical walk W (defined in terms of its set of observed relocation values)—   225

Figure 2: **Current scale-dependent analytical methods for analysing movement paths.** The temporal range is only suggestive and best applied to medium and large vertebrates. Also the image placements are not precise and some methods, such as machine learning (Thessen 2016), can be applied to data at any scale, but here are associated with the scale at which they are likely to be most useful. In addition, deep learning (useful for identifying different types of long-term patterns) is actually a subset of machine learning (where other deep-learning techniques, such as random forests and support vector machines have been applied to accelerometer data; Nathan et al. 2012, Fehlmann et al. 2017). Stochastic walks include correlated and biased random walks (Johnson et al. 2008). Space-time residence analyses represents a family methods that include first-passage-time (FTP; Fauchald & Tveraa 2003, McKenzie et al. 2009) and related approaches (Torres et al. 2017), while recurrence analyses cover a plethora of methods used to identify recursive movement patterns (Berger-Tal & Bar-David 2015, Bar-David et al. 2009). The melange of images is extracted from publications in the literature (Nathan et al. 2012, Panzacchi et al. 2016, Morales et al. 2004, DeCesare et al. 2012, Wittemyer et al. 2008, Lyons et al. 2013, Abrahms et al. 2017, Fleming & Calabrese 2017, Pohle et al. 2017, Torres et al. 2017, Gurarie et al. 2017), as well as created for this publication.

i.e., in 2-D, W is represented by the time series   226

$$W = \left\{ \big(t; x(t), y(t)\big) \big| t = 0, ..., t_f \right\} \tag{1}$$

let us briefly review the current state of path analysis that is relevant for analysing relocation data   227
sampled at scales typically appropriate for identifying CAMs and higher-level patterns (Fig. 2).   228
Note that the temporal resolution of the data will determine whether it is possible to identify short   229
duration CAMs (e.g., vigilance behavior during grazing; Fortin et al. 2004), long duration CAMs   230
(e.g., heading to water; Polansky et al. 2015), or only movement patterns at diel scales and beyond   231
(Spiegel & O'Farrell 2018, Owen-Smith 2013, Giotto et al. 2015).   232

At any time scale, movement time series may be analysed using purely statistical methods ap-   233
plied to the SL and TA time series derived from W, as explicitly described in the next section,   234
to generate various running statistics. These statistics may include the means, variance, autocor-   235
relations of SL and TA time series data and cross correlations between the two (Chatfield 2016,   236
McCulloch & Cain 1989, Bergman et al. 2000, Byers 2001). Running versions of these statistics   237
can then be used to identify points in time where abrupt changes in their values occur, using meth-   238

ods referred to as behavioral change-point analyses (BCPAs; Owen-Smith & Martin 2015, Chen & Gupta 2011, Matteson & James 2014, Killick & Eckley 2014, Gurarie et al. 2009, 2016) (Fig. 2) or, more generally, path segmentation methods (PS; Nams 2014, Edelhoff et al. 2016, Seidel et al. 2018).

Once particular CAM or DAR types of segments have been identified, they can be organized into ensembles of the same type. For clarity in referring to such ensembles, we use upper case calligraphic letters, which contrasts with the italicized uppercase letters used to denote sets of points that are essentially time series, such as W defined in Eq. 1. In particular, $W^{\text{seg-type}}$ will be used to refer to a segment of W of a particular type: e.g., "seg_type"=DAR$_1$ is a diel activity routine segment of type 1 and $\mathcal{W}^{\text{seg-type}}$ is used to refer to an ensemble or set of segments of this type: viz.,

$$R \text{ instances of a particular segment type:} \qquad \mathcal{W}^{\text{seg-type}} = \{W_r^{\text{seg-type}} | r = 1, ..., R\} \qquad (2)$$

Beyond purely statistical characterizations of the SL and TA time series is the notion that some sort of state (typically behavioral; but possibly physiological, disease, etc.) can be associated with each location point in the walk W (Patterson et al. 2008). A very simple example is the stop/move representation of cattle grazing paths (Zhao & Jurdak 2016). The most widely used methods for identifying subsets of points that reflect some underlying state or canonical activity mode (CAM) involve hidden Markov models (HMM) (Michelot et al. 2016, Franke et al. 2004, Langrock et al. 2012, Zucchini et al. 2016) (Fig. 2), although an independent mixture modeling approach has been shown to be equally viable, if not better (Goodall et al. 2017). These methods are particularly germane when covariate data are available at each location, such as accelerometer and temperature data from GPS collars, remotely sensed physiological information (Cross et al. 2016), vegetation structure (Tsalyuk et al. 2019), or physical landscape features that, for example, facilitate or retard movement (Zeller et al. 2012). The task of an HMM is to find a best estimate of which state of a current set of "hidden" states H_i belonging to a set of n possible states $\mathcal{H} = \{H_1, ..., H_n\}$ should be assigned to the individual when at the location $(x(t), y(t))$.

HMMs can be applied at any temporal scale, including the metaFuME scale, depending only on the relocation and covariate data. Other methods, such as step-selection function (SSF) methods, which address the question of where am I likely to move next as a function of landscape covariates, can be applied at any scale (Fortin et al. 2005, Panzacchi et al. 2016, Thurfjell et al. 2014, Avgar et al. 2016). SSFs, however, address different kinds of questions at different scales: they have more of a behavioral interpretation at a 1-minute scale while more of an ecological (e.g., resource use) interpretation at multi-hour scales. On the other hand, habitat or resource-selection function (HSF or RSF) methods (Boyce et al. 2002, McGarigal et al. 2016, DeCesare et al. 2012) are, by their very nature, low resolution models that typically apply at a seasonal level and assume non-autocorrelated location data. When RSFs are evaluated at a life-time level over many individuals in a population, they are called niche models—but such models no longer rely on movement data, only on point location data (Townsend Peterson et al. 2007).

Other general time-series or signal-analysis methods that have been co-opted by movement ecologists include wavelet analysis (Wittemyer et al. 2008, Polansky et al. 2010)—which provides interpretations at the resolution of one to several orders of magnitude greater than the frequency at which data are collected (e.g., daily movement to monthly movement patterns for data collected at the frequency of 1-hour for at least a year)—and machine learning approaches (Thessen 2016, Valletta et al. 2017) that have been applied both to location (Bennison et al. 2018) and accelerometer

data (Nathan et al. 2012) (Fig. 2). These methods, however, do not have an extensive history of application in movement ecology, but are likely to become more ubiquitous as the field of movement ecology matures.

Stochastic Walk Statistics

As discussed earlier, identifying a set of FuMEs is a problem that requires biomechanical (Delp & Loan 2000), audio (Sapir et al. 2010), or other kinds of covariate data (Chakravarty et al. 2019) than relocation data on its own. With relocation data alone, a hierarchical analysis of a path can only be underpinned by elements that are derived statistically from the relocation time series W (Eq.1).

From this time series we can extract a 1-D time series of step lengths and another 1-D time series of turning angles as follows:

- Generate the step-length (SL) time series

$$S = \{(t, s(t))| \ t = 1, ..., t_f\} \tag{3}$$

  using the derived values

$$s(t) = \sqrt{(x(t) - x(t-1))^2 + (y(t) - y(t-1))^2} \tag{4}$$

- Generate the turning-angle (TA) time series

$$A = \{(t, a(t))| \ t = 2, ..., t_f\} \tag{5}$$

  using the derived values

$$a(t) = \arctan\left(\frac{y(t) - y(t-1)}{x(t) - x(t-1)}\right) - \arctan\left(\frac{y(t-1) - y(t-2)}{x(t-1) - x(t-2)}\right) \tag{6}$$

From these two time series we can obtain their means and variances respectively denoted by (μ_x, σ_x), $x = s, a$, and then use them compute the following three variation related time series:

- Generate two normalized (by the appropriate variances) "running-term autocorrelation" time series
$$V^{xx} = \{(t, v^{xx}(t))| \ t = 1, ..., t_f\} \quad \text{for } x = a, s, \tag{7}$$

  where
$$v^{xx}(t) = \frac{(x(t) - \mu_x)(x(t-1) - \mu_x)}{\sigma_x} \tag{8}$$

- Generate a normalized "running-term cross-correlation" time series

$$V^{as} = \{(t, v^{as}(t))| \ t = 1, ..., t_f\} \tag{9}$$

  where
$$v^{as}(t) = \frac{(a(t) - \mu_a)(s(t) - \mu_s)}{\sqrt{\sigma_a \sigma_s}} \tag{10}$$

Figure 3: **Parsing the walk W into a set of metaFuMEs.** Associated with each point t is a set of running values $\big(s(t), a(t), v^{ss}(t), v^{aa}(t), v^{as}(t), \text{other possible covariates}\big)$. Cluster analysis may be used to generate metaFuME categories. For purposes of illustration, we depict three types of metaFuMEs with associated lists of time steps that can then be used to define the ensemble of values associated with each metaFuME type: blue = $\mathcal{T}_1$, red = $\mathcal{T}_2$, and green = $\mathcal{T}_3$. Note that we expect transition points between strings of metaFuMEs of one type (broken line segments), such as $t-6$, $t-1$ and $t+4$, could well be outliers in the clustering process. Outliers may be assigned to the cluster to which they are closest in some suitably defined sense (as indicated by the colors of the broken lines).

We are now challenged with the task of using the five time series above—and, perhaps, other covariate data when available (e.g., acoustic, accelerometer, local environmental)—to parse W into an several, say Φ, different one-step ensembles (Fig. 3). Each of these Φ ensembles can now be regarded as a set of drawings from the same bivariate distribution $D_\phi(s, a)$, $\phi = 1, ..., \Phi$. If the relocation sampling period, except for speed bursts, encompasses around 10-50 FuMEs (i.e., more than around 5 seconds but less than about a minute—fast walking in humans crudely corresponds to 2 steps per second, though most movement is slower than this); then, in the proposed hierarchical segmentation framework, we are in the metaFuME segment zone (Fig. 1, Table 1). If the relocation sampling period is on the order of tens of minutes to several hours, then each relocation likely encompasses hundreds to tens of thousands of FuMEs. In this case, set of states are more appropriately identified as short or long duration CAMs than metaFuMEs.

Once a set of metaFuMEs has been identified from a particular ensemble of segments of the same CAM or DAR type, we are still faced with the task of reconciling various metaFuME sets obtained from different CAM or DAR ensembles into a comprehensive metaFuME set. Such a set will then underlie all LiMPs of a particular type, but not necessarily across a complete LiTT, because we might expect metaFuME sets for, say, juveniles versus adults to be different for same species. Many different types of cluster analyses (McGarigal et al. 2016) can be tried to identify metaFuMEs. Also, multicrietria optimization methods using, for example, genetic algorithms or approximate Bayesian computations (Coello 2003, Odu & Charles-Owaba 2013, Marin et al. 2012)) could be used to reconcile metaFuME sets obtained from ensembles of different types of segments. Such investigations, however, are well beyond the scope of this concepts paper.

Once the time series W and, hence, the corresponding SL and TA time series have been parsed into ensembles and organized into a set of Φ metaFuMEs, we then identify the index time sets T_ϕ, $\phi = 1, ..., \Phi$, associated with each ensemble (Fig. 3). The distributions of all the joint pairs $(s(t), a(t))$, where $t \in \mathbf{T}_\phi$, $\phi = 1, ..., \Phi$, may be fitted to appropriate metaFuME bivariate distributions $D_\phi(s, a)$, $\phi = 1, ..., \Phi$. In essence, we have estimated the a set of bivariate metaFuME

distributions

$$\mathcal{D}_\Phi = \{D_1(s,a), ..., D_\Phi(s,a)\}. \tag{11}$$

The final missing piece is now use the set of switching times among the index sets (this would be switching between modes at the finest possible scale (Patterson et al. 2008) rather than at courser scales Morales et al. 2004, Jonsen et al. 2007)

$$\mathcal{T}_\Phi = \{T_\phi | \phi = 1, ..., \Phi\}$$

to estimate a matrix of probabilities $(\mathbf{P})_{\phi\psi} = p_{\phi\psi}$, ϕ, $\psi = 1, ..., \Phi$ of the probability of sampling values from the distribution $D_\phi(s,a)$ when the previous drawing was from the distribution $D_\psi(s,a)$. In this case, any walk $W(t_f)$ can be modeled as a process generated by first undertaking the following T drawings (the hat notation represents drawings)

$$T \text{ drawings} := \quad \{(\hat{s}_t, \hat{a}_t) \text{ sampled from } D_\psi(s,a) \text{ with probability } p_{\phi\psi} | t = 0, ..., t_f - 1\} \tag{12}$$

and then generating $W(t_f)$ from these drawings using the equations (Patterson et al. 2008)

$$\begin{aligned} x(t+1) &= x(t) + \hat{s}_t \cos \hat{a}_t \\ y(t+1) &= y(t) + \hat{s}_t \sin \hat{a}_t \end{aligned} \tag{13}$$

Thus, in short, the drawings are a Markov process that facilitates the construction of a walk W that is one instantiation of an ensemble of walks that can be generated using Eqs. 12 and 13 from a proposed or identified metaFuME-distribution-set and transition-matrix pair $(\mathcal{D}_\Phi, \mathbf{P})$.

A priori, the best-fit number of distribution pairs Φ in the set $\mathcal{D}_\Phi$ (Eq. 11) is unknown. A first guess at this number might be the number of modes in H^{SL} or H^{TA}, if such modes are evident. The problem of finding the best mixture of unimodal pairs $(D_\phi^{\text{SL}}, D_\phi^{\text{TA}})$ of distributions that best fit the empirical data is a challenging estimation problem that has been approached in several ways, including ensemble Kalman filtering (Dovera & Della Rossa 2011) and assuming the underlying distributions are log-concave (Walther et al. 2009). A prudent approach may be to proceed by first looking for the two best-fitting pairs of bivariate distributions (the tilde used to indicate that they are "best fitting"):

$$\tilde{\mathcal{D}}_2 = \{\tilde{D}_1(s,a), \tilde{D}_2(s,a)\} \tag{14}$$

and then moving on to three, four, and so on, until no improvement in fit is obtained at the next value for Φ in an information theoretic sense (Symonds & Moussalli 2011).

In a nutshell, the challenge we face is the identification of a set of distributions $\tilde{\mathcal{D}}_\Phi$(where *a priori* the size of the set Φ is unknown) and a Markov transition matrix $\tilde{\mathbf{P}}$ that can then be used to generate N walks $\tilde{W}_i(t_f)$, $i = 1, ..., N$ of length T for which our particular empirical walk $W^{\text{data}}(t_f)$ is most likely to belong; or expressed mathematical using uppercase italics to denote individual walks and uppercase calligraphy to denote a set of walks, suppressing the argument in T we have:

$$\tilde{\mathcal{W}}^{\text{ensemble}} = \left\{\tilde{W}_i | i = 1, ..., N\right\} \quad \text{is a "best fitting" ensemble to} \quad W^{\text{data}} \tag{15}$$

This of course is an extremely challenging undertaking that can be made more manageable by breaking it down into a series of separate more manageable tasks, as discussed below, to make the identification of the best-fitting pair $\left(\tilde{\mathcal{D}}_\Phi, \tilde{\mathbf{P}}\right)$ possible.

Hierarchical Segmentation of Walks

From a practical point of view, particularly as the size of relocation data sets increases when collected at frequencies of around 0.01 to 0.1 Hz, the following approach might prove useful to carrying out a hierarchical segmentation analysis of relocation data. The goal is to obtain ensembles of similar segments that can then used to identify sets of metaFuMEs, initially ensemble by ensemble of similar segments types (typically, ensembles of particular CAM types), and then through consolidation of various metaFuME sets to obtain single metaFuME set applicable to segments of a particular LiMP type (since different LiMPs may be age dependent and fundamental movement elements may also be age dependent)

1. From plots of a relocation data set W^{data} on landscapes (i.e., eyeballing it)—or using any appropriate identification method, such as net-square-displacement data (Bunnefeld et al. 2011, Bischof et al. 2012)—select one or more subsets of W^{data} that individually appear to belong to particular types of LiMPs (e.g., sort segments of W^{data} into $\text{LiMP}_{\text{type}}$, where type=migratory, wet season-residence, etc.) (Fig. 4). The occurrences and frequencies of different types of LiMPs in a LiTT can be then used to categorize an individual's LiTT type, thereby providing a less adhoc alternative to other approaches that may be used (Abrahms et al. 2017)).

2. Sort 24-hour segments of $W^{\text{LiMP}_{\text{type}}}$ (e.g., midnight to next midnight for diurnal species or midday to next midday for nocturnal species) into one or more DAR types ($W^{\text{DAR}_{\text{type}}}$, type $= 1, 2, ...$) and generate a set of summary statistics relating to the SL and TA time series associated with these DAR segments.

3. Use SL, TA, and derived correlation statistics statistics (Eqs. 3-10) to cluster (McGarigal et al. 2016) these 24-hour DAR segments into a set of g_n DAR types or use machine learning methods (Thessen 2016) to create such sets: i.e., recalling notational definition Eq. 2, create sets $\mathcal{W}^{\text{DAR}_g}$ and fit their summary statistics to appropriate distributions $D^{\text{DAR}_g}(s, a)$, $g = 1, ..., g_n$. Note that different $W^{\text{LiMP}_{\text{type}}}$ may yield the same set of DARs, but in very different proportions over their extents. Alternatively, some of them may have one or more DAR types not found in other $W^{\text{LiMP}_{\text{type}}}$ segments.

4. After stringing together all the data contained in $\mathcal{W}^{\text{DAR}_g}$, use one or more segmentation methods to obtain a consensus on the best way to segment these strings into an optimal number of CAM types (Fig. 4). Depending on the resolution of the data, these may be long or short duration CAMs or both (Owen-Smith & Martin 2015). If the process is first applied to obtain long duration CAMS (e.g., using BCPA based on a relatively wide sliding window of say 30 minutes; Owen-Smith & Martin 2015), then each of these long duration CAMS may then be further segmented into short duration CAMS (e.g., using BCPA based on a narrower sliding window of say 10 minutes, should the data permit), though care has to be taken that short duration CAMs obtained from different long duration CAMs are reconciled to optimize the number of shorter duration CAMs that, in different proportions, constitute long duration CAMs. The same can be said of optimizing the number of CAMs used in different proportions to build up the different types of DARs (Fig. 4)

5. Using various sets of CAMs and, possibly DARs, of the same type, apply "yet to be devised methods" to identify the sets of metaFuMES, using the general approach outlined in the

next section that will likely drawn upon clustering methods (McGarigal et al. 2016) and   400
multicriteria optimization (Odu & Charles-Owaba 2013).   401

Figure 4: **The structure of a movement path at different scales.** A multi-season lift-time track
(LiTT) W (Eq. 1) can be segmented at various scales: first into different types of life-history movement phases
(LiMPs), each which can be then be parsed into diel segments that are then classified as being of a particular
diel activity routine (DAR) type. These DARs can be parsed into canonical activity mode (CAM) segments,
each of which are composed of different proportions of metaFuMEs (aggregates of correlated fundamental
movement elements), where each metaFuME is characterized by a joint step-length and turning-angle ((s,a);
Eqs 4 and 6) distribution with specific first-order and correlative statistics. Short duration CAMs (e.g.,
CAM 3) can easily be hidden by being averaged into longer duration CAMs (e.g., CAMs 1 and 2), if DAR
segmentation analyses are not done at a sufficient fine scale. For simplicity, only two or three types of path
segments are depicted at each identified scale.

Conclusion   402

Without an underlying framework to organize information, deep comprehension within any scientific   403
field is impossible. Clearly this has been true of the physical sciences with theories of gravity and   404
the structure of matter being central to our current understanding of the physical universe. This   405
is also true of the life sciences where, for example, the Linnaean taxonomic system, with some   406
modifications, has reigned supreme over the past 250 years (Schuh 2003). This reign still holds   407
today, even within the microbial sciences (Rosselló-Móra & Whitman 2019, Moore et al. 2010)   408
where Mayr's biological species concept fails in several critical ways (Gevers et al. 2005).   409

In the context of movement ecology, the classification of movement types at different spatio-   410
temporal scales is of considerable interest, although approaches to date have been somewhat in-   411
formal. At the LiTT level, basic life-history types regarding, *interalia*, dispersal and migration   412
behavior may be identified. In the movement ecology literature, we see many studies interested   413
in diel activity routine (DARs) (Rahimi & Owen-Smith 2007, Owen-Smith 2013, Owen-Smith &   414
Goodall 2014), life-time phase (LiMPs) (Fahr et al. 2015, Marra et al. 2015) and over-all life time   415
tracks (LiTTs) (Abrahms et al. 2017). Contrasting DAR types may include distinctions among   416
nomadic, central-place foraging, or territorial behavior, where a single LiTT may have LiMPs dom-   417
inated by one or other of these DARs, as in male springbok in Etosha National Park, Namibia,   418

exhibiting territorial behavior during the wet season and nomadic behavior during the dry season, with daily excursions to the same water-hole around midday during the wet season (Lyons et al. 2013). Thus DARs are also very likely to be influenced by seasonal factors, as in pandas that change diel levels of activity in response to seasonal drivers (Zhang et al. 2017).

Extensive effort has also been made to parse diel and longer segments into various types of sub-diel activity modes (Owen-Smith et al. 2010, Donati et al. 2016) that, if stably identifiable across different segments, can be organized into a set of canonical activity modes (CAMs; Getz & Saltz 2008). At finer time scales, however, beyond using accelerometer, acoustic and magento data to identify behavioral states (Williams et al. 2017, Hurme et al. 2019, Nathan et al. 2012, Fehlmann et al. 2017, Chakravarty et al. 2019, Sapir et al. 2010, Zucchini et al. 2016), very little work has been undertaken to identify sets of fundamental movement elements (FuMEs; same as FMEs in Getz & Saltz 2008) from which CAMs are constructed. As pointed out in this paper, using relocation data alone, we cannot endeavour to identify an underlying set of FuMEs. The best we can hope for, provide subminute relocation data is available, is the identification of a set of metaFuMEs consisting of repeated or correlated strings of FuMES, and characterized by particular step-length and turning-angle distributions.

We should not underestimate the analytical and computational challenge required to extract a comprehensive and stable (across many segments at various scales) set of metaFuME elements. The best methods to do this still remain to be developed and how well this may be accomplished remains to be seen. It may require many of the ideas presented here to be refined or modified. But there is no denying, however, that if successful, a formalized, widely accepted, hierarchical-path-segmentation framework will greatly facilitate efforts to address outstanding questions in movement ecology, particularly those involving comparative analyses within species (Wittemyer et al. 2019), as well as across species. Within species variation may allow us to assess differences along geographic clines with application to the behavioral adaptation of species under landscape and climate change (Seebacher & Post 2015). It may help us assess fitness in the context of feeding strategies or social behavior (Harel et al. 2017, Harel, Duriez, Spiegel, Fluhr, Horvitz, Getz, Bouten, Sarrazin, Hatzofe & Nathan 2016) or mapping out landscapes in terms of their overall resistance to movement (Zeller et al. 2012) (which, for example, has bearing on the amount of energy it takes to travel a given distance) or areas attractive to populations (Giotto et al. 2015). It may also be diagnostic of changes in movement behavior when individuals are stressed, ill, have genetic defects, or females are pregnant (Spink et al. 2001, Owen-Smith 2013); or it may be predictive in terms of pathogen transmission and the spread of disease (Cross et al. 2007, Tracey et al. 2014, Dougherty et al. 2018, Zidon et al. 2017). Differences in various species' LiMP/DAR/CAM movement profiles may be critical when it comes to taking a multispecies approach to assessing the impacts of ecosystem management on species conservation (Brodie et al. 2015, Runge et al. 2014).

In addition, a formalized, hierarchical-path-segmentation framework of appropriate complexity (Getz et al. 2018, Larsen et al. 2016) can be used, for example, to address the dozen plus questions that were recently posed regarding the movement ecology of marine megafauna Hays et al. (2016)— but are equally applicable to all animal species. These questions include: 1) "Are there simple rules underlying seemingly complex movement patterns and hence common drivers for movement across species?" and 2.) "How will climate change impact animal movements?" An ability to address both of these question, as well as those raised above and others besides, in a comparative way—with a level of consistency and coherency that only a universally accepted classification framework can provide—is needed with great urgency, as the field of movement ecology matures. The framework

presented here is a starting point to developing a more comprehensive one that can be fleshed out   464
in the future, supported by analytical and computational methods, some of which still need to be   465
developed or refined from existing methods.   466

Glossary   467

An expansion of the various acronyms used in the text are provided here in alphabetical order along   468
with a very brief definition/explanation of their meaning.   469

BCPA: Biological Change Point Analysis. Biological change-point analysis refers to a group of   470
methods used to determine how the statistics of a biological variable y (e.g., its mean, variance,   471
autocorrelation, or rate of change in slope or curvature), dependent on a second variable x,   472
switches at threshold points with changes in x (e.g., space, time, or the abscissa in a stress-   473
response relationship) (Morales et al. 2004, Andersen et al. 2009, Chen & Gupta 2011, Jonsen   474
et al. 2007). Note, more generally, x could be vector-valued.   475

CAM: Canonical Activity Mode. This is a stably classifiable subdiel behavioral mode (pattern of   476
movement) such as a foraging bout, resting period, or purposeful heading (i.e., traveling) to   477
a distant target location.   478

DAR: Diel Activity Routine. This is a stably classifiable 24-hour sequence of CAMs that occur   479
at characteristic frequencies and times of the day.   480

FuME: Fundamental Movement Elements (also FME; Getz & Saltz 2008). This is a relatively   481
rapid, highly repeatable, steorotypic set of body movements that forms the basis of the   482
locomotory capacity of an individual (e.g., a walk step, a running step, a wing flap, a jump,   483
etc.)   484

HMM: Hidden Markov Model. A time series of observable values that depend in a probabilistic   485
sense on the values of an associated, but unobservable Markov chain process (a set of states   486
where the transition from one state to another depends only on the value of the current state)   487
(Zucchini et al. 2016)   488

LiMP: Life-history Movement Phase. This is a path segment that typically reflects a life-history   489
relevant movement behavior such as dispersal (episodic), migration (periodic), or other peri-   490
odic behaviors at a greater-than-diel scale.   491

LiTT: Life Time Track. This is the total movement path of an individual from its birth to its   492
death   493

PS: Path Segmentation. This is the process of breaking up a movement path into reliably stable   494
metaFuMEs, CAMs, DARs, and LiMPs using a suite of methods that include BCPA and   495
HMM approaches (Nams 2014, Edelhoff et al. 2016, Seidel et al. 2018).   496

SL: Step Length. The distance between consecutive relocation points, as generated from Eq. 1   497
using Eq. 4   498

TA: Turning Angle. The change in the angle of heading across three consecutive relocation points,   499
as generated from Eq. 1 using Eq. 6.   500

Acknowledgements   501

I thank Briana Abrahms, Shirli Bar-David, Paul Cross, Victoria Goodall, Nir Horvitz, Leif Loe,   502
Ran Nathan, Norman Owen-Smith, Leo Polansky, David Saltz, Dana Seidel, Orr Spiegel, Krti   503
Tallam, Miriam Tsalyuk, George Wittemyer for comments on earlier versions of this paper. All   504
inaccuracies and misconceptions are my own.   505

References   506

Abrahms, B., Seidel, D. P., Dougherty, E., Hazen, E. L., Bograd, S. J., Wilson, A. M., McNutt, J. W.,   507
Costa, D. P., Blake, S., Brashares, J. S. et al. (2017), 'Suite of simple metrics reveals common movement   508
syndromes across vertebrate taxa', *Movement ecology* **5**(1), 12.   509

Allen, R. M., Metaxas, A. & Snelgrove, P. V. (2018), 'Applying movement ecology to marine animals with   510
complex life cycles', *Annual Review of Marine Science* **10**, 19–42.   511

Andersen, T., Carstensen, J., Hernandez-Garcia, E. & Duarte, C. M. (2009), 'Ecological thresholds and   512
regime shifts: approaches to identification', *Trends in Ecology & Evolution* **24**(1), 49–57.   513

Avgar, T., Potts, J. R., Lewis, M. A. & Boyce, M. S. (2016), 'Integrated step selection analysis: bridging the   514
gap between resource selection and animal movement', *Methods in Ecology and Evolution* **7**(5), 619–630.   515

Bar-David, S., Bar-David, I., Cross, P. C., Ryan, S. J., Knechtel, C. U. & Getz, W. M. (2009), 'Meth-   516
ods for assessing movement path recursion with application to African buffalo in South Africa', *Ecology*   517
90(9), 2467–2479.   518

Bartumeus, F., da Luz, M. G. E., Viswanathan, G. M. & Catalan, J. (2005), 'Animal search strategies: a   519
quantitative random-walk analysis', *Ecology* **86**(11), 3078–3087.   520

Bennison, A., Bearhop, S., Bodey, T. W., Votier, S. C., Grecian, W. J., Wakefield, E. D., Hamer, K. C.   521
& Jessopp, M. (2018), 'Search and foraging behaviors from movement data: a comparison of methods',   522
Ecology and evolution **8**(1), 13–24.   523

Berger-Tal, O. & Bar-David, S. (2015), 'Recursive movement patterns: review and synthesis across species',   524
Ecosphere **6**(9), 1–12.   525

Bergman, C. M., Schaefer, J. A. & Luttich, S. (2000), 'Caribou movement as a correlated random walk',   526
Oecologia **123**(3), 364–374.   527

Bischof, R., Loe, L. E., Meisingset, E. L., Zimmermann, B., Van Moorter, B. & Mysterud, A. (2012), 'A   528
migratory northern ungulate in the pursuit of spring: jumping or surfing the green wave?', *The American*   529
Naturalist **180**(4), 407–424.   530

Bork, E. W. & Su, J. G. (2007), 'Integrating lidar data and multispectral imagery for enhanced classification   531
of rangeland vegetation: A meta analysis', *Remote Sensing of Environment* **111**(1), 11–24.   532

Bowler, D. E. & Benton, T. G. (2005), 'Causes and consequences of animal dispersal strategies: relating   533
individual behaviour to spatial dynamics', *Biological Reviews* **80**(2), 205–225.   534

Boyce, M. S., Vernier, P. R., Nielsen, S. E. & Schmiegelow, F. K. (2002), 'Evaluating resource selection   535
functions', *Ecological modelling* **157**(2-3), 281–300.   536

Brodie, J. F., Giordano, A. J., Dickson, B., Hebblewhite, M., Bernard, H., Mohd-Azlan, J., Anderson, J. & Ambu, L. (2015), 'Evaluating multispecies landscape connectivity in a threatened tropical mammal community', *Conservation Biology* **29**(1), 122–132.

Bunnefeld, N., Börger, L., van Moorter, B., Rolandsen, C. M., Dettki, H., Solberg, E. J. & Ericsson, G. (2011), 'A model-driven approach to quantify migration patterns: individual, regional and yearly differences', *Journal of Animal Ecology* **80**(2), 466–476.

Byers, J. A. (2001), 'Correlated random walk equations of animal dispersal resolved by simulation', *Ecology* **82**(6), 1680–1690.

Calabrese, J. M., Fleming, C. H. & Gurarie, E. (2016), 'ctmm: an r package for analyzing animal relocation data as a continuous-time stochastic process', *Methods in Ecology and Evolution* **7**(9), 1124–1132.

Cattarino, L., McAlpine, C. A. & Rhodes, J. R. (2016), 'Spatial scale and movement behaviour traits control the impacts of habitat fragmentation on individual fitness', *Journal of Animal Ecology* **85**(1), 168–177.

Chakravarty, P., Maalberg, M., Cozzi, G., Ozgul, A. & Aminian, K. (2019), 'Behavioural compass: animal behaviour recognition using magnetometers', *Movement Ecology* **7**(1), 28.

Chatfield, C. (2016), *The analysis of time series: an introduction*, Chapman and Hall/CRC.

Chen, J. & Gupta, A. K. (2011), *Parametric statistical change point analysis: with applications to genetics, medicine, and finance*, Springer Science & Business Media.

Coello, C. A. C. (2003), Evolutionary multi-objective optimization: A critical review, *in* 'Evolutionary optimization', Springer, pp. 117–146.

Cross, P. C., Almberg, E. S., Haase, C. G., Hudson, P. J., Maloney, S. K., Metz, M. C., Munn, A. J., Nugent, P., Putzeys, O., Stahler, D. R. et al. (2016), 'Energetic costs of mange in wolves estimated from infrared thermography', *Ecology* **97**(8), 1938–1948.

Cross, P. C., Edwards, W. H., Scurlock, B. M., Maichak, E. J. & Rogerson, J. D. (2007), 'Effects of management and climate on elk brucellosis in the greater yellowstone ecosystem', *Ecological Applications* **17**(4), 957–964.

de Weerd, N., van Langevelde, F., van Oeveren, H., Nolet, B. A., Kölzsch, A., Prins, H. H. & de Boer, W. F. (2015), 'Deriving animal behaviour from high-frequency gps: tracking cows in open and forested habitat', *Plos one* **10**(6), e0129030.

DeCesare, N. J., Hebblewhite, M., Schmiegelow, F., Hervieux, D., McDermid, G. J., Neufeld, L., Bradley, M., Whittington, J., Smith, K. G., Morgantini, L. E. et al. (2012), 'Transcending scale dependence in identifying habitat with resource selection functions', *Ecological Applications* **22**(4), 1068–1083.

Delcourt, J., Denoël, M., Ylieff, M. & Poncin, P. (2013), 'Video multitracking of fish behaviour: a synthesis and future perspectives', *Fish and Fisheries* **14**(2), 186–204.

Delp, S. L. & Loan, J. P. (2000), 'A computational framework for simulating and analyzing human and animal movement', *Computing in Science & Engineering* **2**(5), 46–55.

Donati, G., Campera, M., Balestri, M., Serra, V., Barresi, M., Schwitzer, C., Curtis, D. J. & Santini, L. (2016), 'Ecological and anthropogenic correlates of activity patterns in eulemur', *International Journal of Primatology* **37**(1), 29–46.

Dougherty, E. R., Seidel, D. P., Carlson, C. J., Spiegel, O. & Getz, W. M. (2018), 'Going through the motions: incorporating movement analyses into disease research', *Ecology letters* **21**(4), 588–604.

Dovera, L. & Della Rossa, E. (2011), 'Multimodal ensemble kalman filtering using gaussian mixture models', *Computational Geosciences* **15**(2), 307–323.

Edelhoff, H., Signer, J. & Balkenhol, N. (2016), 'Path segmentation for beginners: an overview of current methods for detecting changes in animal movement patterns', *Movement ecology* **4**(1), 21.

Fahr, J., Abedi-Lartey, M., Esch, T., Machwitz, M., Suu-Ire, R., Wikelski, M. & Dechmann, D. K. (2015), 'Pronounced seasonal changes in the movement ecology of a highly gregarious central-place forager, the african straw-coloured fruit bat (eidolon helvum)', *PloS one* **10**(10), e0138985.

Fauchald, P. & Tveraa, T. (2003), 'Using first-passage time in the analysis of area-restricted search and habitat selection', *Ecology* **84**(2), 282–288.

Fehlmann, G., ORiain, M. J., Hopkins, P. W., OSullivan, J., Holton, M. D., Shepard, E. L. & King, A. J. (2017), 'Identification of behaviours from accelerometer data in a wild social primate', *Animal Biotelemetry* **5**(1), 6.

Fleming, C. H. & Calabrese, J. M. (2017), 'A new kernel density estimator for accurate home-range and species-range area estimation', *Methods in Ecology and Evolution* **8**(5), 571–579.

Fleming, C. H., Calabrese, J. M., Mueller, T., Olson, K. A., Leimgruber, P. & Fagan, W. F. (2014), 'From fine-scale foraging to home ranges: a semivariance approach to identifying movement modes across spatiotemporal scales', *The American Naturalist* **183**(5), E154–E167.

Fortin, D., Beyer, H. L., Boyce, M. S., Smith, D. W., Duchesne, T. & Mao, J. S. (2005), 'Wolves influence elk movements: behavior shapes a trophic cascade in yellowstone national park', *Ecology* **86**(5), 1320–1330.

Fortin, D., Boyce, M. S., Merrill, E. H. & Fryxell, J. M. (2004), 'Foraging costs of vigilance in large mammalian herbivores', *Oikos* **107**(1), 172–180.

Franke, A., Caelli, T. & Hudson, R. J. (2004), 'Analysis of movements and behavior of caribou (rangifer tarandus) using hidden markov models', *Ecological Modelling* **173**(2-3), 259–270.

Getz, W. M., Marshall, C. R., Carlson, C. J., Giuggioli, L., Ryan, S. J., Romañach, S. S., Boettiger, C., Chamberlain, S. D., Larsen, L., DOdorico, P. et al. (2018), 'Making ecological models adequate', *Ecology letters* **21**(2), 153–166.

Getz, W. M., Salter, R. & Tallam, K. (2019), 'A quantitative narrative on movement, disease and patch exploitation in nesting agent groups', *bioRxiv* .
URL: *https://www.biorxiv.org/content/early/2019/10/03/791400*

Getz, W. M. & Saltz, D. (2008), 'A framework for generating and analyzing movement paths on ecological landscapes', *Proceedings of the National Academy of Sciences* **105**(49), 19066–19071.

Gevers, D., Cohan, F. M., Lawrence, J. G., Spratt, B. G., Coenye, T., Feil, E. J., Stackebrandt, E., Van de Peer, Y., Vandamme, P., Thompson, F. L. et al. (2005), 'Re-evaluating prokaryotic species', *Nature Reviews Microbiology* **3**(9), 733.

Giotto, N., Gerard, J.-F., Ziv, A., Bouskila, A. & Bar-David, S. (2015), 'Space-use patterns of the asiatic wild ass (*equus hemionus*): complementary insights from displacement, recursion movement and habitat selection analyses', *PloS one* **10**(12), e0143279.

Goodall, V., Fatti, L. & Owen-Smith, N. (2017), 'Animal movement modelling: independent or dependent models?', *South African Statistical Journal* **51**(2), 295–315.

Grémillet, D. & Boulinier, T. (2009), 'Spatial ecology and conservation of seabirds facing global climate change: a review', *Marine Ecology Progress Series* **391**, 121–137.

Gurarie, E., Andrews, R. D. & Laidre, K. L. (2009), 'A novel method for identifying behavioural changes in animal movement data', *Ecology letters* **12**(5), 395–408.

Gurarie, E., Bracis, C., Delgado, M., Meckley, T. D., Kojola, I. & Wagner, C. M. (2016), 'What is the animal doing? tools for exploring behavioural structure in animal movements', *Journal of Animal Ecology* **85**(1), 69–84.

Gurarie, E., Fleming, C. H., Fagan, W. F., Laidre, K. L., Hernández-Pliego, J. & Ovaskainen, O. (2017), 'Correlated velocity models as a fundamental unit of animal movement: synthesis and applications', *Movement ecology* **5**(1), 13.

Hardin, P. E. & Panda, S. (2013), 'Circadian timekeeping and output mechanisms in animals', *Current opinion in neurobiology* **23**(5), 724–731.

Harel, R., Duriez, O., Spiegel, O., Fluhr, J., Horvitz, N., Getz, W. M., Bouten, W., Sarrazin, F., Hatzofe, O. & Nathan, R. (2016), 'Decision-making by a soaring bird: time, energy and risk considerations at different spatio-temporal scales', *Philosophical Transactions of the Royal Society B: Biological Sciences* **371**(1704), 20150397.

Harel, R., Horvitz, N. & Nathan, R. (2016), 'Adult vultures outperform juveniles in challenging thermal soaring conditions', *Scientific reports* **6**, 27865.

Harel, R. & Nathan, R. (2018), 'The characteristic time-scale of perceived information for decision-making: Departure from thermal columns in soaring birds', *Functional ecology* **32**(8), 2065–2072.

Harel, R., Spiegel, O., Getz, W. M. & Nathan, R. (2017), 'Social foraging and individual consistency in following behaviour: testing the information centre hypothesis in free-ranging vultures', *Proceedings of the Royal Society B: Biological Sciences* **284**(1852), 20162654.

Hays, G. C., Ferreira, L. C., Sequeira, A. M., Meekan, M. G., Duarte, C. M., Bailey, H., Bailleul, F., Bowen, W. D., Caley, M. J., Costa, D. P. et al. (2016), 'Key questions in marine megafauna movement ecology', *Trends in ecology & evolution* **31**(6), 463–475.

Hein, A. M., Carrara, F., Brumley, D. R., Stocker, R. & Levin, S. A. (2016), 'Natural search algorithms as a bridge between organisms, evolution, and ecology', *Proceedings of the National Academy of Sciences* **113**(34), 9413–9420.

Hinton, G., Deng, L., Yu, D., Dahl, G. E., Mohamed, A.-r., Jaitly, N., Senior, A., Vanhoucke, V., Nguyen, P., Sainath, T. N. et al. (2012), 'Deep neural networks for acoustic modeling in speech recognition: The shared views of four research groups', *IEEE Signal processing magazine* **29**(6), 82–97.

Hurme, E., Gurarie, E., Greif, S., Flores-Martínez, J. J., Wilkinson, G. S., Yovel, Y. et al. (2019), 'Acoustic evaluation of behavioral states predicted from gps tracking: a case study of a marine fishing bat', *Movement ecology* **7**(1), 21.

Johnson, C. J., Parker, K. L., Heard, D. C. & Gillingham, M. P. (2002), 'A multiscale behavioral approach to understanding the movements of woodland caribou', *Ecological Applications* **12**(6), 1840–1860.

Johnson, D. S., London, J. M., Lea, M.-A. & Durban, J. W. (2008), 'Continuous-time correlated random walk model for animal telemetry data', *Ecology* **89**(5), 1208–1215.

Jonsen, I. D., Myers, R. A. & Flemming, J. M. (2003), 'Meta-analysis of animal movement using state-space models', *Ecology* **84**(11), 3055–3063.

Jonsen, I. D., Myers, R. A. & James, M. C. (2006), 'Robust hierarchical state–space models reveal diel variation in travel rates of migrating leatherback turtles', *Journal of Animal Ecology* **75**(5), 1046–1057.

Jonsen, I. D., Myers, R. A. & James, M. C. (2007), 'Identifying leatherback turtle foraging behaviour from satellite telemetry using a switching state-space model', *Marine Ecology Progress Series* **337**, 255–264.

Kane, A. S., Salierno, J. D., Gipson, G. T., Molteno, T. C. & Hunter, C. (2004), 'A video-based movement analysis system to quantify behavioral stress responses of fish', *Water Research* **38**(18), 3993–4001.

Kareiva, P. & Shigesada, N. (1983), 'Analyzing insect movement as a correlated random walk', *Oecologia* **56**(2-3), 234–238.

Killick, R. & Eckley, I. (2014), 'changepoint: An r package for changepoint analysis', *Journal of statistical software* **58**(3), 1–19.

Langrock, R., King, R., Matthiopoulos, J., Thomas, L., Fortin, D. & Morales, J. M. (2012), 'Flexible and practical modeling of animal telemetry data: hidden markov models and extensions', *Ecology* **93**(11), 2336–2342.

Larsen, L. G., Eppinga, M. B., Passalacqua, P., Getz, W. M., Rose, K. A. & Liang, M. (2016), 'Appropriate complexity landscape modeling', *Earth-science reviews* **160**, 111–130.

Levin, S. A. (1992), 'The problem of pattern and scale in ecology: the robert h. macarthur award lecture', *Ecology* **73**(6), 1943–1967.

Lyons, A. J., Turner, W. C. & Getz, W. M. (2013), 'Home range plus: a space-time characterization of movement over real landscapes', *Movement Ecology* **1**(1), 2.

Marin, J.-M., Pudlo, P., Robert, C. P. & Ryder, R. J. (2012), 'Approximate bayesian computational methods', *Statistics and Computing* **22**(6), 1167–1180.

Marra, P. P., Cohen, E. B., Loss, S. R., Rutter, J. E. & Tonra, C. M. (2015), 'A call for full annual cycle research in animal ecology', *Biology letters* **11**(8), 20150552.

Matteson, D. S. & James, N. A. (2014), 'A nonparametric approach for multiple change point analysis of multivariate data', *Journal of the American Statistical Association* **109**(505), 334–345.

McCulloch, C. & Cain, M. (1989), 'Analyzing discrete movement data as a correlated random walk', *Ecology* **70**(2), 383–388.

McGarigal, K., Wan, H. Y., Zeller, K. A., Timm, B. C. & Cushman, S. A. (2016), 'Multi-scale habitat selection modeling: a review and outlook', *Landscape ecology* **31**(6), 1161–1175.

McGavin, S. L., Bishop-Hurley, G. J., Charmley, E., Greenwood, P. L. & Callaghan, M. J. (2018), 'Effect of gps sample interval and paddock size on estimates of distance travelled by grazing cattle in rangeland, australia', *The Rangeland Journal* **40**(1), 55–64.

McKenzie, H. W., Lewis, M. A. & Merrill, E. H. (2009), 'First passage time analysis of animal movement and insights into the functional response', *Bulletin of mathematical biology* **71**(1), 107–129.

Michelot, T., Langrock, R. & Patterson, T. A. (2016), 'movehmm: An r package for the statistical modelling of animal movement data using hidden markov models', *Methods in Ecology and Evolution* **7**(11), 1308–1315.

Milner-Gulland, E., Fryxell, J. M. & Sinclair, A. R. (2011), *Animal migration: a synthesis*, Oxford University Press.

Moore, E. R., Mihaylova, S. A., Vandamme, P., Krichevsky, M. I. & Dijkshoorn, L. (2010), 'Microbial systematics and taxonomy: relevance for a microbial commons', *Research in microbiology* **161**(6), 430–438.

Morales, J. M., Haydon, D. T., Frair, J., Holsinger, K. E. & Fryxell, J. M. (2004), 'Extracting more out of relocation data: building movement models as mixtures of random walks', *Ecology* **85**(9), 2436–2445.

Morelle, K., Podgórski, T., Prévot, C., Keuling, O., Lehaire, F. & Lejeune, P. (2015), 'Towards understanding wild boar *sus scrofa* movement: a synthetic movement ecology approach', *Mammal Review* **45**(1), 15–29.

Morellet, N., Bonenfant, C., Börger, L., Ossi, F., Cagnacci, F., Heurich, M., Kjellander, P., Linnell, J. D., Nicoloso, S., Sustr, P. et al. (2013), 'Seasonality, weather and climate affect home range size in roe deer across a wide latitudinal gradient within e urope', *Journal of Animal Ecology* **82**(6), 1326–1339.

Mueller, T. & Fagan, W. F. (2008), 'Search and navigation in dynamic environments–from individual behaviors to population distributions', *Oikos* **117**(5), 654–664.

Myers, J. H. (2018), 'Population cycles: generalities, exceptions and remaining mysteries', *Proceedings of the Royal Society B: Biological Sciences* **285**(1875), 20172841.

Mysterud, A., Stenseth, N. C., Yoccoz, N. G., Langvatn, R. & Steinheim, G. (2001), 'Nonlinear effects of large-scale climatic variability on wild and domestic herbivores', *Nature* **410**(6832), 1096.

Nams, V. O. (2014), 'Combining animal movements and behavioural data to detect behavioural states', *Ecology letters* **17**(10), 1228–1237.

Nathan, R., Getz, W. M., Revilla, E., Holyoak, M., Kadmon, R., Saltz, D. & Smouse, P. E. (2008), 'A movement ecology paradigm for unifying organismal movement research', *Proceedings of the National Academy of Sciences* **105**(49), 19052–19059.

Nathan, R., Spiegel, O., Fortmann-Roe, S., Harel, R., Wikelski, M. & Getz, W. M. (2012), 'Using tri-axial acceleration data to identify behavioral modes of free-ranging animals: general concepts and tools illustrated for griffon vultures', *Journal of Experimental Biology* **215**(6), 986–996.

Northrup, J. M., Avrin, A., Anderson, C. R., Brown, E. & Wittemyer, G. (2019), 'On-animal acoustic monitoring provides insight to ungulate foraging behavior', *Journal of Mammalogy* .

Odu, G. & Charles-Owaba, O. (2013), 'Review of multi-criteria optimization methods–theory and applications', *IOSR Journal of Engineering (IOSRJEN)* **3**(10), 1–14.

Owen-Smith, N. (2013), 'Daily movement responses by african savanna ungulates as an indicator of seasonal and annual food stress', *Wildlife Research* **40**(3), 232–240.

Owen-Smith, N., Fryxell, J. & Merrill, E. (2010), 'Foraging theory upscaled: the behavioural ecology of herbi-vore movement', *Philosophical Transactions of the Royal Society B: Biological Sciences* **365**(1550), 2267–2278.

Owen-Smith, N. & Goodall, V. (2014), 'Coping with savanna seasonality: comparative daily activity patterns of a frican ungulates as revealed by gps telemetry', *Journal of Zoology* **293**(3), 181–191.

Owen-Smith, N. & Goodall, V. (2019), 'Movement informatics of large mammalian herbivores across hier-archical scales: inferring biology from statistics', *Under review* .

Owen-Smith, N. & Martin, J. (2015), 'Identifying space use at foraging arena scale within the home ranges of large herbivores', *PloS one* **10**(6), e0128821.

Panzacchi, M., Van Moorter, B., Strand, O., Saerens, M., Kivimäki, I., St. Clair, C. C., Herfindal, I. & Boitani, L. (2016), 'Predicting the continuum between corridors and barriers to animal movements using step selection functions and randomized shortest paths', *Journal of Animal Ecology* **85**(1), 32–42.

Patterson, T. A., Thomas, L., Wilcox, C., Ovaskainen, O. & Matthiopoulos, J. (2008), 'State–space models of individual animal movement', *Trends in ecology & evolution* **23**(2), 87–94.

Pettorelli, N., Laurance, W. F., O'Brien, T. G., Wegmann, M., Nagendra, H. & Turner, W. (2014), 'Satellite remote sensing for applied ecologists: opportunities and challenges', *Journal of Applied Ecology* **51**(4), 839–848.

Pettorelli, N., Ryan, S., Mueller, T., Bunnefeld, N., Jedrzejewska, B., Lima, M. & Kausrud, K. (2011), 'The normalized difference vegetation index (ndvi): unforeseen successes in animal ecology', *Climate Research* **46**(1), 15–27.

Pohle, J., Langrock, R., van Beest, F. M. & Schmidt, N. M. (2017), 'Selecting the number of states in hidden markov models: pragmatic solutions illustrated using animal movement', *Journal of Agricultural, Biological and Environmental Statistics* **22**(3), 270–293.

Polansky, L., Kilian, W. & Wittemyer, G. (2015), 'Elucidating the significance of spatial memory on movement decisions by african savannah elephants using state–space models', *Proceedings of the Royal Society B: Biological Sciences* **282**(1805), 20143042.

Polansky, L., Wittemyer, G., Cross, P. C., Tambling, C. J. & Getz, W. M. (2010), 'From moonlight to movement and synchronized randomness: Fourier and wavelet analyses of animal location time series data', *Ecology* **91**(5), 1506–1518.

Preisler, H. K., Ager, A. A., Johnson, B. K. & Kie, J. G. (2004), 'Modeling animal movements using stochastic differential equations', *Environmetrics* **15**(7), 643–657.

Rahimi, S. & Owen-Smith, N. (2007), 'Movement patterns of sable antelope in the kruger national park from gps/gsm collars: a preliminary assessment', *African Journal of Wildlife Research* **37**(2), 143–152.

Rosselló-Móra, R. & Whitman, W. B. (2019), 'Dialogue on the nomenclature and classification of prokaryotes', *Systematic and applied microbiology* **42**(1), 5–14.

Runge, C. A., Martin, T. G., Possingham, H. P., Willis, S. G. & Fuller, R. A. (2014), 'Conserving mobile species', *Frontiers in Ecology and the Environment* **12**(7), 395–402.

Sapir, N., Wikelski, M., McCue, M. D., Pinshow, B. & Nathan, R. (2010), 'Flight modes in migrating european bee-eaters: heart rate may indicate low metabolic rate during soaring and gliding', *PLoS One* **5**(11), e13956.

Schuh, R. T. (2003), 'The linnaean system and its 250-year persistence', *The Botanical Review* **69**(1), 59.

Seebacher, F. & Post, E. (2015), 'Climate change impacts on animal migration', *Climate Change Responses* **2**(1), 5.

Seidel, D. P., Dougherty, E., Carlson, C. & Getz, W. M. (2018), 'Ecological metrics and methods for gps movement data', *International Journal of Geographical Information Science* **32**(11), 2272–2293.

Shepard, E. L., Wilson, R. P., Rees, W. G., Grundy, E., Lambertucci, S. A. & Vosper, S. B. (2013), 'Energy landscapes shape animal movement ecology', *The American Naturalist* **182**(3), 298–312.

Siniff, D. B. & Jessen, C. (1969), A simulation model of animal movement patterns, *in* 'Advances in ecological research', Vol. 6, Elsevier, pp. 185–219.

Spiegel, O., Harel, R., Centeno-Cuadros, A., Hatzofe, O., Getz, W. M. & Nathan, R. (2015), 'Moving beyond curve fitting: using complementary data to assess alternative explanations for long movements of three vulture species', *The American Naturalist* **185**(2), E44–E54.

Spiegel, O. & O'Farrell, S. (2018), 'Spatial orientation and time: Methods'.

Spink, A., Tegelenbosch, R., Buma, M. & Noldus, L. (2001), 'The ethovision video tracking systema tool for behavioral phenotyping of transgenic mice', *Physiology & behavior* **73**(5), 731–744.

Symonds, M. R. & Moussalli, A. (2011), 'A brief guide to model selection, multimodel inference and model averaging in behavioural ecology using akaikes information criterion', *Behavioral Ecology and Sociobiology* **65**(1), 13–21.

Takahashi, J. S., Hong, H.-K., Ko, C. H. & McDearmon, E. L. (2008), 'The genetics of mammalian circadian order and disorder: implications for physiology and disease', *Nature reviews genetics* **9**(10), 764.

Thessen, A. (2016), 'Adoption of machine learning techniques in ecology and earth science', *One Ecosystem* **1**, e8621.

Thurfjell, H., Ciuti, S. & Boyce, M. S. (2014), 'Applications of step-selection functions in ecology and conservation', *Movement ecology* **2**(1), 4.

Toledo, S., Kishon, O., Orchan, Y., Shohat, A. & Nathan, R. (2016), Lessons and experiences from the design, implementation, and deployment of a wildlife tracking system, *in* 'Software Science, Technology and Engineering (SWSTE), 2016 IEEE International Conference on', IEEE, pp. 51–60.

Torney, C. J., Hopcraft, J. G. C., Morrison, T. A., Couzin, I. D. & Levin, S. A. (2018), 'From single steps to mass migration: the problem of scale in the movement ecology of the serengeti wildebeest', *Philosophical Transactions of the Royal Society B: Biological Sciences* **373**(1746), 20170012.

Torres, L. G., Orben, R. A., Tolkova, I. & Thompson, D. R. (2017), 'Classification of animal movement behavior through residence in space and time', *PloS one* **12**(1), e0168513.

Townsend Peterson, A., Papeş, M. & Eaton, M. (2007), 'Transferability and model evaluation in ecological niche modeling: a comparison of garp and maxent', *Ecography* **30**(4), 550–560.

Tracey, J. A., Bevins, S. N., VandeWoude, S. & Crooks, K. R. (2014), 'An agent-based movement model to assess the impact of landscape fragmentation on disease transmission', *Ecosphere* **5**(9), 1–24.

Tsalyuk, M., Kilian, W., Reineking, B. & Getz, W. M. (2019), 'Temporal variation in resource selection of african elephants follows long-term variability in resource availability', *Ecological Monographs* **89**(2), e01348.

Turchin, P. (1998), *Quantitative analysis of movement*, Sinauer assoc. Sunderland (mass.).

Valletta, J. J., Torney, C., Kings, M., Thornton, A. & Madden, J. (2017), 'Applications of machine learning in animal behaviour studies', *Animal Behaviour* **124**, 203–220.

Walther, G. et al. (2009), 'Inference and modeling with log-concave distributions', *Statistical Science* **24**(3), 319–327.

Weiser, A. W., Orchan, Y., Nathan, R., Charter, M., Weiss, A. J. & Toledo, S. (2016), Characterizing the accuracy of a self-synchronized reverse-gps wildlife localization system, *in* 'Information Processing in Sensor Networks (IPSN), 2016 15th ACM/IEEE International Conference on', IEEE, pp. 1–12.

Williams, H. J., Holton, M. D., Shepard, E. L., Largey, N., Norman, B., Ryan, P. G., Duriez, O., Scantlebury, M., Quintana, F., Magowan, E. A. et al. (2017), 'Identification of animal movement patterns using tri-axial magnetometry', *Movement ecology* **5**(1), 6.

Williams, H. J., Taylor, L. A., Benhamou, S., Bijleveld, A. I., Clay, T. A., de Grissac, S., Demšar, U., English, H. M., Franconi, N., Gómez-Laich, A. et al. (2019), 'Optimising the use of bio-loggers for movement ecology research', *Journal of Animal Ecology* .

Wilmers, C. C., Nickel, B., Bryce, C. M., Smith, J. A., Wheat, R. E. & Yovovich, V. (2015), 'The golden age of bio-logging: how animal-borne sensors are advancing the frontiers of ecology', *Ecology* **96**(7), 1741–1753.

Wittemyer, G., Northrup, J. M. & Bastille-Rousseau, G. (2019), 'Behavioural valuation of landscapes using movement data', *Philosophical Transactions of the Royal Society B* **374**(1781), 20180046.

Wittemyer, G., Polansky, L., Douglas-Hamilton, I. & Getz, W. M. (2008), 'Disentangling the effects of forage, social rank, and risk on movement autocorrelation of elephants using fourier and wavelet analyses', *Proceedings of the National Academy of Sciences* pp. pnas–0801744105.

Worton, B. (1987), 'A review of models of home range for animal movement', *Ecological modelling* **38**(3-4), 277–298.

Yerushalmi, S. & Green, R. M. (2009), 'Evidence for the adaptive significance of circadian rhythms', *Ecology letters* **12**(9), 970–981.

Zeller, K. A., McGarigal, K. & Whiteley, A. R. (2012), 'Estimating landscape resistance to movement: a review', *Landscape ecology* **27**(6), 777–797.

Zhang, J., Hull, V., Ouyang, Z., He, L., Connor, T., Yang, H., Huang, J., Zhou, S., Zhang, Z., Zhou, C. et al. (2017), 'Modeling activity patterns of wildlife using time-series analysis', *Ecology and evolution* **7**(8), 2575–2584.

Zhang, Z., Geiger, J., Pohjalainen, J., Mousa, A. E.-D., Jin, W. & Schuller, B. (2018), 'Deep learning for environmentally robust speech recognition: An overview of recent developments', *ACM Transactions on Intelligent Systems and Technology (TIST)* **9**(5), 49.

Zhao, K. & Jurdak, R. (2016), 'Understanding the spatiotemporal pattern of grazing cattle movement', *Scientific reports* **6**, 31967.

Zidon, R., Garti, S., Getz, W. M. & Saltz, D. (2017), 'Zebra migration strategies and anthrax in etosha national park, namibia', *Ecosphere* **8**(8), e01925.

Zucchini, W., MacDonald, I. L. & Langrock, R. (2016), *Hidden Markov models for time series: an introduction using R*, Chapman and Hall/CRC.