1 **Low atmospheric CO$_2$ induces nocturnal carbon accumulation in the lycophyte**
2 **genus *Isoëtes***
3

Authors: Jacob S. Suissa[1,2*] & Walton A. Green[1]

Affiliations:
[1]Department of Organismic and Evolutionary Biology, Harvard University, Cambridge, MA
[2]Arnold Arboretum of Harvard University, Boston, MA
*Correspondence to: jsuissa@g.harvard.edu

Abstract
Metabolic shifts play an essential role in the survival of plants in extreme habitats. Certain
desert plant species, for example, avoid water loss by temporally segregating the light and
dark reactions of photosynthesis. By only opening their stomata at night these plants
inhibit significant water loss during the day. At night CO$_2$ is incorporate into 4-carbon
acids for subsequent daytime fixation[1-5]. This behavior is known as Crassulacean Acid
Metabolism (CAM) and leads to noticeable diel cycles in pH of photosynthetic organs[1-5].
Oddly, similar acidity cycles are found in the submerged aquatic lycophyte genus *Isoëtes*[2,6-
12], which are not water limited in any sense. It has long been assumed that their nocturnal
CO$_2$ accumulation is an adaptation to low daytime carbon levels in aquatic ecosystems, but
this has never been empirically tested[2,6-13]. Here, we offer direct evidence that CO$_2$
starvation induces CAM-like nocturnal carbon accumulation in terrestrial *Isoëtes*.
Populations of terrestrial *Isoëtes engelmannii* grown in climate-controlled chambers and
starved of atmospheric CO$_2$ during the day displayed diel acidity cycles similar to those in
both xerophytic CAM plants and submerged *Isoëtes* species. These results substantiate the
hypothesis that carbon starvation provides a selective pressure for nocturnal carbon
accumulation in *Isoëtes*. Furthermore, while aquatic CO$_2$ levels undoubtedly promote
nocturnal carbon accumulation in extant *Isoëtes*, the induction of this behavior in *terrestrial*
plants suggests a possible earlier terrestrial evolution of this metabolism in response to low
***atmospheric* CO$_2$ levels during the Carboniferous period[14-17]. Our findings therefore both**
provide empirical support for a long-standing assumption about nocturnal carbon
accumulation in the lycophyte lineage[2,6-12,18] and suggest an earlier evolution of this
behavior, leading to the notion that CAM in xerophytes may only represent a subset of
metabolisms that employ nocturnal carbon accumulation in response to variable
environmental pressures.

35
Plants inhabit some of the harshest terrestrial environments on earth. In deserts the
metabolic pathway known CAM evolved in some lineages to prevent dessication[1-5,13,18]. This
metabolism is critical for the success of many xerophytic plants and has been regarded as one of
the most important adaptations to dry environments[1-5]. Although CAM in xerophytic plants
allows for water conservation in dry environments, the pathway behind this metabolism (C4,
Hatch/Slack/Korshak pathway) is fundamentally a carbon concentrating mechanism which
increases the selectivity of rubisco by altering the CO$_2$:O$_2$ ratio within the chloroplast[5,18].
43
In 1981, diel acidity cycles similar to CAM were first observed in the aquatic lycophyte
genus *Isoëtes*[6]. This metabolic process was named 'aquatic CAM' to highlight its similarity to
xerophytic CAM acidity cycles. Here we instead will use the term 'nocturnal carbon

accumulation' (NCA) to emphasize the differences between the behavior described in *Isoëtes*
and that in xerophytic CAM plants. Further investigations of NCA in *Isoëtes* have demonstrated
a high degree of plasticity. For example, *Isoëtes howellii* exposed to atmospheric CO_2 levels
while growing terrestrially tend not to accumulate carbon nocturnally, while *I. karstenii*
accumulates carbon regardless of environment, demonstrating that this behavior can be
facultative or constitutive[7-9]. Since CAM serves as a carbon concentrating mechanism, CAM-like
NCA in *Isoëtes* has been assumed to be a response to low carbon availability in aquatic
ecosystems[2,6-9,12,18]. In eutrophic lakes hypocarbia is present only during the day because of
diurnal photosynthesis and nocturnal respiration of aquatic algae and macrophytes; in
oligotrophic lakes hypocarbia is generally continuous[10-12]. But, the theory that nocturnal carbon
accumulation in aquatic plants is an adaptation to carbon limitation has hitherto remained largely
hypothetical and correlative[6-12]. Furthermore, NCA in *Isoëtes* has been regarded as an adaptation
by extant lineages to these selection pressures[7,18], rather than, as we suggest, an adaptation early
in this ancient lineage, inherited by recent descendants.
61
To test the hypothesis that NCA is a direct response to carbon starvation, we grew
terrestrial plants of two *Isoëtes* species in environmentally controlled growth chambers, starved
plants of atmospheric CO_2 during the day and sampled plant pH from multiple plants for 24-
hours. Diurnal hypocarbia induced diel acidity cycles in terrestrial *Isoëtes engelmannii* like those
observed in xerophytic CAM plants and submerged aquatic *Isoëtes* (Fig. 4c). This provides
direct evidence that CO_2 starvation can induce CAM-like NCA in *Isoëtes*, and supports the
hypothesis that CO_2 limitation is a selective pressure on this behavior in extant *Isoëtes* species[7-
11,13,18]. Moreover, the induction of NCA in terrestrial *Isoëtes* by atmospheric CO_2 starvation may
support the hypothesis that CAM-like photosynthesis evolved early in terrestrial Isoetalean
lineages in the Carboniferous in response to globally low atmospheric CO_2 levels, not more
recently in response low aquatic CO_2 levels[2,7-11,13,18].
73
In our first series of field measurements from May to September, we measured morning
and evening pH in *Isoëtes engelmannii* leaves, roots, and corms, monthly. (Fig. 1a; Fig. 2). We
found that in the field, submerged plants of *I. engelmannii* accumulated carbon on a diel cycle
(Fig. 2; Fig 3). The mean morning pH of the leaves throughout the summer was 3.71 and the
mean evening pH was 4.90 (Fig. 2). Non-photosynthetic organs (roots and corms) did not
accumulate carbon nocturnally (Fig. 3); corms: mean morning pH 5.94, mean evening pH 5.97;
roots: mean morning pH 6.32, mean evening pH 6.36 (Fig. 3). Upon recession of the shoreline in
July, many of the plants were left growing terrestrially, exposed to atmospheric conditions (Fig.
1). In the plants growing terrestrially, pH had higher variability (Fig. 1d; Fig 3), the difference
between morning and evening pH was insignificant (Fig. 2). This supports prior observations
that *I. engelmannii* accumulates carbon nocturnally only when submerged[7].
85
For lab manipulations, we collected *I. engelmannii* from the same population measured in
the field as wells as specimens of *I. tuckermanii,* a related species that generally grows
completely submerged. All plants collected were brought into the lab and cultivated in climate-
controlled growth chambers. Under ambient CO_2 levels the diel pH variation of submerged
individuals of *I. engelmannii* was similar to submerged individuals in the field (Fig. 4a); leaf
mean morning pH was 4.58, and mean evening pH was 5.81. Submerged plants of *I. tuckermanii*
had a less dramatic diel shift in pH, compared to *I. engelmannii* (Fig. 4a, 4d), mean morning pH

was 3.89, and mean evening pH was 4.70. When emergent, as expected from the field
experiments, individuals of *I. engelmannii* no longer displayed a diel shift in pH; (Fig 4b; Fig. 2);
mean morning pH was 5.31, mean evening pH was 5.46. *Isoëtes tuckermanii,* however, showed a
more dramatic diel change in pH, when emergent (Fig. 4e); mean morning pH was 4.15, mean
evening pH was 5.84. These results demonstrate different NCA behavior in these two species, as
has been previously documented in *Isoëtes*[7]. *I. engelmannii* induces nocturnal carbon
accumulation when submerged but not when emergent, while *I. tuckermanii* demonstrated
constitutive NCA, with similar diel variation in pH irrespective of water depth, and only a slight
increase in NCA when emergent compared to when submerged.
102
We next grew the plants with diurnal CO2 starvation and nocturnal enrichment to mimic
the CO2 conditions in a eutrophic lake. *Isoëtes engelmannii* individuals grown under these
conditions again showed a diel cycle in pH similar to that observed in the submerged specimens
in lab and field (Fig. 2; 3a; 3c). For instance, in two independent experiments, we measured a
mean morning pH of 4.69, mean evening pH was 6.37 (Exp. 8); mean morning pH 4.85, mean
evening pH 6.18. The magnitude of pH change in the terrestrial plants grown under diurnal CO_2
starvation was similar to that in submerged plants in the field experiments, though slightly
dampened (Fig. 3; Fig 4c). This may be due to the fact that predawn CO_2 concentrations in the
field in eutrophic lakes and ponds can exceed 2500ppm[10], and the maximum CO_2 enrichment
during our experiment approximated ambient atmospheric concentrations of 400 ppm. (Fig. 3c).
113
When grown under these manipulated CO_2 conditions, *I. tuckermanii* continued to show
nocturnal carbon accumulation (Fig. 1c; Fig. 3f) with a mean morning pH of 4.12, and a mean
evening pH of 5.98. The pH fluctuation of *I. tuckermanii* during diurnal hypocarbia did not differ
from that of the terrestrially growing plants in ambient CO_2 (Fig. 3e), suggesting that this species
is obligate in its NCA. Unlike *I. engelmannii*, the individuals of *I. tuckermanii* we examined had
no stomata, a possible explanation of this constitutive behavior. These results provide a possible
explanation for the plasticity of NCA in the genus *Isoëtes* observed in the past[7,12]. Moreover, we
show that laboratory-based experimental modification of natural conditions can replicate field
conditions and that carbon starvation imposed on terrestrial plants in the lab produces a diel
change in pH comparable to the change observed in submerged aquatic plants.
124
Prior discussions of the adaptive significance of NCA in *Isoëtes* have focused on aquatic
CO_2 limitation as a selective force[2,6-12,18]. Our experiments now show that terrestrial carbon
limitation has the same effect. Low atmospheric CO_2 was a notable feature of the
Carboniferous,[15-17] a time when Isoetalean lycopsids formed a substantial part of the terrestrial
flora. In the Carboniferous atmosphere[14-17], a carbon concentrating mechanism that increased
carbon gain and minimized the photorespiratory loss of energy and carbon in a high oxygen/low
CO_2 atmosphere would have been particularly advantageous. Given that we induced NCA in
terrestrial *Isoëtes* (Fig 3c), and that this behavior is observed in many *Isoëtes* species[7,12,18], it
seems likely that NCA evolved in the early Isoetalean lycopsids during the low atmospheric CO_2
conditions of the Carboniferous period. If so, NCA in extant aquatic *Isoëtes* would represent an
exaptation of a much older metabolic behavior rather than a more recent adaptation to low
aquatic CO_2 levels. Final determination of the evolutionary timing of NCA in *Isoëtes* requires
more complete consideration of the temporal variation and selective pressures of local aquatic
and global atmospheric CO_2 levels, as well as the distribution of this behavior in extant *Isoëtes*

species. Quantification of NCA at the species level across the genus *Isoëtes* will require further
phylogenetic and physiological work[7,18,20]. Examining a broader range of *Isoëtes* species and
using phylogenetic comparative methods may reveal varied appearances in NCA throughout the
extant genus[7,18], just as CAM has been found to evolve many times in xerophytic plants[13,19].
143
Our results presented here do support the hypothesis that NCA in *Isoëtes* was an early
evolutionary behavior, not a specialized or derived adaptation[7,18]. NCA appears throughout the
vascular plant tree, including among ferns, lycophytes, and angiosperms and may have evolved
as much as 350 million years before CAM appeared late in the Miocene[13]. Therefore, it seems
reasonable to consider NCA to be the major photosynthetic pathway for carbon concentration of
which CAM is a more recent offshoot representing an adaptation to dry environments. This view
seems to better reflect the evolutionary sequence and significance of these behaviors.

1.   M. Kluge, I. P. Ting, Crassulacean acid metabolism: an ecological analysis. *Ecological*
studies series. **30**, 209 (1978).
2.   Keeley, J. E. Aquatic CAM Photosynthesis. in *Crassulacean Acid Metabolism:*
Biochemistry, Ecophysiology and Evolution (eds. Winter, K. & Smith, J. A. C.) 281–295
(Springer Berlin Heidelberg, 1996).
3.   D. L. Sipes, I. P. Ting, Crassulacean Acid Metabolism and Crassulacean Acid Metabolism
Modifications in Peperomia camptotricha. *Plant Physiol.* **77**, 59–63 (1985).
4.   U. Lüttge, Ecophysiology of Crassulacean Acid Metabolism (CAM). *Ann. Bot.* **93**, 629–
652 (2004).
5.   E. J. Edwards, Evolutionary trajectories, accessibility and other metaphors: the case of C4
and CAM photosynthesis. *New Phytol.* **223**, 1742–1755 (2019).
6.   J. E. Keeley, *Isoetes howellii*: A submerged aquatic CAM plant? *Am. J. Bot.* **68**, 420–424
(1981).
7.   J. E. Keeley, CAM photosynthesis in submerged aquatic plants. *Bot. Rev.* **64**, 121–175
(1998).
8.   J. E. Keeley, Crassulacean acid metabolism in the seasonally submerged aquatic Isoetes
howellii. *Oecologia*. **58**, 57–62 (1983).
9.   J. E. Keeley, R. P. Mathews, C. M. Walker, Diurnal acid metabolism in Isoetes howellii
from a temporary pool and permanent lake. *Am. J. Bot.* **70**, 854–857 (1983).
10.   J. E. Keeley, G. Bowes, Gas Exchange Characteristics of the Submerged Aquatic
Crassulacean Acid Metabolism Plant, *Isoetes howellii*. *Plant Physiol.* **70**, 1455–1458
(1982).
11.   J. E. Keeley, Photosynthetic pathways in freshwater aquatic plants. *Trends Ecol. Evol.* **5**,
330–333 (1990).
12.   J. E. Keeley, Distribution of diurnal acid metabolism in the genus *Isoetes*. *Am. J. Bot.* **69**,
254–257 (1982).

13.   E. J. Edwards, R. M. Ogburn, Angiosperm Responses to a Low-CO2 World: CAM and C4
Photosynthesis as Parallel Evolutionary Trajectories. *Int. J. Plant Sci.* **173**, 724–733
(2012).
14.   W. A. Green, The function of the aerenchyma in arborescent lycopsids: evidence of an
unfamiliar metabolic strategy. *Proc. Biol. Sci.* **277**, 2257–2267 (2010).
15.   D. J. Beerling, Low atmospheric CO2 levels during the Permo- Carboniferous glaciation
inferred from fossil lycopsids. *Proc. Natl. Acad. Sci. U. S. A.* **99**, 12567–12571 (2002).
16.   M. Steinthorsdottir, A. J. Jeram, J. C. McElwain, Extremely elevated CO2 concentrations
at the Triassic/Jurassic boundary. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* **308**, 418–432
(2011).
17.   D. G. Van Der Meer, R. E. Zeebe, D. J. J. van Hinsbergen, A. Sluijs, W. Spakman, T. H.
Torsvik, Plate tectonic controls on atmospheric CO2 levels since the Triassic. *Proc. Natl.*
Acad. Sci. U. S. A. **111**, 4380–4385 (2014).
18.   Keeley, J. E. & Rundel, P. W. Evolution of CAM and C4 carbon-concentrating
mechanisms. *Int. J. Plant Sci.* **164**, S55–S77 (2003).
19.   L. P. Hancock, J. A. M. Holtum, The evolution of CAM photosynthesis in Australian
Calandrinia reveals lability in C3+ CAM phenotypes and a possible constraint to the
evolution of strong CAM. *Integrative and comparative biology.* icz089 (2019)
20.   Pereira, J., Labiak, P. H. & Stützel, T. Origin and biogeography of the ancient genus Isoëtes
with focus on the Neotropics. *Bot. J. Linn. Soc.* **185**, 253–271 (2017).

203
204

Acknowledgments: The authors would like to thank Katharine E. Black, Sylvia P. Kinosian and
Weston L. Testo for their helpful edits on the manuscript. We thank the Edmund Niles Huyck
Preserve for funding, access to plant materials, and lodging during the field experiment, as well as
the growth facilities staff at Weld Hill Research Building of the Arnold Arboretum of Harvard
University for assisting in plant cultivation and growth chamber accommodations.

Author Contributions: JSS developed the project, JSS and WAG executed the experiments,
WAG analyzed the data, and JSS and WAG wrote the manuscript.

213
Methods

Field Experiments: The Edmund Niles Huyck Preserve (hereafter ENHP) is located in
Rensselaerville, NY, 425–525m elevation. Plants of *Isoëtes engelmanii* were studied from May-
September in a population on the north shore of Lake Myosotis[21], 42°31'26.8" N 74°9'7.2" W.
Three to eight samples composed of 5-6 plants were collected five times during the summer of
2018 from May-September (Fig. 2). During May, June, and July samples were collected only in
the morning at 0600 and in the evening at 1800; in August 3-8 samples were collected every
three hours from 0600--1800 and in September collections were made every three hours for a full
24 hours (1800--1800). All plants were cleaned of algae and soil in DI water in the lab, which
was located 1km from the study site.

In the lab, leaves, corms, and roots were separated using scalpels and forceps and organs were
blotted dry. At each time point leaves from 5-6 individuals of *I. engelmanii* were randomized and
separated into 3-8 distinct samples. Leaves from each of the 3-8 samples were macerated using
plastic dowels in 1.7mL Eppendorf tubes and 0.5mL of DI H2O was added to each sample.

Samples were resuspended by mixing using a vortex mixer at maximum intensity for 10 seconds,
followed by centrifugation at 10,000 rpm for 10 seconds. The supernatant was carefully
extracted using a pipette, and placed on the Horiba LAQUAtwin pH-22 meter, for pH readings.
The pH was measured twice from each sample to ensure accuracy of the reading. The pH meter
was cleaned with deionized water and allowed to drt between each sample. The pH meter was
recalibrated between each time point using pH 4 and 7 standardized buffers.
236
Isoëtes engelmannii specimens collected in the field throughout summer 2018 in Lake Myosotis
were cultivated in growth chambers in the Weld Hill Research Building of the Arnold Arboretum
of Harvard University. In addition, populations of *I. tuckermanii* already growing in the
greenhouse (collected from the NW shore of Lake Mattawa, in Orange, MA, 42°34'11.9" N
72°19'34.1" W), were used in the growth chamber experiments. For 2 months, plants were grown
fully submerged at ambient CO_2 levels (~400ppm), at 20˚C, with a 12-hour photoperiod,
15μmolm$_{-2}$ photosynthetically active radiation. While submerged pH was measured on a diel
cycle following the field protocol (see above). Containers were then drained of water, and the
plants were allowed to acclimate for 1-3 days. While plants were terrestrial under ambient CO_2
levels pH was measured every 3-5 hours for 24 hours following the aforementioned protocol.
Plants were then moved to a growth chamber set to a diel cycle of atmospheric CO_2: 100ppm of
CO_2 during the 12-hour photoperiod (the minimum obtainable in the chamber) and 400ppm
(equivalent to ambient atmospheric levels) during dark period. Temperature and light intensity
were not changed from the ambient CO_2 conditions. Starting at 0700, plants were harvested
every 3--5 hours for 24 hours. At each time point leaves from 2-4 individuals of were
randomized and separated into 2-3 distinct samples and pH was measured following the field
protocol.
254
255   **Data and materials availability:** all data can be found in the Supplemental Information.
256
257

258   21.   N. H. Russell, The Vascular Flora of the Edmund Niles Huyck Preserve, New York. *Am.*
259   *Midl. Nat.* **59**, 138–145 (1958).

260
261
262
263

264
265

Figure 1: **a**: Lake Myosotis study site during average water levels for most of the year. **b**: Lake Myosotis in July when water levels drop. **c** exposed plants of *I. tuckermanii* in lab conditions. **d**: exposed plants of *I. engelmanii in situ*.

Figure 2: Field experiments. Diel pH measurements of leaves from field grown plants throughout the summer months. In July some plants were growing terrestrially and did not show significance differences in pH. Each point represents a single pH measurement of 3-8 samples comprised of pooled leaves from 5-6 individuals, pH was measured twice on each sample. During September more than two time points were measured (see Fig. 3).

Figure 3: Field experiments. pH measurements of leaves, corms and roots, for 24-hours in September. A clear cycle in pH is demonstrated in leaves while corms and roots show no diel change. Each point represents a single pH measurement of 3-8 samples comprised of pooled leaves from 5-6 individuals, pH was measured twice on each sample.

 Figure 4: Laboratory experiments. Leaf pH measurements of *I. engelmanii* and *I. tuckermanii* under, **a, d**. submerged ambient CO_2 levels; **b, e**. terrestrial ambient atmospheric CO_2 levels; and **c, f**. terrestrial CO_2 manipulation. A clear diel change in pH was not observed in *I. engelmanii* when emergent but was induced upon CO_2 starvation. Solid circles represent submerged plants and open circles represent terrestrial plants. Each point represents a single pH measurement of 2-3 samples comprised of pooled leaves from 2-4 individuals, pH was measured twice on each sample.
 *Circles in pane **a** are measurements from the field, crosses represent lab experiments. In pane **c** two separate 24-hour experiments were conducted (crosses and circles).