Pollen-feeding *Heliconius* butterflies host distinctive adult-stage microbiomes

Tobin J. Hammer[1*], Jacob C. Dickerson[2,3], W. Owen McMillan[2], Noah Fierer[3,4]

1. Department of Integrative Biology, University of Texas at Austin, Austin, TX 78712, USA

2. Smithsonian Tropical Research Institute, Apartado 0843-03092, Panama City, Panama

3. Department of Ecology & Evolutionary Biology, University of Colorado at Boulder, Boulder, CO 80309, USA

4. Cooperative Institute for Research in Environmental Sciences, University of Colorado at Boulder, Boulder, CO 80309, USA

*Corresponding author: tjhammer@gmail.com

Keywords: symbiosis, gut bacteria, Lepidoptera, microbiota, parasites

Abstract

Evolutionary transitions in animal diets often coincide with shifts in the microbiome, but

the degree to which diet-microbiome feedbacks vary across host taxa and development is

unresolved. We examined these potential feedbacks from the perspective of Lepidoptera

(butterflies and moths), a diverse clade in which little is known about adult-stage microbial

associations. With the exception of *Heliconius* butterflies, most lepidopteran adults are short-

lived and either feed on simple substrates, like nectar, or do not feed at all. *Heliconius* consume

pollen as adults, which provides amino acids and allows the butterflies to have an extended

lifespan. Using 16S rRNA gene sequencing of 214 field-collected individuals, we found that

adult passion-vine butterfly microbiomes exhibited a strong signal of host phylogeny, with a

clear distinction between *Heliconius* and non-pollen-feeding relatives. This pattern was largely

driven by differing relative abundances of bacterial phylotypes shared among host taxa, as

opposed to the presence or absence of host-specific phylotypes. Using shotgun metagenomic

sequencing, we also discovered trypanosomatids and microsporidia to be prevalent in butterfly

guts, suggesting potential interactions with co-localized gut bacteria. Overall, we show that a

major transition in adult-stage lepidopteran diet and life history coincides with a shift in

microbiomes, and our work provides a foundation for future tests of microbiome function in

adult butterflies.

Introduction

A major goal of contemporary microbial ecology is to identify the factors that influence

the structure and function of host-associated microbiomes. In animals, diet has emerged as a

consistently strong predictor of microbiome structure [1–3]. In some cases, microbes are also

known to directly facilitate evolutionary transitions in host diet through their contributions to

digestion, nutrient synthesis, or detoxification [4–6]. However, there are many animal taxa for

which comparative microbiome data are either unavailable or only available for a specific phase

of their development. This makes it difficult to generalize diet-microbiome interactions, as both

microbiomes and diets often change radically during host development [7, 8].

Lepidoptera (butterflies and moths) are a diverse and ecologically important group in

which microbiome roles are likely to be highly specific to certain host taxa and life stages. For

lepidopteran larvae (caterpillars), even major shifts in larval diet do not coincide with

functionally relevant shifts in the microbiome [9–11]. Lepidopteran larvae typically harbor a

very low number of gut microbes, and most of these microbes are transient with the exception of

pathogen infections [9]. However, the microbiomes of insects that undergo complete

metamorphosis, such as Lepidoptera, often exhibit profound shifts in size, composition and

function across host life stages [7]. A recent survey of various butterfly species found that adult-

stage gut bacterial communities were generally abundant (median ~10^8 16S rRNA gene copies

per gut) and distinct from diet-associated bacterial communities [12]. It is possible that

evolutionary transitions in the diet of adult lepidopterans could cause, or be caused by,

concomitant transitions in adult-stage microbiomes.

Arguably, the single most dramatic and consequential transition in adult diet among the

150,000+ described species of Lepidoptera [13] occurred in the ancestors of neotropical

Heliconius butterflies (Nymphalidae: Heliconiini). While adults of other lepidopterans either do

not feed at all, or feed on comparatively nitrogen-poor and/or inconsistently available substrates

[14], *Heliconius* evolved the ability to consume pollen, an abundant resource rich in essential

amino acids [15]. Directly or indirectly, pollen-feeding led to a suite of changes in *Heliconius*

population structure, behavior, chemical ecology, mimicry, coevolutionary interactions with

plants, and life history traits [15–18]. As an example of the latter, *Heliconius* are exceptional

among butterflies in their adult longevity, with a potential lifespan of several months [18–20].

Heliconius have been an influential model system in evolutionary biology for over 150

years [17]. As with adult butterflies in general, however, very little is known about *Heliconius*

microbiomes. Data for a handful of *Heliconius* individuals and species are available [12, 21, 22],

but a comparative context with non-pollen-feeding members of the passion-vine butterfly tribe

Heliconiini has been missing. Nectar- and fruit-feeding butterfly species were recently found to

host distinct gut microbiomes [12], supporting the possibility that other kinds of shifts in adult

butterfly diets could be linked to shifts in microbiomes.

Does the ability to feed on pollen coincide with the presence of novel symbionts, changes

in community composition, or other microbiome differences? To answer this question, we

collected 214 wild adult butterflies representing 23 species and subspecies across the Heliconiini

and characterized their microbiomes. We used amplicon sequencing targeting a portion of the

16S rRNA gene and assessed the distribution of exact sequence variants (ESVs) across host

individuals, species, genera, and the heliconiine phylogeny. We evaluated the reliability of our

amplicon sequencing approach by obtaining shotgun metagenomic data from a subset of

samples. We also used the metagenomes to assemble near-full-length 16S rRNA genes and

evaluate bacterial strain-level patterns that may not be evident from analyses of amplicon

sequence data. Finally, we uncovered prevalent gut-associated trypanosomatids and

microsporidia. These putatively parasitic microeukaryotes have been little-studied in butterflies,

despite their potential interactions with gut bacteria and relevance to host fitness.

Materials and Methods

Field collections

The wild adult butterflies used for whole-body microbiome sequencing were collected

from seven locations in Panama and Ecuador in May-August 2014 (more detail is provided in the

supplemental file "Collection_localities.txt"). Butterflies were euthanized with ethyl acetate and

stored in DMSO after removal of wings, following [23]. We also stored two DMSO-only blanks

to use as negative controls. In June 2016, we collected additional adult butterflies for gut and

head/thorax sequencing from Gamboa and Pipeline Road, Panama. For these specimens, we

dissected the gut (hindgut, midgut, and the distal ~1/2 of the foregut) using sterilized tools prior

to storage in DMSO. The whole head and thorax (including the proximal foregut) were stored

separately. Species or subspecies were identified based on morphology. Butterflies were

collected under permit # SC/A-7-11 from Panama's Autoridad Nacional del Ambiente and #

005-13 IC-FAU-DNB/MA from Ecuador's Ministerio del Ambiente.

Sample processing, PCR and sequencing

We removed whole bodies and head/thorax samples from DMSO and, after

homogenization, used approximately 50 mg subsamples of homogenate for DNA extractions

with the MoBio PowerSoil kit following the manufacturer's instructions. We added entire guts

directly to DNA extraction tubes, in which they were homogenized during the first bead-beating

step of the protocol. Two DMSO blanks and 30 DNA extraction blanks were also processed in

tandem with the butterfly samples and sequenced.

96   PCR amplifications (515F/806R primers, V4 region) and 2 X 150 bp Illumina MiSeq

sequencing of 16S rRNA genes followed standard Earth Microbiome Project protocols

(dx.doi.org/10.17504/protocols.io.nuudeww). For 29 butterfly individuals of nine species, gut

and head/thorax samples were also amplified using primers that target the ITS gene region of

fungi [24]. Amplification success with these fungal-specific primers (as estimated from gel

electrophoresis) was generally very low, suggesting a lack of abundant fungal DNA that was

later corroborated with the shotgun metagenomic data (see below). DNA extracts from a subset

of 15 amplicon-sequenced gut samples were used for shotgun metagenomic sequencing

following the approach described previously [25] with an input DNA concentration of 0.75

ng/ul, KAPA HiFi HotStart ReadyMix and bead cleanup with Ampure XP beads at a 0.9x ratio.

Amplicon data processing

109   Amplicons from the 2014 whole-body samples and 2016 gut and head/thorax samples

were sequenced on separate runs, demultiplexed using idemp (https://github.com/yhwu/idemp),

and combined for futher processing. Cutadapt [26] was used to remove primer sequences. We

then used the DADA2 pipeline [27] to quality-filter (max EE value = 1) and trim (150 bp

forward, 140 bp reverse) reads, infer exact sequence variants (ESVs), merge paired-end reads,

and remove chimeras. We classified ESVs using the RDP Naive Bayesian Classifier algorithm

[28] against the SILVA training set v. 132 [29].

116   Further data processing and analyses were conducted in R v. 3.6.0 [30]. We used

decontam [31] for prevalence-based identification of putative contaminant ESVs based on 34

negative controls (DMSO and DNA extraction blanks and PCR no-template controls). The

median percentage of contaminant sequences across butterfly samples was 0.09%, but two

samples with >10% contaminants were removed from further analysis. ESVs with < 100 total

sequences across all samples (out of a combined total of 5.6 million sequences) were removed,

as were ESVs classified as mitochondria or chloroplast, or bacteria lacking sub-domain

identification. These ESVs combined typically made up a low proportion of reads from the

libraries (median of 2.6% across all samples).

As the resulting 16S rRNA gene amplicon datasets were highly variable in read depth

across samples, we rarefied all libraries to 5,000 sequences, filtering out 13 samples with lower

sequence depth. We also relabeled *Pantoea, Erwinia, Kluyvera, Citrobacter, Klebsiella*, and

Cronobacter taxonomic assignments to *Enterobacter*. This step was taken as genera within

Enterobacteriaceae are often polyphyletic and are difficult to resolve from short 16S rRNA gene

regions (e.g., [32]), and we wanted to avoid spurious separation of ESVs among genera and

resulting idiosyncracies in genus distributions across butterflies.

Amplicon data analysis

Beta diversity statistics and plots are based on Bray-Curtis dissimilarities and/or

UniFrac distances (weighted and unweighted). To obtain a bacterial phylogeny for the latter, we

used the fragment-insertion method [33] to place our ESV sequences into the Greengenes

reference tree [34]. The butterfly phylogeny is from [35] (TreeBASE #Tr77496). [Note that

although this phylogeny nests *aoede* within *Heliconius*, the phylogenetic position of this species

is held by some to be uncertain (J. Mallet, pers. comm.)]. Four of the butterfly species in our

sample set contained specimens from two distinct subspecies (e.g, *Heliconius sara sara* and

Heliconius sara magdalena). To include these in the species-level host phylogeny, we inserted

subspecies tips halfway along the terminal branches to their sister subspecies. Hereafter we refer

to these subspecies as "species" for simplicity.

To test for host-phylogenetic signal in microbiomes, we used Mantel tests with 9999

permutations to calculate the correlation between microbial community dissimilarities/distances

and host phylogenetic distances [36]. Intraspecific variation in microbiomes was handled by

averaging the pairwise dissimilarities/distances between all individuals of one species and all

individuals of another species. We used the phytools package [37] to visualize concordance

between topologies of the host phylogeny and a dendrogram of bacterial community

dissimilarities. Nodes were rotated with the "cophylo" function in phytools to maximize tip

matching between the two trees.

Differences in overall community composition between host genera were tested with

PERMANOVA as implemented in the vegan package [38]. Using the "betadisper" function we

corroborated that significant test results were due to host genus-level differences in location and

not dispersion [39]. We used a nonparametric statistical test (Wilcoxon rank-sum) to identify

bacterial genera that differed in relative abundance between host taxa or between sample types

(gut versus head/thorax) and applied a false discovery rate correction to the resulting p values.

Metagenome data processing and analysis

For 15 gut samples, we obtained shotgun metagenomic data to complement the

bacterial 16S rRNA gene amplicon dataset. We quality-filtered these reads with sickle [40] and

trimmed adapters with cutadapt [26]. We then used Bowtie 2 [41] to filter out reads matching a

given sample's corresponding host species' genome, obtained from Lepbase [42]. The two

Dryadula phaetusa metagenomes were mapped to a genome of the sister species *Dryas iulia* as

no *Dryadula* genome was available. Since there was a high proportion of host-derived reads, we

focused here on describing microbial diversity using ribosomal RNA gene reads present in the

metagenomes. With the host-filtered reads, we used phyloFlash [43] to find and classify

eukaryotic and bacterial SSU rRNA reads. Bacterial community composition was compared

between the amplicon and shotgun metagenomic datasets using a Mantel test. We also used

phyloFlash to assemble 16S rRNA genes from the 150 bp shotgun reads. These longer sequences

allowed us to estimate the phylogeny of *Orbus*, the dominant bacterium in these 15 samples.

Orbus sequences were aligned with MUSCLE [44], curated with Gblocks [45], and used for

maximum likelihood reconstruction with the phylogeny.fr implementation [46] of PhyML [47].

Data availability

Amplicon data, metadata, and R code are available from figshare

(figshare.com/projects/Heliconius_butterfly_microbiomes/70520). Metagenomes are available

from MG-RAST (project no. MGP89563).

Results

Adult heliconiine butterfly microbiomes have a relatively low diversity, with a median

of 26 bacterial ESVs per individual. A median of 11 ESVs accounted for 95% of the reads.

These communities are also fairly consistent among conspecific butterflies. For example, within

our most deeply sampled population (*H. erato petiverana* in Gamboa, Panama; N = 23), a

median of 84% of the 16S rRNA gene reads obtained from a given individual's microbiome was

assigned to 10 dominant bacterial genera (Fig. S1). Some of these genera are present in roughly

similar relative abundances across individuals and even across our two sampling years.

Microbiomes from whole, homogenized butterfly bodies are mainly composed of gut-

associated taxa. Isolated guts are similar to conspecific whole-body microbiomes in their

bacterial community profiles (Fig. S1), with many bacterial genera occurring at similar relative

abundances in guts and head/thorax tissue (Table S1). Some bacteria did differ between whole-

body and gut samples, such as *Spiroplasma*, which is a known hemolymph (blood) associate [48]

and which tends to be enriched in whole bodies (Fig. S1). Likewise, *Acinetobacter* was ~10-fold

more relatively abundant in head/thorax samples than in guts (Table S1). *Orbus*, *Enterobacter*,

Asaia, and some other dominant bacterial genera are clearly enriched in gut tissue (Table S1).

Host taxa	Phylum	Class	Order	Family	Genus	Mean prop. gut +/- SEM	Mean prop. head/thorax +/- SEM	p value
Heliconius	Bacteroidetes	Bacteroidia	Flavobacteriales	Weeksellaceae	*Apibacter*	0.036 +/- 0.009	0.002 +/- 0.001	< 0.001
					Chishuiella	0.007 +/- 0.003	0.095 +/- 0.027	0.001
					Chryseobacterium	0.011 +/- 0.004	0.096 +/- 0.024	0.001
	Firmicutes	Bacilli	Lactobacillales	Enterococcaceae	*Enterococcus*	0.031 +/- 0.008	0.021 +/- 0.006	0.117
				Streptococcaceae	*Lactococcus*	0.02 +/- 0.006	0.033 +/- 0.009	0.793
	Proteobacteria	Alphaproteobacteria	Acetobacterales	Acetobacteraceae	*Asaia*	0.024 +/- 0.007	0.004 +/- 0.002	0.001
					Commensalibacter	0.12 +/- 0.02	0.007 +/- 0.003	< 0.001
					Swaminathania	0.013 +/- 0.01	0.016 +/- 0.014	0.745
		Gammaproteobacteria	Betaproteobacteriales	Burkholderiaceae	*Variovorax*	0.01 +/- 0.004	0.021 +/- 0.006	0.017
			Enterobacteriales	Enterobacteriaceae	*Enterobacter*	0.233 +/- 0.034	0.031 +/- 0.007	< 0.001
					Serratia	0.037 +/- 0.01	0.027 +/- 0.017	0.013
			Orbales	Orbaceae	*Orbus*	0.163 +/- 0.024	0.021 +/- 0.01	< 0.001
			Pseudomonadales	Moraxellaceae	*Acinetobacter*	0.023 +/- 0.007	0.293 +/- 0.041	< 0.001
				Pseudomonadaceae	*Pseudomonas*	0.086 +/- 0.017	0.024 +/- 0.01	0.291
	Tenericutes	Mollicutes	Entomoplasmatales	Spiroplasmataceae	*Spiroplasma*	0.012 +/- 0.007	0.003 +/- 0.002	0.927
Other Heliconiini	Bacteroidetes	Bacteroidia	Flavobacteriales	Weeksellaceae	*Apibacter*	0.139 +/- 0.045	0.008 +/- 0.004	0.083
					Chishuiella	0.009 +/- 0.003	0.138 +/- 0.029	0.016
					Chryseobacterium	0 +/- 0	0.015 +/- 0.012	0.808
	Firmicutes	Bacilli	Lactobacillales	Enterococcaceae	*Enterococcus*	0.056 +/- 0.027	0.031 +/- 0.011	0.843
				Streptococcaceae	*Lactococcus*	0.036 +/- 0.026	0.012 +/- 0.011	0.098
	Proteobacteria	Alphaproteobacteria	Acetobacterales	Acetobacteraceae	*Asaia*	0.153 +/- 0.039	0.004 +/- 0.001	0.01
					Commensalibacter	0.031 +/- 0.015	0.002 +/- 0.002	0.106
					Swaminathania	0.007 +/- 0.004	0 +/- 0	0.226
		Gammaproteobacteria	Betaproteobacteriales	Burkholderiaceae	*Variovorax*	0.005 +/- 0.002	0.016 +/- 0.004	0.292
			Enterobacteriales	Enterobacteriaceae	*Enterobacter*	0.156 +/- 0.042	0.014 +/- 0.006	0.01
					Serratia	0.02 +/- 0.012	0.001 +/- 0	0.075
			Orbales	Orbaceae	*Orbus*	0.139 +/- 0.043	0.007 +/- 0.004	0.042
			Pseudomonadales	Moraxellaceae	*Acinetobacter*	0.046 +/- 0.018	0.45 +/- 0.054	< 0.001
				Pseudomonadaceae	*Pseudomonas*	0.02 +/- 0.017	0.011 +/- 0.004	0.393
	Tenericutes	Mollicutes	Entomoplasmatales	Spiroplasmataceae	*Spiroplasma*	0.068 +/- 0.038	0.052 +/- 0.052	0.843

201   Table S1. Relative abundances and within-body distribution of the top 15 bacterial genera (ranked by mean abundance)

202   across butterflies collected from Gamboa, Panama in 2016. These individuals were dissected to compare microbiomes

203   between isolated gut tissue and the combined head and thorax. Abundances are shown for *Heliconius* (top) and species

204   belonging to other heliconiine genera (bottom) separately. P values are from a nonparametric statistical test of

205   proportions in guts versus head/thorax samples, after FDR correction (see Methods).

Figure S1. Whole-body bacterial communities are consistent among individuals and predominantly represent gut-associated taxa. Shown are the relative abundances of the top 10 bacterial genera (ranked by mean abundance) across *H. erato petiverana* collected from Gamboa, Panama. Remaining white space represents sequences belonging to other genera; these made up a median 16% of sequence libraries across individuals. Samples on the left are isolated guts from individuals collected in 2016, while samples on the right are whole-body homogenates from individuals collected in 2014. Note that *Enterobacter* here includes sequences originally assigned as *Klebsiella* and some other closely related Enterobacteriaceae genera (see Methods).

We used metagenomes (N = 15) to examine the accuracy of amplicon sequencing for describing bacterial community composition in our broader sample set. Interindividual microbiome variation was highly correlated between the amplicon-based dataset and the metagenomic dataset (Mantel *r* = 0.63, p < 0.001). Furthermore, for six of the most abundant bacterial genera, relative abundances in amplicon libraries were highly predictive of relative

220   abundances in metagenomes (Fig. S2). These results support the use of amplicon sequencing for

221   estimating bacterial community composition in adult heliconiine butterflies.

224   Figure S2. Bacterial genus-level relative abundances are highly correlated between the amplicon sequence libraries

225   and metagenomes. Shown are relative abundances of six of the most abundant genera across the 15 butterfly gut

226   samples for which we obtained both amplicon and shotgun sequence data.

228   Compared to other passion-vine butterflies, adult *Heliconius* harbor clearly distinct

whole-body bacterial communities (Fig. 1). Microbiomes clustered by host genera to varying

degrees depending on the distance metric used. The effect was strongest with taxonomic or

phylogenetic metrics that incorporate information on the relative abundances of exact sequence

variants (ESVs) (Bray-Curtis: $R^2 = 0.13$, p = 0.001; weighted UniFrac: $R^2 = 0.18$, p = 0.001) as

opposed to purely presence/absence-based metrics (Jaccard: $R^2 = 0.11$, p = 0.001; unweighted

UniFrac: $R^2 = 0.10$, p = 0.001). Six individuals of the non-pollen-feeding species *Heliconius*

aoede (formerly *Neruda aeode*; see Methods) had bacterial communities more similar to those

found in pollen-feeding *Heliconius* individuals, while two individuals had communities more

similar to those found in other butterfly genera (Fig. 1).

Figure 1. Adult *Heliconius* butterflies host distinct bacterial communities compared with related non-pollen-feeding

genera. Shown is an ordination of microbiome variation (Bray-Curtis dissimilarities) among all whole-body samples

(N = 104 *Heliconius*, 52 other Heliconiini). With the exception of *H. aoede* (dark blue triangles), pollen-feeding is

exclusive to, and ubiquitous within *Heliconius*. However, the phylogenetic placement of *H. aoede* is not fully settled

(see Methods).

As suggested by the host genus-level clustering, we found that variation in butterfly

microbiome composition was associated with host relatedness (phylogenetic distance). More

closely related butterfly lineages harbored more similar microbiomes (Bray-Curtis: Mantel r =

0.40, p = 0.01), and the topologies of the butterfly phylogeny and the dendrogram of microbiome

similarity were moderately congruent (Fig. 2). This pattern was still statistically significant when

microbiome variation was measured using weighted UniFrac distances (Mantel r = 0.31, p =

0.03), but not when using unweighted UniFrac (Mantel r = 0.03, p = 0.41) or Jaccard distances

(Mantel r = 0.19, p = 0.08), which do not incorporate relative abundance information.

Host phylogeny   Microbiome dendrogram

Figure 2. Heliconiine butterfly microbiomes exhibit a strong signal of host relatedness. Shown is the concordance

between the host phylogeny (from [35]) and a dendrogram representation of microbiome variation among species

(Bray-Curtis dissimilarities). Here, all nodes have been rotated to maximize tip matching. Note, however, that sets of

parallel lines connecting tips do not always correspond to matching clades.

We then determined which bacterial taxa contribute to the observed taxonomic (Fig. 1)

and phylogenetic (Fig. 2) structuring in overall community composition. The dominant bacterial

genera *Enterobacter*, *Orbus*, and *Pseudomonas* (or *Serpens* [49]) were proportionally more

abundant in *Heliconius* versus the non-pollen-feeding butterfly genera, while *Asaia* and

Apibacter showed the opposite pattern (Fig. 3). None of these bacterial genera, however, were

exclusively restricted to one host feeding guild, genus, or species, and their relative abundances

were occasionally highly variable (Fig. 3). We also analyzed the data at the ESV-level, including

all individual ESVs that were reasonably abundant within at least one host species ($\geq$ 5% within-

species average). We did not find evidence for prevalent host species- or genus-restricted ESVs

(Fig. 4). Most ESVs are present, albeit with varying relative abundances, across host genus and

species boundaries. However, ESV_22 (*Chryseobacterium*) is notable as being abundant in

nearly all pollen-feeding species and rare or absent from non-pollen-feeding host species,

including *H. aoede*.

Figure 3. Dominant bacterial genera are largely shared among *Heliconius* and other butterflies, although some are

differentially abundant. Shown are the relative abundances of the top 10 bacterial genera, ranked by mean abundance,

in whole-body microbiomes. Dots indicate replicate individuals, and black bars indicate median proportions within a

host species. Starred bacterial genera differed significantly in relative abundance between *Heliconius* and non-

Heliconius butterflies (p < 0.05 after FDR correction). The arrangement of host species on the x axis corresponds to

the phylogeny shown in Fig. 4. Note that *Enterobacter* here includes sequences originally assigned as *Klebsiella* and

some other closely related Enterobacteriaceae genera (see Methods).

Figure 4. Most bacterial exact sequence variants (ESVs) are not host species- or host genus-specific. Shown are all

ESVs in the dataset that had ≥ 5% mean relative abundance across conspecific individuals for one or more host species.

For each ESV, the species-level mean relative abundance in whole-body samples, after log transformation, is indicated

by the color of the cells (white = not detected in that species). Note that bacterial genera (labeled at the top) contained

varying numbers of ESVs that met the aforementioned prevalence threshold.

Amplicon-derived ESVs are limited in their ability to resolve bacterial strains as they

represent only a short fragment of the 16S rRNA gene (here, ~250 bp). To test for potential host

specificity at a finer level of resolution, we obtained near-full-length 16S rRNA gene sequences

from the bacterium *Orbus*, which is highly prevalent across gut and whole-body samples (Fig. 3,

Fig. S1) and almost exclusively composed of a single ESV (Fig. 4). These sequences were

reconstructed from the 12 metagenomes in which *Orbus* was sufficiently abundant. A phylogeny

based on these sequences and other Orbaceae shows that butterfly-associated *Orbus* form a

single, well-supported clade (Fig. 5). Host-phylogenetic or geographic structure was not evident

within this clade. In fact, many of the Panamanian heliconiine butterflies harbored *Orbus* that

have identical or nearly-identical 16S rRNA gene sequences to an *Orbus* strain isolated from an

East Asian butterfly, *Sasakia charonda* [50].

Figure 5. Maximum likelihood phylogenetic reconstruction of the bacterial family Orbaceae (Gammaproteobacteria: Pasteurellales). Some host taxonomic structure is apparent at the order level (i.e. Lepidoptera, beetles, bees, flies) but not within butterflies. 16S rRNA gene sequences from *Agraulis*, *Dryadula*, *Dryas*, and *Heliconius* were assembled from short metagenomic reads. Other 16S rRNA gene sequences are from GenBank. *Haemophilus influenzae* (Pasteurellaceae) was used as the outgroup.

We also used the shotgun metagenomes to search for microbial eukaryotes in butterfly gut samples, and found that microsporidia and trypanosomatids were prevalent (27% and 47% of the 15 individual samples analyzed, respectively) (Fig. S3). When trypanosomatids were detected in a given sample, we also sometimes detected 16S rRNA gene reads in the metagenomic data classified as *Kinetoplastibacterium*, an obligate bacterial endosymbiont of certain trypanosomatids [51]. *Kinetoplastibacterium* was not detected without its corresponding trypanosomatid host (Fig. S3). This observation led us to reexamine the larger, amplicon-based

bacterial dataset. We found that 9% of *Heliconius* individuals and 4% of individuals belonging to other heliconiine genera were infected with a trypanosomatid, as inferred by the presence of *Kinetoplastibacterium*. These proportions were not significantly different (Fisher's Exact Test, p = 0.28). They are also likely underestimates, given the aforementioned prevalence of trypanosomatids among metagenomes (47%) and the potential for false negatives—in five metagenomes, trypanosomatids were detected without *Kinetoplastibacterium* (Fig. S3). These results point to a need for more targeted analyses of butterfly-associated parasite diversity.

Figure S3. Microsporidia and Trypanosomatidae (related to *Crithidia*, *Leishmania*, and *Trypanosoma*) are prevalent in the 15 butterfly individuals with sequenced gut metagenomes. Points show the proportion of each microeukaryotic taxon out of all non-butterfly 18S rRNA sequences identified by phyloFlash. Circles are filled if the proportion of microsporidia or trypanosomatids exceeded 0.01. Asterisks indicate samples in which *Kinetoplastibacterium*, a bacterial endosymbiont of trypanosomatids, was detected. Other microeukaryotes not shown here comprise a variety of very low-abundance taxa (i.e. ≤ 30 total reads per metagenome) classified mainly as coccidia, ascomycete fungi, acanthamoeba, and algae.

Discussion

The evolution of pollen-feeding in *Heliconius* butterflies underlies a number of unique traits and likely facilitated the rapid diversification of this genus. Although *Heliconius* have been a major study system for decades, their microbiomes were unexplored beyond a few species [12, 21, 22]. In particular, it was unknown whether they evolved distinctive microbiomes concomitant with this major shift in diet and life history. This question is particularly interesting in butterflies, in which even major diet shifts in the larval stage (caterpillars) do not depend on microbiomes [9–11]. *Heliconius* is arguably the single lineage of Lepidoptera most reliant on adult-stage nutrition, and thus might be likely to engage in microbial symbioses as adults.

Using extensive field collections of 21 species and subspecies of heliconiine butterflies, we found that adult *Heliconius* have relatively low-diversity bacterial communities that are compositionally distinct from those found in their non-pollen-feeding relatives (Fig. 1, Fig. 3). Most of the dominant bacterial taxa were present in isolated gut tissue, but others were enriched elsewhere in the body, such as the hemolymph or structures in the head and thorax (Fig. S1, Table S1). Beyond the clear separation of *Heliconius* from other heliconiine genera, there was also strong a phylogenetic signal in adult heliconiine butterfly microbiomes (Fig. 2). The strength of this signal was comparable to that observed in mammalian gut microbiomes [3] and other host groups [36].

Host-phylogenetic structure in microbiomes can arise from host-microbe co-diversification and/or contemporary host filtering of environmental microbes [4, 36]. Given that the dominant bacterial taxa appear to be largely shared among host lineages (discussed below), we suggest that host filtering is likely to be particularly important in heliconiine butterflies. For example, the shift to pollen-feeding in adult *Heliconius* represents a major change in nutrient inputs, potentially altering which resident bacteria are able to proliferate in the gut. Pollen itself also harbors microbes [52], some of which may colonize the gut. Community assembly in the adult stage could also be influenced, in part, by dynamics occurring in earlier life stages [7]. Some bacterial taxa present in mature, adult *H. erato* are already present in freshly emerged (unfed) adults, pupae and larvae, and are potentially carried over from the larval host plant through metamorphosis [22]. Moreover, Heliconiini larvae show marked variation in the species

370   identity and the age (young vs. old foliage) of host plants they consume [17], traits reported to

371   structure leaf microbiomes [53–55].

372   Certain bacterial taxa were enriched in *Heliconius* butterflies and their abundances may

be associated with with the evolution of pollen-feeding. For example, *Orbus* was the most

dominant bacterial genus among *Heliconius* and was more than twice as proportionally abundant

in *Heliconius* as compared with other heliconiine butterflies. The biology of the Orbaceae is

largely unknown, excepting the specialized bee gut symbionts *Gilliamella* and *Frischella* [56,

57]. In honey bees, which also consume pollen, some *Gilliamella* strains can degrade and

ferment structural compounds in pollen walls [58, 59]. Adding pollen to honey bee diets also

increases gut colonization by *Gilliamella* and *Frischella* [60]. However, a key difference is that

pollen digestion in *Heliconius*, unlike bees, occurs extra-orally. Pollen grains collected on the

proboscis are steeped in exuded saliva, and amino acids and other compounds released from the

grains are then ingested [61]. Host-encoded enzymes that likely play a role in pollen digestion

have been identified [62, 63]. Whether microbes contribute to this process is still unclear, but we

identified candidate bacterial taxa that are overrepresented in the head and thorax (Table S1),

structures that contain the proboscis and salivary gland. The exact localization and role of these

microbes is worth more detailed investigation.

Based on the amplicon-based data, we found that microbiome differences across

heliconiine butterflies were largely driven by shifts in the relative abundance of shared bacterial

genera and ESVs, as opposed to the presence or absence of host-specific bacterial taxa. These

bacteria might be widely distributed among larval host plants or adult nectar sources, and

independently acquired by different butterflies. Sympatric butterfly species might also exchange

bacteria, given their frequent co-occurrence at nectar sources. Flowers can act as hubs for the

horizontal transmission of microbes among pollinators [64, 65].

The amplicon data were based on a short region of the 16S rRNA gene and might obscure

finer-scale diversity and bacterial distribution patterns among hosts. We examined this

possibility using near-full-length 16S rRNA gene sequences from the bacterium *Orbus*. As

inferred from the amplicon data, this genus was represented by a single ESV that was nearly

ubiquitous across all butterflies (Fig. 4). We found additional diversity among the longer

sequences, but within butterflies, different *Orbus* phylotypes did not segregate by butterfly

phylogeny or geography (Fig. 5). At a broader scale, the Orbaceae do appear to exhibit some

host-taxonomic specificity (Fig. 5; also see [49]). Potentially, the Orbaceae have diversified into

loosely insect-order-specific lineages which at finer scales (e.g. within butterflies) frequently

disperse among host taxa and regions. However, we note that the 16S rRNA gene evolves

slowly, making it difficult to differentiate environmental transmission from co-diversification in

a young host radiation such as the Heliconiini [35] without additional information [4].

Adult butterflies clearly structure the composition of their bacterial communities, but the

reciprocal effect of these microbes on adult butterfly biology is completely unknown. A recent

study found nectar- and fruit-feeding butterflies to host gut microbes with different abilities to

metabolize substrates *in vitro* [12], but whether these differences are expressed *in vivo* and

whether they would matter for host nutrition remains unclear. A subsequent experiment on the

butterfly *Speyeria mormonia* using antibiotics did not find a strong relationship between gut

bacterial loads and measures of adult fitness [66]. However, antibiotic-based studies could fail to

test a role of the microbiome in resistance to parasites and pathogens if the latter are also

suppressed by antibiotics.

Colonization resistance is a common feature of many animal gut microbiomes [4, 67, 68]

and could be important in adult butterflies. Here, we discovered a high prevalence of putative

eukaryotic parasites (trypanosomatids and microsporidia) across adult heliconiine butterflies.

Butterfly-associated microeukaryotes are almost unknown, and there is very little research on

any adult-stage pathogens or parasites (with the exception of the neogregarine *Ophryocystis*

elektroscirrha in monarch butterflies [69]). *Serratia* were also widespread among heliconiine

butterflies and are frequently pathogenic in insects [70, 71]. Given their co-localization in

butterfly guts, we expect direct interactions and potential antagonism between these putative

parasites and other gut microbes. For example, in some pyrrhocorid bugs, tsetse flies and social

bees, gut bacteria provide an important layer of defense against trypanosomatid parasites and

Serratia [72–75].

Conclusions

This study provides the first in-depth characterization of *Heliconius* and other passion-

vine butterfly microbiomes, adding a new dimension to a classic model system in evolutionary

biology. Our finding that a major diet shift (pollen-feeding) coincides with a shift in adult-stage

microbiomes advances previous work on Lepidoptera feeding ecology and microbial interactions, most of which has focused on the larval stage. The ecological dynamics of microbiomes appear to be very different between larval and adult life stages, at least for lepidopteran groups that feed as adults. We also identified a strong host phylogenetic signal in passion-vine butterfly microbiomes, and speculate that it arises from host filtering of environmental microbes with conserved dietary or behavioral traits, as opposed to co-diversification.

The unique microbiome in *Heliconius* likely represents a response to the nutritional and microbial inputs from pollen, but may also represent a functional contribution to host pollen digestion. However, despite these microbial community-level differences, individual bacterial taxa were typically shared across host species and diet guilds. We suggest there is also overlap in microbiome function among hosts. In particular, the common occurrence of putative parasites in guts of various host species points to colonization resistance as a candidate function of adult microbiomes in passion-vine butterflies. Manipulative experiments can now be performed, informed by our characterization of natural microbial community structure across butterfly hosts. Such studies would fill a critical gap in our understanding of butterflies, a diverse group of considerable ecological, societal, and scientific importance [76].

Acknowledgments

This material is based on work supported by the Cooperative Institute for Research in Environmental Sciences, the Ecology and Evolutionary Biology Department at the University of Colorado Boulder, a National Science Foundation Graduate Research Fellowship (1144083), and a postdoctoral fellowship from the USDA National Institute of Food and Agriculture (2018-08156), to TJH. This work was further supported by grants from the Simons Foundation (429440) and by funding from the Smithsonian Tropical Research Institute. We also thank Jessica Henley for assistance with sequencing, the Smithsonian Tropical Research Institute staff for administrative support and the Panamanian and Ecuadorian Authorities for permission to collect butterflies.

References

1. Muegge BD, Kuczynski J, Knights D, Clemente JC, González A, Fontana L, et al. Diet drives convergence in gut microbiome functions across mammalian phylogeny and within humans. *Science* 2011; **332**: 970–974.

2. Delsuc F, Metcalf JL, Wegener Parfrey L, Song SJ, González A, Knight R. Convergence of gut microbiomes in myrmecophagous mammals. *Mol Ecol* 2014; **23**: 1301–1317.

3. Groussin M, Mazel F, Sanders JG, Smillie CS, Lavergne S, Thuiller W, et al. Unraveling the processes shaping mammalian gut microbiomes over evolutionary time. *Nature Communications* 2017; **8**: 14319.

4. Moran NA, Ochman H, Hammer TJ. Evolutionary and Ecological Consequences of Gut Microbial Communities. *Annu Rev Ecol Evol Syst* 2019; **50**.

5. Engel P, Moran NA. The gut microbiota of insects - diversity in structure and function. *FEMS Microbiology Reviews* 2013; **37**: 699–735.

6. McFall-Ngai M, Hadfield MG, Bosch TCG, Carey HV, Domazet-Lošo T, Douglas AE, et al. Animals in a bacterial world, a new imperative for the life sciences. *Proceedings of the National Academy of Sciences* 2013; **110**: 3229–3236.

7. Hammer TJ, Moran NA. Links between metamorphosis and symbiosis in holometabolous insects. *Phil Trans R Soc B* 2019; **374**: 20190068.

8. Koenig JE, Spor A, Scalfone N, Fricker AD, Stombaugh J, Knight R, et al. Succession of microbial consortia in the developing infant gut microbiome. *Proceedings of the National Academy of Sciences* 2011; **108**: 4578–4585.

9. Hammer TJ, Janzen DH, Hallwachs W, Jaffe SP, Fierer N. Caterpillars lack a resident gut microbiome. *Proceedings of the National Academy of Sciences* 2017; **114**: 9641–9646.

10. Szenteczki MA, Pitteloud C, Casacci LP, Kešnerová L, Whitaker MRL, Engel P, et al. Bacterial communities within *Phengaris (Maculinea) alcon* caterpillars are shifted following transition from solitary living to social parasitism of *Myrmica* ant colonies. *Ecol Evol* 2019; **9**: 4452–4464.

11. Whitaker M, Pierce N, Salzman S, Kaltenpoth M, Sanders J. Microbial communities of lycaenid butterflies do not correlate with larval diet. *Frontiers in Microbiology* 2016; **7**: 1920.

494   12.   Ravenscraft A, Berry M, Hammer T, Peay K, Boggs C. Structure and function of the
495   bacterial and fungal gut microbiota of Neotropical butterflies. *Ecological Monographs*
496   2019; **89**: e01346.

497   13.   Stork NE. How Many Species of Insects and Other Terrestrial Arthropods Are There on
498   Earth? *Annu Rev Entomol* 2018; **63**: 31–45.

499   14.   Scoble MJ. The Lepidoptera. 1992. Oxford University Press.

500   15.   Gilbert LE. Pollen Feeding and Reproductive Biology of *Heliconius* Butterflies.
501   *Proceedings of the National Academy of Sciences* 1972; **69**: 1403–1407.

502   16.   Cardoso MZ, Gilbert LE. Pollen feeding, resource allocation and the evolution of chemical
503   defence in passion vine butterflies. *Journal of Evolutionary Biology* 2013; **26**: 1254–1260.

504   17.   Jiggins CD. The ecology and evolution of *Heliconius* butterflies. 2016. Oxford University
505   Press.

506   18.   Ehrlich PR, Gilbert LE. Population Structure and Dynamics of the Tropical Butterfly
507   *Heliconius ethilla*. *Biotropica* 1973; **5**: 69–82.

508   19.   Dunlap-Pianka H, Boggs CL, Gilbert LE. Ovarian Dynamics in Heliconiine Butterflies:
509   Programmed Senescence versus Eternal Youth. *Science* 1977; **197**: 487–490.

510   20.   Cook LM, Thomason EW, Young AM. Population Structure, Dynamics and Dispersal of
511   the Tropical Butterfly *Heliconius charitonius*. *The Journal of Animal Ecology* 1976; **45**:
512   851–863.

513   21.   van Schooten B, Godoy-Vitorino F, McMillan WO, Papa R. Conserved microbiota among
514   young *Heliconius* butterfly species. *PeerJ* 2018; **6**: e5502.

515   22.   Hammer TJ, McMillan WO, Fierer N. Metamorphosis of a Butterfly-Associated Bacterial
516   Community. *PLoS ONE* 2014; **9**: e86995.

517   23.   Hammer TJ, Dickerson JC, Fierer N. Evidence-based recommendations on storing and
518   handling specimens for analyses of insect microbiota. *PeerJ* 2015; **3**: e1190.

519   24.   Gardes M, Bruns TD. ITS primers with enhanced specificity for basidiomycetes -
520   application to the identification of mycorrhizae and rusts. *Mol Ecol* 1993; **2**: 113–118.

521   25.   Baym M, Kryazhimskiy S, Lieberman TD, Chung H, Desai MM, Kishony R. Inexpensive
522   Multiplexed Library Preparation for Megabase-Sized Genomes. *PLOS ONE* 2015; **10**:
523   e0128036.

26.   Martin M. Cutadapt removes adapter sequences from high-throughput sequencing reads.

EMBnet.journal 2011; **17**: 10–12.

27.   Callahan BJ, McMurdie PJ, Rosen MJ, Han AW, Johnson AJA, Holmes SP. DADA2:

High-resolution sample inference from Illumina amplicon data. *Nat Methods* 2016; **13**:

581–583.

28.   Wang Q, Garrity GM, Tiedje JM, Cole JR. Naive Bayesian classifier for rapid assignment

of rRNA sequences into the new bacterial taxonomy. *Applied and Environmental*

Microbiology 2007; **73**: 5261–5267.

29.   Quast C, Pruesse E, Yilmaz P, Gerken J, Schweer T, Yarza P, et al. The SILVA ribosomal

RNA gene database project: improved data processing and web-based tools. *Nucleic Acids*

Research 2012; **41**: D590–D596.

30.   R Core Team. R: A language and environment for statistical computing. 2016. R

Foundation for Statistical Computing, Vienna, Austria.

31.   Davis NM, Proctor DM, Holmes SP, Relman DA, Callahan BJ. Simple statistical

identification and removal of contaminant sequences in marker-gene and metagenomics

data. *Microbiome* 2018; **6**: 226.

32.   Brady C, Cleenwerck I, Venter S, Coutinho T, De Vos P. Taxonomic evaluation of the

genus *Enterobacter* based on multilocus sequence analysis (MLSA). *Systematic and*

Applied Microbiology 2013; **36**: 309–319.

33.   Janssen S, McDonald D, Gonzalez A, Navas-Molina JA, Jiang L, Xu ZZ, et al.

Phylogenetic Placement of Exact Amplicon Sequences Improves Associations with Clinical

Information. *mSystems* 2018; **3**: e00021-18.

34.   McDonald D, Price MN, Goodrich J, Nawrocki EP, DeSantis TZ, Probst A, et al. An

improved Greengenes taxonomy with explicit ranks for ecological and evolutionary

analyses of bacteria and archaea. *The ISME Journal* 2012; **6**: 610–618.

35.   Kozak KM, Wahlberg N, Neild AF, Dasmahapatra KK, Mallet J, Jiggins CD. Multilocus

Species Trees Show the Recent Adaptive Radiation of the Mimetic *Heliconius* Butterflies.

Systematic Biology 2015; **64**: 505–524.

36.   Mazel F, Davis KM, Loudon A, Kwong WK, Groussin M, Parfrey LW. Is Host Filtering

the Main Driver of Phylosymbiosis across the Tree of Life? *mSystems* 2018; **3**: e00097-18.

37. Revell LJ. phytools: An R package for phylogenetic comparative biology (and other
things). *Methods in Ecology and Evolution* 2012; **3**: 217–223.

38. Oksanen J, Kindt R, Legendre P. vegan: Community Ecology Package. R package version
2.0-7. 2013.

39. Anderson MJ, Walsh DCI. PERMANOVA, ANOSIM, and the Mantel test in the face of
heterogeneous dispersions: What null hypothesis are you testing? *Ecological Monographs*
2013; **83**: 557–574.

40. Joshi, NA, Fass, JN. Sickle: A sliding-window, adaptive, quality-based trimming tool for
FastQ files. https://github.com/najoshi/sickle.

41. Langmead B, Salzberg SL. Fast gapped-read alignment with Bowtie 2. *Nat Methods* 2012;
9: 357–359.

42. Challis RJ, Kumar S, Dasmahapatra KK, Jiggins CD, Blaxter M. Lepbase: the Lepidopteran
genome database. *bioRxiv* 2016; http://dx.doi.org/10.1101/056994.

43. Gruber-Vodicka HR, Seah BK, Pruesse E. phyloFlash — Rapid SSU rRNA profiling and
targeted assembly from metagenomes. *bioRxiv* 2019; https://doi.org/10.1101/521922.

44. Edgar RC. MUSCLE: Multiple sequence alignment with high accuracy and high
throughput. *Nucleic Acids Research* 2004; **32**: 1792–1797.

45. Castresana J. Selection of Conserved Blocks from Multiple Alignments for Their Use in
Phylogenetic Analysis. *Molecular Biology and Evolution* 2000; **17**: 540–552.

46. Dereeper A, Guignon V, Blanc G, Audic S, Buffet S, Chevenet F, et al. Phylogeny.fr:
robust phylogenetic analysis for the non-specialist. *Nucleic Acids Research* 2008; **36**:
W465–W469.

47. Guindon S, Delsuc F, Dufayard JF, Gascuel O. Estimating maximum likelihood phylgenies
with PhyML. *Methods in Molecular Biology* 2009; **537**: 113–137.

48. Blow F, Douglas AE. The hemolymph microbiome of insects. *Journal of Insect Physiology*
2019; **115**: 33–39.

49. Martinson VG, Carpinteyro-Ponce J, Moran NA, Markow TA. A Distinctive and Host-
Restricted Gut Microbiota in Populations of a Cactophilic *Drosophila* Species. *Applied and*
Environmental Microbiology 2017; **83**: e01551–17.

50. Kim JY, Lee J, Shin N-R, Yun J-H, Whon TW, Kim M-S, et al. *Orbus sasakiae* sp. nov., a
bacterium isolated from the gut of the butterfly *Sasakia charonda*, and emended description

of the genus *Orbus*. *International Journal of Systematic and Evolutionary Microbiology*
2012; **63**: 1766–1770.

51. De Souza W, Motta MCM. Endosymbiosis in protozoa of the Trypanosomatidae family.
FEMS Microbiology Letters 1999; **173**: 1–8.

52. Ambika Manirajan B, Ratering S, Rusch V, Schwiertz A, Geissler-Plaum R, Cardinale M,
et al. Bacterial microbiota associated with flower pollen is influenced by pollination type,
and shows a high degree of diversity and species-specificity: The bacterial microbiota of
flower pollen. *Environmental Microbiology* 2016; **18**: 5161–5174.

53. Redford AJ, Fierer N. Bacterial succession on the leaf surface: a novel system for studying
successional dynamics. *Microbial Ecology* 2009; **58**: 189–98.

54. Bulgarelli D, Schlaeppi K, Spaepen S, van Themaat EVL, Schulze-Lefert P. Structure and
Functions of the Bacterial Microbiota of Plants. *Annual Review of Plant Biology* 2013; **64**:
807–838.

55. Lindow SE, Brandl MT. Microbiology of the Phyllosphere. *Applied and Environmental*
Microbiology 2003; **69**: 1875.

56. Kwong WK, Moran NA. Cultivation and characterization of the gut symbionts of honey
bees and bumble bees: description of *Snodgrassella alvi* gen. nov., sp. nov., a member of
the family Neisseriaceae of the Betaproteobacteria, and *Gilliamella apicola* gen. nov., sp.
nov., a member of Orbaceae fam. nov., Orbales ord. nov., a sister taxon to the order
'Enterobacteriales' of the Gammaproteobacteria. *International Journal of Systematic and*
Evolutionary Microbiology 2013; **63**: 2008–2018.

57. Engel P, Kwong WK, Moran NA. *Frischella perrara* gen. nov., sp. nov., a
gammaproteobacterium isolated from the gut of the honey bee, *Apis mellifera*. *International*
Journal of Systematic and Evolutionary Microbiology 2013; **63**: 3646–3651.

58. Engel P, Martinson VG, Moran NA. Functional diversity within the simple gut microbiota
of the honey bee. *Proceedings of the National Academy of Sciences* 2012; **109**: 11002–
11007.

59. Kesnerova L, Mars RAT, Ellegaard KM, Troilo M, Sauer U, Engel P. Disentangling
metabolic functions of bacteria in the honey bee gut. *PLoS Biology* 2017; **15**: e2003467.

60. Kešnerová L, Emery O, Troilo M, Erkosar B, Engel P. Gut microbiota structure differs
between honey bees in winter and summer. *bioRxiv* 2019; https://doi.org/10.1101/703512.

61.   Eberhard SH, Hikl AL, Boggs CL, Krenn HW. Saliva or Regurgitated Nectar? What
617   *Heliconius* Butterflies (Lepidoptera: Nymphalidae) Use for Pollen Feeding. *Annals of the*
618   *Entomological Society of America* 2009; **102**: 1105–1108.

619   62.   Harpel D, Cullen DA, Ott SR, Jiggins CD, Walters JR. Pollen feeding proteomics: Salivary
620   proteins of the passion flower butterfly, *Heliconius melpomene. Insect Biochemistry and*
621   *Molecular Biology* 2015; **63**: 7–13.

622   63.   Smith G, Macias-Muñoz A, Briscoe AD. Gene Duplication and Gene Expression Changes
623   Play a Role in the Evolution of Candidate Pollen Feeding Genes in *Heliconius* Butterflies.
624   *Genome Biology and Evolution* 2016; **8**: 2581–2596.

625   64.   McArt SH, Koch H, Irwin RE, Adler LS. Arranging the bouquet of disease: floral traits and
626   the transmission of plant and animal pathogens. *Ecology Letters* 2014; **17**: 624–636.

627   65.   Graystock P, Goulson D, Hughes WOH. Parasites in bloom: flowers aid dispersal and
628   transmission of pollinator parasites within and between bee species. *Proceedings of the*
629   *Royal Society B: Biological Sciences* 2015; **282**: 20151371.

630   66.   Ravenscraft A, Kish N, Peay K, Boggs C. No evidence that gut microbiota impose a net
631   cost on their butterfly host. *Molecular Ecology* 2019; **28**: 2100–2117.

632   67.   Kamada N, Chen GY, Inohara N, Núñez G. Control of pathogens and pathobionts by the
633   gut microbiota. *Nature Immunology* 2013; **14**: 685–690.

634   68.   Ducarmon QR, Zwittink RD, Hornung BVH, van Schaik W, Young VB, Kuijper EJ. Gut
635   Microbiota and Colonization Resistance against Bacterial Enteric Infection. *Microbiol Mol*
636   *Biol Rev* 2019; **83**: e00007-19.

637   69.   Altizer SM, Oberhauser KS. Effects of the Protozoan Parasite *Ophryocystis elektroscirrha*
638   on the Fitness of Monarch Butterflies (*Danaus plexippus*). *Journal of Invertebrate*
639   *Pathology* 1999; **74**: 76–88.

640   70.   Raymann K, Coon KL, Shaffer Z, Salisbury S, Moran NA. Pathogenicity of *Serratia*
641   *marcescens* Strains in Honey Bees. *mBio* 2018; **9**: e01649-18.

642   71.   Grimont F, Grimont PAD. The Genus *Serratia*. In: Dworkin M, Falkow S, Rosenberg E,
643   Schleifer K-H, Stackebrandt E (eds). *The Prokaryotes*. 2006. Springer New York, pp 219–
644   244.

72.   Onchuru TO, Martinez AJ, Kaltenpoth M. The cotton stainer's gut microbiota suppresses
infection of a cotransmitted trypanosomatid parasite. *Molecular Ecology* 2018; **27**: 3408–
3419.
73.   Raymann K, Shaffer Z, Moran NA. Antibiotic exposure perturbs the gut microbiota and
elevates mortality in honeybees. *PLoS Biology* 2017; **15**: e2001861.
74.   Koch H, Schmid-Hempel P. Socially transmitted gut microbiota protect bumble bees
against an intestinal parasite. *Proceedings of the National Academy of Sciences* 2011; **108**:
19288–19292.
75.   Weiss BL, Maltz MA, Vigneron A, Wu Y, Walter KS, O'Neill MB, et al. Colonization of
the tsetse fly midgut with commensal *Kosakonia cowanii* Zambiae inhibits trypanosome
infection establishment. *PLOS Pathogens* 2019; **15**: e1007470.
76.   Boggs CL, Watt WB, Ehrlich PR. Butterflies: ecology and evolution taking flight. 2003.
The University of Chicago Press.