Heliconius butterflies host characteristic and phylogenetically structured adult-stage microbiomes

Tobin J. Hammer[1*], Jacob C. Dickerson[2,3], W. Owen McMillan[2], Noah Fierer[3,4]

1. Department of Integrative Biology, University of Texas at Austin, Austin, TX 78712, USA

2. Smithsonian Tropical Research Institute, Apartado 0843-03092, Panama City, Panama

3. Department of Ecology & Evolutionary Biology, University of Colorado at Boulder, Boulder, CO 80309, USA

4. Cooperative Institute for Research in Environmental Sciences, University of Colorado at Boulder, Boulder, CO 80309, USA

*Corresponding author: tobin.hammer@utexas.edu

Keywords: symbiosis, gut bacteria, Lepidoptera, microbiota, parasites

Abstract

Lepidoptera (butterflies and moths) are diverse and ecologically important, yet we know

little about how they interact with microbes as adults. Due to metamorphosis, the form and

function of their adult-stage microbiomes might be very different from microbiomes in the larval

stage (caterpillars). We studied adult-stage microbiomes of *Heliconius* and closely related

passion-vine butterflies (Heliconiini), which are an important model system in evolutionary

biology. To characterize the structure and dynamics of heliconiine microbiomes, we used field

collections of wild butterflies, 16S rRNA gene sequencing, quantitative PCR, and shotgun

metagenomics. We found that *Heliconius* harbor simple and abundant bacterial communities that

are moderately consistent among conspecific individuals and over time. Heliconiine

microbiomes also exhibited a strong signal of host phylogeny, with a major distinction between

Heliconius and other butterflies. These patterns were largely driven by differing relative

abundances of bacterial phylotypes shared among host species and genera, as opposed to the

presence or absence of host-specific phylotypes. We suggest that phylogenetic structure in

heliconiine microbiomes arises from conserved host traits that differentially filter microbes from

the environment. While the relative importance of different traits remains unclear, our data

indicate that pollen-feeding (unique to *Heliconius*) is not a primary driver. Using shotgun

metagenomics, we also discovered trypanosomatids and microsporidia to be prevalent in

butterfly guts, raising the possibility of antagonistic interactions between eukaryotic parasites

and co-localized gut bacteria. Our discovery of characteristic and phylogenetically structured

microbiomes provides a foundation for tests of adult-stage microbiome function, a poorly

understood aspect of lepidopteran biology.

Importance

Many insects host microbiomes with important ecological functions. However, the

prevalence of this phenomenon is unclear, because in many insect taxa microbiomes have only

been studied in part of the life cycle, if at all. A prominent example is the butterflies and moths,

in which the composition and functional role of adult-stage microbiomes are largely unknown.

We comprehensively characterized microbiomes in adult passion-vine butterflies. Butterfly-

associated bacterial communities are generally abundant in guts, consistent within populations,

and composed of taxa widely shared among hosts. More closely related butterflies harbor more

similar microbiomes, with the most dramatic shift in microbiome composition occurring in

tandem with a suite of ecological and life history traits unique to the genus *Heliconius*.

Butterflies are also frequently infected with previously undescribed eukaryotic parasites, which

may interact with bacteria in important ways. These findings advance our understanding of

butterfly biology and of insect-microbe interactions generally.

Introduction

Insect microbiome research has historically focused on hosts with highly stable, specific,

and functionally important microbiomes (1). For example, the obligate nutritional endosymbionts

of sap- and blood-feeding insects, which exemplify this scenario, have long served as important

model systems. More recently, however, wider microbial explorations have shown that this kind

of association does not apply to all insect taxa. In some insect groups, microbiomes are more

variable, and the constituent microbes have low or no specificity to their hosts (2, 3). Moreover,

the functional importance of these microbiomes is often unclear—and possibly variable, low, or

nonexistent in some groups (4). Understanding the nature of microbiomes in these insects is not

only important to their biology, but also for a broader understanding of insect microbiome

evolution.

The Lepidoptera (butterflies and moths) are a group in which microbiome form and

function have been particularly difficult to resolve. Recent work indicates that larvae

(caterpillars) typically harbor microbiomes that are variable, low-abundance, and transient (5–9).

However, a crucial consideration for Lepidoptera and other insects that undergo complete

metamorphosis is that microbial associations may change between life stages (10). In

Lepidoptera, larvae and adults do often differ in the composition and absolute abundance of their

microbiomes (6, 11–15). Beyond this pattern, however, the ecology of adult-stage microbiomes

remains poorly understood.

There are three major unknowns about microbiomes of adult Lepidoptera that we sought

to address in this study. First, it is unclear how variable microbiomes are among host individuals

and taxa, and which ecological or evolutionary factors (e.g., host phylogeny) are associated with

microbiome variation. Understanding these patterns could give insights into whether

microbiomes are important for certain ecological traits, such as feeding, and into the likelihood

of host-microbe coevolution. Second, it is unclear whether butterfly-associated microbes have

any specificity to their host species or to higher taxonomic levels; tight specificity is a hallmark

of obligate insect-microbe symbioses (3). Previous studies of adult Lepidoptera microbiomes

have largely used 97% operational taxonomic units (OTUs) based on short fragments of the 16S

rRNA gene. OTUs can encompass substantial strain-level diversity (16), hindering analyses of

specificity. Third, the full diversity of eukaryotic microbes that may be associated with adult

Lepidoptera has not been explored, as previous studies have exclusively focused on bacteria or

fungi.

Our study focuses on the neotropical passion-vine butterflies (Nymphalidae: Heliconiini).

This tribe, which includes the genus *Heliconius*, is a foundational system in evolutionary biology

with over 150 years of scientific study (17). Given the scientific relevance of heliconiines (18),

and the myriad roles that microbes can play in animal biology (19), it is important to know the

structure and function of heliconiine-associated microbiomes. Heliconiine butterflies are also a

potentially useful system for the field of host-microbe interactions, given the wealth of genomic

data available (20) and the ability to rear and genetically manipulate members of this group (21).

A handful of amplicon sequencing-based microbiome studies have included adult heliconiines,

but these were limited to a small number of individuals and species (11, 22, 23). Consequently,

we lack even basic information about the microbes harbored by heliconiine butterflies and how

they may vary across individuals and the host phylogeny.

In addition to the practical advantages mentioned above, heliconiines are also an

interesting system to address how host ecological traits evolve in concert with microbiomes.

Within Heliconiini, the genus *Heliconius* exhibits unique feeding and life history traits relative to

other genera: larvae feed on young passion-vine shoots and leaves as opposed to old foliage,

adults of most species (besides *H. aoede*) feed on pollen as well as nectar, and they have

distinctive defensive chemistry and a greatly extended adult-stage lifespan of several months (18,

24). These traits could influence and be influenced by microbiomes in many ways. For example,

we hypothesized that pollen-feeding might be associated with a distinctive microbiome, either

because it is a source of microbes (25) or of nutrients that shift resident microbiome composition

(26), or, because the trait itself depends on metabolic contributions from novel microbes (11). A

useful starting point to uncovering these kinds of interactions is to examine whether *Heliconius*

have microbiomes that are internally consistent yet distinct from those of other heliconiine

genera. If so, then we can zero in on the specific ecological and evolutionary drivers of variation,

and the potential functional roles of microbiomes in *Heliconius* biology.

Here we addressed the following main questions: *i*) How variable are heliconiine

microbiomes among host individuals, species, genera, and across the heliconiine phylogeny? *ii*)

Are host traits, such as pollen-feeding, associated with microbiome variation? *iii*) How host-

specific are butterfly-associated microbes? At the same time, we sought to facilitate future

microbiome research on heliconiine butterflies by answering some additional questions: *iv*) How

are microbes distributed within the butterfly body? *v*) How abundant are these microbes? *vi*)

How reliable is amplicon sequencing of 16S rRNA genes for studying butterfly-associated

bacterial communities, and *vii*) are there eukaryotic microbes we might be missing with targeted

amplicon sequencing?

We collected 214 wild adult butterflies, representing 23 species and subspecies of

Heliconiini, and characterized their microbiomes with 16S rRNA gene sequencing. While many

adult Lepidoptera microbiome studies have used reared, captive specimens (6, 12, 13, 22, 27),

these may give a biased picture of microbial community structure in the wild (11). We also used

shotgun metagenomic sequencing on a subset of butterflies both to provide an untargeted

assessment of microbial diversity (including eukaryotes) and to provide a finer-resolution picture

of microbial specificity than is available from amplicon sequencing of short 16S rRNA gene

regions. Furthermore, we used quantitative PCR to estimate the absolute abundance of bacteria

across different heliconiine taxa and tissue types. Our work illustrates ecological and

evolutionary dynamics of microbiomes in adult heliconiine butterflies and advances our general

understanding of Lepidoptera-microbe interactions.

Materials and Methods

Field collections

123   The wild adult butterflies used for whole-body microbiome sequencing were collected

from seven locations in Panama and Ecuador in May-August 2014 (more detail is provided in the

supplemental file "Collection_localities.txt"). Butterflies were euthanized with ethyl acetate and

stored in DMSO after removal of wings, following (28). We also stored two DMSO-only blanks

to use as negative controls. In June 2016, we collected additional adult butterflies for gut and

head/thorax sequencing from Gamboa and Pipeline Road, Panama. For these specimens, we

dissected the gut (hindgut, midgut, and the distal ~1/2 of the foregut) using sterilized tools prior

to storage in DMSO. The whole head and thorax (including the proximal foregut) were stored

separately. Species or subspecies were identified based on morphology. Butterflies were

collected under permit # SC/A-7-11 from Panama's Autoridad Nacional del Ambiente and #

005-13 IC-FAU-DNB/MA from Ecuador's Ministerio del Ambiente.

Sample processing, qPCR, PCR and sequencing

137   We removed whole bodies and head/thorax samples from DMSO and, after

homogenization, used approximately 50 mg subsamples of homogenate for DNA extractions

with the MoBio PowerSoil kit following the manufacturer's instructions. We added entire guts

directly to DNA extraction tubes, in which they were homogenized during the first bead-beating

step of the protocol. Two DMSO blanks and 30 DNA extraction blanks were also processed in

tandem with the butterfly samples and sequenced. For dissected gut and head/thorax samples, we

estimated total bacterial abundance using quantitative PCR (qPCR) with 16S rRNA gene primers

(515F/806R) following the protocol described in ref. (5).

PCR amplifications (515F/806R primers, V4 region) and 2 X 150 bp Illumina MiSeq

sequencing of 16S rRNA genes followed standard Earth Microbiome Project protocols

(dx.doi.org/10.17504/protocols.io.nuudeww). For gut and head/thorax samples of 29 butterfly

individuals of nine species, we also attempted PCR amplification with primers that target the ITS

gene region of fungi (29). However, amplification success with these fungal-specific primers (as

assessed by gel electrophoresis) was very low, suggesting a lack of abundant fungal DNA that

was later corroborated with the shotgun metagenomic data (see below). DNA extracts from a

subset of 15 amplicon-sequenced gut samples were used for shotgun metagenomic sequencing

following the approach described previously (30) with an input DNA concentration of 0.75

ng/ul, KAPA HiFi HotStart ReadyMix and bead cleanup with Ampure XP beads at a 0.9x ratio.

Amplicon data processing

Amplicons from the 2014 whole-body samples and 2016 gut and head/thorax samples

were sequenced on separate runs, demultiplexed using idemp (https://github.com/yhwu/idemp),

and combined for further processing. Cutadapt (31) was used to remove primer sequences. We

then used the DADA2 pipeline (32) to quality-filter (max EE value = 1) and trim (150 bp

forward, 140 bp reverse) reads, infer exact sequence variants (ESVs), merge paired-end reads,

and remove chimeras. We classified ESVs using the RDP Naive Bayesian Classifier algorithm

(33) against the SILVA training set v. 132 (34).

Further data processing and analyses were conducted in R v. 3.6.0 (35). We used

decontam (36) for prevalence-based identification of putative contaminant ESVs based on 34

negative controls (DMSO and DNA extraction blanks and PCR no-template controls). The

median percentage of contaminant sequences across butterfly samples was 0.09%, but two

samples had >10% contaminants and were removed from further analysis. ESVs with < 100 total

sequences across all samples (out of a combined total of 5.6 million sequences) were removed,

as were ESVs classified as mitochondria or chloroplast, or bacteria lacking sub-domain

identification. These ESVs combined typically made up a low proportion of reads from the

libraries (median of 2.6% across all samples). We also used these data to correct qPCR-based

absolute abundances for non-bacterial amplification. Specifically, we multiplied the proportion

of bacteria in sequence libraries by the total number of 16S rRNA gene copies to obtain

estimates of bacterial absolute abundance.

As the 16S rRNA gene amplicon libraries were highly variable in read depth across

samples, we rarefied all samples to 5,000 sequences, filtering out 13 samples with lower

sequence depth. We relabeled *Pantoea*, *Erwinia*, *Kluyvera*, *Citrobacter*, *Klebsiella*, and

Cronobacter taxonomic assignments to *Enterobacter*. This step was taken as genera within

Enterobacteriaceae are often polyphyletic and are difficult to resolve from short 16S rRNA gene

regions (e.g., (37)), and we wanted to avoid spurious separation of ESVs among genera and

resulting idiosyncrasies in genus distributions across butterflies. Another case of ambiguous

taxonomy occurred with sequences classified as *Pseudomonas* by the SILVA reference database,

which we label below as "*Entomomonas/Pseudomonas*". *Entomomonas* is a recently described

genus of apparently insect-specialized Pseudomonadaceae and includes several sequences

retrieved from *Heliconius* genome assemblies (38).

Amplicon and qPCR data analysis

Beta diversity statistics and plots are based on Bray-Curtis dissimilarities and/or

UniFrac distances (weighted and unweighted). To obtain a bacterial phylogeny for the latter, we

used the fragment-insertion method (39) to place our ESV sequences into the Greengenes

reference tree (40). The butterfly phylogeny is from (41) (TreeBASE #Tr77496). Four of the

butterfly species in our sample set contained specimens from two distinct subspecies (e.g.,

Heliconius sara sara and *Heliconius sara magdalena*). To include these in the species-level host

phylogeny, we inserted subspecies tips halfway along the terminal branches to their sister

subspecies. Hereafter we refer to these subspecies as "species" for simplicity.

To test for host-phylogenetic signal in microbiomes, we used Mantel tests with 9999

permutations to calculate the correlation between microbial community dissimilarities/distances

and host phylogenetic distances (42). Intraspecific variation in microbiomes was handled by

averaging the pairwise dissimilarities/distances between all individuals of one species and all

individuals of another species. We used the phytools package (43) to visualize concordance

between topologies of the host phylogeny and a dendrogram of bacterial community

dissimilarities. Nodes were rotated with the "cophylo" function in phytools to maximize tip

matching between the two trees.

Differences in overall community composition between host genera were tested with

PERMANOVA as implemented in the vegan package (44). Using the "betadisper" function we

corroborated that significant test results were due to host genus-level differences in location and

not dispersion (45). We used a nonparametric statistical test (Wilcoxon rank-sum) to identify

bacterial genera that differed in relative abundance between host taxa or between sample types

(gut versus head/thorax) and applied a false discovery rate correction to the resulting p values.

We tested whether total bacterial abundance differed between *Heliconius* and other

butterfly genera using qPCR data from gut samples and head/thorax samples. Each sample type

was analyzed separately using log-transformed counts of bacterial 16S rRNA gene copies. We

verified that residuals were approximately normally distributed and used a linear mixed-effects

model as implemented in the nlme package (46), treating host genus (*Heliconius* vs. others) as a

fixed effect and host species as a random effect.

Metagenome data processing and analysis

For 15 gut samples, we obtained shotgun metagenomic data to complement the

bacterial 16S rRNA gene amplicon dataset. We quality-filtered these reads with sickle (47) and

trimmed adapters with cutadapt (31). We then used Bowtie 2 (48) to filter out reads matching a

given sample's corresponding host species' genome, obtained from Lepbase (20). The two

Dryadula phaetusa metagenomes were mapped to a genome of the sister species *Dryas iulia* as

no *Dryadula* genome was available. Since there was a high proportion of host-derived reads, we

focused here on describing microbial diversity using ribosomal RNA gene reads present in the

metagenomes. With the host-filtered reads, we used phyloFlash (49) to find and classify

eukaryotic and bacterial SSU rRNA reads. Bacterial community composition was compared

between the amplicon and shotgun metagenomic datasets using a Mantel test. We also used

phyloFlash to assemble 16S rRNA genes from the 150 bp shotgun reads. These longer sequences

allowed us to estimate the phylogeny of *Orbus*, the dominant bacterium in these 15 samples.

Orbus sequences were aligned with MUSCLE (50), curated with Gblocks (51), and used for

maximum likelihood reconstruction with the phylogeny.fr implementation (52) of PhyML (53).

Data availability

Amplicon data, metadata, and R code are available from figshare

(figshare.com/projects/Heliconius_butterfly_microbiomes/70520). Metagenomes are available

from MG-RAST (project no. MGP89563).

Results

Adult heliconiine butterflies host whole-body microbiomes that are typically low in

diversity and evenness. Across all individuals, the median ESV richness was 26, of which only

11 accounted for 95% or more of the sequences. The composition of these communities is also

reasonably consistent within host species. For example, within our most deeply sampled

population (*H. erato demophoon* in Gamboa, Panama; N = 23), a median of 84% of the 16S

rRNA gene reads obtained from a given individual's microbiome belonged to a core set of 10

bacterial genera (Fig. S1). Some of these genera are present in roughly similar relative

abundances across individuals and across our two sampling years (Fig. S1). Applying the

aforementioned calculation to all heliconiine species from which five or more individuals were

sequenced, we found that this intraspecific consistency extends beyond *H. erato demophoon*. For

most host species, over 80% of the constituent individuals' microbiome reads belonged to the 10

most abundant bacterial genera for that species (median = 88%, range = 77-94%).

Microbiomes from whole, homogenized butterfly bodies are mainly composed of gut-

associated taxa. Isolated guts are similar to conspecific whole-body microbiomes in their

bacterial community profiles (Fig. S1), with many bacterial genera occurring at similar relative

abundances in guts and head/thorax tissue (Table S1). Some bacteria did differ between whole-

body and gut samples, such as *Wolbachia*, which usually inhabit reproductive tissues (54) (Fig.

S1). Likewise, in both *Heliconius* and other heliconiine genera, the relative abundance of

Acinetobacter was ~10-fold higher in head/thorax samples as compared with guts (Table S1). In

contrast, *Orbus*, *Enterobacter*, *Asaia*, and some other dominant bacterial genera are enriched in

gut tissue (Table S1).

We used metagenomes (N = 15) to examine the accuracy of amplicon sequencing for

describing bacterial community composition in our broader sample set. Interindividual

microbiome variation was highly correlated between the amplicon-based dataset and the

metagenomic dataset (Mantel r = 0.63, p < 0.001). Furthermore, for six of the most abundant

bacterial genera, relative abundances in amplicon libraries were highly predictive of relative

abundances in metagenomes (Fig. S2). These results support the use of amplicon sequencing for

evaluating bacterial community composition in adult heliconiine butterflies.

Adult *Heliconius* butterflies generally harbor a high absolute abundance of gut

bacteria, though titers sometimes varied substantially even among conspecific individuals

collected from the same location (Fig. 1A). On average, *Heliconius* harbored significantly higher

(by ~5.5-fold) gut bacterial titers than species belonging to other passion-vine butterfly genera (p

= 0.039). Bacterial communities associated with head and thorax tissue did not significantly

differ in titer between *Heliconius* and other genera (p = 0.38; Fig. S3).

Heliconius also harbor compositionally distinct whole-body bacterial communities as

compared with other passion-vine butterfly genera (Fig. 1B). Microbiomes clustered by host

genera to varying degrees depending on the distance metric used. The effect was strongest with

taxonomic or phylogenetic metrics that incorporate information on the relative abundances of

exact sequence variants (ESVs) (Bray-Curtis: $R^2 = 0.13$, p = 0.001; weighted UniFrac: $R^2 = 0.18$,

p = 0.001) as opposed to purely presence/absence-based metrics (Jaccard: $R^2 = 0.11$, p = 0.001;

unweighted UniFrac: $R^2 = 0.10$, p = 0.001). This separation of *Heliconius* microbiomes from

related butterfly genera is not fully explicable by adult-stage diet, as individuals of the non-

pollen-feeding species *Heliconius aoede* mostly clustered with pollen-feeding *Heliconius* species

(Fig. 1B). An analysis of species-by-species Bray-Curtis dissimilarities confirmed that *H. aoede*

microbiomes are not uniquely distinct from those of pollen-feeding *Heliconius* species (Fig. S4).

As suggested by the host genus-level clustering, we found that variation in butterfly

microbiome composition was associated with host relatedness (phylogenetic distance). More

closely related butterfly lineages harbored more similar microbiomes (Bray-Curtis: Mantel r =

0.40, p = 0.01), and the topologies of the butterfly phylogeny and the dendrogram of microbiome

similarity were moderately congruent (Fig. 2). This pattern was still statistically significant when

microbiome variation was measured using weighted UniFrac distances (Mantel r = 0.31, p =

0.03), but not when using unweighted UniFrac (Mantel r = 0.03, p = 0.41) or Jaccard distances

(Mantel r = 0.19, p = 0.08), which do not incorporate relative abundance information.

We then determined which bacterial taxa contribute to the observed taxonomic (Fig.

1B) and phylogenetic (Fig. 2) structuring in overall community composition. The dominant

bacterial genera *Enterobacter*, *Orbus*, and *Entomomonas*/*Pseudomonas* were proportionally

more abundant in *Heliconius* versus the non-pollen-feeding butterfly genera, while *Asaia* and

Apibacter showed the opposite pattern (Fig. 3). None of these bacterial genera, however, were

exclusively restricted to one host feeding guild, genus, or species, and their relative abundances

were occasionally highly variable (Fig. 3). We also analyzed the data at the ESV-level, including

all individual ESVs that were reasonably abundant within at least one host species ($\geq$ 5% within-

species average). We did not find evidence for prevalent host species- or genus-restricted ESVs

(Fig. 4). Most ESVs are present, albeit with varying relative abundances, across host genus and

species boundaries.

Amplicon-derived ESVs are limited in their ability to resolve bacterial strains as they

represent only a short region of the 16S rRNA gene (here, ~250 bp). To test for potential host

specificity at a finer level of resolution, we obtained near-full-length 16S rRNA gene sequences

from the bacterium *Orbus*, which is highly prevalent across gut and whole-body samples (Fig. 3,

Fig. S1) and almost exclusively composed of a single ESV (Fig. 4). These sequences were

reconstructed from the 12 metagenomes in which *Orbus* was sufficiently abundant. A phylogeny

based on these sequences and other Orbaceae shows that butterfly-associated *Orbus* form a

unique clade (Fig. 5). Host-phylogenetic or geographic structure was not evident within this

clade. In fact, many of the Panamanian heliconiine butterflies harbored *Orbus* that have identical

or near-identical 16S rRNA gene sequences to an *Orbus* strain isolated from an East Asian

butterfly, *Sasakia charonda* (55).

We also used the shotgun metagenomes to search for microbial eukaryotes in butterfly

gut samples and found that microsporidia and trypanosomatids were prevalent (27% and 47% of

the 15 individual samples analyzed, respectively) (Fig. S5). Fungi and other microeukaryotic

taxa were only sporadically detected, and at very low relative abundances. When

trypanosomatids were detected in a given sample, we also sometimes detected 16S rRNA gene

reads in the metagenomic data classified as *Kinetoplastibacterium*, an obligate bacterial

endosymbiont of certain trypanosomatids (56). *Kinetoplastibacterium* was not detected without

its corresponding trypanosomatid host (Fig. S5). This observation led us to reexamine the larger,

amplicon-based bacterial dataset. We found that 9% of *Heliconius* individuals and 4% of

individuals belonging to other heliconiine genera were infected with a trypanosomatid, as

inferred by the presence of *Kinetoplastibacterium*. These proportions were not significantly

different (Fisher's Exact Test, $p = 0.28$). They are also likely underestimates, given the

aforementioned prevalence of trypanosomatids among metagenomes (47%) and the potential for

false negatives—in five metagenomes, trypanosomatids were detected without

Kinetoplastibacterium (Fig. S5).

Discussion

We found that adult-stage microbiomes in heliconiine butterflies are generally similar in

composition among conspecific individuals and over time, and are also abundant within gut

tissue (especially in *Heliconius*). Our estimates of total bacterial abundances (via qPCR), in

parallel with those of ref. (23), support the hypothesis that bacteria actively colonize adult

butterflies as opposed to passing through the gut transiently. These features contrast with the

typical situation in lepidopteran larvae (5), reinforcing the idea that insect-microbe associations

can differ strongly between life stages (10).

Although stable relative to larvae, dominant butterfly-associated microbes do exhibit

higher levels of interindividual variability as compared with obligate nutritional endosymbionts

or gut microbes of some other insect groups. For example, in honey bees, >98% of sequences

belong to a honey bee-specific set of 5-9 bacterial species (57). We found that ~80-90% of

sequences in a given butterfly species' microbiome belonged to a set of 10 core bacterial genera.

This contrast suggests that microbiomes are more facultative for butterflies or that there is

greater functional redundancy among butterfly-associated microbes.

We also discovered that *Heliconius* butterfly microbiomes are distinct from those of other

heliconiine genera, in terms of overall community structure (Fig. 1B), relative abundances of

specific bacterial genera (Fig. 3), and total numbers of gut bacteria (Fig. 1A). This finding opens

the possibility that one or more of the ecological traits specific to *Heliconius* influences, and may

be influenced by, the microbiome. As these traits evolved in tandem (18) it is difficult to

disentangle their potential links to microbes, but our analysis of the species *Heliconius aoede*

suggests that pollen-feeding is not a primary driver. *H. aoede* does not pollen-feed (58, 59), yet

its microbiomes were not uniquely distinct from pollen-feeding *Heliconius* species (Fig. 1B, Fig.

S4). This result contrasts with previous work on butterflies finding a strong association between

adult-stage feeding ecology (nectivory versus frugivory) and adult gut microbiomes (23).

Larval feeding ecology, however, warrants further investigation as a potential driver of

variation in adult gut microbiomes. Microbiomes of butterfly larvae and adults are not fully

decoupled: in *H. erato*, some adult-stage microbes appear to be carried over from the larval stage

(11). All *Heliconius* (including *H. aoede*) feed on young passion-vine foliage while other

heliconiine genera feed on mature foliage (18). Related variation in diet-derived microbes, or in

host processes that determine which microbes persist through metamorphosis (10, 60), could

potentially underlie variation in adult-stage microbiomes.

368   In addition to the clear separation of *Heliconius* from other heliconiine genera (Fig.

1B), there was also a strong signal of the host phylogeny in heliconiine butterfly microbiomes

(Fig. 2). The strength of this signal was comparable to that observed in gut microbiomes of

mammals (61) and other host groups (42). This result is also in agreement with a recent survey of

neotropical butterflies, in which microbiomes were found to be phylogenetically structured

across a broad diversity of hosts (six different families) (23). Thus, even in a comparatively

young radiation such as the one studied here (41), host phylogenetic history is clearly an

important factor shaping the composition of adult butterfly microbiomes. The question then

becomes, how has this pattern (also known as phylosymbiosis (62)) arisen, and what does it

signify?

378   There are at least two, non-mutually exclusive processes that can lead to

phylosymbiosis: host-microbe co-diversification, and contemporary host filtering of

environmental microbes (42, 63). In the former process, there is a shared evolutionary history of

host and microbial lineages. This does not necessarily apply to the latter process, which occurs

when host traits that influence microbial colonization, such as diet preference, are conserved to

some degree across the host phylogeny. We suggest that co-diversification is unlikely in this

system given the broad distribution of bacterial phylotypes across host species and genera

(discussed below). Rather, host filtering of environmental microbes likely explains phylogenetic

structure in heliconiine microbiomes, as is the case in a wide variety of other animal groups (42).

A priority for future work on butterflies is to identify the specific traits underlying host filtering.

In Heliconiini, two candidates are larval host plant use (i.e., *Passiflora* species identity and age

of tissue consumed) and adult foraging behavior (especially whether pollen is collected, and

from which plant species), both of which exhibit phylogenetic signal (18, 64, 65).

391   Bacterial community-level variation among butterfly genera and across the phylogeny

was largely driven by shifts in the relative abundance of shared bacterial genera and ESVs, as

opposed to the presence or absence of host-specific bacterial taxa (Figs. 3, 4). Moreover, a finer-

resolution analysis of the nearly ubiquitous bacterium *Orbus* did not find evidence for specificity

within butterflies (Fig. 5). This pattern is notable in part because it weighs against the co-

diversification model described above, and in part because it provides a contrast to the high

degree of host specificity documented in a number of other insect-microbe symbioses (66–69).

While not highly host-specific, at least some of the dominant bacteria in butterflies appear to be

insect specialists as opposed to cosmopolitan environmental taxa. For example, phylogenetic

evidence supports a general insect association for *Entomomonas* (38), *Apibacter* (70), and *Orbus*

(Fig. 5) (71). The ecology of these groups is poorly understood; one important unknown is how

they are transmitted among butterflies, which may be via flowers (72) or other shared resources.

403   Shotgun metagenomes allowed us to corroborate the amplicon-based bacterial data (Fig.

S2) and led to the discovery of putative eukaryotic parasites (trypanosomatids and microsporidia)

in gut tissue of many adult heliconiine butterflies (Fig. S5). Beyond fungi (12, 23), which do not

appear to be abundant in heliconiines, microeukaryotes are almost unknown from adult-stage

Lepidoptera—with the exception of the neogregarine *Ophryocystis elektroscirrha* in monarch

butterflies (73). Given their potential interactions with gut bacteria and relevance to host fitness,

more targeted analyses of butterfly-associated microeukaryote diversity are clearly warranted.

410   While effects of endosymbionts such as *Wolbachia* and *Spiroplasma* have been

documented (74), a major open question is what functional role gut microbes play in the biology

of adult heliconiines and Lepidoptera generally (75). In *Heliconius*, our hypothesis that microbes

mediate pollen-feeding is not strongly supported. To directly influence pollen digestion—which

occurs extraorally using saliva exuded from the proboscis (76)—microbes would likely need to

inhabit the proboscis or salivary gland. Yet we saw no clear signal of elevated microbial

abundances or unique microbial taxa in these tissues in *Heliconius* as compared with non-pollen-

feeding butterflies (Fig. S3). Endogenous pollen-digestion mechanisms (59, 77, 78) may be

sufficient.

Other microbial roles in host nutrition are possible, although a recent experiment on the

butterfly *Speyeria mormonia* did not find evidence for nutrition-related functions (79). We

suggest that colonization resistance (i.e., protection from pathogens and parasites), which is a

feature common to many symbioses (63, 80), could be a primary ecological function of adult

butterfly gut microbiomes. We found that heliconiines are frequently infected by

trypanosomatids and microsporidia, as well as *Serratia* (Fig. 4), which are common opportunistic

bacterial pathogens in insects (81, 82). Adult butterflies may be similar to pyrrhocorid bugs,

tsetse flies and social bees, in which gut bacteria provide an important layer of defense against

related parasites and pathogens (83–87). Importantly, colonization resistance can readily evolve

even in symbioses lacking strong host-microbe specificity (88), such as that between heliconiines

and their adult-stage microbiomes.

Conclusions

This study provides an in-depth characterization of *Heliconius* and other passion-vine

butterfly microbiomes, adding a new dimension to a classic model system in evolutionary

biology. The characteristics of adult-stage microbiomes we report contrast with larval

Lepidoptera, emphasizing that holometabolous insects are able to interact with microbes in very

different ways across life stages (10). However, many Lepidoptera do not feed or even have a

digestive tract as adults (89), and we do not yet know whether patterns from heliconiines or other

butterflies are generalizable to these other groups. We further suggest that heliconiines may serve

as a useful system for exploring how animal ecology and life history relate to microbes.

Heliconius host phylogenetically structured microbiomes that are markedly distinct from those of

their close relatives, and this finding sets the stage for experiments to test the specific host traits

that may be involved. Finally, we uncovered novel microeukaryote diversity in butterflies, and

hypothesize that they are parasites that interact antagonistically with gut bacteria. Further

research on adult-stage microbiomes will help advance our understanding of both insect-

microbiome evolution and of the biology of Lepidoptera, a diverse group of considerable

ecological, societal, and scientific importance (89, 90).

Acknowledgments

451   This material is based on work supported by the Cooperative Institute for Research in

Environmental Sciences, the Ecology and Evolutionary Biology Department at the University of

Colorado Boulder, a National Science Foundation Graduate Research Fellowship (1144083), and

a postdoctoral fellowship from the USDA National Institute of Food and Agriculture (2018-

08156), to TJH. This work was further supported by grants from the Simons Foundation

(429440) and by funding from the Smithsonian Tropical Research Institute. We thank Jessica

Henley for assistance with sequencing, the Smithsonian Tropical Research Institute staff for

administrative support and the Panamanian and Ecuadorian Authorities for permission to collect

butterflies. We also acknowledge the reviewers for helpful comments that improved the

manuscript.

References

1. Buchner P. 1965. Endosymbiosis of animals with plant microorganisms. John Wiley &

Sons.

2. Jones RT, Sanchez LG, Fierer N. 2013. A Cross-Taxon Analysis of Insect-Associated

Bacterial Diversity. PLoS ONE 8:e61218.

3. Engel P, Moran NA. 2013. The gut microbiota of insects – diversity in structure and

function. FEMS Microbiology Reviews 37:699–735.

4. Hammer TJ, Sanders JG, Fierer N. 2019. Not all animals need a microbiome. FEMS

Microbiology Letters 366:fnz117.

5. Hammer TJ, Janzen DH, Hallwachs W, Jaffe SP, Fierer N. 2017. Caterpillars lack a

resident gut microbiome. Proceedings of the National Academy of Sciences 114:9641–

9646.

6. Staudacher H, Kaltenpoth M, Breeuwer JAJ, Menken SBJ, Heckel DG, Groot AT. 2016.

Variability of bacterial communities in the moth *Heliothis virescens* indicates transient

association with the host. PLoS ONE 11:e0154514.

7.   Priya NG, Ojha A, Kajla MK, Raj A, Rajagopal R. 2012. Host Plant Induced Variation in

479   Gut Bacteria of *Helicoverpa armigera*. PLoS ONE 7:e30768.

480   8.   Whitaker M, Pierce N, Salzman S, Kaltenpoth M, Sanders J. 2016. Microbial communities

481   of lycaenid butterflies do not correlate with larval diet. Frontiers in Microbiology 7:1920.

482   9.   Phalnikar K, Kunte K, Agashe D. 2019. Disrupting butterfly caterpillar microbiomes does

483   not impact their survival and development. Proc R Soc B 286:20192438.

484   10.   Hammer TJ, Moran NA. 2019. Links between metamorphosis and symbiosis in

485   holometabolous insects. Phil Trans R Soc B 374:20190068.

486   11.   Hammer TJ, McMillan WO, Fierer N. 2014. Metamorphosis of a Butterfly-Associated

487   Bacterial Community. PLoS ONE 9:e86995.

488   12.   Harrison JG, Urruty DM, Forister ML. 2016. An exploration of the fungal assemblage in

489   each life history stage of the butterfly, *Lycaeides melissa* (Lycaenidae), as well as its host

490   plant *Astragalus canadensis* (Fabaceae). Fungal Ecology 22:10–16.

491   13.   Paniagua Voirol LR, Weinhold A, Johnston PR, Fatouros NE, Hilker M. 2020. Legacy of a

492   Butterfly's Parental Microbiome in Offspring Performance. Appl Environ Microbiol

493   86:e00596-20.

494   14.   Phalnikar K, Kunte K, Agashe D. 2018. Dietary and developmental shifts in butterfly-

495   associated bacterial communities. Royal Society Open Science 5:171559.

15.   Wang X, Sun S, Yang X, Cheng J, Wei H, Li Z, Michaud JP, Liu X. 2020. Variability of

Gut Microbiota Across the Life Cycle of *Grapholita molesta* (Lepidoptera: Tortricidae).

Front Microbiol 11:1366.

16.   Chase AB, Karaoz U, Brodie EL, Gomez-Lunar Z, Martiny AC, Martiny JBH. 2017.

Microdiversity of an Abundant Terrestrial Bacterium Encompasses Extensive Variation in

Ecologically Relevant Traits. mBio 8:e01809-17.

17.   Bates HW. 1861. Contributions to an Insect Fauna of the Amazon Valley. Lepidoptera:

Heliconidae. Transactions of the Linnean Society of London 23:495–566.

18.   Jiggins CD. 2016. The ecology and evolution of *Heliconius* butterflies. Oxford University

Press.

19.   McFall-Ngai M, Hadfield MG, Bosch TCG, Carey HV, Domazet-Lošo T, Douglas AE,

Dubilier N, Eberl G, Fukami T, Gilbert SF, Hentschel U, King N, Kjelleberg S, Knoll AH,

Kremer N, Mazmanian SK, Metcalf JL, Nealson K, Pierce NE, Rawls JF, Reid A, Ruby

EG, Rumpho M, Sanders JG, Tautz D, Wernegreen JJ. 2013. Animals in a bacterial world,

a new imperative for the life sciences. Proceedings of the National Academy of Sciences

110:3229–3236.

20.   Challis RJ, Kumar S, Dasmahapatra KK, Jiggins CD, Blaxter M. 2016. Lepbase: the

Lepidopteran genome database. bioRxiv http://dx.doi.org/10.1101/056994.

21.   Concha C, Wallbank RWR, Hanly JJ, Fenner J, Livraghi L, Rivera ES, Paulo DF, Arias C,

Vargas M, Sanjeev M, Morrison C, Tian D, Aguirre P, Ferrara S, Foley J, Pardo-Diaz C,

Salazar C, Linares M, Massardo D, Counterman BA, Scott MJ, Jiggins CD, Papa R, Martin

A, McMillan WO. 2019. Interplay between Developmental Flexibility and Determinism in

the Evolution of Mimetic Heliconius Wing Patterns. Current Biology 29:3996-4009.e4.

22. van Schooten B, Godoy-Vitorino F, McMillan WO, Papa R. 2018. Conserved microbiota

among young *Heliconius* butterfly species. PeerJ 6:e5502.

23. Ravenscraft A, Berry M, Hammer T, Peay K, Boggs C. 2019. Structure and function of the

bacterial and fungal gut microbiota of Neotropical butterflies. Ecological Monographs

89:e01346.

24. Sculfort O, Castro ECP, Kozak KM, Bak S, Elias M, Nay B, Llaurens V. 2020. Variation of

chemical compounds in wild Heliconiini reveals ecological factors involved in the

evolution of chemical defenses in mimetic butterflies. Ecol Evol 10:2677–2694.

25. Ambika Manirajan B, Ratering S, Rusch V, Schwiertz A, Geissler-Plaum R, Cardinale M,

Schnell S. 2016. Bacterial microbiota associated with flower pollen is influenced by

pollination type, and shows a high degree of diversity and species-specificity: The bacterial

microbiota of flower pollen. Environmental Microbiology 18:5161–5174.

26. Kešnerová L, Emery O, Troilo M, Liberti J, Erkosar B, Engel P. 2020. Gut microbiota

structure differs between honeybees in winter and summer. ISME J 14:801–814.

27. Kingsley VV. 1972. Persistence of intestinal bacteria in the developmental stages of the

monarch butterfly (*Danaus plexippus*). Journal of Invertebrate Pathology 20:51–58.

28. Hammer TJ, Dickerson JC, Fierer N. 2015. Evidence-based recommendations on storing

and handling specimens for analyses of insect microbiota. PeerJ 3:e1190.

29. Gardes M, Bruns TD. 1993. ITS primers with enhanced specificity for basidiomycetes -

application to the identification of mycorrhizae and rusts. Mol Ecol 2:113–118.

30. Baym M, Kryazhimskiy S, Lieberman TD, Chung H, Desai MM, Kishony R. 2015.

Inexpensive Multiplexed Library Preparation for Megabase-Sized Genomes. PLOS ONE

10:e0128036.

31. Martin M. 2011. Cutadapt removes adapter sequences from high-throughput sequencing

reads. EMBnet.journal 17:10–12.

32. Callahan BJ, McMurdie PJ, Rosen MJ, Han AW, Johnson AJA, Holmes SP. 2016.

DADA2: High-resolution sample inference from Illumina amplicon data. Nat Methods

13:581–583.

33. Wang Q, Garrity GM, Tiedje JM, Cole JR. 2007. Naive Bayesian classifier for rapid

assignment of rRNA sequences into the new bacterial taxonomy. Applied and

Environmental Microbiology 73:5261–5267.

34. Quast C, Pruesse E, Yilmaz P, Gerken J, Schweer T, Yarza P, Peplies J, Glöckner FO.

2012. The SILVA ribosomal RNA gene database project: improved data processing and

web-based tools. Nucleic Acids Research 41:D590–D596.

35. R Core Team. 2016. R: A language and environment for statistical computing. R

Foundation for Statistical Computing, Vienna, Austria.

36.   Davis NM, Proctor DM, Holmes SP, Relman DA, Callahan BJ. 2018. Simple statistical

identification and removal of contaminant sequences in marker-gene and metagenomics

data. Microbiome 6:226.

37.   Brady C, Cleenwerck I, Venter S, Coutinho T, De Vos P. 2013. Taxonomic evaluation of

the genus *Enterobacter* based on multilocus sequence analysis (MLSA). Systematic and

Applied Microbiology 36:309–319.

38.   Wang J, Su Q, Zhang X, Li C, Luo S, Zhou X, Zheng H. 2020. *Entomomonas moraniae*

gen. nov., sp. nov., a member of the family Pseudomonadaceae isolated from Asian honey

bee gut, possesses a highly reduced genome. International Journal of Systematic and

Evolutionary Microbiology 70:165–171.

39.   Janssen S, McDonald D, Gonzalez A, Navas-Molina JA, Jiang L, Xu ZZ, Winker K, Kado

DM, Orwoll E, Manary M, Mirarab S, Knight R. 2018. Phylogenetic Placement of Exact

Amplicon Sequences Improves Associations with Clinical Information. mSystems

3:e00021-18.

40.   McDonald D, Price MN, Goodrich J, Nawrocki EP, DeSantis TZ, Probst A, Andersen GL,

Knight R, Hugenholtz P. 2012. An improved Greengenes taxonomy with explicit ranks for

ecological and evolutionary analyses of bacteria and archaea. The ISME Journal 6:610–

618.

41.   Kozak KM, Wahlberg N, Neild AF, Dasmahapatra KK, Mallet J, Jiggins CD. 2015.

Multilocus Species Trees Show the Recent Adaptive Radiation of the Mimetic *Heliconius*

Butterflies. Systematic Biology 64:505–524.

42.   Mazel F, Davis KM, Loudon A, Kwong WK, Groussin M, Parfrey LW. 2018. Is Host

Filtering the Main Driver of Phylosymbiosis across the Tree of Life? mSystems 3:e00097-

18.

43.   Revell LJ. 2012. phytools: An R package for phylogenetic comparative biology (and other

things). Methods in Ecology and Evolution 3:217–223.

44.   Oksanen J, Kindt R, Legendre P. 2019. vegan: Community Ecology Package. R package

version 2.5-6.

45.   Anderson MJ, Walsh DCI. 2013. PERMANOVA, ANOSIM, and the Mantel test in the face

of heterogeneous dispersions: What null hypothesis are you testing? Ecological

Monographs 83:557–574.

46.   Pinheiro J, Bates D, DebRoy S, Sarkar D, R Core Team. 2019. Linear and Nonlinear Mixed

Effects Models.

47.   Joshi, NA, Fass, JN. Sickle: A sliding-window, adaptive, quality-based trimming tool for

FastQ files. https://github.com/najoshi/sickle.

48.   Langmead B, Salzberg SL. 2012. Fast gapped-read alignment with Bowtie 2. Nat Methods

9:357–359.

49.   Gruber-Vodicka HR, Seah BK, Pruesse E. 2019. phyloFlash — Rapid SSU rRNA profiling

and targeted assembly from metagenomes. bioRxiv https://doi.org/10.1101/521922.

50.   Edgar RC. 2004. MUSCLE: Multiple sequence alignment with high accuracy and high

throughput. Nucleic Acids Research 32:1792–1797.

51.   Castresana J. 2000. Selection of Conserved Blocks from Multiple Alignments for Their Use

597   in Phylogenetic Analysis. Molecular Biology and Evolution 17:540–552.

598   52.   Dereeper A, Guignon V, Blanc G, Audic S, Buffet S, Chevenet F, Dufayard J-F, Guindon

599   S, Lefort V, Lescot M, Claverie J-M, Gascuel O. 2008. Phylogeny.fr: robust phylogenetic

600   analysis for the non-specialist. Nucleic Acids Research 36:W465–W469.

601   53.   Guindon S, Delsuc F, Dufayard JF, Gascuel O. 2009. Estimating maximum likelihood

602   phylgenies with PhyML. Methods in Molecular Biology 537:113–137.

603   54.   Saridaki A, Bourtzis K. 2010. Wolbachia: more than just a bug in insects genitals. Current

604   Opinion in Microbiology 13:67–72.

605   55.   Kim JY, Lee J, Shin N-R, Yun J-H, Whon TW, Kim M-S, Jung M-J, Roh SW, Hyun D-W,

606   Bae J-W. 2012. *Orbus sasakiae* sp. nov., a bacterium isolated from the gut of the butterfly

607   *Sasakia charonda*, and emended description of the genus *Orbus*. International Journal of

608   Systematic and Evolutionary Microbiology 63:1766–1770.

609   56.   De Souza W, Motta MCM. 1999. Endosymbiosis in protozoa of the Trypanosomatidae

610   family. FEMS Microbiology Letters 173:1–8.

611   57.   Zheng H, Steele MI, Leonard SP, Motta EVS, Moran NA. 2018. Honey bees as models for

612   gut microbiota research. Lab Animal 47:317–325.

613   58.   Brown KS. 1981. The Biology of *Heliconius* and Related Genera. Annual Review of

614   Entomology 26:427–456.

59.   Gilbert LE. 1972. Pollen Feeding and Reproductive Biology of *Heliconius* Butterflies.

Proceedings of the National Academy of Sciences 69:1403–1407.

60.   Johnston PR, Rolff J. 2015. Host and Symbiont Jointly Control Gut Microbiota during

Complete Metamorphosis. PLOS Pathogens 11:e1005246.

61.   Groussin M, Mazel F, Sanders JG, Smillie CS, Lavergne S, Thuiller W, Alm EJ. 2017.

Unraveling the processes shaping mammalian gut microbiomes over evolutionary time.

Nature Communications 8:14319.

62.   Lim SJ, Bordenstein SR. 2020. An introduction to phylosymbiosis. Proc R Soc B

287:20192900.

63.   Moran NA, Ochman H, Hammer TJ. 2019. Evolutionary and Ecological Consequences of

Gut Microbial Communities. Annu Rev Ecol Evol Syst 50:451-475.

64.   Estrada C, Jiggins CD. 2002. Patterns of pollen feeding and habitat preference among

Heliconius species. Ecological Entomology 27:448–456.

65.   Boggs CL, Smiley JT, Gilbert LE. 1981. Patterns of pollen exploitation by *Heliconius*

butterflies. Oecologia 48:284–289.

66.   Ohkuma M. 2008. Symbioses of flagellates and prokaryotes in the gut of lower termites.

Trends in Microbiology 16:345–352.

67.   Kikuchi Y, Hosokawa T, Nikoh N, Meng X-Y, Kamagata Y, Fukatsu T. 2009. Host-

symbiont co-speciation and reductive genome evolution in gut symbiotic bacteria of

acanthosomatid stinkbugs. BMC Biol 7:2.

68.   Kwong WK, Moran NA. 2015. Evolution of host specialization in gut microbes: The bee

636   gut as a model. Gut Microbes 6:214–220.

637   69.   Kaltenpoth M, Roeser-Mueller K, Koehler S, Peterson A, Nechitaylo TY, Stubblefield JW,

638   Herzner G, Seger J, Strohm E. 2014. Partner choice and fidelity stabilize coevolution in a

639   Cretaceous-age defensive symbiosis. Proceedings of the National Academy of Sciences

640   111:6359–64.

641   70.   Kwong WK, Moran NA. 2016. *Apibacter adventoris* gen. nov., sp. nov., a member of the

642   phylum Bacteroidetes isolated from honey bees. International Journal of Systematic and

643   Evolutionary Microbiology 66:1323–1329.

644   71.   Martinson VG, Carpinteyro-Ponce J, Moran NA, Markow TA. 2017. A Distinctive and

645   Host-Restricted Gut Microbiota in Populations of a Cactophilic *Drosophila* Species.

646   Applied and Environmental Microbiology 83:e01551–17.

647   72.   Graystock P, Goulson D, Hughes WOH. 2015. Parasites in bloom: flowers aid dispersal and

648   transmission of pollinator parasites within and between bee species. Proceedings of the

649   Royal Society B: Biological Sciences 282:20151371.

650   73.   Altizer SM, Oberhauser KS. 1999. Effects of the Protozoan Parasite *Ophryocystis*

651   *elektroscirrha* on the Fitness of Monarch Butterflies (*Danaus plexippus*). Journal of

652   Invertebrate Pathology 74:76–88.

653   74.   Duplouy A, Hornett EA. 2018. Uncovering the hidden players in Lepidoptera biology: the

654   heritable microbial endosymbionts. PeerJ 6:e4629.

75. Paniagua Voirol LR, Frago E, Kaltenpoth M, Hilker M, Fatouros NE. 2018. Bacterial

Symbionts in Lepidoptera: Their Diversity, Transmission, and Impact on the Host. Frontiers

in Microbiology 9:556.

76. Eberhard SH, Hikl AL, Boggs CL, Krenn HW. 2009. Saliva or Regurgitated Nectar? What

Heliconius Butterflies (Lepidoptera: Nymphalidae) Use for Pollen Feeding. Annals of the

Entomological Society of America 102:1105–1108.

77. Harpel D, Cullen DA, Ott SR, Jiggins CD, Walters JR. 2015. Pollen feeding proteomics:

Salivary proteins of the passion flower butterfly, *Heliconius melpomene*. Insect

Biochemistry and Molecular Biology 63:7–13.

78. Smith G, Macias-Muñoz A, Briscoe AD. 2016. Gene Duplication and Gene Expression

Changes Play a Role in the Evolution of Candidate Pollen Feeding Genes in *Heliconius*

Butterflies. Genome Biology and Evolution 8:2581–2596.

79. Ravenscraft A, Kish N, Peay K, Boggs C. 2019. No evidence that gut microbiota impose a

net cost on their butterfly host. Molecular Ecology 28:2100–2117.

80. Kamada N, Chen GY, Inohara N, Núñez G. 2013. Control of pathogens and pathobionts by

the gut microbiota. Nature Immunology 14:685–690.

81. Raymann K, Coon KL, Shaffer Z, Salisbury S, Moran NA. 2018. Pathogenicity of *Serratia*

marcescens Strains in Honey Bees. mBio 9:e01649-18.

82.   Grimont F, Grimont PAD. 2006. The Genus *Serratia*, p. 219–244. *In* Dworkin, M, Falkow,

S, Rosenberg, E, Schleifer, K-H, Stackebrandt, E (eds.), The Prokaryotes. Springer New

York.

83.   Onchuru TO, Martinez AJ, Kaltenpoth M. 2018. The cotton stainer's gut microbiota

suppresses infection of a cotransmitted trypanosomatid parasite. Molecular Ecology

27:3408–3419.

84.   Raymann K, Shaffer Z, Moran NA. 2017. Antibiotic exposure perturbs the gut microbiota

and elevates mortality in honeybees. PLoS Biology 15:e2001861.

85.   Koch H, Schmid-Hempel P. 2011. Socially transmitted gut microbiota protect bumble bees

against an intestinal parasite. Proceedings of the National Academy of Sciences 108:19288–

19292.

86.   Weiss BL, Maltz MA, Vigneron A, Wu Y, Walter KS, O'Neill MB, Wang J, Aksoy S.

2019. Colonization of the tsetse fly midgut with commensal *Kosakonia cowanii* Zambiae

inhibits trypanosome infection establishment. PLOS Pathogens 15:e1007470.

87.   Wu Y, Zheng Y, Chen Y, Chen G, Zheng H, Hu F. 2020. *Apis cerana* gut microbiota

contribute to host health though stimulating host immune system and strengthening host

resistance to *Nosema ceranae*. R Soc Open Sci 7:192100.

88.   McLaren MR, Callahan BJ. 2020. Pathogen resistance may be the principal evolutionary

advantage provided by the microbiome. Phil Trans R Soc B 375:20190592.

89.   Scoble MJ. 1992. The Lepidoptera. Oxford University Press.

90. Boggs CL, Watt WB, Ehrlich PR. 2003. Butterflies: ecology and evolution taking flight.

The University of Chicago Press.

Figures

Figure 1. A) Adult *Heliconius* butterflies have high, though often variable, titers of gut bacteria. Shown
are quantitative PCR-derived estimates of bacterial abundances in terms of the number of bacterial 16S
rRNA gene copies per individual gut. These individuals were collected from Gamboa, Panama in 2016 (N
= 42 *Heliconius*, 16 other Heliconiini). B) *Heliconius* host distinct adult-stage bacterial communities
compared with related genera. Shown is an ordination of microbiome variation (Bray-Curtis
dissimilarities) among all whole-body samples collected at various sites in Panama and Ecuador in 2014
(N = 104 *Heliconius*, 52 other Heliconiini). With the exception of *H. aoede* (dark blue triangles), pollen-
feeding is exclusive to, and ubiquitous within *Heliconius*.

Host phylogeny   Microbiome dendrogram

708
709
710 Figure 2. Variation in heliconiine butterfly microbiomes is correlated with host relatedness. Shown is the
711 concordance between the host phylogeny (from (41)) and a dendrogram representation of microbiome
712 variation among species (Bray-Curtis dissimilarities). Here, all nodes have been rotated to maximize tip
713 matching. Note, however, that sets of parallel lines connecting tips between the phylogeny and the
714 dendrogram do not always signify congruent branching structure.

Figure 3. Dominant bacterial genera are largely shared among *Heliconius* and other butterflies, although some are differentially abundant. Shown are the relative abundances of the top 10 bacterial genera, ranked by mean abundance, in whole-body microbiomes. Dots indicate replicate individuals, and black bars indicate median proportions within a host species. Starred bacterial genera differed significantly in relative abundance between *Heliconius* and non-*Heliconius* butterflies (p < 0.05 after FDR correction). The arrangement of host species on the x axis corresponds to the phylogeny shown in Fig. 4. Note that *Enterobacter* here includes sequences originally assigned as *Klebsiella* and some other closely related Enterobacteriaceae genera (see Methods).

725
726
Figure 4. Most bacterial exact sequence variants (ESVs) are not host species- or host genus-specific. Shown
are all ESVs in the dataset that had ≥ 5% mean relative abundance across conspecific individuals for one
or more host species. For each ESV, the species-level mean relative abundance in whole-body samples,
after log transformation, is indicated by the color of the cells (white = not detected in that species). Note
that bacterial genera (labeled at the top) contained varying numbers of ESVs that met the aforementioned
prevalence threshold.

733
734
Figure 5. Maximum likelihood phylogenetic reconstruction of the bacterial family Orbaceae
(Gammaproteobacteria: Pasteurellales). Some host taxonomic structure is apparent at the order level (i.e.
Lepidoptera, beetles, bees, flies) but not within butterflies. 16S rRNA gene sequences from *Agraulis*,
Dryadula, *Dryas*, and *Heliconius* were assembled from short metagenomic reads. Other 16S rRNA gene
sequences are from GenBank. Branch support values are shown next to nodes. *Haemophilus influenzae*
(Pasteurellaceae) was used as the outgroup.

Supplementary Figures

743
744
Figure S1. Whole-body bacterial communities are consistent among individuals and predominantly
represent gut-associated taxa. Shown are the relative abundances of the top 10 bacterial genera (ranked by
mean abundance) across *H. erato demophoon* collected from Gamboa, Panama. Remaining white space
represents sequences belonging to other genera; these made up a median 16% of sequence libraries across
individuals. Samples on the left are isolated guts from individuals collected in 2016, while samples on the
right are whole-body homogenates from individuals collected in 2014. Note that *Enterobacter* here
includes sequences originally assigned as *Klebsiella* and some other closely related Enterobacteriaceae
genera (see Methods).
753

754
755
756 Figure S2. Bacterial genus-level relative abundances are highly correlated between the amplicon sequence
757 libraries and metagenomes. Shown are relative abundances of six of the most abundant genera across the
758 15 butterfly gut samples for which we obtained both amplicon and shotgun metagenomic sequence data.
759

760
761
762 Figure S3. Absolute abundances of bacteria in the head and thorax of adult heliconiine butterflies, derived
763 from qPCR. Shown are the number of bacterial 16S rRNA gene copies per subsample (approx. 50 mg) of
764 homogenized, combined head and thorax tissue. These individuals were collected from Gamboa, Panama
765 in 2016 (N = 28 *Heliconius*, 16 other Heliconiini).
766
767

768
769
770 Figure S4. The non-pollen-feeding species *Heliconius aoede* is not uniquely distinct from pollen-feeding
771 *Heliconius* species. Whole-body microbiomes from each focal species (x axis) are compared to all other
772 species using Bray-Curtis dissimilarities. For a given comparison of two species' microbiomes,
773 intraspecific replication was handled by averaging dissimilarities among all pairs of individuals.
774

775
776
Figure S5. Microsporidia and Trypanosomatidae (related to *Crithidia*, *Leishmania*, and *Trypanosoma*) are
prevalent in the 15 butterfly individuals with sequenced gut metagenomes. Points show the proportion of
each microeukaryotic taxon out of all non-host 18S rRNA sequences identified by phyloFlash. Circles are
filled if the proportion of microsporidia or trypanosomatids exceeded 0.01. Asterisks indicate samples in
which *Kinetoplastibacterium*, a bacterial endosymbiont of trypanosomatids, was detected in metagenomes.
Other microeukaryotes not shown here comprise a variety of very low-abundance taxa (i.e. ≤ 30 total reads
per metagenome) classified mainly as coccidia, ascomycete fungi, acanthamoeba, and algae.

Supplementary Table

Host taxon	Phylum	Class	Order	Family	Genus	Mean prop. gut +/- SEM	Mean prop. head/thorax +/- SEM	p value
Heliconius	Bacteroidetes	Bacteroidia	Flavobacteriales	Weeksellaceae	*Apibacter*	0.036 +/- 0.009	0.002 +/- 0.001	< 0.001
					Chishuiella	0.007 +/- 0.003	0.095 +/- 0.027	0.001
					Chryseobacterium	0.011 +/- 0.004	0.096 +/- 0.024	0.001
	Firmicutes	Bacilli	Lactobacillales	Enterococcaceae	*Enterococcus*	0.031 +/- 0.008	0.021 +/- 0.006	0.117
				Streptococcaceae	*Lactococcus*	0.02 +/- 0.006	0.033 +/- 0.009	0.793
	Proteobacteria	Alphaproteobacteria	Acetobacterales	Acetobacteraceae	*Asaia*	0.024 +/- 0.007	0.004 +/- 0.002	0.001
					Commensalibacter	0.12 +/- 0.02	0.007 +/- 0.003	< 0.001
					Swaminathania	0.013 +/- 0.01	0.016 +/- 0.014	0.745
		Gammaproteobacteria	Betaproteobacteriales	Burkholderiaceae	*Variovorax*	0.01 +/- 0.004	0.021 +/- 0.006	0.017
			Enterobacteriales	Enterobacteriaceae	*Enterobacter*	0.233 +/- 0.034	0.031 +/- 0.007	< 0.001
					Serratia	0.037 +/- 0.01	0.027 +/- 0.017	0.013
			Orbales	Orbaceae	*Orbus*	0.163 +/- 0.024	0.021 +/- 0.01	< 0.001
			Pseudomonadales	Moraxellaceae	*Acinetobacter*	0.023 +/- 0.007	0.293 +/- 0.041	< 0.001
				Pseudomonadaceae	*Entomomonas/Pseudomonas*	0.086 +/- 0.017	0.024 +/- 0.01	0.291
	Tenericutes	Mollicutes	Entomoplasmatales	Spiroplasmataceae	*Spiroplasma*	0.012 +/- 0.007	0.003 +/- 0.002	0.927
Other Heliconiini	Bacteroidetes	Bacteroidia	Flavobacteriales	Weeksellaceae	*Apibacter*	0.139 +/- 0.045	0.008 +/- 0.004	0.083
					Chishuiella	0.009 +/- 0.003	0.138 +/- 0.029	0.016
					Chryseobacterium	0 +/- 0	0.015 +/- 0.012	0.808
	Firmicutes	Bacilli	Lactobacillales	Enterococcaceae	*Enterococcus*	0.056 +/- 0.027	0.031 +/- 0.011	0.843
				Streptococcaceae	*Lactococcus*	0.036 +/- 0.026	0.012 +/- 0.011	0.098
	Proteobacteria	Alphaproteobacteria	Acetobacterales	Acetobacteraceae	*Asaia*	0.153 +/- 0.039	0.004 +/- 0.001	0.01
					Commensalibacter	0.031 +/- 0.015	0.002 +/- 0.002	0.106
					Swaminathania	0.007 +/- 0.004	0 +/- 0	0.226
		Gammaproteobacteria	Betaproteobacteriales	Burkholderiaceae	*Variovorax*	0.005 +/- 0.002	0.016 +/- 0.004	0.292
			Enterobacteriales	Enterobacteriaceae	*Enterobacter*	0.156 +/- 0.042	0.014 +/- 0.006	0.01
					Serratia	0.02 +/- 0.012	0.001 +/- 0	0.075
			Orbales	Orbaceae	*Orbus*	0.139 +/- 0.043	0.007 +/- 0.004	0.042
			Pseudomonadales	Moraxellaceae	*Acinetobacter*	0.046 +/- 0.018	0.45 +/- 0.054	< 0.001
				Pseudomonadaceae	*Entomomonas/Pseudomonas*	0.02 +/- 0.017	0.011 +/- 0.004	0.393
	Tenericutes	Mollicutes	Entomoplasmatales	Spiroplasmataceae	*Spiroplasma*	0.068 +/- 0.038	0.052 +/- 0.052	0.843

Table S1. Relative abundances and within-body distribution of the top 15 bacterial genera (ranked by mean abundance) across butterflies collected from Gamboa, Panama in 2016. These individuals were dissected to compare microbiomes between isolated gut tissue and the combined head and thorax. Abundances are shown for *Heliconius* (top) and species belonging to other heliconiine genera (bottom) separately. P values are from a nonparametric statistical test of proportions in guts versus head/thorax samples, after FDR correction (see Methods).