Stability of Two-quartet G-quadruplexes and Their Dimers in Atomistic Simulations

Barira Islam,[1] Petr Stadlbauer,[1,2] Michaela Vorlíčková,[1] Jean-Louis Mergny,[1, 3, 4] Michal Otyepka[1,2,5] and Jiří Šponer[1]*

[1]Institute of Biophysics of the Czech Academy of Sciences, Královopolská 135, 612 65 Brno, Czech Republic

[2]Regional Centre of Advanced Technologies and Materials, Faculty of Science, Palacký University, tř. 17 listopadu 12, 771 46, Olomouc, Czech Republic

[3]Institut Curie, PSL Research University, CNRS-UMR 9187, INSERM U1196, F-91405 Orsay, France

[4]Université Paris Sud, Université Paris-Saclay, CNRS-UMR 9187, INSERM U1196, F-91405 Orsay, France

[5]Department of Physical Chemistry, Faculty of Science, Palacký University, tř. 17 listopadu 12, 771 46, Olomouc, Czech Republic

KEYWORDS. G-quadruplex, two-quartet stem, quadruplex stability, MD simulations

ABSTRACT

G-quadruplexes (GQ) are four-stranded non-canonical DNA and RNA architectures that can be formed by guanine-rich sequences. The stability of GQs increases with the number of G-quartets and three G-quartets generally form stable GQs. However, stability of two-quartet GQs is an open issue. To understand the intrinsic stability of two-quartet GQ stems, we have carried out a

series of unbiased molecular dynamics (MD) simulations (~470 µs in total) of two- and four-quartet DNA and RNA GQs, with attention paid mainly to parallel-stranded arrangements. We used AMBER DNA parmOL15 and RNA parmOL3 force fields and tested different ion and water models. DNA two-quartet parallel-stranded GQs unfolded in all the simulations while the equivalent RNA GQ was stable in most of the simulations. GQs composed of two stacked units of two-quartet GQs were, however, stable for both DNA and RNA. The simulations suggest that a minimum of three quartets are needed to form an intrinsically stable all-*anti* parallel-stranded DNA GQ in the presence of Na^+ or K^+ ions. Parallel two-quartet DNA GQ may exist if substantially stabilized by another molecule/structural element (including multimerisation). On the other hand, we predict that isolated RNA two-quartet parallel GQs may form, albeit being weakly stable. We also show that ionic parameters and water models should be chosen with caution, because some parameter combinations can cause spurious instability of GQ stems in simulations. Some in-so-far unnoticed limitations of force-field description of multiple ions inside the GQs are discussed, which appear to compromise capability of simulations to fully capture the effect of the increase of the number of quartets on the GQ stability.

INTRODUCTION

DNA and RNA can form diverse stable secondary structures depending on their sequence and environment.[1, 2] Guanine-rich sequences can adopt G-quadruplex (GQ) structures by forming guanine-guanine Hoogsteen base pairing.[3] Potential GQ-forming sequences have been found at telomeres, gene promoters, immunoglobulin heavy chain switch regions, hypervariable repeats and in genomically unstable and recombination-prone regions.[4] The abundance of GQ-forming sequences at functional and gene controlling sites suggests their potential role in gene

expression.[5] Therefore, there is considerable interest in studying and modulating the formation of GQs as a means for controlling gene expression and other cellular processes.[6]

GQs are formed by stacking of planar guanine quartets, creating the G-stem. The O6 atoms of all the guanines face to form a central electron-rich channel within the GQ where cations bind and coordinate with the O6 atoms. This coordination is essential for GQ stability.[7] Bases not involved in G-stem belong to bulges, flanking nucleotides or participate in the loops linking the G-strands of the GQs.[3]

The GQs can be monomolecular, bimolecular or tetramolecular depending on the number of DNA or RNA molecules that form the GQ.[8, 9] The topologies of GQs are highly variable as the G-strand orientation can be parallel or antiparallel and based on that GQs can fold into parallel, antiparallel or hybrid topology.[10, 11] The guanines in the DNA GQs can adopt *syn* or *anti* orientation.[10, 12, 13] In the RNA GQs, the 2 -hydroxyl group in ribose allows for more interactions with the bases, cations and water molecules thereby increasing their stability, as common in RNA molecules.[14-19] It also restricts the orientation of the guanine bases favouring the *anti*-orientation.[14] Therefore, the majority of the RNA GQs have been observed in the parallel-stranded topology with all bases in the *anti*-orientation. [15-18]

The loops further add to the diversity as the GQs can have propeller, lateral or diagonal loops.[8, 20-27] The parallel-stranded GQ have propeller (double-chain reversal) loops as they are necessary to orient the following and preceding G-strands in a parallel arrangement.[8] The antiparallel and hybrid GQs can have various combinations of the three loop types.[20-27] The overall GQ topology dictates the loop combination as well as the allowable *syn-anti* patterns of the guanines. Na^+ ions tend to coordinate within the plane while K^+ ions coordinate between the planes of the G-

quartets.[28, 29] The cation-quadruplex interactions can induce kinetic and equilibrium differences between the different topologies.[28, 30, 31]

Stability of GQs is affected by the number of G-quartets in the stem.[14, 32-34] It is commonly accepted that increasing the number of G-quartets usually stabilizes the GQ structure.[12, 14, 34, 35] In DNA, stable GQ structures generally involve three quartets or more, although stable antiparallel two-quartet GQs with loop alignments or stacked triads have also been observed.[21, 36 27] In such GQs, additional interactions formed by the loops or triads above and/or below the G-stems contribute to their stabilization.[21, 27, 36] The solution structure of a two-quartet stacked dimer of RNA GQs has been solved, arguing that stabilization is provided by the dimerization of the two GQ units.[37] Various experiments have suggested increased stability of RNA GQs over their DNA counterparts.[14, 38] On the other hand, it was shown that substitution of even one guanine in the three-quartet telomere DNA quadruplex by abasic site or by any non-G base prevents its parallel folding under dehydrating conditions.[39, 40] However, the intrinsic stability of isolated two-quartet G-stems cannot be fully clarified by the experiments, due to difficulty to separate the G-stem stability from the influence of additional interactions as well as due to structural resolution limits of the experiments. Still, understanding stability of two-quartet G-stems is important. They may for example transiently occur during folding processes of larger GQs, in the initial fast phase of folding as well as in later transitions between different GQ folds.[41] Thus, they can have impact on the folding processes. They also may provide regulatory functions due to possible equilibrium between folded and unfolded state at physiological temperatures. In addition, the existence of these two-quartet GQs is of fundamental importance for all bioinformatics searches as their inclusion greatly increases the number of putative GQ-forming sequences in the genome.

The goal of this study was to assess the stability of the smallest two-quartet G-stems, with a special emphasis on the parallel arrangement. We have carried out series of unbiased explicit-solvent molecular dynamics (MD) simulations (total of 470 μs). The two-quartet DNA and RNA G-stems are studied in isolation as well as in the context of GQ dimers. MD methodology, when wisely applied, can provide useful insights into various aspects of GQ structural dynamics and folding.[42] While the contemporary MD is not robust enough to quantitatively evaluate the absolute stability of the GQs in a thermodynamic sense, the simulations should fairly well reflect different relative structural stabilities of GQ arrangements, provided the simulation time scale is sufficiently long to see structural perturbations and unfolding in the simulations, *i.e.*, to assess the lifetimes ($1/k_{off}$) of the studied arrangements. We have carried out the simulations with both Na^+ and K^+ as the stabilizing cations to analyse and compare the cation dynamics. We suggest that the all-*anti* parallel stranded two-quartet DNA G-stem is not stable *per se*, and explain the reason for this behaviour. Nevertheless, such G-stems may still be involved in the formation of larger GQ structures. We also suggest that the RNA two-quartet G-stem is more stable. Simulations were further done for two-quartet DNA antiparallel G-stem to explain the difference in stability of various two-quartet G-stems. To obtain additional insights into the balance of forces in the G-stems and the consequences of the force-field approximations, we have also carried out simulations of tetrameric parallel stranded all-*anti* DNA three and four-quartet G-stem. We compared performance of several water models and ionic parameters. Strikingly, our simulations show that several water-ion combinations lead to visible destabilization of G-stems. Together with our previous study on three-quartet G-stems,[29] these results illustrate the complexity of the choice of water models and ionic parameters in simulations of nucleic acids; it appears that no combination could be generally recommended as the best one. In addition, our

simulations indirectly suggest that the pair-additive molecular mechanics (MM) description generally does not properly capture the stabilizing effect due to increasing the number of G-quartets in the stem.

METHODS

Starting structures

We took the 28-mer two-quartet stacked DNA GQ (PDB id 2N3M[43], first model; Figure 1a), which is a solution structure of a synthetic construct d[(TGG)$_3$TTGTTG(TGG)$_4$T] that exhibits anti-proliferative activity *in vitro*. It consists of a dimer of two-quartet parallel-stranded GQ units stacked by their 3□-5□ interface and connected by a linker nucleotide. The simulations were carried out with the whole 2N3M structure, 2N3M with no bond between G15 and T16 of first and second GQ unit and also for its individual two-quartet GQ units (Figure 1a, 1b and 1c). The individual GQ units are called first and second GQ starting from the 5'-end of 2N3M. The model of parallel-stranded DNA G-stem d[GG]$_4$ was made using the first two quartets of PDB structure 1KF1 (Figure 1d). The model of parallel-stranded DNA G-stem d[GG]$_4$ with all 5'-guanines in *syn* orientation (Figure 1e) was made using the first two quartets of PDB structure 3TVB.[44] The antiparallel DNA G-stem d[GG]$_4$ was obtained from the solution structure of the thrombin binding aptamer (TBA) sequence d[(GGTTGGTGTGGTTGG)] GQ (PDB id 148D, first frame; Figure 1f).[21]

The coordinates for parallel- and antiparallel-stranded three-quartet system d[GGG]$_4$ were obtained from the human telomeric DNA quadruplex (PDB ids 1KF1, chain A and 143D, model 1; Figure 1g and 1h).[8, 22] For four-quartet tetramolecular DNA G-stem d[GGGG]$_4$ simulations, we took the coordinates of guanine core from the X-ray structure of the d(TGGGGT) sequence

(PDB id 4R44, chains A-D; Figure 1i).[45] It is a tetramolecular GQ with three K^+ in the GQ as the stabilizing ions; the structure is virtually identical to other available d[TGGGGT]$_4$ structures.[46]

The coordinates for the dimer and monomer of two-quartet RNA GQ were taken from a solution structure of a dimer of two-RNA GQ units formed by the sequence 5'-r(GGA)$_4$-3' stacked through the 5'-5' interface (PDB id 2RQJ, first model; Figure 1j).[37] The head-to-head stacking means that quartets of the two subunits are of opposite directionality. For simulations of the monomer, we took the first GQ unit (Figure 1k). The antiparallel RNA G-stem r[GG]$_4$ was prepared directly from the DNA variant d[GG]$_4$ (extracted from TBA, see above) by changing deoxyribose into ribose.

MD simulations

The GQs were solvated in a truncated octahedral box with a minimal distance of 10 Å of solute from the box border, mostly in either SPC/E[47] or OPC[48] water model; some simulations were carried out in TIP4P-D water model.[49] The simulations were carried out in 0.15 M excess NaCl or KCl, using mostly SPC/E specific (for SPC/E water model) or TIP4Pew specific (for OPC and TIP4P-D) Joung and Cheatham (JC) parameters.[50] A minority of simulations was performed with Amber-adapted CHARMM22 parameters[51] (for TIP4P-D), SPC/E specific (for SPC/E) and HFE or IOD OPC specific (for OPC) 12-6 LJ parameters developed by Li, Merz and coworkers (LM).[52] The channel cations, when intended, were manually placed between the quartets and in the cavity between the stacked GQs. System preparation was done in xleap module of AMBER16.[53] The list of structures and specific conditions of each simulation are presented in Table 1 and Tables S1 and S2 in the Supporting Information.

We used OL15[54] and OL3 (known also as χ_{OL3})[55] AMBER force field for DNA and RNA GQ, respectively. These force-field versions include also earlier essential DNA[56-58] and RNA[56] modifications of the original AMBER Cornell et al. nucleic acids force-field;[59] for more details see refs.[42, 60]

The starting structures were equilibrated using standard protocols described in the Supporting Information. The simulations were performed with PMEMD CUDA version of AMBER16.[53] The electrostatic interactions were calculated using Particle mesh Ewald method.[61] The cut-off distance for non-bonded interactions was set to 9 Å. Covalent bonds involving hydrogen atoms were constrained using the SHAKE algorithm with a tolerance of 0.0001 Å.[62] Hydrogen mass repartitioning of solute atoms was utilized allowing for an integration time step of 4 fs.[63] Berendsen weak coupling thermostat and barostat were used to maintain constant temperature and pressure of 300 K and 1 atm, respectively.[64] Langevin thermostat with a friction coefficient of 2 ps^{-1} was used in a few simulations. The final production runs without restraints were carried out for continuous length specified in Tables 1-3. Analyses of trajectory were performed using cpptraj[65] module of AMBER and VMD[66] program was used for visualization. Grace (version 5.1.22) program was used to produce the graphs.

Justification of the sampling strategy

We have carried out 114 individual simulations. Rather than trying to repeat small number of equivalent simulations multiple times to get standard statistics of events, we performed as diverse set of simulations as possible. We suggest that although a few individual simulations in the dataset may exhibit some rare behaviour (though we have no indication of that), all the

reported trends consistently emerge from the overall set of simulations and are backed up by multiple simulations.

RESULTS

Dimer of DNA GQs is stable in simulations (Simulations 1a - 1e)

We first performed simulations of the complete 2N3M structure. The sequence d[(TGG)$_3$TTGTTG(TGG)$_4$T] forms a dimer of two-quartet parallel-stranded DNA GQs that are stacked by their 3'-5' region (Figure 1a). The residues G2:G5:G8:G12 and G3:G6:G9:G15 form the guanine quartets of the first GQ unit. T4 and T7 form single nucleotide propeller loops and T10 and T11 form a dinucleotide propeller loop. The residues T13 and T14 form a bulge between the two quartets, while T16 links the two GQ units. In the second GQ unit, G17:G20:G23:G26 and G18:G21:G24:G27 form the guanine quartets and T19, T22 and T25 form single-nucleotide propeller loops. Three cation binding sites in between the subsequent quartets are described further as Site 1, 2 and 3 starting from the 5'-end. We carried out 10 μs long simulations of 2N3M in SPC/E and OPC water models with both Na$^+$ and K$^+$ as the stabilizing ions (see Figure S1 for the RMSD developments). The simulations were initiated with three ions inside the GQ.

The structure of 2N3M was maintained in the 10 μs SPC/E simulations with both Na$^+$ and K$^+$ (Simulations 1a and 1b). In the Simulation 1a, the Na$^+$ ions were present in the three sites (Figure 2a). Two ion exchange events were observed at 9.25 and 9.55 μs. In both events Na$^+$ ion entered from the solvent through the bottom of the stem (the fourth quartet) into the Site 3 (Figure S2a). Specifically, the approach of new cation below the quartet was followed by a linear concerted upward movement of internal ions leading to escape of Na$^+$ ion through the first quartet into the

solvent and movement of the approaching cation into the channel. In the Simulation 1b, the K^+ ion from Site 1 moved up at 1.2 μs and stayed on the top of the first quartet for further 300 ns and then escaped into the solvent. K^+ from Site 2 moved to Site 1. Subsequently, Site 2 remained vacant and no ion intake from the solvent was observed in this simulation (Figure S2b). Transient ion sites were formed just above and below the GQ during the simulation (Figure 2b).

The 2N3M structure was also maintained in the simulation carried out in the Na^+ ions with OPC water model (Simulation 1c). An ion exchange event was observed at 80 ns as a new Na^+ ion entered from the bottom and occupied the Site 3 causing linear movement of cations within the GQ expelling the cation from the top quartet. The second ion exchange event was observed at 6.71 μs as a new Na^+ ion entered into the Site 1 concurrently with linear movement of cations and escape of a cation through the bottom quartet. Within 70 ns of this event, Na^+ ion escaped from Site 3 and Na^+ ion from Site 2 moved to occupy Site 3. No further ion entered the G-stem and only two Na^+ ions remained in the GQ (Figure S2c). However, in one simulation carried out with K^+ as the stabilizing ion (Simulation 1d) the ion from Site 1 escaped into the solvent at 350 ns which was accompanied by perturbation of the first quartet (Figure S2d). There was no concerted linear movement of K^+ ions. At 1.45 μs, the first quartet was disrupted as G5 oriented in the solvent and the top quartet had only three guanines remaining (Figure S1f). The Site 1 remained vacant during the 10 μs long simulation. We carried out additional 10 μs long simulation of 2N3M in OPC/K^+ condition (Simulation 1e). The K^+ ion from Site 1 moved to align in plane with the first quartet within few ns of the start of the simulation and escaped into the solvent at 1.1 μs. This was followed by a subsequent movement of K^+ ion from Site 2 to Site 1. The GQ was stable and Site 2 remained vacant till the simulation end (Figures S1 and S2e).

The results show several things. MD simulations with K^+, at least with the used pair-additive force fields, tend to leave the ion-binding cavity between the two stacked GQs empty. Consistently with literature data,[29, 67] observed ion exchanges with Na^+ are typically concerted, *i.e.*, an ion approach from the bulk to the channel entrance facilitates ion expulsion on the other side of GQ stem accompanied with a concerted shift of all ions. Loss of one quartet in the K^+ OPC simulation, although not confirmed by the second simulation, is consistent with our recent tests showing, for model $d[GGG]_4$ systems, that the OPC water model may understabilize the GQs.[29] In the present case we observe perturbation of GQ core of an experimentally determined structure, which is exceptionally rare in AMBER GQ simulations. [68-72]

Cations can quickly enter the two-quartet dimer through both top and bottom of GQ (Simulations 2a and 2b)

We also carried out 1 μs long simulations of 2N3M in the SPC/E water model with K^+ and Na^+ ions, but initially with no internally bound ions. In the simulation with Na^+ ions in the bulk, an ion was trapped at Site 2 during equilibration. A new ion entered Site 1 after 6 ns of the start of the simulation and pushed ion from Site 2 to Site 3. No further ion exchange was observed. In the K^+ simulation, two K^+ ions positioned near both ends of the G-stem during the equilibration. These ions entered the G-stem within the first ns of the simulation and no further ion exchange was observed. The G-stem was stable in both simulations. While Sites 1 and 3 were occupied, Site 2 remained ion-free (Figure S3).

The linker is not required for the stability of GQ dimer (**Simulations 3a-3b**)

We carried out Na^+ and K^+ SPC/E simulations of 2N3M upon cutting the bond between the G15 and T16, with standard 3'-OH and 5'-OH chain termini and with initially three bound ions. The

GQ dimer stacking was stable. As for the simulations of the complete 2N3M structure, Na$^+$ ions were more dynamic than K$^+$ (Figure S4).[29, 68-70, 73] In the Simulation 3a, three Na$^+$ ion exchange events were observed from 440 ns - 2.28 μs (Figure S4a). These ion exchanges were linked to the entry of cation from the bottom quartet concurrent with a concerted movement leading to the escape of ion from the top quartet. All three sites remained occupied (Figure S4a). In the Simulation 3b, the K$^+$ ion initially resided just below the fourth quartet rather than in Site 3. At 1.08 μs, this ion escaped into the solvent followed by the movement of ion from Site 2 into Site 3 (Figure S4b) and Site 2 remained unoccupied till the end of the simulation. The behaviour of cations was similar thus as in the complete 2N3M GQ.

DNA two-quartet monomer GQs are unstable in simulations (Simulations 4a - 4h)

We carried out simulations of both the GQ units. The two-quartet GQs were unstable and unfolded in both the SPC/E and OPC water models between 80 ns - 7 μs (Table 1). Several unfolding pathways were observed. This may be partly caused by the differences in the ions and water models, though most likely it primarily reflects the known multidimensionality of the free-energy landscape.[42] The first GQ unit unfolded by strand slippage and rotation leading to the formation of cross like structure in Na$^+$ and K$^+$ ions in SPC/E water model and K$^+$ ions in OPC water model (Table 1 and Figure 3). Strand slippage is a movement of a strand by one step up or down, reducing the number of full quartets by one.[74, 75] By cross-like structure we mean a perpendicular or at least significantly inclined arrangement of the G-strands.[74-76] In the simulation with Na$^+$ and OPC water model, unfolding was initiated by disintegration of the second quartet after excessive buckling. The first quartet then disintegrated within 2-3 ns leading to the complete unfolding of GQ.

In the simulations of second GQ unit as well, unfolding was marked by strand slippage and strand rotation in all the simulations (Table 1, Figures S5 and S6). Further details of the simulations are presented in the Supporting Information.

These simulations reflected that *parallel two-quartet DNA GQs are unstable per se as they unfolded in all the ion (K^+ or Na^+) and water conditions.* Obviously, the result can be affected by the force-field approximations (see the Discussion). Still, the unfolding seen in our simulations is so decisive that we predict that all-*anti* parallel-stranded two-quartet DNA GQs are intrinsically unstable in K^+ and Na^+. Hallmarks of the initial stages of unfolding events have been strand slippage and strand rotations.

Tetramolecular two-quartet DNA GQs with 5'-*syn* guanines are more stable than all-*anti* two-quartet DNA GQ (Simulations 5a - 5d, 6a - 6d and 7a - 7k)

We carried out two independent SPC/E simulations of two-quartet parallel stranded d[GG]$_4$ G-stems with all guanines in *anti* orientation in both Na^+ and K^+ ion conditions. The G-stems were highly unstable as strand slippage events followed by structure loss were observed at 5 and 6 ns in the Na^+ ion simulations and at 76 and 228 ns in the two K^+ ion simulations (Figure S7). The strand slippage events were not linked to any ion exchange events.

Simulations were then carried out for parallel-stranded d[GG]$_4$ G-stems with all four 5'-guanines in *syn* orientation (Simulations 6a - 6d). The GQ was maintained in all the simulations (Figure 4a). As many as twenty ion exchange events were observed in the simulations in Na^+ ions while there was no cation exchange in the K^+ ions simulations (Figures 4b-4e). The terminal *syn*-specific 5'-OH – G(N3) H-bond was dominantly sampled in all the simulations (Figure S8), as expected.[12, 77, 78]

We also carried out simulations of two-quartet antiparallel d[GG]$_4$ G-stem which maintained its structure in all five 5 µs long simulations in Na$^+$ ions and SPC/E water model (Simulations 7a - 7e), although there were several (>10 per microsecond) ion-exchange events (Figure 5). The bases in the quartets were staggered for most of the simulation time but the two-quartet arrangement was maintained. In the two 5 µs long simulations in K$^+$ ions and SPC/E water model (Simulations 7f-7g), no cation exchange was observed and GQ was maintained although the quartets were again not entirely planar (Simulations 7a- 7e) (Figure S9).

The two-quartet antiparallel G-stem was also stable in simulations in OPC water model (Simulations 7h-7k). In the 5 µs long simulation in Na$^+$ ions there were continuous ion exchange events (Figure S10) but in the 2 µs long simulation in K$^+$ ions no ion exchange was observed (Figure S11). This behaviour of Na$^+$ and K$^+$ ions was thus consistent across SPC/E and OPC water models (Figures 5, S9-S11).

In summary, all-*anti* parallel-stranded GQs unfolded in all the simulations (Simulations 4a-4h, 5a-5d). The greater stability of both parallel and antiparallel d[GG]$_4$ G-stem with 5□-*syn* bases (Simulations 6a-6d and 7a-7k) suggests that the intramolecular 5□-OH – G(N3) H-bond and the opposite directionality of quartets, hindering the strand slippage stabilize the G-stem.

Dimer of RNA two-quartet GQs is stable (Simulations 8a - 8f, 9a - 9b, 10)

We have carried out six 6-10 µs SPC/E K$^+$ simulations, two SPC/E Na$^+$ simulations and one OPC Na$^+$ simulation (the latter is the potentially least stable combination of parameters) of dimer of two-quartet RNA GQ represented by 2RQJ. Various initial ion positions were considered (Table S1). Universally, all simulations were stable, the ions quickly occupied the Sites 1 and 3, Site 2 was vacant, and no ion exchanges occurred in the whole dataset once the final distribution of the

ions was obtained (Figures S12 - S16). The results are in details described in Supporting Information.

Two-quartet all-*anti* RNA GQ is more stable than the equivalent DNA GQ (Simulations 11a -11d, 12a-12d, 13, 14a-14c, 15a-15c and 16a-16d)

We carried out four 5.5 - 10 μs K$^+$ SPC/E simulations of a monomer of 2RQJ (Simulations 11a-11d). The two-quartet RNA GQ was stable in all the simulations and no ion exchange was observed (Figure 6). The terminal adenine (A12) was stacked below the strand 4 of the GQ as in the starting structure. The loop bases showed some dynamics but could not form any stable alignment with the G-stem. They either aligned in the groove of the GQ by forming transient hydrogen bonds with the G-stem bases or were exposed to the solvent.

We also carried out simulations of a monomer of 2RQJ without the 3' flanking base in Na$^+$ and K$^+$ ions and in both SPC/E and OPC water models (simulation series 12 - 15). The GQ was maintained in three out of four 5 μs long Simulations 12a-12d which included Na$^+$ and SPC/E water model (Figure S17). The GQ unfolded in one simulation at 4.27 μs (Figure S17b). The χ angle of G1 attained *syn* orientation at 4.27 μs and G2 moved towards the solvent. The first propeller loop was lost as G2 moved to stack over G1 but G1 could not re-attain *anti* orientation till the end of the simulation. G1 also sampled the terminal *syn*-specific 5'-OH – G(N3) H-bond when it was in *syn* orientation. More than 20 ion exchange events were observed in all four of these simulations (Figure S18). The GQ also was stable in the 1 μs long K$^+$ SPC/E simulation (Simulation 13). No ion exchange was observed in this simulation and the loop bases showed similar dynamics as in the Simulations 11a-11d (Figure S19).

We carried out three 1 μs long simulations of RNA GQ in Na^+ ions and OPC water model. In Simulation 14a, GQ was stable and loop bases behaved similarly to the equivalent stable SPC/E simulations (Figure S20). An ion exchange event was observed at 750 ns. However, the GQ unfolded in Simulations 14b and 14c (Figure S20). In Simulation 14b, a cross-like structure was formed due to the buckling and rotation of strands 1 and 4 at 400 ns. In Simulation 14c, strand 4 formed a cross-like structure at 130 ns leading to the unfolding of GQ. The unfolding was not linked to any cation exchange movement in these simulations. In contrast, the RNA GQ was stable in three 1 μs K^+ OPC simulations and no cation exchange event was observed (Figure S21).

In brief, simulation series 11-15 suggests *universally enhanced stability of parallel-stranded all anti two-quartet RNA GQs over DNA GQs*. Previous works have shown that 2□-hydroxyl group in ribose sugar allows for more interactions with the bases, cations and water molecules thereby increasing the stability of the RNA GQ.[14, 18] The GQ was stable in all K^+ simulations. The two-quartet RNA GQ was able to maintain its structure in Na^+ SPC/E simulations despite the multiple ion exchange events. The simulations nevertheless reveal that the combination of OPC water model with JC TIP4Pew Na^+ ions destabilizes the RNA GQ compared to the other water-ion combinations. The cation exchange events may alter the energy of GQ due to change in solvation, ion-coordination and structural rearrangements within the GQ.[79] However, the unfolding of GQs in the present OPC Na^+ simulations could not be linked to any cation movements.

The two-quartet RNA GQ represented by monomer of 2RQJ was also a good test system for the TIP4P-D water model. It was stable in three out of four TIP4P-D Simulations 16a-16d (Figure S22). Multiple cation exchanges were observed in the Simulation 16a carried out in JC Na^+ ions

but GQ was maintained throughout the 5 µs simulation (Figures S22a and S23a). The GQ completely unfolded in the Simulation 16b carried out with CHARMM22 Na^+ ions (Figure S22b). Excessive buckling of quartets was observed in this simulation from the start. One cation escaped through the first quartet at 1.8 µs and the GQ then completely unfolded within few ns (Figure S23b). In the Simulations 16c and 16d carried out in JC and CHARMM22 K^+ ions, respectively the GQ was stable and no cation exchange with the solvent was observed (Figures S22c-S22d and S23c-S23d).

Substitution of thymine by adenine loop bases does not stabilize the two-quartet DNA GQ (Simulation 17)

To confirm that the difference in structural stability of DNA and RNA two-quartet GQ was not due to the loop bases, we substituted the thymine by adenine bases in the second GQ monomer of 2N3M and carried out one simulation in K^+, SPC/E simulation (Simulation 17). The GQ unfolded at 450 ns (Figure S24).

Tetramolecular antiparallel RNA G-stem model was unstable in the simulations (Simulations 18a - 18d)

The model of hypothetical tetramolecular two-quartet antiparallel RNA G-stem unfolded in all the simulations in Na^+ and K^+ ions in SPC/E water model. The quartets were not planar and the guanines were buckled from the start of the simulations. The GQ unfolded in the two Na^+ simulations at 10 and 50 ns by strand slippage followed by strand rotation (Figure S25a and S25b).The GQ unfolded at ~180 ns and ~510 ns in the two K^+ simulations (Figures S25c and S25d).

Tetramolecular three-quartet antiparallel DNA G-stem is more stable than parallel-stranded G-stem (Simulations 19a - 19b, 20a - 20b and 21a - 21d)

In order to see the effect on adding one quartet (and ion), we have carried out a set of parallel-stranded d[GGG]$_4$ simulations. In the two Na$^+$ SPC/E simulations, strand slippage was observed at 1.6 μs and 530 ns, respectively (Figures S26a and S26b). However, the G-stem was entirely stable in the two 5 μs long K$^+$ SPC/E simulations (Figures S26c and S26d). In Simulation 20a, a channel K$^+$ was exchanged at the 5'-end at 4.8 μs while in the Simulation 20b it was exchanged at the 3'-end at 273 ns (Figures S27b). In both cases, the cation-leaving site was vacant for ~25-30 ns followed by entry of new cation from the bulk at the same site. Thus, using the SPC/E water model the G-stem was stable in K$^+$ ions but have shown a strand slippage in Na$^+$ ion simulations.

The d[GGG]$_4$ antiparallel G-stem was fully stable in SPC/E simulations with both ion types (Figure S28). Four ion exchange events were observed in the Na$^+$ simulation while no ion exchange was observed in the K$^+$ simulation (Figure S29). In the Na$^+$ OPC simulation (Simulation 21c), ion between the middle and top quartet escaped into the solvent at 65 ns. This resulted in instability of the quartets and G12 from the first quartet moved towards the solvent at 185 ns (Figure S28a). However, the G-stem was maintained in the Simulation 21d with K$^+$ ions in OPC water model (Figure 28b).

To summarize the simulations, the all-*anti* parallel stranded three-quartet G-stem was more stable than similar two-quartet G-stem as no structural change was observed in the K$^+$ SPC/E simulations of d[GGG]$_4$ compared to the rapid complete unfolding of all the parallel d[GG]$_4$ simulations. There is, however, a tendency for strand slippage in presence of Na$^+$. The

antiparallel d[GGG]$_4$ is evidently more stable than the all-*anti* parallel-stranded d[GGG]$_4$. Additional 30 µs of simulations of the all parallel all-*anti* [GGG]$_4$ can be found in ref.;[29] the results are consistent with the present simulations.

Simulations of tetramolecular DNA four-quartet parallel stranded all-*anti* G-stem show excessive ion dynamics (Simulations 22a - 22e, 23a - 23e, 24a - 24e, 25a - 25e and 26a - 26g)

In order to obtain more insights into the balance of forces in GQs and also to better understand the force-field limitations, we have carried out a series of simulations of parallel stranded all-*anti* d[GGGG]$_4$ tetrameric G-stem with three ions in the channel. The d[GGGG]$_4$ simulations can be compared with the 2N3M simulations (simulation series 1-3) and with d[GGG]$_4$ simulations (simulation series 19-21). The d[GGGG]$_4$ all parallel all-*anti* G-stem has not been studied extensively before by long time-scale MD and, intuitively, one would expect it to be very stable. However, the results show a surprisingly complex picture.

The G-stem was maintained in all ten 1 µs long simulations in the SPC/E water model with either Na$^+$ (Simulations 22a-22e) or K$^+$ (Simulations 23a-23e) ascertaining that four quartets provide sufficient stabilization, consistently with the literature. However, in four out of five Na$^+$ simulations, three to seven ion exchange events were observed (Figure S30a and S30c-30d) while no ion exchange was seen in one simulation (Figure S30b). All the ion movements were concerted. The Na$^+$ ions in the channel were mobile, confirming that the stems sample diverse subtly different heterogeneous substates, interconnecting the detailed ion dynamics/distribution with modest differences in the internal G-stem conformation.[77] The Na$^+$ ions were mostly residing close to the quartet planes rather than in between the quartet planes (Figure S31).

Although the G-stems were stable, in all the SPC/E simulations with K^+ there has been a visible tendency to reduce the number of the ions in the channel to two, indicating a structural stress associated with the three closely-spaced cations (see Discussion). This has not been observed in the above (simulation series 19-21) and earlier simulations of the d[GGG]$_4$ with the same force-field parameters,[29] *i.e.*, two K^+ are stable in d[GGG]$_4$. In early stages of the d[GGGG]$_4$ K^+ simulations, the ion between quartet 1 and 2 (Site 1) was pushed above the stem (above its 5'-end). After the first loss of cation from the Site 1, K^+ ion could re-occupy Site 1 in only one simulation out of five (Figure S32e). Otherwise the cations stayed above the first quartet throughout the simulations (Figures 7 and S32). As this newly formed "outer" ion-binding site was exposed to the solvent, four to >20 cation exchanges with bulk ions were observed in the simulations (Figure S32). Additionally, in two simulations, the bottom ion was lost to the solvent. Then the ion from the central cavity Site 2 moved to Site 3 and the middle site remained vacant for most of the Simulations 23d and 23e (Figures S32d and S32e), thus resulting in only two ions in the stem. The K^+ ions that remained within the G-stems were stable and stayed in the cavities between two quartets, as expected for the K^+ ions.

We also carried out five 1 μs long simulations of d[GGGG]$_4$ in OPC with Na^+ as the stabilizing cation (Figure S33). The systems were visibly distorted in three out of five simulations as strand slippage events were observed. In one simulation the GQ was stable and no ion exchange event was observed while in the last simulation, four cation exchange events were observed but the stem was stable. The strand slippage events in this series of simulations were initiated by melting of the terminal quartet. One guanine left the quartet, stacked on the remaining G-triad, and it eventually enforced strand slippage, as the lone stacked guanine pulled the whole G-strand by one level (Figure S34). Substantially non-planar G-quartets were sampled from the onset of the

simulations (because of the apparently imbalanced G(O6)-Na$^+$ attraction resulting in buckling of Gs), but that was likely a prelude to the described strand slippage. Considering the short simulation time scale, the simulations thus suggest a substantial volatility of the d[GGGG]$_4$ system compared to what one would expect for a four-quartet G-stem (see Discussion).

In the 1 μs long simulations of d[GGGG]$_4$ in K$^+$ and OPC water model, the G-stem was stable in only two out of five simulations (Figure S35). In the Simulations 25a - 25c, Site 1 became vacant and transient cation binding site over the first quartet was observed. The first (5'-end) quartet in these simulations was disrupted at 180 ns, 280 ns and 350 ns, respectively by movement of one or two guanines over the remaining bases of the quartet. The other three quartets of the G-stem did not fully disintegrate but exhibited substantial fluctuations. The cations at Sites 2 and 3 were retained in these simulations (Figure S36a - S36c). In the Simulation 25d, the number of ions in the stem was reduced to two, the Site 2 became vacant and additional transient cation binding sites were formed above and below the G-stem (Figure S36d). In the Simulation 25e, the ion from Site 1 was pushed above the first quartet. Re-entry of this cation back into the Site 1 pushed the cation from Site 3 below the G-stem and when the latter ion returned back to the Site 3 the ion from Site 1 moved again above the stem (Figure 36e). An ion exchange was also observed at Site 3. This all indicates difficulty to stable incorporate three ions in the channel.

We also tested LM Na$^+$ and K$^+$ ions in the SPC/E water model in the Simulations 26a-26c for the d[GGGG]$_4$ G-stem. The G-stem was stable in the Na$^+$ ion simulation but disruption of a quartet was observed in one of the two simulations in K$^+$ ions, (Simulation 26b, Figure S37a). The SPC/E adapted LM Na$^+$ ions showed similar behaviour as the JC Na$^+$ ions as three ions stabilized the G-stem and ~20 ion exchange events were observed (Figure S38a). In the Simulation 26b with the SPC/E adapted LM K$^+$ ions, ion from Site 3 was lost to the solvent at 90 ns and guanine

from the last quartet moved towards the solvent, disrupting the quartet. The other three quartets were maintained albeit with substantial fluctuations. In the second simulation in SPC/E adapted LM K$^+$ ions (Simulation 26c), ion from Site 1 was pushed over the first quartet until the movement of ion from Site 2 to Site 1 at 0.7 μs (Figure S38b). Cations at Sites 1 and 3 stabilized the G-stem and no cation was present in Site 2 for the remaining of the simulation time. Thus, the LM Na$^+$ and K$^+$ ions in Simulations 26a and 26c behaved similarly to JC ions.

In the Simulation 26d with OPC water model and HFE Na$^+$ ions, GQ was stable but various ion exchange events were observed (Figures S37b and S38c) and the fourth quartet was buckled. In the OPC water model and HFE K$^+$ ions (Simulation 26e), 5'-quartet was disrupted at 1.7 μs (Figures S37b and S38d). Disruption of 5'-quartet was also observed in the Simulations 26f and 26g with OPC water model and IOD Na$^+$ and K$^+$ ions at 1.0 μs and 400 ns, respectively (Figure S37c).

Additional simulations for testing various parameters (Simulations 27a-27d and 28a-28f)

Further parallel-stranded d[GGGG]$_4$ simulations were carried out with TIP4P-D water with JC and CHARMM22 Na$^+$ and K$^+$ ions (simulation series 27). The d[GGGG]$_4$ G-stem was fully stable in only two out of four simulations in TIP4P-D water model. Strand slippage was observed in the Simulation 27a in JC Na$^+$ ions at ~3 μs (Figure S39). However, the G-stem was stable in the Simulations 27b and 27c in CHARMM22 Na$^+$ ions and JC K$^+$ ions, respectively. In the simulation with CHARMM22 Na$^+$ ions, cations were present in all the three sites and ion exchanges were observed while in the simulation with JC K$^+$ ions, cations were retained at Sites 1 and 3 and Site 2 was vacant for majority of the simulation time. Disruption of 5'-quartet was observed in the Simulation 27d with CHARMM22 K$^+$ ions at ~1.3 μs as cation escaped from the

top quartet (Figure S40d). The quartet reassembled at 3.6 μs and full four-quartet G-stem was formed again (Figure S39d). Na^+ ions were more dynamic than K^+ ions (Figure S40). The perturbation of quartets and excessive ion dynamics suggest that combination of cations and TIP4P-D water parameter tested here is probably less suitable for GQ simulations than SPC/E with JC ions.

We also carried out additional simulations of two quartet parallel and antiparallel GQ (Simulations 28a -28f) using the Langevin thermostat. The GQ system behaved similarly as with the Berendsen thermostat, within the limits of sampling (Figure S41 and S42). Further details of these simulations are presented in the Supporting Information.

Complex dynamics of water molecules inside the channel of four-quartet GQs (Simulation series 1-3, 8-10, 22-27)

The trajectories of both RNA and DNA four-quartet GQs showed that water molecules may enter the channel via its 5' or 3' end. These waters may then be pushed further into the channel by an incoming cation. Then they may leave the channel concurrently with another ion exchange through the channel ends. In case of four-quartet GQs with discontinuous strands (2RQJ, 2N3M), we rarely observed that waters could escape from Site 2 (the middle one) into the groove. This process is facilitated by incorporation of the leaving water molecule into a G-quartet, which then hosts one water-mediated G:G base pair (Figure S43). Such a G-quartet perturbation may last for microseconds, but it apparently does not destabilize the GQ. In one case we saw the process in reverse, a water molecule penetrated into the channel from the groove. In tetramolecular GQs with continuous strands (d[GGGG]$_4$), water molecules could escape the channel via incorporation into a quartet, but only the terminal ones; we did not see escape through the middle

quartets into the groove. The analysis again confirms that the ensembles of the simulated G-stem structures are not homogeneous[77] and contain diverse long-living substates. Interaction of long-residency water molecules in G-triads participating in a G-stem core has been characterized by NMR and MD simulations.[80] However, incorporation of water molecules into a G-quartet appears to be a new observation. It may lead to rare conformational substates of the G-stems which might affect their interactions with other molecules.

DISCUSSION

Potential roles of two-quartet GQs

GQs are often assumed to need three consecutive G-quartets for stabilization. On the other hand, the formation of stable two-quartet DNA or RNA GQs would imply that GQs could be far more widespread in the genome.[81] In addition, RNA is single-stranded and in principle could form GQ whenever the right conditions are met.[82-88] It has been shown that RNA GQs formed by 5'-untranslated regions of genes and mRNA could modulate translation *in vitro*.[89-93] The stability of two-quartet RNA GQs would implicate that shorter RNA molecules such as microRNAs may form GQs under physiological conditions and modulate the cellular functions.[94] In addition, these less stable GQs could have interesting dynamic properties, as they can quickly fold and unfold, and sensitively respond to the changes in cellular conditions. Various two-quartet GQs could be transiently formed during folding of larger GQs. The current experimental data are not sufficient to unambiguously assess the intrinsic stability of two-quartet GQs, albeit they show that very stable two-quartet GQs can form if stabilized by additional structural features such as favourable loop alignments or adjacent G-triads. Two most known cases are the DNA Thrombin binding aptamer (TBA) stabilized by the loops and the two-quartet form of the DNA human

telomeric GQ stabilized by a G-triad,[21, 27] both being anti-parallel GQs. More recent two-quartet DNA GQs have also been discovered in anti-parallel conformation.[36, 95]

Stability of two-quartet GQs as captured by MD simulations

In the present work, we investigated the potential to form two-quartet GQs with minimal or no support by additional stabilizing elements. We have performed series of standard 1-10 μs long atomistic explicit solvent MD simulations assessing the intrinsic structural stability of two-quartet DNA and RNA GQs starting from the folded state. Such computations aim to capture the unfolding process, i.e., the lifetime ($1/k_{off}$) of a given GQ when folded, as described by the simulation force field. Although the computations do not address directly the thermodynamic stability, it is plausible to assume that species with too short lifetimes cannot significantly contribute to the folding landscape. Due to many force-field approximations discussed below we assume that the simulations underestimate the lifetimes of the studied species, but should provide fairly good estimate of their relative structural stabilities. Most of our efforts were devoted to the more elusive parallel-stranded all-*anti* arrangements.

Our simulations predict that *two-quartet parallel stranded all-anti DNA G-stems are intrinsically very unstable*, and cannot be formed (except of being very transient) without extensive support by other structural elements (Simulations 4a-4h and 5a-5d, Figures 3, S5 and S7). Parallel-stranded G-stems with 5'-*syn* bases and antiparallel two-quartet DNA G-stem are evidently more stable (Simulations 6a-6d and 7a-7k, Figures 4, 5 and S9-S11). This is consistent with the fact that the solution structures of antiparallel two-quartet DNA GQs with alignments on both sides of the G-stems are known.[21] Similarly, the two-quartet RNA GQ is predicted to be more stable than the parallel DNA two-quartet GQ, which is consistent with experiments suggesting larger

stability of RNA over DNA GQs (simulation series 11 -15, Figure 6 and S17-S21).[38, 82, 85, 96] Mergny and colleagues studied the kinetics and thermodynamics of tetramolecular DNA and RNA quadruplexes, and concluded that RNA quadruplexes are more stable than their DNA counterparts as a result of both faster association and slower dissociation.[34] Note, however, that all these comparisons were based on tetramolecular structures involving at least three quartets.

The simulations clearly reveal the origin of short lifetimes of unsupported two-quartet all-*anti* DNA G-stems (Table 1). They are volatile due to the ability of unobstructed vertical movements (slippage) of the strands caused by common thermal fluctuations.[74, 75] In contrast to longer stems, for two-quartet stems any strand slippage leads to a weak structure with only one quartet that is prone to immediate disintegration. In case of antiparallel G-stems, mutual vertical movements of the strands are hindered by the mixture of *syn* and *anti* residues in the stem. Additionally, the terminal *syn*-specific 5'-OH – G(N3) H-bond may also provide stability to the antiparallel G-stem. In case of RNA GQs, we suggest that their larger rigidity due to the presence of the 2 □-OH hydroxyl group reduces their propensity to vertical strand slippage.

Force-field approximations and their effect on G-stem simulations

The simulations are influenced by diverse force-field imbalances.[60] Generally, outcomes of simulation studies of fully folded cation-stabilized GQ structures are often unaffected by these imbalances, due to the extraordinary stability and stiffness of GQs.[42] Simulations with appropriate force fields starting from fully folded cation-stabilized GQs are usually entirely stable, as their life-times, even as described by the force field, extend far beyond the simulation time scale. In contrast, any larger perturbation of the quartets in simulations of GQ structures derived from experiments indicate that an inadequate force field has been used.[42, 97] Still, even

with appropriate force fields, subtle details of the structures (details of quartet's H-bonds,[29, 72] cation distribution,[29] backbone substates[57] and loop conformations [68, 69, 71]) may be sampled imperfectly.

Force-field imbalances may impact considerably more simulations of GQ folding intermediates[42, 76] and other weak structures such as the two-quartet stems studied here. Such simulations may be already visibly affected by force-field approximations. Our above-presented analysis of different stabilities of two-quartet GQ stems has been made based on primary MD data with consideration of the known systematic errors in the force-field description. Thus, we suggest that our interpretation of the simulation data is justified. Let us now explicitly comment on the imbalances, explain how they influence the presented primary simulation results and what new insights were learnt from the simulations about force-field limitations.

i) Stability of the GG H-bonding is generally assumed to be underestimated[42] with the presently-used non-bonded terms.[60] This problem can be partly corrected, for example, by adding a specific term to stabilize the H-bonds.[98] Note that it does not mean that the H-bonding is uniformly underestimated in MD simulations of nucleic acids; there are other H-bonds which are over-stabilized by the force field, especially in RNA.[60, 98] No means to enhance stability of the GG base pairing was used in this study.

ii) There are many indications that the base stacking is often over-stabilized in simulations of nucleic acids with the presently-used non-bonded terms, though this effect is likely context-dependent and non-uniform (reviewed in ref.[60]). For GQs, some over-stabilization of G-quartet stacking may be helpful since it compensates for some other force-field deficiencies. Stability of

H-bonding and stacking in nucleic acids simulations is determined not only by the solute-solute interactions, but mainly by the balance with the water model.[99-101]

iii) The force fields are suspected to provide inaccurate description of the propeller loops.[68, 70, 71] They are unable to sample the experimentally-determined (if known unambiguously)[68] propeller loop geometries in otherwise stable simulations of fully folded GQs and are assumed to excessively destabilize the parallel G-hairpins and G-triplexes which affects simulations of folding intermediates.[41, 76] The exact origin and magnitude of this imbalance remains unknown.[68] The description of lateral and diagonal loops appears to be more realistic. [69]

iv) With commonly used force-field parameters, the pair additive force field underestimates the strength of the O6-cation interaction and thus also direct cation binding to the G-quartets.[71, 102] In addition, when using monovalent cations that have parameters optimized to reproduce solvation energies with current water models, the cations inside the G-stem channel appear as too large (having a too large radius or too steep onset of the van der Waals repulsion).[71, 102] These inaccuracies, together with approximate description of the base-base H-bonds, probably contribute to common sampling of bifurcated GG H-bonds in the simulated G-stems.[29, 72]

v) When two or more cations are lining up in the G-stem, yet another and likely severe imbalance arises, which has been noticed only quite recently.[103] It is manifested as an excessive inter-cation repulsion, because the ions are modelled as having fixed +1 charges. In real GQs, the inter-cation repulsion is weakened due to a complex interplay of polarization effects between the ions and the G-quartets, which is strongly geometry-dependent.[103] This problem thus cannot be fixed when using the formalism of pair-additive force fields.

Due to the latter two imbalances, we remain rather sceptical with respect to MD simulation studies aiming at quantitative (free-energy) description of ion binding inside GQs, including studies attempting to explain the Na^+ *vs*. K^+ stabilization difference. In addition, as shown here and elsewhere,[77] binding of ions in GQs includes diverse local substates which have a dramatic impact on results of free-energy calculations using the approximate MM-PBSA (or similar) methods. These methods magnify the free-energy differences among the substates while the simulation time scale is not sufficient to provide their converged populations.[77] Recent literature data suggest that polarizable force fields could substantially improve the description of ions in GQs,[97] though they may still require a careful calibration and it is not yet clear if they do not introduce other hitherto not identified problems. However, G-stems stabilized by cations are genuine systems for application of the polarizable force fields.

The above-listed imbalances explain why we suggest that the structural stability of the two-quartet GQs is underestimated by the present simulations. It is affected by under-stabilization of the ion-quartet interactions and the GG base pairs, which is only partly compensated for by the likely overestimated stacking. Further, the intra- and bimolecular parallel GQs may be destabilized by the description of the propeller loops.

Our simulations of the larger systems are further affected by the imbalanced description of the inter-cation repulsion. This can contribute to the observation that in the simulations of the 2N3M structure, the ion Site 2 is often not occupied (Table 1). This specific prediction should thus be taken with care.

Consideration of the excessive inter-cation repulsion also explains the quite frequent and likely excessive exchanges and ion shifts seen in simulations of the d[GGGG]$_4$ all parallel tetrameric

GQ (Figure 7). It also explains why the ion exchanges are not attenuated in the present d[GGGG]$_4$ simulations compared to the earlier simulations of d[GGG]$_4$.[29] While at first sight one might expect a major stabilization/rigidification of the system when moving from three quartets to four quartets, the force-field imbalance of the inter-cation interaction is getting more severe for the latter system. *In other words we suggest that the fact that we see more ion dynamics, exchanges and expulsions for the d[GGGG]$_4$ than for d[GGG]$_4$ is a simulation artefact.* Note that in some of our simulations we have even seen modest perturbations of the stems. It indicates that simulations with pair-additive force-fields do not properly capture the increase of stability of GQs upon the increase of the number of G-quartets.[34, 104] Although direct calculation of free-energy differences between GQs with different number of quartets are not realistic due to sampling limitations, even structural-dynamics simulations can be affected when smooth lining up of the multiple ions in the channel is disturbed, as evidenced above. Fortunately, there are indications that the ion dynamics is attenuated in simulations of complete GQs compared to simulations of just the G-stems.[29, 77] We did not see any preference for cation exchanges through the 5'-end and 3'-end quartets. Previous MD studies also indicated that in all-*anti* parallel-stranded GQ, the energy barriers for cations escaping the G-stem is approximately equal for both the 5'- and 3'-end.[105]

The above-noted imbalances also provide clues about the less satisfactory performance of the four-site OPC and TIP4P-D water models in our simulations. The OPC water model is becoming very popular in simulations of RNA molecules, since it appears to weaken the base stacking and also spurious H-bonds in RNA simulations;[60, 98-100] overstabilization of these interactions is a result of the balance of solute-solvent and solvent-solvent interactions. TIP4P-D enhances dispersion interactions and it was developed to balance the populations of compacted and loosely

ordered states of disordered proteins, because the widely used three-site TIP3P model overstabilizes the compacted state.[49] However, as shown in the present work, the OPC and TIP4P-D water models (at least with most of the used ion parameters) are risky for GQ simulations and we do not recommend them. They can lead to severe destabilization of at least some GQ systems, probably due to the loss of fortunate compensation of errors present with some more simple water models such as SPC/E. It appears that finding a universal water model for nucleic acids simulations is a complex problem and that the SPC/E water model with JC ions is a reasonable albeit not ideal compromise for GQ simulations.

CONCLUSIONS

Nearly 250 GQ structures are currently accessible in the PDB database. The majority of the crystallised GQs are parallel-stranded and have three quartets. Solution structures of antiparallel two-quartet DNA GQs have also been observed.[21, 36] The antiparallel two-quartet DNA GQs have extensive loop alignments above and below the guanine quartets or other stabilizing factors which could provide stability to the GQ. However, it remains to be explored if independent parallel-stranded two-quartet DNA GQs are unstable *per se* or are yet to be discovered.

Various experimental evidences suggest larger stability of RNA over DNA GQs. The 2□ - hydroxyl group in ribose sugar allows for more interactions with the bases, cations and water molecules thereby increasing the stability of the RNA GQ.[14] It also restricts the orientation of the base favouring the *anti*-orientation. Therefore, the majority of the RNA GQs have been observed in the parallel-stranded topology with all bases in the *anti*-orientation.

Based on the simulations we suggest that parallel-stranded all-*anti* two-quartet DNA G-stems are intrinsically very unstable due to the propensity of strand slippage, leading to very short life-

times. Such stems can be stably formed only with the help of additional stabilizing interactions, such as dimerization and linked two-quartet parallel-stranded DNA GQs.[43] Nevertheless, it does not preclude occurrence of parallel-stranded two-quartet stems in transitory ensembles during folding processes. Antiparallel DNA two-quartet stems as well as RNA two-quartet stems are structurally considerably more stable than the parallel DNA two-quartet stems, although even these species likely require additional stabilizing elements to achieve a detectable thermodynamic stability.

On the methodological side, we identify a visible consequence of exaggerated inter-cation repulsion in the G-stems. This leads to enhanced rate of cation exchanges and even expulsions of ions in the d[GGGG]$_4$ parallel all-*anti* stem compared to the d[GGG]$_4$. This has been an entirely unexpected and counterintuitive result, considering the experimentally known sharp increase of GQ stability with the number of G-quartets.[34] Although there is no direct experimental evidence of ion dynamics in these constructs, we assume that on µs time scale the ion-exchange dynamics in d[GGGG]$_4$ should be minimal and the stem should be fully occupied by ions. The simulations also show differences among the tested water and ion parameters, confirming earlier observation[29] that improved water models such as OPC may be non-optimal for simulations of GQs.

Figure 1. Starting structures used in the simulations. (a) 2N3M GQ with nucleotide numbering of guanines, (b) two-quartet GQ extracted from upper half of 2N3M, (c) two-quartet GQ extracted from lower half of 2N3M, (d) all *anti* parallel-stranded d[GG]₄ G-stem, (e) parallel-stranded d[GG]₄ with 5'-guanines in *syn* orientation, (f) antiparallel d[GG]₄ extracted from 148D, (g) parallel-stranded d[GGG]₄ from 1KF1, (h) antiparallel d[GGG]₄ from 143D and (i) parallel-stranded d[GGGG]₄ from 4R44 with cation binding sites 1, 2 and 3 shown by red, blue and

green circles, respectively. The RNA simulations were carried out using the dimer of (j) 2RQJ with nucleotide numbering of guanines and (k) monomer of 2RQJ GQ. The *anti* and *syn* guanines are shown in yellow and orange, respectively. The 5'-end of RNA structures is shown in tan. The loops are shown in thin lines while dark arrows show the backbone of GQ. The yellow, magenta and orange circles show guanine, thymine and adenine bases, respectively. The solid red lines represent characteristic Hoogsteen base pairing.

Figure 2. Cation binding sites (densities) observed in the MD simulations of 2N3M. (a) Three Na$^+$ ions stabilized the GQ in the Simulation 1a while (b) two K$^+$ ions stabilized the GQ for the majority of the Simulation 1b. The Na$^+$ and K$^+$ binding sites are shown in orange and pink, respectively. The backbone of the GQ is shown in tan tube. The guanine nucleotides of GQ are shown in silver ribbons and blue lines. The backbone of 5'-end of the GQ is shown in cyan. In the simulation in K$^+$ ions, transient cation binding sites were developed above and below the GQ. A 2500 Å^{-3} threshold was used for density representation.

Figure 3. Simulations of the first GQ unit of 2N3M. The two-quartet structure unfolded in all the Simulations 4a-4d irrespective of the water model and stabilizing ion suggesting the instability of this two-quartet DNA GQ. The (a) starting structure and main transient structure occurring during the unfolding in (b) Na$^+$/SPC/E, (c) K$^+$/SPC/E, (d) Na$^+$/OPC and (e) K$^+$/OPC are shown. See legend to Figure 1 for further details. The solid red lines represent characteristic Hoogsteen base pairing while dash red lines indicate any other hydrogen bonds.

Figure 4. Backbone RMSD (a) and distance plots monitoring the ion binding (d-e) in parallel-stranded d[GG]$_4$ with all 5'- bases in *syn* configuration in the Simulations 6a - 6d. Simulations 6a

and 6b were carried out with Na^+ ions and SPC/E water model while Simulations 6c and 6d were carried out with K^+ ions and SPC/E water model. The distances were measured between centre of mass of G-stem and the cations in Simulations (b) 6a, (c) 6b, (d) 6c and (e) 6d. In the panels (b)-(e) each color represents different cation in the simulations. The centrally bound cation shows < 1 Å distance from the centre of mass of the G-stem. Frequent cation exchange events were observed in the simulations carried out with Na^+ ions while there was no cation exchange in the simulations carried out with K^+ ions. In the panels b and c cations around 3 Å represent cations just above or below the G-stem.

Figure 5. Backbone RMSD (a) and distance plots monitoring ion binding (b-f) of antiparallel d[GG]$_4$ G-stem in the Simulations 7a – 7e in the presence of Na$^+$ ions and the SPC/E water

model. Distance plot between cations and centre of mass of the G-stem reveal very fast cation exchanges in all the simulations. The cation within < 1 Å represents the internally bound cation. Each color represents a different cation within the plot. In the panels b - f cations around 3 Å represent cations just above or below the G-stem.

Figure 6. Distance/backbone RMSD plots showing stability of two-quartet RNA GQ 2RQJ in the Simulations 11a - 11d carried out in the presence of K^+ and SPC/E water model. The backbone RMSD of the GQ is shown in black while the red graph indicates distance between the centre of mass of the GQ and the channel cation between the two quartets. No significant fluctuation in RMSD and no exchange of channel cation were observed. In panel a, the minor fluctuation observed in backbone RMSD at ~4.5 µs is due to the movement of the loops.

Site 1
Site 2
Site 3

Figure 7. Cation binding sites (densities) observed in the Simulations (a) 23a, (b) 23b, (c) 23c, (d) 23d and (e) 23e of d[GGGG]$_4$ in the presence of K$^+$ ions and SPC/E water model. The backbone of the GS is shown in tan tube and the 5'-end of the GQ backbone is shown in cyan. The guanine nucleotides are shown in silver lines and the K$^+$ binding sites within the G-stem are shown in blue. The cation densities were developed above the GS due to the movement of cation from Site 1 to above the 5'-quartet. Cation densities were also developed below the GS as solvent cations were transiently trapped below the 3'-quartet. The figure shows that mostly two cations were present in the channel due to repulsion between closely spaced cations. A 300 Å^{-3} threshold was used for density representation.

Table 1. List of 2N3M and two-quartet DNA GQ simulations carried out in the present study

Simulation Number	subcategory	Structure	Length of simulation (μs)	Conditions (force field /ion/ water model, initial ion position)[a]	Summary
1	a	2N3M	10	OL15/Na$^+$/SPC/E	three ions stabilized the GQ, two ion exchange events seen after 9 μs.
	b		10	OL15/K$^+$/SPC/E	two ions stabilized the GQ, the middle ion is missing
	c		10	OL15/Na$^+$/OPC	three ions stabilized GQ from 0-6.71 μs and two ions from 6.71-10 μs (middle ion missing)
	d		10	OL15/K$^+$/OPC	first quartet was disrupted as there was no ion in between quartets 1 and 2 and therefore a guanine from the first quartet moved towards the solvent at 1.45 μs.
	e		10	OL15/K$^+$/OPC	K$^+$ from Site 1 escaped into the solvent at 1.1 μs and ion from Site 2 moved to Site 1. Two ions stabilized GQ for remaining time, middle ion missing.
2	a	2N3M	1	OL15/Na$^+$/SPC/E; initially no ions inside	Ion trapped at Site 2 during equilibration. New ion entered at Site 1 at 6 ns of the simulation and pushed ion from Site 2 to Site 3. No further ion exchange was observed; Site 2 remained vacant.
	b		1	OL15/K$^+$/SPC/E; initially no ions inside	Ions positioned near the G-stem (both above and below) during the equilibration. These two ions entered the G-stem within the first ns of the simulation and no further ion exchange was observed. Site 2 remained vacant.
3	a	2N3M without linker	5	OL15/Na$^+$/SPC/E	GQ stable; three Na$^+$ ion exchange events were observed from 440 ns - 2.28 μs.

	b		5	OL15/K$^+$/SPC/E	GQ stable; no ion exchange. Two K$^+$ ions stabilized the GQ for majority of the simulation time.
4	a	first GQ unit of 2N3M	10	OL15/Na$^+$/SPC/E	unfolded at ~400 ns
	b		10	OL15/K$^+$/SPC/E	unfolded at 1.83 μs
	c		10	OL15/Na$^+$/OPC	unfolded at 1.75 μs
	d		10	OL15/K$^+$/OPC	unfolded at 6.58 μs
	e	second GQ unit of 2N3M	10	OL15/Na$^+$/SPC/E	unfolded at ~880 ns
	f		10	OL15/K$^+$/SPC/E	unfolded at ~4.65 μs
	g		10	OL15/Na$^+$/OPC	unfolded at ~85 ns
	h		10	OL15/K$^+$/OPC	unfolded at ~1.3 μs
5	a-d	parallel stranded all *anti* d[GG]$_4$	2 (2x simulations)	OL15/Na$^+$/SPC/E	strand slippage at 5 ns and 6 ns, then structure loss
			2 (2x simulations)	OL15/K$^+$/SPC/E	strand slippage at 76 ns and 228 ns, then structure loss
6	a-d	parallel stranded d[GG]$_4$ with 5'-*syn*	1 (2x simulations)	OL15/Na$^+$/SPC/E	G-stem was stable but frequent ion exchanges were observed in both the simulations
			1 (2x simulations)	OL15/K$^+$/SPC/E	G-stem was stable; no ion exchange
7	a-k	antiparallel d[GG]$_4$ (from PDB 148D)	5 (5x simulations)	OL15/Na$^+$/SPC/E	Several ion exchange events but G-stem was stable.
			5 (2x simulations)	OL15/K$^+$/SPC/E	G-stem stable; no ion exchange
			5 (2x simulations)	OL15/Na$^+$/OPC	Several ion exchange events but G-stem was stable.
			2 (2x simulations)	OL15/K$^+$/OPC	G-stem stable; no ion exchange

[a] fully occupied channel at the beginning if not mentioned otherwise. JC ions used in all simulations.

ASSOCIATED CONTENT

Supporting Information. Equilibration protocol, detailed results of Simulations 4a - 4h, simulation series 8 - 10 and Simulations 28a - 28f are presented in the Supporting Information. Tables S1 - S2 and Figures S1 - S43 supporting the results are also presented in the Supporting Information. The following files are available free of charge. Supporting_Information.pdf

AUTHOR INFORMATION

Corresponding Author

*Prof. Jiri Sponer. Tel: 420 541 517 133; Fax: +420 541 212 179; Email: sponer@ncbr.muni.cz

Author Contributions

The manuscript was written through contributions of all authors. All authors have given approval to the final version of the manuscript.

Funding Sources

This work was supported by the project SYMBIT reg. number: CZ.02.1.01/0.0/0.0/15_003/0000477 financed by the ERDF. JS also acknowledges support by Praemium Academiae.

ACKNOWLEDGMENT

B.I. and P.S. greatly appreciate access to storage facilities owned by parties and projects contributing to the National Grid Infrastructure MetaCentrum, provided under the programme

"Projects of Large Research, Development, and Innovations Infrastructures" (CESNET LM2015042).

REFERENCES

1.   Bochman, M. L.; Paeschke, K.; Zakian, V. A., DNA secondary structures: stability and function of G-quadruplex structures. *Nature reviews. Genetics* **2012**, 13, 770-80.
2.   Gilbert, D. E.; Feigon, J., Multistranded DNA structures. *Current opinion in structural biology* **1999**, 9, 305-14.
3.   Burge, S.; Parkinson, G. N.; Hazel, P.; Todd, A. K.; Neidle, S., Quadruplex DNA: sequence, topology and structure. *Nucleic acids research* **2006**, 34, 5402-15.
4.   Sen, D.; Gilbert, W., Formation of parallel four-stranded complexes by guanine-rich motifs in DNA and its implications for meiosis. *Nature* **1988**, 334, 364-6.
5.   Mao, S. Q.; Ghanbarian, A. T.; Spiegel, J.; Martinez Cuesta, S.; Beraldi, D.; Di Antonio, M.; Marsico, G.; Hansel-Hertsch, R.; Tannahill, D.; Balasubramanian, S., DNA G-quadruplex structures mold the DNA methylome. *Nature structural & molecular biology* **2018**, 25, 951-957.
6.   Guilbaud, G.; Murat, P.; Recolin, B.; Campbell, B. C.; Maiter, A.; Sale, J. E.; Balasubramanian, S., Local epigenetic reprogramming induced by G-quadruplex ligands. *Nature chemistry* **2017**, 9, 1110-1117.
7.   Gellert, M.; Lipsett, M. N.; Davies, D. R., Helix formation by guanylic acid. *Proceedings of the National Academy of Sciences of the United States of America* **1962**, 48, 2013-8.
8.   Parkinson, G. N.; Lee, M. P.; Neidle, S., Crystal structure of parallel quadruplexes from human telomeric DNA. *Nature* **2002**, 417, 876-80.
9.   Sket, P.; Plavec, J., Tetramolecular DNA quadruplexes in solution: insights into structural diversity and cation movement. *Journal of the American Chemical Society* **2010**, 132, 12724-32.
10.   Webba da Silva, M., Geometric formalism for DNA quadruplex folding. *Chemistry* **2007**, 13, 9738-45.
11.   Phan, A. T., Human telomeric G-quadruplex: structures of DNA and RNA sequences. *The FEBS journal* **2010**, 277, 1107-17.
12.   Cang, X.; Sponer, J.; Cheatham, T. E., 3rd, Explaining the varied glycosidic conformational, G-tract length and sequence preferences for anti-parallel G-quadruplexes. *Nucleic acids research* **2011**, 39, 4499-512.
13.   Dvorkin, S. A.; Karsisiotis, A. I.; Webba da Silva, M., Encoding canonical DNA quadruplex structure. *Science advances* **2018**, 4, eaat3007.
14.   Sacca, B.; Lacroix, L.; Mergny, J. L., The effect of chemical modifications on the thermal stability of different G-quadruplex-forming oligonucleotides. *Nucleic acids research* **2005**, 33, 1182-92.
15.   Xu, Y.; Kaminaga, K.; Komiyama, M., G-quadruplex formation by human telomeric repeats-containing RNA in Na+ solution. *Journal of the American Chemical Society* **2008**, 130, 11179-84.
16.   Martadinata, H.; Phan, A. T., Structure of propeller-type parallel-stranded RNA G-quadruplexes, formed by human telomeric RNA sequences in K+ solution. *Journal of the American Chemical Society* **2009**, 131, 2570-8.
17.   Martadinata, H.; Phan, A. T., Structure of human telomeric RNA (TERRA): stacking of two G-quadruplex blocks in K(+) solution. *Biochemistry* **2013**, 52, 2176-83.

18. Collie, G. W.; Haider, S. M.; Neidle, S.; Parkinson, G. N., A crystallographic and modelling study of a human telomeric RNA (TERRA) quadruplex. *Nucleic acids research* **2010**, 38, 5569-80.

19. Sweeney, B. A.; Roy, P.; Leontis, N. B., An introduction to recurrent nucleotide interactions in RNA. *Wiley Interdisciplinary Reviews: RNA* **2015**, 6, 17-45.

20. Dai, J.; Punchihewa, C.; Ambrus, A.; Chen, D.; Jones, R. A.; Yang, D., Structure of the intramolecular human telomeric G-quadruplex in potassium solution: a novel adenine triple formation. *Nucleic acids research* **2007**, 35, 2440-50.

21. Schultze, P.; Macaya, R. F.; Feigon, J., Three-dimensional solution structure of the thrombin-binding DNA aptamer d(GGTTGGTGTGGTTGG). *Journal of molecular biology* **1994**, 235, 1532-47.

22. Wang, Y.; Patel, D. J., Solution structure of the human telomeric repeat d[AG3(T2AG3)3] G-tetraplex. *Structure* **1993**, 1, 263-82.

23. Lim, K. W.; Ng, V. C.; Martin-Pintado, N.; Heddi, B.; Phan, A. T., Structure of the human telomere in Na+ solution: an antiparallel (2+2) G-quadruplex scaffold reveals additional diversity. *Nucleic acids research* **2013**, 41, 10556-62.

24. Dai, J.; Carver, M.; Punchihewa, C.; Jones, R. A.; Yang, D., Structure of the Hybrid-2 type intramolecular human telomeric G-quadruplex in K+ solution: insights into structure polymorphism of the human telomeric sequence. *Nucleic acids research* **2007**, 35, 4927-40.

25. Luu, K. N.; Phan, A. T.; Kuryavyi, V.; Lacroix, L.; Patel, D. J., Structure of the human telomere in K+ solution: an intramolecular (3 + 1) G-quadruplex scaffold. *Journal of the American Chemical Society* **2006**, 128, 9963-70.

26. Phan, A. T.; Kuryavyi, V.; Luu, K. N.; Patel, D. J., Structure of two intramolecular G-quadruplexes formed by natural human telomere sequences in K+ solution. *Nucleic acids research* **2007**, 35, 6517-25.

27. Lim, K. W.; Amrane, S.; Bouaziz, S.; Xu, W.; Mu, Y.; Patel, D. J.; Luu, K. N.; Phan, A. T., Structure of the human telomere in K+ solution: a stable basket-type G-quadruplex with only two G-tetrad layers. *Journal of the American Chemical Society* **2009**, 131, 4301-9.

28. Schultze, P.; Hud, N. V.; Smith, F. W.; Feigon, J., The effect of sodium, potassium and ammonium ions on the conformation of the dimeric quadruplex formed by the Oxytricha nova telomere repeat oligonucleotide d(G(4)T(4)G(4)). *Nucleic acids research* **1999**, 27, 3018-28.

29. Havrila, M.; Stadlbauer, P.; Islam, B.; Otyepka, M.; Šponer, J., Effect of Monovalent Ion Parameters on Molecular Dynamics Simulations of G-Quadruplexes. *Journal of chemical theory and computation* **2017**, 13, 3911-3926.

30. Marchand, A.; Gabelica, V., Folding and misfolding pathways of G-quadruplex DNA. *Nucleic acids research* **2016**, 44, 10999-11012.

31. Hardin, C. C.; Henderson, E.; Watson, T.; Prosser, J. K., Monovalent cation induced structural transitions in telomeric DNAs: G-DNA folding intermediates. *Biochemistry* **1991**, 30, 4460-4472.

32. Payet, L.; Huppert, J. L., Stability and structure of long intramolecular G-quadruplexes. *Biochemistry* **2012**, 51, 3154-61.

33. Bardin, C.; Leroy, J. L., The formation pathway of tetramolecular G-quadruplexes. *Nucleic acids research* **2008**, 36, 477-88.

34. Mergny, J.-L.; De Cian, A.; Ghelab, A.; Saccà, B.; Lacroix, L., Kinetics of tetramolecular quadruplexes. *Nucleic acids research* **2005**, 33, 81-94.

35.   Lee, J. Y.; Yoon, J.; Kihm, H. W.; Kim, D. S., Structural diversity and extreme stability of unimolecular Oxytricha nova telomeric G-quadruplex. *Biochemistry* **2008**, 47, 3389-96.
36.   Lenarčič Živković, M.; Rozman, J.; Plavec, J., Adenine-Driven Structural Switch from a Two- to Three-Quartet DNA G-Quadruplex. *Angewandte Chemie International Edition* **2018**, 57, 15395-15399.
37.   Mashima, T.; Matsugami, A.; Nishikawa, F.; Nishikawa, S.; Katahira, M., Unique quadruplex structure and interaction of an RNA aptamer against bovine prion protein. *Nucleic acids research* **2009**, 37, 6249-58.
38.   Arora, A.; Maiti, S., Differential biophysical behavior of human telomeric RNA and DNA quadruplex. *The journal of physical chemistry. B* **2009**, 113, 10515-20.
39.   Kejnovska, I.; Bednarova, K.; Renciuk, D.; Dvorakova, Z.; Skolakova, P.; Trantirek, L.; Fiala, R.; Vorlickova, M.; Sagi, J., Clustered abasic lesions profoundly change the structure and stability of human telomeric G-quadruplexes. *Nucleic acids research* **2017**, 45, 4294-4305.
40.   Bednářová, K.; Kejnovská, I.; Vorlíčková, M.; Renčiuk, D., Guanine Substitutions Prevent Conformational Switch from Antiparallel to Parallel G-Quadruplex. *Chemistry – A European Journal* **2019**, 0.
41.   Stadlbauer, P.; Kührová, P.; Vicherek, L.; Banáš, P.; Otyepka, M.; Trantírek, L.; Šponer, J., Parallel G-triplexes and G-hairpins as potential transitory ensembles in the folding of parallel-stranded DNA G-Quadruplexes. *Nucleic acids research* **2019**, 47, 7276-7293.
42.   Šponer, J.; Bussi, G.; Stadlbauer, P.; Kührová, P.; Banáš, P.; Islam, B.; Haider, S.; Neidle, S.; Otyepka, M., Folding of guanine quadruplex molecules–funnel-like mechanism or kinetic partitioning? An overview from MD simulation studies. *Biochimica et Biophysica Acta (BBA) - General Subjects* **2017**, 1861, 1246-1263.
43.   Do, N. Q.; Chung, W. J.; Truong, T. H. A.; Heddi, B.; Phan, A. T., G-quadruplex structure of an anti-proliferative DNA sequence. *Nucleic acids research* **2017**, 45, 7487-7493.
44.   Clark, G. R.; Pytel, P. D.; Squire, C. J., The high-resolution crystal structure of a parallel intermolecular DNA G-4 quadruplex/drug complex employing syn glycosyl linkages. *Nucleic acids research* **2012**, 40, 5731-8.
45.   Mandal, P. K.; Collie, G. W.; Kauffmann, B.; Huc, I., Racemic DNA crystallography. *Angewandte Chemie* **2014**, 53, 14424-7.
46.   Laughlan, G.; Murchie, A. I.; Norman, D. G.; Moore, M. H.; Moody, P. C.; Lilley, D. M.; Luisi, B., The high-resolution crystal structure of a parallel-stranded guanine tetraplex. *Science* **1994**, 265, 520-4.
47.   Berendsen, H. J. C.; Grigera, J. R.; Straatsma, T. P., The missing term in effective pair potentials. *The Journal of Physical Chemistry* **1987**, 91, 6269-6271.
48.   Izadi, S.; Anandakrishnan, R.; Onufriev, A. V., Building Water Models: A Different Approach. *The Journal of Physical Chemistry Letters* **2014**, 5, 3863-3871.
49.   Piana, S.; Donchev, A. G.; Robustelli, P.; Shaw, D. E., Water Dispersion Interactions Strongly Influence Simulated Structural Properties of Disordered Protein States. *The Journal of Physical Chemistry B* **2015**, 119, 5113-5123.
50.   Joung, I. S.; Cheatham, T. E., 3rd, Determination of alkali and halide monovalent ion parameters for use in explicitly solvated biomolecular simulations. *The journal of physical chemistry. B* **2008**, 112, 9020-41.
51.   MacKerell, A. D.; Bashford, D.; Bellott, M.; Dunbrack, R. L.; Evanseck, J. D.; Field, M. J.; Fischer, S.; Gao, J.; Guo, H.; Ha, S.; Joseph-McCarthy, D.; Kuchnir, L.; Kuczera, K.; Lau, F. T. K.; Mattos, C.; Michnick, S.; Ngo, T.; Nguyen, D. T.; Prodhom, B.; Reiher, W. E.; Roux, B.;

Schlenkrich, M.; Smith, J. C.; Stote, R.; Straub, J.; Watanabe, M.; Wiórkiewicz-Kuczera, J.; Yin, D.; Karplus, M., All-Atom Empirical Potential for Molecular Modeling and Dynamics Studies of Proteins. *The Journal of Physical Chemistry B* **1998**, 102, 3586-3616.

52.   Li, P.; Song, L. F.; Merz, K. M., Systematic Parameterization of Monovalent Ions Employing the Nonbonded Model. *Journal of chemical theory and computation* **2015**, 11, 1645-1657.

53.   Case, D.; Betz, R.; Cerutti, D. S.; Cheatham, T.; Darden, T.; Duke, R.; Giese, T. J.; Gohlke, H.; Götz, A.; Homeyer, N.; Izadi, S.; Janowski, P.; Kaus, J.; Kovalenko, A.; Lee, T.-S.; LeGrand, S.; Li, P.; Lin, C.; Luchko, T.; A. Kollman, P., *Amber 16, University of California, San Francisco*. 2016.

54.   Zgarbova, M.; Sponer, J.; Otyepka, M.; Cheatham, T. E., 3rd; Galindo-Murillo, R.; Jurecka, P., Refinement of the Sugar-Phosphate Backbone Torsion Beta for AMBER Force Fields Improves the Description of Z- and B-DNA. *Journal of chemical theory and computation* **2015**, 11, 5723-36.

55.   Zgarbová, M.; Otyepka, M.; Sponer, J.; Mládek, A.; Banáš, P.; Cheatham, T. E., 3rd; Jurečka, P., Refinement of the Cornell et al. Nucleic Acids Force Field Based on Reference Quantum Chemical Calculations of Glycosidic Torsion Profiles. *Journal of chemical theory and computation* **2011**, 7, 2886-2902.

56.   Perez, A.; Marchan, I.; Svozil, D.; Sponer, J.; Cheatham, T. E., 3rd; Laughton, C. A.; Orozco, M., Refinement of the AMBER force field for nucleic acids: improving the description of alpha/gamma conformers. *Biophysical journal* **2007**, 92, 3817-29.

57.   Krepl, M.; Zgarbova, M.; Stadlbauer, P.; Otyepka, M.; Banas, P.; Koca, J.; Cheatham, T. E., 3rd; Jurecka, P.; Sponer, J., Reference simulations of noncanonical nucleic acids with different chi variants of the AMBER force field: quadruplex DNA, quadruplex RNA and Z-DNA. *Journal of chemical theory and computation* **2012**, 8, 2506-2520.

58.   Zgarbova, M.; Luque, F. J.; Sponer, J.; Cheatham, T. E., 3rd; Otyepka, M.; Jurecka, P., Toward Improved Description of DNA Backbone: Revisiting Epsilon and Zeta Torsion Force Field Parameters. *Journal of chemical theory and computation* **2013**, 9, 2339-2354.

59.   Cornell, W. D.; Cieplak, P.; Bayly, C. I.; Gould, I. R.; Merz, K. M.; Ferguson, D. M.; Spellmeyer, D. C.; Fox, T.; Caldwell, J. W.; Kollman, P. A., A Second Generation Force Field for the Simulation of Proteins, Nucleic Acids, and Organic Molecules. *J. Am. Chem. Soc.* **1995**, 117, 5179-5197.

60.   Sponer, J.; Bussi, G.; Krepl, M.; Banas, P.; Bottaro, S.; Cunha, R. A.; Gil-Ley, A.; Pinamonti, G.; Poblete, S.; Jurecka, P.; Walter, N. G.; Otyepka, M., RNA Structural Dynamics As Captured by Molecular Simulations: A Comprehensive Overview. *Chemical reviews* **2018**, 118, 4177-4338.

61.   Darden, T.; York, D.; Pedersen, L., Particle mesh Ewald: An N·log(N) method for Ewald sums in large systems. *J. Chem. Phys.* **1993**, 98, 10089-10092.

62.   Ryckaert, J.-P.; Ciccotti, G.; Berendsen, H. J. C., Numerical integration of the cartesian equations of motion of a system with constraints: molecular dynamics of n-alkanes. *Journal of Computational Physics* **1977**, 23, 327-341.

63.   Hopkins, C. W.; Le Grand, S.; Walker, R. C.; Roitberg, A. E., Long-Time-Step Molecular Dynamics through Hydrogen Mass Repartitioning. *Journal of chemical theory and computation* **2015**, 11, 1864-74.

64. Berendsen, H. J. C.; Postma, J. P. M.; van Gunsteren, W. F.; DiNola, A.; Haak, J. R., Molecular dynamics with coupling to an external bath. *The Journal of Chemical Physics* **1984**, 81, 3684-3690.

65. Roe, D. R.; Cheatham, T. E., 3rd, PTRAJ and CPPTRAJ: Software for Processing and Analysis of Molecular Dynamics Trajectory Data. *Journal of chemical theory and computation* **2013**, 9, 3084-95.

66. Humphrey, W.; Dalke, A.; Schulten, K., VMD: Visual molecular dynamics. *Journal of Molecular Graphics* **1996**, 14, 33-38.

67. Akhshi, P.; Wu, G., Umbrella sampling molecular dynamics simulations reveal concerted ion movement through G-quadruplex DNA channels. *Physical Chemistry Chemical Physics* **2017**, 19, 11017-11025.

68. Islam, B.; Stadlbauer, P.; Gil-Ley, A.; Pérez-Hernández, G.; Haider, S.; Neidle, S.; Bussi, G.; Banas, P.; Otyepka, M.; Sponer, J., Exploring the Dynamics of Propeller Loops in Human Telomeric DNA Quadruplexes Using Atomistic Simulations. *Journal of chemical theory and computation* **2017**, 13, 2458-2480.

69. Islam, B.; Stadlbauer, P.; Krepl, M.; Havrila, M.; Haider, S.; Sponer, J., Structural Dynamics of Lateral and Diagonal Loops of Human Telomeric G-Quadruplexes in Extended MD Simulations. *Journal of chemical theory and computation* **2018**, 14, 5011-5026.

70. Islam, B.; Stadlbauer, P.; Krepl, M.; Koca, J.; Neidle, S.; Haider, S.; Sponer, J., Extended molecular dynamics of a c-kit promoter quadruplex. *Nucleic acids research* **2015**, 43, 8673-93.

71. Fadrna, E.; Spackova, N.; Sarzynska, J.; Koca, J.; Orozco, M.; Cheatham, T. E., 3rd; Kulinski, T.; Sponer, J., Single Stranded Loops of Quadruplex DNA As Key Benchmark for Testing Nucleic Acids Force Fields. *Journal of chemical theory and computation* **2009**, 5, 2514-30.

72. Špačková, N. a.; Berger, I.; Šponer, J., Nanosecond Molecular Dynamics Simulations of Parallel and Antiparallel Guanine Quadruplex DNA Molecules. *Journal of the American Chemical Society* **1999**, 121, 5519-5534.

73. Akhshi, P.; Mosey, N. J.; Wu, G., Free-Energy Landscapes of Ion Movement through a G-Quadruplex DNA Channel. *Angewandte Chemie International Edition* **2012**, 51, 2850-2854.

74. Stadlbauer, P.; Krepl, M.; Cheatham, T. E., III; Koča, J.; Šponer, J., Structural dynamics of possible late-stage intermediates in folding of quadruplex DNA studied by molecular simulations. *Nucleic acids research* **2013**, 41, 7128-7143.

75. Stefl, R.; Cheatham, T. E., 3rd; Spacková, N. a.; Fadrná, E.; Berger, I.; Koca, J.; Sponer, J., Formation pathways of a guanine-quadruplex DNA revealed by molecular dynamics and thermodynamic analysis of the substates. *Biophysical journal* **2003**, 85, 1787-1804.

76. Havrila, M.; Stadlbauer, P.; Kührová, P.; Banáš, P.; Mergny, J.-L.; Otyepka, M.; Šponer, J., Structural dynamics of propeller loop: towards folding of RNA G-quadruplex. *Nucleic acids research* **2018**, 46, 8754-8771.

77. Islam, B.; Stadlbauer, P.; Neidle, S.; Haider, S.; Sponer, J., Can We Execute Reliable MM-PBSA Free Energy Computations of Relative Stabilities of Different Guanine Quadruplex Folds? *The Journal of Physical Chemistry B* **2016**, 120, 2899-2912.

78. Sponer, J.; Mladek, A.; Spackova, N.; Cang, X.; Cheatham, T. E., 3rd; Grimme, S., Relative stability of different DNA guanine quadruplex stem topologies derived using large-scale quantum-chemical computations. *Journal of the American Chemical Society* **2013**, 135, 9785-96.

79. Largy, E.; Mergny, J. L.; Gabelica, V., Role of Alkali Metal Ions in G-Quadruplex Nucleic Acid Structure and Stability. *Metal ions in life sciences* **2016**, 16, 203-58.

80.   Heddi, B.; Martín-Pintado, N.; Serimbetov, Z.; Kari, Teuku Mahfuzh A.; Phan, A. T., G-quadruplexes with (4n - 1) guanines in the G-tetrad core: formation of a G-triad·water complex and implication for small-molecule binding. *Nucleic acids research* **2015**, 44, 910-916.

81.   Kwok, C. K.; Marsico, G.; Sahakyan, A. B.; Chambers, V. S.; Balasubramanian, S., rG4-seq reveals widespread formation of G-quadruplex structures in the human transcriptome. *Nature Methods* **2016**, 13, 841.

82.   Fay, M. M.; Lyons, S. M.; Ivanov, P., RNA G-Quadruplexes in Biology: Principles and Molecular Mechanisms. *Journal of molecular biology* **2017**, 429, 2127-2147.

83.   Malgowska, M.; Czajczynska, K.; Gudanis, D.; Tworak, A.; Gdaniec, Z., Overview of the RNA G-quadruplex structures. *Acta biochimica Polonica* **2016**, 63, 609-621.

84.   Ji, X.; Sun, H.; Zhou, H.; Xiang, J.; Tang, Y.; Zhao, C., Research Progress of RNA Quadruplex. *Nucleic Acid Therapeutics* **2011**, 21, 185-200.

85.   Zhang, D. H.; Fujimoto, T.; Saxena, S.; Yu, H. Q.; Miyoshi, D.; Sugimoto, N., Monomorphic RNA G-quadruplex and polymorphic DNA G-quadruplex structures responding to cellular environmental factors. *Biochemistry* **2010**, 49, 4554-63.

86.   Pandey, S.; Agarwala, P.; Maiti, S., Effect of Loops and G-Quartets on the Stability of RNA G-Quadruplexes. *The Journal of Physical Chemistry B* **2013**, 117, 6896-6905.

87.   Biffi, G.; Di Antonio, M.; Tannahill, D.; Balasubramanian, S., Visualization and selective chemical targeting of RNA G-quadruplex structures in the cytoplasm of human cells. *Nature chemistry* **2014**, 6, 75-80.

88.   Malgowska, M.; Gudanis, D.; Kierzek, R.; Wyszko, E.; Gabelica, V.; Gdaniec, Z., Distinctive structural motifs of RNA G-quadruplexes composed of AGG, CGG and UGG trinucleotide repeats. *Nucleic acids research* **2014**, 42, 10196-10207.

89.   Kumari, S.; Bugaut, A.; Huppert, J. L.; Balasubramanian, S., An RNA G-quadruplex in the 5' UTR of the NRAS proto-oncogene modulates translation. *Nature chemical biology* **2007**, 3, 218-221.

90.   Murat, P.; Marsico, G.; Herdy, B.; Ghanbarian, A. T.; Portella, G.; Balasubramanian, S., RNA G-quadruplexes at upstream open reading frames cause DHX36- and DHX9-dependent translation of human mRNAs. *Genome biology* **2018**, 19, 229.

91.   Wolfe, A. L.; Singh, K.; Zhong, Y.; Drewe, P.; Rajasekhar, V. K.; Sanghvi, V. R.; Mavrakis, K. J.; Jiang, M.; Roderick, J. E.; Van der Meulen, J.; Schatz, J. H.; Rodrigo, C. M.; Zhao, C.; Rondou, P.; de Stanchina, E.; Teruya-Feldstein, J.; Kelliher, M. A.; Speleman, F.; Porco, J. A., Jr.; Pelletier, J.; Rätsch, G.; Wendel, H.-G., RNA G-quadruplexes cause eIF4A-dependent oncogene translation in cancer. *Nature* **2014**, 513, 65-70.

92.   Patel, D. J.; Phan, A. T.; Kuryavyi, V., Human telomere, oncogenic promoter and 5'-UTR G-quadruplexes: diverse higher order DNA and RNA targets for cancer therapeutics. *Nucleic acids research* **2007**, 35, 7429-7455.

93.   Beaudoin, J.-D.; Perreault, J.-P., 5'-UTR G-quadruplex structures acting as translational repressors. *Nucleic acids research* **2010**, 38, 7022-7036.

94.   Chan, K. L.; Peng, B.; Umar, M. I.; Chan, C.-Y.; Sahakyan, A. B.; Le, M. T. N.; Kwok, C. K., Structural analysis reveals the formation and role of RNA G-quadruplex structures in human mature microRNAs. *Chemical Communications* **2018**, 54, 10878-10881.

95.   Kotar, A.; Rigo, R.; Sissi, C.; Plavec, J., Two-quartet kit* G-quadruplex is formed via double-stranded pre-folded structure. *Nucleic acids research* **2018**, 47, 2641-2653.

96.   Kankia, B., Stability Factors of the Parallel Quadruplexes: DNA Versus RNA. *The journal of physical chemistry. B* **2019**, 123, 1060-1067.

97.   Salsbury, A. M.; Lemkul, J. A., Molecular Dynamics Simulations of the c-kit1 Promoter G-Quadruplex: Importance of Electronic Polarization on Stability and Cooperative Ion Binding. *The Journal of Physical Chemistry B* **2019**, 123, 148-159.

98.   Kuhrova, P.; Mlynsky, V.; Zgarbova, M.; Krepl, M.; Bussi, G.; Best, R. B.; Otyepka, M.; Sponer, J.; Banas, P., Improving the Performance of the Amber RNA Force Field by Tuning the Hydrogen-Bonding Interactions. *Journal of chemical theory and computation* **2019**, 15, 3288-3305.

99.   Bergonzo, C.; Cheatham, T. E., Improved Force Field Parameters Lead to a Better Description of RNA Structure. *Journal of chemical theory and computation* **2015**, 11, 3969-3972.

100.   Bergonzo, C.; Henriksen, N. M.; Roe, D. R.; Cheatham, T. E., 3rd, Highly sampled tetranucleotide and tetraloop motifs enable evaluation of common RNA force fields. *Rna* **2015**, 21, 1578-90.

101.   Hase, F.; Zacharias, M., Free energy analysis and mechanism of base pair stacking in nicked DNA. *Nucleic acids research* **2016**, 44, 7100-8.

102.   Sponer, J.; Spackova, N., Molecular dynamics simulations and their application to four-stranded DNA. *Methods* **2007**, 43, 278-90.

103.   Gkionis, K.; Kruse, H.; Platts, J. A.; Mládek, A.; Koča, J.; Šponer, J., Ion Binding to Quadruplex DNA Stems. Comparison of MM and QM Descriptions Reveals Sizable Polarization Effects Not Included in Contemporary Simulations. *Journal of chemical theory and computation* **2014**, 10, 1326-1340.

104.   Gros, J.; Rosu, F.; Amrane, S.; De Cian, A.; Gabelica, V.; Lacroix, L.; Mergny, J.-L., Guanines are a quartet's best friend: impact of base substitutions on the kinetics and stability of tetramolecular quadruplexes. *Nucleic acids research* **2007**, 35, 3064-3075.

105.   Zhu, H.; Xiao, S.; Wang, L.; Liang, H., Communication: Asymmetrical cation movements through G-quadruplex DNA. *The Journal of Chemical Physics* **2014**, 141, 041103.

For Table of Contents Use Only

Synopsis: We have carried out extended unbiased atomistic simulations using different ion and water parameters to study the stability of two and four quartet G-quadruplexes (GQs). The all-*anti* parallel-stranded two-quartet DNA GQs are highly unstable compared to their RNA counterpart in all the ion and water conditions. GQs composed of two stacked units of two-quartet GQs were, however, stable for both DNA and RNA.