**Cross-kingdom regulation of tRNAs/tRFs derived from Chinese yew**

Kai-Yue Cao[1,2†], Tong-Meng Yan[1,2†], Ji-Zhou Zhang[3], Ting-Fung Chan[3], Jie Li[4],

Chong Li[4], Elaine Lai-Han Leung[1,2], Zhi-Hong Jiang[1,2]*

[1]State Key Laboratory of Quality Research in Chinese Medicine, Macau University of

Science and Technology, Macau SAR, China.

[2]Macau Institute for Applied Research in Medicine and Health, Macau University of

Science and Technology, Macau SAR, China.

[3]School of Life Sciences, and State Key Laboratory of Agrobiotechnology, The

Chinese University of Hong Kong, Shatin, Hong Kong SAR, China.

[4]College of Pharmaceutical Sciences, Southwest University, Chongqing 400715, P. R.

China.

**\*Corresponding author**: Dr. Zhi-Hong Jiang, State Key Laboratory of Quality

Research in Chinese Medicine, Macau University of Science and Technology,

Avenida Wai Long, Taipa, Macau SAR, China.

**E-mail**: zhjiang@must.edu.mo (Zhi-Hong Jiang)

†These authors contributed equally to this work.

Plants containing countless chemical constituent have benefited mankind since

the origin of life. Although secondary metabolites in plants, such as morphine,

artemisinin and taxol, have been developed as therapeutic drugs for clinical

therapy, few study focuses on the pharmacological activities of plant small RNAs

with function of cross-kingdom regulations. Yew is widely considered as a

"superstar" plant due to the discovery of paclitaxel, or taxol, which is a

well-known natural drug for the treatment of multiple types of cancer[1]. Here we

show the surprising finding that an RNA fragment, named tRF-T11, derived

from $tRNA^{His(GUG)}$ of Chinese yew strongly suppressed human ovarian cancer

progression. In A2780 cells, tRF-T11 mimic (a double-stranded RNA with

tRF-T11 as antisense chain) exhibited potent cytotoxicity comparable to that of

taxol, but no significant cytotoxicity to normal ovarian surface epithelial cells.

Moreover, the cytotoxicity of tRF-T11 mimic is 80-fold stronger than that of

taxol in taxol-resistant A2780 cells. Bioinformatic and molecular biological

studies revealed that tRF-T11 targets transient receptor potential cation channel

subfamily A member 1 (*TRPA1*) to inhibit its expression levels. In a further *in*

**vivo investigation, the growth rate of ovarian tumor xenografts in nude mice was**

**significantly reduced by treatment with tRF-T11 mimic, and the *TRPA1* protein**

**expression in tumors treated with tRF-T11 mimic was also down-regulated. Our**

**findings are the first to provide evidence that plant-derived tRFs can regulate**

**the expression of target genes *in vitro* and *in vivo*, indicating that they may**

**become as a new source of druggable siRNA. Moreover, this discovery**

**demonstrated a pilot example of an innovative approach for not only identifying**

**pharmacologically-active tRFs from plants, but also for improving the efficiency**

**and possibilities of discovering new drug target.**

Plants have provided mankind with an important source of medicinal products

for millennia. Throughout ancient history, humans learned to harness herbal

decoctions for disease prevention and treatment. Today, thousands of secondary

metabolites, such as flavonoids, terpenes and alkaloids, from this chemical treasure

have been identified as active compounds. Intriguingly, crude plant extracts were

more effective than the synthetic versions of phytochemicals, implying that whole

plant extracts contain a cocktail of products in which substances like macromolecules

other than small molecules may contribute to the overall pharmacological effects[2].

However, few study has been performed on the pharmacological activity of plant

macromolecules excluding polysaccharides because they are considered inactive,

nonabsorbable and unstable in human body, such as RNA, DNA and proteins. Small

RNAs (<200 nt) were found to play critical roles in a wide range of biological

processes in microorganisms, plants, animals and human, and to participate in a

variety of human diseases by regulating protein-coding genes[3,4]. Since the first

evidence that plant-derived miRNAs can regulate gene expression in mammalian

systems was provided by Zhang *et al*[5] in 2012, many other miRNAs from small RNAs

in plants have been extensively studied for their cross-kingdom regulatories[6-9]. These

findings suggested the promise of small RNAs from plants to be developed as

therapeutic agents. A recent report by Shekhawat[10] showed that the small RNA of

corn exhibited appreciable antiproliferative activity toward HeLa cells, implying the

possibility that small RNA species with larger molecular weight than miRNA play

regulatory roles. In fact, as the most abundant small RNA species expressed in all

living organisms, transfer RNAs (tRNAs) play a well-defined role in protein

translation and perform diverse regulatory functions in some important biological

processes, such as cell signaling, apoptosis modulation and stress response programs[11].

Recently, tRNA-derived fragments (tRFs) have also been recognized to play

regulatory roles[12]. This class of small RNAs is non-randomly generated from tRNAs,

and many are specifically and abundantly produced under particular conditions, such

as cancer and viral infection[13,14]. Some studies have also revealed that the expression

of several tRFs is positively correlated with cell proliferation in different cancer cell

lines[15], suggesting that tRFs and their related signaling pathways could be novel

targets for intervention in the treatment of cancers. These findings prompted us to

consider the further possibility that in addition to well-established players such as

miRNAs and small molecules in plants, tRNAs and their fragments, which account

for overwhelming majority of small RNA species in plants, may hold great potential

as biological regulators in mammals. Below, we describe the first study of the

pharmacological activities of tRNAs and tRFs derived from Chinese yew (*Taxus*

*chinensis*) and identification of their molecular target.

Total RNA with high integrity (RIN=7.7) and purity (A260/280=2.08,

A260/230=2.00; Supplementary Fig 1a) was isolated from fresh branches of *T.*

*chinensis* by using a well-developed CTAB method (Supplementary Fig 1b, c), in

which small RNA was separated from large RNA with high purity (<200 nt,

Supplementary Fig 2). The tRNAs were then gel fractionated and purified

(Supplementary Fig 3a) to successfully delete 5S and 5.8S rRNA and other RNA

species (Supplementary Fig 3b). Sequence information and the proportion of each

type of RNA in this tRNA-enriched fraction (tEF) were determined by performing

next-generation sequencing (NGS). As shown in Fig 1c and Supplementary Table 1,

over 1.7 million clean reads distributed mainly from 60 to 80 mer were generated, and

over 20 individual tRNA sequences were identified, which accounted for

approximately 14% of the overall number of reads in the tEF library (Fig 1d). The

cytotoxicity of the tEF was investigated using an MTT assay on the A2780 human

ovarian cancer (OVCA) cell line. The $IC_{50}$ value for dose-dependent growth

inhibition of A2780 cells *in vitro* was 128.8 nM when the tEF was well encapsulated

and delivered via liposomes, while no significant cytotoxicity of the tEF was observed

without transfection (Fig 1a, b), implying that tRNAs or their fragments in the tEF

might act as effectors. Furthermore, when rRNA degradation fragments (Fig 1e) were

removed from the tEF samples via a Ribo-Zero rRNA removal kit (Illumina), the

retained tRNA species induced the death of A2780 cells with increased efficiency

(Fig 1f), indicating the major contribution of tRNA species to the overall cytotoxic

effects of the tEF. Subsequently, the five most abundant (>500 reads per million)

individual tRNAs were purified from small RNA of *T. chinensis* by using a magnetic

solid-phase DNA probe method[16] and were identified by ultra-high performance

liquid chromatography coupled with quadrupole time-of-flight mass spectrometry

(Fig 1g-i). Cytotoxicity of purified tRNAs was then investigated on A2780 cells,

HepG2 of liver cancer and MCF-7 of breast cancer cells. As a result, A2780 and

HepG2 cells exhibited a proliferation reduction of over 75% when treated with

individual tRNAs at 25 nM (Fig 1j, k), while MCF-7 cells were much less sensitive

(reduction of less than 30%) to these tRNA molecules. This pattern corresponded to

an approximate $IC_{50}$ value of 14.3 nM for the growth inhibition of A2780 cells caused

by treatment with $tRNA^{Trp(CCA)}$ for 48 h, which is close to that of taxol (7.4 nM) used

as a positive control drug (Fig 1l). This provides the first evidence of pharmacological

activities of plant-derived tRNAs, suggesting that they are a new class of therapeutic

agents from medicinal plants.

Meanwhile, since the previous study has revealed that the tRFs could regulate

endogenous gene in breast cancer cells[15], we hypothesized that tRF from the above

cytotoxic tRNAs of Chinese yew may be pharmacologically-active. Indeed, tRFs

from the 5' terminal and 3' ends (5'-tRF and 3'-tRF, respectively) are the most

abundant tRFs associated with the AGO protein, which is a key component of the

RNA interference pathway[17]. A similar report showed that a 5'-tRF of 19 nucleotides

is the most abundant tRF in land plants[18]. Thus, 36 tRFs derived from the top 9

abundant tRNAs of Chinese yew, including 5'-tRF and 3'-tRF with lengths of 22 and

19 nt, respectively, were selected as antisense strands of tRF mimics in siRNA form

(Fig 2a, Supplementary Table 2), and the antiproliferative activities of these tRF

mimics against A2780 and its taxol-resistant strain (A2780T) cells were screened.

The results showed that two of these 36 tRF mimics significantly inhibited the

proliferation of the A2780 cells by over 50% at 50 nM (Fig 2b). Among these tRF

mimics, tRF-T11 mimic (a double strand RNA with a 22 nt 5'-tRF derived from

$tRNA^{His(GUG)}$ as antisense RNA) showed the highest antiproliferative activity against

both cell lines. As shown in Fig 2c, the $IC_{50}$ value of tRF-T11 mimic for inhibiting the

growth of A2780 cells was 31 nM, which is comparable to that of taxol (10.2 nM).

Surprisingly, high sensitivity to tRF-T11 mimic was also observed in A2780T cells,

corresponding to an $IC_{50}$ value of 32 nM, which is 80-fold lower than that of taxol

(2600 nM). In addition, no significant cytotoxicity of tRF-T11 mimic to IOSE80

normal ovarian surface epithelial cells was observed. Furthermore, the cytotoxicity of

tRF-T11 mimic was confirmed by visualizing and tracking fluorescent dye

(FAM)-labeled tRF-T11 mimic in A2780 cells. The images shown in Fig 2d indicated

that tRF-T11 mimic was efficiently transfected into the cytoplasm of A2780 cells,

where they exhibited functions involved in the biological processes, such as

suppressing the expression of endogenous targets[19]. These results indicated that

*Taxus*-derived tRFs are promising therapeutic agents for drug-resistant OVCA.

The divergent activity of tRF mimics derived from individual tRNAs observed in

this study suggested that plant-derived tRFs can exhibit bioactivities in a

sequence-specific manner in mammalian cells. It has been reported that functionality

of exogenous tRFs possibly overlaps that of miRNAs, a characteristic that has been

widely documented for endogenous tRFs in both plants and mammalian cells[20,21]. To

further identify potential target genes of tRF-T11, we performed an *in silico*

prediction using two independent bioinformatic tools, IntaRNAv2[22] and

psRNATarget[23], followed by differential expression analysis based on the gene

expression profiles in normal and tumor ovarian tissue to prioritize the candidate

targets. As a result, the top three differentially expressed genes, namely transient

receptor potential cation channel subfamily A member 1 (*TRPA1*), tetraspanin 17

(*TSPAN17*) and secretogranin V (*SCG5*), with the minimum required energy to

hybridize with tRF-T11 at their 3' UTR regions of mRNA, were considered as the

potential targets for further experimental validation (Fig 2e). Among these three genes,

*TRPA1*, a new target recently unveiled by Takahashi *et al*[24] for cancer therapies,

which acts by triggering a noncanonical oxidative stress defense mechanism of tumor

suppression, was dramatically downregulated in A2780 cells (by approximately 75%)

in a dose-dependent manner by tRF-T11 mimic treatment (Fig 2j, g), while the

mRNA levels of the other two targets did not vary significantly (Fig 2f). Furthermore,

a marked decrease of over 50% in the protein expression level of *TRPA1* was also

observed in A2780 cells treated with tRF-T11 mimic (Fig 2h, i), which may act as a

translational repressor of the *TRPA1* gene by acting like a miRNA in human OVCA

cells. Indeed, previous reports revealed that *TRPA1* overexpression is associated with

poor survival probability in patients (Fig 2k). Unfortunately, no clinical therapeutic

drug targeting *TRPA1* is available, and our findings provide a novel RNA sequence

derived from plant can potently silence human *TRPA1*.

Recently, advances in RNA therapeutics have been facilitated by breakthroughs

in pharmaceutical delivery systems, including liposomes, nanoparticles, dendrimers

and carbon nanotubes have been shown to provide promising application *in vitro* and

*in vivo*[25]. To evaluate the antitumor activity *in vivo*, a histidine-lysine polymer (HKP)

nanoparticle system was applied to deliver tRF-T11 mimic to a well-established

xenograft mouse model of human OVCA cells. The mass ratio of HKP and tRF-T11

mimic was optimized (Fig 3a, b) with an appropriate particle size (164.2 nm) and

good dispersion (41.2 mV, Fig 3c). As shown in Fig 3d to g, compared to vehicle

control group, tRF-T11 mimic treatment of xenografts for 20 days significantly

diminished the tumor growth rate by approximately 50% at a dose of only 177

nmol/kg, which is more than 16-fold lower than that of taxol (2928 nmol/kg) required

to achieve comparable tumor-suppressive effects without significant adverse impacts

on body weight. Compared to those from control mice, the protein expression levels

of *TRPA1* in the tumors with intratumoral injection of high-dose tRF-T11 mimic

exhibited significant reduction (Fig 3h, i), which is consistent with the western blot

results *in vitro*. Although taxol is a commonly used first-line drug for cancer patients,

chemoresistance is usually observed. Since tRF-T11 mimic exhibited comparable

tumor suppression efficacy to taxol at a much lower dose, our study may indicate an

alternative treatment option.

In summary, this study provides the first evidence that plant-derived tRNAs and

tRFs from Chinese yew exhibit potent antitumor effects against human OVCA, which

will greatly expand our knowledge on the active ingredients in natural sources, and

provide additional clues for evaluating the active components of medicinal plants or

herbal medicines. Although cross-kingdom regulation in human by miRNA uptake

from food and medicinal plants has been widely studied, this report emphasizes the

potential regulatory roles of plant-derived small RNAs from an abundance of

housekeeping noncoding RNA species, such as tRNAs and tRFs. This infinitely vast

pool filled with RNA sequences and chemical modifications provided by nature can

serve as a new sequence library for the development of novel RNAi-based

therapeutics. Furthermore, since chemical modified oligonucleotides could be simply

and efficiently synthesized in large amount using the phosphoramidite

methodology[26,27], this discovery demonstrated a pilot strategy for identifying

pharmacologically active tRFs from plants, such as those from the genus *Taxus*.

As a promising and rapidly advancing frontier, siRNA can be readily designed

based on a well-validated target mRNA. Often, the process of unraveling the

pathogenesis of many human diseases to find key molecular targets for drug treatment

is time-consuming and difficult without a functional approach with bottom-up

guidance. As a result, the effective design of RNAi therapeutic drug candidates will

be difficult or impossible due to the lack of reliable targets. As a benefit of the high

target specificity and selectivity of RNA drugs, pharmacologically active RNA

molecules identified from medicinal plants could, in turn, lead to the improvement of

the efficiency and possibilities of discovering new drug target, and shed light on

disease etiology.

**Acknowledgements**

We thank Novogene Bioinformatics Institute (Beijing, China) for technical

support in next-generation sequencing; Y. L. (Sirnaomics, Inc., U.S.A.) for technical

advice and the gift of HKP for RNA delivery *in vivo*. This work was financially

funded by the CUHK Direct Grant 4053364, the Innovation and Technology

Commission, Hong Kong Government to the State Key Laboratory, and grant from

The Science and Technology Development Fund, Macau SAR (File no.

015/2017/AFJ).

**Contributions**

Z.-H.J. and T.-M.Y. conceived the study. K.-Y.C., T.-M.Y. and T.-F.C. designed

the experiments. K.-Y.C., T.-M.Y., J.-Z.Z. and J. L. conducted the experiments.

K.-Y.C., J.-Z.Z. and C. L. analyzed the data. K.-Y.C., T.-M.Y. and E.L.L. drafted the

manuscript. All authors approved the manuscript.

**Methods**

**Plant materials and RNA extraction of *Taxus chinensis*.** Plants of *Taxus chinensis*

*(Pilger) Rehd. var. mairei* (Lemee et Levl.) Cheng et L. K. Fu used in this study were

collected from Sanming City, Fujian Province, China. Plant materials were freshly cut,

frozen and stored in liquid nitrogen until use.

Plant tissues were ground into a fine powder in liquid nitrogen and were then

homogenized (IKA, Germany) in CTAB buffer. After incubation for 2 min at 65°C,

the tissue lysate was cooled immediately on ice for 10 min, followed by

centrifugation at 12,000 ×g for 15 min at 4°C. The supernatant was collected and

extracted with an equal volume of phenol:chloroform:isopentanol (50:48:1). Phases

were separated at 4°C by centrifugation at 12,000 ×g for 15 min. The supernatant was

extracted again as described above with chloroform:isopentanol (24:1) and mixed

with an equal volume of 6 M guanidinium thiocyanate solution (Tokyo Chemical

Industry, Japan) and ethanol (Anaqua Chemicals Supply, U.S.A.) to a final

concentration of 55%. The mixture was passed through a silicon-membrane column

and washed with 80% ethanol. Finally, total RNA was eluted with RNase-free water,

and small RNA species were enriched by using a mirVana miRNA isolation kit

(Invitrogen, U.S.A.) following the manufacturer's instructions.

**tEF preparation.** Small RNA was separated by 6% urea-polyacrylamide gel

electrophoresis (PAGE). Gel fractions containing the tEF, as visualized by SYBR

Gold nucleic acid gel staining (Thermo, U.S.A.), were excised and electroeluted in 3

kD molecular weight cut off dialysis tubing (Spectrum, U.S.A.) at 100 V for 1.5 h.

The tEF was finally desalted and recovered on a silicon-membrane column.

**tEF library construction and sequencing.** Sequencing libraries were generated by

using a TruSeq small RNA Library Preparation Kit (Illumina, U.S.A.), followed by a

round of adaptor ligation, reverse transcription and PCR. PCR products were then

purified, and libraries were quantified on an Agilent Bioanalyzer 2100 system

(Agilent Technologies, U.S.A.). The library preparations were sequenced at the

Novogene Bioinformatics Institute (Beijing, China) on an Illumina HiSeq platform

using the 150-bp paired-end (PE150) strategy to generate over 15 million raw paired

reads. A total of 1,729,438 clean reads were obtained by removing low quality

regions and adaptor sequences. The tRNA genes were identified by using the

tRNAscan-SE 2.0 program (http://lowelab.ucsc.edu/tRNAscan-SE/) and annotated by

searching the Nucleotide Collection (nr/nt) database using the Basic Local Alignment

Search Tool (BLAST) program (https://blast.ncbi.nlm.nih.gov/Blast.cgi).

**Purification of individual tRNAs.** Synthetic single-stranded DNA oligonucleotides

(BGI, China; Supplementary Table 3) complementary to the target tRNA were mixed

with *Taxus* small RNA and denatured at 95°C for 5 min, followed by incubation at a

temperature 5°C lower than the melting temperature (Tm) for 1.5 h. Streptavidin

magnetic beads (Beaverbio, China) were then added to the mixture and incubated for

30 min at the annealing temperatures. Then, the biotinylated DNA/tRNA-coated

beads were separated with a magnet and washed at 40°C. Subsequently, the magnetic

beads were resuspended in RNase-free water, and the immobilized tRNA with probes

was released by incubation at 65°C for 5 min. The solutions were directly injected

using an ACQUITY UPLC OST $C_{18}$ column (2.1×100 mm i.d., 1.7 μm; Waters,

U.S.A.) maintained at 60°C into a UHPLC Agilent 1290 system (Agilent, U.S.A.)

equipped with a diode array detector. The flow rate was set at 0.2 mL/min, and a

gradient elution was performed with (A) 100 mM HFIP+15 mM TEAA and (B) 50%

MeOH in A as follows: 0-1.5 min, 2% B; 1.5-8.3 min, 2%-28% B; and 8.3-16.5 min,

28%-34% B. MS data were acquired on an Agilent 6545 Q-TOF mass spectrometer

with a gas temperature of 320°C, spray voltage of 3.5 kV, and sheath gas flow and

temperature of 12 L/min and 350°C, respectively. For the MS experiment, samples

were analyzed in negative polarity over an m/z range of 500 to 3200. Fractions from

chromatographic peaks of individual tRNAs were collected and freeze dried.

**RNA transfection and cytotoxicity determination.** RNA samples dissolved in

RNase-free water were stored at -20°C before use. Transfection of RNA samples was

performed according to the manufacturer's instructions for Lipofectamine RNAiMAX

Transfection Reagents (Thermo, U.S.A.). After treatment for 48 h, an MTT assay was

performed, and absorbance values were calorimetrically determined at 570 nm using a

SpectraMax 190 microplate reader (Molecular Devices, Sunnyvale, U.S.A). The

dose-dependent $IC_{50}$ values were calculated by GraphPad Prism 5.0 (GraphPad,

U.S.A.). A siRNA (Forward: 5'-GGAGCGAUUUAGCCAAGAATT-3', Sense:

5'-UUCUUGGC UAAAUCGCUCCTT-3') was used as positive control. Each

experiment was carried out three times. The results are expressed as the means $\pm$

standard deviations.

**Acquisition of fluorescence images of transfected FAM-labeled tRF-T11 mimic**

**in A2780 cells.** Briefly, $1\times10^5$ A2780 cells were seeded in glass bottom cell culture

dishes (Nest Biotechnology, China). The medium was discarded after 20 h, and 50

nM FAM-labeled tRF-T11 mimic (GenePharma, China) was transfected into the cells.

After 6 h, the transfection medium was discarded, and the cells were rinsed 3 times

with RPMI 1640 medium. For nuclear staining, cells were incubated with 1 mL of

Hoechst 33258 (Sigma, U.S.A.) for 10 min at 37°C, followed by washing with RPMI

1640 medium. For mitochondrial staining, cells were incubated with 1 mL of

MitoBeacon Red (GeneCopoeia, U.S.A.) for 30 min at 37°C, followed by washing

with RPMI 1640 medium. Fluorescence images were acquired with an API

DeltaVision Live-cell Imaging System (Applied Precision Inc., GE Healthcare Co.,

U.S.A.) and were captured at x60 magnification, scale bar, 25 µm.

**tRF-T11 target prediction and prioritization.** Previously described small RNA or

tRF target prediction methods and restriction criteria were applied in this study to

identify functional target genes of tRF-T11[28]. Specifically, the 3' UTR sequences of

all mRNAs retrieved from GENCODE (v23) and the sequence of tRF-T11 derived

from *Taxus* were firstly loaded into IntaRNAv2 and psRNATarget to predict the

potential targets. The top 10 overlapping targets predicted were then prioritized and

filtered by crosschecking their expression in tumor and normal ovarian tissues.

Accordingly, we first obtained the gene expression profiles for 379 OVCA tissues and

88 normal ovarian tissues from UCSC Xena, which contains curated comprehensive

batch-effect-adjusted pancancer genomic data[29]. Subsequently, EBSeq[30] was

employed to filter the target genes differentially expressed between tumor and normal

ovarian tissues. The predicted targets satisfying two criteria, fold change

(tumor/normal) $> 2$ and FDR $< 0.1$.

**Quantitative real-time PCR**. Total RNA was extracted from A2780 cells treatment

with the tRF-T11 mimic for 48 h using TRIzol (Invitrogen, U.S.A.) on a

silicon-membrane column. RNA was then reverse transcribed using the GoScript

Reverse Transcription System (Promega, U.S.A.). Quantitative real-time PCR was

performed on a high-productivity real-time quantitative PCR ViiA$^{TM}$ 7 system (Life

Technologies, U.S.A.) using GoTaq$^{®}$ qPCR Master Mix (Promega, U.S.A.) according

to the manufacturers' instructions. The PCR primers were synthesized commercially

(BGI, China, Supplementary Table 4). All data are presented as the mean fold change

values duplicated and normalized to $\beta$-actin expression using the $2^{-\Delta\Delta CT}$ method. Each

experiment was carried out three times. The results are expressed as the means ±

standard deviations.

**Western blot analysis**. A2780 cell lysates were collected after treatment with

tRF-T11 mimic for 48 h, and protein extraction was carried out using RIPA buffer

(Cell Signaling Technology, U.S.A.) supplemented with protease and phosphatase

inhibitor cocktails (Roche, Switzerland). Protein concentrations were then determined

by a BCA protein assay (Thermo, U.S.A.). Denatured proteins were separated by 8%

SDS-PAGE and transferred to nitrocellulose membranes, followed by blocking with 5%

bovine serum albumin (Thermo, U.S.A.) for 2 h. Membranes containing proteins were

incubated with primary antibody against *TRPA1* (1:500 dilution, Novus, U.S.A.) or

*β*-actin (1:1000 dilution, Santa Cruz, U.S.A.) overnight, followed by washing in

TBST and incubation with a fluorescent secondary antibody (Li-cor, U.S.A.) for

detection. Band intensities were measured using ImageJ and were normalized to those

of *β*-actin. The sequences of TRPA1_siRNA01, TRPA1_siRNA02 and the scrambled

control (GenePharma) are shown in Supplementary Table 5. Each experiment was

carried out three times. The results are expressed as the means ± standard deviations.

**Preparation of HKP-tRF-T11 mimic nanoparticles**. HKP (a gift from Dr. Yang Lu,

Sirnaomics, Inc., U.S.A.) dissolved in RNase-free water (240 µg/mL) were first

balanced overnight at 4℃. Equal volumes of tRF-T11 mimic solutions at a

concentration of 80 µg/mL were then mixed and incubated at room temperature for 30

min. The particle sizes and zeta potential of the resulting nanoparticles were then

evaluated by a Malvern Zetasizer Nano ZS90 (Malvern Instruments, U.K.).

***In vivo* models for tumor therapy**. A2780 cells $(4.0 \times 10^6)$ were injected

subcutaneously into the armpits of 8-week-old BALB/c nude mice (Shanghai SLAC

Laboratory Animal Co., Ltd., China). When the tumors reached 50 $mm^3$, the

tumor-bearing nude mice were treated with different doses (71 nmol/kg, 177 nmol/kg)

of tRF-T11 mimic (GeneDesign, Inc. Osaka, Japan) encapsulated with HKP and

different doses of taxol (2928 nmol/kg or 5855 nmol/kg) via intratumoral injection

once every three days. A group treated with HKP alone was established as a control.

The animals were sacrificed on day 20. Tumor diameters were measured at the points

of maximum length and maximum width with digital calipers. Tumor volumes were

calculated by the following formula: volume=$(width)^2 \times length/2$. The data were

statistically analyzed using GraphPad Prism 5.0. Animal care and use protocols were

performed in compliance with the guidelines of the Ethics Committee of the College

of Pharmaceutical Sciences, Southwest University.

**References**

1. Wani, M.C., Taylor, H.L., Wall, M.E., Coggon, P. & McPhail, A.T. Plant

antitumor agents. VI. The isolation and structure of taxol, a novel

antileukemic and antitumor agent from *Taxus brevifolia*. *J. Am. Chem. Soc.* 93,

2325-2327 (1971).

2. Yarnell E. Synergy in herbal medicines: part 1. *J. Restor. Med.* 4, 60–73

(2015).

3. Wu, L., Fan, J., & Belasco, J.G. MicroRNAs direct rapid deadenylation of

mRNA. *Proc. Natl. Acad. Sci.* 103, 4034–4039 (2006).

4. Standart, N. & Jackson, R.J. MicroRNAs repress translation of

$m^7$Gppp-capped target mRNAs in vitro by inhibiting initiation and promoting

deadenylation. *Genes Dev.* 21, 1975–1982 (2007).

5. Zhang, L. *et al.* Exogenous plant MIR168a specifically targets mammalian

LDLRAP1: evidence of cross-kingdom regulation by microRNA. *Cell Res.* 22,

273-274 (2012).

6.    Chin, A.R. *et al.* Cross-kingdom inhibition of breast cancer growth by plant

miR159. *Cell Res.* 26, 217-228 (2016).

7.    Zhou, Z. *et al.* Honeysuckle-encoded atypical microRNA2911 directly targets

influenza A viruses. *Cell Res.* 25, 39-49 (2015).

8.    Du, J.C. *et al*. Plant-derived phosphocholine facilitates cellular uptake of

anti-pulmonary fibrotic HJT-sRNA-m7. *Sci. China-Life Sci*. 60, 1-11 (2017).

9.    Teng, Y. *et al*. Plant-derived exosomal microRNAs shape the gut microbiota.

*Cell Host Microbe* 24, 1-16 (2018).

10.    Shekhawat M., Jahagirdar D., Yadav S. & Sharma N.K. Induction of apoptosis

in HeLa by corn small RNAs. *Nutr. Cancer* 71, 1-11 (2019).

11.    Barciszewska M.Z., Perrigue P.M. & Barciszewski J. tRNA – the golden

standard in molecular biology. *Mol. Biosyst.* 12, 12-17 (2016).

12.    Anderson P. & Ivanov P. tRNA fragments in human health and disease. *FEBS*

*Lett.* 588, 4297-4304 (2014).

13.    Yeung, M.L. *et al.* Pyrosequencing of small non-coding RNAs in HIV-1

infected cells: evidence for the processing of a viral-cellular double-stranded

RNA hybrid. *Nucleic Acids Res.* 37, 6575-6586 (2009).

14.    Saikia, M. *et al.* Angiogenin-cleaved tRNA halves interact with cytochrome c,

protecting cells from apoptosis during osmotic stress. *Mol. Cell. Biol.* 34,

2450-2463 (2014).

15.    Goodarzi, H. *et al.* Endogenous tRNA-derived fragments suppress breast

cancer progression via YBX1 displacement. *Cell* 161, 790-802 (2015).

16.    Tsurui, H. *et al*. Batchwise purification of specific tRNAs by a solid-phase

DNA probe. *Anal. Biochem.* 221, 166–172 (1994).

17.    Martinez, G., Choudury, S.G. & Slotkin, R.K. tRNA-derived small RNAs

target transposable element transcripts. *Nucleic Acids Res.* 45, 5142-5152

(2017).

18.    Loss-Morais, G., Waterhouse, P.M. & Margis, R. Description of plant

tRNA-derived RNA fragments (tRFs) associated with argonaute and

identification of their putative targets. *Biol. Direct* 8, 1-6 (2013).

19.    Keam, S.P. & Hutvagner, G. tRNA-derived fragments (tRFs): Emerging new

417           roles for an ancient RNA in the regulation of gene expression. *Life* 5,

418           1638-1651 (2015).

419    20.    Maute, R.L. *et al.* tRNA-derived microRNA modulates proliferation and the

420           DNA damage response and is down-regulated in B cell lymphoma. *Proc. Natl.*

421           *Acad. Sci.* 110, 1404-1409 (2013).

422    21.    Deng, J.F. *et al.* Respiratory syncytial virus utilizes a tRNA fragment to

423           suppress antiviral responses through a novel targeting mechanism. *Mol. Ther.*

424           23, 1622-1629 (2015).

425    22.    Mann, M., Wright, P.R. & Backofen, R. IntaRNA 2.0: enhanced and

426           customizable prediction of RNA-RNA interactions. *Nucleic Acids Res.* 45,

427           W435-W439 (2017).

428    23.    Dai, X., Zhuang, Z. & Zhao, P.X. psRNATarget: a plant small RNA target

429           analysis server (2017 release). *Nucleic Acids Res.* 46: W49-W54 (2018).

24.  Takahashi, N. *et al*. Cancer cells co-opt the neuronal redoxsensing channel

TRPA1 to promote oxidative stress tolerance. *Cancer Cell* 33, 985–1003

(2018).

25.  Kesharwani, P., Gajbhiye, V. & Jain, N.K. A review of nanocarriers for the

delivery of small interfering RNA. *Biomaterials* 33, 7138-7150 (2012).

26.  Michelson, A.M. & Todd, A.R. Synthesis of a dithymidine dinucleotide

containing a 3' : 5'-internucleotidic linkage. *J. Chem. Soc.* 2632–2638 (1955).

27.  Beaucage, S.L. & Iyer, R.P. Advances in the synthesis of oligonucleotides by

the phosphoramidite approach. *Tetrahedron*. 48, 2223-2311 (1992).

28.  Babski, J. *et al*. Small regulatory RNAs in Archaea. *RNA Biol.* 11, 484-493

(2014).

29.  Goldman, M. *et al*. The UCSC Xena Platform for cancer genomics data

visualization and interpretation. *bioRxiv* 326470 (2018).

30.  Leng, N. *et al*. EBSeq: an empirical Bayes hierarchical model for inference in

RNA-seq experiments. *Bioinformatics* 29, 1035-1043 (2013).

**Figure Legends**

**Figure 1.** *Taxus*-derived tRNAs exhibit cytotoxicity *in vitro*. **a**, *Taxus* tEF with

liposomal transfection exhibited strong cytotoxicity on A2780 cells. **b**, Representative

MTT image of the *Taxus* tEF with or without liposomal transfection. **c**, Read length

distribution of the tEF determined by NGS was primarily between 60 and 80 mer. **d**,

NGS read content analysis showed that 14% of the tEF was tRNA. **e**, Urea-PAGE

analysis of the tEF, rRNA fragment and purified tEF. **f**, The cytotoxicity of purified

tEF increased after rRNA fragment deletion. **g**, Representative UHPLC method for

purification of tRNA$^{His(GUG)}$. **h**, Deconvolution mass spectrometry and urea-PAGE

analysis of purified tRNA$^{His(GUG)}$. **i**, Deconvolution mass results of tRNA$^{Glu(UUC)}$,

tRNA$^{Trp(CCA)}$. tRNA$^{Leu(CAA)}$ and tRNA$^{Arg(ACG)}$. **j**, MTT assay of the top five abundant

tRNAs from Chinese yew on A2780, HepG2 and MCF-7 cells. **k**, Representative

MTT image of individual tRNAs on A2780 cells. **l**, Dose-dependent investigation of

tRNA$^{Trp(CCA)}$ on A2780 cells. The data are shown as the means $\pm$ SDs of 3

independent experiments. **P<0.01*, ***P<0.001*, ****P<0.0001*.

**Figure 2**. *Taxus*-derived tRF-T11 mimic inhibits human OVCA cell proliferation by

targeting *TRPA1 in vitro*. **a,** Design of tRF mimic from 5' and 3' end of tRNA. **b,**

Heatmap of the inhibition rate of 36 synthesized tRF mimics against A2780 and

A2780T cells. **c,** Investigation of the dose dependency of tRF-T11 mimic and taxol

against A2780, A2780T and IOSE80 cells. **d,** Fluorescent images capture of FAM

labeled tRF-T11 mimic in A2780 cells. **e**, Summary of hybridization energies and

differential expression analysis of predicted target genes identified by the IntaRNA

and psRNATarget tools. **f**, qPCR validation of the mRNA levels of predicted targets

of tRF-T11 in A2780 cells. **g**, tRF-T11 mimic decreased the mRNA levels of *TRPA1*

in a dose-dependent manner. **h**, Western blot assay for the evaluation of *TRPA1*

expression in A2780 cells treated with tRF-T11 mimic, siRNA against *TRPA1*

(TRPA1_siRNA01, TRPA1_siRNA02) and scrambled control at 25 nM. Band

intensities were measured using ImageJ and were normalized to those of *β*-actin (**i**). **j,**

Putative tRF-T11 binding sites in the *TRPA1* 3' UTR. **k**, Overall survival probability

of patients in the TCGA OVCA cohort. The data are shown as the means ± SDs of 3

independent experiments. **P<0.01, ***P<0.001, ****P<0.0001.

**Figure 3.** *Taxus*-derived tRF-T11 mimic decreases OVCA tumor growth via targeting

*TRPA1 in vivo*. **a,** Structure of HKP. **b**, Optimization of the mass ratio of tRF-T11

mimic and HKP. **c**, The particle sizes and zeta potential of the HKP-tRF-T11 mimic.

**d**, Growth rate of tumors treated with tRF-T11 mimic, taxol and HKP alone. **e**,

Quantification of body weights in each group. **f**, Images of each tumor. **g**, No

significant difference in tumor weight was observed during treatment in the high-dose

tRF-T11 mimic group and the low-dose taxol group. **h**, Representative western blot

image of *TRPA1* protein levels in tumors of different treatment. **i**, Band intensity ratio

of *TRPA1*/$\beta$-actin in tumors treated with different doses of tRF-T11 mimic (n=6 per

group). The data are shown as the means $\pm$ SDs of 3 independent experiments

excluding those shown in **c** ($\pm$ SEM). *$P<0.05$, **$P<0.01$, ***$P<0.001$.

491                              **Figure 1**

493                                              **Figure 2**

496 **Figure 3**

**a**

R=[KHHHKHHHKHHHKHHHK]
K=L-Lysine
H=L-Histidine

**b**

HKP-tRF-T11 mimic

Naked tRF-T11 mimic

1:3   1:2   1:1   1:0.5   tRF-T11   HKP

**c**

Size Potential Distribution

41.2 mv

Size Distribution by Intensity

164.2 nm

**d**

Ratio of Tumor Volume vs Time (Day)

- HKP Control
- tRF-T11 mimic (71 nmol/kg)
- tRF-T11 mimic (177 nmol/kg)
- Taxol (2928 nmol/kg)
- Taxol (5855 nmol/kg)
- Injection day

51%

$P$ >0.1

**e**

Body Weight (g) vs Time (Day)

- HKP Control
- tRF-T11 mimic (71 nmol/kg)
- tRF-T11 mimic (177 nmol/kg)
- Taxol (2928 nmol/kg)
- Taxol (5855 nmol/kg)
- Injection day

**f**

HKP Control
tRF-T11 mimic (71 nmol/kg)
tRF-T11 mimic (177 nmol/kg)
Taxol (2928 nmol/kg)
Taxol (5855 nmol/kg)

**g**

Tumor Weight (g)

$P$ =0.138

HKP Control
tRF-T11 mimic (71 nmol/kg)
tRF-T11 mimic (177 nmol/kg)
Taxol (2928 nmol/kg)
Taxol (5855 nmol/kg)

**h**

TRPA1   110 kDa
$\beta$-actin   43 kDa

HKP Control   tRF-T11 mimic (71 nmol/kg)   tRF-T11 mimic (177 nmol/kg)

**i**

Fold change from HKP control group

$P$ =0.0054

HKP Control
tRF-T11 mimic (71 nmol/kg)
tRF-T11 mimic (177 nmol/kg)