Broadband Signal Rather than Frequency-Specific Rhythms Underlies Prediction Error in the Primate Auditory Cortex

Andrés **Canales-Johnson**[1,2,3,*,#], Ana Filipa **Teixeira Borges**[4,#], Misako **Komatsu**[5], Naotaka **Fujii**[5,,]
Johannes Jacobus **Fahrenfort**[6,7], Kai Joshua **Miller**[8], Valdas **Noreika**[1,N,*]

[1] Department of Psychology, University of Cambridge, Cambridge, United Kingdom.

[2] Center for Social and Cognitive Neuroscience (CSCN), Universidad Adolfo Ibáñez, Santiago, Chile

[3] The Neuropsychology and Cognitive Neurosciences Research Center (CINPSI Neurocog), Universidad Catolica del Maule, Talca, Chile

[4] Department of Integrative Neurophysiology, Center for Neurogenomics and Cognitive Research, VU University Amsterdam, Amsterdam, The Netherlands

[5] Laboratory for Adaptive Intelligence, RIKEN Brain Science Institute, Wako, Saitama, Japan

[6] Department of Psychology, University of Amsterdam, Amsterdam, The Netherlands

[7] Department of Experimental and Applied Psychology, Vrije Universiteit, Amsterdam, The Netherlands

[8] Department of Neurosurgery, Mayo Clinic, Rochester, United States of America.

[N] Lead Contact

[#] These authors contributed equally to this work

[*] Correspondence: afc37@cam.ac.uk (A.C.-J.), vn261@cam.ac.uk (V.N.)

Summary

Detection of statistical irregularities, measured as a prediction error response, is fundamental to the perceptual monitoring of the environment. We studied whether prediction error response is generated by neural oscillations or asynchronous neuronal firing. Electrocorticography (ECoG) was carried out in three monkeys, who passively listened to the auditory roving oddball stimuli. Local field potentials (LFP) recorded over the auditory cortex underwent spectral principal component analysis, which decoupled broadband and rhythmic components of LFP signal. We found that broadband component generated prediction error response, whereas none of the rhythmic components encoded statistical irregularities of sounds. The broadband component displayed more stochastic, asymmetrical multifractal properties than the rhythmic components, which revealed more self-similar dynamics. We thus conclude that the prediction error response is encoded by asynchronous neuronal populations, defined by irregular dynamical states which, unlike oscillatory rhythms, appear to enable the neural representation of auditory prediction error response.

Keywords
Auditory cortex; Broadband component; Local field potentials; Mismatch negativity; Multiscale multifractal analysis; Prediction error; Rhythmic components; Scaling behaviour; Scale-free dynamics; Spectral principal component analysis

Introduction

Detection of novel sensory information enables adaptive interaction with the surrounding environment (Clark, 2013; Whitmire and Stanley, 2016). In the predictive coding framework of brain functioning, this interaction is characterized by a reciprocal loop between sensory predictions and prediction error signals (Bastos et al., 2012; Friston and Kiebel, 2009). Neural mechanisms of prediction error are typically studied by presenting a series of "standard" stimuli with intermittently occurring deviant stimuli, also called "oddballs", and by contrasting brain responses between these stimuli categories (Chennu et al., 2013; Lumaca et al., 2019; Parras et al., 2017). This way, event-related potentials (ERP) and a range of neural oscillations have been identified as neural markers of prediction error. The most widely studied deviance ERP is the auditory mismatch negativity (MMN) – a negative deflection of electrical event-related potential recorded on the scalp or using intracranial electrodes (Halgren et al., 1995; Näätänen et al., 1978; 2007). MMN originates from the primary auditory cortex (Alain et al., 1998; Alho, 1995; Edwards et al., 2005), and it peaks around 150-200 ms in humans, whilst the peak latencies below 100 ms are typically reported in monkeys (Javitt et al., 1992; Komatsu et al., 2015). In addition to MMN, prediction error responses are observed in neural oscillations in a variety of frequency ranges including theta (3-8 Hz) (Choi et al., 2013; Fuentemilla et al., 2008; Hsiao et al., 2009; Ko et al., 2012; MacLean et al., 2014), alpha (8-12 Hz) (Ko et al., 2012; MacLean et al., 2014), beta (14-30 Hz) (Haenschel et al., 2000; MacLean et al., 2014) and gamma (>30 Hz) (Edwards et al., 2005; Eliades et al., 2014; Haenschel et al., 2000; MacLean et al., 2014; Marshall et al., 1996) ranges.

Several interpretations could be formulated aiming to explain the abundance of prediction error responses in the frequency dimension. First of all, there could be multiple independent neural mechanisms sensitive to stimulus deviance. This suggestion, however, does not explain why there would be so many distinct mechanisms with an identical functional role. Alternatively, frequency-specific detectors of prediction error might be only partially independent, forming hierarchical cross-frequency interactions. For instance, rhythms of different frequency bands could drive each other, e.g. delta phase could modulate theta amplitude and theta phase could modulate gamma amplitude in the auditory cortex (Lakatos et al., 2005). Yet another possibility – which we pursue in the present study – is that a broad frequency range of deviance responses, including theta, alpha, beta and gamma bands, points to a *broadband* prediction error response, which is not restricted to a particular frequency band, but instead is driven by an arrhythmic or asynchronous neural signal. In fact, a large number of studies reported deviance effects to run across several frequency bands (Chao et al., 2018; Haenschel et al., 2000; Hsiao et al., 2009; Ko et al., 2012; MacLean et al., 2014), arguably alluding to arrhythmic processing of unexpected stimuli.

The electrophysiological signal recorded by scalp EEG or local field potentials (LFP) is a summed
activity of both postsynaptic and action potentials. Post-synaptic potentials contribute to the rhythmic
oscillations of different frequency bands (Buzsaki et al., 2012), reflecting neural synchrony at specific
timescales. Contrary to this, empirical data analysis and modelling suggest that the average input
firing rate produces asynchronous, broadband changes across a wide frequency range (Miller et al.,
2009a, 2009b; Manning et al., 2009). Such rhythmic and broadband components of LFP signal can
be decomposed using spectral principal component analysis (spectral PCA) (Miller et al., 2009a,
2009b, 2017), this way separating synchronous and asynchronous neural activity. Broadband
component of the LFP power spectrum is commonly characterized by a power-law function (Freeman
and Zhai, 2009; He, 2014; Hermes et al., 2019), which reflects the lack of any specific temporal beat
(e.g. 10 Hz) in the signal. Contrary to this, rhythmic components produce frequency-specific spectral
peaks that deviate from the power law. In fact, the electrocorticography power is characterized by at
least three different power-law regions of which the transitions vary across individuals and recordings
in human (Chaudhuri et al., 2017; He et al., 2010) and non-human primates. The functional relevance
of this heterogeneous scaling is discernible as, for instance, levels of arousal across a gradual
progression from awake to grades of anaesthesia (Gifani et al., 2007) or to deep sleep (Ma et al.,
2006; Weiss et al., 2009) can manifest selectively within power-law changes at different timescales.
Such complex dynamics across different LFP timescales can be characterized by multiscale
multifractal analysis (MMA; Gierałtowski et al., 2012), developed to analyse signal fluctuations on a
wide range of timescales like those observed in LFP signals.

In the present study, we aimed to assess whether such broadband neural dynamics rather than
frequency-specific rhythms underlie prediction error in the auditory cortex in the primate brain. We
hypothesized that the broadband component of LFP has a multiscale scaling dynamics, distinct from
that of the rhythmic LFP components.

Results

Using epidurally implanted electrodes, we recorded electrocorticograms (ECoGs) from three awake
common marmosets, who passively listened to the stream of varying tones (see Fig 1A and Suppl.
Fig 1A-D). By contrasting neural responses to standard and deviant tones, which were physically
matched across expectancy conditions, we first identified MMN deviance response from the auditory
cortex (see Fig 1B). Afterwards, we decomposed the raw LFP signal into broadband and rhythmic
spectral components (following Miller et al., 2009a, 2009b, 2017; see Fig 1C). Spectral decomposition
allowed us to assess whether MMN is driven by the broadband rather than oscillatory components of
the LFP signal. In the following, we report a single-trial analysis that was carried out separately for

each monkey (referred to as Fr, Go and Kr), using electrodes located in the auditory cortex (see Fig
1B and Suppl. Fig 1E-H).

Figure 1 | Experimental design, classical ERP and spectral decoupling analysis. (A) Using a
roving oddball paradigm, 20 different single-tones were presented in the trains of 3, 5 or 11 identical
stimuli. Any two subsequent trains consisted of different tones. This way, while the adjacent standard
(depicted in black) and deviant (depicted in green) tones deviated in the frequency due to the transition
between the trains, the two expectancy conditions were physically matched, as the first and the last
tones of the same train were treated as deviant and standard tones in the analysis of the adjacent
stimuli pairs. **(B)** Time courses of ERP waveforms of the standard (black) and deviant (green) stimuli
conditions. 0 ms time point indicates the onset of a given tone. Error shades represent the standard
error of the mean (SEM), calculated across all trials at each time point. Data in this and other subplots
were recorded by the electrode marked in red in the ECoG montage insets. Each subplot represents
a different monkey. **(C)** Spectral decoupling. Temporally adjacent raw LFP segments of the standard
tone (i.e. the last stimulus of the previous train) and the deviant tone (i.e. the first stimulus of the
subsequent train) were extracted for the spectral PCA. First, Fast Fourier transform was used to
calculate log power (1-250 Hz) of the raw LFP signal, which was afterwards normalized across all
trials within a given expectancy condition. Normalized spectral snapshots were input into spectral
PCA, which separated broadband and rhythmic components. Principal spectral components were
reconstructed back to the time-series for the subsequent contrast between the expected and
unexpected stimuli conditions.

Auditory ERP in the raw LFP signal

First, we confirmed that perturbation of auditory cortex with a deviant tone compared to a preceding
standard tone increased the amplitude of auditory evoked potentials in the MMN time window (Fr:
$t(1,719) = -7.37$, $p < 0.001$, Cohen's $d = 0.275$; Go: $t(1,1439) = -4.60$, $p < 0.001$, Cohen's $d = 0.121$;
Kr: $t(1,1439) = -9.27$, $p < 0.001$, Cohen's $d = 0.244$; see Fig 1B). Latency of ERP peaks (58-66 ms)
was consistent with the previous MMN studies of non-human primates (Javitt et al., 1992; Komatsu
et al., 2015).

Auditory evoked responses reconstructed with broadband and rhythmic components

Aiming to differentiate broadband component of LFP signal from rhythmic sources, we carried out
spectral principal component analysis that decouples the power spectrum density (PSD) into
components reflecting different underlying neural dynamics (see Fig 1C and Methods). Using this
technique, a broadband component can be identified by a uniform power increase, i.e. a component
without clear peaks in the PSD, across a wide range of frequencies (see Fig 2A, H, O, red lines). In
addition to broadband spectral changes, the technique also reveals a diverse set of narrow-band
oscillatory components, revealed by peaks in the PSD (see Fig 2A, H, O, blue and black lines).

This way, three major principal spectral components (PSCs), one representing a broadband
component and two representing rhythmic components, were identified from the auditory LFP signal.
PSCs were highly consistent across three monkeys (see Fig 2A, H, O), matching tightly with the
original depiction of spectral principal component analysis (PCA) (see Fig 1A in Miller et al., 2009b).
In order to assess which of these three major PSCs encode auditory prediction error response,
components were back-projected to the time dimension.

We found that the Broadband PSC carried a characteristic auditory event-related broadband (ERBB)
response, reminiscent of auditory ERP, compared to largely flat responses derived from the rhythmic
PSCs with alpha (Rhythmic 1) and delta (Rhythmic 2) peaks. The ERBB response was evident in the
average of individual - standard and deviant - responses (see Fig 2B, I, P) as well as along the whole
sequence of 11 identical tones as compared to the tone sequences reconstructed from the Rhythmic
components (see Fig 2C-E,J-L,Q-S). Repeated measures ANOVA between the PSC (Broadband,
Rhythmic 1, Rhythmic 2) and the stimulus expectancy (standards, deviants) factors revealed the main
effects for the PSC and the stimulus expectancy, and the interaction between the PSC and the
stimulus expectancy factors (see Fig 2F,M,T). Post-hoc comparisons showed that the ERBB response
locked to the deviant tones had a larger amplitude compared to the ERBB response locked to the
standard tones in the Broadband PSC contrast, but not in the Rhythmic 1 PSC nor the Rhythmic 2

 PSC contrasts (see Fig 2G,N,U). We thus conclude that MMN response recorded by the ECoG of the

 auditory cortex is driven by broadband rather than rhythmic components of LFP signal.

Figure 2 | Event-related broadband response of stimulus expectancy. (A,H,O) Element magnitude of the major principal spectral components (PSCs) in the frequency domain (1-240 Hz). In this and other subplots, the Broadband PSC is depicted in red, the Rhythmic 1 PSC (alpha) in blue, and the Rhythmic 2 PSC (delta/theta) in black. **(B,I,P)** A narrow window of back-reconstructed time series of the broadband and rhythmic PSCs, locked to the onset of tones (0 ms). Standard and deviant stimuli are averaged together. **(C-E,J-L,Q-S)** Back-reconstructed time series of the Broadband and Rhythmic PSCs along a sequence of 11 identical tones. 0 ms indicates the onset of the deviant tone. **(F,M,T)** ANOVA results of the stimulus expectancy (standard, deviant) and the spectral component (Broadband, Rhythmic 1, Rhythmic 2) contrast. Significant main effects were observed for the PSC (Fr: $F_{(2,1438)}=341.70$, $p<0.001$, eta-squared = 0.322; Kr: $F_{(2,2878)}=113.00$, $p<0.001$, eta-squared = 0.073; Go: $F_{(2,2878)}=78.60$, $p<0.001$, eta-squared = 0.052) and the stimulus expectancy (Fr: $F_{(1,719)}=14.1$, $p<0.001$, eta-squared = 0.01; Kr: $F_{(1,1439)}=23.60$, $p<0.001$, eta-squared = 0.016; Go: $F_{(1,1439)}=4.81$, $p<0.029$, eta-squared = 0.003) factors, and the interaction between the PSC and the stimulus expectancy (Fr: $F_{(2,1438)}=17.20$, $p<0.001$, eta-squared = 0.02; Kr: $F_{(2,2878)}=20.80$, $p<0.001$, eta-squared = 0.014; Go: $F_{(2,2878)}=15.49$, $p<0.001$, eta-squared = 0.011). Error bars indicate the standard error of the mean (SEM). '=' refers to the main effects, 'x' refers to the interaction. **(G,N,U)** Stimuli locked waveforms show post-hoc comparisons between the standard and deviant stimuli in the broadband and rhythmic PSCs, which revealed larger amplitude for the deviant stimuli in the Broadband PSC contrast (Fr: $t=6.96$, $p_{Bc}<0.001$; Kr: $t=7.84$, $p_{Bc}<0.001$; Go: $t=5.48$, $p_{Bc}<0.001$), but not in the Rhythmic 1 (Fr: $t=0.378$, $p_{Bc}=1.00$; Kr: $t=0.612$, $p_{Bc}=1.00$; Go: $t=0.397$, $p_{Bc}=0.99$) nor the Rhythmic 2 (Fr: $t=0.812$, $p_{Bc}=1.00$; Kr: $t=-0.033$, $p_{Bc}=1.00$; Go: $t=-1.567$, $p_{Bc}=1.00$) PSC contrasts.

Cross-individual decoding of stimulus expectancy with broadband and rhythmic components

While the single-subject results of ERBB response were highly consistent across all three monkeys (see Fig 2), we wanted to establish whether the broadband prediction-error response of an individual monkey can be extrapolated to other individuals of the same species. This would indicate that the prediction error information generated in the auditory cortex is implemented similarly across monkeys. We thus assessed the cross-individual generalizability of the ERBB response by decoding the stimuli expectancy using the Broadband and Rhythmic PSCs. Using all trials of a respective PSC of one monkey, we trained a linear discriminant (LDA) classifier to learn stimuli categories (standard vs. deviant) in the auditory cortex electrode (see Fig 1B and S1). Afterwards, we decoded stimuli categories using the same PSC in a different monkey. Using Broadband PCS, we obtained significant decodability in all six pairs of comparisons, i.e. cross-individual decoding between 3 monkeys (see Fig 3A). The time windows of significant decoding above chance level (50% AUC) were consistent with MMN and BRBB responses (see Fig 1B and Fig 2G, N, U). Contrary to this, no significant cross-individual decodability was observed using Rhythmic 1 and 2 PCSs (see Fig 3B, C). These findings confirm the cross-individual generalizability of the broadband PSC encoding of stimulus expectancy.

Figure 3 | Decoding of stimulus expectancy across monkeys with broadband and rhythmic components. Classification of stimulus expectancy conditions (standard, deviant) was carried out in one of the monkeys (plotted in green). Afterwards, the classifier was tested on the other two monkeys

264 (plotted in purple). Time points of significant decoding of stimulus categories above chance level (50%
265 of AUC), estimated with a cluster-based permutation test, are depicted in red. **(A)** Decoding was
266 successful in all six pairs using Broadband PSC. **(B)** Decoding did not exceed chance level using the
267 Rhythmic 1 PSC. **(C)** Decoding did not exceed chance level using the Rhythmic 2 PSC.
268

Multiscale multifractal analysis of broadband and rhythmic neural components

We hypothesized that the Broadband component has a distinct multiscale temporal organization

within the milliseconds of the evoked response than the one from the rhythmic components. In

particular, we sought to characterize the scale-free temporal properties of the segregated neural

components. These properties relate to the functional state of neural systems (Papo, 2014; He, 2014;

Werner, 2010). We further hypothesized that the broadband component–-the neural signal subserving

oddball detection–-has a more stochastic multiscale temporal organization which allows greater

dynamical flexibility. The scale-free nature of the neuronal population firing rate, manifested in the

broadband PSC (Miller, 2009; Manning et al., 2010), is usually estimated by determining the slope of

the log-log function of PSD (power vs. frequency), also referred to as 1/f (fractal) scaling. However,

often the PSD is not characterized by a single exponent and may show a scale-dependence (Miller,

2009; Chaudhuri et al, 2017) and/or different scaling depending on the statistical moment and hence

exhibit multifractality (Nagy et al. 2017). Indeed, the single-trial auditory responses, standards and

deviants, revealed a piecewise linear decay of power with frequency in each marmoset (Fig 4A),

suggesting that the dynamics of the underlying processes may have scale-free properties but also a

heterogeneous scaling dependent on frequency (timescale). This is noticeable by the different slopes

which characterize the 1/f-like PSD depending on the frequency range (Fig. 4A), precluding the fitting

of a unique line to estimate the slope across the whole spectrum. Thus, to fully characterize the scale-

free properties of the three components, we sought to test for the presence of scale-dependent

multifractality in the series of increments of neural activity in the marmoset auditory cortex.

Multifractality requires the presence of different scaling exponents (h) of different moments of the

fluctuations (q) over a wide range of timescales (s) (Kantelhardt et al., 2002). Using multiscale

multifractal analysis (MMA; Gierałtowski et al., 2012) (Fig 4B), which is particularly suited to analyze

complex systems which exhibit fluctuations of activity on a wide range of timescales and a broad

distribution of values, we found that all the three PSCs show considerable variability in the values of

the generalized q and s-dependent scaling exponents ($h(q,s)$ – *Hurst surface*) (Fig 4D). The Rhythmic

components displayed similar surfaces, distinct from the nonlinear profile of h across $q \in [-5, 5]$ of

the Broadband PSC activity. These results were consistent across monkeys (see Fig S5). The

average tendency across scales revealed a nearly linear dependence of h with q for both Rhythmic

components suggesting their underlying dynamics appears multifractal. Conversely, although the

dynamics of the Broadband PSC is also multifractal (in the sense that its fractal properties depend on

q), the profile is nonlinear and distinct for small (*q* < 0) and large (*q* > 0) fluctuations (Fig 4E). We note

that the conventional Hurst scaling analysis (*q* = 2 results) did not provide a clear distinction between

the Broadband and Rhythmic 2 components. Furthermore, averaged surface values of *h* suggest the

Broadband fluctuations can be quasi-stochastic (*h* ~ 0.5) or persistent without obeying strictly a power-

law (*h* ~ 1.1), depending on if large (*q* >0) or small fluctuations (*q* < 0) are considered (Fig 4F).

Conversely, Rhythmic 1 and Rhythmic 2 fluctuations ranged from being close to Brownian motion

(integrated white noise, *h* ~ 1.5) to scale-free. There was a qualitative agreement on the values across

monkeys (Fig 4E, F). Thus, while all three PSC components showed scale-free properties, there were

significant differences in the apparent stochasticity, expressed as *h(q)*, between the components (Go:

RANOVA $F_{(2,20)}$=103, $p<0.001$, eta-squared=0.339; Kr: RANOVA $F_{(2,20)}$=134, $p<0.001$, eta-

squared=0.404; Fr: RANOVA $F_{(2,20)}$=40.2, $p<0.001$, eta-squared=0.228). For all three monkeys, the

Broadband component exhibited lower *h(q)* values compared to the Rhythmic 1 (Go: t=-14.05,

$p<0.001$; Kr: t=-13.39, $p<0.001$; Fr: t=-8.54, $p<0.001$) and Rhythmic 2 (Go: t=-9.54, $p<0.001$; Kr: t=-

14.88, $p<0.001$; Fr: t=-6.64, $p<0.001$) components.

In order to determine whether the multifractality, depicted in the Hurst surfaces (Fig 4D), is caused by

the temporal correlations of the signal distribution, we created a distribution of shuffled surrogates,

i.e. copies of the original data with identical mean, variance and histogram distribution but no temporal

structure. While the mean Hurst surfaces of the surrogates distribution showed for all monkeys a

decrease in multifractality ($p<0.001$) (Fig 4D; Suppl. Figure 5), the averaged Hurst exponent values

indicated that the neural dynamics approached randomness ($h = 0.5$) for all monkeys (Fig 4H).

Therefore, the multifractality is caused mostly by the temporal correlations but also by a fat-tailed

probability distribution. We subsequently computed the multifractal spectrum f(α). Analogously to a

Fourier analysis, i.e. the decomposition of a signal into a sum of components with fixed frequencies,

f(α) can be understood as decomposition of a signal into a set of exponents α (Mandelbrot, 2003).

Their relative presence in the signal is weighted by the f(α) function. The Broadband activity

interweaved more densely sets of singularities that are less self-similar than those of the Rhythmic

components and displayed a lower degree of multifractality and a more asymmetrical $f(\alpha)$ (Fig 4G

,H), suggesting its dynamics differs from simple multiplicative cascades. The shape of the multifractal

spectra for the Broadband activity also displayed a right-truncation (Fig 4C,G and Suppl. Figure 5),

which is expected due to the leveling of *h(q,s)* for *q* < 0 (Ilhen, 2012).

To sum up, MMA analysis revealed that the generalized scale-dependent Hurst exponent *h(q,s)* and

the derived *f(α)* curves of the dynamics of Broadband and Rhythmic components show multifractality

as well as marked differences of this property. Importantly, the Broadband component more closely

approached a stochastic asymmetrical multifractal distribution.

Figure 4 | Multifractal characterization of the Broadband and Rhythmic dynamics. (A) Double logarithmic plots of the power spectral densities of the Broadband (blue), Rhythmic 1 (pink) and Rhythmic 2 (green) components during all trials of the auditory MMN paradigm reveal a piecewise approximately linear decay of power with frequency. The average scaling (fractal) properties of the power spectral densities (last column) are distinct across frequencies, spectral components and

marmosets. **(B)** Multiscale Multifractal Analysis (MMA) method. *Left.* Log-log plots of the fluctuation
functions $F_q(s)$ for each $q \in [-5,5]$, color-coded from dark blue (q = -5) to dark red (q = 5) and scale
s (in ms) of the time series correspondent to the Broadband activity of monkey Go. The Hurst (scaling)
exponent ($h_{1,2,...,11}$) is obtained by determining the slope of a linear fit within a window lasting the
period ($s_{1,2,...,11}$) marked with vertical dashed lines. Three example scales are displayed: $s_1 \in [10,50]$,
$s_2 \in [20,100]$ and $s_{11} \in [120,600]$. *Right.* Computed Hurst exponents $h(q,s)$ are displayed in a (Hurst)
surface plot grid. As an example, the cells corresponding to $h_{1,2,...,11}$ (q = 2; s = 1, 2 or 11) are
highlighted with their respective colors (light grey, lilac, green). **(C)** The Hurst surface can also be
converted into a multifractal spectrum, $f(\alpha)$, which describes how densely the singularities (i.e. scaling
exponents, α) are distributed in a signal. The parabolic vertex shows the central tendency, a measure
of the regularity present, and the width, the degree of multifractality ($\Delta\alpha$). **(D)** Hurst surfaces ($h(q,s)$
) of the component activities (each column) and the $< h(q,s) >$ of a distribution of 50 shuffled
surrogates. Monkey Go is shown here; for the other monkeys, see Figure Suppl. 5. **(E)** Scaling
properties averaged for all scales. The Hurst exponent h dependency on q is evident for all
components, suggesting their multifractality. **(F)** Mean (+/-SD) of the Hurst surfaces ($< h(q,s) >$)
suggests that the Broadband activity has an overall more random profile. Each group of 3 dots with
error bars refers respectively to $< h(q,s) >$ across all scales (s) for negative, positive and all values
of q. Individual results for the Broadband (BB), Rhythmic 1 (R1) and Rhythmic 2 (R2) PSCs are
displayed in variations of blue, pink and green colors, respectively. Bottom row shows the values
obtained for the distribution of surrogates. **(G)** Multifractal spectrum of the three PSCs of Monkey Go,
the lightness of the colors represents the results for different scales (s) (light $\rightarrow$ dark with increasing
scales $s_{1,2,...11}$). **(H)** Central tendency of the multifractal spectrum (α_{max}), degree of multifractality ($\Delta\alpha$
) and asymmetry of the spectrum ($as_{\Delta\alpha}$) for the three types of activity ((Broadband (BB): blue;
Rhythmic 1 (R1): pink; Rhythmic 2 (R2): green)). Each monkey is displayed in a different shade of the
colors.
370
371

Discussion

In the present study, we compared two alternative views of prediction error processing, namely

whether LFP oscillatory vs. broadband components of neural activity encode deviant sensory stimuli.

We found that auditory MMN response, a classical marker of prediction error, is primarily driven by

the broadband component of LFP signal. Given that broadband PSC reflects the mean firing rate of

neuronal populations (Hermes et al., 2014, 2017; Manning et al, 2009; Miller, 2010), and that neuronal

spiking correlates tightly with the high-frequency LFP in the auditory cortex (Mukamel et al., 2005),

our findings indicate that prediction error response depends on the asynchronous neuronal firing rate

rather than oscillatory neuronal encoding of incoming stimuli.

We first replicated previous research by showing that auditory MMN is generated in the auditory

cortex. Afterwards, we separated the LFP signal into Broadband and Rhythmic components and

repeated MMN analyses separately for each component. While the main two rhythmic components

present in the data were not able to distinguish between the standard and deviant tones, the

broadband component indexed the stimuli difference in the auditory cortex. The findings were highly

consistent across all three marmosets, and the cross-individual decoding successfully classified

stimuli category (standard or deviant) when data were trained on one monkey and tested on a different

one. Importantly, significant decoding was observed only with the Broadband PSC, as the decoding

was unsuccessful with the Rhythmic PSCs.

Our study challenges the generality of low-frequency neural oscillations as an instrument for

enhancing (Schroeder and Lakatos, 2009) and gating information in the auditory cortex (Lakatos et

al., 2013). In particular, it has been argued that a phase reset of slow frequencies in the range of delta

and theta oscillations may underlie prediction error response (Arnal et al, 2015; Fuentemilla et al.,

2008; Ko et al., 2012). However, we show that the Rhythmic 2 component with a distinctive delta peak

and a considerable contribution from theta range activity (Fig 2A, H, O) does not discriminate between

standard and deviant tones. Likewise, the Rhythmic 1 component representing alpha range activity

did not encode prediction error response, challenging previous studies that linked MMN to alpha band

power (Ko et al., 2012; MacLean et al., 2014). This contrast is striking given that these ideas were

deemed most relevant in the context of rhythmic as opposed to continuous stimulation (Schroeder

and Lakatos, 2009), and rhythmicity is prevalent in the current oddball paradigm. However, it is

important to emphasize that we do not claim that low-frequency neuronal counterparts of neural

activity do not contribute to predictive coding: long-term dependencies are relevant in sensory

prediction in the auditory cortex (Rubin et al., 2016). Rather, we suggest they enable information

processing through more flexible, dynamical unstable codes than an oscillatory code.

Our results demonstrate that prediction error processing is subsumed by an asynchronous broadband

activity with dynamical properties very distinct from that of the rhythmic components. Importantly, this

difference is unveiled when a multiscale approach is used to characterize fluctuations with several

degrees of resolution (multiple fractal hierarchies) and it is patent in the surfaces and multifractal

spectrum; the difference is equivocal by simply observing the power spectral densities or doing a

classical Hurst analysis. The broadband component is distinctive from the other components by its

lower level of self-similarity and multifractality and also by its asymmetric multifractal spectrum. The

presence of multifractality in the broadband and rhythmic electrocorticographic activity suggests it

may be a generic feature of neuronal networks and cognition may operate by modulations of this

property (Papo, 2014). Arguably, spike trains represent information with a multifractal temporal coding

(Fetterhoff et al., 2015) and the integrated multifractal spectrum permit to infer the tuning curve of

spiking activity in primates (Fayyaz et al, 2019). This could be a more effective mechanism of how

information is encoded in neuronal assemblies than the one provided by oscillatory rhythms. This

hypothesis is bolstered by ideas that synchronization *per se* only arises in collective states where no

new information can be created. In contrast, adaptive behaviour emerges from more subtle forms of

coordination, e.g. through the metastability or asynchronous coupling of spatiotemporal patterns of

neural activity (Friston, 2000; Tognoli and Kelso, 2014). The multifractality present in the recordings

reveals how the macroscopic neural dynamics is intermittent, its spectral density changes with time,
which has been hypothesized as a facet of temporal metastability (Friston, 1997; Tognoli and Kelso,
2014); at the core of metastability is the broken symmetry of spatiotemporal patterns (Kelso and Kelso,
1995) which was only present in the broadband activity. In fact, the more asymmetrical multifractal
spectra of the broadband activity suggest this feature may be a proxy of a dynamical regime which
allows the breakdown of symmetry, characteristic of systems that can perceptibly or meaningfully
react to afferent inputs (Freeman and Vitiello, 2006).

Furthermore, the prediction error processing by neural assemblies in the auditory cortex is sustained
by an irregular broadband component with small fluctuations lying in a tight range of the non-ergodic
dynamical regime $\underline{h} > 1$, which has been proposed as an explanation for the $1/f$ noise of cognitive
processes (Grigolini et al., 2009), and large fluctuations with stochastic-like properties. Altogether,
this result emphasizes the importance of asynchronous chaotic irregular states for optimal
responsiveness to external stimuli (Beaman et al., 2017; Renart et al., 2010; Zerlaut and Destexhe,
2017).
439
Our findings were enabled by a novel approach to quantify these complex dynamics of neural
systems, the so-called brain's "stochastic chaos" (Freeman et al., 2001). Future studies are
anticipated to extend MMA analysis of MMN to wider frequency ranges (>100 Hz), with a fine-grained
resolution to arguably uncover the spike tuning underlying sensory-state discrimination (Fayyaz et al.,
2019). The broadband prediction error response should be further studied using hierarchical auditory
prediction paradigms that can discriminate sensory and top-down prediction error responses
(Bekinschtein et al., 2009; Chennu et al., 2013, 2016). Developed in human studies, such paradigms
have been recently successfully applied in the common marmosets (Chao et al., 2018). Furthermore,
while the marmoset model of MMN deemed successful and very stable, as indicated by cross-
individual decoding, the current study should be replicated using LFP recordings in humans.
450
Importantly, our findings provide a unifying framework for the micro- to macro-level neural
mechanisms of prediction error response. While most of the auditory MMN studies are carried out at
the macro-level using scalp EEG recordings or meso-level LFP, auditory prediction error responses
have also been identified using single-neuron recordings (Nieto-Diego and Malmierca, 2016; Parras
et al., 2017; Pérez- González et al., 2005; Solomon and Kohn, 2014; Ulanovsky et al., 2003, 2004).
In particular, individual neurons located in the primary auditory cortex increase spiking rate following
presentation of oddball stimuli, which has been observed in different mammal species, including cat
(Ulanovsky et al., 2003, 2004), rat and mouse (Nieto-Diego and Malmierca, 2016; Parras et al., 2017).
Similar responses have also been identified in sub-cortical neurons (Parras et al., 2017; Pérez-
González et al., 2005). In particular, a subclass of neurons located in the dorsal and external cortices

of the inferior colliculus of the rat respond selectively to novel auditory stimuli, while muting their
response to repetitive stimuli (Pérez- González et al., 2005). A recent study of single-neuron activity
recorded from different auditory centers in rats and mice suggests that prediction error response is
organized hierarchically along the non-lemniscal auditory pathway comprising of inferior colliculi,
medial geniculate bodies and the primary auditory cortex with sensitivity to the deviant tones
increasing along the pathway (Parras et al., 2017). MMN-like deviance sensitivity of firing rate
increases further in the non-primary regions of auditory cortex (Nieto-Diego and Malmierca, 2016).
How do such micro-level single-neuron responses relate to the MMN potentials recorded with ECoG
and/or EEG? Are different neuronal mechanisms at different levels of measurement, such as single
neuron spiking rate vs. neuronal oscillations recorded using ECog/EEG?

Our study indicates that increased neuronal firing rate underlies prediction error responses not only
at the micro-level of single-neuron recordings, but also at the higher meso-level LFP measurements.
In particular, we show that MMN prediction error response is driven by the Broadband component of
the meso-level LFP signal. Given that the Broadband PSC reflects largely stochastic neuronal firing
rate, as suggested by previous modeling studies (Miller et al., 2007; Miller et al., 2009a), our findings
indicate that auditory prediction error response is indeed encoded at a single action potential level
within neuronal populations, which generate broadband signal at the meso- and most likely macro-
level electrophysiology. Broadband LFP activity provides indirect access to the total spiking output of
neurons, as shown by a growing number of experiments and simulations (Crone et al., 2011;
Freeman, 2004; Rash et al., 2008). Thus, the reported Broadband activity in this study provides a
'proxy' for investigating the neuronal mechanisms underlying auditory prediction error. As such, the
mesoscopic information of the Broadband LFP component represents a crucial link between
macroscopic-level EEG and the microscopic-level spiking activity of neural populations (Buzsaki et
al., 2012).

How could our LFP-based broadband results be reconciled with an abundant literature on frequency-
specific MMN results, mostly derived from EEG experiments that do not find broadband MMN
response across all frequencies? Miller (2010) argues that low-frequency range of broadband effects
can be obscured by coincident changes in specific rhythmic phenomena. We further suggest that
EEG artefacts may decrease signal-to-noise ratio in frequency-specific segments of the broadband
signal, in which case only relatively clean segments would survive as significant detectors of
prediction error response. For instance, blink artifacts may distort neural signal in the delta and theta
frequency range (Gasser et al., 1992), whereas muscular artifacts are likely to interfere with the beta
and gamma range activity (van de Velde et al., 1998). Similarly, spontaneous fluctuation of alertness
level, which is likely to occur during passive "oddball" paradigms, would interfere with neural
processing in the theta and alpha frequency range (Noreika et al., 2019a, 2019b). Thus, depending

on the experimental demands, the selection and training of participants, and the data preprocessing
steps, certain segments of the broadband signal may be occluded by artifactual or irrelevant signals
when contrasting standard and novel stimuli, yielding frequency-specific deviance responses that in
fact originate from scale-free broadband component of neuronal signal. The suggested role of EEG
artifacts in the preclusion of broadband response could be tested using simultaneous EEG and LFP
recordings. Alternatively, too often classical frequency bands are loosely equated to specific rhythms
(Lopes da Silva, 2013) and the views of collective neural network activity as oscillations lend too much
emphasis on "rhythmicity" (Cole and Voytek, 2017) when in reality, in those narrow-band analyses
perhaps no characteristic frequency oscillation was present and/or may even be spurious and caused
by filtering (de Cheveigné and Nelken, 2017).

To conclude, we show that in a well-studied paradigm of auditory prediction error, oscillations do not
constitute a means to temporally constrain information processing. They are perhaps the tips of the
iceberg, the latter being an arrhythmic broadband component with asymmetric multifractal stochastic
properties at several timescales. Our paper establishes the relevance of the broadband activity to
encode relatively low-level auditory patterns and provides a theoretical background and empirical
tools to probe which predictive values lie under the "noisy" surface in other paradigms and sensory
modalities.

515
516 **Methods**
517
518 *Subjects*
519
We used three adult male common marmosets (*Callithrix jacchus*) that weighed 320–380 g. Monkeys
were implanted with ECoG electrode array under general anaesthesia, and all efforts were made to
minimize suffering. All surgical and experimental procedures were performed in accordance with the
National Institutes of Health Guidelines for the Care and Use of Laboratory Animals and approved by
the RIKEN Ethical Committee (No. H26-2-202). ERP data of one monkey (Fr) was reported previously
(Komatsu et al., 2015), whereas datasets of monkeys Go and Kr are new.

526
527 *Implantation of ECoG arrays*
528
Chronically implanted, customized multichannel ECoG electrode arrays (Fig. S1) (Cir-Tech Inc.,
Japan) were used for neural recordings (Komatsu et al., 2015; 2017). We implanted 32 (the left
hemisphere of monkey Fr), 64 (the right hemisphere of monkey Go), and 62 (the right hemisphere of
monkey Kr) electrodes in the epidural space. For 32 electrode array, each electrode contact was 1mm

in diameter and had an inter-electrode distance of 2.5–5.0 mm (Komatsu et al., 2015). For 64
electrode array, each electrode contact was 0.6mm in diameter and had an inter-electrode distance
of 1.4mm in a bipolar pair (Komatsu et al., 2017). The electrode-array covered the frontal, parietal,
temporal, and occipital lobes. The additional 4 electrodes of monkey Fr covered part of the right frontal
lobe. The animals were initially sedated with butorphanol (0.2 mg/kg i.m.), and surgical anaesthesia
was achieved with ketamine (30 mg/kg i.m.) and medetomidine (350 µg/kg i.m.). The animals were
then positioned in a stereotaxic frame (Narishige, Japan) and placed on a heating pad during surgery.
Vital signs were monitored throughout surgery. Implantation of the electrode-arrays involved the
removal of a bone flap (~2 cm along the anterior-posterior axis and ~1 cm along the mediolateral axis)
over the parietal cortex. The array was advanced into the epidural space. After positioning the
electrode-array, connectors were attached to the bone using dental acrylic and titanium (size 1.0 x
0.1mm) or PEEK (size 1.4 x 2.5 mm) screws. The reference electrodes were placed in the epidural
space and the ground electrodes in the episkull space. The anti-inflammatory corticosteroid
dexamethasone (1.25mg/kg, i.m.) was administered after surgery to prevent brain swelling. The
animals were given antibiotics and analgesics daily for 5 days after surgery. Following the animals'
recovery, the position of each electrode in the arrays was identified based on computer tomography,
and then co-registered to a template T1-weighted anatomical magnetic resonance image (MRI)
(http://brainatlas.brain.riken.jp/marmoset/; Hikishima et al., 2011) (monkey Fr) or pre-acquired MRI
(monkeys Go and Kr) using MRIcron software (http://www.mricro.com; Rorden et al., 2007). In all
monkeys, the electrode-array covered the frontal, parietal, occipital, and temporal cortices, including
the primary auditory area (Fig. 2A-C and S2).
554
555   ***Stimuli and task***
556
We adopted a roving oddball paradigm (Cowan et al., 1993; Haenschel et al., 2005; Garrido et al.,
2008). The trains of 3, 5, or 11 repetitive single-tones of 20 different frequencies (250–6727 Hz with
intervals of 1/4 octave) were pseudo-randomly presented. Tones were identical within each tone-train,
but differed between tone-trains. Because tone-trains followed on from one another continuously, the
first tone of a train was considered to be an unexpected deviant tone, because it was of a different
frequency from that of the preceding train. The final tone was considered to be an expected standard
tone, because it was preceded by several repetitions of this same tone. To avoid analytical artefacts
stemming from differences in the number of standard and deviant stimuli, we considered only the last
tone of a train as standard. Standards and deviants were presented 240 times in a single recording
session. Pure sinusoidal tones lasted 64 ms (7 ms rise/fall), and stimulus onset asynchrony was 503
ms. Stimulus presentation was controlled by MATLAB (MathWorks Inc., Natick, MA, USA) using the
Psychophysics Toolbox extensions (Pelli, 1997; Brainard and Vision, 1997). Tones were presented

through two audio speakers (Fostex, Japan) with an average intensity of 60 dB SPL around the
animal's ear.

ECoG recording and preprocessing

ECoG recordings were taken in the passive listening condition while monkeys were awake. In each
recording session, the monkey Fr was held in a drawstring pouch, which was stabilized in a dark
room, and the monkeys Go and Kr sat on a primate chair in a dimly lit room. The length of a single
session was about 15 min: the first 3 min of data were used for many standard stimuli (data are not
shown in this paper) and the remaining 12 min of data were used for the roving oddball sequences.
For monkey Fr, data from 3 sessions were used for analysis, which resulted in 720 (=240 × 3) standard
and deviant presentations. For monkeys Go and Kr, data from 6 sessions were used for analysis,
which resulted in 1440 (=240 × 6) standard and deviant presentations.

ECoG signals were recorded at a sampling rate of 1 kHz per channel. In the signal preprocessing,
those signals were re-referenced using an average reference montage, and high-pass filtered above
1 Hz. We segmented datasets from −903 to 400 ms relative to the onset of the unexpected tone, so
that each segment would include a pair of a deviant and a standard immediately preceding the
deviant, as well as a baseline of 400 ms preceding the standard tone. The segments were then divided
into standard epochs and deviant epochs (-400 ms to 400 ms). Parts of the dataset are shared in the
public server Neurotycho.org (http://neurotycho.org/; Nagasaka et al., 2011).

ECoG electrode-of-interested was identified functionally by contrasting time-frequency charts
between standard and deviant stimuli (0-350 ms), separately for each electrode (see Suppl. Fig. 2-4).
The Hilbert transform was applied every 10 Hz and z-scored with respect to the baseline period (-100
ms to 0 ms). One electrode with the largest high-gamma difference between the standard and deviant
tones (Edwards et al., 2005; Eliades et al., 2014; Haenschel et al., 2000; MacLean et al., 2014;
Marshall et al., 1996) was selected for each monkey for further analyses. In all three monkeys, the
selected electrode-of-interest was located in the auditory cortex (see Suppl. Fig. 1E-H).

Event-related potentials

For ERP analysis of the raw signal, a low-pass filter of 40 Hz was used. ECoG recordings were re-
referenced with respect to the common average reference across all electrodes. Data were then
epoched around the onset of tones (-100 ms to 350 ms), and baseline correction was applied by
subtracting the mean of the 100 ms period before the stimulus onset. MMN was assessed by
comparing the standard ERP and deviant ERP.

Decoupling the cortical spectrum to isolate Broadband and Rhythmic spectral components

To extract the course of broadband spectral activity, we carried out the spectral decoupling of raw LFP signal (Miller et al. 2009a, 2009b, 2017). As for the ERP analysis, ECoG potentials were re-referenced with respect to the common average reference across all electrodes. For the selected electrodes-of-interest (see above), discrete samples of power spectral density (PSD) were calculated using -200 to 400 ms epochs centered at the stimuli onset. With trials from both conditions (standards and deviants) grouped together, individual PSDs were normalized with an element-wise division by the average power at each frequency, and the obtained values were log-transformed. An inner product matrix of these normalized PSDs was diagonalized with a singular value decomposition, and was then applied to identify components of stimulus-related changes in the PSD. The eigenvectors (Principal Spectral Components or PSCs) from this decomposition revealed distinct components of cortical processing. Continuous time-frequency power charts were calculated using complex Morlet wavelets. These power charts were then normalized in the same way as the discrete spectra and projected onto the first PSC (broadband), second PSC (alpha rhythm, ~ 10 Hz), and third PSC (delta rhythm, ~2 Hz), separately. The raw time series were smoothed with an 80-ms Gaussian envelope (SD 80 ms), z-scored per trial, using a pre-stimulus period between -100 to 0 ms, and exponentiated, and then a value of 1 was subtracted (setting the mean at 0). The first PSC allowed to obtain the "broadband time course" which has been shown to reflect a power law in the cortical PSD (Miller et al., 2009a), and the second and third PSCs uncovered the "rhythmic time courses" with distinct frequency peaks.

Cross-individual decoding

To assess cross-individual generalizability of our findings, a univariate temporal decoding model was applied on each individual PSC time-courses on the selected auditory cortex electrodes, aiming to decode the stimuli expectancy categories, i.e. standards vs deviants. (Figure 3). The ADAM-toolbox was used on the Broadband and Rhythmic PSC time-courses with epochs from -100 ms to 400 ms (Fahrenfort et al., 2018). Crucially, and for each individual component, we trained a linear discriminant (LDA) classifier in one monkey and tested in a separate monkey for obtaining cross-individual decodability of stimuli expectancy category, i.e. standard vs deviant trials. As decoding algorithms are known to be time-consuming, data were downsampled to 250 Hz. Next, a backward decoding algorithm, using either stimulus expectancy category was applied according to a tenfold cross-validation scheme. A linear discriminant analysis (LDA) was used to discriminate between stimulus classes (e.g. deviant versus standard trials) after which classification accuracy was computed as the area under the curve (AUC), a measure derived from the Signal Detection Theory. AUC scores were tested per time-point with double-sided t-tests per subjects against a 50% chance-level. These t-tests

643   were corrected for multiple comparisons over time, using cluster-based permutation tests (p<0.05,
644   1000 iterations). This procedure yields time clusters of significantly above-chance classifier accuracy.

646   ***Dynamical characterization of the scaling behaviour***
647

To characterize the scaling properties of the neural activities of all PSCs of all monkeys, we combined
all individual trials for each PSC after removing their baseline – the resulting series had a length of
1327920 (Monkey Fr) and 2655840 samples (Monkeys Kr and Go) – and quantified the relationship
between ln(power) and ln(frequency).

Continuous power spectral densities. The power spectral density (band: 1–100 Hz) of each combined
time series for the principal components studied was computed by applying the modified Welch
periodogram method as implemented in Matlab's *pwelch()* function. We used 50% overlapping Hann
windows of 1.024 s.

Multiscale Multifractal Analysis (MMA). The nonstationarity of neural dynamics (Paluš, 1996) and, in
particular, the existence of changing points (crossovers) in the scaling laws in the marmoset
electrocorticographic data, which vary across individual and PSCs, precluded a pre-defined selection
of the scales of interest and called for a data-driven scaling analysis robust to nonstationarity. Thus,
to characterize the scaling behaviour, we used a method designated Multiscale Multifractal Analysis
(MMA) (Gierałtowski et al., 2012). MMA is an extension of the Detrended Fluctuation Analysis (DFA)
(Peng *et al.*, 1995), an established method to quantify the monofractal scaling behaviour of
nonstationary signals, robust to some extrinsic trends (Hu *et al.*, 2001). DFA is essentially a modified
root mean square (RMS) analysis of a random walk (Peng *et al.*, 1995). Briefly, for a given time series
x_k of length N, the profile $y(k)$ is determined by integrating the time series, then $y(k)$ is split into non-
overlapping segments with length s which are detrended by subtracting the local least-squares line fit,
$y_s(k)$. Since N/s is often not an integer, to avoid discarding data samples, a second splitting is
performed starting from the end of the time series; a total of $2N$ segments are considered. The root-
mean-square fluctuation of integrated and detrended time series is given by:

$$F(s) = \sqrt{\frac{1}{2N}\sum_{k=1}^{2N}[y(k) - y_s(k)]^2}$$

675   A generalized version of this method—the Multifractal Detrended Fluctuation Analysis (MF-DFA)
676   (Kantelhardt *et al.,* 2002)—permits to further characterize the fluctuations by inspecting
677   simultaneously small and large fluctuations q dependent on the multifractal parameter :

$$F_q(s) \equiv \left\{ \frac{1}{2N_s} \sum_{k=1}^{2N} [y(k) - y_s(k)]^{\frac{q}{2}} \right\}^{\frac{1}{q}}$$

The conventional DFA corresponds to the situation in which $q = 2$, and it is for stationary random

processes equivalent to the Hurst exponent (Hurst, 1951). The process is repeated for several lengths

of s; typically, $F_q(s)$ increases with s and displays the asymptotic behaviour $F(s) \sim n^{h(q)}$. The

generalized Hurst exponent $(h(q))$ is estimated by extracting the slope of a linear least-square

regression of $F_q(s)$ on a log-log plot for a given set of s values. The MMA algorithm's advantage is that

instead of prefixing a unique range of scales to estimate the scaling behaviour, it allows to scan for

several scale-ranges yielding a quasi-continuous characterization of the scaling behaviour $(h(q))$,

which may vary along scales (frequencies of the power spectrum): the result is a scaling exponent

depending on both *q* and *s*—$h(q,s)$. It can be visualized in a grid, the Hurst surface, each cell of which

corresponds to a value of *q* and a given range of scales *s*.

We applied MMA to the PSCs of the 3 marmosets within a range of $q \in [-5, 5]$ and fixed the lower

scale limit to 10 samples and the upper to 600, the first being the minimum required to avoid arithmetic

underflow (Gierałtowski et al., 2012), and the second to not include scales above the length of a single

continuous trial. We computed MMA along 12 scales, comprising the range $s \in [10, 600]$ ms which is

equivalent to ~1.67-100 Hz. The first scale integrated the scales $s_1 \in [10, 50]$ (20-100 Hz) and for

$s_{2,3,\dots,12}$ this window was progressively slid 10 ms and expanded ($s_2 \in [20, 100]$, $s_3 \in [30, 150]$ and so

forth). This permitted a nearly continuous coverage of the whole spectrum, allowing to identify any

crossover areas. For the detrending, we used a polynomial of order 2.

The values of $h(q,s)$ are interpreted in the following way (Gierałtowski et al., 2012): if $h(q,s) = 0.5$

the signal is constituted by uncorrelated randomness (white noise), $h(q,s) \in]0.5, 1]$ indicates

persistent long-range correlations and scale-free properties, if $h(q,s) \in [0, 0.5[$ the signal has anti-

correlations, $h(q,s) = 1.5$ indicates Brownian motion (integrated white noise) and, finally,

$h(q,s) > 2$ indicates black noise. Further, monofractal signals will have identical h for all q values

while multifractal time series display different exponent values depending on whether q is negative

(short fluctuations) or positive (large fluctuations) (Kantelhardt et al., 2002). Within the regime of

persistent long-range correlations, there is also a straightforward correspondence between *h* and the

spectral exponent β obtained from the slope of the power spectral density ($S(f) \sim \dfrac{1}{\beta}$, where *f* is the

frequency); according to the Wiener-Khintchine theorem: $\beta = 2h - 1$. A full description of MMA is

available at (Gierałtowski et al., 2012) and we used the original code available at Physionet,

(https://physionet.org/physiotools/mma/; (Goldberger et al., 2000)).

Multifractal spectrum. A complementary way of identifying the scaling properties of the different PCSs is to consider their singularities characterized by the parameters α (Lipschitz-Hölder exponent) and $f(\alpha)$, respectively associated with the strength of the singularities and how densely they are distributed (Halsey et al., 1986). Thus, the multifractal (or singularity) spectrum is described by the possible range of α values and the function $f(\alpha)$. There is an established connection between these latter measures and the exponents obtained with MMA through the known relationship between MF-DFA and the general multifractal formalism (Kantelhardt et al., 2002; Kantelhardt, 2011). Using fractal geometry and specifically the box-counting method, one can obtain an estimate of dimension by relating a linear distance (*s*) to the mass of a given object. If in an *E*-dimensional space of the observations is partitioned into (hyper-)cubes with side *s*, and one counts the number *N(s)* of cubes that contain at least one point of a set *S*, one obtains a very crude measure of this set (Feder, 1988) without any information from its structure. A better estimate applies a weighted sum that takes into consideration the number of points in each hyper-cube. For a set *S* consisting of *N* points, N_i will be the number of points in each partition *i* and the mass or probability $\mu_i = \dfrac{N_i}{N}$. It follows that the weighted number of boxes, N (q,s) is defined by:

$$N(q,s) = \sum i(\mu_i)^q \sim s^{-\tau(q)}$$

Where *q* is the moment order and $\tau(q)$ is the mass exponent and $\tau(0) = D_0$ is the fractal dimension of the set. If there is a linear dependency of $\tau(q)$ with *q* then, a set is monofractal, otherwise, it is multifractal. In (Kantelhardt et al., 2002), it was derived how the mass exponent relates to the *h(q)* of the method used in this report:

$$\tau(q) = qh(q) - 1$$

It follows that $f(\alpha)$ is derived from $\tau(q)$ via a Legendre transform (Halsey et al., 1986):

$$\alpha = \frac{d\tau(q)}{dq}$$

$$f(\alpha) = q\alpha - \tau(q)$$

Then by simply replacing Eq. 4 and 5 in Eq. 6, one obtains the singularities strength (α) and the dimensions of the subset of the time series that is characterized by those singularities ($f(\alpha)$):

$$\alpha = h(q) - qh'(q) \text{ and } f(\alpha) = q[\alpha - h(q)] + 1$$

The function $f(\alpha)$ is always convex upward and its left-hand branch corresponds to $q > 0$ and the
right-hand branch to $q < 0$ (Theiler, 1990). Furthermore, the central tendency (peak α_0) of the
multifractal spectrum is closely related to the Hurst exponent derived from the monofractal ($q = 2$)
DFA, thus representing the average fractal structure in the signal. The multifractal spectrum width (
$\Delta(\alpha)$) indicates the deviation from monofractality, or conversely, the degree of multifractality (Ihlen,
2012). The ($\Delta(\alpha)$) is defined as the difference between the maximum (α_{max}) and minimum (α_{min})
values of the Lipschitz-Hölder exponent:

757
758
$$\Delta\alpha = \alpha_{max} - \alpha_{min}$$
759

760 Of note, $f(\alpha)$ is not forcefully a symmetric function and can differ from the shape like the symbol ∩
761 characteristic of the most trivial multifractals, which are not strictly self-similar (scale-free), but have a
762 multiplicative rescaling structure, i.e. a scale-dependent self-similarity (Riedi, 1999). Therefore, we
763 also computed a rough estimate of the degree of asymmetry:

764
$$as_{\Delta\alpha} = \frac{\alpha_{max} - \alpha_0}{\alpha_0 - \alpha_{min}}$$
765
766

767 We computed these parameters of the multifractal spectrum for all scales (*s*) analyzed with MMA and
768 for all PCS activities of the marmosets.

769
Surrogate data. We created 50 shuffled surrogates by randomly permuting in temporal order the
samples of the original time series of each marmoset's PSCs. If the shuffling procedure yields time
series exhibiting simple random behaviour ($h = 0.5$), one can conclude that the multifractality present
is due to different long-range correlations of small and large fluctuations (Kantelhardt et al., 2002). On
the contrary, if shuffling does not affect the values of $h(q,s)$, the multifractality originates in a broad
probability density function (PDF) of the values in the time series. If the multifractality originates both
from correlations and broad PDF, the shuffling version will display weaker multifractality than the
original one. All analyses were carried out in Matlab® (v. 2018a, The MathWorks).

778
779 ***Statistics***

780
For ERP MMN (Fig 1) pairwise comparisons were used by comparing a pair of adjacent standard (i.e.
the last tone of the N train) and deviant (i.e. the first tone of the N+1 train) stimuli. Similarly, for the
spectrally-decoupled time series (Fig 2F,M,T), we performed separate repeated-measures ANOVA
(RANOVA) for each individual monkey between PSC (Broadband, Rhythmic 1, Rhythmic 2) and
stimulus (standards, deviants), *using* Bonferroni correction for *post hoc* comparisons. Similarly, in the
case of the MMA analyses (Fig 4F), Hurst exponents were compared using RANOVA for each monkey

between PSC (Broadband, Rhythmic 1 and Rhythmic 2) and *post hoc* comparisons were Bonferroni
corrected. Statistical analyses were performed using open-source statistical software *jamovi* (Version
0.9; Jamovi project, 2019).
790
791
792 **Acknowledgments**
793

794 We thank Robin Ince, Laura Imperatori and Daniel Bor for contributing to valuable discussion. This
795 manuscript is dedicated to the memory of Prof. Walter J. Freeman (1927 - 2016) whose pioneering
796 work on Neurodynamics has inspired countless meaningful insights during the execution of this
797 project.

References

Alain, C., Woods, D. L., and Knight, R. T. (1998). A distributed cortical network for auditory sensory memory in humans. Brain Research *812*, 23-37.

Alho, K. (1995). Cerebral generators of mismatch negativity (MMN) and its magnetic counterpart (MMNm) elicited by sound changes. Ear and Hearing *16*, 38-51.

Arnal, L.H., Doelling, K.B., and Poeppel, D. (2015). Delta-beta coupled oscillations underlie temporal prediction accuracy. Cerebral Cortex *25*, 3077–3085.

Bastos, A. M., Usrey, W. M., Adams, R. A., Mangun, G. R., Fries, P., and Friston, K. J. (2012). Canonical microcircuits for predictive coding. Neuron *76*, 695-711.

Beaman, C. B., Eagleman, S. L., and Dragoi, V. (2017). Sensory coding accuracy and perceptual performance are improved during the desynchronized cortical state. Nature Communications *8*: 1308. doi: 10.1038/s41467-017-01030-4

Bekinschtein, T. A., Dehaene, S., Rohaut, B., Tadel, F., Cohen, L., and Naccache, L. (2009). Neural signature of the conscious processing of auditory regularities. Proceedings of the National Academy of Sciences *106*, 1672-1677.

Brainard, D. H., and Vision, S. (1997). The psychophysics toolbox. Spatial Vision *10*, 433-436.

Buzsaki, G., Anastassiou, C.A., and Koch, C. (2012). The origin of extracellular fields and currents—EEG, ECoG, LFP and spikes. Nature Reviews Neuroscience *13*, 407–420.

Chao, Z. C., Takaura, K., Wang, L., Fujii, N., and Dehaene, S. (2018). Large-scale cortical networks for hierarchical prediction and prediction error in the primate brain. Neuron *100*, 1252-1266.

Chaudhuri, R., He, B. J., and Wang, X.-J. (2017). Random recurrent networks near criticality capture the broadband power distribution of human ECoG dynamics. Cerebral Cortex *28*, 3610–3622.

Chennu, S., Noreika, V., Gueorguiev, D., Blenkmann, A., Kochen, S., Ibáñez, A., Owen, A. M., and Bekinschtein, T. A. (2013). Expectation and attention in hierarchical auditory prediction. Journal of Neuroscience *33*, 11194-11205.

861 Chennu, S., Noreika, V., Gueorguiev, D., Shtyrov, Y., Bekinschtein, T. A., and Henson, R. (2016).
862 Silent expectations: dynamic causal modeling of cortical prediction and attention to sounds that
863 weren't. Journal of Neuroscience *36*, 8305-8316.
864
865 Choi, J. W., Lee, J. K., Ko, D., Lee, G. T., Jung, K. Y., and Kim, K. H. (2013). Fronto-temporal
866 interactions in the theta-band during auditory deviant processing. Neuroscience Letters *548*, 120-
867 125.
868
869 Clark, A. (2013). Whatever next? Predictive brains, situated agents, and the future of cognitive
870 science. Behavioral and Brain Sciences *36*, 181-204.
871
872 Cole, S.R., and Voytek, B. (2017). Brain oscillations and the importance of waveform shape. Trends
873 in Cognitive Sciences *21*, 137–149.
874
875 Cowan, N., Winkler, I., Teder, W., and Näätänen, R. (1993). Memory prerequisites of mismatch
876 negativity in the auditory event-related potential (ERP). Journal of Experimental Psychology:
877 Learning, Memory, and Cognition *19*, 909-921.
878
879 Crone, N. E., Korzeniewska, A., and Franaszczuk, P. J. (2011). Cortical gamma responses:
880 searching high and low. *Int. J. Psychophysiol. 79*, 9–15.
881
882 de Cheveigné, A., Nelken, I. (2017). Filters: When, Why, and How (Not) to Use Them. Neuron *102*,
883 280-293.
884
885 Edwards, E., Soltani, M., Deouell, L. Y., Berger, M. S., and Knight, R. T. (2005). High gamma
886 activity in response to deviant auditory stimuli recorded directly from human cortex. Journal of
887 Neurophysiology *94*, 4269-4280.
888
889 Eliades, S. J., Crone, N. E., Anderson, W. S., Ramadoss, D., Lenz, F. A., and Boatman-Reich, D.
890 (2014). Adaptation of high-gamma responses in human auditory association cortex. Journal of
891 Neurophysiology *112*, 2147-2163.
892
893 Fahrenfort, J. J., Van Driel, J., Van Gaal, S., and Olivers, C. N. (2018). From ERPs to MVPA using
894 the Amsterdam decoding and modeling toolbox (ADAM). Frontiers in Neuroscience *12*: 368. doi:
895 10.3389/fnins.2018.00368
896

897 Fayyaz, Z., Bahadorian, M., Doostmohammadi, J., Davoodnia, V., Khodadadian, S., and Lashgari,
898 R. (2019). Multifractal detrended fluctuation analysis of continuous neural time series in primate
899 visual cortex. Journal of Neuroscience Methods *312*, 84-92.
900
901 Feder, J. (1988). *Fractals.* 4th edn. New York: Springer Science & Business Media.
902
903 Fetterhoff, D., Opris, I., Simpson, S.L., Deadwyler, S.A., Hampson, R.E., and Kraft, R.A. (2015).
904 Multifractal analysis of information processing in hippocampal neural ensembles during working
905 memory under Δ_9-tetrahydrocannabinol administration. Journal of Neuroscience Methods *244*, 136–
906 153.
907
908 Freeman, W. J. (2004). Origin, structure, and role of background EEG activity. Part 1. Analytic
909 amplitude. *Clin. Neurophysiol. 115*, 2077–2088.
910
911 Freeman, W. J., Kozma, R., and Werbos, P. J. (2001). Biocomplexity: adaptive behavior in complex
912 stochastic dynamical systems. Biosystems 59, 109–123.
913
914 Freeman, W. J. and Vitiello, G. (2006). Nonlinear brain dynamics as macroscopic manifestation of
915 underlying many-body field dynamics. Physics of Life Reviews *3*, 93–118.
916
917 Freeman, W. J., and Zhai, J. (2009). Simulated power spectral density (PSD) of background
918 electrocorticogram (ECoG). Cognitive Neurodynamics *3*, 97-103.
919
920 Friston, K.J. (1997). Transients, metastability, and neuronal dynamics. Neuroimage *5*, 164–171.
921
922 Friston, K. J. (2000). The labile brain. I. Neuronal transients and nonlinear coupling. Philosophical
923 Transactions of the Royal Society of London. Series B: Biological Sciences *355*, 215-236.
924
925 Friston, K., and Kiebel, S. (2009). Predictive coding under the free-energy principle. Philosophical
926 Transactions of the Royal Society B: Biological Sciences *364*, 1211-1221.
927
928 Fuentemilla, L. L., Marco-Pallarés, J., Münte, T. F., and Grau, C. (2008). Theta EEG oscillatory
929 activity and auditory change detection. Brain Research *1220*, 93-101.
930
Garrido, M. I., Friston, K. J., Kiebel, S. J., Stephan, K. E., Baldeweg, T., and Kilner, J. M. (2008).
The functional anatomy of the MMN: a DCM study of the roving paradigm. Neuroimage *42*, 936-944.

Gasser, T., Ziegler, P., and Gattaz, W. F. (1992). The deleterious effect of ocular artefacts on the

quantitative EEG, and a remedy. European Archives of Psychiatry and Clinical Neuroscience *241*,

352-356.

Gierałtowski, J., Żebrowski, J. J., and Baranowski, R. (2012). Multiscale multifractal analysis of

heart rate variability recordings with a large number of occurrences of arrhythmia. Physical Review

E *85*: 021915. doi: 10.1103/physreve.85.021915

Gifani, P., Rabiee, H. R., Hashemi, M. H., Taslimi, P., and Ghanbari, M. (2007). Optimal fractal-

scaling analysis of human EEG dynamic for depth of anesthesia quantification. Journal of the

Franklin Institute *344*, 212-229.

Goldberger, A. L., Amaral, L. A., Glass, L., Hausdorff, J. M., Ivanov, P. C., Mark, R. G., Mietus, J.E.,

Moody, G.B., Peng, C.-K., and Stanley, H. E. (2000). PhysioBank, PhysioToolkit, and PhysioNet:

components of a new research resource for complex physiologic signals. *Circulation 101*, e215-

e220.

Grigolini, P., Aquino, G., Bologna, M., Luković, M., and West, B. J. (2009). A theory of 1/f noise in

human cognition. Physica A: Statistical Mechanics and its Applications *388*, 4192-4204.

Haenschel, C., Baldeweg, T., Croft, R. J., Whittington, M., and Gruzelier, J. (2000). Gamma and

beta frequency oscillations in response to novel auditory stimuli: a comparison of human

electroencephalogram (EEG) data with in vitro models. Proceedings of the National Academy of

Sciences *97*, 7645-7650.

Haenschel, C., Vernon, D. J., Dwivedi, P., Gruzelier, J. H., and Baldeweg, T. (2005). Event-related

brain potential correlates of human auditory sensory memory-trace formation. Journal of

Neuroscience *25*, 10494-10501.

Halgren, E., Baudena, P., Clarke, J. M., Heit, G., Liégeois, C., Chauvel, P., and Musolino, A. (1995).

Intracerebral potentials to rare target and distractor auditory and visual stimuli. I. Superior temporal

plane and parietal lobe. Electroencephalography and Clinical Neurophysiology *94*, 191-220.

Halsey, T. C., Jensen, M. H., Kadanoff, L. P., Procaccia, I., and Shraiman, B. I. (1986). Fractal

measures and their singularities: The characterization of strange sets. Physical Review A *33*: 1141.

doi: 10.1103/PhysRevA.33.1141.

He, B. J., Zempel, J. M., Snyder, A. Z., and Raichle, M. E. (2010). The temporal structures and
functional significance of scale-free brain activity. Neuron *66*, 353-369.

He, B. J. (2014). Scale-free brain activity: past, present, and future. *Trends in Cognitive Sciences*
18, 480-487.

Hermes, D., Miller, K. J., Wandell, B. A., and Winawer, J. (2014). Stimulus dependence of gamma
oscillations in human visual cortex. Cerebral Cortex *25*, 2951-2959.

Hermes, D., Nguyen, M., and Winawer, J. (2017). Neuronal synchrony and the relation between the
blood-oxygen-level dependent response and the local field potential. PLoS Biology *15*: e2001461.
doi: 10.1371/journal.pbio.2001461

Hermes, D., Petridou, N., Kay, K., and Winawer, J. (2019). An image-computable model for the
stimulus selectivity of gamma oscillations. bioRxiv *583567*. doi: 10.1101/583567

Hikishima, K., Quallo, M. M., Komaki, Y., Yamada, M., Kawai, K., Momoshima, S., Okano, H.J.,
Sasaki, E., Tamaoki, N., Lemon, R.N., Iriki, A., and Okano, H. (2011). Population-averaged
standard template brain atlas for the common marmoset (Callithrix jacchus). Neuroimage *54*, 2741-
2749.

Hsiao, F. J., Wu, Z. A., Ho, L. T., and Lin, Y. Y. (2009). Theta oscillation during auditory change
detection: an MEG study. Biological Psychology *81*, 58-66.

Hu, K., Ivanov, P. C., Chen, Z., Carpena, P., and Stanley, H. E. (2001). Effect of trends on
detrended fluctuation analysis. Physical Review E *64*, 011114. doi: 10.1103/PhysRevE.64.011114

Hurst, H. E. (1951). Long-term storage capacity of reservoirs. Transactions of the American Society
of Civil Engineers *116*, 770-799.

1000

1001   Lakatos, P., Musacchia, G., O'Connel, M.N., Falchier, A.Y., Javitt, D.C., and Schroeder, C.E.
1002   (2013). The spectrotemporal filter mechanism of auditory selective attention. Neuron *77*, 750–761.

1003

1004   Lopes da Silva, F. (2013). EEG and MEG: relevance to neuroscience. Neuron *80*, 1112–1128.

1005 Ihlen, E. A. F. (2012). Introduction to multifractal detrended fluctuation analysis in Matlab. Frontiers

1006 in Physiology *3*, 141. doi: 10.3389/fphys.2012.00141

1007

1008 Jamovi project (2019). *jamovi* (Version 0.9) [Computer Software]. Retrieved from

1009 https://www.jamovi.org

1010

1011 Javitt, D. C., Schroeder, C. E., Steinschneider, M., Arezzo, J. C., and Vaughan Jr, H. G. (1992).

1012 Demonstration of mismatch negativity in the monkey. Electroencephalography and Clinical

1013 Neurophysiology *83*, 87-90.

1014

1015 Kantelhardt, J. W., Zschiegner, S. A., Koscielny-Bunde, E., Havlin, S., Bunde, A., and Stanley, H. E.

1016 (2002). Multifractal detrended fluctuation analysis of nonstationary time series. Physica A: Statistical

1017 Mechanics and its Applications *316*, 87-114.

1018

1019 Kantelhardt, J. W. (2011). Fractal and multifractal time series. In Mathematics of Complexity and

1020 Dynamical Systems (ed. R.A. Meyers), pp. 463–487. New York, NY: Springer.

1021

1022 Kelso, J.A.S. (1995). Dynamic patterns: the self-organization of brain and behavior. MIT Press,

1023 Cambridge, Mass.

1024

1025 Ko, D., Kwon, S., Lee, G. T., Im, C. H., Kim, K. H., and Jung, K. Y. (2012). Theta oscillation related

1026 to the auditory discrimination process in mismatch negativity: oddball versus control paradigm.

1027 Journal of Clinical Neurology *8*, 35-42.

1028

1029 Komatsu, M., Takaura, K., and Fujii, N. (2015). Mismatch negativity in common marmosets: Whole-

1030 cortical recordings with multi-channel electrocorticograms. Nature Scientific Reports *5*, 15006. doi:

1031 10.1038/srep15006

1032

1033 Komatsu, M., Sugano, E., Tomita, H., and Fujii, N. (2017). A chronically implantable bidirectional

1034 neural interface for non-human primates. Frontiers in Neuroscience *11*, 514. doi:

1035 10.3389/fnins.2017.00514

1036

1037 Lakatos, P., Shah, A. S., Knuth, K. H., Ulbert, I., Karmos, G., and Schroeder, C. E. (2005). An

1038 oscillatory hierarchy controlling neuronal excitability and stimulus processing in the auditory cortex.

1039 Journal of Neurophysiology *94*, 1904-1911.

1040

1041   Lumaca, M., Trusbak Haumann, N., Brattico, E., Grube, M., and Vuust, P. (2019). Weighting of
1042   neural prediction error by rhythmic complexity: A predictive coding account using mismatch
1043   negativity. European Journal of Neuroscience *49*, 1597-1609.

1044

1045   Ma, Q., Ning, X., Wang, J., and Bian, C. (2006). A new measure to characterize multifractality of
1046   sleep electroencephalogram. Chinese Science Bulletin *51*, 3059-3064.

1047

1048   MacLean, S. E., and Ward, L. M. (2014). Temporo-frontal phase synchronization supports
1049   hierarchical network for mismatch negativity. Clinical Neurophysiology *125*, 1604-1617.

1050

1051   Mandelbrot, B. B. (2003). Multifractal power law distributions: Negative and critical dimensions and
1052   other "anomalies," explained by a simple example. Journal of Statistical Physics *110*, 739-774.

1053

1054   Manning, J. R., Jacobs, J., Fried, I., and Kahana, M. J. (2009). Broadband shifts in local field
1055   potential power spectra are correlated with single-neuron spiking in humans. Journal of
1056   Neuroscience *29*, 13613-13620.

1057

1058   Marshall, L., Mölle, M., and Bartsch, P. (1996). Event-related gamma band activity during passive
1059   and active oddball tasks. Neuroreport *7*, 1517-1520.

1060

1061   Miller, K. J., Leuthardt, E. C., Schalk, G., Rao, R. P., Anderson, N. R., Moran, D. W., Miller, J.W.,
1062   and Ojemann, J. G. (2007). Spectral changes in cortical surface potentials during motor movement.
1063   Journal of Neuroscience *27*, 2424-2432.

1064

1065   Miller, K. J., Sorensen, L. B., Ojemann, J. G., and Den Nijs, M. (2009a). Power-law scaling in the
1066   brain surface electric potential. PloS Computational Biology *5*, e1000609. doi:
1067   10.1371/journal.pcbi.1000609

1068

1069   Miller, K. J., Zanos, S., Fetz, E. E., Den Nijs, M., and Ojemann, J. G. (2009b). Decoupling the
1070   cortical power spectrum reveals real-time representation of individual finger movements in humans.
1071   Journal of Neuroscience *29*, 3132-3137.

1072

1073   Miller, K. J. (2010). Broadband spectral change: evidence for a macroscale correlate of population
1074   firing rate? Journal of Neuroscience *30*, 6477-6479.

1075

1076   Miller, K. J., Hermes, D., Pestilli, F., Wig, G. S., and Ojemann, J. G. (2017). Face percept formation
1077   in human ventral temporal cortex. Journal of Neurophysiology *118*, 2614-2627.

1078   Mukamel, R., Gelbard, H., Arieli, A., Hasson, U., Fried, I., and Malach, R. (2005). Coupling between
1079   neuronal firing, field potentials, and FMRI in human auditory cortex. Science *309*, 951-954.
1080

1081   Nagasaka, Y., Shimoda, K., and Fujii, N. (2011). Multidimensional recording (MDR) and data
1082   sharing: an ecological open research and educational platform for neuroscience. *PloS One*, *6*,
1083   e22561. doi: 10.1371/journal.pone.0022561
1084

1085   Nagy, Z., Mukli, P., Herman, P., and Eke, A. (2017). Decomposing multifractal crossovers. Frontiers
1086   in Physiology *8*, 533. doi: 10.3389/fphys.2017.00533
1087

1088   Näätänen, R., Gaillard, A. W., and Mäntysalo, S. (1978). Early selective-attention effect on evoked
1089   potential reinterpreted. Acta Psychologica *42*, 313-329.
1090

1091   Näätänen, R., Paavilainen, P., Rinne, T., and Alho, K. (2007). The mismatch negativity (MMN) in
1092   basic research of central auditory processing: a review. Clinical Neurophysiology *118*, 2544-2590.
1093

1094   Nieto-Diego, J., and Malmierca, M. S. (2016). Topographic distribution of stimulus-specific
1095   adaptation across auditory cortical fields in the anesthetized rat. PLoS Biology *14*, e1002397. doi:
1096   10.1371/journal.pbio.1002397
1097

1098   Noreika, V., Canales-Johnson, A., Johnson, A., Arnatkevičiūtė, A., Koh, J., Chennu, S., and
1099   Bekinschtein, T. A. (2019a). Wakefulness state modulates conscious access: Suppression of
1100   auditory detection in the transition to sleep. bioRxiv *155705*. doi: 10.1101/155705
1101

1102   Noreika, V., Kamke, M. R., Canales-Johnson, A., Chennu, S., Bekinschtein, T. A., and Mattingley, J.
1103   B. (2019b). Alertness fluctuations during task performance modulate cortical evoked responses to
1104   transcranial magnetic stimulation. bioRxiv *155754*. doi: 10.1101/155754
1105

1106   Paluš, M. (1996). Nonlinearity in normal human EEG: cycles, temporal asymmetry, nonstationarity
1107   and randomness, not chaos. Biological Cybernetics *75*, 389-396.
1108

1109   Papo, D. (2014). Functional significance of complex fluctuations in brain activity: from resting state
1110   to cognitive neuroscience. Frontiers in Systems Neuroscience *8*, 112. doi:
1111   10.3389/fnsys.2014.00112
1112

1113   Parras, G. G., Nieto-Diego, J., Carbajal, G. V., Valdés-Baizabal, C., Escera, C., and Malmierca, M.
1114   S. (2017). Neurons along the auditory pathway exhibit a hierarchical organization of prediction error.
1115   Nature Communications *8*, 2148. doi: 10.1038/s41467-017-02038-6
1116
1117   Pelli, D. G. (1997). The VideoToolbox software for visual psychophysics: Transforming numbers into
1118   movies. Spatial Vision *10*, 437-442.
1119
1120   Peng, C. K., Havlin, S., Stanley, H. E., and Goldberger, A. L. (1995). Quantification of scaling
1121   exponents and crossover phenomena in nonstationary heartbeat time series. Chaos: An
1122   Interdisciplinary Journal of Nonlinear Science *5*, 82-87.
1123
1124   Pérez- González, D., Malmierca, M. S., and Covey, E. (2005). Novelty detector neurons in the
1125   mammalian auditory midbrain. European Journal of Neuroscience *22*, 2879-2885.
1126
1127   Rasch, M. J., Gretton, A., Murayama, Y., Maass, W., and Logothetis, N. K. (2008). Inferring spike
1128   trains from local field potentials. Journal of Neurophysiology *99*, 1461-1476.
1129
1130   Renart, A., De La Rocha, J., Bartho, P., Hollender, L., Parga, N., Reyes, A., and Harris, K. D.
1131   (2010). The asynchronous state in cortical circuits. Science *327*, 587-590.
1132
1133   Riedi, R. H. (1999) Multifractal processes. In Long Range Dependence: Theory and Applications
1134   (Eds. P. Doukhan, M. S. Taqqu, and G. Oppenheim), pp 625-715. Birkhäuser.
1135
1136   Rorden, C., Karnath, H. O., and Bonilha, L. (2007). Improving lesion-symptom mapping. Journal of
1137   Cognitive Neuroscience *19*, 1081-1088.
1138
1139   Rubin, J., Ulanovsky, N., Nelken, I., and Tishby, N. (2016). The representation of prediction error in
1140   auditory cortex. PLoS Computational Biology *12*, e1005058. doi: 10.1371/journal.pcbi.1005058
1141
1142   Schroeder, C. E., and Lakatos, P. (2009). The gamma oscillation: master or slave? Brain
1143   Topography *22*, 24-26.
1144
1145   Schroeder, C.E., and Lakatos, P. (2009). Low-frequency neuronal oscillations as instruments of
1146   sensory selection. Trends in Neurosciences *32*, 9–18.
1147

Solomon, S. G., and Kohn, A. (2014). Moving sensory adaptation beyond suppressive effects in single neurons. Current Biology *24*, R1012-R1022.

Theiler, J. (1990). Estimating fractal dimension. Journal of the Optical Society of America A 7, 1055-1073.

Tognoli, E., and Kelso, J.A.S. (2014). The metastable brain. Neuron *81*, 35–48.

Ulanovsky, N., Las, L., and Nelken, I. (2003). Processing of low-probability sounds by cortical neurons. Nature Neuroscience *6*, 391-398.

Ulanovsky, N., Las, L., Farkas, D., and Nelken, I. (2004). Multiple time scales of adaptation in auditory cortex neurons. Journal of Neuroscience *24*, 10440-10453.

van de Velde, M., van Erp, G., and Cluitmans, P. J. (1998). Detection of muscle artefact in the normal human awake EEG. Electroencephalography and Clinical Neurophysiology *107*, 149-158.

Weiss, B., Clemens, Z., Bódizs, R., Vágó, Z., and Halász, P. (2009). Spatio-temporal analysis of monofractal and multifractal properties of the human sleep EEG. Journal of Neuroscience Methods *185*, 116-124.

Whitmire, C. J. and Stanley, G. B. (2016). Rapid sensory adaptation redux: A circuit perspective. Neuron 92, 298–315.

Zerlaut, Y. and Destexhe, A. (2017. Enhanced responsiveness and low-level awareness in stochastic network states. Neuron *94*, 1002–1009.