Anthropogenic impact on lake bacterial communities across Eastern Canada

S.A. Kraemer[1*], N. Barbosa da Costa[2], B.J. Shapiro[2], Y. Huot[3], D. Walsh[1]

[1]Concordia University, Department of Biology, Canada
[2]Université de Montréal, Department of Biology, Canada
[3]Université de Sherbrooke, Department of Applied Geomatics, Canada

Authorship contributions: SAK conducted the sample processing, analyses and wrote the first draft. NBC and BJS contributed to the molecular processing, analyses and manuscript preparation. YH and DW designed the experiment and contributed to analyses and manuscript preparation. All authors contributed to and approved the final version of the manuscript.

Competing interests: The authors declare no competing interests

Running title: Anthropogenic impact structures lake communities

Keywords: Anthropogenic impact, bacterial communities, community assembly, networks, indicator species

Number of words in abstract: 145
Number of words in main text: 4972
Number of references: 61
Number of figures, tables and text boxes: 5

Contact: Susanne Kraemer;
Concordia University, Department of Biology, L-GE 300.03,
7141 Sherbrooke W., Montreal, QC, H4B 1R6, Canada,
(+1)-514-848-2424 ext. 5848,
Susanne.kraemer@mail.concordia.ca

Abstract

Lakes play a pivotal role in ecological and biogeochemical processes and have been described as 'sentinels' of environmental change. Assessing 'lake health' across large geographic scales is critical to predict the stability of their ecosystem services and their vulnerability to anthropogenic disturbances. Here, we utilized surface water samples from 220 lakes located across Eastern Canada to assess the land use impact within watersheds on bacterial community composition using 16S rRNA gene sequencing. Alteration of communities was found on every level examined. Specifically, increasing anthropogenic impact within the watershed lowered richness mediated by changes in chloride and nitrogen concentrations. Likewise, community composition was significantly altered by differential land use within a watershed, resulting in decreased interaction network complexity and fewer keystone taxa in highly impacted lakes. Together, these findings point to vast bacterial community changes of largely unknown consequences induced by human activity within lake watersheds.

Introduction

Even though lakes only cover a small fraction of the Earth's surface, lake ecosystems have garnered a large amount of attention from researchers and policy makers in recent years as 'sentinels' of climate change and human impacts (Adrian *et al.* 2009) owing to their central importance in biogeochemical cycles (Tranvik *et al.* 2009). This pivotal role of lakes arises from their central position within watersheds from which they receive nutrients and environmental contaminants. Consequently, many anthropogenic activities in the watershed threaten the ecosystem and freshwater services provided by lakes. For example, intense agricultural activity, and specifically the use of fertilizers within watersheds, has been connected to lake eutrophication (Arbuckle & Downing 2001; Taranu & Gregory-Eaves 2008), in conjunction with the bloom of certain species of cyanobacteria (Heisler *et al.* 2008), decreases in oxygen conditions (Scavia *et al.* 2014), and increased methane emissions (Bastviken *et al.* 2004), while watershed urban development has been connected to overall water degradation caused by road salt runoff (Novotny *et al.* 2008; Dugan *et al.* 2017) and phosphorus export (Hobbie *et al.* 2017).

While the complex biogeochemical processes occurring within lakes have traditionally been treated as occurring within a black box, developments in high-throughput sequencing and analytical tools have linked ecosystem-level responses to the bacterial communities performing many of the cycling and mineralization reactions (Shade *et al.* 2007; Kara *et al.* 2013). As bacterial communities shape lake biogeochemical cycles and thus the flow of energy and nutrients within and between lakes, downstream waters, atmosphere and land, these communities will in turn be impacted by substrates and products of cycling reactions, as well as other processes occurring within the watershed. Likewise, watershed effects may also impact bacterial communities by altering species interactions, as well as overall community properties such as resilience to disturbance and ecological redundancy (Peura *et al.* 2015), but little is known about the impact of watershed-scale environmental changes on freshwater bacterial communities (but see Sher et al. 2019).

Canada is responsible for the stewardship of nearly 1,000,000 lakes which make up 20% of the world's freshwater stocks and provide the main source of drinking water for many major Canadian cities (Environment Canada 2011; Huot *et al.* 2019). In this context, the LakePulse network is tackling a large-scale assessment of the health and susceptibility status of lakes across Canada in connection to watershed land use as determined from satellite imaging, as well as environmental variables collected in situ for each lake (Figure 1 A). As defined in Huot et al. (2019), 'health' is a qualitative term and is defined in LakePulse as the departure of the ecological state of a lake from the pristine state whether historical or contemporary; the further removed from the pristine state the less healthy a lake is. During the project's first sampling season in the summer of 2017 over 200 lakes located across Eastern Canada were sampled. Here, we utilize DNA of the water column of these lakes to connect bacterial community structure to watershed anthropogenic activities such as agriculture, forestry, pasture and urban development. Moreover, we aim to untangle the underlying proximate environmental variables that cause the community shifts observed as well as describe how anthropogenic impact changes bacterial interaction networks. Lastly, we use the scope of our dataset to investigate the relative strengths of different processes of community assembly such as drift and selection. We found that communities were markedly altered with respect to both their richness and community composition by watershed anthropogenic activity, specifically agriculture and urban development, which increased lake total nitrogen concentrations and lake salinity respectively.

Moreover, highly impacted watersheds harboured more fragmented bacterial communities with fewer keystone taxa, which could affect ecosystem function and resilience.

Methods

Lake selection

Lakes were selected across Eastern Canada as described in detail in Huot *et al.* (2019). Briefly, lakes were picked in a random sampling design with ecozone, lake size and human impact index as stratifying factors. Canadian ecozones (represented by AH: Atlantic Highlands, AM: Atlantic Maritimes, BS: Boreal Shield and MP: Mixedwood Plains in south-eastern Canada) represent regions with distinct geological, climatic and ecological features (Ecological Stratification Working Group 1996). For each lake within 1 km of a road the watershed was delineated as described in Huot *et al.* (2019). Within the watershed, each pixel was assigned a human impact value between 0 and 1 depending on their category (Urban development/Road: 1; Mines/Oil: 1, Agriculture: 1, Pasture: 0.5, Forestry (recent clear-cuts): 0.5, Natural landscapes: 0) and the average Human Impact Index (HII) for the lake calculated across the watershed. Across the ecozones of Eastern Canada, the Mines/Oil category was found to be negligible in magnitude (only five lakes total with > 1% mining within their watershed and only one lake with > 5% mining) and was thus not explicitly considered in the analyses of specific land use classes herein.

Surface water sampling

The surface water of 220 lakes (Fig. 1A) was sampled during the time of maximum summer lake stratification between July and September 2017 at the deepest point of each lake (as measured on site using a sonar). The depth of the euphotic zone was determined as twice the Secchi disk depth and an integrated epilimnion sampler (sampling tube) was used to sample to that depth, or to a depth of 2 m, whichever was shallower. Water samples were immediately transferred and stored in the cold and dark, subsequently filtered through a Durapore 0.22 µm membrane filter (Sigma-Aldrich, St. Louis, USA) on site and immediately frozen at -80°C until analysis. In addition, for each lake we used the same water sample to determine dissolved organic and inorganic carbon (DOC and DIC (mg/L), supplemental material), total nitrogen (TN (mg/L), (Patton & Kryskalla 2003)), concentrations of the ions magnesium, potassium, sodium (U.S. Environmental Protection Agency 1994), chloride and sulfate (U.S. Environmental Protection Agency 1997) (all in mg/L). The average concentration of dissolved oxygen (DO, mg/mL) over the same sampling depth were measured by averaging the values from a multiparameter probe profile (RBR Maestro[3] profile, RBR Ltd., Ottawa, Canada; with a fast oxygen probe Rinko III, JFE Advantech Co., Nishinomiya, Japan) over that depth.

Processing of environmental and spatial data

Distances between the lakes (km) were used to calculate Moran's eigenvector maps (MEMs, *adespatial* R package; Dray *et al.* 2016). We visually inspected the MEM eigenvalue distribution and chose the first six MEMs to describe spatial structure within our system. We extracted the six MEMs' coordinates and combined them with data on the lake area, lake depth, DOC, DIC, TN, DO, magnesium, potassium, sodium, chloride and sulfate. Lakes with missing data for any of these variables were removed and a PCA (after scaling of the environmental variables) performed for the remaining 167 lakes (*rda* function, *vegan* R package, Oksanen *et al.* 2016). The first eight principle component axes (PCs) were chosen for further analysis as they

represented more variation than the mean. Factor loadings for these PCs are shown in Supplemental Table 1 and we considered factor loadings >=|0.43| to be significant (Hair *et al.* 1998). Ecozone, HII and size class membership, as well as environmental data for each lake can be found in Supplemental Table 2.

DNA extraction and sequencing

DNA was extracted from filters with PowerWater kits (Mobio Technologies Inc., Vancouver, Canada) and the V4 region of the 16S rRNA gene PCR-amplified using the standard primers U515_F and E786_R before sequencing on an Illumina MiSeq machine. DNA extraction and sequencing succeeded for 212 of the 220 samples. Details of PCR and sequencing reactions can be found in the supplemental material.

Sequence data processing

Read files were demultiplexed using idemp (Wu 2014) and Illumina adapters removed with cutadapt (Martin 2011). Reads were processed using the *DADA2* package in R (Callahan *et al.* 2016): Reads were trimmed, merged, chimeras removed and taxonomy of Amplicon Sequence Variants (ASVs) assigned to the genus level using the Fresh Train taxonomic database (Rohwer *et al.* 2018) and the SILVA database (version v128align) (Quast *et al.* 2013) to obtain an ASV table. Further analysis of the ASV table was done using the *phyloseq* package in R (McMurdie & Holmes 2013). ASVs from chloroplasts were removed before rarifying the data to a common read depth of 15000 reads after removing two lakes with less than 15000 reads. A phylogeny of the remaining 11510 ASVs was created using FastTree (Price *et al.* 2010). Bacterial richness of each lake was calculated using the Shannon index as implemented within the *phyloseq* package.

Linear modeling of richness and community composition analyses

We modeled the impact of the Human Impact Index (HII) variable and of percent land use types in the watershed on bacterial richness using both generalized least squares (GLS models; *nlme* R package Pinheiro *et al.* 2019)) and linear mixed models (LMM; *lme4* package (Bates *et al.* 2015)). Linear mixed models additionally contained ecozone as a random effect, but either fitted worse than the corresponding GLS model or failed to converge.

We performed structural equation modeling (*lavaan* R package (Rosseel 2012)). We considered the impact of HII onto the first eight environmental PCs, as well as the impact of the PCs and HII directly onto ranked richness. Additionally, we modeled the impact of land use classes on the environmental PCs directly (supplemental material). Models were only considered to fit better than the null model if the chi-square *p*-value was > 0.05 and the comparative fit index > 0.95.

We utilized a PERMANOVA test to investigate the impact of the environmental PCs onto community composition (*adonis* function; *vegan* R package (Oksanen *et al.* 2016)). In addition, we performed forward selection db-RDA to determine how MEMs and watershed land use impact community composition (*capscale* function; *vegan* R package (Oksanen *et al.* 2016)).

Networks of lake bacterial communities

Lakes were assigned to either a low (HII 0-0.1, 91 lakes), medium (HII>0.1&≤0.5, 96 lakes), or high (HII>0.5, 23 lakes) class. Within each network dataset, we excluded ASVs that were not present in at least 10 % of the samples, resulting in 826 ASVs in low-HII lakes, 775

ASVs in medium-HII, and 764 ASVs in high-HII lakes. We followed a combinatorial approach to determine significant connections between ASVs using both the Maximum Information Criterion (Albanese *et al.* 2013) and Sparse Inverse Covariance Estimation (Kurtz *et al.* 2015) as detailed in the Supplemental material.

We determined Order influence within high, moderate and low-impact networks by dividing the total number of edges of each Order by the number of nodes belonging to the Order (excluding Orders only represented by a single ASV or absent in any of the networks) (Banerjee *et al.* 2019). Moreover, we dropped all ASVs that could not be identified to the Order level.

Indicator species of lake health

We investigated potential indicator species of high and low impact lakes based on abundance changes of taxa in pairs of lakes that were geographically similar but varied with respect to their HII. The log-ratio of ASV abundance change between paired lake communities was compared to control lake pairs. Additionally, we screened all ASVs for rank changes of at least 33% between paired impacted and pristine lakes as described in the supplemental material.

Community assembly processes of lake bacterial communities

We defined generalist and specialist taxa within our dataset as follows: taxa with an occupancy $\geq$ 158 lakes (75% of the data set) were considered generalists, while ASVs in $\leq$ ten lakes (5%) and with a relative abundance of higher than 2% were considered specialists (Barberán *et al.* 2012). We investigated distance-decay curves between community dissimilarity (determined as the Bray-Curtis dissimilarity as calculated using the *vegdist* function in vegan (Oksanen *et al.* 2016), and the distance between the lakes (in km), using Mantel tests with 9999 permutations.

We followed the null model method as proposed by Stegen et al. (2013) to determine quantitative estimates for the strength of selection, dispersal limitation and ecological drift in the lake communities on a subsetted dataset in which rare taxa (abundance <500) were removed (679 ASVs remaining) as detailed in the supplemental material.

Results

Land use impact on bacterial richness, community composition and interactions

We firstly investigated the interaction between the human impact index (HII) and bacterial richness (here measured as the Shannon-Weaver index) of lake communities. Overall, we detected a significantly negative impact of HII on richness (slope: -0.31, $p<0.05$, Fig. 1B). When modeling richness as a function of different land use classes, we found the percentage of forestry within a watershed to have a significantly positive impact (slope: 1.26, $p<0.01$), while the percentage of urban development had a marginally significant negative impact (slope: -0.67, $p=0.057$).

To further investigate the relationship between HII and richness we performed structural equation modeling (SEM) to investigate the relationship between HII and the first eight environmental PCs and in turn the impact of the PCs onto ranked richness (Fig 2). The SEM ($\chi^2=2.81$, DF=28, $p=1.00$, CFI=1.00), showed a significant negative impact of PC1 on richness ($p=0.048$). PC1 carried significant positive loadings for the cations Ca^{2+}, K^+ and Mg^{2+}. We also found a significant negative impact on richness of PC7, which carried a significant loading for SO_4^{2-} ($p=0.019$), as well as a significant positive impact of PC3 ($p<0.001$), which carried a

significant negative loading for lake depth. In addition, we detected marginally significant impacts of the PC 2 (p=0.076), which was positively loaded with chloride and total nitrogen concentrations and PC6 (p=0.069) which carried loadings for MEM6 and DO. The HII variable also directly negatively impacted ranked richness (p=0.036) and strongly positively impacted PC2 (p>0.001). Thus, HII was associated with increased lake chloride and nitrogen concentrations, which in turn led to a decrease in bacterial richness (Fig. 2). Moreover, richness was impacted or correlated with cation concentrations, lake depth, spatial factors and available oxygen. Lastly, HII also negatively impacted richness independent of the environmental PCs considered here, suggesting unknown environmental variables.

To understand the cause of the impact of the HII variable, we examined the underlying land use classes and investigated their relationship with ranked richness and the eight environmental PCs in an additional SEM (Supplemental material, Supp. Fig. 1). We found PC2 to be significantly positively associated with agriculture, urban development and pasture within the watershed.

We investigated factors structuring community composition via PERMANOVA with the first eight environmental PCs as explanatory variables. All PCs except PC7 and PC8 were found to significantly impact community composition (all p<0.05). PC2, carrying chloride and nitrogen concentration loadings, showed the highest R^2 value in the model (0.06), but all factors included in the model only explained a small amount of the total variation (~9%).

We utilized db-RDA to select watershed-scale variables impacting community composition. A full model containing the spatial eigenvectors, lake morphometric parameters (lake depth and area) and land use data was found to be significant (p=0.001) so we proceeded with forward selection. Overall, a model containing the land use classes: natural landscapes, forestry, agriculture, pasture and urban development in the watershed, as well as the first four MEMs, and lake depth was selected as the best model (Fig 3). However, the model only explained ~15% of the variation observed.

To determine how anthropogenic impact may alter the structure of bacterial interactions, we constructed networks of high, moderate, and low HII lake communities. Low HII lake communities showed significant co-occurrence patterns for 179 nodes (taxa) with a total of 193 edges (co-occurrence connections). Moderate impact communities had significant co-occurrence networks for 145 nodes and 163 edges, while high impact communities had 220 nodes and 174 edges. Networks were more fragmented under high human impact (low: 24 components, moderate: 13 components, high: 59 components) and the clustering coefficient as well as the centralization value decreased with increasing HII (Fig. 4). Both low and moderate impact lake communities were characterized by ASVs having similar average numbers of neighbors (low: 2.16, medium: 2.25) and keystone taxa (with over five connections each: low: 13, moderate: 11), whereas ASVs in highly impacted lake communities had on average fewer neighbours (1.58). Lastly, highly impacted lakes only had two keystone taxa (taxa with >5 edges, Fig. 4).

We determined whether specific bacterial Orders varied in their influence in high, moderate or low HII lakes by dividing their number of total edges by the number of nodes of a given Order. We identified 17 Orders which were represented in all three sets of networks (Supp. Table 3). Four of these Orders changed consistently between low, moderate and high lake networks. Burkholderiales ASVs, Frankiales ASVs and Rhizobiales ASVs had decreasing influence from low to moderate to high impacted networks, while verrucomicrobial OPB35'ASVs' influence increased. Furthermore, Acidimicrobiales and Planctomycetales ASVs

were of relatively low influence in high impact networks, but of higher influence in moderate and low impacted networks.

We investigated potential indicator species, i.e. species that can be used as ecological indicators for specific environmental conditions due to their niche preferences, for high or low impact lakes using log-ratio and rank abundance changes of ASVs. Overall, no strong indicator patterns emerged, demonstrating the high variability of freshwater communities under both pristine and impacted conditions. Log-ratio changes in treatment (pristine-impacted), but not control (pristine-pristine or impacted-impacted), lake pairs were detected for six ASVs in AH, two in AM, nine in BS and ten in MP. In total, 26 ASVs showed significant log-ratio changes, but these ASVs rarely overlapped between ecozones. Taking into account ASV taxonomy, we found ASVs belonging to the Burkholderiales *betI-A* clade and of the cyanobacterial *Family I* clade to be associated with low impact lakes across three and two ecozones, respectively (Supp. Table 4).

95 ASVs were found to show significant rank abundance changes between high and low impact lakes, but not between control lake pairs (17 ASVs in AH, 19 in AM, 46 in BS, 25 in MP, Supp. Table. 5). Specific taxa were associated with changing lake conditions across ecozones. For example, the Burkholderiales *betI-A* clade (represented by six ASVs) was associated with low impact lakes in all four ecozones, except for one ASV, which was a high impact indicator in AH.

Evolutionary and ecological processes shaping lake bacterial communities

We did not detect any low-occupancy, high-abundance specialists within our dataset. In contrast, we identified 25 generalists, which were present in 75% or more of lakes. The most common generalist lineage was the actinobacterial *acI* lineage (9 ASVs), followed by the betaproteobacterial *betI* lineage (5 ASVs). Other lineages represented included *acIV* (2 ASVs), *betIV* (2 ASVs) and bacteroidal *bacI* (2 ASVs) (Supp. Table 6).

Overall, we found bacterial communities to be less similar the further apart in space they were sampled (Mantel test, $p<0.001$). Specifically, across all communities investigated, lakes sampled up to approximately 350 km apart were significantly more similar than expected by chance (Mantel correlogram analysis, $p=0.004$, Fig. 5A). To determine the causes for this structure, we performed a quantitative analysis of the community assembly processes shaping lake communities to estimate the relative strength of selection and drift processes in our dataset. Firstly, we determined whether we could detect a phylogenetic signal within our data (i.e. whether closely related species had similar niches) by testing for correlation of phylogenetic distance, based on part of the 16S rRNA gene, with ecological distance (differences in abundance patterns expressed as Bray-Curtis dissimilarity). Overall, we found a significant decay of phylogenetic distance with ecological distance (Mantel test: $p<0.05$, Fig. 5B). Specifically, and as expected, we detected a phylogenetic signal over relatively short phylogenetic distances (cophenetic distance>0.365), indicating that the beta nearest taxon index (βNDI), which indicates the impact of selection on community composition, is an appropriate metric in our dataset to measure phylogenetic turnover (Stegen *et al.* 2013).

Within the 21945 community comparisons possible with our data, we found selection to be responsible for 12.3% of the observed community turnover (βNTI>|2|). Most selection increases community similarity (homogeneous selection βNTI<-2: 9.6%), while only 2.7% of community turnover was consistent with heterogeneous selection (βNTI>2). Of the remaining 87.7% variation, 16% was governed by dispersal limitation (βNTI<|2| and $RC_{Bray}>0.95$), whereas

39.1% was attributable to homogenizing dispersal (βNTI<|2| and RC$_{Bray}$<0.95). This leaves 32.6% of community turnover attributable to ecological drift (e.g. stochastic processes) alone (Fig. 5C). Environmental PCs associated with either selection or drift processes are described in the supplemental material.

Discussion

Our dataset, encompassing over 200 lakes located across Eastern Canada, showed that watershed anthropogenic activity strongly influences surface water bacterial community composition. Bacterial communities were altered on all levels investigated, ranging from richness, to community composition, to community interactions, to indicator species of high and low lake quality.

Human impact structures bacterial richness and community composition

The human impact index (HII) variable was found to be associated with surface water communities with significantly lower richness. Specifically, high HII values were associated with high chloride and nitrogen values, which in turn were associated with reduced richness. Salinity has been shown to be a major factor structuring bacterial diversity in a range of environments (Benlloch et al. 2002; Abed et al. 2007; Lozupone & Knight 2007), but the effect of salinity on diversity in lake systems is less clear (Wu et al. 2006; Wang et al. 2011). Importantly, these findings point to a significant impact of salt contamination on the bacterial community even at relatively low levels of salinity (and well below the chronic pollution thresholds for chloride ions at 230 mg/L) and thus suggests that bacterial ecosystem services in lakes may be highly vulnerable to salt contamination.

Total nitrogen concentration was equally implicated in lower richness. Environmental nitrogen concentrations have been mostly discussed with reference to the application of fertilizer in agriculture (Kaushal et al. 2011) and have been linked to lake eutrophication and blooms of cyanobacteria and associated community shifts (Paerl & Scott 2010; Glibert 2017). Corresponding to our findings, previous studies have also found an association between the application of nitrogen and decreased bacterial community diversity in soil (Zeng et al. 2016, but see Fierer et al. 2012), lakes (Cao et al. 2017) and sediments (Zeng et al. 2016).

PC2 was found to be impacted by agriculture, pasture and urban development within the watershed, indicating that road salt and fertilizer are likely sources for the environmental variables loaded onto the PC. High nitrogen within watersheds has previously been linked to anthropogenic activity (Van Breemen et al. 2002), and agricultural landscapes have been shown to have lower N retention rates than suburban or forested areas (Kaushal et al. 2011). Likewise, chloride concentrations in streams and lakes have been previously shown to be driven by the application of road salt (Kelly et al. 2008; Corsi et al. 2010) and even relatively low road cover within a watershed (~1%) has been linked to increased chloride concentrations within lakes (Dugan et al. 2017).

Community composition was also structured by landscape-scale variables. Lake communities were impacted by nearly every factor explored here, ranging from lake morphometry to most land use classes and spatial components. However, although most factors that we included in our models were significant, our models were only able to explain a small proportion of lake community composition overall (~9-15%). Thus, despite the inclusion of over 30 partially nested variables in the models, we acknowledge that we still fall short of describing the majority of the prokaryotic community diversity observed in this system.

Human impact weakens bacterial interaction networks

Overall, anthropogenic activity is associated with strong shifts in bacterial community composition. To relate these shifts to community stability and functionality, we investigated how communities in lakes with high, moderate or low human impacted watersheds were altered in their interactions and co-occurrences. As we were interested in overall community structure rather than specific group interactions (e.g. Eiler *et al.* 2012), we treated all interactions (positive and negative) as undirected, to facilitate analysis. Significant co-occurrence may indicate that taxa are directly interacting but may also show taxa as co-occurring due to overall niche similarity (or competitive exclusion in the case of negative co-occurrence). While lake networks constructed from communities from low and moderately impacted watersheds were similar, we detected altered structure and topology in networks from highly impacted lakes. Even though communities from highly impacted lakes had the highest number of taxa involved in significant co-occurrences (220 taxa), the number of significant connections did not scale up proportionally, resulting in, on average, less neighbours per node in these communities than in moderate or low impacted communities. In turn, this resulted in less highly connected keystone taxa and thus a more fragmented network with a lower clustering coefficient (the average number of actual three-node connections passing through each node versus the potential number of three-node connections) and lower network centralization. Higher levels of centralization (i.e. average number of nodes connected to each node or how 'star-shaped' a network structure is) have been linked to increased system stability in root microbiomes (Banerjee *et al.* 2019) and lake systems (Peura *et al.* 2015).

Interestingly, moderate HII communities seemed to be only marginally impacted with respect to their co-occurrence network in comparison with low impact communities. This finding may indicate a buffering effect in which microbiomes are able to maintain most ecosystem functions despite shifts in the underlying taxa performing them, for example via functional redundancy. The altered pattern in high impacted communities may indicate that once a threshold is reached, the system's buffering capacities are exhausted, and that specifically the loss of highly connected and influential keystone taxa may lead to cascading effects of community fragmentation (Peura *et al.* 2015). Alternatively, the patterns could indicate that while low and moderately impacted lakes are quite similar to each other, highly impacted lakes may be overall more variable or that, depending on the exact nature of the anthropogenic impact, multiple high impact communities may exist.

Ecological and evolutionary processes structuring lake communities

In addition to investigating the effect of watershed anthropogenic activity, the large spatial scale of the study allowed us to assess the ecological and evolutionary processes structuring lake bacterial communities in Eastern Canada in general, as has been previously done in a range of bacterial communities including aquifers (Stegen *et al.* 2013) and the ocean (Logares *et al.* 2018). Firstly, we investigated whether we could identify generalist and specialist taxa within our dataset (Barberán *et al.* 2012). We were able to identify 25 generalist taxa with an occupancy of over 75% in our dataset, including well-known cosmopolitan freshwater lineages such as the actinobacterial *acI* clade ASVs, betaproteobacterial *betI* ASVs and an alphaproteobacterial LD12 ASV. In contrast, we were unable to identify any specialist (low occupancy but relatively high abundance) ASVs, indicating that taxa have either low abundance or wider distributions.

We also utilized the dataset to investigate patterns in community assembly. In accordance with the absence of specialists, we found heterogenizing selection to be only responsible for a very small amount of community variation. Rather, lake ecosystems seemed to be extensively connected by homogenizing dispersal and successful lineages were thus able to establish themselves in other communities (homogenizing selection). Consequently, dispersal limitation was not found to be an important factor structuring lake bacterial communities and instead, ecological drift processes were prevalent. We are not aware of any comparable study of community assembly in lakes, but it was noticeable that, in our system, selection was a much less powerful force shaping communities than in subsurface or marine ecosystems (Stegen *et al.* 2013; Logares *et al.* 2018).

Caveats and conclusion

Data obtained within the first sampling season of the LakePulse project allows an unprecedented insight into how watershed anthropogenic impact shape bacterial communities on landscape scales. However, the large-scale nature of the project and the sampling effort involved also lead to several caveats. Due to technical difficulties we were unable to measure pH and phosphate for a large quantity of the lakes and were thus not able to include these variables in our models. pH has been previously shown to be a key driver of bacterial community structure in a range of environments (Lindström *et al.* 2005; Lauber *et al.* 2009; Xiong *et al.* 2012). pH is a complex variable and correlated to some degree with both the concentrations of ions such as chloride and ammonium within the water, as well as lake geography and morphometry. Likewise, phosphate concentrations have been shown to impact lake bacterial community compositions (Tong *et al.* 2005; Romina Schiaffino *et al.* 2011) and sediments (Zeng *et al.* 2009) but were not taken into account in this model. As phosphate, in combination with nitrate, is often used in agricultural settings, we expect some positive correlation between phosphate and nitrogen-species but are unable to directly investigate its role in bacterial community diversity. We suspect that at least some of the unexplained variation in our system is connected to these missing environmental variables. Likewise, we limited our analysis to only include abiotic factors as explanatory variables of bacterial community structure. However, biotic interactions are not only likely to be altered by environmental factors, as shown here via network analysis, but can also exert strong biological control mechanisms for example via anticompetitor compounds (Kraemer *et al.* 2017), predator-prey or mutualistic relationships across trophic levels. Future work within the project is aimed at incorporating bacterial diversity into a food web framework utilizing phytoplankton and zooplankton data also collected from the lakes, as well as deepening our functional understanding of lake communities using metagenomics. Moreover, current sampling efforts are underway that will allow us to include more lakes to extend our frameworks to all of Canada, including regions of extremely high human impact within watersheds and associated environmental decay due to agriculture and mining.

In conclusion, we used an extensive dataset of lake bacterial communities to model the impact of anthropogenic activity across Eastern Canada. Human impact, and specifically variables related to urban and agricultural development within watersheds, had a pronounced effect on lake communities in such a way that high human watershed impact may lead to the fragmentation of bacterial communities and ultimately the decline of the ecosystem services provided by them.

Data accessibility statement
We confirm that the data supporting the results will be made publicly available on Figshare and the data DOI included in the manuscript upon acceptance.

Acknowledgements
        We thank the coordinators and field team members of the LakePulse 2017 sampling campaign for their efforts. We also would like to thank members of the network, and specifically B. Beisner and V. Fugere for helpful discussions during the manuscript preparation.

References
Abed, R.M.M., Kohls, K. & De Beer, D. (2007). Effect of salinity changes on the bacterial diversity, photosynthesis and oxygen consumption of cyanobacterial mats from an intertidal flat of the Arabian Gulf. *Environ. Microbiol.*

Adrian, R., O'Reilly, C.M., Zagarese, H., Baines, S.B., Hessen, D.O., Keller, W., *et al.* (2009). Lakes as sentinels of climate change. *Limnol. Oceanogr.*

Albanese, D., Filosi, M., Visintainer, R., Riccadonna, S., Jurman, G. & Furlanello, C. (2013). Minerva and minepy: A C engine for the MINE suite and its R, Python and MATLAB wrappers. *Bioinformatics*.

Arbuckle, K.E. & Downing, J.A. (2001). The influence of watershed land use on lake N : P in a predominantly agricultural landscape. *Limnol. Oceanogr.*

Banerjee, S., Walder, F., Büchi, L., Meyer, M., Held, A.Y., Gattinger, A., *et al.* (2019). Agricultural intensification reduces microbial network complexity and the abundance of keystone taxa in roots. *ISME J.*

Barberán, A., Bates, S.T., Casamayor, E.O. & Fierer, N. (2012). Using network analysis to explore co-occurrence patterns in soil microbial communities. *ISME J.*

Bastviken, D., Cole, J., Pace, M. & Tranvik, L. (2004). Methane emissions from lakes: Dependence of lake characteristics, two regional assessments, and a global estimate. *Global Biogeochem. Cycles*.

Bates, D., Mächler, M., Bolker, B.M. & Walker, S.C. (2015). Fitting linear mixed-effects models using lme4. *J. Stat. Softw.*

Benlloch, S., López-López, A., Casamayor, E.O., Øvreås, L., Goddard, V., Daae, F.L., *et al.* (2002). Prokaryotic genetic diversity throughout the salinity gradient of a coastal solar saltern. *Environ. Microbiol.*

Van Breemen, N., Boyer, E.W., Goodale, C.L., Jaworski, N.A., Paustian, K., Seitzinger, S.P., *et al.* (2002). Where did all the nitrogen go? Fate of nitrogen inputs to large watersheds in the northeastern U.S.A. *Biogeochemistry*.

Callahan, B.J., McMurdie, P.J., Rosen, M.J., Han, A.W., Johnson, A.J.A. & Holmes, S.P. (2016). DADA2: High-resolution sample inference from Illumina amplicon data. *Nat. Methods*.

Cao, X., Wang, J., Liao, J., Gao, Z., Jiang, D., Sun, J., *et al.* (2017). Bacterioplankton community responses to key environmental variables in plateau freshwater lake ecosystems: A structural equation modeling and change point analysis. *Sci. Total Environ.*

Corsi, S.R., Graczyk, D.J., Geis, S.W., Booth, N.L. & Richards, K.D. (2010). A fresh look at road salt: Aquatic toxicity and water-quality impacts on local, regional, and national scales. *Environ. Sci. Technol.*

Dray, S., Blanchet, G., Borcard, D., Guenard, G., Jombart, T., Legendre, P., *et al.* (2016). Package "adespatial": Multivariate Multiscale Spatial Aanalysis.

Dugan, H.A., Bartlett, S.L., Burke, S.M., Doubek, J.P., Krivak-Tetley, F.E., Skaff, N.K., *et al.* (2017). Salting our freshwater lakes. *Proc. Natl. Acad. Sci. U. S. A.*

Ecological Stratification Working Group. (1996). *A National Ecological Framework for Canada*.

Eiler, A., Heinrich, F. & Bertilsson, S. (2012). Coherent dynamics and association networks among lake bacterioplankton taxa. *ISME J.*

Environment Canada. (2011). *2011 Municipal Water Use Report*.

Fierer, N., Lauber, C.L., Ramirez, K.S., Zaneveld, J., Bradford, M.A. & Knight, R. (2012). Comparative metagenomic, phylogenetic and physiological analyses of soil microbial communities across nitrogen gradients. *ISME J.*

Glibert, P.M. (2017). Eutrophication, harmful algae and biodiversity — Challenging paradigms in a world of complex nutrient changes. *Mar. Pollut. Bull.*

Hair, J., Tatham, R., Anderson, R. & Black, W. (1998). *Multivariate Data Analysis*. Fifth Edit. Prentice-Hall, London.

Heisler, J., Glibert, P.M., Burkholder, J.M., Anderson, D.M., Cochlan, W., Dennison, W.C., *et al.* (2008). Eutrophication and harmful algal blooms: A scientific consensus. *Harmful Algae*.

Hobbie, S.E., Finlay, J.C., Janke, B.D., Nidzgorski, D.A., Millet, D.B. & Baker, L.A. (2017). Contrasting nitrogen and phosphorus budgets in urban watersheds and implications for managing urban water pollution. *Proc. Natl. Acad. Sci.*

Huot, Y., Brown, C.A., Potvin, G., Antoniades, D., Baulch, H.M., Beisner, B.E., *et al.* (2019). The NSERC Canadian Lake Pulse Network: A national assessment of lake health providing science for water management in a changing climate. *Sci. Total Environ.*

Kara, E.L., Hanson, P.C., Hu, Y.H., Winslow, L. & McMahon, K.D. (2013). A decade of seasonal dynamics and co-occurrences within freshwater bacterioplankton communities from eutrophic Lake Mendota, WI, USA. *ISME J.*

Kaushal, S.S., Groffman, P.M., Band, L.E., Elliott, E.M., Shields, C.A. & Kendall, C. (2011). Tracking nonpoint source nitrogen pollution in human-impacted watersheds. *Environ. Sci. Technol.*

Kelly, V.R., Lovett, G.M., Weathers, K.C., Findlay, S.E.G., Strayer, D.L., Burns, D.J., *et al.* (2008). Long-term sodium chloride retention in a rural watershed: Legacy effects of road salt on streamwater concentration. *Environ. Sci. Technol.*

Kraemer, S.A., Soucy, J.P.R. & Kassen, R. (2017). Antagonistic interactions of soil pseudomonads are structured in time. *FEMS Microbiol. Ecol.*

Kurtz, Z.D., Müller, C.L., Miraldi, E.R., Littman, D.R., Blaser, M.J. & Bonneau, R.A. (2015). Sparse and Compositionally Robust Inference of Microbial Ecological Networks. *PLoS Comput. Biol.*

Lauber, C.L., Hamady, M., Knight, R. & Fierer, N. (2009). Pyrosequencing-based assessment of soil pH as a predictor of soil bacterial community structure at the continental scale. *Appl. Environ. Microbiol.*

Lindström, E.S., Kamst-Van Agterveld, M.P. & Zwart, G. (2005). Distribution of typical freshwater bacterial groups is associated with pH, temperature, and lake water retention time. *Appl. Environ. Microbiol.*

Logares, R., Tesson, S.V.M., Canback, B., Pontarp, M., Hedlund, K. & Rengefors, K. (2018). Contrasting Prevalence of Selection and Drift in the Community Structuring of Bacteria and Microbial Eukaryotes. *Environ. Microbiol.,* 20, 2231-2240.

Lozupone, C.A. & Knight, R. (2007). Global patterns in bacterial diversity. *Proc. Natl. Acad.*

Sci., 104, 11436–11440.

Marmen, S., Blank, L. Al-Ashhab, A., Malik, A., Ganzert, L. Lalzar, M., Grossart, H.-P. & Sher, D. (2019). The Role of Land Use Types and Water Chemical Properties in Structuring the Microbiome of a Connected lake System. *bioRxiv*.

Martin, M. (2011). Cutadapt removes adapter sequences from high-throughput sequencing reads. *EMBnet.journal*.

McMurdie, P.J. & Holmes, S. (2013). Phyloseq: An R Package for Reproducible Interactive Analysis and Graphics of Microbiome Census Data. *PLoS One*.

Novotny, E. V., Murphy, D. & Stefan, H.G. (2008). Increase of urban lake salinity by road deicing salt. *Sci. Total Environ.*

Oksanen, J., Blanchet, F.G., Friendly, M., Kindt, R., Legendre, P., Mcglinn, D., *et al.* (2016). Vegan: Community Ecology Package. *URL https//cran.r-project.org, https//github.com/vegandevs/vegan*.

Paerl, H.W. & Scott, J.T. (2010). Throwing fuel on the fire: Synergistic effects of excessive nitrogen inputs and global warming on harmful algal blooms. *Environ. Sci. Technol.*

Patton, C. & Kryskalla, J. (2003). Methods of Analysis by the U.S. Geological Survey National Water Quality Laboratory - Evaluation of Alakline Digestion as an Alternative to Kjedahl Digestion for Determination of Total and Dissolved Nitrogen and Phosphorous. *Water-Resources Investig.*, 03.

Peura, S., Bertilsson, S., Jones, R.I. & Eiler, A. (2015). Resistant microbial cooccurrence patterns inferred by network topology. *Appl. Environ. Microbiol.*

Pinheiro, J., Bates, D., DebRoy, S., Sarkar, D. & R Development Core Team, T. (2019). nlme: Linear and Nonlinear Mixed Effect Models. R package version 3.1-141.

Price, M.N., Dehal, P.S. & Arkin, A.P. (2010). FastTree 2 - Approximately maximum-likelihood trees for large alignments. *PLoS One*.

Quast, C., Pruesse, E., Yilmaz, P., Gerken, J., Schweer, T., Yarza, P., *et al.* (2013). The SILVA ribosomal RNA gene database project: Improved data processing and web-based tools. *Nucleic Acids Res.*

Rohwer, R.R., Hamilton, J.J., Newton, R.J. & McMahon, K.D. (2018). TaxAss: Leveraging a Custom Freshwater Database Achieves Fine-Scale Taxonomic Resolution. *mSphere*.

Romina Schiaffino, M., Unrein, F., Gasol, J.M., Massana, R., Balagué, V. & Izaguirre, I. (2011). Bacterial community structure in a latitudinal gradient of lakes: The roles of spatial versus environmental factors. *Freshw. Biol.*

Rosseel, Y. (2012). Lavaan: An R package for structural equation modeling. *J. Stat. Softw.*

Scavia, D., David Allan, J., Arend, K.K., Bartell, S., Beletsky, D., Bosch, N.S., *et al.* (2014). Assessing and addressing the re-eutrophication of Lake Erie: Central basin hypoxia. *J. Great Lakes Res.*

Shade, A., Kent, A.D., Jones, S.E., Newton, R.J., Triplett, E.W. & McMahon, K.D. (2007). Interannual dynamics and phenology of bacterial communities in a eutrophic lake. *Limnol. Oceanogr.*, 52, 487–494.

Stegen, J.C., Lin, X., Fredrickson, J.K., Chen, X., Kennedy, D.W., Murray, C.J., *et al.* (2013). Quantifying community assembly processes and identifying features that impose them. *ISME J.*

Taranu, Z.E. & Gregory-Eaves, I. (2008). Quantifying relationships among phosphorus, agriculture, and lake depth at an inter-regional scale. *Ecosystems*.

Tong, Y., Lin, G., Ke, X., Liu, F., Zhu, G., Gao, G., *et al.* (2005). Comparison of microbial

community between two shallow freshwater lakes in middle Yangtze basin, East China. *Chemosphere*.

Tranvik, L.J., Downing, J.A., Cotner, J.B., Loiselle, S.A., Striegl, R.G., Ballatore, T.J., *et al.* (2009). Lakes and reservoirs as regulators of carbon cycling and climate. *Limnol. Oceanogr.*, 54, 2298–2314.

U.S. Environmental Protection Agency. (1994). *Method 200.7: Determiantion of Metals and Trace Eelements in Water and Wastes by Inductively Coupled Plasma-Atomic Emission Spectrometry*.

U.S. Environmental Protection Agency. (1997). *Method 300.1: Determination of Inorganic Anions in Drinking Water by Ion Chromatography*.

Wang, J., Yang, D., Zhang, Y., Shen, J., van der Gast, C., Hahn, M.W., *et al.* (2011). Do Patterns of Bacterial Diversity along Salinity Gradients Differ from Those Observed for Macroorganisms? *PLoS One*.

Wu, Q.L., Zwart, G., Schauer, M., Kamst-Van Agterveld, M.P. & Hahn, M.W. (2006). Bacterioplankton community composition along a salinity gradient of sixteen high-mountain lakes located on the Tibetan Plateau, China. *Appl. Environ. Microbiol.*

Wu, Y. (2014). Barcode Demultiplex for Illumina I1, R1, R2 fastq.gz files.

Xiong, J., Liu, Y., Lin, X., Zhang, H., Zeng, J., Hou, J., *et al.* (2012). Geographic distance and pH drive bacterial distribution in alkaline lake sediments across Tibetan Plateau. *Environ. Microbiol.*

Zeng, J., Liu, X., Song, L., Lin, X., Zhang, H., Shen, C., *et al.* (2016). Nitrogen fertilization directly affects soil bacterial diversity and indirectly affects bacterial community composition. *Soil Biol. Biochem.*

Zeng, J., Yang, L., Li, J., Liang, Y., Xiao, L., Jiang, L., *et al.* (2009). Vertical distribution of bacterial community structure in the sediments of two eutrophic lakes revealed by denaturing gradient gel electrophoresis (DGGE) and multivariate analysis techniques. *World J. Microbiol. Biotechnol.*

Figure 1: A) Map of Eastern Canada showing 210 sampled lakes coloured by their Human Impact Index. B) Scatterplot showing the relationship between Shannon diversity and the Human Impact Index variable for 210 lakes.

Figure 2: Graphical representation of the structural equation model. All paths were tested, but only those showing significant ($p<0.05$, solid lines) or marginally significant ($p<0.1$, dashed lines) correlations are shown. Positive interactions and loadings are shown in green, negative in red.

Figure 3: Plot of the db-rda coordinates of lake communities. Arrows show the significant environmental variables after forward selection.

Figure 4: Co-occurrence networks and statistics of high, moderate and low impact lake communities. Nodes are colored by the number of their edges.

Figure 5: A) Distance-decay plot showing that communities were less similar (based on BC) the further apart they occur. B) Phylogenetic distance decay plot showing that communities were

less similar with greater phylogenetic distance. C) Relative contributions of community assembly processes across the sampled lakes.

Low
HII: 0-0.1

Centralization: 0.056
Clustering coef.: 0.119
Av. # neighbors: 2.16
24 fragments
13 keystone taxa

Moderate
HII: 0.1-0.5

Centralization: 0.041
Clustering coef.: 0.09
Av. # neighbors: 2.25
13 fragments
11 keystone taxa

High
HII: > 0.5

Centralization: 0.03
Clustering coef.: 0.031
Av. # neighbors: 1.58
59 fragments
2 keystone taxa

A

B

C