PTP-MEG2 regulates quantal size and fusion pore opening through two distinct structural bases and substrates

Yun-Fei Xu*[#1], Xu Chen[#2], Zhao Yang[1], Peng Xiao[1], Chun-Hua Liu[3], Kang-Shuai Li[1], Xiao-Zhen Yang[4], Yi-Jing Wang[1], Zhong-Liang Zhu[5], Zhi-Gang Xu[6], Sheng Zhang[7], Chuan Wang[8], Wei-Dong Zhao[9], Chang-He Wang[10], Zhi-Liang Ji[4], Zhong-Yin Zhang[7], Min Cui*[2], Jin-Peng Sun*[1] and Xiao Yu*[&2]

Affiliations:

[1]Key Laboratory Experimental Teratology of the Ministry of Education and Department of Biochemistry and Molecular Biology, Shandong University School of Medicine, 44 Wenhua Xi Road, Jinan, Shandong, 250012, China

[2]Key Laboratory Experimental Teratology of the Ministry of Education and Department of Physiology, Shandong University School of Medicine, 44 Wenhua Xi Road, Jinan, Shandong, 250012, China

[3]Department of Physiology, Taishan Medical University, Taian, Shandong 271000, China.

[4]State Key Laboratory of Cellular Stress Biology, School of Life Sciences, Xiamen University, Xiamen 361102, Fujian, China

[5]School of Life Sciences, University of Science and Technology of China, 96 Jinzhai Road, Hefei, Anhui 230026, China

[6]Shandong Provincial Key Laboratory of Animal Cells and Developmental Biology, School of Life Sciences, Shandong University, Jinan, China

[7]Departments of Medicinal Chemistry and Molecular Pharmacology and of Chemistry, Center for Cancer Research, and Institute for Drug Discovery, Purdue University, 575 Stadium Mall Drive, West Lafayette, IN 47907, USA.

[8]Department of Pharmacology, Hebei Medical University, Shijiazhuang, China

[9]Department of Developmental Cell Biology, China Medical University, 77 Puhe Road, Shenbei New District, 110122 Shenyang, China.

[10]Center for Mitochondrial Biology and Medicine, The Key Laboratory of Biomedical Information Engineering of Ministry of Education, School of Life Science and Technology, Xi'an Jiaotong University, Xi'an, China

[#] These authors contributed equally to this work

[&] Lead author

* Corresponding author: Jin-Peng Sun (corresponding author)

                E-mail: sunjinpeng@sdu.edu.cn

                Xiao Yu (corresponding author)

                E-mail: yuxiao@sdu.edu.cn

                Min Cui (corresponding author)

40   E-mail: cuimin@sdu.edu.cn

41   Yun-Fei Xu (corresponding author)

42   E-mail:xuyunfei1988@126.com

ABSTRACT

Tyrosine phosphorylation of secretion machinery proteins is a crucial regulatory mechanism for exocytosis. However, the participation of protein tyrosine phosphatases (PTPs) in different exocytosis stages has not been defined. Here we demonstrated that PTP-MEG2 controls multiple steps of catecholamine secretion. Biochemical and crystallographic analyses revealed key residues that govern the PTP-MEG2 and NSF-pY^{83} site interactions, specify PTP-MEG2 substrate selectivity and modulate the fusion of catecholamine-containing vesicles. Unexpectedly, delineation of PTP-MEG2 mutants along with the NSF binding interface revealed that PTP-MEG2 controls the fusion pore opening through non-NSF dependent mechanisms. Utilizing bioinformatics search and biochemical and electrochemical screening approaches, we uncovered that PTP-MEG2 regulates the opening and extension of the fusion pore by dephosphorylating the DYNAMIN-2-pY^{125} and MUNC18-1-pY^{145}, which is associated with epileptic encephalopathy. The crystal structure of the PTP-MEG2/phospho-MUNC18-1-pY^{145} segment confirmed the interaction of PTP-MEG2 with MUNC18-1 through a distinct structural basis. Our studies extended mechanistic insights in complex exocytosis processes.

KEY WORDS:

exocytosis, tyrosine phosphorylation, structure, catecholamine, PTP-MEG2

INTRODUCTION

Secretion via vesicle exocytosis is a fundamental biological event involved in almost all physiological processes (Dittman and Ryan, 2019; Neher and Brose, 2018; Wu et al., 2014). The contents of secreted vesicles include neuronal transmitters, immune factors and other hormones (Alvarez de Toledo et al., 1993; Magadmi et al., 2019; Sudhof, 2013). There are three main exocytosis pathways in secretory cells, namely, full-collapse fusion, kiss-and-run, and compound exocytosis, which possess different secretion rates and release amount (Sudhof, 2004). It was previously reported that phosphorylation of critical proteins at serine/threonine or tyrosine residues participated in stimulus–secretion coupling in certain important exocytosis processes, for example, the secretion of insulin from pancreatic β cells and

secretion of catecholamine from the adrenal medulla (Laidlaw et al., 2017; Ortsater et al., 2014; Seino et al., 2009). However, the exact roles of protein tyrosine phosphatases (PTPs) in the regulation of key hormone secretion procedures are not fully understood.

The 68-kDa PTP-MEG2, encoded by *ptpn9*, is a non-receptor classical PTP encompassing a unique N-terminal domain with homology to the human CRAL/TRIO domain and yeast Sec14p (Alonso et al., 2004; Cho et al., 2006; Gu et al., 1992; Huynh et al., 2004; Zhang et al., 2016; Zhang et al., 2012). The N-terminal Sec14p homology domain of PTP-MEG2 recognizes specific phospholipids in the membrane structure and is responsible for its specific subcellular location. In secretory immune cells, PTP-MEG2 has been suggested to regulate vesicle fusion via directly dephosphorylating the pY[83] site of N-ethylmaleimide-sensitive fusion protein (NSF) (Huynh et al., 2004). However, many key issues regarding PTP-MEG2-regulated cell secretion remain controversial or even unexplored. For example, it is uncertain whether PTP-MEG2 regulates vesicle exocytosis only inside immune cells (Zhang et al., 2016) and play only insignificant roles in other hormone secretion processes. It remains elusive whether PTP-MEG2 regulates vesicle trafficking pathways other than the NSF-mediated vesicle fusion.

Functional characterization of PTP-MEG2 in vivo normally requires a knockout model; however, PTP-MEG2-deficient mice show neural tube and vascular defects, and the lack of PTP-MEG2 is embryonic lethal (Wang et al., 2005). Alternatively, a specific small molecule inhibitor of PTP-MEG2 has the properties of fast action and no compensatory effect, which enable it to serve as a powerful tool to investigate PTP-MEG2 functions (Yu and Zhang, 2018; Zhang et al., 2012). Recently, we developed a potent and selective PTP-MEG2 inhibitor, Compound 7, that has a Ki of 34 nM and shows at least 10-fold selectivity for PTP-MEG2 over more than 20 other PTPs (Zhang et al., 2012). The application of this selective PTP-MEG2 inhibitor in combination with electrochemical approaches enabled us to reveal that PTP-MEG2 regulates multiple steps of catecholamine secretion from the adrenal medulla by controlling the vesicle size, the release probabilities of individual vesicles and the initial opening of the fusion pore during exocytosis. Further crystallographic study of the PTP-MEG2 protein in complex with the pY[83]-NSF fragment and enzymological kinetic studies captured the transient interaction between PTP-MEG2 and NSF and provided the structural basis for PTP-MEG2 substrate specificity. Interestingly, by delineating the substrate specificity of deficient PTP-MEG2 mutants in the study of catecholamine secretion from primary chromaffin cells, our results suggested that PTP-MEG2 regulated the initial opening of the fusion pore during exocytosis by regulating substrates other than the known NSF through distinct structural bases. We therefore took advantage of this key knowledge and utilized bioinformatics analysis, GST pull-down screening, enzymology and electrochemistry to identify the potential key PTP-MEG2 substrates involved in fusion pore opening. These

experiments led to the identification of several new PTP-MEG2 substrates in the adrenal medulla, of which DYNAMIN-2-pY[125] and MUNC18-1-pY[145] are the crucial dephosphorylation sites of PTP-MEG2 in the regulation of initial pore opening and expansion. Further crystallographic analysis and functional assays with MUNC18-1 Y[145] revealed the structural basis of the recognition of MUNC18-1 by PTP-MEG2 and how Y[145] phosphorylation regulates fusion pore opening. Our results provide new information and mechanistic insights regarding how dynamic tyrosine phosphorylation and PTP-MEG2 explicitly regulate different processes of vesicle fusion and hormone secretion, both of which may lead to human disease when dysregulated.

RESULTS

Phosphatase activity of PTP-MEG2 is required for catecholamine secretion from adrenal glands

Endogenous PTP-MEG2 expression was readily detected in mouse adrenal gland and chromaffin cell line PC12 cells (Supplemental Fig. 1A-B). PTP-MEG2 knockout is embryonic lethal (Wang et al., 2005). To investigate the functional role of PTP-MEG2 in catecholamine secretion from adrenal glands, we therefore applied our newly developed active-site directed and specific PTP-MEG2 inhibitor (Compound 7) (Supplemental Fig. 1C). Compound 7 is a potent and cell permeable PTP-MEG2 inhibitor, with a Ki value of 34 nM and extraordinary selectivity against other phosphatases (Zhang et al., 2012). The administration of either high concentrations of potassium chloride (70 mM) or 100 nM Angiotensin II (AngII) significantly increased the secretion of both epinephrine (EPI) and norepinephrine (NE) from adrenal medulla, as previously reported (Liu et al., 2017; Teschemacher and Seward, 2000), and this effect was specifically blocked by pre-incubation with Compound 7 (400 nM) for 1 hour (Fig. 1A-D). Notably, basal catecholamine secretion was also decreased after pre-incubation with Compound 7 (Fig. 1A-D). However, the intracellular catecholamine contents were not changed in response to Compound 7 incubation (Supplemental Fig. 1 D-E). These results indicated that PTP-MEG2 plays an essential role in catecholamine secretion from the adrenal medulla.

PTP-MEG2 inhibition reduced the quantal size and the release probabilities of catecholamine secretion from individual vesicles.

We used the carbon fiber electrode (CFE) to characterize the effects of PTP-MEG2

inhibition on the kinetics of catecholamine secretion from primary cultured chromaffin cells (Chen et al., 2005; Harada et al., 2015) (Fig. 1E-F and Supplemental Fig. 1 F-G). The Angiotensin II-induced catecholamine secretion was gradually attenuated by increasing the concentration of Compound 7 after pre-incubation with the primary chromaffin cells, from 20% at 100 nM Compound 7 to 80% at 2 μM Compound 7 (Fig. 1E-F and Supplemental Fig. 1 F-H). We then compared isolated amperometric spikes of chromaffin cells pre-incubated with different concentrations of Compound 7 to determine the effect of the inhibition of PTP-MEG2 activity on quantal size (total amperometric spike charge) and vesicle release probability. The application of PTP-MEG2 inhibitor reduced quantal size, as indicated by statistical analysis of the quantal size distribution and averaged amperometric spike amplitude (Fig. 1G and Supplemental Fig. 1I). Specifically, the peak of the amperometric spikes was reduced from 1.1 pC to 0.5 pC after incubation with 400 nM Compound 7 (Fig. 1G). The number of AngII-induced amperometric spikes was also significantly decreased in the presence of Compound 7(Supplemental Fig. 1 J), suggesting that either the release probabilities of individual vesicles or the size of the readily released pool was affected.

We next used transmission electron microscopy to examine the location of the large-dense-core vesicles (LDCVs) in the adrenal gland medulla after incubation with AngII and Compound 7 (Fig. 1H-1I and Supplemental Fig. 2 A-B). The intracellular distribution of the LDCVs were summarized by 50 nM bins according to their distance from the chromaffin plasma membrane (Fig. 1J). Vesicles with distance less than 50nm was generally considered as in docking stage. Here we found that the application of 400 nM Compound 7 significantly decreased the number of docking vesicles in contact with the plasma membrane (Fig. 1J). Moreover, Compound 7 significantly increased the numbers of LDCVs with diameters less than 50 nm, and decreased the LDCVs numbers with diameters larger than 150 nm (Fig. 1K), which is consistent with the statistical analysis of the total amperometric spike charge obtained by electrochemical measurement (Supplemental Fig. 1 H-M). This change of vesicle size in adrenal medulla recalled the functional studies of PTP-MEG2 in immunocytes , which demonstrated that vesicle was excessively larger if PTP-MEG2 was overexpressed (Huynh et al., 2004).

PTP-MEG2 regulates the initial opening of the fusion pore

Generally, the presence of pre-spike foot (PSF) is a common phenomenon preceding large amperometric spikes of catecholamine secretion in chromaffin cells, while stand-alone foot (SAF) is considered to represent "kiss-and-run" exocytosis (Chen et al., 2005). Both PSF

 and SAF are generally considered indications of the initial opening of the fusion pore

 (Alvarez de Toledo et al., 1993; Zhou et al., 1996). In this study, Compound 7 substantially

 decreased the PSF frequency from 25% to 3%, whereas attenuated SAF frequency from 18%

 to 4% in response to AngII stimulation (Fig. 1L). The inhibitor Compound 7 had no

 significant effect on the PSF duration (Supplemental Fig. 2 C-D) but markedly reduced the

 amplitude and charge of PSF (Supplemental Fig. 2 E-F). Intriguingly, the duration of SAF

 was also decreased with increasing Compound 7 concentration, in contrast to the PSF

 (Supplemental Fig. 2 G-H). Additionally, the average amplitude and charge of SAF decreased

 from 14 pA to 5 pA and from 150 fC to 65 fC, respectively (Supplemental Fig. 2 I-J). These

 results suggested that PTP-MEG2 activity was likely required for the initial opening of the

 fusion pore, or influenced the endocytosis rate of partially fused vesicles.

 The crystal structure of the PTP-MEG2/phospho-NSF complex revealed significant

 structural rearrangement in the WPD loop and β3-loop-β4

 PTP-MEG2 is known to modulate interleukin-2 secretion in macrophages via

 de-phosphorylation of NSF, a key regulator in vesicle fusion (Huynh et al., 2004). In response

 to stimulation with either high potassium chloride or AngII, the two stimulators for

 catecholamine secretion from the adrenal medulla, the tyrosine phosphorylation of NSF

 increased, indicating that NSF phosphorylation actively participates in catecholamine

 secretion (Fig. 2A and Supplemental Fig. 3 A-B). Moreover, a significant portion of NSF

 co-localized with PTP-MEG2 in adrenal medulla upon AngII stimulation, and the substrate

 trapping mutant PTP-MEG2-D^{470}A interacted with the tyrosine-phosphorylated NSF from the

 adrenal medulla treated with peroxide (Fig. 2B-D and Supplemental Fig. 3C-D), indicating

 that PTP-MEG2 regulates catecholamine secretion in chromaffin cells through direct

 dephosphorylation of NSF.

 We therefore co-crystallized PTP-MEG2/phospho-NSF-E^{79}-pY^{83}-K^{87}, and the structure

 was solved at 1.7 Å resolution (Table 1). The 2Fo-Fc electron density map allowed

 unambiguous assignment of the phospho-NSF-E^{79}-pY^{83}-K^{87} in the crystal structure (Fig.

 2E-F). Importantly, the binding of phospho-NSF-E^{79}-pY^{83}-K^{87} induced substantial

 conformational changes at both the WPD loop and β3-loop-β4 compared to the crystal

 structure of PTP-MEG2 alone (Barr et al., 2009) (Fig. 2G-H). Specifically, the interaction of

 the phosphate of the pY^{83} of the NSF with the guanine group of the R^{521} of the PTP-MEG2

 induced a rotation of approximately 90 degrees, which resulted in the movement of the W^{468}

 and a traverse of 7 Å of the WPD loop for a closed state (Fig. 2I-J). Unique to the

PTP-MEG2/substrate complex, the movement of L^{466} in the WPD loop by 1.7 Å enabled the

formation of a new hydrogen bond between its main chain carbonyl group and the side chain

of R^{403} (Fig. 2I-J). The disruption of the salt bridge between E^{406} and R^{521} also contributed to

the new conformational state of the N-terminal of β3-loop-β4 (Fig. 2K-L). The presence of

phosphate in the PTP-MEG2 active site C-terminal to β3-loop-β4 caused a 180 degree

rotation of the side chain of S^{516}, allowing its side chain oxygen to form a new hydrogen bond

with the main chain carbonyl of the K^{411} (Fig. 2K-L). This structural rearrangement altered the

side chain conformation of K^{411}, which pointed to the solvent region and formed new polar

interactions with E^{406} and the carbonyl group of R^{409} (Fig. 2K-L). Moreover, the presence of

the phospho-NSF-E^{79}-pY^{83}-K^{87} peptide between R^{409} and D^{335} disrupted their charge

interaction, which enabled a movement of the main chain from G^{408} to K^{411}, accompanied by a

side chain movement of R^{410} and the formation of new polar interactions with the main chain

of P^{378} and C^{412} (Fig. 2M-N). The structural rearrangements that occurred at WPD and

β3-loop-β4 enabled the accommodation of the phospho-substrate of PTP-MEG2 and may be

important for its appropriate interaction with its physiological substrates/partners and

subsequent activation.

Structural basis of the PTP-MEG2-NSF interaction

The structural analysis identified critical residues for phospho-substrate recognition by

PTP-MEG2 (Fig. 3A). PTP-MEG2 Y^{333} forms extensive hydrophobic interaction with the

phenyl ring of pY^{83} in NSF. Mutation of this residue caused a significant decrease of more

than 8-fold in activity for both p-nitrophenyl phosphate (pNPP) and the phospho-NSF-peptide

(Fig. 3B-C), suggesting that this residue is important for PTP-MEG2 recognition of all

substrates with phenyl rings. N-terminal to pY^{83}, the side chain oxygen of S^{81} forms a

hydrogen bond with the carbonyl oxygen of the main chain of R^{332} of PTP-MEG2. The

carbonyl oxygen of the main chain of S^{81} forms a hydrogen bond with the amide of G^{334}, and

L^{82} forms hydrophobic interactions with the side chain of D^{335} (Fig. 3A). Specifically,

mutation of G^{334} to R impaired the activity of PTP-MEG2 towards the NSF-pY^{83}

phospho-peptide but had no effect on its intrinsic phosphatase activity as measured by using

pNPP as a substrate, suggesting that G^{334} plays an important role in the recognition of the

N-terminal conformation of the peptide substrate (Fig. 3B-C and Supplemental Fig. 4 A-B).

The D^{335} in the pY binding loop of PTP-MEG2 is also critical for determining the

peptide orientation of the substrate by forming important hydrogen bonds with the main chain

amide and carbonyl groups of pY^{83} and T^{84} in NSF. Residues C-terminal to NSF-pY^{83}, T^{84} and

F^{85} form substantial hydrophobic interactions with V^{336}, F^{556}, Q^{559} and Y^{471} (Fig. 3A).

Accordingly, mutation of D^{335}A or Q^{559}A showed no significant effect on pNPP activity but

substantially decreased their activities towards the phospho-NSF peptide (Fig. 3B-C and

Supplemental Fig. 4 A-B). Mutation of I^{519}A caused a decrease in the intrinsic activity of

PTP-MEG2 and a further decrease of approximately 256 -fold in its ability to

dephosphorylate phospho-NSF-peptide. Moreover, Y^{471} forms extensive hydrophobic

interactions with T^{84} and F^{85} and a hydrogen bond with the carboxyl group of pY^{83}. Mutation

of Y^{471} to either A or F greatly reduced its activity towards phospho-NSF- E^{79}-pY^{83}-K^{87}

peptide but had little effect on pNPP dephosphorylation. Taken together, the structural

analysis and enzymology studies identified G^{334}, D^{335}, Y^{471}, I^{519} and Q^{559} as critical residues

for the substrate recognition of NSF by PTP-MEG2. Importantly, although none of these

residues are unique to PTP-MEG2, the combination of these residues is not identical across

the PTP superfamily but is conserved between PTP-MEG2 across different species,

highlighting the important roles of these residues in mediating specific PTP-MEG2 functions

(Fig. 3D and Supplemental Fig. 5).

Molecular determinants of Q^{559}:D^{335} for the substrate specificity of PTP-MEG2

The pY+1 pocket is an important determinant of substrate specificity in different PTP

superfamily members (Barr et al., 2009; Li et al., 2016; Wang et al., 2014; Yu et al., 2011).

The pY+1 pocket of PTP-MEG2 was found to consist of D^{335}, V^{336}, F^{556} and Q^{559} (Fig. 4A).

Unique to the PTP-MEG2/NSF-E^{79}-pY^{83}-K^{87} complex structure, a relatively small T^{84} residue

occurs at the pY+1 position, in contrast to L^{993} in the PTP1B/EGFR-pY^{992} complex structure,

D^{76} in   the   LYP/SKAP-HOM-pY^{75} complex   structure   and   L^{1249} in   the

PTPN18/phospho-HER2-pY^{1248} complex structure (Fig. 4A). Although several PTPs have an

equivalent D:Q pair similar to D^{335}:Q^{559} of PTP-MEG2 determining the entrance of the pY+1

residue into the pY+1 pocket, such as PTP1B, LYP, PTPN18, STEP, PTP-MEG1, SHP1,

PTPH1 and SHP2, structural analysis indicated that the Cβ between Q^{559} and D^{335} is the

smallest in PTP-MEG2, at least 1 Å narrower than in the other PTP structures examined (Fig.

4B). The narrower pY+1 pocket entrance could be a unique feature of substrate recognition

by PTP-MEG2.

PTP-MEG2 regulates two different steps of catecholamine secretion through processing

of different substrates

We   next   infected   primary   chromaffin   cells   with   lentivirus   encoding   wild-type

PTP-MEG2 or different mutants for electrochemical investigation of the structure-function

relationship of PTP-MEG2 in the regulation of catecholamine secretion (Fig. 5A,

Supplemental Fig. 6 A-G and Supplemental Fig. 7 A-D). In addition to wild-type PTP-MEG2,

we chose 6 PTP-MEG2 mutants, including $G^{334}R$, $D^{335}A$, $Y^{471}A$, $Y^{471}F$, $I^{519}A$ and $Q^{559}A$,

whose positions are determinants of the interactions between PTP-MEG2 and NSF from the

pY^{83}-1 position to the pY^{83}+2 position (Fig. 5B). Cells with approximately 15 folds of

overexpressed exogenous PTP-MEG2 wild type or mutants than the endogenous PTP-MEG2

were selected for electrophysiology studies (Supplemental Fig. 6 E-F). The PTP-MEG2

mutations did not significantly affect its interaction with NSF, suggesting that residues not

locating within the active site of PTP-MEG2 may also participate in NSF associations

(Supplemental Fig. 6G). The overexpression of wild-type PTP-MEG2 significantly increased

both the number and amplitude of the amperometric spikes, which are indicators of the

release probabilities of individual vesicles and quantal size, respectively (Fig. 5C-E and

Supplemental Fig. 7 E). In contrast, expression of $G^{334}R$, $D^{335}A$, $Y^{471}A$, $Y^{471}F$, $I^{519}A$ and $Q^{559}A$

all significantly decreased the quantal size, the release probabilities of individual vesicles, the

half width and the rise rate of each spike (Fig. 5C-5E, Supplemental Fig. 7 E-F and

Supplemental Fig. 7 H). However, there was no difference on the rise time between these

mutations and the wild type (Supplemental Fig. 7 G). These results suggested that the

interaction of PTP-MEG2 with NSF is important for controlling vesicle size and the release

probability of catecholamine secretion.

Unexpectedly, the PTP-MEG2 mutants showed different effects on the probability of the

occurrence of the foot, including both PSF and SAF, which is an indicator of the initial

opening of the fusion pore (Fig. 5F-G). $G^{334}R$ mutation which disrupted the recognition of the

N-terminal conformation of the peptide substrate of PTP-MEG2, and $D^{335}A$ and $Q^{559}A$, which

are the determinants of pY+1 substrate specificity, significantly reduced the AngII -induced

foot probability. The mutations of I^{519} and Y^{471}, which form specific interactions with the T^{84}

and F^{85} of the NSF and are determinants of the C-terminal region of the central

phospho-tyrosine involved in the substrate specificity of PTP-MEG2, showed no significant

effect (Fig. 5B, Fig. 5F and 5G and Supplemental Fig. 7 I-N). These results indicated that

PTP-MEG2 regulated the initial opening of the fusion pore via a distinct structural basis from

that of vesicle fusion, probably through dephosphorylating other unknown substrates. As

$D^{335}A$ and $Q^{559}A$ of PTP-MEG2 maintained the occurrence of the foot probability, the

unknown PTP-MEG2 substrate that regulates the fusion pore opening should have a small

residue, such as G, A, S or T at the pY+1 position. Conversely, the unknown PTP-MEG2

substrate should have a less hydrophobic residue at the pY+2 position because $Y^{471}F$, $Y^{471}A$

and $I^{519}A$ of PTP-MEG2 had no significant effect on the foot probability.

Identification of new PTP-MEG2 substrates that contributed to the initial opening of the fusion pore

The effects of PTP-MEG2 mutations along the PTP-MEG2/NSF-phospho-segment interface on catecholamine secretion indicated that a PTP-MEG2 substrate other than NSF with distinct sequence characteristics contributes to the regulation of "foot probability" (Fig. 5). We therefore utilized this key information to search for new potential PTP-MEG2 substrates by bioinformatics methods (Fig. 6A). First, we searched for the keywords "fusion pore", "secretory vesicle" and "tyrosine phosphorylation" using the functional protein association network STRING and the text mining tool PubTator, which resulted in a candidate list of 54 proteins. Second, we applied UniProt by selecting proteins located only in the membrane or vesicle, which limited the candidates to 28 members. Third, as our experiments were carried out in the adrenal gland, we used the Human Protein Atlas database for filtering to excluded the proteins which is not detectable in adrenal gland, which narrowed the candidate list to 23 proteins. Finally, we exploited the post-translational-motif database PhosphoSitePlus to screen candidate proteins with potential phospho-sites that matched the sequence requirements at both the pY+1 position and the pY+2 position, which are "G, S, A, T, V, P" and "G, A, S, T, C, V, L, P, D, H", respectively. These positions were further evaluated by surface exposure if a structure was available. The combination of these searches produced 12 candidate lists with predicted pY positions (Fig. 6B).

To biochemically characterize whether these proteins are substrates of PTP-MEG2, we transfected the plasmids encoding the cDNAs of these candidate proteins into PC12 cells, stimulated the cells with AngII, and performed a pull-down assay with the GST-PTP-MEG2-D[470]A trapping mutant or GST controls. The known PTP-MEG2 substrate NSF was used as a positive control. Notably, six candidates, including PACSIN1, MUNC18-1, VAMP7, SNAP25, DYNAMIN 1 and DYNAMIN 2 showed specific interactions with PTP-MEG2 after AngII stimulation in PC12 cells (Fig. 6C and Supplemental Fig. 8A). Whereas the protein of MUNC18-1, VAMP7, DYNAMIN 2 and SNAP25 were readily detected in the adrenal medulla, DYNAMIN 1 and PACSIN1 showed substantially lower expression than that in the liver and brain (Supplemental Fig. 8B). Moreover, whereas MUNC18-1, VAMP7 and DYNAMIN 2 strongly co-localized with PTP-MEG2 in the adrenal medulla (Fig. 6D-F), the co-localization of SNAP25 and PACSIN1 with PTP-MEG2 was relatively weak (Supplemental Fig. 8C-D). MUNC18-1, VAMP7 and DYNAMIN 2 are candidate PTP-MEG2 substrates which were further strengthened by the fact that the high potassium chloride- or AngII-stimulated tyrosine phosphorylation of these proteins in the

adrenal medulla was significantly dephosphorylated by PTP-MEG2 in vitro (Supplemental

Fig. 9 A-C, E), whereas the tyrosine phosphorylation of SNAP25 had no obvious change

(Supplemental Fig. 9 D, E).

.

PTP-MEG2 regulated the initial opening of the fusion pore through dephosphorylating

MUNC18-1- pY^{145}

The predicted PTP-MEG2 dephosphorylation site on MUNC18-1 (also called STXBP1)

is Y^{145}, which is localized on the β sheet linker and forms extensive hydrophobic interactions

with surrounding residues (Hu et al., 2011; Yang et al., 2015) (Fig. 7A and Supplemental Fig.

10A). Moreover, the phenolic-oxygen of Y^{145} forms specific hydrogen bonds with the main

chain amide of F^{540} and the main chain carbonyl oxygens of I^{539} and G^{568} (Fig. 7B). These key

interactions might be involved in regulating the arrangement of the arc shape of the three

domains of MUNC18-1, by tethering the interface between domain 1 and domain 2. The

phosphorylation of Y^{145} likely abolishes this H-bond network and changes its ability to

associate with different snare complexes participating in vesicle fusion procedures (Fig. 7B).

Interestingly, a missense mutation of MUNC18-1 Y^{145}H was found to be associated with early

infantile epileptic encephalopathy (Stamberger et al., 2017) (Fig. 7C).

Notably, the Y^{145}H missense mutation may not be phosphorylated properly. We therefore

overexpressed wild-type MUNC18-1, Y^{145}A, a non-phosphorylable mutant, and the

disease-related Y^{145}H mutant in PC12 cells stimulated with AngII and then examined their

abilities to interact with PTP-MEG2 trapping mutant. The GST pull-down results indicated

that all MUNC18-1 Y^{145} mutations significantly decreased their ability to associate with

PTP-MEG2 (Fig. 7D, supplemental Fig. 9G). In contrast, mutation of the predicted

phosphorylation sites of SNAP25, the SNAP25-Y^{101}A, had no significant effect on their

interactions with PTP-MEG2 (Supplemental Fig. 9 F, G). These results confirmed that pY^{145}

is the major site of MUNC18-1 regulated by PTP-MEG2 (Fig. 7D).

We next infected primary chromaffin cells with a lentivirus encoding wild-type

MUNC18-1, the MUNC18-1 Y^{145}-tyrosine phosphorylation-deficient mutant Y^{145}F, the

MUNC18-1 Y^{145}-tyrosine phosphorylation mimic mutant Y^{145}E and the disease-related mutant

Y^{145}H. We examined the effects of these mutants using primary chromaffin cells harbouring

approximately 20~30 folds MUNC18-1 overexpression related to that of endogenous

expression levels (Supplemental Fig. 10 B-D). Interestingly, either Y^{145}E or Y^{145}H

significantly reduced the PSF and SAF percentage of catecholamine secretion in response to

AngII stimulation, whereas the phosphorylation-deficient mutant Y^{145}F showed no significant

effects on the PSF or SAF percentage compared to that of the wild type (Fig. 7E-G, and Supplemental Fig. 10 and 11). These results suggested that the tyrosine phosphorylation of Y^{145} impaired the initial opening of the fusion pore in agonist-induced catecholamine secretion in primary chromaffin cells.

To further dissect the mechanism underlying the phosphorylation of MUNC18-1 Y^{145} as well as the disease-related mutant $Y^{145}H$ in the regulation of hormone secretion, we compared the interactions of wild-type and mutant MUNC18-1 with the binding partner SYNTAXIN1 (Lim et al., 2013). Importantly, both the phosphorylation mimic mutant MUNC18-1-$Y^{145}E$ and the disease-related mutant $Y^{145}H$ significantly impaired the interaction between MUNC18-1 and SYNTAXIN1 (Fig. 7H). These results suggested that the phosphorylation of Y^{145} of MUNC18-1 decreases the binding of MUNC18-1 to SYNTAXIN1, which plays an important role in the formation of the foot during catecholamine secretion. In contrast, the dephosphorylation of pY^{145} of MUNC18-1 by PTP-MEG2 promoted initial pore opening and fusion.

PTP-MEG2 regulated the initial opening of the fusion pore through dephosphorylating DYNAMIN 2-pY^{125}

Except for MUNC18-1, two additional substrates of PTP-MEG2, which are the Y^{45} site of VAMP7 and the Y^{125} site of DYNAMIN 2, were confirmed by GST-pull down assay together with specific mutating effects on the corresponding phosphorylation sites (Fig. 7D, supplemental Fig. 9F, G). It was well known that both DYNAMIN1 and DYNAMIN2 are active regulators of pore fusion dynamics, involved in the pore establishment, stabilization, constriction and fission processes (Jones et al., 2017; Shin et al., 2018; Zhao et al., 2016). The dephosphorylation site of DYNAMIN 2 by PTP-MEG2, the Y^{125}, is located in a β-sheet of DYNAMIN GTPase domain according to the solved DYNAMIN-1 structure (PDB ID: 5D3Q). Previous studies have identified that phosphorylation of DYNAMIN close to the G domain significantly increased its GTPase activity, thus promoting endocytosis (Kar et al., 2017). We therefore hypothesized that the phosphorylation of the DYNAMIN-2 pY^{125} site increased pore fission by up-regulating its GTPase activity, whereas dephosphorylation of DYNAMIN-2 pY^{125} site by PTP-MEG2 reversed it. Consistent with such hypothesis, the primary chromaffin cells transduced by lentivirus encoding DYNAMIN-2 wild type and the $Y^{125}F$ mutant significantly decreased percentage of the PSF and SAF of catecholamine secretion elicited by AngII, whereas the phosphorylation mimic mutant DYNAMIN-2 $Y^{125}E$ mutant exhibited significantly more reduction for the PSF and SAF percentage (Fig. 7I-K and Supplemental Fig. 12). The DYNAMIN mediated pore fission process is closely correlated

 with the receptor endocytosis. Whereas overexpression of the DYNAMIN-2 wild type or

 Y125E phospho-mimic mutant markedly increased the AngII induced AT1aR endocytosis, the

 Y125F mutant has no significant effect, suggesting DYNAMIN-2 $Y^{125}E$ is a gain of function

 mutant to promote the membrane fission (Fig. 7L) (Liu et al., 2017). Consistent with these

 cellular results, the in vitro GTPase activity further suggested that the phosphorylation of

 DYNAMIN-2 at pY^{125} site up-regulated its activity (Fig. 7M).

Molecular mechanisms of the PTP-MEG2-MUNC18-1 and PTP-MEG2-DYNAMIN-2 interaction

The k_{cat}/K_m of PTP-MEG2 towards a phospho-segment derived from MUNC18-1-pY^{145} is very similar to that obtained with a phospho-segment derived from the known substrate pY^{83} site of NSF (Supplemental Fig. 4A-B and 8A). We therefore crystallized the PTP-MEG2 trapping mutant with the MUNC18-1-E^{141}-pY^{145}-S^{149} phospho-segment and determined the complex structure at 2.2 Å resolution (Table 1). The 2Fo-Fc electro-density map allowed the unambiguous assignment of seven residues of the phospho-MUNC18-1-E^{141}-pY^{145}-S^{149} segment in the crystal structure (Fig. 8A). Importantly, comparing with the phospho-NSF-E^{79}-pY^{83}-K^{87} segment, the phospho-MUNC18-1-E^{141}-pY^{145}-S^{149} displayed different interaction patterns with the residues in the PTP-MEG2 active site, forming new interactions with R^{409} and R^{410} but lost interactions with Y^{471} and I^{519} (Fig. 8B). Whereas PTP-MEG2 Y^{471} formed extensive hydrophobic interactions with NSF-T^{84} and NSF-F^{85}, as well as a well-defined hydrogen bond (2.4 Å) with the carbonyl oxygen of NSF-pY^{83}, PTP-MEG2 Y^{471} formed only a weaker H-bond with the carbonyl oxygen of MUNC18-1-pY^{145} due to a 0.62 Å shift of Y^{471} away from the central pY substrate (Fig. 8C and Supplemental Fig. 13A). Similarly, PTP-MEG2 I^{519} did not form specific interactions with the MUNC18-1-E^{141}-pY^{145}-S^{149} segment except for the central pY. In contrast, PTP-MEG2 I^{519} formed specific hydrophobic interactions with NSF-T^{84} (Fig. 8D and Supplemental Fig. 13B). Consistently, mutations of PTP-MEG2 $Y^{471}A$, $Y^{471}F$ or $I^{519}A$ significantly decreased the phosphatase activity towards the phospho-NSF-E^{79}-pY^{83}-K^{87} segment (Fig. 3C) but had no significant effect on the phospho-MUNC18-1-E^{141}-pY^{145} -S^{149} segment (Fig. 8E and Supplemental Fig.14). Meanwhile, $R^{409}A$ and $R^{410}A$ mutations substantially decreased the catalytic activity toward phospho-segements derived from both MUNC18-1-pY^{145} and DYNAMIN-2 pY^{125} sites, indicating that recognition of these substrates was mediated by common residues (Fig. 8F and Supplemental Fig. 14). Notably, the PTP-MEG2 catalytic domain mutations shows no significant effects on the interactions between PTP-MEG2 with the NSF, MUNC18-1 and DYNAMIN-2, indicating other domain of PTP-MEG2 participated into the association of PTP-MEG2 with these proteins (Supplemental Fig. 6G and Supplemenatl Fig. 13C). Combined with the results that

PTP-MEG2 $Y^{471}A$, $Y^{471}F$ and $I^{519}A$ affect only the spike number and amount but show not the

foot probability (Fig. 5B-G), these data suggested that PTP-MEG2 regulated intracellular

vesicle fusion by modulating the NSF-pY^{83} phospho-state but regulated the process of vesicle

fusion pore initiation by dephosphorylating MUNC18-1 at pY^{145} site and DYNAMIN 2 at

pY^{125} site (Fig. 8G).

DISCUSSION

Posttranslational modifications of secretion machinery proteins are known as powerful

ways to regulate exocytosis. In contrast to the well-characterized serine/threonine

phosphorylation, the importance of tyrosine phosphorylation in exocytosis has only recently

begun to be appreciated (Cijsouw et al., 2014; Gabel et al., 2019; Jewell et al., 2011; Laidlaw

et al., 2017; Meijer et al., 2018; Seino et al., 2009). In addition to the phosphorylation of NSF

at its pY^{83} site, recent studies have shown that tyrosine phosphorylation of MUNC18-3 at the

pY^{219} and pY^{527} sites, Annexin-A2 at pY^{23}, and MUNC18-1 at pY^{473} actively participates in

the vesicle release machinery to explicitly regulate exocytosis processes (Gabel et al., 2019;

Jewell et al., 2011; Meijer et al., 2018). The tyrosine phosphorylation at specific sites of the

signalling molecule is precisely regulated by protein tyrosine kinases and protein tyrosine

phosphatases (Tonks, 2006; Yu and Zhang, 2018). Although tyrosine kinases such as the

insulin receptor, Src and Fyn are acknowledged to play critical roles in hormone secretion

(Jewell et al., 2011; Meijer et al., 2018; Oakie and Wang, 2018; Soares et al., 2013), only a

very few PTPs that regulate the vesicle release machinery have been identified, and the

structural basis of how these PTPs selectively dephosphorylate the key tyrosine

phosphorylation sites governing exocytosis was unknown. In the current study, we

demonstrated that PTP-MEG2 is an important regulator of hormone secretion from the

chromaffin cell, using a selective PTP-MEG2 inhibitor in combination with cellular and

electrochemical amperometric recording. The current study extended the regulatory role of

PTP-MEG2 in various steps of exocytosis in hormone secretion beyond the previously known

simple vesicle fusion step of the immune system (Huynh et al., 2004). We then determined the

crystal structure of PTP-MEG2 in complex with the pY^{83} phospho-segment of the NSF, the

key energy provider for disassembling fusion-incompetent cis SNARE complexes in the

process of vesicle fusion in immunocytes (Huynh et al., 2004). The complex structure not

only revealed the structural rearrangement in PTP-MEG2 upon binding the NSF substrate,

identifying Q^{559}:D^{335} as the key pair for substrate specificity of the pY+1 site, but also

provided clues that PTP-MEG2 regulates the initial opening of the fusion pore through

another unknown substrate. Fortunately, we were able to deduce the signature of the pY+1

and pY+2 positions of this unknown substrate by carefully inspecting the

PTP-MEG2/phospho-NSF-E[79]-pY[83]-K[87] complex structure and analysing the functional data of the PTP-MEG2 interface mutants. Further bioinformatics studies and cellular and physiological experiments enabled us to discover that PTP-MEG2 regulates the initial opening of the fusion pore by modulating the tyrosine phosphorylation states of the MUNC18-1 at the pY[145] site and the DYNAMIN-2 at the pY[125] site. Therefore, we have revealed that PTP-MEG2 regulates different steps of the exocytosis processes via dephosphorylating distinct substrates. PTP-MEG2 regulates the vesicle size and vesicle-vesicle fusion step by dephosphorylating NSF at its NSF-pY[83] site, whereas it regulates the process of LDCV fusion pore initiation and expansion by controlling specific phosphorylation of MUNC18-1 and DYNAMIN-2. Moreover, our studies highlight that the combination of structural determination and functional delineation of the interface mutants of the protein complex is a powerful approach to characterizing the signalling events and identifying unknown downstream signalling molecules.

Fusion pore opening and expansion is thought to be a complex process requiring the docking of apposed lipid bilayers and involvement of multiple proteins to form a hemi-fusion diaphragm including SNAREs, MUNC18-1, and DYNAMIN (Baker and Hughson, 2016; Hernandez et al., 2012; Jones et al., 2017; Mattila et al., 2015; Sudhof and Rothman, 2009; Zhao et al., 2016). Dissection of the molecular mechanism underlying pore fusion dynamics in exocytosis is challenging because direct observation of this process is difficult to achieve due to the short expansion time and the tiny size of the pore (Baker and Hughson, 2016; Gaisano, 2017; Hong and Lev, 2014). Importantly, the DYNAMINs are key terminators of the fusion pore expansion by scissoring vesicles from the cell membrane (Eitzen, 2003)(Jones et al., 2017; Shin et al., 2018; Zhao et al., 2016). The large scale phosphoproteomic screens have identified multiple tyrosine phosphorylation sites of DYNAMIN, such as that occurring at pY[80], pY[125], pY[354] and pY[597] (Ballif et al., 2008) (Mallozzi et al., 2013). Among them, the Y[597] was a well-studied phosphorylation site of DYNAMIN, which was previously reported to increase self-assembly and GTP hydrolysis in both DYNAMIN 1 and 2 (Kar et al., 2017). However, the Y[597] of DYNAMIN was probably not the target of PTP-MEG2 because the interaction of Y[597] or A[597] with PTP-MEG2 exhibited no significant differences. However, the regulation of other tyrosine phosphorylation site, such as pY[125], as well as how this particular tyrosine phosphorylation participated in a selective secretion process, has not been elucidated. Here, by using pharmacological approach with high selectivity, functional alanine mutagenesis according to structural characterization, combined with bioinformatics, electrophysiology and enzymology, we have demonstrated that PTP-MEG2 negatively regulated the pY[125] phosphorylation state during the catecholamine secretion, which promoted the cessation of the fusion pore expansion through increasing its GTP activity.

In addition to DYNAMIN, MUNC18-1 and its closely related subfamily members have

been demonstrated to participate in several processes during vesicle secretion by interacting

with SNAREs, such as docking, priming and vesicle fusion (Chai et al., 2016; Cijsouw et al.,

2014; Fisher et al., 2001; Gulyas-Kovacs et al., 2007; He et al., 2017; Korteweg et al., 2005;

Ma et al., 2013; Ma et al., 2015; Meijer et al., 2018; Sitarska et al., 2017). This was also in

consistent with our results that vesicle docking was impaired with PTP-MEG2 inhibitor

incubation, since MUNC18-1 was a pivotal modulator of vesicle docking. Importantly, the

tyrosine phosphorylation of MUNC18-1 at Y^{473} was recently reported as a key step in

modulating vesicle priming by inhibiting synaptic transmission and preventing SNARE

assembly (Meijer et al., 2018). In addition, the tyrosine phosphorylation of MUNC18- on Y^{521}

was essential for the dissociation of MUNC18-3 and SYNTAXIN4 (Umahara et al., 2008).

Moreover, the dephosphorylation of MUNC18-1-Y^{145} was suggested to be essential in

maintaining the association between MUNC18-1 and SYNTAXIN1 (Lim et al., 2013). In the

present study, we demonstrated that the MUNC18-1 $Y^{145}E$ phospho-mimic mutation, but not

the non-phosphorylated mutation $Y^{145}F$, significantly decreased the PSF and the SAF

probability. It's very likely that the association of the MUNC18-1 with the SYNTAXIN1 is

required for initial pore opening and expansion. Future studies within the fusion pore

dynamics using in vitro reconstitution system could verify such hypothesis. Consistently, only

the specific PTP-MEG2 mutations that affect its activity towards MUNC18-1 reduced the

PSF and SAF probability. The structural analysis of PTP-MEG2 in complex with

MUNC18-1-pY^{145} and the enzymatic analysis further confirmed these observations.

Collectively, these results indicate that the tyrosine phosphorylation of MUNC18-1-Y^{145},

DYNAMIN 2-Y^{125} and their dephosphorylation by PTP-MEG2 play essential roles in the

regulation of the fusion pore opening and collapse processes.

Notably, the MUNC18-1 $Y^{145}H$ mutation is a known SNP that is associated with epileptic

encephalopathy (Stamberger et al., 2017). $Y^{145}H$ behaves similarly to the MUNC18-1

phosphorylation mimic mutant $Y^{145}E$ by disrupting its interaction with SYNTAXIN1 and

reducing the probability of PSF elicited by AngII in primary chromaffin cells. This

observation provided a clue for the pathological effects of the MUNC18-1 $Y^{145}H$ mutation. In

addition to MUNC18-1, we found that VAMP7 interacted with PTP-MEG2 via their Y^{45}

tyrosine phosphorylation site, respectively. DYNAMIN 1, PASCIN1 and SNAP25 are also

potential PTP-MEG2 substrates depending on specific cellular contexts. The functions of

VAMP7 phosphorylation at the Y^{45} site and its dephosphorylation by PTP-MEG2, the

interaction of PTP-MEG2 with DYNAMIN 2, etc. in the exocytosis process await further

investigation.

Finally, by solving the two crystal structures of PTP-MEG2 in complex with two

substrates, the phospho-NSF-E^{79}-pY^{83}-K^{87} segment and the phospho-MUNC18-1-E^{141}-pY^{145}

-S^{149} segment, we revealed that PTP-MEG2 recognized these functionally different substrates

through distinct structural bases. Whereas K^{411}, Y^{471} and I^{519} contributed most to the selective interaction of PTP-MEG2 with NSF, another set of residues, including R^{409} and R^{410}, mediated the specific binding of PTP-MEG2 to MUNC18-1 (Fig. 8B). Most importantly, mutating Y^{471} and I^{519} to A significantly decreased the activity of PTP-MEG2 towards the phospho-NSF-E^{79}-pY^{83}-K^{87} segment but not the phospho-MUNC18-1-E^{141}-pY^{145}-S^{149} segment. The biochemical data agreed well with the functional data that PTP-MEG2 Y^{471}A and I^{519}A of PTP-MEG2 affected only the vesicle fusion procedure but not the fusion pore opening and expansion processes. These data not only indicate that PTP-MEG2 regulates different steps of exocytosis through different substrates in an explicit temporal and spatial context but also afforded important guidance for the design of selective PTP-MEG2 inhibitors according to the different interfaces between PTP-MEG2 and its substrates to explicitly regulate specific physiological processes, supporting the hypothesis of "substrate-specific PTP inhibitors" (Doody and Bottini, 2014). The design of such inhibitors will certainly help to delineate specific roles of PTP-MEG2 in different physiological and pathological processes.

In conclusion, we have found that PTP-MEG2 regulates two different processes of exocytosis during catecholamine secretion, namely, vesicle fusion and the opening and extension of the fusion pore, through two different substrates with distinct structural bases. We achieved this knowledge by determining the complex structure and performing functional delineation of the protein complex interface mutants. The present study supports the hypothesis that the tyrosine phosphorylation of secretion machinery proteins is an important category of regulatory events for hormone secretion and is explicitly regulated by protein tyrosine phosphatases, such as PTP-MEG2. Dissecting the molecular and structural mechanisms of such modulation processes will provide an in-depth understanding of the exocytosis process and guide further therapeutic development for exocytosis-related diseases, such as epileptic encephalopathy (Stamberger et al., 2017).

METHODS

KEY RESOURCES TABLE

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies		
NSF Antibody	Proteintech	Cat # 21172-1-AP

NSF Antibody	Santa Cruz	Cat # sc-515043
MUNC18-1 Antibody	Proteintech	Cat # 11459-1-AP
MUNC18-1 Antibody	Proteintech	Cat # 67137-1-lg
VAMP7 Antibody	Proteintech	Cat # 22268-1-AP
VAMP7 Antibody	Santa Cruz	Cat # sc-166394
SNAP25 Antibody	Proteintech	Cat # 60159-1-lg
SNAP25 Antibody	Proteintech	Cat # 14903-1-AP
PACSIN1 Antibody	Proteintech	Cat # 13219-1-AP
SYNTAXIN1a Antibody	Proteintech	Cat # 66437-1-lg
pTyr(PY20) Antibody	Santa Cruz	Cat # sc-508
pTyr(PY99) Antibody	Santa Cruz	Cat # sc-7020
FLAG Antibody	Cell Signaling Technology	Cat # 2368
GAPDH Antibody	Cell Signaling Technology	Cat # 5174
PTP-MEG2 Antibody	R&D Systems	Cat # MAB2668
DYNAMIN 1 Antibody	Abcam	Cat # AB52611
DYNAMIN 2 Antibody	Abcam	Cat # AB65556
DYNAMIN 2 Antibody	Santa Cruz	Cat # sc-166669

Chemicals, Peptides, and Recombinant Proteins

Anti-GST-tag beads	TransGen Biotech Co., Ltd	Cat # DP201
Anti-GFP-tag beads	AlpaLife by KangTi Co., Ltd.	Cat # KTSM1301
Protein A/G-PLUS-Agarose	Santa Cruz	Cat # sc-2003
Ni-NTA Agarose	Thermo Fisher scientific	Cat # R90101
The phospho-peptide of NSF "EVSLpYTFDK"	China Peptides Co., Ltd	N/A
The phospho-peptide of MUNC18-1 "ESQVpYSLDS"	China Peptides Co., Ltd	N/A
The phospho-peptide of DYNAMIN 2 "NLRVpYSPHV"	China Peptides Co., Ltd	N/A

Mouse EPI ELISA Kit	Shanghai Jianglai Co.,Ltd	JL11194-48T
Mouse NE ELISA Kit	Shanghai Jianglai Co.,Ltd	JL13969-96T
Angiotensin II (HDRVYIHPF-OH)	China Peptides Co., Ltd	N/A
Compound 7(PTP-MEG2 inhibitor)	Synthesized as previously described (Zhang et al., 2012)	

Experimental Models: Cell Lines

293	ATCC	CRL-1573
293T	ATCC	CRL-3216
PC12	ATCC	CRL-1721

Software and Algorithms

PyMol	Schrödinger	https://www.pymol.org/
CCP4i	Instruct Associate Centre for Integrated Structural Biology	http://www.ccp4.ac.uk/
Phenix	The PHENIX Industrial Consortium	https://www.phenix-online.org
Igor	Fourier Transform	https://www.wavemetrics.com/
Prism	GraphPad	http://www.graphpad.com

621 **CONTACT FOR REAGENT AND RESOURCE SHARING**

622 Further information and requests for resources and reagents should be directed to and will be

623 fulfilled by the Lead Contact, Professor Xiao Yu. (yuxiao@sdu.edu.cn) or Jinpeng Sun

624 (sunjinpeng@bjmu.edu.cn)

 EXPERIMENTAL MODEL AND SUBJECT DETAILS

Cell Culture

The HEK293 cell lines, the 293T cell lines and PC12 cell lines were originally obtained from

the American Type Culture Collection (ATCC). The HEK293 cell lines and 293T cell lines

were grown in DMEM with 10% FBS (Gibco, Grand Island, NY, US) and 1%

penicillin/streptomycin at 37 °C. PC12 cells were maintained at 37 °C in DMEM medium

containing 10% FBS (Gibco, US), 5% donor equine serum (Gibco, US) and 1%

penicillin/streptomycin.

Constructs

Sequences of PTP-MEG2 catalytic domain were subcloned into PET-15b expression vector

with an N-terminal His tag or PGEX-6P-2 expression vector containing an N-terminal GST

tag. The mutations of PTP-MEG2 were constructed by the Quikchange kit from Stratagene .

Plasmid	Vehicle information	Tag information
NSF	PCDNA3.1	FLAG
His-PTPMEG2-WT	pET-15b	His
His-PTPMEG2-333Y-A	pET-15b	His
His-PTPMEG2-334G-R	pET-15b	His
His-PTPMEG2-335D-A	pET-15b	His
His-PTPMEG2-471Y-A	pET-15b	His
His-PTPMEG2-471Y-F	pET-15b	His
His-PTPMEG2-519I-A	pET-15b	His
His-PTPMEG2-559Q-A	pET-15b	His
His-PTPMEG2-409R-A	pET-15b	His
His-PTPMEG2-410R-A	pET-15b	His
PCDH-PTPMEG2-WT	PCDH	GFP
PCDH-PTPMEG2-334G-R	PCDH	GFP
PCDH-PTPMEG2-335D-A	PCDH	GFP
PCDH-PTPMEG2-471Y-A	PCDH	GFP
PCDH-PTPMEG2-471Y-F	PCDH	GFP
PCDH-PTPMEG2-519I-A	PCDH	GFP
PCDH-PTPMEG2-559Q-A	PCDH	GFP
PCDH-PTPMEG2-409R-A	PCDH	GFP

PCDH-PTPMEG2-410R-A	PCDH	GFP	
SYN1	PEGFP-N1	FLAG	GFP
MUNC18-3	PEGFP-N1	FLAG	GFP
PACSIN1	PEGFP-N1	FLAG	GFP
SCAMP1	PEGFP-N1	FLAG	GFP
MUNC18-1	PEGFP-N1	FLAG	GFP
PPP3CA	PEGFP-N1	FLAG	GFP
STX17	PEGFP-N1	FLAG	GFP
VAMP7	CMV10	FLAG	
SYT7	PEGFP-N1	FLAG	GFP
SYT11	PEGFP-N1	FLAG	GFP
SNAP25	PEGFP-N1	FLAG	GFP
DYNAMIN 1	CMV10	FLAG	YFP
DYNAMIN 2	pMYC-C2	FLAG	MYC
MUNC18-1-145Y-A	PEGFP-N1	FLAG	GFP
MUNC18-1-145Y-H	PEGFP-N1	FLAG	GFP
MUNC18-1-145Y-E	PEGFP-N1	FLAG	GFP
MUNC18-1-145Y-F	PEGFP-N1	FLAG	GFP
FLAG-VAMP7-45Y-A	CMV10	FLAG	
FLAG-VAMP7-45Y-C	CMV10	FLAG	
FLAG-DYNAMIN 2-125Y-A	pMYC-C2	FLAG	MYC
FLAG-DYNAMIN 2-125Y-E	pMYC-C2	FLAG	MYC
FLAG-DYNAMIN 2-125Y-F	pMYC-C2	FLAG	MYC
His-DYNAMIN 2	pET-28a	His	
His-DYNAMIN 2-125Y-E	pET-28a	His	
His-DYNAMIN 2-125Y-F	pET-28a	His	
SNAP25-101Y-A	PEGFP-N1	FLAG	GFP
PCDH-MUNC18-1-WT	PCDH	GFP	
PCDH-MUNC18-1-145Y-A	PCDH	GFP	
PCDH-MUNC18-1-145Y-H	PCDH	GFP	
PCDH-MUNC18-1-145Y-E	PCDH	GFP	
PCDH-MUNC18-1-145Y-F	PCDH	GFP	
His-MUNC18-1-WT	Pet15b	His	
His-MUNC18-1-145Y-H	Pet15b	His	
His-MUNC18-1-145Y-E	Pet15b	His	
His-MUNC18-1-145Y-F	Pet15b	His	
SYNTAXIN1	PEGFP-N1	FLAG	GFP
PCDH-DYNAMIN 2-WT	PCDH	GFP	
PCDH-DYNAMIN 2-125Y-E	PCDH	GFP	
PCDH-DYNAMIN 2-125Y-F	PCDH	GFP	

Plasmid	Primer sequence

His-PTPMEG 2-333Y-A	F:5'GAAACCTAGAGAAAAACCGTGCGGGGGATGTACCCTGCCTGGAC 3' R:5'GTCCAGGCAGGGTACATCCCCCGCACGGTTTTTCTCTAGGTTTC 3'
His-PTPMEG 2-334G-R	F: 5' CTAGAGAAAAACCGTTATCGTGATGTACCCTGCCTGGAC 3' R: 5' GTCCAGGCAGGGTACATCACGATAACGGTTTTTCTCTAG 3'
His-PTPMEG 2-335D-A	F: 5' GAAAAACCGTTATGGGGCGGTACCCTGCCTGGACC 3' R: 5' GGTCCAGGCAGGGTACCGCCCCATAACGGTTTTTC 3'
His-PTPMEG 2-471Y-A	F: 5' AGTTCTTGAGCTGGCCAGACGCGGGTGTCCCTTCCTCA 3' R: 5' TGAGGAAGGGACACCCGCGTCTGGCCAGCTCAAGAACT 3'
His-PTPMEG 2-471Y-F	F: 5' GCTGGCCAGACTTTGGTGTCCCTTC 3' R: 5' GAAGGGACACCAAAGTCTGGCCAGC 3'
His-PTPMEG 2-519I-A	F: 5' ATTGCAGTGCAGGCGCGGGCAGGACAGGT 3' R: 5' ACCTGTCCTGCCCGCGCCTGCACTGCAAT 3'
His-PTPMEG 2-559Q-A	F: 5' AGAGGGCCTTCAGCATCGCGACCCCTGAGCAGTACTA 3' R: 5' TAGTACTGCTCAGGGGTCGCGATGCTGAAGGCCCTCT 3'
PCDH-PTPMEG2	F: 5' CCATAGAAGATTCTAGAATGGAGCCCGCGACC 3' R: 5' CGTCGACTGCAGAATTCTTACAGATCCTCTTC 3'
MUNC18-1-GFP-145Y-H	F:5'TTCTCCCCTATGAGTCCCAGGTGCATTCCCTGGACTCCGCTGACTCT 3' R:5'AGAGTCAGCGGAGTCCAGGGAATGCACCTGGGACTCATAGGGGAGAA 3'
MUNC18-1-GFP-145Y-A	F:5'TCTCCCCTATGAGTCCCAGGTGGCTTCCCTGGACTCCGCTGACTCTT 3' R:5'AAGAGTCAGCGGAGTCCAGGGAAGCCACCTGGGACTCATAGGGGAGA 3'
GST-MUNC18-1-145Y-H	F:5'TCCAAAATCGGATCTGGTTCCGCGTGGATCCATGGATTACAAGGATGACGACGATA 3' R:5'CGAGGCAGATCGTCAGTCAGTCACGATGCGGCCGCTCACTCCATTGTTGGAGCCTGATCCTCAAA 3'
GST-MUNC18-1-145Y-E	F:5'TCTCCCCTATGAGTCCCAGGTGGAGTCCCTGGACTCCGCTGACTCTTTCC 3' R:5'GTCAGCGGAGTCCAGGGACTCCACCTGGGACTCATAGGGGAGAAACGC 3'
GST-MUNC18-1-145Y-F	F:5'TCTCCCCTATGAGTCCCAGGTGTTTTCCCTGGACTCCGCTGACTCTTC 3' F:5'GAAAGAGTCAGCGGAGTCCAGGGAAAACACCTGGGACTCATAGGGG 3'
FLAG-VAMP7-45Y-A	F: 5' CTGAAAATAATAAACTAACTGCCTCACATGGCAATTATTTGT 3' R: 5' ACAAATAATTGCCATGTGAGGCAGTTAGTTTATTATTTTCAG 3'
FLAG-VAMP7-45Y-C	F: 5' CTGAAAATAATAAACTAACTTGCTCACATGGCAATTATTTG 3' R: 5' CAAATAATTGCCATGTGAGCAAGTTAGTTTATTATTTTCAG 3'
FLAG-MYC-DYNAMIN2	F:5'TAGCAAGATCTCGAGCTCAAGCTTCGAATTCGATTACAAGGACGACGATGACAAGATGGGCAACCGCGGGATGGAAGAGCTCATC 3'

	R:5'CGGTGGATCCCGGGCCCGCGGTACCGTCGACCTAGTCGAGCAGGGACGGC 3'
FLAG-MYC-DYNAMIN2--125Y-A	F:5'GGTGCTCGAGTGCGGCCGCAAGCTTGTCGACCTAGTCGAGCAGGGACGGCTC 3' R:5'GAGGGTCAAGTTCAACACGTGTGGTGAGGCGACCCGAAGGTTGATGGGCACAGG 3'
FLAG-MYC-DYNAMIN2-RFP-125Y-E	F:5'TGCCCATCAACCTTCGGGTCGAGTCACCACACGTGTTGAACTTGAC 3' R:5'CAGCTGCCGGTAATCCTTAGCGACATTCCTCTGCTCAGTGTTGAA 3'
FLAG-MYC-DYNAMIN2-RFP-125Y-F	F:5'TGCCCATCAACCTTCGGGTCTTCTCACCACACGTGTTGAACTTGAC 3' R:5'TCAAGTTCAACACGTGTGGTGAGAAGACCCGAAGGTTGATGGGCACAGG 3'
His-DYNAMIN2	F:5'CCCTGTGCCCATCAACCTTCGGGTCTACTCACCACACGTGTTGAACTTGACC 3' R:5'GGTCAAGTTCAACACGTGTGGTGAGTAGACCCGAAGGTTGATGGGCACAGGG 3'
His-DYNAMIN2-125Y-E	F:5'CAGCGGCCTGGTGCCGCGCGGCAGCCATATGGGCAACCGCGGGATGGAAGAGCTC 3' R:5'GGTGCTCGAGTGCGGCCGCAAGCTTGTCGACCTAGTCGAGCAGGGACGGCT 3'
His-DYNAMIN2-125Y-F	F:5'CAGCGGCCTGGTGCCGCGCGGCAGCCATATGGGCAACCGCGGGATGGAAGAGCTCA 3' R:5'GGTGCTCGAGTGCGGCCGCAAGCTTGTCGACCTAGTCGAGCAGGGACGGCTC 3'
SNAP25-101Y-A	F:5'TCGCCACCATGGATTACAAGGATGACGACGATAAGATGGCCGAGGACGCAGACATGCGTA 3' R:5'TACGCATGTCTGCGTCCTCGGCCATCTTATCGTCGTCATCCTTGTAATCCATGGTGGCGA 3'

Q-rtPCR Primer sequence

PTP-MEG2	F:5' AGTTTGATGTGCTCCGTGC 3' R:5' AGAGGGCAATGGAGGCTCC 3'
MUNC18-1	F: 5' TGGAAGGTGCTGGTGGTGGA 3' R: 5' GCAGTCGGCGGGTCCTTAA 3'
DYNAMIN-2	F: 5' TTCGGTGCTCGAGAACTTCG 3' R: 5' CTCTGCTTCGATCTCCTGCC 3'
β -ACTIN	F: 5' AGGGCTATGCTCTCCCTCAC 3' R: 5' CTCTCAGCTGTGGTGGTGAA 3'

Recombinant lentivirus construction and lentivirus infection

For recombinant lentivirus packaging, the construction and infection was carried out as previously reported (Zhang et al., 2018). Plasmids carrying different genes including pCDH-PTP-MEG2-G^{334}R/D^{335}A/Y^{471}A/Y^{471}F/I^{519}A/Q^{559}A/WT-GFP, pCDH-MUNC18-1-Y^{145}H/Y^{145}E/Y^{145}F/WT-GFP and pCDH-DYNAMIN 2-Y^{125}E/Y^{125}F/WT-GFP were transfected into 293T cells using Lipofectamine TM 2000 (Thermo Fisher, Waltham, MA, USA) according to the manufacturer's instructions. Three days after transfection, the supernatant of virus encoding PTP-MEG2, MUNC18-1 or DYNAMIN2 were collected and filtered. The PTP-MEG2, MUNC18-1 or DYNAMIN2 lentivirus (1×10^6 TU/ml) was used to infect the primary chromaffin cells in later experiments.

ELISA

Freshly isolated adrenal medullas from adult female mice (6–8 weeks) were cultured in DMEM medium containing 1% penicillin/streptomycin and 10% FBS. After 2 hours starvation, adrenal medullas were stimulated with high KCl (70mM), AngII (100nM) for 1 min, with or without pre-incubation of a specific PTP-MEG2 inhibitor (400nM) for 2 hours. The supernatants were collected and the epinephrine or norepinephrine secretion were determined by using the Epinephrine or norepinephrine ELISA kit (Shanghai Jianglai Co.,Ltd, JL11194-48T/JL13969-96T) according to the manufacture's protocol.

Quantitative real-time PCR

Freshly isolated adrenal medullas from adult female mice (6–8 weeks). The PTP-MEG2, MUNC18-1 or DYNAMIN 2 lentivirus (1×10^6 TU/ml) was used to infect the primary chromaffin cells. On the third day, about 10 to 20 pheochromoffin cells with the same brightness were found in each plate under the fluorescence microscope. The cells were glued with the tips of ultra-fine glass tubes, and the tips of glass tubes with the cells were directly broken in the PCR tube. The PCR tube was preloaded with 5ul RNA enzyme inhibitors

(Invitrogen, 10777019). 1ul DNA scavenger was added to the PCR tube, which was left at

room temperature for half an hour and then terminated with a terminator. The purpose of this

step is to remove the interference of genomic DNA. All the samples were reverse-transcribed

using the Revertra Ace qPCR RT Kit (TOYOBO FSQ-101) to obtain cDNA. Then the

quantitative real-time PCR was performed with the LightCycler apparatus (Bio-Rad) using

the FastStart Universal SYBR Green Master (Roche).

Electrochemical amperometry

5-mm glass carbon fiber electrode (CFE) was used to measure quantal CA released from the

mouse adrenal medulla chromaffin cell as previously described (Liu et al., 2017). We used a

Multiclamp 700B amplifier (2012, Axon, Molecular Devices, USA) to perform

electrochemical amperometry, which interfaced to Digidata 1440A with the pClamp 10.2

software (Liu et al., 2017). The holding potential of 780 mV was used to record the

amperometric current (Iamp). All experiments were performed at room temperature (20–

25℃). The CFE surface was positioned in contact with the membrane of clean primary

chromaffin cells to monitor the quantal release of the hormone containing catecholamine

substances. In our kinetic analysis of single amperometric spike, we only used the

amperometric spikes with S/N > 3 (signal/noise). The standard external solution for our

amperometry measurement is as follows: 5 mM KCl, 10 mM glucose, 10 mM HEPES pH 7.4,

2 mM $CaCl_2$, 150 mM NaCl and 2 mM $MgCl_2$. We analysed all data using Igor (WaveMetrix,

Lake Oswego, Oregon) and a custom-made macro programme. Statistical data were given as

the mean ± SEM and analyzed with one-way ANOVA.

Electron microscopy

The female mice (6–8 weeks) were decapitated, and the adrenal medullas were freshly

isolated and cut to 150-μm-thick sections. The sections were immersed in Ringer's saline

(125mM NaCl, 2.5mM KCl, 1.25mM NaH_2PO_4, 26mM $NaHCO_3$, 10mM D-Glucose, 2mM

$CaCl_2$, 1mM $MgCl_2$) for 40 minutes at room temperature. During this period, continuous

gases of 5% CO_2 and 95% O_2 were offered to the saline to ensure the survival of the tissue

slice. After 40 minutes of starvation, the sections were stimulated with different conditions

(control; only 100nM Ang II agonists for 1 min; only 400nM PTP-MEG2 inhibitor for 45 min;

100nM AngII agonists and 400nM PTP-MEG2 inhibitor for 1 min or 45 min) at 37°C

respectively. These sections were firstly immersed in precooled 3% glutaraldehyde and fixed

at 4°C for 2 hours, and then rinsed in PBS isotonic buffer, with repeated liquid exchanges and

cleaning overnight, so that the samples were thoroughly rinsed and soaked in the buffer. After

rinsing, the sample was fixed at 4°C with 1% osmium acid for 2 hours. It was rinsed with

isosmotic buffer solution at 0.1M PBS for 15 minutes. The sections were dehydrated with

ethanol at concentrations of 50%, 70%, 90%, then ethanol at concentration of 90% and

acetone at concentration of 90%, at last only acetone at concentrations of 90%, 100%. We

then replaced the acetone with the Epon gradually. The sections were added to Epon and

polymerized at 60°C for 36 hours. Ultra-thin sections were performed at the thickness of

60nm by the LKB-1 ultra microtome, and then the ultra-thin sections were collected with the

single-hole copper ring attached with formvar film. The sample was stained with 2% uranium

acetate for 30 minutes, and then stained with 0.5% lead citrate for 15 minutes. These prepared

samples were examined by JEM-1200EX electron microscope (Japan).

Western

Cells or medulla sections were lysed in lysis buffer (50 mM Tris pH 8.0, 150 mM NaCl, 1

mM NaF, 1% NP-40, 2mM EDTA, Tris-HCl pH 8.0, 10% glycerol, 0.25% sodium

deoxycholate, 1mM Na_3VO_4, 0.3μM aprotinin, 130μM bestatin, 1μM leupeptin, 1μM

repstatin and 0.5% IAA) after rinsing with the pre-chilled PBS on ice. Cell and tissue

lysates were kept on ice for 35 min and then sediment via centrifugation for 15 min at 4 °C.

The whole-cell and tissue protein lysates samples (30μg) were prepared for SDS-PAGE. The

proteins in the gel were ransferred to a nitrocellulose filter membrane by electro blotting, then

probed with the appropriate primary and secondary antibodies. Antibody binding was

detected by an HRP system.

Immunofluorescence.

For the acquisition of tissue cells used in immunofluorescence, the mice were decapitated,

and the the adrenal medullas were freshly isolated (female mice, 6–8 weeks).The isolated

adrenal medullas was immersed in 4% paraformaldehyde for fixation overnight at 4°C. Then

the fixed tissues were washed for 4 hours in PBS containing 10% sucrose at 4°C for 8 hours

in 20% sucrose, and in 30% sucrose overnight. Then these adrenal medullas were imbedded

in Tissue-Tek OCT compound and then mounted and frozen them at -25°C. Subsequently, the

adrenal medulla were cut to 4-μm-thick coronal serial sections. The adrenal medullas sections

were blocked with 1% (vol/vol) donkey serum, 2.5% (wt/vol) BSA and 0.1% (vol/vol) Triton

X-100 in PBS for 1.5 h. Then, the slides were incubated with primary antibodies against

PTP-MEG2 (1:100), NSF (1:50), MUNC18-1 (1:50), VAMP7 (1:50), DYNAMIN 2 (1:100),

SNAP25 (1:100), PACSIN1 (1:50) at 4°C overnight. After washing with PBS for 3 times, the

slides were incubated with the secondary antibody (1:500) for 1 h at room temperature. The

slides were stained with DAPI (1:2000). Images were captured using a confocal microscope

(ZEISS, LSM780).　　The Pearson's co-localization coefficients were analyzed with

Image-Pro Plus.

K_m and k_{cat} measurements.

Enzymatic activity measurement was carried out as previously reported (Li et al., 2016; Wang

et al., 2014). The standard solution (DMG buffer) for our enzymatic reactions is following: 50

mM 3, 3-dimethyl glutarate pH 7.0, 1 mM EDTA, 1 mM DTT. The ionic strength was

maintained at 0.15 M (adjusted by NaCl). For the pNPP activity measurement, 100 μl reaction

mixtures were set up in a total volume in a 96-well polystyrene plate (Thermo Fisher

Scientific, Waltham, MA, US). The substrate concentration ranging from 0.2 to 5 K_m was used

to determine the k_{cat} and K_m values. Reactions were started by the addition of an appropriate

amount of His-PTP-MEG2-CD-WT or corresponding mutants, such as $Y^{333}A$, $G^{334}R$, $D^{335}A$,

$Y^{471}A$, $Y^{471}F$, $I^{519}A$, $R^{409}A$, and $R^{410}A$. The dephosphorylation of pNPP was terminated by

adding 120μl 1M NaOH, and the enzymatic activity was monitored by measuring the

absorbance at 405 nm. The activities toward phospho-peptide segment derived from NSF or

MUNC18-1 were measured as following: In the first column, 90μl diluted

NSF/MUNC18-1/DYNAMIN 2 phospho-peptide substrate (100μM) was added. The

successive columns were diluted by 1.5 times. The phospho-peptide substrates were

preincubated at 37°C for 5 min. Reactions were started by the addition of an appropriate

amount of enzymes. The dephosphorylation of NSF/MUNC18-1/DYNAMIN2 was

terminated by adding 120μl Biomol green, and the enzymatic activities were monitored by

measuring the absorbance at 620 nm. The steady-state kinetic parameters were determined

from a direct fit of the data to the Michaelis-Menten equation using GraphPad Prism 5.0.

GST-pull down

To screen the candidate proteins interacting with PTP-MEG2, the GST beads were washed

five times by cold binding buffer (20mM HEPES pH7.5, 1mM DTT, 1mM EDTA, and

100mM NaCl) and incubated with 5μg purified GST-PTP-MEG2-CD-D^{470}A protein for 2

hours at 4°C. PC12 cells were transfected with FLAG-SYN1-GFP, FLAG-MUNC18-1-GFP,

FLAG-MUNC18-1-Y^{145}A-GFP,   FLAG-MUNC18-1-Y^{145}H-GFP   ,

FLAG-MUNC18-1-Y^{145}E-GFP, FLAG-MUNC18-1-Y^{145}F-GFP, FLAG-MUNC18-3-GFP,

FLAG-PACSIN1-GFP, FLAG-SCAMP1-GFP, FLAG-PPP3CA-GFP, FLAG-STX17-GFP,

FLAG-VAMP7,   FLAG-VAMP7-Y^{45}A,   FLAG-VAMP7-Y^{45}C,   FLAG-SYT7-GFP,

FLAG-SYT11-GFP, FLAG-SNAP25-GFP, FLAG-SNAP25-Y^{101}A-GFP, FLAG-DYNAMIN

1-GFP, FLAG-DYNAMIN 2, FLAG-DYNAMIN 2-Y^{125}A, FLAG-DYNAMIN 2-Y^{125}E,

FLAG-DYNAMIN 2-Y^{125}F. After stimulation with 100nM AngII for 5 minutes at 37 °C, the

cells were washed and then lysed in lysis buffer [20 mM HEPES pH 7.5, 100 mM NaCl, 0.5%

NP-40, 5 mM iodoacetic acid, and a protease inhibitor mixture (final concentrations, 10μg of

leupeptin, 1μg of aprotinin, 1μg of pepstatin, 1μg of antipain, and 20μg of

phenylmethylsulfonyl fluoride per ml), 10mM DTT, 2mM EDTA] on ice for 30 minutes, then

centrifuged at 12000rpm for 15 minutes at 4 °C. 20μl GST beads-PTP-MEG2-D^{470}A protein

was added into 500μl supernatants and the mixtures were subjected to end-to-end rotation at

4 °C for 2 hours. The GST beads and their binding proteins were washed five times with cold

binding buffer to exclude the unspecific binding proteins.

The pull down experiment of MUNC18-1 and SYNTAXIN1 were described similar to above

description. The GST beads were washed five times by cold binding buffer (described as

above). After that, 5μg purified GST-MUNC18-1-WT or its Y^{145}H, Y^{145}E or Y^{145}F mutant

proteins were added into 20μl GST-agarose, and incubated at for 4 °C 2 hours with end to end

rotation. PC12 cells transfected with FLAG-SYNTAXIN1 were lysed in lysis buffer

(described as above) and centrifuged to remove the pallets. The supernatants were added with

20μl GST beads/GST-fusion protein and the mixtures were subjected to end-to-end rotation at

4 °C for 2 hours. The FLAG-SYNTAXIN1 was detected with FLAG antibody.

GFP-pull down

PC12 cells were infected with lentiviruses encoding GFP-tagged PTP-MEG2-D^{470}A of wild type

and different mutant types. After stimulation with 100nM AngII for 5 minutes at 37 °C, the

cells were washed and then lysed in lysis buffer [20 mM HEPES pH 7.5, 100 mM NaCl, 0.5%

NP-40, 5 mM iodoacetic acid, and a protease inhibitor mixture (final concentrations, 10μg of

leupeptin, 1μg of aprotinin, 1μg of pepstatin, 1μg of antipain, and 20μg of

phenylmethylsulfonyl fluoride per ml), 10mM DTT, 2mM EDTA] on ice for 30 minutes, then

centrifuged at 12000rpm for 15 minutes at 4 °C. The GFP beads were washed five times by

cold binding buffer (20mM HEPES pH7.5, 1mM DTT, 1mM EDTA, and 100mM NaCl). 25μl

anti-GFP-antibody conjugated beads were added into 500μl supernatants and the mixtures

were subjected to end-to-end rotation at 4 °C for 1 hours. The PTP-MEG2-D^{470}A-GFP

proteins were pulled down and washed five times with cold binding buffer to exclude the

unspecific binding proteins. The associated PTP-MEG2 substrates in the GFP-pull-down

beads were then examined by western blott.

Immunoprecipitation and in vitro dephosphorylation

Rat adrenal medullas were isolated and cut into pieces in D-hanks buffer, and stimulated with

70mM KCl or 100nM AngII for 5 minutes. The tissues were then lysed and grinded on ice in

lysis buffer supplemented with proteinase inhibitor as described before. The lysates were

centrifuged at 12000 rpm for 20 minutes at 4 °C and the pallets were removed. Before

incubation with the lysates in the later step, the 1μg primary antibodies of NSF, MUNC18-1,

VAMP7, SNAP25 or DYNAMIN 2 were incubated with 20μl Protein A/G beads by

end-to-end rotation overnight at 4 °C. Then the supernatants were incubated with primary

antibody pre-coated with Protein A/G beads and rotated for 2 hours. 5μg purified

PTP-MEG2-WT protein or control solution were added into the lysates for the in vitro

dephosphorylation. A pan-phospho-tyrosine antibody pY[20]/ pY[99] was used in Western blotting

to detect the phosphorylated tyrosine of NSF/MUNC18-1/VAMP7/SNAP25/DYNAMIN 2 in

adrenal medullar with or without incubation with the PTP-MEG2.

Protein Expression and Purification

The wide type and mutant proteins of His-tagged PTP-MEG2-catalytic domain were

expressed in BL21-DE3 Escherichia coli as previously described (Pan et al., 2013). In brief,

0.4 mM isopropyl1-thio-D-galactopyranoside (IPTG) was used to induce the expression of

His-PTP-MEG2, and the bacteria lysates were centrifuged at 12000 rpm for 1 hour. Ni-NTA

Agrose was applied to bind the His-tagged PTP-MEG2 and an imidazole gradation was used

to elute the binding proteins. His-PTP-MEG2 was then purified with gel filtration

chromatography to achieve at least 95% purity. The wide type and mutant proteins of

GST-PTP-MEG2 and GST-MUNC18-1 were also expression in E.coli in presence of 0.4 mM

IPTG for 16 hours at 25°C. After centrifugation and lysis, the proteins were purified by

binding with GST-Sepharose for 2 hours and eluted by GSH.

Crystallization and Data Collection

For crystallization, His-PTP-MEG2-C[515]A/D[470]A protein (concentration at 15mg/ml) was

mixed with NSF-pY[83] peptide (EVSLpYTFDK) or MUNC18-1-pY[145] peptide

(ESQVpYSLDS) with molar ratio as 1:3 in buffer A (pH 7.2, 20 mM Hepes, 350 mM NaCl,

and 1 mM DTT). 1μl mixed protein was blended with 1μl buffer B (pH 6.4, 20% PEG 4000,

0.2M KSCN, 10% ethelene glycol, 0.1M bis-tris propane) at 4°C for 3 days before crystals

appears. The cubic crystals were preserved in liquid nitrogen very quickly dipped in storage

buffer (buffer B supplemented with 10% glycerol). The data were collected at Shanghai

Synchrotron Radiation Facility beamline BL17U1 using 0.98Å X-ray wave length and

analyzed by HKL2000.

Structural determination and refinement

The crystals of PTP-MEG2-NSF-pY[83] and PTP-MEG2-MUNC18-1-pY[145] peptides belong to

the $P2_12_12$ space group. In each asymmetric unit, both PTPMEG2-NSF-pY[83] and

PTPMEG2-MUNC18-1-pY[145] contain one monomer in one unit. Molecular replacement with

Phaser in the CCP4 software package, with PTP-MEG2 catalytic domain (PDB code: 2PA5,

water deleted) as the initial search model. Further refinements were carried out using the

PHENIX program with iterative manual building in COOT. The data of the final refined

structures are shown in Supplementary information Table 1.

BRET assay

The BRET experiment was performed to monitor the mutation effects of DYNAMIN2 on

AT1aR endocytosis in response to AngII (1μM) stimulation as previously described. Plasmids

encoding DYNAMIN2-WT, DYNAMIN2-Y[125]E or DYNAMIN2-Y[125]F and LYN-YFP,

AT1aR-C-RLUC were co-transferred into HEK293 cells at 1:1:1 for 48 h. After 12 h of

starvation, cells were digested with trypsin and harvested. Then the transfected cells were

washed with PBS at least three times and evenly distributed into a 96-well plate (Corning

Costar Cat. # 3912). Then cells were stimulated with AngII (1μM) for 25 min at 37 °C.

Afterwords, we incubated the cells with Coelenterazine h at 25 °C (Promega S2011, final

concentration, 5μM). We used two different light emissions for measurement (530/20 nm for

yellow fluorescent protein and 480/20 nm for luciferase).

DYNAMIN 2 GTPase activity detection

To prepare His-tagged DYNAMIN-2-WT, DYNAMIN-2-Y[125]E, DYNAMIN-2-Y[125]F (His-

DYNAMIN-2-WT, His-DYNAMIN-2-Y[125]E, His-DYNAMIN-2-Y[125]F), DNA fragments

were generated by PCR using MYC-DYNAMIN-2 as the template and subcloned into

pET-28a (+) plasmid. The resulting plasmids were transformed into bacterial

BL21-DE3-RIPL cells for protein expression. We measured the DYNAMIN-2 GTPase

activity by using GTPase-Glo assay test kit (V7681) from Promega Corporation. The three

DYNAMIN2 enzymes were diluted to 500ng with GTPase/GAP Buffer of the kit, and

dispense 12.5μl into each well of a 96-well plate (Corning Costar Cat. # 3912). Then Prepare

a 2X GTP solution containing 10μM GTP and 1mM DTT in GTPase/GAP Buffer, initiate the

GTPase reaction by adding 12.5μl of the 2X GTP solution prepared to each well. The total

GTPase reaction volume is 25µl. Incubate the reaction with shaking at room temperature

(25°C) for 90 minutes. Configuration 1ml the Reconstituted GTPase-Glo™ solution with 2µl

GTPase-Glo™ Reagent (500X), 0.5µl 10mM ADP, 998µl GTPase-Glo™ Buffer. Add 25µl of

reconstituted GTPase-Glo™ Reagent to the completed GTPase reaction, mix briefly and

incubate with shaking for 30 minutes at room temperature (25°C). Add 25µl of Detection

Reagent, and incubate the plate for 7 minutes at room temperature (25°C). Measure

luminescence.

Bioinformatic search of PTP-protein interactions

We identified the PTP-protein interactions by two independent bioinformatic analyses. On

one site, we extracted the potential PTP-interacting proteins from the STRING database

(Szklarczyk et al., 2015) by setting the parameters of Homo sapiens. On the other site, we

used Pubtator (Wei et al., 2013) to text-mine potential PTP-protein associations from PubMed

literature (by 2018.9) via the search of combinational keywords such as "tyrosine

phosphorylation mutation, vesicular, fusion, and human". We compared these two protein lists

to produce a consensus PTP-interacting protein list. Subsequently, we narrowed down the

protein list by satisfying following constraints: (a) The protein should express in the adrenal

gland as documented in the TissueAtlas (Thul et al., 2017); (b) The protein should expose

tyrosine residue(s) on surface for phosphorylation as simulated by the Molecular Operating

Environment (MOE) (Vilar et al., 2008); (c) On the protein, at least one predicted tyrosine

phosphorylation site predicted by the PhosphoSitePlus (Hornbeck et al., 2015) is also

experiment-validated. The refined PTP-protein pairs were then ready for later experimental

analyses.

Statistical analysis

All data are presented as mean ± SEM. The statistical significance between different groups

was compared with ANOVA tests, and the statistical significance between two different

groups was generated by ANOVA tests. All of the Western films were scanned, and band

intensity was quantified with ImageJ software (National Institutes of Health, Bethesda MD). P

< 0.05 was considered as statistically significant.

DATA AVAILABILITY

The coordinates and density map for the PTP-MEG2-NSF-pY[83] (PDB ID: 6KZQ) and

PTP-MEG2-MUNC18-1-pY[145] (PDB ID: 6L03) peptides has been deposited in protein data

bank. All other data are available upon request to the corresponding authors.

REFERENCE

Alonso, A., Sasin, J., Bottini, N., Friedberg, I., Friedberg, I., Osterman, A., Godzik, A., Hunter, T., Dixon, J., and Mustelin, T. (2004). Protein tyrosine phosphatases in the human genome. Cell *117*, 699-711.

Alvarez de Toledo, G., Fernandez-Chacon, R., and Fernandez, J.M. (1993). Release of secretory products during transient vesicle fusion. Nature *363*, 554-558.

Baker, R.W., and Hughson, F.M. (2016). Chaperoning SNARE assembly and disassembly. Nat Rev Mol Cell Biol *17*, 465-479.

Ballif, B.A., Carey, G.R., Sunyaev, S.R., and Gygi, S.P. (2008). Large-scale identification and evolution indexing of tyrosine phosphorylation sites from murine brain. Journal of proteome research *7*, 311-318.

Barr, A.J., Ugochukwu, E., Lee, W.H., King, O.N., Filippakopoulos, P., Alfano, I., Savitsky, P., Burgess-Brown, N.A., Muller, S., and Knapp, S. (2009). Large-scale structural analysis of the classical human protein tyrosine phosphatome. Cell *136*, 352-363.

Chai, Y.J., Sierecki, E., Tomatis, V.M., Gormal, R.S., Giles, N., Morrow, I.C., Xia, D., Gotz, J., Parton, R.G., Collins, B.M.*, et al.* (2016). Munc18-1 is a molecular chaperone for alpha-synuclein, controlling its self-replicating aggregation. J Cell Biol *214*, 705-718.

Chen, X.K., Wang, L.C., Zhou, Y., Cai, Q., Prakriya, M., Duan, K.L., Sheng, Z.H., Lingle, C., and Zhou, Z. (2005). Activation of GPCRs modulates quantal size in chromaffin cells through G(betagamma) and PKC. Nat Neurosci *8*, 1160-1168.

Cho, C.Y., Koo, S.H., Wang, Y., Callaway, S., Hedrick, S., Mak, P.A., Orth, A.P., Peters, E.C., Saez, E., Montminy, M.*, et al.* (2006). Identification of the tyrosine phosphatase PTP-MEG2 as an antagonist of hepatic insulin signaling. Cell Metab *3*, 367-378.

Cijsouw, T., Weber, J.P., Broeke, J.H., Broek, J.A., Schut, D., Kroon, T., Saarloos, I., Verhage, M., and Toonen, R.F. (2014). Munc18-1 redistributes in nerve terminals in an activity- and PKC-dependent manner. J Cell Biol *204*, 759-775.

Dittman, J.S., and Ryan, T.A. (2019). The control of release probability at nerve terminals. Nat Rev

Neurosci *20*, 177-186.

Doody, K.M., and Bottini, N. (2014). "PEST control": regulation of molecular barcodes by tyrosine

phosphatases. Cell Res *24*, 1027-1028.

Eitzen, G. (2003). Actin remodeling to facilitate membrane fusion. Biochimica et biophysica acta *1641*,

175-181.

Fisher, R.J., Pevsner, J., and Burgoyne, R.D. (2001). Control of fusion pore dynamics during exocytosis

by Munc18. Science *291*, 875-878.

Gabel, M., Delavoie, F., Royer, C., Tahouly, T., Gasman, S., Bader, M.F., Vitale, N., and Chasserot-Golaz,

S. (2019). Phosphorylation cycling of Annexin A2 Tyr23 is critical for calcium-regulated exocytosis in

neuroendocrine cells. Biochimica et biophysica acta Molecular cell research *1866*, 1207-1217.

Gaisano, H.Y. (2017). Recent new insights into the role of SNARE and associated proteins in insulin

granule exocytosis. Diabetes Obes Metab *19 Suppl 1*, 115-123.

Gu, M., Warshawsky, I., and Majerus, P.W. (1992). Cloning and expression of a cytosolic megakaryocyte

protein-tyrosine-phosphatase with sequence homology to retinaldehyde-binding protein and yeast

SEC14p. Proc Natl Acad Sci U S A *89*, 2980-2984.

Gulyas-Kovacs, A., de Wit, H., Milosevic, I., Kochubey, O., Toonen, R., Klingauf, J., Verhage, M., and

Sorensen, J.B. (2007). Munc18-1: sequential interactions with the fusion machinery stimulate vesicle

docking and priming. J Neurosci *27*, 8676-8686.

Harada, K., Matsuoka, H., Miyata, H., Matsui, M., and Inoue, M. (2015). Identification of muscarinic

receptor subtypes involved in catecholamine secretion in adrenal medullary chromaffin cells by

genetic deletion. Br J Pharmacol *172*, 1348-1359.

He, E., Wierda, K., van Westen, R., Broeke, J.H., Toonen, R.F., Cornelisse, L.N., and Verhage, M. (2017).

Munc13-1 and Munc18-1 together prevent NSF-dependent de-priming of synaptic vesicles. Nat

Commun *8*, 15915.

Hernandez, J.M., Stein, A., Behrmann, E., Riedel, D., Cypionka, A., Farsi, Z., Walla, P.J., Raunser, S., and

Jahn, R. (2012). Membrane fusion intermediates via directional and full assembly of the SNARE

complex. Science *336*, 1581-1584.

Hong, W., and Lev, S. (2014). Tethering the assembly of SNARE complexes. Trends Cell Biol *24*, 35-43.

Hornbeck, P.V., Zhang, B., Murray, B., Kornhauser, J.M., Latham, V., and Skrzypek, E. (2015).

PhosphoSitePlus, 2014: mutations, PTMs and recalibrations. Nucleic Acids Res *43*, D512-520.

Hu, S.H., Christie, M.P., Saez, N.J., Latham, C.F., Jarrott, R., Lua, L.H., Collins, B.M., and Martin, J.L.

(2011). Possible roles for Munc18-1 domain 3a and Syntaxin1 N-peptide and C-terminal anchor in

SNARE complex formation. Proc Natl Acad Sci U S A *108*, 1040-1045.

Huynh, H., Bottini, N., Williams, S., Cherepanov, V., Musumeci, L., Saito, K., Bruckner, S., Vachon, E.,

Wang, X., Kruger, J.*, et al.* (2004). Control of vesicle fusion by a tyrosine phosphatase. Nat Cell Biol *6*,

831-839.

Jewell, J.L., Oh, E., Ramalingam, L., Kalwat, M.A., Tagliabracci, V.S., Tackett, L., Elmendorf, J.S., and

Thurmond, D.C. (2011). Munc18c phosphorylation by the insulin receptor links cell signaling directly to

SNARE exocytosis. The Journal of cell biology *193*, 185-199.

Jones, D.M., Alvarez, L.A., Nolan, R., Ferriz, M., Sainz Urruela, R., Massana-Munoz, X., Novak-Kotzer, H.,

Dustin, M.L., and Padilla-Parra, S. (2017). Dynamin-2 Stabilizes the HIV-1 Fusion Pore with a Low

Oligomeric State. Cell reports *18*, 443-453.

Kar, U.P., Dey, H., and Rahaman, A. (2017). Regulation of dynamin family proteins by post-translational

modifications. Journal of biosciences *42*, 333-344.

Korteweg, N., Maia, A.S., Thompson, B., Roubos, E.W., Burbach, J.P., and Verhage, M. (2005). The role
of Munc18-1 in docking and exocytosis of peptide hormone vesicles in the anterior pituitary. Biol Cell
97, 445-455.

Laidlaw, K.M.E., Livingstone, R., Al-Tobi, M., Bryant, N.J., and Gould, G.W. (2017). SNARE
phosphorylation: a control mechanism for insulin-stimulated glucose transport and other regulated
exocytic events. Biochem Soc Trans *45*, 1271-1277.

Li, H., Yang, F., Liu, C., Xiao, P., Xu, Y., Liang, Z., Liu, C., Wang, H., Wang, W., Zheng, W.*, et al.* (2016).
Crystal Structure and Substrate Specificity of PTPN12. Cell Rep *15*, 1345-1358.

Lim, S.H., Moon, J., Lee, M., and Lee, J.R. (2013). PTPRT regulates the interaction of Syntaxin-binding
protein 1 with Syntaxin 1 through dephosphorylation of specific tyrosine residue. Biochemical and
biophysical research communications *439*, 40-46.

Liu, C.H., Gong, Z., Liang, Z.L., Liu, Z.X., Yang, F., Sun, Y.J., Ma, M.L., Wang, Y.J., Ji, C.R., Wang, Y.H.*, et al.*
(2017). Arrestin-biased AT1R agonism induces acute catecholamine secretion through TRPC3 coupling.
Nature communications *8*, 14335.

Ma, C., Su, L., Seven, A.B., Xu, Y., and Rizo, J. (2013). Reconstitution of the vital functions of Munc18
and Munc13 in neurotransmitter release. Science *339*, 421-425.

Ma, L., Rebane, A.A., Yang, G., Xi, Z., Kang, Y., Gao, Y., and Zhang, Y. (2015). Munc18-1-regulated
stage-wise SNARE assembly underlying synaptic exocytosis. Elife *4*.

Magadmi, R., Meszaros, J., Damanhouri, Z.A., and Seward, E.P. (2019). Secretion of Mast Cell
Inflammatory Mediators Is Enhanced by CADM1-Dependent Adhesion to Sensory Neurons. Front Cell
Neurosci *13*, 262.

Mallozzi, C., D'Amore, C., Camerini, S., Macchia, G., Crescenzi, M., Petrucci, T.C., and Di Stasi, A.M.
(2013). Phosphorylation and nitration of tyrosine residues affect functional properties of
Synaptophysin and Dynamin I, two proteins involved in exo-endocytosis of synaptic vesicles.
Biochimica et biophysica acta *1833*, 110-121.

Mattila, J.P., Shnyrova, A.V., Sundborger, A.C., Hortelano, E.R., Fuhrmans, M., Neumann, S., Muller, M.,
Hinshaw, J.E., Schmid, S.L., and Frolov, V.A. (2015). A hemi-fission intermediate links two
mechanistically distinct stages of membrane fission. Nature *524*, 109-113.

Meijer, M., Dorr, B., Lammertse, H.C., Blithikioti, C., van Weering, J.R., Toonen, R.F., Sollner, T.H., and
Verhage, M. (2018). Tyrosine phosphorylation of Munc18-1 inhibits synaptic transmission by
preventing SNARE assembly. EMBO J *37*, 300-320.

Neher, E., and Brose, N. (2018). Dynamically Primed Synaptic Vesicle States: Key to Understand
Synaptic Short-Term Plasticity. Neuron *100*, 1283-1291.

Oakie, A., and Wang, R. (2018). beta-Cell Receptor Tyrosine Kinases in Controlling Insulin Secretion
and Exocytotic Machinery: c-Kit and Insulin Receptor. Endocrinology *159*, 3813-3821.

Ortsater, H., Grankvist, N., Honkanen, R.E., and Sjoholm, A. (2014). Protein phosphatases in pancreatic
islets. The Journal of endocrinology *221*, R121-144.

Pan, C., Liu, H.D., Gong, Z., Yu, X., Hou, X.B., Xie, D.D., Zhu, X.B., Li, H.W., Tang, J.Y., Xu, Y.F.*, et al.* (2013).
Cadmium is a potent inhibitor of PPM phosphatases and targets the M1 binding site. Scientific reports
3, 2333.

Seino, S., Takahashi, H., Fujimoto, W., and Shibasaki, T. (2009). Roles of cAMP signalling in insulin
granule exocytosis. Diabetes Obes Metab *11 Suppl 4*, 180-188.

Shin, W., Ge, L., Arpino, G., Villarreal, S.A., Hamid, E., Liu, H., Zhao, W.D., Wen, P.J., Chiang, H.C., and
Wu, L.G. (2018). Visualization of Membrane Pore in Live Cells Reveals a Dynamic-Pore Theory

Governing Fusion and Endocytosis. Cell *173*, 934-945 e912.

Sitarska, E., Xu, J., Park, S., Liu, X., Quade, B., Stepien, K., Sugita, K., Brautigam, C.A., Sugita, S., and Rizo, J. (2017). Autoinhibition of Munc18-1 modulates synaptobrevin binding and helps to enable Munc13-dependent regulation of membrane fusion. Elife *6*.

Soares, H., Henriques, R., Sachse, M., Ventimiglia, L., Alonso, M.A., Zimmer, C., Thoulouze, M.I., and Alcover, A. (2013). Regulated vesicle fusion generates signaling nanoterritories that control T cell activation at the immunological synapse. J Exp Med *210*, 2415-2433.

Stamberger, H., Weckhuysen, S., and De Jonghe, P. (2017). STXBP1 as a therapeutic target for epileptic encephalopathy. Expert Opin Ther Targets *21*, 1027-1036.

Sudhof, T.C. (2004). The synaptic vesicle cycle. Annual review of neuroscience *27*, 509-547.

Sudhof, T.C. (2013). Neurotransmitter release: the last millisecond in the life of a synaptic vesicle. Neuron *80*, 675-690.

Sudhof, T.C., and Rothman, J.E. (2009). Membrane fusion: grappling with SNARE and SM proteins. Science *323*, 474-477.

Szklarczyk, D., Franceschini, A., Wyder, S., Forslund, K., Heller, D., Huerta-Cepas, J., Simonovic, M., Roth, A., Santos, A., Tsafou, K.P.*, et al.* (2015). STRING v10: protein-protein interaction networks, integrated over the tree of life. Nucleic Acids Res *43*, D447-452.

Teschemacher, A.G., and Seward, E.P. (2000). Bidirectional modulation of exocytosis by angiotensin II involves multiple G-protein-regulated transduction pathways in chromaffin cells. J Neurosci *20*, 4776-4785.

Thul, P.J., Akesson, L., Wiking, M., Mahdessian, D., Geladaki, A., Ait Blal, H., Alm, T., Asplund, A., Bjork, L., Breckels, L.M.*, et al.* (2017). A subcellular map of the human proteome. Science *356*.

Tonks, N.K. (2006). Protein tyrosine phosphatases: from genes, to function, to disease. Nature reviews Molecular cell biology *7*, 833-846.

Umahara, M., Okada, S., Yamada, E., Saito, T., Ohshima, K., Hashimoto, K., Yamada, M., Shimizu, H., Pessin, J.E., and Mori, M. (2008). Tyrosine phosphorylation of Munc18c regulates platelet-derived growth factor-stimulated glucose transporter 4 translocation in 3T3L1 adipocytes. Endocrinology *149*, 40-49.

Vilar, S., Cozza, G., and Moro, S. (2008). Medicinal chemistry and the molecular operating environment (MOE): application of QSAR and molecular docking to drug discovery. Curr Top Med Chem *8*, 1555-1572.

Wang, H.M., Xu, Y.F., Ning, S.L., Yang, D.X., Li, Y., Du, Y.J., Yang, F., Zhang, Y., Liang, N., Yao, W.*, et al.* (2014). The catalytic region and PEST domain of PTPN18 distinctly regulate the HER2 phosphorylation and ubiquitination barcodes. Cell Res *24*, 1067-1090.

Wang, Y., Vachon, E., Zhang, J., Cherepanov, V., Kruger, J., Li, J., Saito, K., Shannon, P., Bottini, N., Huynh, H.*, et al.* (2005). Tyrosine phosphatase MEG2 modulates murine development and platelet and lymphocyte activation through secretory vesicle function. J Exp Med *202*, 1587-1597.

Wei, C.H., Kao, H.Y., and Lu, Z. (2013). PubTator: a web-based text mining tool for assisting biocuration. Nucleic Acids Res *41*, W518-522.

Wu, L.G., Hamid, E., Shin, W., and Chiang, H.C. (2014). Exocytosis and endocytosis: modes, functions, and coupling mechanisms. Annu Rev Physiol *76*, 301-331.

Yang, X., Wang, S., Sheng, Y., Zhang, M., Zou, W., Wu, L., Kang, L., Rizo, J., Zhang, R., Xu, T.*, et al.* (2015). Syntaxin opening by the MUN domain underlies the function of Munc13 in synaptic-vesicle priming. Nat Struct Mol Biol *22*, 547-554.

Yu, X., Chen, M., Zhang, S., Yu, Z.H., Sun, J.P., Wang, L., Liu, S., Imasaki, T., Takagi, Y., and Zhang, Z.Y.
(2011). Substrate specificity of lymphoid-specific tyrosine phosphatase (Lyp) and identification of Src
kinase-associated protein of 55 kDa homolog (SKAP-HOM) as a Lyp substrate. The Journal of biological
chemistry *286*, 30526-30534.

Yu, Z.H., and Zhang, Z.Y. (2018). Regulatory Mechanisms and Novel Therapeutic Targeting Strategies for
Protein Tyrosine Phosphatases. Chemical reviews *118*, 1069-1091.

Zhang, D., Marlin, M.C., Liang, Z., Ahmad, M., Ashpole, N.M., Sonntag, W.E., Zhao, Z.J., and Li, G.
(2016). The Protein Tyrosine Phosphatase MEG2 Regulates the Transport and Signal Transduction of
Tropomyosin Receptor Kinase A. J Biol Chem *291*, 23895-23905.

Zhang, D.L., Sun, Y.J., Ma, M.L., Wang, Y.J., Lin, H., Li, R.R., Liang, Z.L., Gao, Y., Yang, Z., He, D.F.*, et al.*
(2018). Gq activity- and beta-arrestin-1 scaffolding-mediated ADGRG2/CFTR coupling are required for
male fertility. Elife *7*.

Zhang, S., Liu, S., Tao, R., Wei, D., Chen, L., Shen, W., Yu, Z.H., Wang, L., Jones, D.R., Dong, X.C.*, et al.*
(2012). A highly selective and potent PTP-MEG2 inhibitor with therapeutic potential for type 2
diabetes. J Am Chem Soc *134*, 18116-18124.

Zhao, W.D., Hamid, E., Shin, W., Wen, P.J., Krystofiak, E.S., Villarreal, S.A., Chiang, H.C., Kachar, B., and
Wu, L.G. (2016). Hemi-fused structure mediates and controls fusion and fission in live cells. Nature
534, 548-552.

Zhou, Z., Misler, S., and Chow, R.H. (1996). Rapid fluctuations in transmitter release from single
vesicles in bovine adrenal chromaffin cells. Biophysical journal *70*, 1543-1552.

1103
1104
1105 **ACKNOWLEDGEMENTS**
1106

We thank Dr Michael Xi Zhu for stimulating discussions and critical reading of the
manuscript. We thank Yanmei Lu from Shandong jiaotong hospital, for her help with
Transmission electron microscopy analysis. We thank Daolai Zhang and Mingliang Ma for
their technical assistance in lentivirus packaging. We thank Yujing Sun and Zhixin Liu for
their technical assistance in electrochemical recording. We acknowledge support from the
National Key Basic Research Program of China Grant 2018YFC1003600 (to X.Y. and J.-P.S.),
the National Natural Science Foundation of China Grant 81773704 (to J.-P.S.) and Grant
81601668 (to Y.-F.X.), the National Science Fund for Distinguished Young Scholars Grant
81825022 (to J.-P.S.), the National Science Fund for Excellent Young Scholars Grant
81822008 (to X. Y.) and the Rolling program of ChangJiang Scholars and Innovative
Research Team in University Grant IRT_17R68 (to Y. S.). S.Z. and Z.-Y. Z. are supported by
NIH RO1 CA69202.

1119
1120
AUTHOR CONTRIBUTIONS:
J.-P.S. and X.Y. conceived the whole research and initiated the project. J.-P.S., X.Y.,

Y.-F.X. and X.C. designed all of the experiments. J.-P.S. and X.Y. supervised the overall

project design and execution. X.C., Y.-F.X., Z.Y., X.Y. and J.-P.S. participated in data

analysis and interpretation. C.-H.L. and Y.-J.W. performed electrochemical experiments. Z.Y.,

P.X., Z.-L.Z. helped us collect crystal data and analyze crystal structure. M.C., K.-S.L.

performed cell biology, molecular biology and biochemistry experiments. Z.-G.X, provided

DYNAMIN2 plasmid. Z.-Y.Z. and S.Z. synthesized and purified Compound 7 (PTP-MEG2

inhibitor). X.-Z.Y. and Z.-L.J. performed substrate bioinformatics screening. Z.-Y.Z.,

W.-D.Z., C.-H.W. and C.W. provided insightful idea and experimental designs. J.-P.S,

Y.-F.X., X.C. and X.Y. wrote the manuscript. All of the authors have seen and commented on

the manuscript.

1133

1134

1135 **ADDITIONAL INFORMATION**

1136 **Competing financial interests**   The authors declare no competing financial interests.

1137

1138

1139 **FIGURE LEGENDS**

1140

Figure 1. Inhibition of PTP-MEG2 reduced the spike amplitude and 'foot' probability of

catecholamine secretion.

(A-D). The epinephrine (A-B) or norepinephrine (C-D) secreted from the adrenal medulla

was measured after stimulation with high KCl (70 mM) (A,C) or Angiotensin II (AngII, 100

nM) (B,D) for 1 min, with or without pre-incubation with a specific PTP-MEG2 inhibitor

(400 nM) for 1 hours. Data were obtained from 6 independent experiments and displayed as

the mean ± SEM. ***, P<0.001: cells stimulated with KCl or AngII compared with

un-stimulated cells. #, P<0.05; ##, P<0.01; ###, P<0.001: cells pre-incubated with

PTP-MEG2 inhibitors compared with control vehicles.

(E-F). Ang II (100 nM) induced amperometric spikes by primary mouse chromaffin cells after

incubation with PTP-MEG2 inhibitor at different concentrations.

(G). The distribution of the quantal size of AngII (100 nM)-induced amperometric spikes by

primary chromaffin cells after incubation with different concentrations of PTP-MEG2

inhibitor. Histograms show the number of amperometric spikes of different quantal sizes.

(H and I). Secretory vesicles of primary chromaffin cells were examined by transmission

electron microscopy before or after 100 nM AngII stimulation, with or without pre-incubation

with 400 nM Compound 7.   Scale bars: H, I, 150 nm. Red arrows indicate morphologically

docked LDCVs and green arrows stands for undocked LDCVs.

(J). Vesicle numbers according to different distances from the plasma membrane were

calculated in the presence or absence of 100 nM AngII or 400 nM Compound 7. PM stands

for plasma membrane. * and *** indicate P<0.05 and P<0.001, respectively, compared with

the control group. ### indicate P<0.001 between incubated with the control vehicles and cells

incubated with Compound 7.

(K). The percentage of vesicles with different diameters was measured after 100 nM AngII

stimulation or incubation with 400 nM Compound 7. Compound 7 significantly decreased

vesicle size under AngII stimulation. *, ** and *** indicate P<0.05, P<0.01 and P<0.001,

respectively, between cells stimulated with AngII compared with un-stimulated cells. ###

indicate P<0.001 between incubated with the control vehicles and cells incubated with

Compound 7. N.S. means no significant difference.

(L). The percentage of pre-spike foot (PSF) (left panel) and stand-alone foot (SAF) (right

panel) were calculated after incubation with the indicated concentrations of Compound 7. #

indicates incubation with the indicated concentrations of Compound 7 compared with the

subgroup with AngII and without Compound 7. # indicates P<0.05 and ### indicates P<0.001.

All statistical significance was calculated with one-way ANOVA.

1175

Figure 2 Interaction of PTP-MEG2 and tyrosine-phosphorylated NSF in the medulla

and the crystal structure of the PTP-MEG2/phospho-NSF-segment complex.

(A). NSF was phosphorylated after stimulation with AngII or KCl. Adrenal medulla cells

were stimulated with 100 nM AngII or 70 mM KCl for 5 min and lysed. NSF was

immunoprecipitated with a specific NSF antibody coated with Protein A/G beads. A

pan-phospho-tyrosine antibody pY^{20} was used in Western blotting to detect the

tyrosine-phosphorylated NSF in the adrenal medulla under different conditions.

(B). NSF was co-localized with PTP-MEG2 in the adrenal medulla. After stimulation with

100 nM AngII, NSF and PTP-MEG2 in the adrenal medulla were visualized with

immunofluorescence. White arrow stands for co-localization of the PTP-MEG2 and NSF.

(C). Analysis of PTP-MEG2 and NSF fluorescence intensities by Pearson's correlation

analysis. The Pearson's correlation coefficient was 0.7.

(D). Phosphorylated NSF interacted with PTP-MEG2. PC12 cells were transfected with

FLAG-NSF. After stimulation of the cells with 100nM AngII, the potential PTP-MEG2

substrate in cell lysates was pulled down with GST-PTP-MEG2-$D^{470}A$ or GST control.

(E). The overall structure of PTP-MEG2-$C^{515}A/D^{470}A$ in complex with the NSF-pY^{83}

phospho-segment.

(F). The 2Fo-Fc annealing omit map (contoured at 1.0σ) around the NSF-pY^{83}

phospho-segment.

(G). Plot of distance RMSDs of individual residues between the crystal structures of the

PTP-MEG2/NSF-pY^{83} phospho-peptide complex and the PTP-MEG2 native protein

(PDB:2PA5).

(H). Superposition of the PTP-MEG2/NSF-pY^{83} phospho-peptide complex structure (red) on

the PTP-MEG2 native protein structure (PDB:2PA5, blue). The structural rearrangement of

the WPD loop and β3-loop-β4 is highlighted.

(I and J). The closure of the WPD loop and corresponding conformational changes in the

inactive state (I) and the active state (J) of PTP-MEG2. The rotation of R^{521} leads to the

movement of W^{468} and a corresponding 7 Å movement of the WPD loop.

(K and L). The structural rearrangement of β3-loop-β4 of PTP-MEG2 in the active state (L)

compared with the inactive state (K). The disruption of the salt bridge between E^{406} and R^{521}

contributed to the new conformational state of the N-terminal of β3-loop-β4.

(M and N). The conformational change of the P-loop of the inactive state (M) and the active

state of PTP-MEG2 in response to NSF-pY^{83} segment binding (N). The disruption of the

charge interaction between R^{409} and D^{335} resulted in the movement of the main chain from

G^{408} to K^{411}.

1211

Figure 3. Molecular determinants of PTP-MEG2 interaction with the NSF-pY^{83} site.

(A). Schematic representation of interactions between PTP-MEG2 and the NSF-pY^{83} site.

(B). Relative values of the decreased phosphatase activities of different PTP-MEG2 mutants

towards pNPP (p-Nitrophenyl phosphate) compared with wild-type PTP-MEG2.

(C). Fold-change decreases in the phosphatase activities of different PTP-MEG2 mutants

towards the NSF-pY^{83} phospho-segment compared with wild-type PTP-MEG2. The red

column indicates that the mutation site has greater effect on the enzyme activity toward

phosphor-NSF segement than the PNPP. The green column indicates that the mutation site

only has significant effect on the enzyme activity toward NSF phospho-segement but not

pNPP White column stands for the mutant has similar significant effects on both pNPP and

NSF phospho-segement.

(D). Sequence alignment of PTP-MEG2 from different species and with other PTP members.

Important structural motifs participating in phosphatase catalysis or the recognition of the

NSF-pY^{83} site are shown, and key residues contributing to the substrate specificity are

highlighted.

B-C, the statistical significance was calculated with one-way ANOVA. *** P<0.001;

PTP-MEG2 mutants compared with the control. Data were obtained from 3 independent

experiments. N.S. means no significant difference. All statistical significance was calculated

 with one-way ANOVA.

Figure 4. The tip opening of the pY+1 pocket is critical for PTP-MEG2 substrate specificity.

(A). Surface representations of the complex structures of PTP-MEG2-C^{515}A/D^{470}A/NSF-pY^{83}, PTP1B-C^{215}A/IRK-pY$^{1162/1163}$ (PDB:1EEN), LYP-C^{227}S/SKAP-HOM-pY^{75} (PDB:3OMH), PTPN18-C^{229}S/HER2-pY^{1248} (PDB:4GFU), and PTP-STEP/pY (PDB:2CJZ). Crystal structures of PTP-MEG1 (PDB:2I75), SHP1 (PDB:4HJQ), PTPH1 (PDB:4QUN) and SHP2 (PDB: 3B7O). The pY+1 sites are highlighted.

(B). Summary of the distance between the Cβ atoms of D^{335} and Q^{559} (corresponding to PTP-MEG2 number) of the pY+1 pocket in PTP-MEG2 and other classical non-receptor PTPs bearing the same residues at similar positions.

Figure 5. Effects of different PTP-MEG2 mutants on properties of catecholamine secretion from primary chromaffin cells.

(A). Primary mouse chromaffin cells were transduced with a lentivirus containing the gene for wild-type PTP-MEG2 or different mutants with a GFP tag at the C-terminus. Positive transfected cells were confirmed with green fluorescence and selected for electrochemical analysis. Typical amperometric current traces evoked by AngII (100 nM for 10 seconds) in the control (transduced with control vector) (top left panel), WT (top right panel), G^{334}R (bottom left panel), and D^{335}A (bottom right panel) are shown.

(B). The schematic diagram shows key residues of PTP-MEG2 defining the substrate specificity adjacent to the pY^{83} site of NSF.

(C). The distribution of the quantal size of chromaffin cells transduced with lentivirus containing the genes encoding different PTP-MEG2 mutants.

(D). Statistical diagram of the quantal size in Fig. 5A. The secretion amount of each group was standardized with respect to the control group.

(E-G). Calculated parameters of secretory dynamics, including the spike amplitude (E), PSF frequency (F), and SAF frequency (G).

D-G, * indicates PTP-MEG2 mutants overexpression group compared with the WT overexpression group. # indicates the overexpression group compared with the control group. **, P<0.01; *** P<0.001 and #, P<0.05; ##, P<0.01; ### P<0.001. N.S. means no significant difference. All statistical significance was calculated with one-way ANOVA.

Figure 6. Identification of potential candidate substrates of PTP-MEG2 that participate in fusion pore initiation and expansion.

(A). Flowchart for the workflow to predict the candidate substrates of PTP-MEG2 during fusion pore initiation and expansion. A total of 54 proteins were enriched with the functional protein association network STRING and the text mining tool PubTator by searching the keywords "fusion pore", "secretory vesicle" and "tyrosine phosphorylation". These proteins were filtered with UniProt by selecting proteins located only in the membrane or vesicle, which resulted in 28 candidates. The Human Protein Atlas database was then applied to exclude proteins with no expression in the adrenal gland. Finally, we used the post-translational-motif database PhosphoSitePlus to screen candidate proteins with potential phospho-sites that matched our sequence motif prediction at the pY+1 or pY+2 positions.

(B). After the bioinformatics analysis, a total of 12 candidate PTP-MEG2 substrates that may participate in fusion pore initiation and expansion and their potential phospho-sites were displayed.

(C). The GST pull-down assay suggested that PACSIN1, MUNC18-1, VAMP7, SNAP25, DYNAMIN 1 and DYNAMIN 2 directly interact with PTP-MEG2. PC12 cells were transfected with plasmids of candidate substrates, including SYN1, MUNC18-3, PACSIN1, SCAMP1, MUNC18-1, PPP3CA, STX17, VAMP7, SYT7, SYT11, SNAP25, DYNAMIN 1, and DYNAMIN 2 stimulated with 100 nM AngII. The tyrosine phosphorylation of these proteins was verified by specific anti-pY antibodies (Supplemental Fig. 9 A-E). The potential substrates of PTP-MEG2 in cell lysates were pulled down with a GST-PTP-MEG2-D^{470}A trapping mutant and then detected by Western blotting. White arrow stands for western band consistent with the predicted molecular weight of the potential PTP-MEG2 substrate.

(D-F). Co-immunostaining assays of PTP-MEG2 with potential substrates in the adrenal medulla. MUNC18-1, VAMP7 and DYNAMIN 2 all showed strong co-localization with PTP-MEG2 after 100 nM AngII stimulation in the adrenal medulla. White arrow stands for co-localization of PTP-MEG2 with MUNC18-1 or VAMP7. The Pearson's correlation coefficients for D, E and F were 0.61, 0.65 and 0.79 respectively. The co-immunostaining results of PTP-MEG2 with other potential substrates are shown in Supplemental Fig. 8.

Figure 7. Dephosphorylation of MUNC18-1 at the pY^{145} site by PTP-MEG2 determines the "foot" probability of catecholamine secretion from chromaffin cells.

(A). Structural representation and location of MUNC18-1-Y^{145} in the complex structure of MUNC18-1-SYNTAXIN1 (PDB: 3PUJ).

(B). Detailed structural representation of Y^{145} and its interacting residues. Y^{145} of MUNC18-1 interacts with the main chain carboxylic oxygen of residues I^{539} and G^{568} and forms hydrophobic interactions with L^{6}, I^{539}, L^{147}, L^{179}, L^{183}, and L^{230} to tether the arc shape of the three domains of MUNC18-1 (PDB: 3PUJ).

(C). Association analysis of SNPs of MUNC18-1 with human disease.

(D). Interactions of the PTP-MEG2-trapping mutants with the MUNC18-1-Y^{145} mutants and DYNAMIN2-Y^{125} mutants. PC12 cells were transfected with FLAG-MUNC18-1-Y^{145}, FLAG-DYNAMIN2-Y^{125} and different mutations of the FLAG-MUNC18-1-Y^{145}A, Y^{145}H, Y^{145}E or Y^{145}F mutants, FLAG-DYNAMIN2-Y^{125}A, Y^{125}E, Y^{125}F mutants, 24 hours before stimulation with 100 nM AngII, respectively. The cell lysates were then incubated with GST-PTP-MEG2-D^{470}A for 4 hours with constant rotation. The potential PTP-MEG2 substrates were pulled down by GST-beads, and their levels were examined by the FLAG antibody with Western blotting.

(E). Primary chromaffin cells were transduced with lentivirus containing the gene encoding wild-type MUNC18-1 or different mutants. These cells were stimulated with 100 nM AngII. The amperometric spikes were detected with CFE. Typical amperometric traces are shown.

(F). The percentages of pre-spike foot for wild-type MUNC18-1 or different mutants were calculated.

(G). The percentages of stand-alone foot for wild-type MUNC18-1 or different mutants were calculated.

(H). The MUNC18-1-Y^{145} mutation decreased the interaction between MUNC18-1 and SYNTAXIN1. PC12 cells were transfected with plasmid carrying SYNTAXIN-1. The proteins in cell lysates were pulled down with purified GST-MUNC18-1-Y^{145} and the GST-MUNC18-1-Y^{145}H/E/F, and detected with SYNTAXIN1 antibody. The right histogram shows the quantified protein levels.

(I). Primary chromaffin cells were transduced with lentivirus containing the gene encoding wild-type DYNAMIN2 or different mutants. These cells were stimulated with 100 nM AngII. The amperometric spikes were detected with CFE. Typical amperometric traces are shown.

(J). The percentages of pre-spike foot for wild-type DYNAMIN2 or different mutants were calculated.

(K). The percentages of stand-alone foot for wild-type DYNAMIN2 or different mutants were calculated.

(L). Plasmids encoding DYNAMIN2-WT, DYNAMIN2-Y^{125}E or DYNAMIN2-Y^{125}F and LYN-YFP, AT1aR-C-RLUC were co-transferred into HEK293 cells at 1:1:1. The BRET experiment was performed to monitor the mutation effects of DYNAMIN2 on AT1aR endocytosis in response to AngII (1μM) stimulation as previously described.

(M). A GTPase assay was performed on His-DYNAMIN-2-WT (500ng), His-DYNAMIN-2-Y^{125}E (500ng), His-DYNAMIN-2-Y^{125}F (500ng). Purified DYNAMIN-2 proteins were incubated with 5μM GTP in the presence of 500μM DTT for 90 minutes, followed by addition of GTPase-Glo™ Reagent and detection reagent. The luminescence were measured after 37-minute incubation. Bar representation of the results from three

independent experiments.

F-H, J-M * indicates MUNC18-1, DYNAMIN2 mutants overexpression group compared with

the WT overexpression group. # indicates MUNC18-1, DYNAMIN2 overexpression group

compared with the control group. *, $P<0.05$; **, $P<0.01$; *** $P<0.001$ and #, $P<0.05$; ##,

$P<0.01$; ### $P<0.001$. N.S. means no significant difference. All statistical significance was

calculated with one-way ANOVA.

1345

Figure 8. The structural details of the PTP-MEG2-MUNC18-1-Y^{145} interaction and

catalytic bias of PTP-MEG2 towards NSF-pY^{83} and MUNC18-1-pY^{145}

(A). The 2Fo-Fc annealing omit map (contoured at 1.0σ) around MUNC18-1-pY^{145}

phospho-segment.

(B). Comparison of residues of PTP-MEG2 interacting with NSF and MUNC18-1. Amino

acid residues of NSF and MUNC18-1 are coloured as follows: green, residues interacting

with both NSF and MUNC18-1; red, residues specifically contributing to NSF recognition;

blue, residues selectively contributing to MUNC18-1 interaction.

(C). The structural alteration of the interactions surrounding Y^{471} of PTP-MEG2 with the

MUNC18-1-pY^{145} site.

(D). The structural alteration of the interactions surrounding I^{519} of PTP-MEG2 with the

MUNC18-1-pY^{145} site.

(E). Relative phosphatase activities of different PTP-MEG2 mutants towards the

MUNC18-1-pY^{145} phospho-segment compared with wild-type PTP-MEG2.

(F). Relative phosphatase activities of different PTP-MEG2 mutants towards the DYNAMIN

2-pY^{125} phospho-segment compared with wild-type PTP-MEG2. E-F Residues are coloured

according to Figure 8B.

(G). Schematic illustration of the PTP-MEG2-regulated processes of vesicle fusion and

secretion in chromaffin cells via the dephosphorylation of different substrates with distinct

structural basis. PTP-MEG2 regulates vesicle fusion and vesicle size during the

catecholamine secretion of adrenal gland by modulating the phosphorylation state of the pY^{83}

site of NSF, which relies on the key residues G^{334}, D^{335} (pY loop), Y^{471} (WPD loop), I^{519}

(P-loop) and Q^{559} (Q loop). PTP-MEG2 regulates the fusion pore initiation and expansion

procedures of catecholamine secretion by the adrenal gland (also designated as foot

probability) by modulating the newly identified substrate MUNC18-1 at its pY^{145} site and

DYNAMIN 2 at its pY^{125}, through distinct structural basis from that of its regulation of NSF

phosphorylation. E-F, * $P<0.05$, *** $P<0.001$; PTP-MEG2 mutants compared with the control.

N.S. means no significant difference. Data were obtained from 3 independent experiments.

All statistical significance was calculated with one-way ANOVA.

1375

Table -1 Crystallographic Data and Refinement Statistics

[a] Each dataset was collected from a single crystal.

The values in parentheses correspond to the highest resolution shell.

1379

Data Collection	PTP-MEG2·NSF pY[83] Peptide	PTP-MEG2·MUNC18-1 pY[145] Peptide
Space group	$p2_12_12$	$p2_12_12$
Cell Dimensions		
a (Å)	71.447	71.048
b (Å)	83.708	84.074
c (Å)	48.615	48.753
α (deg)	90	90
β (deg)	90	90
γ (deg)	90	90
Resolution (Å)	80-1.70(1.76-1.70)[a]	54.3-2.08(2.08-2.20)[a]
Unique observations	32748(3167)[a]	18019(1773)[a]
Completeness (%)	99.6(98.3)[a]	99.8(99.8)[a]
Redundancy	3.8(3.8)[a]	6.1(6.3)[a]
$<I>/<\sigma>$	22.22(2.34)[a]	6.5(2.3)[a]
R_{merge}	0.064(0.477)[a]	0.191(0.927)[a]
Structure Refinement	80-1.70	
Resolution (Å)	1.70	2.08
Reflections used for R_{work}/R_{free}	31212/1536	17995/896
R_{work}/R_{free}[b] (%)	0.1900/0.2173	0.2102/0.2304
Average B-factor		
Protein	20.98	35.36
Peptide	31.525	53.25
RMSD ideal bonds (Å)	0.008	0.008
RMSD ideal angles (deg)	1.212	0.968
RamachandranPlot (%)		
Most favored	93.73	93.56
Allowed	4.88	6.10
Generously allowed	1.39	0.34
Disallowed	0	0

Figure 1

Figure 2

Figure 3

Figure 4

Complex	PDB code	pY+1	Distance(Cβ)	
PTP-MEG2/NSF(pY83)		T	10.42	D335:Q559
PTP1B/EGFR(pY992)	1EEN	L	11.46	D48:Q262
LYP/SKAP-HOM(pY75)	3OMH	D	11.32	D63:Q274
PTPN18/HER2(pY1248)	4GFU	L	11.89	D64:Q276
PTP-STEP/pY	2CJZ		12.43	N312:Q516
PTP-MEG1	2I75		11.16	D687:Q896
SHP1	4HJQ		12.27	N278:Q500
PTPH1	4QUN		12.35	D678:Q886
SHP2	3B7O		12.44	N281:Q506

Figure 5

Figure 6

B

Protein	Tyrosine phosphorylation site	The sequence of phosphorylation sites
CAV1.2	Tyr 14	EGHLYTVPI
DYNAMIN 1	Tyr 273	LQKKYPSLA
SYNTAXIN 17(STX17)	Tyr 157	LTQIYALPE
MUNC18-3	Tyr 148	ESQVYTLDV
SYNAPSIN 1	Tyr 301	LTKTYATAE
TI-VAMP/VAMP7	Tyr 45	NKLTYSHGN
MUNC18-1	Tyr 145	ESQVYSLDS
DYNAMIN 2（DYN2)	Tyr 125	NLRVYSPHVL
PPP3CA	Tyr 83	PDVAYGATG
THE DYNAMIN-BINDING-PROTEIN ACTIN	Tyr 430	KSQYYSTVG
SYNAPTOTAGMIN 7(SYT7)	Tyr 371	IGKIYLSWK
INSULIN RECEPTOR	Tyr 94	LFRVYGLES
	Tyr 401	IRRSYALVS

Figure 7

Figure 8