# Roles of adenine methylation and genetic mutations in adaptation to different temperatures in *Serratia marcescens*

Matthieu Bruneaux[1][*][†], Ilkka Kronholm[1][*], Roghaieh Ashrafi[1],
Tarmo Ketola[1]

March 11, 2021

[1]Department of Biological and Environmental Science, University of Jyväskylä, Finland
[*]Joint first authors
[†]Author for correspondence: Matthieu Bruneaux, Department of Biological and Environmental Science, University of Jyväskylä, Finland (`matthieu.bruneaux@ens-lyon.org`).

Running title: "Roles of genetics and epigenetics in adaptation"

## Abstract

Epigenetic modifications can contribute to adaptation, but the relative contributions of genetic and epigenetic variation are unknown. Previous studies on the role of epigenetic changes in adaptation in eukaryotes have nearly exclusively focused on cytosine methylation (m5C), while prokaryotes exhibit a richer system of methyltransferases targetting adenines (m6A) or cytosines (m4C, m5C). DNA methylation in prokaryotes has many roles, but its potential role in adaptation still needs further investigation. We collected phenotypic, genetic, and epigenetic data using single molecule real-time sequencing of clones of the bacterium *Serratia marcescens* that had undergone experimental evolution in contrasting temperatures to investigate the relationship between environment and genetic, epigenetic, and phenotypic changes. This data provided a detailed description of the methylation landscape of *S. marcescens* and allowed us to examine the potential contributions of genetic and epigenetic changes to phenotypic adaptation. The genomic distribution of GATC motifs, which are the main target for m6A methylation, and of partially methylated epiloci pointed to a link between m6A methylation and regulation of gene expression in *S. marcescens*. Evolved strains, while genetically homogeneous, exhibited many polymorphic m6A epiloci. There was no strong support for a genetic control of methylation changes in our experiment, and no clear evidence of parallel environmentally-induced or environmentally-selected methylation changes at specific epiloci was found. Both some genetic and epigenetic variants were associated with some phenotypic traits. Overall, our results suggest that both genetic and adenine methylation changes have potential to contribute to phenotypic adaptation in *S. marcescens*, but that any environmentally-induced epigenetic change occurring in our experiment would probably have been quite labile.

**Keywords:** Experimental evolution, methyltransferase, single molecule real-time sequencing.

# Introduction

While the traditional view of evolution is that adaptation proceeds via changes in DNA sequences, any heritable phenotypic variation can in principle be subject to natural se-

lection. In recent years, some epigenetic changes such as DNA methylation changes have been found to be heritable and could thus theoretically play a role in evolution

(Jablonka and Raz, 2009; Day and Bonduriansky, 2011; Danchin et al., 2011; Kronholm and Collins, 2016; Danchin et al., 2019). However, even if convincing cases of epige-

netic inheritance have been documented (Cortijo et al., 2014), the role of epigenetic variation in evolution has been met with some skepticism (Charlesworth et al., 2017),

the main argument for caution being that despite frequent observations we know very little about the relative contributions of genetic and epigenetic variation to adaptation.

Addressing this question requires the simultaneous characterization of genetic, epigenetic and phenotypic changes in a population, which remains extremely challenging in

a natural system, and so far empirical results have lagged behind theoretical models. Experimental evolution under controlled laboratory conditions with bacteria that have

short generation times and large population sizes provides the opportunity to study this question in a practical way.

Non-genetic inheritance encompasses various mechanisms, such as structural inheritance via propagating templates (e.g. prions), self-sustaining regulation loops and

DNA methylation (Jablonka and Raz, 2009). DNA methylation is particularly well studied and exists both in eukaryotes and in prokaryotes. While DNA methylation

can have phenotypic effects in both of those taxonomic groups, important differences exist between them. In eukaryotes, C5-methylcytosine (m5C) is the best known mod-

ification and occurs in CpG-rich regions. Genome methylation patterns are maintained after DNA replication via maintenance methyltransferases but DNA methy-

lation can be reset in the germ line. In prokaryotes, N4-methylcytosine (m4C) and N6-methyladenine (m6A) are known in addition to m5C (Ratel et al., 2006; Casadesús

and Low, 2006). Methylation is lost on the newly synthetized DNA strand during

replication, and methylation restoration depends on the activity of sequence-specific

58  methyltransferases after passage of the replication fork.

DNA methylation systems in prokaryotes are very diverse and include restriction-

60  modification (RM) systems and orphan methyltransferases (MTases). RM systems

associate an MTase activity and a complementary endonuclease activity: methylation

62  of the target sequence protects DNA from restriction. Orphan MTases do not have

an associated restriction enzyme but are thought to be evolutionary derived from RM

64  systems. DNA methylation in bacteria has a wide range of well-documented roles:

protection against foreign DNA via RM systems, regulation of DNA replication and

66  mismatch repair based on hemimethylated DNA, control of mobile elements and reg-

ulation of gene expression (Casadesús and Low, 2006; Sánchez-Romero et al., 2015).

68  Crucially, DNA methylation patterns can be inherited in prokaryotes: orphan MTases

can compete with other regulatory proteins for access to cognate DNA sequences, for

70  example in promoter and regulatory regions, resulting in self-regulating loops that can

be maintained across cell division (Braaten et al., 1994).

72  We used *Serratia marcescens* as a bacterial model in evolution experiments and

characterized genetic, epigenetic and phenotypic differences between strains evolved

74  from a common ancestor culture. *S. marcescens* is an opportunistic pathogen which

can live in a variety of environments, including inside insect and mammalian hosts,

76  with an optimal growth temperature close to 31 °C (Grimont and Grimont, 1978; Flyg

et al., 1980; Ketola et al., 2013). *S. marcescens* possesses a Dam MTase targeting

78  adenines in 5′-GATC-3′ motifs, and it has been shown that this enzyme is important

to enable the mismatch repair machinery to distinguish between the original and the

80  new DNA strand during replication (Ostendorf et al., 1999; Blow et al., 2016).

While examples of inherited epigenetic changes and their involvement in bacterial

82  adaptation exist (Adam et al., 2008; Atack et al., 2015), we do not fully understand

yet the relative importance of epigenetic and genetic variation in evolution. Here, we

84  studied the role of epigenetics in evolution in a prokaryotic system by (1) characteriz-

ing the epigenetic variation related to m6A in *Serratia marcescens* after evolution in

86   different conditions, (2) determining if the observed epigenetic variation was associated

with specific environmental conditions, and (3) comparing the potential contributions

88   of epigenetic and genetic variation to phenotypic variation.

We addressed these questions by phenotyping and single-molecule real-time (SMRT)

90   sequencing of bacterial strains from an evolution experiment conducted at three con-

trasting thermal regimes: 31 °C constant, 38 °C constant, and 24–38°C fluctuating

92   environments (Ketola et al., 2013). By introducing a common garden step before

phenotypic measurements and DNA sequencing, we focused on the identification of

94   evolutionary differences in methylation patterns which would be stable enough to be

propagated across several generations after their establishment once selection pres-

96   sure was relieved and we ignored epigenetic mutations with fast back-mutation rates

which would be more akin to phenotypic plasticity. Additionally, to investigate the

98   mechanism by which adenine methylation could affect cell processes in *S. marcescens*,

we examined if the genomic distribution of the GATC cognate motif exhibited any

100  informative pattern.

## Materials and Methods

### 102  Experimental evolution and phenotypic measurements

We used bacterial strains that were obtained from a previous evolution experiment

104  (Ketola et al., 2013). To summarize this experiment briefly, we let populations of

*Serratia marcescens* initiated from a single common ancestor colony evolve under either

106  constant or fluctuating temperatures during three weeks (treatments: constant 31 °C,

constant 38 °C or alternating daily between 24 °C and 38 °C) (Ketola et al., 2013)

108  (Supplementary Figure S1). Note that lines evolved in 38 °C were not reported in

Ketola et al. (2013). The experiment lasted approximately 70 generations (based on

110  1/10 daily subcultures for 3 weeks). Given that population sizes were of the order of

$8 \times 10^6$ cells in each $400\,\mu\text{L}$ culture well at plateau, and that the fraction transferred to the next generation was $1/10$ of the previous culture, the effective population size is estimated to be $N_e = 2.6 \times 10^6$ during the experiment (using $N_e = N_0 g$ from Lenski et al. (1991), where $N_0 = 8 \times 10^5$ cells is the bottleneck size and $g = 3.3$ is the number of generations between transfers). After experimental evolution, individual colonies were isolated from each population by plating and stored at $-80\,°\text{C}$ in $50\,\%$ glycerol.

We randomly selected one frozen clone from each population for sequencing (10 from the $31\,°\text{C}$ treatment, 8 from the $38\,°\text{C}$ treatment and 10 from the fluctuating treatment, i.e. 28 evolved strains sequenced in total). As a reference and since the frozen stocks for the common ancestor itself could not be revived successfully prior to sequencing, we also sequenced the stock strain from ATCC (ATCC 13880) from which the common ancestor colony was derived (Supplementary Figure S1). This strain is referred to as the "reference strain" in the rest of this manuscript. We used the phenotypic data from Ketola et al. (2013) (Supplementary Figure S2), which included measurements of growth rate and yield at constant $24\,°\text{C}$, $31\,°\text{C}$ and $38\,°\text{C}$ for the evolved strains. In addition, growth rate and yield had been measured in a series of novel environments: under redox balance stress ($1\,\text{mg/ml}$ dithiotreitol), in the presence of the ciliate predator *Tetrahymena thermophila* and in the presence of a virus, the lytic bacteriophage PPV. For further details on these phenotypic measurements, see Ketola et al. (2013).

## Sequencing and genome annotation

**Sequencing**   We used single molecule real-time sequencing (SMRT) with the PacBio platform to sequence the evolved strains and the reference strain. Since no template amplification takes place prior to sequencing on a PacBio platform in order to detect base modifications, relatively large amounts of DNA per strain are necessary with this method. Evolved strains were thawed and grown overnight in 150 ml of liquid medium and DNA was extracted using the Wizard Genomic DNA Purification Kit

138 from Promega (WI, USA). One DNA sample (20 to 60 μg) per strain was sequenced

by the DNA Sequencing and Genomics Laboratory of the University of Helsinki on a

140 PacBio RS II sequencing platform using P6-C4 chemistry. Two single-molecule real-

time sequencing (SMRT) cells were run per DNA sample.

142     PacBio software and recommended protocols were used with default parameters for

the assembly and modification calling pipeline. For each strain, reads from the RS

144 II instrument were assembled with PacBio `RS_HGAP_Assembly.3`, as implemented in

SMRTportal 2.3.0. The resulting assembly for each strain was processed with the Gap4

146 program to generate *de novo* a first draft sequence for this strain and to circularize it.

PacBio `RS_Resequencing.1` protocol was then run 2 to 3 times for each sample to map

148 the reads to the draft sequence and generate a consensus sequence for each strain. The

average coverage of the draft chromosome per strain was high (from 102 to 413, average

150 268 across all strains). We used Samtools (version 1.7, Li et al. (2009)) to examine

PacBio reads mapped back to their respective assemblies in order to check for evidence

152 of polymorphism within each sequenced strain. Such polymorphism would indicate

that a given strain would be comprised of several genetic clones, which is possible if

154 an isolated colony grew from more than a single isolated cell. Based on the Samtools

pileup files, the distribution of proportions of mismatches for each position within a

156 strain was compatible with the expected base error rate for PacBio SMRT sequencing

(about 11-15%, Rhoads and Au (2015)), except for a short ca. 700bp region comprised

158 of repeated sequences and which was ignored in downstream analyses. Given the high

coverage of the chromosome sequences across all strains, the base calling was considered

160 accurate outside of this repeat region and genetic variants were directly called from an

alignment of the 29 assembled chromosomes built with Mugsy (Angiuoli and Salzberg,

162 2011).

**Estimation of methylated fractions**   Inter-pulse duration (IPD) ratios were used

164 with the PacBio `RS_Modification and Motif_Analysis` protocol to detect modified

bases and identify methylation sequence motifs. Methylated adenines (m6A) are the

166  modified bases with the strongest kinetic signal (Clark et al., 2012) and we only anal-

ysed those modifications in downstream analyses. Before downstream analysis, we

168  applied a quality value threshold of 20 (i.e. $p$-value $\leq 0.01$) for the event detection

score, which is a Phred-like quality score defined as $Q = -10\log_{10}(pval)$. For detected

170  events, a second quality score is available for modification identification (m6A or other

modification type). Since the identification quality score depends on the methylated

172  fraction (lower fractions yielding lower scores), we accepted identifications as m6A for

a given position in all strains if the identification quality score at that position was $\geq$

174  20 in at least one strain. We did this to avoid being over-conservative and underesti-

mating the methylated fractions for positions which were likely to be truly methylated

176  but with a low methylation fraction in some strains. We checked that the ability to

estimate low methylation fractions for m6A was not impaired in strains with the lowest

178  coverages by examining the relationship between coverage and m6A fraction in GATC

motifs (Supplementary Figure S3).

180  **Genome annotation**  We used the NBCI Prokaryotic Genome Annotation Pipeline

(PGAP, Tatusova et al. (2016), version 2020-09-24.build4894) to annotate the assem-

182  bled genome of our reference strain. The location of the origin of replication was

determined using the DoriC server (Gao et al., 2012). The CDS predicted by PGAP

184  were assembled into operons using the Operon Mapper server (`http://biocomputo.`
`ibt.unam.mx/operon_mapper/`, Taboada et al. (2018), run on 2020-12-07). Gene

186  ontology annotations for the CDS predicted by PGAP were pooled from two ap-

proaches: (i) submitting the predicted CDS to Interproscan (Jones et al. (2014), ver-

188  sion 5.48-83.0) and (ii) running a tblastn search of the predicted CDS against all

UniProtKB entries related to *Serratia marcescens* (UniProtKB database searched via

190  `https://www.uniprot.org/` on 2019-01-31) and keeping the hits with the highest bit

scores.

192  **Identification of putative methyltransferase genes**    To identify putative MTase

genes in our reference strain, we downloaded the nucleotide sequences of all the entries

194  available for the *Serratia* genus in the REBASE database (Roberts et al., 2015) (`http:`

`//rebase.neb.com/rebase/rebase.html` accessed on 2020-03-07) and performed a

196  local nucleotide-to-nucleotide blast search of those sequences against the assembled

genome of our reference strain. In order to identify putative MTases missing from the

198  REBASE database, we also downloaded all the protein entries from the NCBI Identical

Protein Groups (IPG) database using the search terms "DNA adenine methyltrans-

200  ferase" and restricting the search to the "bacteria" taxon (`https://www.ncbi.nlm.`

`nih.gov/ipg/` accessed on 2020-03-11). Those protein sequences were then matched

202  against our reference strain genome using a local protein-to-translated nucleotide blast

search. The IPG database was accessed programmatically using the `rentrez` R package

204  (Winter, 2017).

## Calculation of tetramer composition bias along the genome

In order to test if the main target sequence identified for adenine methylation (5′-
GATC-3′) was under differential selection depending on the genomic context, we cal-
culated the tetramer composition bias in two sets of chromosome segments: coding
sequences (CDS) and promoter regions (defined as 200-bp-long regions immediately
upstream of an operon-leading CDS). Sequences for CDS were taken from their (+)
strand, and sequences for promoters were taken from the strand corresponding to the
(+) strand of the downstream CDS. For each of the 256 possible tetramers, we counted
the total number of occurrences observed in those sets, $N_{\text{tet}}$, separately for CDS and
for promoter sets. We compared those observed values with the expected number of
occurrences based on two models: a zero-order model based on the frequencies of each
nucleotide (A, T, G, C) in each segment and assuming that tetramers were generated
from a random combination of those, and a Markov chain model taking into account
the underlying biases that might exist in the frequencies of dimers and trimers compris-

ing each tetramer in a given segment. In the zero-order model, the expected number of occurrences of a given tetramer in a sequence segment $i$ is (Pride et al., 2003):

$$E_{tet,i}^{zero} = (p_A)^a (p_T)^t (p_G)^g (p_C)^c \times L$$

where $p_A$, $p_T$, $p_G$ and $p_C$ are the observed frequencies of each nucleotide in the segment, $a$, $t$, $g$ and $c$ are the count of each nucleotide in the considered tetramer (between 0 and 4), and $L$ is the segment length. In the Markov chain approach, the expected number of occurrences of a given tetramer of composition $b_1 b_2 b_3 b_4$ in a sequence segment $i$ is (Rocha et al., 1998; Pride et al., 2003):

$$E_{tet,i}^{MC} = \frac{N_{b_1 b_2 b_3} \times N_{b_2 b_3 b_4}}{N_{b_2 b_3}}$$

where $N_{b_1 b_2 b_3}$, $N_{b_2 b_3 b_4}$ and $N_{b_2 b_3}$ are the numbers of occurrences of the trimers $b_1 b_2 b_3$, $b_2 b_3 b_4$ and of the dimer $b_2 b_3$ in segment $i$, respectively. For the CDS and promoter sets separately, we calculated the expected total number of occurrences as the sum of expected numbers for each segment $i$: $E_{tet}^{zero} = \Sigma_i E_{tet,i}^{zero}$ and $E_{tet}^{MC} = \Sigma_i E_{tet,i}^{MC}$. Deviations in the usage of each tetramer from expectation were then calculated as:

$$D_{tet}^{zero} = \frac{N_{tet} - E_{tet}^{zero}}{E_{tet}^{zero}} \quad \text{and} \quad D_{MC} = \frac{N_{tet} - E_{tet}^{MC}}{E_{tet}^{MC}}$$

206 for the zero-order and the Markov chain models, respectively.

## Identification of partially methylated m6A loci in GATC motifs

208 GATC motifs targetted by Dam are usually methylated on both strands, but hemi-methylated or non-methylated locations can be inheritable features important for the
210 regulation of various cellular processes (Braaten et al., 1994; Casadesús and Low, 2006). To identify GATC loci which were not fully methylated in our dataset, we considered
212 the estimated fractions of methylated adenines at each GATC locus separately for the

plus and the minus strands. In any given strain, loci for which no methylated fraction

214    was reported by the PacBio pipeline (event detection score < 20) were assigned a value

of 0 for that strain (i.e. non-methylated). Additionally, loci for which the modification

216    identity score was < 20 in all strains were also considered not to be methylated as m6A.

Assuming that the distribution of modified fractions on the plus and minus strands for

218    fully methylated loci would follow a truncated bivariate distribution centered around

(1, 1) (i.e. both strands fully methylated), we defined the set of partially methylated

220    GATC loci of interest for downstream analyses as the loci which presented modified

fractions on the plus or on the minus strand which deviated from the point of full

222    methylation at coordinates (1, 1) more than four times the average observed quadratic

distance to (1, 1), in at least one strain (Supplementary Figure S4). We checked that

224    low estimated values of methylated fractions were not a sequencing artifact due to low

coverage of the corresponding GATC loci by examining the relationship between cover-

226    age bins and estimated methylated fraction (Supplementary Figure S3). We tested for

association between genetic variants and methylated fractions of partially methylated

228    m6A epiloci using Wilcoxon rank-sum tests. For each genetic variant present in more

than two sequenced genomes, we ran Wilcoxon tests against all partially methylated

230    m6A epiloci and applied a Benjamini-Hochberg false discovery rate (FDR) correction

(Benjamini and Hochberg, 1995) on those tests p-values.

## Effect of evolutionary treatments on genetic and epigenetic changes

234    To test for an association between evolutionary treatments and genetic variants, we

ran Fisher's exact tests on the contingency tables relating genotypes and evolutionary

236    treatments, for each genetic variant or linked variants (haplotypes) and for pooled

variants (as defined in Figure 3a), with a Benjamini-Hochberg FDR correction to the

238    tests p-values.

To determine if some adenine methylation changes had been selected by the evo-

240  lutionary treatments and were stable enough to persist after several generations in
common garden conditions, we tested for association between evolutionary treatments

242  and methylated fractions of partially methylated m6A with a methylated fraction range
over the sequenced clones $> 0.2$ using Kruskal-Wallis rank sum tests with Benjamini-

244  Hochberg FDR correction. In addition to those individual tests, and to investigate
whether the genes related to m6A epiloci which tended to be associated with evolu-

246  tionary treatments were consistently related to some biological processes, we performed
a gene ontology enrichment analysis. Each partially methylated m6A epiloci was as-

248  signed a gene if it was contained in its CDS or, alternatively, if it was located less than
500bp upstream of it. A gene assigned to an m6A epiloci was assigned this epiloci's

250  Kruskal-Wallis test's uncorrected p-value. Genes assigned to more than one m6A epiloci
were assigned their lowest p-values. The GO enrichment analysis was performed using

252  the topGO R package (Alexa and Rahnenfuhrer, 2020). The reference list of genes
comprised all the genes assigned a Kruskal-Wallis test's p-value as described above,

254  and the subset of interest was the genes which had an uncorrected p-value $< 0.05$. GO
enrichment test was performed using Fisher's exact test and the "weight01" algorithm

256  (Alexa et al., 2006; Alexa and Rahnenfuhrer, 2020).

## Association between phenotypes and genetic and epigenetic
258  data

We explored the association between genetic variants present in at least two sequenced
260  genomes and phenotypic traits using Wilcoxon rank sum tests with FDR correction. We
investigated the association between methylated fraction of the partially methylated

262  m6A epiloci and phenotypic traits using Spearman's $\rho$. We analyzed only partially
methylated epiloci with a methylated fraction range $> 0.2$ and we assigned each epiloci

264  its overlapping gene or, alternatively, the closest downstream gene within 500bp (if
any).

# Results

## Genome sequences and genetic variants

We sequenced ten strains from the 31 °C treatment, eight from the 38 °C treatment and ten from the 24–38°C fluctuating treatment, as well as the original reference stock strain used in the experimental evolution (Supplementary Figure S1). The reference strain chromosome was 5 117 300 bp long, with a GC content of 59.8%. The genome annotation by NCBI's PGAP predicted 4795 CDS, organized in 2686 operons according to Operon Mapper. 52 mutations were identified from the aligned genomes of the reference and evolved strains, of which 35 were located inside coding regions and 32 were non-synonymous (Table 1 and Supplementary Figure S5). Some mutations always occurred together in the sequenced strains: we use the term "haplotype" to refer to a set of linked mutations in our dataset (Table 1). Almost all identified haplotypes were strain-specific mutations. However, a striking feature of the variant map is the presence of 11 genetic mutations associated into a single haplotype and for which the minor allele is observed in 5 of the evolved strains from 38 °C and in the reference genome, but in no other evolved strain (haplotype $a$ in Table 1). This suggests that the ancestor colony used to initiate the replicated populations in the evolution experiment, and which was itself derived from the reference strain sequenced here, actually contained at least two lineages which were preserved in some populations at 38 °C but in no other evolutionary treatments. Even excluding variants from haplotype $a$, multiple parallel substitutions were observed among the evolved strains (Figure S5). Some genes exhibited several independent mutations in different strains: two independent mutations occurred in a deacetylase, three in a galactokinase and four in a glycosyltransferase (Table 1).

| ID | Haplotype | Freq. | Distrib. (31/fl./38) | Pos. (bp) | Type | Region | Effect | Overlapping or closest (≤500bp) gene | |
|---|---|---|---|---|---|---|---|---|---|
| | | | | | | | | Name | Function |
| 01 | f | 1/28 | 0/1/0 | 31 753 | SNP | CDS | non-syn. | DUF3053 domain-containing protein | unknown |
| 02 | a | *5/28 | 0/0/5 | 40 239 | SNP | CDS | non-syn. | HAP resistance protein | Unknown |
| 03 | d | 1/28 | 0/1/0 | 70 546 | indel | CDS | frameshift | Icd-like protein | host cell division inhibitor |
| 04 | | 6/28 | 2/3/1 | 92 159 | indel | non-CDS | - | - | - |
| 05 | b | 1/28 | 0/0/1 | 108 315 | indel | non-CDS | - | PTS beta-glucoside transporter | carbohydrate import |
| 06 | a | *5/28 | 0/0/5 | 131 841 | SNP | CDS | non-syn. | cellulose biosynthesis protein BcsG | biofilm? |
| 07 | | 1/28 | 1/0/0 | 173 551 | indel | non-CDS | - | protoheme IX biogenesis protein HemY | heme metabolism |
| 08 | a | *5/28 | 0/0/5 | 328 601 | SNP | CDS | non-syn. | condensation protein | non-ribosomal peptide synthesis |
| 09 | | 1/28 | 0/0/1 | 391 159 | indel | CDS | frameshift | RNA chaperone Hfq | regulation of stress transcription factors |
| 10 | g | 1/28 | 1/0/0 | 429 888 | indel | non-CDS | - | transcriptional regulator | similar to regulator of E. coli phage Mu |
| 11 | | 4/28 | 0/4/0 | 522 878 | indel | CDS | frameshift | integrase | prophage DNA Integration/excision |
| 12 | b | 1/28 | 0/0/1 | 751 961 | indel | CDS | frameshift | DNA polymerase II | DNA elongation |
| 13 | b | 1/28 | 0/0/1 | 770 532 | indel | CDS | frameshift | leu operon leader peptide | leucine biosynthesis |
| 14 | | 1/28 | 1/0/0 | 914 534 | indel | non-CDS | - | hydroxyacylglutathione hydrolase | lactate metabolism/response to heat |
| 15 | | 1/28 | 0/1/0 | 979 119 | indel | non-CDS | - | hypothetical protein | unknown |
| 16 | f | 1/28 | 0/1/0 | 1 093 517 | indel | CDS | frameshift | competence protein ComEA | cell surface DNA binding |
| 17 | | 1/28 | 1/0/0 | 1 185 019 | indel | CDS | frameshift | hypothetical protein | unknown |
| 18 | c | 1/28 | 0/0/1 | 1 311 662 | indel | CDS | no frameshift | galactokinase | galactose metabolism |
| 19 | b | 1/28 | 0/0/1 | 1 311 735 | SNP | CDS | non-syn. | galactokinase | galactose metabolism |
| 20 | h | 1/28 | 0/0/1 | 1 311 996 | SNP | CDS | syn. | galactokinase | galactose metabolism |
| 21 | a | *5/28 | 0/0/5 | 1 317 345 | SNP | CDS | syn. | Mo-dependent transcriptional regulator | regulation of transcription |
| 22 | | 5/28 | 1/2/2 | 1 421 879 | indel | non-CDS | - | acyl carrier protein | Fatty acid/polyketide biosynthesis |
| 23 | d | 1/28 | 0/1/0 | 1 609 697 | indel | non-CDS | - | transcriptional regulator GalS | regulation of galactose transport/catabolism |
| 24 | | 1/28 | 0/0/1 | 1 610 617 | indel | CDS | frameshift | galactose/galactoside ABC transporter MglB | galactose import |
| 25 | | 1/28 | 0/1/0 | 1 611 529 | SNP | CDS | non-syn. | galactose/galactoside ABC transporter MglA | galactose import |
| 26 | | 1/28 | 1/0/0 | 1 612 777 | indel | CDS | no frameshift | galactose/galactoside ABC transporter MglC | galactose import |
| 27 | | 4/28 | 0/3/1 | 1 665 941 | indel | CDS | frameshift | glycosyltransferase | LPS biosynthesis |
| 28 | | 1/28 | 0/1/0 | 1 670 147 | SNP | CDS | non-syn. | glycosyltransferase | LPS biosynthesis |
| 29 | | 2/28 | 0/1/1 | 1 670 356 | SNP | CDS | non-syn. | glycosyltransferase | LPS biosynthesis |
| 30 | e | 1/28 | 0/1/0 | 1 670 370 | SNP | CDS | non-syn. | glycosyltransferase | LPS biosynthesis |
| 31 | a | *5/28 | 0/0/5 | 1 861 227 | indel | non-CDS | - | putative transcriptional regulator | regulation of transcription |
| 32 | | 1/28 | 0/0/1 | 2 144 682 | indel | non-CDS | - | hypothetical protein | unknown |

(Continued on next page.)

| ID | Haplotype | Freq. | Distrib. (31/fl./38) | Pos. (bp) | Type | Region | Effect | Overlapping or closest (≤500bp) gene | |
|---|---|---|---|---|---|---|---|---|---|
| | | | | | | | | Name | Function |
| 33 | | 1/28 | 0/1/0 | 2 282 483 | indel | non-CDS | - | MATE family efflux transporter | Na+/H+ driven multidrug efflux pump |
| 34 | h | 1/28 | 0/0/1 | 2 353 326 | indel | CDS | frameshift | fumarase C (iron independent) | TCA cycle |
| 35 | b | 1/28 | 0/0/1 | 2 384 093 | indel | CDS | frameshift | HlyD (haemolysin secretion system) | haemolysin/cutinase excretion |
| 36 | a | *5/28 | 0/0/5 | 2 456 338 | SNP | CDS | non-syn. | peptidoglycan synthase | peptidoglycan biosynthesis |
| 37 | | 2/28 | 2/0/0 | 2 466 586 | SNP | CDS | non-syn. | MmgE/PrpD family protein | propionate metabolism/TCA cycle? |
| 38 | e | 1/28 | 0/1/0 | 2 941 884 | indel | non-CDS | - | VOC family protein | unknown |
| 39 | | 2/28 | 2/0/0 | 3 161 361 | SNP | CDS | syn. | serine/threonine protein kinase | regulation of cell processes |
| 40 | e | 1/28 | 0/1/0 | 3 408 594 | indel | non-CDS | - | nucleoside diphosphate hydrolase | regulation of cell processes |
| 41 | a | *5/28 | 0/0/5 | 3 477 366 | SNP | CDS | non-syn. | transcriptional regulator RcsB | capsule synthesis/cell division/biofilm/motility |
| 42 | | *0/28 | - | 3 600 509 | indel | non-CDS | - | phospholipid-binding lipoprotein MlaA | Outer membrane maintenance |
| 43 | a | *5/28 | 0/0/5 | 3 607 617 | SNP | CDS | non-syn. | heme exporter protein CcmB | cytochrome c biogenesis |
| 44 | a | *5/28 | 0/0/5 | 3 869 219 | SNP | CDS | syn. | cytochrome c | energy metabolism |
| 45 | | 1/28 | 1/0/0 | 4 025 724 | indel | non-CDS | - | acetyl-CoA carboxylase alpha subunit | lipid metabolism |
| 46 | | 1/28 | 1/0/0 | 4 337 062 | indel | non-CDS | - | tRNA-Phe | translation |
| 47 | | 1/28 | 0/0/1 | 4 362 753 | indel | non-CDS | - | glycoporin | carbohydrate import |
| 48 | c | 1/28 | 0/0/1 | 4 845 837 | indel | CDS | frameshift | peptidylprolyl isomerase | protein folding chaperone |
| 49 | g | 1/28 | 1/0/0 | 4 872 989 | indel | CDS | frameshift | short chain dehydrogenase | oxidoreductase |
| 50 | d | 1/28 | 0/1/0 | 4 924 755 | SNP | CDS | non-syn. | threonine dehydratase | amino acid metabolism |
| 51 | a | *5/28 | 0/0/5 | 5 010 850 | indel | CDS | frameshift | deacetylase | LPS biosynthesis |
| 52 | a | *5/28 | 0/0/5 | 5 010 868 | SNP | CDS | non-syn. | deacetylase | LPS biosynthesis |

Table 1: Summary of the genetic variants observed in the sequenced strains. Haplotype: letters denote groups of mutations for which alleles are associated together. Freq.: minor allele frequency among the 28 evolved strains. An asterisk denotes loci for which the reference strain carries the minor allele. Distribution of mutations per strains is shown in Supplementary Figure S5.

## Adenine methylation landscape in *Serratia marcescens*

### Methylated bases and methylation motifs

Out of $2\,057\,542$ adenine bases present in the bacterial chromosome, $81\,434$ ($4.0\,\%$) were detected as m6A (identification quality score $\geq 20$ in at least one of the evolved strains or in the reference strain). Most of those m6A were in GATC motifs: out of the $81\,434$ positions detected as m6A, $76\,228$ ($93.6\,\%$) were in a GATC context. Since a total of $76\,300$ adenines belong to a GATC context in the genome, this corresponds to a very high rate of adenine methylation in GATC motifs: $99.9\,\%$ of GATC adenines are detected as m6A in at least one evolved strain or the reference strain and $99.7\,\%$ were detected in all sequenced genomes. When detected as modified, m6A bases had an average methylated fraction of $70.3\,\%$ (s.d. $28.3\,\%$) outside GATC motifs and of $97.5\,\%$ (s.d. $5.0\,\%$) inside GATC motifs. We did not find any evidence of a problematic coverage effect on the ability to estimate low methylated fraction for m6A (Supplementary Figure S3).

The PacBio pipeline identified two sets of motifs for adenine methylation (Table 2). One motif set for m6A was the GATC palindrome mentioned above and the other was the much rarer pair AAAGNNNNNNTCG / TTTCNNNNNNAGC. For both sets, almost all genomic occurrences ($> 99\,\%$) were detected as modified.

| Motif | Occurrences on chromosome | Detected as methylated | Methylated fraction |
|---|---|---|---|
| 5′−G<u>A</u>TC−3′ | 38150 | | |
| 3′−CT<u>A</u>G−5′ | 38150 | 99.9 % | 97.5 % |
| 5′−AA<u>A</u>GNNNNNNTCG−3′ | 882 | 100 % | 93.8 % |
| 3′−TTTCNNNNNN<u>A</u>GC−5′ | 882 | 100 % | 96.3 % |

Table 2: Cognate sequence motifs for m6A modification. Methylated positions are underlined. Percentages of occurrences detected as methylated are reported as mean across the 29 sequenced samples; same for methylated fractions (calculated considering only methylated positions).

Three candidate genes for adenine MTases were identified using REBASE, and no additional candidates were identified using NCBI's IPG. One candidate was similar to M.Sma36365I, which is annotated as a type I MTase in REBASE and modifies the motif set AAGNNNNNGTTC / TTCNNNNNCAAG. It was located close to a restriction endonuclease in our reference strain genome, and is likely to be part of a restriction-modification system. A second candidate was identified as a type II MTase targetting GATC and was located inside a predicted prophage on our reference strain chromosome. Finally, the third candidate was similar to the orphan MTase M.Sma13880DamP, which also targets GATC motifs.

**Genomic methylation profiles**

We investigated m6A methylation profiles around the boundary between the first CDS of each predicted operon and the immediate upstream regions, which are likely to contain regulatory and promoter regions (Figure 1). From a base-composition perspective, upstream regions have lower GC content that coding regions, which can be related to constraints on DNA curvature or double helix stability in promoter regions (Bohlin et al., 2008) (Figure 1a). Average methylation of adenines into m6A bases was clearly lowered in upstream regions, in particular within 200 bp of the translation start sites, and reached background level again after a few tens of bp once in the operon coding regions (Figure 1b). This mostly mirrored the pattern observed for the proportion of adenines located in GATC motifs (Figure 1c).

## GATC usage bias in *Serratia marcescens* genome

Oligonucleotide usage bias in prokaryotic genomes exists as a result of evolutionary constraints, such as codon usage bias and palindrome avoidance (Rocha et al., 1998). To investigate if any such evolutionary constraint dictated the distribution of the main *S. marcescens* adenine methyltransferase cognate motif in promoter regions, we searched for evidence of differential usage bias of the 5′-GATC-3′ tetramer between

Figure 1: Profiles of nucleotide composition and adenine methylation around the start positions of operon-leading CDS. Red lines: LOESS regression (in (b) and (c), span = 0.75) and binned values (in (d), width = 20*bp*). Data points are averaged over each position relative to the leading CDS initiation codon, based on the 2686 operons predicted in the reference genome. Horizontal dashed lines and black triangles show genome-wide average values. Values for the three first bases on the coding sequences (usually ATG) are dropped from the plot to keep the y-scale reasonably narrow.

Figure 2: Tetramer usage bias in coding sequences (CDS) and promoters. (a) bias calculation based on observed single-nucleotide frequencies (zero-order approach). (b) bias calculation taking into account the observed frequencies of dimers and trimers (Markov chain approach). In the upper panels, the distance between a data point and its projection on the identity line measures how imbalanced the usage biases between CDS and promoters are (i.e. the differential usage bias). In the lower panels, those differential usage biases are sorted across all tetramers.

coding sequences (CDS) and promoters, defined here as 200-bp long regions upstream

334 of operon-leading CDS. In practice, we compared the usage bias of GATC with the

usage bias of all other nucleotide tetramers (Figure 2). The usage bias for a tetramer

336 in a given genomic region is positive if the tetramer is more abundant than expected by

chance, and negative if it is rarer. Overall, the range of tetramer usage bias in CDS and

338 promoters was larger when measured from a zero-order approach (deviation ranging

from -0.97 to +1.95) than from a Markov chain approach (-0.71 to +1.12). The cor-

340 relation between values obtained from the two approaches was moderate (Spearman's

$\rho = 0.20$), indicating that tetramer usage biases are strongly related to biases in dimers

342 and trimers usage which are taken into account by the Markov chain approach but not

by the zero-order approach. Most of the variation in tetramer usage bias was posi-

344 tively correlated between CDS and promoters (Spearman's $\rho = 0.58$ for Markov chain

estimates), but some tetramers exhibited large differences between their in-CDS and

346 in-promoter usage biases (Figure 2). Remarkably, 5′-GATC-3′ showed one of the largest

distortions in usage bias between CDS and promoter regions among all tetramers, in

348 both approaches. The 5′-GATC-3′ tetramer was rarer in promoters and more frequent

in CDS than expected by chance, even when taking into account underlying biases in

350 dimer and trimer usage (Figure 2).

## Identification of partially methylated m6A epiloci

352 Non-methylated and hemimethylated GATC loci are usually rare in the genome of

bacteria species possessing an adenine MTase targetting this motif. Such GATC loci

354 can be involved in the regulation of DNA replication and of DNA-protein interactions

and their methylation status can potentially be transmitted to the next generation

356 (Braaten et al., 1994; Casadesús and Low, 2006). Here, most adenines in GATC motifs

were observed close to full methylation but we identified 892 partially methylated

358 adenines in GATC motifs, i.e. 1.2 % of all adenines in GATC motifs. They were located

in 451 distinct GATC palindromes. We considered those partially methylated m6A as

360   the m6A epiloci of interest and used them for downstream analyses of association

with genetic and phenotypic data. Taking into account the genomic distribution of all

362   GATC motifs, partially methylated m6A were more frequent than expected in promoter

regions ($\chi^2 = 1417.3$, $df = 1$, $p < 0.001$) and rarer than expected in operons ($\chi^2 =$

364   $2908.5$, $df = 1$, $p < 0.001$), as is visible in Figure 1d. Finally, very little evidence of

association between genetics and epigenetics was found in this experiment: based on

366   FDR-corrected Wilcoxon rank-sum tests, associations with p-values $\leq 0.05$ were only

observed between haplotype $a$ and five epiloci.

## Effect of evolutionary treatments on genetic and epigenetic changes

370   Before FDR correction, only haplotype $a$ ($p < 0.001$), pooled variants for glycosyl-

transferase ($p = 0.007$) and for galactokinase ($p = 0.017$) and variant 11 ($p = 0.024$)

372   were tentatively associated with evolutionary treatments (p-values $<0.05$). After FDR

correction, only haplotype $a$ ($p = 0.018$) remained significantly associated with evolu-

374   tionary treatments, being only observed in evolved strains from the $38\,°C$ treatment

and in the reference strain (Table 1 and Supplementary Figure S5).

376        Concerning the association between evolutionary treatments and partially methy-

lated m6A epiloci, no association was significant at the $p < 0.05$ level after FDR

378   correction. We examined if some gene ontology terms were significantly enriched in

the genes which were the closest to the m6A epiloci which were tentatively associ-

380   ated with evolutionary treatments (uncorrected p-value for association $< 0.05$). Using

Fisher's exact test and the "weight01" algorithm (Alexa et al., 2006; Alexa and Rah-

382   nenfuhrer, 2020), five biological process (BP) categories were enriched in this set of

genes (topGO's $p < 0.05$): "cellular response to stress", "lipopolysaccharide biosyn-

384   thetic process", "negative regulation of cellular process", "L-amino acid transport" and

"cellular component biogenesis".

## Association between phenotypes and genotypes

386  We explored the association between genetic variants present in at least two sequenced
388  genomes and phenotypic traits using Wilcoxon rank sum tests with FDR correction
     (Figure 3a). The only genetic variants associated with phenotypic traits at the level
390  $p < 0.05$ were those comprising haplotype $a$, which were associated with growth rate
     and yield in the presence of a virus and yield at $24\,°C$. Those trait values decreased
392  in the presence of the minor allele (Figure 3b). Pooled variants for a galactokinase
     and a glycosyltransferase were tentatively associated with some traits ($p < 0.1$): for
394  example, the minor alleles related to the glycosyltransferase were potentially associated
     with higher growth rate at $31\,°C$, in the presence of DTT or in the presence of a virus.

## Association between phenotypes and partially methylated epi-loci

398  We examined the association between methylated fraction of the partially methylated
     m6A epiloci and phenotypic traits using Spearman's $\rho$ and assigning the m6A epiloci
400  to the closest genes (Figure 4). Forty genes were related to a partially methylated
     m6A epiloci tentatively associated with at least one trait, using an uncorrected p-value
402  threshold of 0.005 for Spearman's correlation. We did not apply FDR correction in this
     analysis, as the large number of tests run (10 traits $\times$ 892 partially methylated epiloci $=$
404  8920 tests) and the relatively small sample size of each test (28 evolved strains) resulted
     in all FDR-adjusted p-values to be $> 0.25$, even when the original uncorrected p-values
406  were $< 10^{-4}$. Gene ontology enrichment tests performed with topGO on a trait-by-
     trait basis revealed only very broad functional categories for genes associated with a
408  given trait, such as metabolism, molecule transport, regulation of cellular processes, cell
     communication and response to chemical stimulus (Supplementary Table S1). Based
410  on those functions, we performed a literature search for each of those forty genes to
     determine which had ever been found related to metabolism, nutrient uptake, cell

Figure 3: Association between genetic variants observed in at least two sequenced genomes and phenotypes. (a) Distributions of variants across evolutionary treatments and association between alleles and phenotypic traits (GR, growth rate; Y, yield). "Pooled" variants are variants affecting the same enzyme or function. (b) Detailed illustration of the association between major (M) and minor (m) alleles and phenotypes for variant *a* and for the "pooled" variants related to the galactokinase and to the glycosyltransferase. Data point colors correspond to the evolutionary treatment of each strain.

Figure 4: Association between adenine methylation and phenotypes. The heatmap shows Spearman's correlation between methylated fractions of partially methylated m6A epiloci (rows) and phenotypes (columns; GR, growth rate; Y, yield) for partially methylated m6A epiloci exhibiting a methylation fraction range ≥0.2 across sequenced samples and an uncorrected p-value ≤0.005 for Spearman's ρ with at least one trait. Overlapping or closest (≤500 bp) downstream genes are assigned to each epiloci. Gene functions are based on manual literature search for each gene product. Examples of relationships between m6A methylated fractions and phenotypes are given for the heatmap cells annotated with a letter.

412  motility and biofilm formation, or regulation of transcription, and found that most of
those genes were related to at least one such category (Figure 4).

# Discussion

We have shown that substantial variation in adenine methylation existed in *Serratia*

416  *marcescens* strains after experimental evolution followed by common garden conditions, and that GATC motifs and partially methylated m6A epiloci were not randomly

418  distributed along the genome. Sequenced strains were genetically clonal but they exhibited polymorphism in a small fraction of their m6A epiloci. In addition to de

420  novo mutations, genetic variants were dominated by 11 linked loci (haplotype *a*) which
were only observed in strains evolving at 38 °C and suggested the existence of some

422  degree of genetic polymorphism in the ancestor strain and differential selection between temperature treatments during the experiment. No clear effect of evolutionary

424  treatments on specific m6A epiloci was found, suggesting that parallel environmentally-induced or environmentally-selected changes on particular epiloci were rare or labile

426  in our experiment. Both genetic and epigenetic data were associated with some phenotypic traits to some extent. Surprisingly, the involved genes were not related to

428  temperature response in particular but rather related to metabolism, nutrient uptake,
motility/biofilm/adhesiveness and cell wall structure.

## Methylation landscape in *S. marcescens*

*E. coli* and several other Gammaproteobacteria possess a Dam enzyme responible for

432  m6A methylation of GATC motifs (Hattman et al., 1978; Casadesús, 2016), and GATC
adenine methylation in those species is involved in important functions such as protec-

434  tion against foreign DNA, timing of DNA replication and regulation of gene expression
(Casadesús, 2016). *S. marcescens* is known to also harbour a Dam enzyme targetting

436  GATC motifs (Ostendorf et al., 1999). The detailed *S. marcescens* adenine methylation

data we collected in our experiment indeed allowed us to identify two cognate motifs in

this species, including GATC. Putative adenine methyltransferases were identified in

the genome, two of which could be orphan MTases responsible for GATC methylation.

About 4 % of observed m6A positions were outside the identified cognate motifs, which

suggests either off-target methylation by specific MTases (Murray et al., 2012), the

existence of an unspecific DNA adenine MTase (Murray et al., 2018), or that some of

those detected methylation events could possibly be false positives (McIntyre et al.,

2019).

Dam activity in *S. marcescens* has been shown to play a role in mutation avoid-

ance by keeping track of the reference strand during post-replication mismatch repair

(Ostendorf et al., 1999). Here, several lines of evidence based on the genomic distri-

bution of GATC motifs and of the partially methylated epiloci indicate that GATC

methylation in *S. marcescens* is likely to be involved in other cell functions than solely

recognition of foreign DNA or help in mismatch repair, such as gene expression reg-

ulation. Firstly, average adenine methylation was lower in regulatory and promoter

regions than in the rest of the genome (Figure 1b), which was primarily due to the

lower frequency of GATC motifs in those regions (Figure 1c). Secondly, this unbal-

anced distribution of GATC motifs was not fully accounted for by the underlying

distributions of constituting di- and trinucleotides: the Markov chain model showed

that GATC were rarer than expected in promoters but more frequent in CDS, sug-

gesting that evolutionary constraints on the GATC motif differ between those genomic

regions (Figure 2). Finally, partially methylated sites were not randomly distributed

among the GATC sites but were more frequent in promoters and rarer in coding re-

gions (Figure 1d). Those observations are consistent with previous results in *E. coli*,

where GATC motifs are also more frequent in coding rather than non-coding regions

(Barras and Marinus, 1988) and GATC located in 5′ upstream regions of CDS are more

frequently under-methylated (Tavazoie and Church, 1998). In *E. coli*, GATC methy-

lation in gene regulatory regions is responsible for phase variation (Blyn et al., 1990)

(but not all methylation in regulatory regions lead to regulation of gene expression - see for example van der Woude et al. (1998)). Oshima et al. (2002) suggested that GATC in upstream regions could modulate gene expression by interacting with some regulatory proteins, while Riva et al. (2004) argued that the regulation of expression was due to clusters of GATC situated inside the coding regions and which would affect DNA stability and thus expression based on their methylation status.

Finally, while each sequenced strain in this experiment appeared genetically clonal, it is important to note that the methylated fractions observed within a given strain show that most of the partially methylated m6A epiloci were heterogeneously methylated in the culture: about 55 % of the methylated fraction values for those epiloci were between 0.2 and 0.8, showing that cells in a single culture are differently methylated at those epiloci. Methylated fraction must thus be considered a culture-level property, rather than a cell-level characteristic, in the same way that bacterial phenotypic traits such as growth rate and yield are measured for a culture, not for an individual cell.

## Effect of evolutionary treatments on adenine methylation

The origin of epigenetic variation and its stability are two fundamental evolutionary questions. Changes in methylation at a genomic locus can be either genetically determined, spontaneous, or environmentally induced. In eukaryotic systems, for example, most methylation changes observed in natural plant populations generally seem to be under genetic control (Dubin et al., 2015; Hagmann et al., 2015) but only a small proportion of methylation changes were under genetic control during experimental evolution of *Chlamydomonas* algae (Kronholm et al., 2017). In eukaryotes again, both spontaneous epigenetic changes (van der Graaf et al., 2015) or environmentally induced changes (Jiang et al., 2014; Wibowo et al., 2016) have also been observed. In prokaryotes, even though the mechanisms involved in methylation processes are markedly different from the eukaryotic ones, those three types of methylation changes are also possible: genetically determined changes (e.g. mutations affecting the MTase

492  enzyme itself or mutations in MTase target sites or in their flanking regions; Coffin and

Reich (2008)), spontaneous (stochastic competition between MTase and DNA binding

494  protein heritable via self-regulating loops), and environmentally induced (when other

DNA binding proteins influencing MTase access to DNA are responding to environ-

496  ment, like in the case of the *pap* operon or the *agn43* gene in *E. coli*; Wallecha et al.

(2002); Casadesús and Low (2013)). Importantly, the stability of methylation states in

498  prokaryotes can vary from a regular resetting during DNA replication to bistable epige-

netic switches underpinning phase variation. For a given epigenetic switch, OFF-to-ON

500  and ON-to-OFF rates can differ, allowing for some relative stability after a switching

event (Kaminska and van der Woude, 2010; Olivenza et al., 2019). Epigenetic changes

502  in bacteria can thus be involved in both short-term, partially heritable phenotypic

plasticity and in long-term response, possibly allowing for later genetic assimilation.

504  Given our experimental design, our conclusions below apply only to epigenetic changes

that can remain stable for several generations rather than to the more labile ones.

506     We did not find any strong evidence to support a genetic control of adenine methy-

lation in our experiment, as only five m6A epiloci were found associated with haplotype

508  *a*. While the statistical power to detect association from our data is limited since there

is no segregation among the bacterial clones, extensive genetic control would require

510  assuming that each genetic mutation controls multiple different epiloci in order for ge-

netic mutations to explain the observed epigenetic variation. Furthermore, the limited

512  association observed between m6A epiloci and haplotype *a* could be due to the

shared history of those epiloci with haplotype *a* prior to the initiation of the evolution

514  experiment (assuming that haplotype *a* was part of some standing genetic variation in

the ancestor culture at the initiation of the experimental populations).

516     Any consistent methylation pattern associated with multiple clones from a given

evolutionary treatment can in principle be due to methylation states induced by a

518  common environment or explained by spontaneous changes that were environmentally

selected in multiple populations, thus reflecting parallel evolution. It is also possible

520   that some epiloci have extremely high forward- and back-mutations rates, so that

some polymorphism is always present, or that the optimal fitness of a given culture

522   is composed of a mix of subpopulations heterogeneously methylated. We did not

observe any partially methylated m6A epiloci whose methylated fraction would be

524   significantly associated with evolutionary treatments, which suggest that no parallel

epigenetic changes were either induced nor selected by the evolutionary treatments, or

526   that changes would be too labile to persist over a few generations in common garden

conditions. Most of the observed epigenetic variation is thus likely to be spontaneous,

528   stochastic fluctuations, for which our experimental design does not allow to determine

if they affect the phenotype or are neutral. Highly labile epigenetic fluctuations could

530   possibly reflect some sort of epigenetic bet-hedging mechanism (Veening et al., 2008).

Our observations do not preclude, however, that environmental conditions could exert

532   selective pressure on stochastic epigenetic variation at different epiloci that would affect

a common biological process. Indeed, the results of the gene ontology enrichment

534   analysis of the genes located by the m6A epiloci most associated with evolutionary

treatments suggest that environmental pressure could act as a sieve selecting distinct

536   epigenetic changes related to common, important biological functions.

## Distribution of genetic variants among evolutionary treatments

538   Most of the genetic variants observed in our experiment were present in single strains

and we cannot determine if those variants provide an advantage in a specific evolution-

540   ary treatment or in the laboratory conditions in general. The only variant that was

statistically associated with a particular treatment was haplotype $a$, which comprised

542   11 linked loci and was observed only in evolved strains from the 38 °C treatment. Hap-

lotype $a$ was also found in the reference strain from which the ancestor used for the

544   evolution experiment was derived. The probability of 11 mutations arising indepen-

dently and in succession in several strains in our dataset is low given the duration of

546   the evolution experiment. It is thus likely that the ancestor culture used to initiate

all the populations of the experiment exhibited some genetic diversity in relation with

548  haplotype $a$, possibly due to cell aggregation occurring during the preparation of the

ancestor colony. No sign of this diversity is observed in the sequenced strains evolved

550  at 31 °C and 24–38°C, indicating that haplotype $a$ was driven to low frequencies or

extinction in 31 °C and 24–38°C conditions. This indicates that evolving at lower av-

552  erage temperature might impose stronger selective constraints on the genetic variants

in our experimental setup while higher temperature allowed for more diverse genetic

554  trajectories.

Interestingly, we observed mutations occurring independently at different positions

556  within several genes and being selected in parallel. This happened independently of

haplotype $a$ for a galactokinase and for a glycosyltransferase. Given the large effective

558  population size in our experiment ($N_e = 2.6 \times 10^6$, see Methods), genetic drift cannot

explain the fixation of multiple parallel mutations in independent populations, sug-

560  gesting instead that selection favoured a modified function for those particular genes.

Remarkably, those two genes and the deacetylase gene (with two linked mutations

562  belonging to haplotype $a$) are all involved in the biosynthesis of lipopolysaccharide.

Mutations in those three genes were observed in the 24–38°C and 38 °C treatments,

564  but not in the 31 °C treatment.

## Effect of genetic and epigenetic variation on phenotypes

566  Our main objective was to determine the potential contributions of genetics and epi-

genetics to adaptation in rapidly changing environments. In the original evolution

568  experiment during which the evolved strains were generated, 12 colonies were isolated

from each replicate population, thus providing good statistical power to detect an effect

570  of evolutionary treatment on phenotypic traits and enabling to show that strains from

the fluctuating 24–38°C treatment outperformed strains that evolved at constant 31 °C

572  (Ketola et al., 2013). In the subset of strains we sequenced here, phenotypic variabil-

ity results in overlapping phenotype ranges across treatments (Supplementary Figure

S2). Given this overlap, trying to link genetic and epigenetic data directly to phenotypic trait values rather than to evolutionary treatments might give insight into how

genetic variants and partially methylated m6A epiloci can influence phenotype. Note that given our experimental design, where evolved clones were isolated and cultured

in common garden conditions before phenotypic measurements and DNA extraction, short-lived epigenetic changes involved in rapid phenotypic plasticity would be lost and

only more persistent changes, stable over several generations and more amenable to playing a role in evolution, would be carried over from the evolution experiment.

Evidence supporting a potential effect on phenotypes was found both for genetic variants and for partially methylated m6A epiloci. Given that most genetic variants

were present in only one strain, the ability to detect statistically significant associations was limited, but haplotype $a$ was associated with some reduced performance in

the strains evolved at 38 °C and independent genetic variants associated with galactokinase and glycosyltransferase were tentatively associated with phenotypic changes.

From a general biological perspective, those genes and the genes affected by haplotype $a$ are somehow related to cell wall structure, metabolism and regulation of transcription,

which are all crucial cell functions likely to be impacted when environmental conditions change, but not to thermal stress in particular, and were associated both with

temperature-related traits and with performances in novel environments (exposure to DTT, ciliate or virus). Similarly, partially methylated m6A epiloci which were found

to be associated with phenotypes were related to genes involved in important, broad functional categories (metabolism, nutrient uptake, motility and biofilm formation).

Metabolism and nutrient uptake are both likely to be crucial in determining growth rate and yield, especially after evolving in a nutrient-limiting medium such as SPL 1%.

## Conclusion

We have shown that substantial epigenetic variation existed in adenine methylation

600  in *Serratia marcescens*, both across and within cultures of evolved strains. The dis-

tribution of methylated fractions and of the main adenine MTase cognate motif along

602  the genome pointed to a probable connection between adenine methylation and gene

expression regulation, as is the case in other Gammaproteobacteria. No clear evidence

604  of a genetic control of epigenetic variation, nor of stable environmentally-induced or

environmentally-selected epigenetic changes were found in our experiment. However,

606  this result does not rule out the existence of labile, short-term epigenetic responses

in particular in fluctuationg environmental conditions, which could be explored with

608  more massive methylome sequencing of evolved strains grown in their treatment condi-

tions and challenged in new conditions. Genetic variants from unexpected pre-existing

610  standing genetic variation at the beginning of the evolution experiment seemed to

be responsible for the majority of divergent phenotypic adaptation between the 38 °C

612  treatment and the others, but indications of both genetic and epigenetic effects on

some of the phenotypic traits observed in our experiment were found.

614  **Acknowledgements.** We acknowledge the Academy of Finland (Project 278751)

and the Centre of Excellence in Biological Interactions for funding and facilities, and

616  CSC-IT center for Science for computational resources.

**Authors contributions.** TK and IK conceptualized the study. Experimental design

618  for PacBio sequencing was done by TK and MB. DNA extraction for PacBio sequencing

was done by RA. MB, IK and TK analyzed the data and wrote the original draft with

620  later review by all co-authors.

**Conflict of interest.** The authors declare no competing financial interests.

622  **Data availability.** PacBio sequencing data (HDF5 files) were submitted to the Euro-

pean Nucleotide Archive's Sequence Read Archive (ENA-SRA) (https://www.ebi.ac.

624  uk/ena, project PRJEB40306) and assembled genomes to NCBI's GenBank (biosam-

ples SAMEA7301478 to SAMEA7301506). Genetic, epigenetic and phenotypic datasets

626  used for analysis will be submitted to the Dryad repository (https://datadryad.org).

# References

628  Adam M, Murali B, Glenn NO, Potter SS. 2008. Epigenetic inheritance based evolution
      of antibiotic resistance in bacteria. *BMC Evolutionary Biology*. 8(1):52.

630  Alexa A Rahnenfuhrer J. topGO: Enrichment Analysis for Gene Ontology, 2020. R
      package version 2.42.0.

632  Alexa A, Rahnenführer J, Lengauer T. 2006. Improved scoring of functional groups
      from gene expression data by decorrelating GO graph structure. *Bioinformatics*. 22
634  (13):1600–1607.

      Angiuoli SV Salzberg SL. 2011. Mugsy: fast multiple alignment of closely related
636  whole genomes. *Bioinformatics*. 27(3):334–342.

      Atack JM, Srikhanta YN, Fox KL, Jurcisek JA, Brockman KL, Clark TA, Boitano
638  M, Power PM, Jen FEC, McEwan AG, Grimmond SM, Smith AL, Barenkamp SJ,
      Korlach J, Bakaletz LO, Jennings MP. 2015. A biphasic epigenetic switch con-
640  trols immunoevasion, virulence and niche adaptation in non-typeable *Haemophilus
      influenzae*. *Nat Commun*. 6:7828.

642  Barras F Marinus MG. 1988. Arrangement of Dam methylation sites (GATC) in the
      Escherichia coli chromosome. *Nucleic Acids Research*. 16(20):9821–9838.

644  Benjamini Y Hochberg Y. 1995. Controlling the False Discovery Rate: A Practical
      and Powerful Approach to Multiple Testing. *Journal of the Royal Statistical Society.
646  Series B (Methodological)*. 57(1):289–300.

      Blow MJ, Clark TA, Daum CG, Deutschbauer AM, Fomenkov A, Fries R, Froula J,
648  Kang DD, Malmstrom RR, Morgan RD, Posfai J, Singh K, Visel A, Wetmore K,

Zhao Z, Rubin EM, Korlach J, Pennacchio LA, Roberts RJ. 2016. The epigenomic
650    landscape of prokaryotes. *PLOS Genet.* 12(2):e1005854.

Blyn LB, Braaten BA, Low DA. 1990. Regulation of *pap* pilin phase variation by a
652    mechanism involving differential dam methylation states. *The EMBO Journal.* 9
       (12):4045–4054.

654  Bohlin J, Skjerve E, Ussery DW. 2008. Investigations of Oligonucleotide Usage Variance
       Within and Between Prokaryotes. *PLOS Computational Biology.* 4(4):e1000057.
656    Publisher: Public Library of Science.

Braaten BA, Nou X, Kaltenbach LS, Low DA. 1994. Methylation patterns in pap
658    regulatory DNA control pyelonephritis-associated pili phase variation in *E. coli. Cell.*
       76(3):577–588.

660  Casadesús J. 2016. Bacterial DNA Methylation and Methylomes. *Advances in Exper-
       imental Medicine and Biology.* 945:35–61.

662  Casadesús J Low D. 2006. Epigenetic Gene Regulation in the Bacterial World. *Micro-
       biology and Molecular Biology Reviews.* 70(3):830–856.

664  Casadesús J Low DA. 2013. Programmed Heterogeneity: Epigenetic Mechanisms in
       Bacteria. *Journal of Biological Chemistry.* 288(20):13929–13935.

666  Charlesworth D, Barton NH, Charlesworth B. 2017. The sources of adaptive variation.
       *Proceedings of the Royal Society of London B: Biological Sciences.* 284(1855).

668  Clark TA, Murray IA, Morgan RD, Kislyuk AO, Spittle KE, Boitano M, Fomenkov A,
       Roberts RJ, Korlach J. 2012. Characterization of DNA methyltransferase specificities
670    using single-molecule, real-time DNA sequencing. *Nucleic Acids Research.* 40(4):
       e29–e29.

672  Coffin SR Reich NO. 2008. Modulation of *Escherichia coli* DNA Methyltransferase

Activity by Biologically Derived GATC-flanking Sequences. *Journal of Biological Chemistry.* 283(29):20106–20116.

Cortijo S, Wardenaar R, Colomé-Tatché M, Gilly A, Etcheverry M, Labadie K, Cail-lieux E, Hospital F, Aury JM, Wincker P, Roudier F, Jansen RC, Colot V, Johannes F. 2014. Mapping the epigenetic basis of complex traits. *Science.* 343(6175):1145–1148.

Danchin E, Charmantier A, Champagne FA, Mesoudi A, Pujol B, Blanchet S. 2011. Beyond DNA: integrating inclusive inheritance into an extended theory of evolution. *Nat Rev Genet.* 12(7):475–486.

Danchin E, Pocheville A, Rey O, Pujol B, Blanchet S. 2019. Epigenetically facili-tated mutational assimilation: epigenetics as a hub within the inclusive evolutionary synthesis. *Biological Reviews of the Cambridge Philosophical Society.* 94(1):259–282.

Day T Bonduriansky R. 2011. A unified approach to evolutionary consequences of genetic and nongenetic inheritance. *American Naturalist.* 178:E18–E36.

Dubin MJ, Zhang P, Meng D, Remigereau MS, Osborne EJ, Paolo Casale F, Drewe P, Kahles A, Jean G, Vilhjálmsson B, Jagoda J, Irez S, Voronin V, Song Q, Long Q, Rätsch G, Stegle O, Clark RM, Nordborg M. 2015. DNA methylation in *Arabidopsis* has a genetic basis and shows evidence of local adaptation. *eLife.* 4:e05255.

Flyg C, Kenne K, Boman HG. 1980. Insect pathogenic properties of *Serratia* mar-cescens: phage-resistant mutants with a decreased resistance to *Cecropia* immunity and a decreased virulence to *Drosophila*. *Journal of General Microbiology.* 120(1): 173–181.

Gao F, Luo H, Zhang CT. 2012. DoriC 5.0: an updated database of oriC regions in both bacterial and archaeal genomes. *Nucleic Acids Research.* 41(D1):D90–D93.

Grimont PAD Grimont F. 1978. The genus *Serratia*. *Annual Review of Microbiology*. 32(1):221–248.

Hagmann J, Becker C, Müller J, Stegle O, Meyer RC, Wang G, Schneeberger K, Fitz J, Altmann T, Bergelson J, Borgwardt K, Weigel D. 2015. Century-scale methylome stability in a recently diverged *Arabidopsis thaliana* lineage. *PLoS Genetics*. 11(1): e1004920.

Hattman S, Brooks JE, Masurekar M. 1978. Sequence specificity of the P1 modification methylase (M.Eco P1) and the DNA methylase (M.Eco dam) controlled by the Escherichia coli dam gene. *Journal of Molecular Biology*. 126(3):367–380.

Jablonka E Raz G. 2009. Transgenerational epigenetic inheritance: prevalence, mechanisms, and implications for the study of heredity and evolution. *Quarterly Review of Biology*. 84:131–176.

Jiang C, Mithani A, Belfield EJ, Mott R, Hurst LD, Harberd NP. 2014. Environmentally responsive genome-wide accumulation of de novo *Arabidopsis thaliana* mutations and epimutations. *Genome Research*. 24(11):1821–1829.

Jones P, Binns D, Chang HY, Fraser M, Li W, McAnulla C, McWilliam H, Maslen J, Mitchell A, Nuka G, Pesseat S, Quinn AF, Sangrador-Vegas A, Scheremetjew M, Yong SY, Lopez R, Hunter S. 2014. InterProScan 5: genome-scale protein function classification. *Bioinformatics*. 30(9):1236–1240.

Kaminska R van der Woude MW. 2010. Establishing and maintaining sequestration of Dam target sites for phase variation of agn43 in Escherichia coli. *Journal of Bacteriology*. 192(7):1937–1945.

Ketola T, Mikonranta L, Zhang J, Saarinen K, Örmälä AM, Friman VP, Mappes J, Laakso J. 2013. Fluctuating temperature leads to evolution of thermal generalism and preadaptation to novel environments. *Evolution*. 67:2936–2944.

722    Kronholm I Collins S. 2016. Epigenetic mutations can both help and hinder adaptive
       evolution. *Molecular Ecology.* 25:1856–1868.

724    Kronholm I, Bassett A, Baulcombe D, Collins S. 2017. Epigenetic and genetic contri-
       butions to adaptation in *Chlamydomonas. Molecular Biology and Evolution.* 34(9):
726    2285–2306.

       Lenski RE, Rose MR, Simpson SC, Tadler SC. 1991. Long-Term Experimental Evo-
728    lution in *Escherichia coli.* I. Adaptation and Divergence During 2,000 Generations.
       *The American Naturalist.* 138(6):1315–1341.

730    Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G,
       Durbin R, 1000 Genome Project Data Processing Subgroup. 2009. The Sequence
732    Alignment/Map format and SAMtools. *Bioinformatics (Oxford, England).* 25(16):
       2078–2079.

734    McIntyre ABR, Alexander N, Grigorev K, Bezdan D, Sichtig H, Chiu CY, Mason
       CE. 2019. Single-molecule sequencing detection of N 6-methyladenine in microbial
736    reference materials. *Nature Communications.* 10(1):1–11. Number: 1 Publisher:
       Nature Publishing Group.

738    Murray IA, Clark TA, Morgan RD, Boitano M, Anton BP, Luong K, Fomenkov A,
       Turner SW, Korlach J, Roberts RJ. 2012. The methylomes of six bacteria. *Nucleic*
740    *Acids Research.* 40(22):11450–11462.

       Murray IA, Morgan RD, Luyten Y, Fomenkov A, Corrêa IR, Dai N, Allaw MB, Zhang
742    X, Cheng X, Roberts RJ. 2018. The non-specific adenine DNA methyltransferase
       M.EcoGII. *Nucleic Acids Research.* 46(2):840–848.

744    Olivenza DR, Nicoloff H, Sánchez-Romero MA, Cota I, Andersson DI, Casadesús J.
       2019. A portable epigenetic switch for bistable gene expression in bacteria. *Scientific*
746    *Reports.* 9(1):1–9. Number: 1 Publisher: Nature Publishing Group.

Oshima T, Wada C, Kawagoe Y, Ara T, Maeda M, Masuda Y, Hiraga S, Mori H. 2002. Genome-wide analysis of deoxyadenosine methyltransferase-mediated control of gene expression in *Escherichia coli*. *Molecular Microbiology*. 45(3):673–695.

Ostendorf T, Cherepanov P, Vries Jd, Wackernagel W. 1999. Characterization of a dam mutant of *Serratia marcescens* and nucleotide sequence of the dam region. *Journal of Bacteriology*. 181(13):3880–3885.

Pride DT, Meinersmann RJ, Wassenaar TM, Blaser MJ. 2003. Evolutionary implications of microbial genome tetranucleotide frequency biases. *Genome Research*. 13(2):145–158.

Ratel D, Ravanat JL, Berger F, Wion D. 2006. N6-methyladenine: the other methylated base of DNA. *Bioessays*. 28(3):309–315.

Rhoads A Au KF. 2015. PacBio Sequencing and Its Applications. *Genomics, Proteomics & Bioinformatics*. 13(5):278–289.

Riva A, Delorme MO, Chevalier T, Guilhot N, Hénaut C, Hénaut A. 2004. The difficult interpretation of transcriptome data: the case of the GATC regulatory network. *Computational Biology and Chemistry*. 28(2):109–118.

Roberts RJ, Vincze T, Posfai J, Macelis D. 2015. REBASE—a database for DNA restriction and modification: enzymes, genes and genomes. *Nucleic Acids Research*. 43(D1):D298–D299. Publisher: Oxford Academic.

Rocha EP, Viari A, Danchin A. 1998. Oligonucleotide bias in *Bacillus subtilis*: general trends and taxonomic comparisons. *Nucleic Acids Research*. 26(12):2971–2980.

Sánchez-Romero MA, Cota I, Casadesús J. 2015. DNA methylation in bacteria: from the methyl group to the methylome. *Current Opinion in Microbiology*. 25:9–16.

Taboada B, Estrada K, Ciria R, Merino E. 2018. Operon-mapper: a web server for

precise operon identification in bacterial and archaeal genomes. *Bioinformatics*. 34 (23):4118–4120.

Tatusova T, DiCuccio M, Badretdin A, Chetvernin V, Nawrocki EP, Zaslavsky L, Lomsadze A, Pruitt KD, Borodovsky M, Ostell J. 2016. NCBI prokaryotic genome annotation pipeline. *Nucleic Acids Research*. 44(14):6614–6624.

Tavazoie S Church GM. 1998. Quantitative whole-genome analysis of DNA-protein interactions by in vivo methylase protection in E. coli. *Nature Biotechnology*. 16(6): 566–571.

van der Graaf A, Wardenaar R, Neumann DA, Taudt A, Shaw RG, Jansen RC, Schmitz RJ, Colomé-Tatché M, Johannes F. 2015. Rate, spectrum, and evolutionary dynamics of spontaneous epimutations. *Proceedings of the National Academy of Sciences*. 112(21):6676–6681.

van der Woude M, Hale WB, Low DA. 1998. Formation of DNA Methylation Patterns: Nonmethylated GATC Sequences in gut and pap Operons. *Journal of Bacteriology*. 180(22):5913–5920.

Veening JW, Smits WK, Kuipers OP. 2008. Bistability, epigenetics, and bet-hedging in bacteria. *Annual Review of Microbiology*. 62(1):193–210.

Wallecha A, Munster V, Correnti J, Chan T, van der Woude M. 2002. Dam- and OxyR-dependent phase variation of agn43: essential elements and evidence for a new role of DNA methylation. *Journal of Bacteriology*. 184(12):3338–3347.

Wibowo A, Becker C, Marconi G, Durr J, Price J, Hagmann J, Papareddy R, Putra H, Kageyama J, Becker J, Weigel D, Gutierrez-Marcos J. 2016. Hyperosmotic stress memory in *Arabidopsis* is mediated by distinct epigenetically labile sites in the genome and is restricted in the male germline by DNA glycosylase activity. *eLife*. 5:e13546.

796  Winter DJ. 2017. rentrez: an R package for the NCBI eUtils API. *The R Journal.* 9: 520–526.

# Supplementary Tables

| GO ID | GO term | Annotated | Significant | Expected | P-value |
|-------|---------|-----------|-------------|----------|---------|
| Yield at 24 °C | | | | | |
| GO:1901607 | Alpha-amino acid biosynthetic process | 10 | 5 | 1.24 | 0.003 |
| GO:0048519 | Negative regulation of biological process | 20 | 4 | 2.47 | 0.015 |
| GO:0044272 | Sulfur compound biosynthetic process | 8 | 3 | 0.99 | 0.063 |
| Yield at 38 °C | | | | | |
| GO:0048523 | Negative regulation of cellular process | 17 | 3 | 2.61 | 0.025 |
| GO:0001510 | RNA methylation | 5 | 3 | 0.77 | 0.027 |
| GO:0006508 | Proteolysis | 5 | 3 | 0.77 | 0.027 |
| GO:1905039 | Carboxylic acid transmembrane transport | 5 | 3 | 0.77 | 0.027 |
| GO:0065009 | Regulation of molecular function | 6 | 3 | 0.92 | 0.048 |

Supplementary Table S1: Results of the gene ontology (GO) enrichment tests based on the association between m6A methylation and phenotypes. Partially methylated m6A epiloci exhibiting a methylation fraction range $\geq$0.2 across sequenced samples were considered and the overlapping or closest ($\leq$500 bp) downstream gene, if any, was assigned to each of those epiloci. For each phenotypic trait and each epiloci, Spearman's $\rho$ was calculated between the trait values and the epiloci methylated fractions, and the corresponding uncorrected p-value was assigned to the gene associated to the epiloci. We then defined the set of genes of interest based on a threshold of Spearman's $p = 0.05$ and performed a GO enrichment test using `topGO` with the Fisher's exact test and the "weight01" algorithm (Alexa et al., 2006; Alexa and Rahnenfuhrer, 2020). GO terms with enrichment test $p \leq 0.1$ are reported for each trait (table continued on next page).

| GO ID | GO term | Annotated | Significant | Expected | P-value |
|---|---|---|---|---|---|
| Yield with ciliate | | | | | |
| GO:0048523 | Negative regulation of cellular process | 17 | 3 | 2.29 | 0.019 |
| GO:0006310 | DNA recombination | 6 | 3 | 0.81 | 0.034 |
| GO:0065009 | Regulation of molecular function | 6 | 3 | 0.81 | 0.034 |
| GO:0006974 | Cellular response to DNA damage stimulus | 17 | 5 | 2.29 | 0.062 |
| | | | | | |
| Yield with virus | | | | | |
| GO:0048523 | Negative regulation of cellular process | 17 | 5 | 1.72 | 0.009 |
| GO:0065009 | Regulation of molecular function | 6 | 3 | 0.61 | 0.015 |
| GO:0007154 | Cell communication | 14 | 4 | 1.42 | 0.026 |
| GO:0006790 | Sulfur compound metabolic process | 12 | 3 | 1.21 | 0.028 |
| GO:0006974 | Cellular response to DNA damage stimulus | 17 | 4 | 1.72 | 0.079 |
| GO:0006508 | Proteolysis | 5 | 2 | 0.51 | 0.081 |
| GO:0043648 | Dicarboxylic acid metabolic process | 5 | 2 | 0.51 | 0.081 |
| GO:1901564 | Organonitrogen compound metabolic process | 66 | 8 | 6.67 | 0.099 |

(Continued from previous page and on next page.)

| GO ID | GO term | Annotated | Significant | Expected | P-value |
|-------|---------|-----------|-------------|----------|---------|
| Growth rate at 24 °C | | | | | |
| GO:0044282 | Small molecule catabolic process | 14 | 4 | 1.26 | 0.017 |
| GO:0070887 | Cellular response to chemical stimulus | 7 | 3 | 0.63 | 0.018 |
| GO:0015672 | Monovalent inorganic cation transport | 6 | 2 | 0.54 | 0.093 |
| GO:0051301 | Cell division | 6 | 2 | 0.54 | 0.093 |
| GO:0006081 | Cellular aldehyde metabolic process | 6 | 2 | 0.54 | 0.093 |
| Growth rate at 31 °C | | | | | |
| GO:0042180 | Cellular ketone metabolic process | 5 | 3 | 0.41 | 0.004 |
| GO:0006081 | Cellular aldehyde metabolic process | 6 | 3 | 0.49 | 0.008 |
| GO:0016051 | Carbohydrate biosynthetic process | 14 | 3 | 1.15 | 0.035 |
| GO:0051301 | Cell division | 6 | 2 | 0.49 | 0.080 |
| GO:0006793 | Phosphorus metabolic process | 32 | 3 | 2.64 | 0.085 |
| GO:0044282 | Small molecule catabolic process | 14 | 3 | 1.15 | 0.097 |
| Growth rate at 38 °C | | | | | |
| GO:0044085 | Cellular component biogenesis | 23 | 5 | 1.98 | 0.053 |
| GO:0030031 | Cell projection assembly | 5 | 2 | 0.43 | 0.061 |
| GO:0042180 | Cellular ketone metabolic process | 5 | 2 | 0.43 | 0.061 |
| GO:0044238 | Primary metabolic process | 149 | 15 | 12.84 | 0.076 |
| GO:1903509 | Liposaccharide metabolic process | 10 | 2 | 0.86 | 0.085 |
| GO:0006396 | RNA processing | 11 | 2 | 0.95 | 0.086 |

(Continued from previous page and on next page.)

| GO ID | GO term | Annotated | Significant | Expected | P-value |
|---|---|---|---|---|---|
| Growth rate with DTT | | | | | |
| GO:0006091 | Generation of precursor metabolites and energy | 12 | 4 | 1.35 | 0.020 |
| GO:0006518 | Peptide metabolic process | 14 | 3 | 1.57 | 0.035 |
| GO:0009103 | Lipopolysaccharide biosynthetic process | 8 | 3 | 0.90 | 0.049 |
| GO:0044237 | Cellular metabolic process | 163 | 23 | 18.31 | 0.076 |
| GO:0009314 | Response to radiation | 10 | 3 | 1.12 | 0.089 |
| GO:0006979 | Response to oxidative stress | 5 | 2 | 0.56 | 0.099 |
| GO:0072528 | Pyrimidine-containing compound biosynthetic process | 5 | 2 | 0.56 | 0.099 |
| Growth rate with ciliate | | | | | |
| GO:0072528 | Pyrimidine-containing compound biosynthetic process | 5 | 3 | 0.77 | 0.027 |
| GO:0009263 | Deoxyribonucleotide biosynthetic process | 5 | 3 | 0.77 | 0.027 |
| GO:0006260 | DNA replication | 6 | 3 | 0.92 | 0.048 |
| GO:0006259 | DNA metabolic process | 13 | 5 | 2.00 | 0.070 |
| Growth rate with virus | | | | | |
| GO:0006508 | Proteolysis | 5 | 3 | 0.47 | 0.006 |
| GO:0009314 | Response to radiation | 10 | 3 | 0.94 | 0.068 |
| GO:0006400 | TRNA modification | 5 | 2 | 0.47 | 0.071 |
| GO:0006091 | Generation of precursor metabolites and energy | 12 | 4 | 1.12 | 0.090 |
| GO:0015980 | Energy derivation by oxidation of organic compounds | 6 | 2 | 0.56 | 0.092 |

(Continued from previous page.)

# Supplementary Figures

Supplementary Figure S1: Setup of the evolution experiment from which sequenced clones were isolated. Open arrows after experimental evolution indicates steps where evolved clones were grown under common garden conditions.

Supplementary Figure S2: Phenotypic traits measured in Ketola et al. (2013) and used in the present study. Each data point represents one evolved strain sequenced in this study. Bars and whiskers are mean values +/- 1.96 × s.e. for all sequenced strains within an evolutionary treatment.

Supplementary Figure S3: Relationship between coverage and estimated fraction of methylated adenines in GATC motifs. All adenines present in GATC motifs, from all sequenced strains, are included. Data is binned by intervals of methylated fraction of 0.1 width, as indicated on the $x$ axis. There is no correlation between coverage and estimated methylated fraction (Spearman's $\rho$ = -0.01). However, the slight increase in average coverage for the (0.1, 0.2) methylated fractions compared to the (0, 0.1) fractions suggests that at very low methylation levels (below 0.2), higher coverage is needed to estimate a methylated fraction other than 0.

Supplementary Figure S4: Detection of partially methylated GATC loci. Distribution of methylated fractions of adenines on boths DNA strands for GATC palindromes (showing data for all strains together). Left panel, all GATC palindromes shown; right panel, only GATC palindromes qualified as low methylation sites shown. The red arc in the left panel delimits the observations which are less four times the average quadratic distance to full methylation (point at (1,1)) away from full methylation. GATC palindromes are considered as low methylation sites if they lay outside this area (right panel).

Supplementary Figure S5: Distribution of genetic variants among sequenced *S. marcescens* clones. Each vertical lane represents the chromosome sequence of one clone. The locations along the chromosome are approximate in order to keep close loci visually distinct. Minor alleles of genetic variants are depicted. The numerical IDs of variants observed in more than one genome are shown in bold. Details about each variant are available in Table 1.