Cross-Infectivity of *Vorticella* across Genera of Mosquitoes for Development of

Biological Mosquito Control Strategies

Shelby Durden[1,2], Anthony Cruz [1], Wayne B. Hunter [2], Mustapha Debboun [3], Dagne Duguma[1,3]

[1]University of Florida/ IFAS, Florida Medical Entomology Laboratory, Vero Beach, Florida, USA.

[2]United States Department of Agriculture, Agricultural Research Service, U.S. Hort. Res. Lab.,

2001 South Rock Road, Fort Pierce, Florida, 34945. USA. Wayne.hunter@usda.gov ,
Orchid ID: 0000-0001-8603-3337

9 [3]Harris County Public Health, Mosquito and Vector Control Division, Houston, Texas, USA.

10 **Contact:** Dagne.Dugma@phs.hctx.net
11
12
13
Corresponding Author:

Shelby Durden,

United States Department of Agriculture/ ARS, 2001 South Rock Road, Fort Pierce, Florida

(772) 708-3756

Sdurden97@gmail.com

Keywords:

Vorticella, biocontrol, *Culex nigripalpus, Culex quinquefasciatus, Aedes albopictus*

ABSTRACT

Protozoans in general comprise about one-third of the parasitic species infecting arthropod vectors,

the role of free-living ciliates on mosquitoes have been insufficiently studied either due to their

low pathogenicity or being facultative parasites. Studies have shown that exposure of Paramecium

ciliate protists, like *Vorticella* species, to first instar Cx. nigripalpus larvae delayed larval

development and reduced biomass of emerged adults due to competition for food sources like

bacteria and other microbes essential to mosquito growth and survival. Thus, we report on the

capacity of a *Vorticella* protozoan's ability to cross-infect host species and parasitize multiple

mosquito larvae. The unique adapted behavior with the ability to remain on the exuviae in tree

hole habitats provides a novel delivery system to develop products for target species-specific

mosquitocides, larvicides, or viricides to be applied and sustained in aquatic systems.

Introduction

Millions of deaths occur annually due to mosquito-borne illnesses such as malaria with the heaviest

burden occurring in sub-Saharan Africa (WHO 2014a). Increasingly, emergence and reemergence

of mosquito-borne viral illnesses such as Dengue, West Nile virus (WNV), and Zika virus

epidemics, around the world are occurring (Gubler 2002; WHO 2014a,b; Liu and Zhou, 2017). To

reduce the mortality and economic losses caused by mosquito-borne diseases, there is a high

demand for new methods of mosquito control. Current mosquito control involves the use of

chemicals and biopesticides that target the larval and adult life stages of the mosquito. These

chemicals can range from organophosphates, insect growth regulators, or pyrethroids (Benelli

2015). However, past and current chemicals in use are having unintended negative effects on the

ecosystem such as the emergence of chemically-resistant mosquito populations. Resistance within

the population reduces the already diminished number of pesticides available for effective

mosquito control (Brogdon and McAllister 1998). Stahl (2002) also compiled a report that

presented the negative health and environmental risks of the four common pesticides used for

mosquito control: Scourge, Anvil, Permethrin, and Malathion. Therefore, there is a renewed focus

for alternative, more effective and environmentally friendly control strategies, including use of

biocontrol agents or strategies which provide two avenues of attack: directly reduce mosquito

populations or improve the efficacy of existing pesticides against mosquitoes.

During a field study of microbial and mosquito community dynamics, larvae of *Aedes* aegypti (L.),

Ae. albopictus (Skuse), *Culex nigripalpus* Theobald and *Cx. quinquefasciatus* Say (Diptera:

Culicidae) collected from various habitats, including artificial containers, tree holes, and

waterways in Florida were found to be infected with species of *Vorticella* (Duguma et al. 2017).

Over the course of four months during winter and early spring, *Vorticella* infectivity rates of

mosquito larvae were found to vary across time and the type of substrate found in the larval habitat

in Florida (D. Duguma, *unpublished data*). *Vorticella* is a genus of ciliates commonly found in

aquatic habitats in association with mosquito larvae and other zooplankton. This ciliate protest has

a contractile stalk used for attaching itself to substances *via* the means of a biopolymer glue (Cabral

et al. 2017). Attachment to mobile organisms, such as the mosquito larvae, gives *Vorticella* a

competitive advantage over other microbes in finding food (Kankaala and Eloranta 1987). Some

studies have shown a parasitic relationship between *Vorticella* and the mosquito larval host. For

example, Patil et al. (2016) demonstrated that *Anopheles stephensi* L. and *Ae. aegypti* larvae

inoculated with *Vorticella* showed reduced larval growth, slower development and adult

emergence. These findings indicated a potential use of *Vorticella* as a mosquito biocontrol agent

to augment chemical insecticides in use. While suggested by Patil et al. (2016) that *Vorticella* may

infect *Anopheles* and *Aedes* species, in this study we examined whether *Vorticella* from *Aedes*

genus infects *Cx. nigripalpus* and *Cx. quinquefasciatus* and whether it impacts the two species

differently. *Culex nigripalpus* and *Cx. quinquefasciatus* are prominent disease vectors in Florida

and southern United States for Saint Louis encephalitis and WNV (King et al. 1960, Day and Curtis

1994, Mores et al. 2007). The larval mosquito community was examined to identify major

Vorticella sites of attachment on the integument of the mosquitoes and whether it can transfer from

larval stages to adults.

Materials and Methods

Mosquito collection: Twenty *Vorticella*- infected *Ae. albopictus* late (3^{rd} to 4^{th}) instar larvae were

collected from a tree hole located on the property of Indian River Mosquito Control District

(coordinates 27.6661° N, 80.4438° W) and used as source of *Vorticella* for investigation in the

laboratory. The uninfected *Cx. nigripalpus* and *Cx. quinquefasciatus* larvae were hatched from egg

rafts collected from mesocosms located at the University of Florida, Institute of Food and

Agriculture Sciences, Florida Medical Entomology Laboratory (coordinates 27.58672, -

80.371069). Egg rafts were contained individually in wells of tissue culture plates in distilled water

until they hatched. The larvae were identified to species under a dissection microscope using

taxonomic keys from Darsie and Morris (2003). The larvae of each of the mosquito species were

transferred to plastic trays. Larvae were fed a diet of ground brewer's yeast and liver powder (1:1)

and kept in an incubator at 27± 1 °C.

Vorticella isolation and inoculation: The infected *Ae. albopictus* larvae collected from the field

were individually dissected using sterile forceps and vortexed in one mL of 1X Phosphate Buffer

Solution (PBS) solution (pH 7.2) to promote *Vorticella* suspension. The suspensions were each

dispensed among two groups: Group one was: 10 *Cx. nigripalpus*/container; and Group two was:

10 *Cx. quinquefasciatus* /container. Each group had 10 plastic containers, 250 mL each, with 200

mL of deionized water. Five untreated plastic buckets each containing 10 *Cx. nigripalpus*, or 10

Cx. quinquefasciatus were included to serve as control treatments. The experiment was carried out

in an incubator at 27 ± 1 °C and larval observations were made over a one month period for

Vorticella infection. At first sighting of larval pupation, the remaining larvae were observed for

infection by visual observation under a dissecting microscope, and monitored until their emergence

to adults. Sixteen infected larval samples (9 *Cx. quinquefasciatus* and 7 *Cx. nigripalpus)* were

placed individually in 1.0 ml of 200 proof ethanol for imaging. Light microscopy images were

taken using a Keyence VHX-5000 Digital Microscope (Keyence Corporation of America, Itasca,

IL, USA). Scanning Electron Microscope (SEM) images were taken using Hitachi S-4800

Scanning Transmission Electron Microscopy (STEM) (Hitachi High Technologies America, Inc.,

Peasanton, CA, USA) to closely examine parasitic relationship between *Vorticella* and *Ae.*

albopictus larval samples. All microscopy imaging was performed at the USDA-ARS, U.S.

Horticultural Research Lab, Electron Microscope Unit, Fort Pierce, FL, USA.

Data analysis: To determine differences between *Vorticella* infection rate of *Cx. nigripalpus* and

Cx. quinquefasciatus larvae, unpaired t-test was conducted. Non-parametric Kruskal-Wallis test

followed by Dunn's post hoc mean comparison was performed to determine differences in

mortality rate between larval mosquitoes subjected to *Vorticella* and mosquitoes subjected to

untreated control. A similar analysis was conducted for differences in pupation among different

larval groups. The difference in total number of adults emerged between *Cx. nigripalpus* and *Cx.*

quinquefasciatus larvae was evaluated using unpaired t-test. All statistical analyses and graphs

were conducted using GraphPad Prism 7 (GraphPad Prism Software Inc. San Diego, CA, USA).

Results

We examined *Vorticella's* ability to cross-infect across genera of mosquitoes and documented the

presence of *Vorticella* on different parts of larval mosquito hosts (**Fig. 1**). Using the procedure

described in a previous study (Patil et al. 2016), *Vorticella* isolated from *Ae. albopictus* larvae was

successfully transferred to two species of *Culex* mosquito larvae (i.e., *Cx. nigripalpus* and *Cx.*

quinquefasciatus). Upon microscopic examination, major sites of infestation were determined to

be along the larva's abdominal segments, the thorax and the siphon (**Fig. 1A**). Both dorsal and

ventral body parts of the second to fourth larval instars and exuviae were seen infected with

Vorticella but not pupae or adults. *Vorticella* remained on the exuviae of the host larva during its

metamorphosis to pupa suggesting that this *Vorticella* species might be restricted to immature

mosquito life stages. Their attachment to the exuviae may facilitate their transition to new larval

mosquito cohorts or other zooplankton.

The individual larvae were examined for *Vorticella* infestation at first sight of larval pupation. The

infection rate differed between the two *Culex* species with significantly greater mean number of

Cx. quinquefasciatus larvae infected with *Vorticella* than *Cx. nigripalpus* (t=3.4, df =18, *P>0.05*,

Fig. 2A). On average, 52% of *Cx. quinquefasciatus* larvae were infected with *Vorticella*, whereas,

only 22% of *Cx. nigripalpus* was infected with *Vorticella*. Larval mortality differed significantly

between *Vorticella* -infected and non-infected mosquito larvae (Kruskal-Wallis test, $P \leq 0.05$, **Fig.**

2B). *Culex nigripalpus* experienced greater mortality compared to *Cx. quinquefasciatus*. Greater

proportion (~20%) of *Cx. nigripalpus* larvae incurred larval mortality compared to only 10%

observed in *Cx. quinquefasciatus*. Mortality in the larvae of the non-infected *Cx. nigripalpus* larvae

was significantly lower than the larvae exposed to *Vorticella* suggesting that *Vorticella* may have

negative impact on this species. The mean proportion of mortality rates in unexposed control (i.e.,

without *Vorticella* treatment) of *Cx. nigripalpus* and *Cx. quinquefasciatus* was 0 and 13%,

respectively (**Fig. 2D**).

The difference in pupation rate among the four treatment groups was not statistically significant

although slightly greater pupation rate occurred in both infected and non-infected *Cx.*

quinquefasciatus and non-infected *Cx. nigripalpus* than infected *Cx. nigripalpus* larvae (Kruskal-

Wallis test, *P>0.05*, **Fig. 2C**). However, significantly a greater number (74%) of *Cx.*

quinquefasciatus emerged from the *Vorticella*-exposed larvae compared to 52% emergence rate

observed in *Cx. nigripalpus* ($P \leq 0.05$, **Fig. 2D**). These findings may suggest mosquito species-

specific effects of the *Vorticella* infestation and warrant further investigation.

Discussion

Vorticella is naturally occurring in freshwater environments which are also breeding sites for

mosquitoes. Although the exact biological relationship is unknown, *Vorticella* appears to have a

symbiotic relationship with the mosquito larvae to provide mobility to the ciliate and a competitive

survival advantage. Micks's (1955) study showed stunted growth and higher mortality rates in

Vorticella-infected *An. atroparvus* van Thiel larvae. The effects may be due to the *Vorticella*

secreting products toxic to the larvae which can cause pore formation in the larvae's bodies or that

the larvae are unable to remain on the water surface for tracheal breathing due to high levels of

Vorticella infestation (Micks 1955).

While protozoans in general comprise about one-third of the parasitic species infecting arthropod

vectors (Jenkins 1964), the role of free-living ciliates on mosquitoes have been insufficiently

studied either due to their low pathogenicity, or being facultative parasites. A previous study by

Duguma et al. (2017) found that exposure of *Paramecium* ciliate protists to first instar *Cx.*

nigripalpus larvae delayed larval development and reduced biomass of emerged adults due to

competition for food sources such as bacteria and other similar-sized microbes found essential to

the growth of mosquitoes (Duguma et al., 2019). In the absence of *Paramecium*, the mosquito

larvae harvested bacterial and other similar small-sized organic particles more efficiently than

when found in association with the protists. Other studies have shown a severe competition for

food between Vorticellid epibionts and Daphinids (Kankaala and Eloranta 1987). In addition, the

heavy infestation of *Vorticella* hampers mobility of its host subjecting the host susceptible to

predation (Kankaala and Eloranta 1987). Although we have not measured growth performances of

larvae in our study, the higher mortality observed in *Cx. nigripalpus* larvae may be attributed to

competition by *Vorticella* for bacterial food sources in addition to their possible physiological

effects on the larvae. However, a similar effect was not observed in *Cx. quinquefasciatus*

suggesting that their effect might be species-specific. These findings show promise in the

utilization of ciliates in mosquito population control.

The protozoan's ability to cross-infect and parasitize multiple mosquito larvae and its ability to

remain on the exuviae provides a unique delivery system for novel species-specific mosquitocides,

or viricides to be applied and sustained in aquatic systems. The need for studies to evaluate these

advantages of *Vorticella* as a potential delivery system, as a form of biological control to reduce

mosquitoes and the spread of vector-borne pathogens, such as Dengue, Zika and West Nile Virus

are being pursued as a sustainable mechanism for mosquito control.

Acknowledgements

We thank Maria T. Gonzalez, Senior Technician, USDA, STEM Microscopy unit, Fort Pierce,

FL., Jacqueline L. Metz, technical assistance, USDA, and Matthew Hentz for instruction on the

VHX-5000 digital microscope, and Dr. Joseph Krystel for his valuable comments. This study was

supported by funding from Florida Department of Agriculture and Consumer Services.

This research was supported in part by the U.S. Department of Agriculture, Agricultural Research

Service." Mention of trade names or commercial products in this publication is solely for the

purpose of providing specific information and does not imply recommendation or endorsement by

the U.S. Department of Agriculture. USDA is an equal opportunity provider and employer.

References

Benelli, G. Plant-borne ovicides in the fight against mosquito vectors of medical and veterinary
importance: a systematic review. *Parasitol Res* 2015; *114*: 3201–3212.
199

200   Brogdon, W.G., McAllister, J.C. Insecticide resistance and vector control. *Emerging Infect Dis*
201   1998; *4*: 605-613.
202

203   Cabral, A.F., Utz, L.R.P., Velho, L.F.M. Structure and distribution of ciliate epibiont
204   communities in a tropical floodplain. *Hydrobiologia* 2017*; 787***: 167-180.
205

206   Day, J.F., Curtis, G.A. When it rains, they soar--and that makes *Culex nigripalpus* a
207   dangerous mosquito. *American Entomologist* 1994; *40*:162-167.
208

209   Darsie, R.F., Morris, C.D. Keys to the adult females and fourth Instar Larvae of the
210   mosquitoes of Florida (Diptera, Culicidae). *Tech Bull Fla Mosq Control Assoc* 2003; *1*: 1-159.
211

Duguma, D., Kaufman, M.G. Domingos, S., Arthur, B. Aquatic microfauna alter larval
food resources and affect development and biomass of West Nile and Saint Louis encephalitis

vector, *Culex nigripalpus* (Diptera: Culicidae). *Ecology and Evolution* 2017; *7*: 3507-3519.

Duguma, D, Hall, M.W., Smartt, C.T, Debboun, M., Neufeld, J.D. Microbiota variations in *Culex*
nigripalpus disease vector mosquito of West Nile virus and Saint Louis Encephalitis from different
geographic origins 2019, *PeerJ* 6:e6168 Doi 10.7717/peerj.6168.

Gubler, D.J. The global emergence/resurgence of arboviral diseases as public health
problems. *Arch Med Res* 2002; *33*: 330-342.

Jenkins, D.W. Pathogens, parasites, and predators of medically important arthropods. *Bull*
World Health Organization 1964; *30*: 1-50.

Kankaala, P., Eloranta, P. Epizooic ciliates (*Vorticella* sp.) complete for food with their
host *Daphnia-Longispina* in a small polyhumic lake. *Oecologia* 1987; *73***:** 203-206.

King, W.V., Bradley, G.H., Smith, C.N., McDuffie, W.C. A handbook of the mosquitoes of
the Southern United States. *United States Department Agriculture* 1960; *173*: 118-119.

Liu, Z., Zhou, T. Competence of *Aedes aegypti*, *Ae. albopictus*, and *Culex*
quinquefasciatus mosquitoes as Zika virus vectors, China. *Emerging Infectious Dis* 2017;
23:1085-1091.

Mores, C.N., Turell, M.J., Dohm, D.J.. Blow, J.A., Carranza, M.T,. Quintana, M. Experimental
transmission of West Nile virus by *Culex nigripalpus* from Honduras. *Vector Borne Zoonotic*
Diseases 2007; *7*: 279-84.

Micks, D. *Vorticella* infestation of *Anopheles atroparvus* larvae. *Journal of Economic Entomology*
1955*;48*: 215–216.

Patil, C.D., Narkhede, C.P., Suryawanshi, R.K., Patil, S.V. *Vorticella* sp: Prospective
mosquito biocontrol agent. *Journal of Arthropod-Borne Diseases* 2016*; 10*: 602-607.

Stahl, A. The health effects of pesticides used for mosquito control. *Citizens Campaign for the*
Environment and Citizens Environmental Research Institute 2002*; 1*: 1-15.

WHO. World Malaria Report. *World Health Organization* 2014a.

WHO. A global brief on vector-borne diseases. *World Health Organization* 2014b.

Figure 1. *Aedes albopictus* **(A & D)** larval samples collected from tree holes were found to be highly infested with clusters of *Vorticella*. Image **D** shows a close-up of major *Vorticella* infestation along the abdominal segments of the mosquito larva. Scanning Electorn Microsopy (SEM) images of *Vorticella* on mosquito larvae indicate polyp colony **(B)** or a singles **(C)**. A single stalk form averaged 31 µm in width and 55 µm in length **(E)**. Images of early instar *Cx. nigripalpus* **(D)** and *Cx. quinquefasciatus* **(F)** larvae infected with *Vorticella* in the laboratory. The *red arrow* in image **C,** depicts *Vorticella* attachment as a single stalk on *Cx. nigripalpus* larvae.

Fig. 2. Box-and Whisker Tukey plot of mosquito *Vorticella*-infection experiment showing **(A)** infection susceptibility difference between larvae of *Cx. quinquefasciatus* and *Cx. nigripalpus*, **(B)** proportion of larvae died in both *Vorticella*-infected and control (non-infected) larvae of the two species, **(C)** proportion of mosquito larvae developed to pupae from larvae infected with *Vorticella* and control larvae, and **(D)** proportion of larvae emerged to adults following *Vorticella* infection. ***** Indicates statistically significant differences at p ≤ 0.05, whereas **ns**= not statistically significant (p>0.05).