Whole-genome duplication and host genotype affect rhizosphere microbial communities

Julian C. B. Ponsford [a], Charley J. Hubbard [a,b], Joshua G. Harrison[a], Lois Maignien[c,d] C. Alex

Buerkle [a,b], Cynthia Weinig[a,b,e]#

a.   Department of Botany, University of Wyoming, Laramie, WY, USA

b.   Program in Ecology, University of Wyoming, Laramie, WY, USA

c.   Marine Biological Laboratory, Josephine Bay Paul Center, Woods Hole, MA, USA

d.   Laboratory of Microbiology of Extreme Environments, UMR 6197, Institut Européen de

8   la Mer, Université de Bretagne Occidentale, Plouzane, France

e.   Department of Molecular Biology, University of Wyoming, Laramie, WY, USA

Corresponding author: Cynthia Weinig, cweinig@uwyo.edu

Keywords: *Arabidopsis thaliana,* whole genome duplication, multinomial modeling, plant

microbe interactions

Funding: National Science Foundation grant: IOS-1444571 to LM and CW & EPS-16557226 to

CW and CAB.

Abstract

The composition of complex microbial communities found in association with plants is influenced in part by host phenotype. Yet, the salient genetic architecture is often unknown. Genome duplication events are common in the evolutionary history of plants, influence many important plant traits, and may affect associated microbial communities. Using experimentally induced whole genome duplication (WGD), we tested the effect of WGD on rhizosphere bacterial communities in *Arabidopsis thaliana*. Specifically, we performed 16S rRNA amplicon sequencing to characterize differences between microbiomes associated with specific host genotypes (Columbia *vs*. Landsberg) and ploidy levels (diploid *vs*. tetraploid). We modeled abundances of individual bacterial taxa by utilizing a hierarchical Bayesian framework, based on the Dirichlet and multinomial distributions. We found that host genotype and host ploidy level affected rhizosphere community composition, for instance, the microbiome of the tetraploid Columbia genotype differed from that of other host genotypes. We then tested to what extent microbiomes derived from a given host genotype or ploidy level affected plant performance by inoculating sterile seedlings of each genotype with microbial communities harvested from a prior generation. We found a negative effect of the tetraploid Columbia microbiome on growth of all four plant genotypes. The findings suggest that while both host genotype and ploidy affect microbial community assembly, bacterial communities found in association with only some host genotypes may affect growth of subsequent plant generations.

Importance

Plants influence the composition of their associated microbial communities; yet the underlying

host genetic factors are often unknown. Genome duplication events are common in the

evolutionary history of plants and affect many plant traits, including the quality and quantity of

compounds exuded into the root zone, which can affect root-bound microbes. In *Arabidopsis*

thaliana, we characterized how whole-genome duplication affected the composition of

rhizosphere bacterial communities, and how bacterial communities associated with two host

plant genotypes and ploidy levels affected subsequent plant growth. We observed an interaction

in which ploidy level within one host genotype affected both bacterial community composition

and function. This research reveals how genome duplication, a widespread genetic feature of

both wild and crop plant species, influences the coexistence of bacterial taxa and affects plant

growth.

Introduction

Plant-microbe interactions can exhibit a complete feedback cycle, in which changes in

the microbiome affect plant performance and the genetics of the plant host alter microbial

community composition (1–3). The traits of bacterial taxa in plant-associated communities

affect both host fitness and ecological function (4). Rhizosphere bacteria, in particular, affect

many aspects of plant performance, such as increasing access to nutrients (5), relieving abiotic

and biotic stress (6), and promoting growth (7). Even slight changes in the rhizosphere

microbiome can affect host plant performance (8). For instance, Korir *et al.* (2017) found that

increased abundance of one taxon, *Bacillus megaterium*, led to enhanced nitrogen access and

growth of *Phaseolus vulgaris* in field conditions (9). Wholesale changes in the abundance of taxa

comprising rhizosphere microbiomes also affects plant performance (10, 11). For example, in

Arabidopsis thaliana, differences in microbial community composition attributable to past or

nearby plant communities strongly influenced host growth (12, 13).

A plant host's genetic background can also affect the composition of microbiomes

consisting of thousands of taxa (14–16). For instance, over a range of environmental conditions,

host genotype in maize explains on average ~19% of the variance in relative abundance of root

microbial taxa (17). Among host genotypes, allelic variation segregating at loci with diverse

functions could potentially contribute to differences in rhizosphere community composition. The

contribution of specific host-plant loci and pathways has been demonstrated experimentally

using genetic knock-outs or transgenic overexpression (18, 19). Genetic manipulation to shift the

plant circadian clock by $\pm$ 4 hours explained ~22% of the variance in rhizosphere bacterial

communities among experimental *Arabidopsis* lines (13). However, the number of studies

characterizing causal genetic factors is small, and the extent to which specific host genetic and

genomic features alter the assembly and function of microbial communities remains largely

unknown, despite the demonstrably important effect of the rhizosphere microbiome on host plant

performance.

Plants influence rhizosphere microbial communities via root exudation of small

molecular-weight organic compounds (20–22). For simple symbioses consisting of one plant and

one microbial taxon, such as between legumes and *Rhizobia,* the host genetic mechanisms that

underlie these interactions are increasingly well characterized (23–25). How more complex

genetic mechanisms affect associated root microbial communities is less clear. Whole-genome

duplication (WGD) is one genetic feature of host plants that could potentially influence

microbial community composition via root exudation. WGD can lead to changes in cell size, life-

history, and physiology, which may influence numerous other phenotypes (26, 27). WGD also

occurs naturally in wild populations and is ubiquitous in the evolutionary history of plants and in

the domestication history of many crop species (28).

In laboratory settings, colchicine is used to induce autopolyploidization and create stable

lines of tetraploids in *A. thaliana.* This mutation theoretically produces no genetic changes

besides those associated with genome duplication (29). Therefore, the comparison of rhizosphere

microbiomes between *A. thaliana* tetraploid genotypes developed from inbred diploid lines

provides a basis for testing the effects of host ploidy level on microbial community composition,

without the confounding effects of allelic variation or fixed heterozygosity between tetraploids

and diploid progenitors. Tetraploidy can induce changes in core metabolic pathways such as the

tricarboxylic acid cycle (TCA), malate and citrate concentrations, and potassium uptake in *A.*

thaliana (30, 31). These phenotypic shifts may alter root exudate profiles or other phenotypes

that consequentially change microbial colonization and community composition.

In the current study, we were interested in characterizing the effects of host WGD on

rhizosphere bacterial communities and testing how changes in bacterial community composition

affected plant performance. We hypothesized that WGD would affect bacterial communities due

to changes in host physiology induced by polyploidization. Further, if tetraploid plants assemble

disparate communities compared to diploid hosts, then we predicted that differences in these

communities could affect plant performance.

Material and Methods

Plant material and growth conditions

To test the effects of whole genome duplication and plant genotype on rhizosphere

bacterial community composition and plant host performance, we selected two *A. thaliana*

diploid genotypes: Columbia (Col-2x) and Landsberg *erecta* (Ler-2x) and their tetraploid

counterparts (Col-4x CS922178) and (Ler-4x CS3900). Tetraploid genotypes were multiple

generations removed from initial colchicine treatment, and therefore can be assumed to be

mutationally stable (32). In our first experiment, each genotype was planted into sterilized

potting mix and inoculated with a microbial community from the Catsburg region of Durham,

North Carolina, USA (36.0622^0N, -78.8496^0W), a site with a history of *A. thaliana* growth. The

microbiomes associated with each host plant genotype were characterized by 16S rRNA gene

amplicon sequencing in the first experiment and tested for effects on growth of a second

generation of plants in a second experiment.

In both experiments, seeds were surface sterilized using a solution of 15% bleach, 0.1%

Tween, and autoclave sterilized reverse osmosis purified water (RO H_2O). Seeds were

thoroughly rinsed in RO H_2O to remove any remaining detergents. Seeds were placed in 1.5mL

tubes containing 1mL RO H_2O and stored at 4°C for seven days and then placed on greenhouse

benches under natural 14 hr photoperiods to induce synchronous germination (33). On the day

root radicles were observed, seeds were transferred to 2-inch diameter net pots filled with a

mixture of autoclaved Redi-Earth Potting Mix (Sungro Horticulture, Agawam, MA, USA) and 2

mL of liquid inoculant (described below). To ensure sterility, potting mix was autoclaved on a

wet cycle for 60 min at 121°C, allowed to rest for at least one hour and autoclaved for another 60

min. No microbes could be cultured on tryptic soy agar media using serial dilutions of

autoclaved soil as inoculum. To create inoculants for the first experiment, 60 g of the Catsburg

soil was mixed with 540 mL of autoclaved RO H_2O and sieved through 1000 µm, 212 µm and

45 µm sterile sieves to remove nematodes that could potentially affect plant performance (34).

The Catsburg site not only has a well-documented occurrence of naturalized *A. thaliana*

populations but also has a soil pH approximately identical to the potting substrate, which

minimized potential selection by the common soil matrix on microbial community composition

(35–37).

For the second experiment, we used four randomly selected plant-conditioned soil

samples from each host genotype to create inoculums. Soil was combined by genotype, manually

homogenized, and used to create inoculum as above. In all experiments, after ten days of growth,

seedlings were thinned to one plant per pot. All experiments were performed at the Williams

Conservatory or Agricultural Experiment Station at the University of Wyoming.

<u>Experimental design</u>

Experiment 1: Effects of host plant genotype and ploidy on microbial community

To measure the influence of host ploidy on rhizosphere bacterial community composition, 20

replicates of Col-2x, Col-4x, Ler-2x, and Ler-4x were planted in a fully randomized four block

design. To avoid confounding the effects of ploidy with developmental stage, eight rhizosphere

samples from each genotype were collected on day 24, by which time all plants displayed visible

elongation of the primary inflorescence from the apical meristem (N = 32). Unharvested plants

were allowed to grow until senescence to quantify additional traits. To account for variation in

microbial communities due to greenhouse conditions and intrinsic variation in soil independent

of plant host effects, we collected soil from empty pots containing only soil that were potted

simultaneously with our experiment (N = 7).

Experiment 2: Microbial effects on plant performance

To investigate how host performance is affected by microbiomes associated with specific

host genotypes, we used a fully factorial design with four host plant genotypes grown in four

host-influenced microbiomes (harvested from the host genotypes in expt. 1). 40 replicates of

each plant genotype were grown in sterilized potting mix inoculated with one of the four

rhizosphere microbiomes from plant-conditioned soil in a fully randomized block design. Plants

were checked daily for bolting and flowering. Upon the first plant bolting, a subset of plants was

collected for above- and belowground biomass (N = 382) to measure plant growth and resource

allocation. A subset of plants were allowed to senesce and seeds harvested to quantify seed mass

(N = 152). All phenotypic measurements were performed following Rubin et al (38).

DNA extraction and sequencing

Rhizosphere samples were collected from the root surface as described in (39) by first

uprooting plants, manually agitating plants, removing loosely adhering soil particles, placing

roots with closely adhering soil in a buffer of sterile PBS with 0.1% Silwet in a 15 mL tube, and

vortexing for 10 min on maximum speed. Rhizosphere samples were centrifuged at 3714 rcf, and

a total soil mass of no more than 250 mg was transferred from each tube into a sterile Qiagen

PowerSoil (Qiagen, Valencia, CA) bead tube. DNA was extracted following manufacturer's

instructions. Extracted DNA was shipped to the Marine Biological Laboratories (Woods Hole,

MA, USA) for amplification and sequencing of the V4-V5 (primers 518f and 926r) region of the

16S rRNA gene (40) on an Illumina MiSeq (Illumina, San Diego, CA, USA).

Sequence and data analysis

We used the R package *dada2* (ver. 1.5.8) to filter and trim reads based on quality,

estimate the error rate using 1 000 000 reads, dereplicate reads, infer amplicon sequence variants

(ASV), merge paired end reads, remove chimeras, and assign unique sequences to taxa using the

Silva 16S database (ver. 128) (41, 42). Next, we ascertained the relative abundance of

operational taxa by quantifying unique sequences and amplicon sequence variants using the

default settings in the *microbiome* R package (43). Raw sequences were uploaded to the NIH

NCBI Short Read Archive under project ID PRJNA474006.

Dissimilarity analyses identify coarse-grained changes in community composition across

treatments, which arise from the combined effects of all community members. Yet, significant

differences in relative abundance may exist when comparing individual taxa between treatments

that are not discernible when aggregating effects of whole communities. To test for differences

in the relative abundances of individual taxa between treatment levels (contrasts of host ploidy

and host genotype), we used a hierarchical Bayesian modelling approach reliant upon the

Dirichlet and multinomial distributions (44, 45). The Dirichlet-multinomial model (DMM)

explicitly accounts for the compositional nature of community sampling (whether in sequencing

data or any finite number of taxa observations from an assemblage), readily allows information-

sharing among replicates within a category and obtains estimates of the relative abundance of

each microbial taxon, while propagating uncertainty in those estimates. We estimated differences

in the relative abundance of each microbial taxon between experimental treatments through

comparison of parameter estimates as per Harrison *et al.* (2020) (44).

Briefly, DMM estimates the multinomial parameters describing the relative abundances

of each taxon in a replicate, denoted as a vector of parameters (p). These multinomial parameters

were informed by a Dirichlet distribution, with parameters characterizing the expected

frequencies of each taxon ($\pi * \Theta$), where π is a vector describing the expected relative abundance

of each taxon in the sampling group and Θ is an intensity parameter that models among-replicate

variation (44, 46). To quantify differences in the relative abundance of taxa between groups, we

calculated the posterior probability distribution for the difference in π_i parameters between

groups for some taxon *i*. The probability of an effect of treatment on a focal taxon can be

determined by the location of zero in this distribution of differences. Following convention, if

95% or more of the distribution did not overlap zero, then there was little evidence that the

relative abundance of the focal taxon differed between treatment groups.

The DMM offers several advantages over existing analytical methodologies. First,

parameters were estimated while propagating uncertainty, thus avoiding cumbersome multiple

comparison correction, and precluding the use of p-values. Second, information was shared

among replicates and treatment groups. Third, rarefaction is unnecessary because analyses are

performed on taxon frequencies, which are estimated using information from all replicates within

a sampling group. Fourth, the Dirichlet and multinomial distributions have interdependent

parameters that reflect the compositional nature of sequencing data, thus we model the

composition as a whole as opposed to estimating the relative abundance of each taxon separately,

as in DESeq2 (47).

DMM was specified in the Stan probabilistic programming language and implemented

through the rstan *(*ver. 2.18.2) package in the R statistical computing environment (ver. 3.6.0)

(48, 49). For each of four chains, the sampler was run for 1500 steps as a burn-in period and was

followed by an additional 1000 steps (a total of 4000 samples were drawn from the posterior

distributions of focal parameters [π from each sampling group]). The Gelman-Rubin statistic was

computed to measure convergence among chains (50). Separate models were constructed for the

following comparisons: all diploid versus all tetraploid hosts, Columbia versus Landsberg *erecta*,

and Columbia tetraploid versus all other treatments (pooled).

For those taxa that differed credibly among our treatment groups, we used NCBI BLAST

to confirm 16S rRNA identifications generated using the SILVA database (51, 52). We report

species level identification in cases where reads align 100% with entries in the 16S ribosomal

RNA sequences database, and report bacterial family if genus level resolution is not corroborated

by SILVA (53).

To assess effects on plant performance in experiment 2, we used fixed effect, three-way

ANOVAs, where inoculum, genotype, and block were the explanatory variables. For purposes of

data representation, residuals were plotted after statistically accounting for the effect of genotype

and block. Finally, we used planned comparisons to contrast plant performance between plants

grown in the Col-2x versus Col-4x inoculums and Ler-2x versus Ler-4x inoculums.

Results

Sequencing results

After quality filtering and chimera removal in *dada2* as well as removal of bacterial taxa

occurring in plant-less pots and removal of 16S rRNA sequences from chloroplasts,

mitochondria, and Eukaryotes, 1 647 741 processed reads remained from an initial 3 282 410

raw reads. We retained 36 033 reads to 143 254 reads per plant rhizosphere (**Supplemental**

Figure 1). In total, there were 2689 microbial taxa (ASVs) present.

We utilized DMM to identify individual taxa with significantly different relative

abundance due to host genotype and host ploidy level. This model allowed us to quantify

differences in the abundance of individual taxa among treatments that might not be evident in

whole community dissimilarity analyses. Based on this approach we identified 25 taxa that were

significantly enriched in both Ler genotypes and 29 taxa that were more abundant in the Col

genotypes (**Figure 1a**). A taxon of genus *Pedobacter* was the most highly enriched in Ler

associated bacterial communities with an 8-fold increase. In Col bacterial communities, the most

enriched taxon was *Pseudomonas* at a 4.8-fold increase in abundance. DMM also identified 17

taxa that were significantly enriched in all diploid genotypes. Of these taxa, the most enriched

taxon was a member of the genus *Bacillus,* enriched 3.1-fold relative to tetraploid communities.

Tetraploid communities' most enriched member was *Mucilaginibacter*, which was present at 4-

fold greater relative abundance compared to rhizosphere microbiomes in diploids. 23 taxa were

significantly enriched across both tetraploid genotypes (**Figure 1b**). Our model identified 16 taxa

as higher abundance in the Columbia tetraploid rhizosphere and 30 taxa enriched across the

bacterial microbiomes of the other three host genotypes. In Col-4x, a bacterium identified as

Sphingomonas had the highest change in abundance at 6-fold. Conversely, a taxon of genus

Pedobacter was enriched in all other microbiomes. This taxon appeared at 5.8-fold greater

frequency in these microbiomes than Col-4x. Across all comparisons, ASVs within the phylum

Proteobacteria were more commonly differentially abundant than taxa within any other phylum.

Effects of rhizosphere microbes on plant performance

When microbiomes from plant-conditioned soil were harvested following experiment 1,

and used as inoculant in experiment 2, plants grown in soils inoculated with the Col-4x

microbiome had significantly lower aboveground and belowground biomass compared to plants

grown in soils inoculated with Col-2x microbiome ($P < 0.001$ & $P = 0.012$) or microbiomes

influenced by either of the two Ler genotypes (**Figure 2ab**; $P < 0.001$ & $P < 0.001$). The soil

microbiome had no effect on phenological characteristics or fruit number (**Supplemental Table**

2).

Discussion

Whole genome duplication is estimated to have occurred across 30-70% of the

angiosperm phylogeny over its evolutionary history and is a common genetic feature of many

economically important plants (54). We found that host genotype and WGD influenced

numerous individual taxa within the rhizosphere microbiome. Furthermore, host-genotype

specific effects on rhizosphere microbiomes influenced plant growth. In contrast to prior studies

(55), our experiments also utilized a soil with a history of *Arabidopsis* occurrence as inoculant.

This should, in principle, allow genotypes to assemble root microbiota from the existing soil

microbiomes.

Previous studies examining the root microbiota found in association with common lab

accessions of *A. thaliana* have found a link between genotype and rhizosphere bacteria (15, 20).

These studies report a < 10% shift in composition between the Ler and Col genotypes, which we

corroborate here. As noted above, we identified many individual taxa associated with host

ploidy level (**Figure 1a**). This provides support for the hypothesis that these specific taxa are

directly responsive to genome duplication. Although the causal physiological mechanisms

remain unclear in the current study, one candidate mechanism is root exudation. Previous studies

suggest that carbon root exudates strongly affect the occurrence and abundance of microbial taxa

in the rhizosphere (2, 56), and these exudates can be influenced by genome duplication (30).

Consequently, shifts in root exudates, or other metabolites produced by the plant, in response to

WGD could be partially responsible for the association between ploidy and the relative

abundance of microbial taxa that we observed (57, 58). Given that the number of microbial taxa

with a demonstrably significant response to WGD is in the 10's rather than 100's or 1000's, we

conclude that while host ploidy level affects the abundance of some taxa, it may not lead to

broad-scale changes in the rhizosphere microbiome.

We found that differences in the rhizosphere bacterial microbiome lead to differences in

host performance [**Figure 2**: 23, 61, 62]. Here, plants grown in soils inoculated with microbial

rhizosphere communities harvested from Col-4x plants had reduced vigor compared to plants

grown in soils inoculated with microbial communities from all other genotypes (**Figure 2**). One

bacterial species, *Novaherbaspirillium autotrophicum*, was identified as being significantly

enriched in the Columbia tetraploid genotype. *N. autotrophicum* was identified as a facultative

autotroph capable of denitrification, isolated from rice paddy fields (61). The greater abundance

of this microbe in the Columbia tetraploid community may point to changes in rhizobial

community in plants inoculated with this microbiome. The greater abundance of *N.*

autotrophicum may result in a decrease in available nitrogen in the rhizosphere causing lower

overall growth in the plant host. This loss of biomass may have effects on nutrient cycling or

survivability in stressful environments. Further, the effect of taxa such as *N. autotrophicum* may

be especially impactful in the context of mixed populations of diploid and tetraploid individuals,

where stressful conditions could be compounded or mitigated by host-associated microbes (62).

Conclusions

In sum, we have shown that whole genome duplication influences rhizosphere bacterial

community composition by affecting the relative abundance of certain bacterial taxa. In some

genetic backgrounds, plant ploidy level affects microbial function, as indicated by enhanced

above and belowground plant biomass in the Col-2x versus Col-4x genotypes. Multiple bacterial

taxa are affected by host ploidy level, irrespective of host genetic background. The ASVs

enriched (or reduced) in the rhizosphere of tetraploid hosts could help explain the advantages

tetraploids appear to have in stressful environments (63, 64). Given the prevalence of WGD

among plants in both natural and agricultural systems, our results highlight a novel mechanism

by which plant evolutionary history influences the root-associated microbiome, thereby affecting

plant phenotype.

Acknowledgments

Computing was performed in the Teton Computing Environment at the Advanced Research

Computing Center, University of Wyoming, Laramie (https://doi.org/10.15786/M2FY47). This

work was supported by the National Science Foundation grants IOS-1444571 to CW, LM and

EPS-16557226 to CW and CAB. We would like to thank Lindsay Leverett for collecting soil,

and Meredith Pratt and Ryan Pendleton for help with experimental and greenhouse maintenance

Conflict of Interests

The authors report no conflicts of interest.

References

1.   Schlaeppi K, Bulgarelli D. 2015. The Plant Microbiome at Work. Mol Plant-Microbe
     Interact MPMI 212:212–217.

2.   Chaparro JM, Badri D V, Vivanco JM. 2013. Rhizosphere microbiome assemblage is
     affected by plant development. ISME J 8:790–803.

3.   Van Nuland ME, Bailey JK, Schweitzer JA. 2017. Divergent plant–soil feedbacks could
     alter future elevation ranges and ecosystem dynamics. Nat Ecol Evol 1:0150.

4.   Talbot JM, Bruns TD, Taylor JW, Smith DP, Branco S, Glassman SI, Erlandson S,
     Vilgalys R, Liao H, Smith ME, Peay KG. 2014. Endemism and functional convergence
     across the North American soil mycobiome. Proc Natl Acad Sci U S A 111:6341–6.

5.   Richardson AE, Simpson RJ. 2011. Soil Microorganisms Mediating Phosphorus
     Availability Update on Microbial Phosphorus. Plant Physiol 156:989–996.

6.   Mendes R, Kruijt M, de Bruijn I, Dekkers E, van der Voort M, Schneider JHM, Piceno
     YM, DeSantis TZ, Andersen GL, Bakker PAHM, Raaijmakers JM. 2011. Deciphering the
     Rhizosphere Microbiome for Disease-Suppressive Bacteria. Science (80-) 332:1097–
     1100.

7.   Bulgarelli D, Schlaeppi K, Spaepen S, van Themaat EVL, Schulze-Lefert P. 2013.

Structure and Functions of the Bacterial Microbiota of Plants. Annu Rev Plant Biol

64:807–838.

8.   Henning JA, Weston DJ, Pelletier DA, Timm CM, Jawdy SS, Classen AT. 2016. Root

bacterial endophytes alter plant phenotype, but not physiology. PeerJ 4:e2606.

9.   Korir H, Mungai NW, Thuita M, Hamba Y, Masso C. 2017. Co-inoculation Effect of

Rhizobia and Plant Growth Promoting Rhizobacteria on Common Bean Growth in a Low

Phosphorus Soil. Front Plant Sci 08:141.

10.   Wagner MR, Lundberg DS, Coleman-Derr D, Tringe SG, Dangl JL, Mitchell-Olds T.

2014. Natural soil microbes alter flowering phenology and the intensity of selection on

flowering time in a wild Arabidopsis relative. Ecol Lett 17:717–726.

11.   Zolla G, Badri D V., Bakker MG, Manter DK, Vivanco JM. 2013. Soil microbiomes vary

in their ability to confer drought tolerance to Arabidopsis. Appl Soil Ecol 68:1–9.

12.   Sugiyama A, Bakker MG, Badri D V., Manter DK, Vivanco JM. 2013. Relationships

between Arabidopsis genotype-specific biomass accumulation and associated soil

microbial communities. Botany 91:123–126.

13.   Hubbard CJ, Brock MT, van Diepen LT, Maignien LL, Ewers BE, Weinig C. 2017. The

plant circadian clock influences rhizosphere community structure and function. ISME J

doi:in press.

14.   Wagner MR, Lundberg DS, Del Rio TG, Tringe SG, Dangl JL, Mitchell-Olds T. 2016.

Host genotype and age shape the leaf and root microbiomes of a wild perennial plant. Nat

Commun 7:1–15.

15.   Lundberg DS, Lebeis SL, Paredes SH, Yourstone S, Gehring J, Malfatti S, Tremblay J,

367   Engelbrektson A, Kunin V, Rio TG del, Edgar RC, Eickhorst T, Ley RE, Hugenholtz P,

368   Tringe SG, Dangl JL. 2012. Defining the core Arabidopsis thaliana root microbiome.

369   Nature 488:86–90.

370   16.   Marques JM, da Silva TF, Vollu RE, Blank AF, Ding G-C, Seldin L, Smalla K. 2014.

371   Plant age and genotype affect the bacterial community composition in the tuber

372   rhizosphere of field-grown sweet potato plants. FEMS Microbiol Ecol 88:424–435.

373   17.   Peiffer JA, Spor A, Koren O, Jin Z, Tringe SG, Dangl JL, Buckler ES, Ley RE. 2013.

374   Diversity and heritability of the maize rhizosphere microbiome under field conditions.

375   Proc Natl Acad Sci 110:6548–6553.

376   18.   Berendsen RL, Pieterse CMJ, Bakker PAHM. 2012. The rhizosphere microbiome and

377   plant health. Trends Plant Sci 17:478–486.

378   19.   Agler MT, Ruhe J, Kroll S, Morhenn C, Kim S-T, Weigel D, Kemen EM. 2016. Microbial

379   Hub Taxa Link Host and Abiotic Factors to Plant Microbiome Variation. PLoS Biol

380   14:e1002352.

381   20.   Jones DL, Nguyen C, Finlay RD. 2009. Carbon flow in the rhizosphere: Carbon trading at

382   the soil-root interface. Plant Soil.

21.   Walker TS. 2003. Root Exudation and Rhizosphere Biology. PLANT Physiol 132:44–51.

22.   Dennis PG, Miller AJ, Hirsch PR. 2010. Are root exudates more important than other

sources of rhizodeposits in structuring rhizosphere bacterial communities? FEMS

Microbiol Ecol 72:313–327.

23.   Bravo A, York T, Pumplin N, Mueller LA, Harrison MJ. 2016. Genes conserved for

arbuscular mycorrhizal symbiosis identified through phylogenomics. Nat Plants 2:15208.

24.   Sanchez-Contreras M, Bauer WD, Gao M, Robinson JB, Allan Downie J. 2007. Quorum-

sensing regulation in rhizobia and its role in symbiotic interactions with legumes. Philos

Trans R Soc B Biol Sci 362:1149–1163.

25.   Udvardi M, Poole PS. 2013. Transport and Metabolism in Legume-Rhizobia Symbioses.

Annu Rev Plant Biol 64:781–805.

26.   Garbutt K, Bazzaz FA. 1983. Leaf demography, flower production and biomass of diploid

and tetraploid populations of phlox-drummondii hook on a soil-moisture gradient. New

Phytol. WileyNew Phytologist Trust.

27.   Stebbins GL. 1971. Chromosomal Evolution in Higher PlantsContemporary Biology.

Edward Arnold Ltd., London.

28.   Ramsey J, Ramsey TS. 2014. Ecological studies of polyploidy in the 100 years following

its discovery. Philos Trans R Soc B Biol Sci 369:20130352–20130352.

29.   Breuer C, Stacey NJ, West CE, Zhao Y, Chory J, Tsukaya H, Azumi Y, Maxwell A,

Roberts K, Sugimoto-Shirasu K. 2007. BIN4, a Novel Component of the Plant DNA

Topoisomerase VI Complex, Is Required for Endoreduplication in Arabidopsis. PLANT

CELL ONLINE 19:3655–3668.

30.   Vergara F, Kikuchi J, Breuer C. 2016. Artificial Autopolyploidization Modifies the

Tricarboxylic Acid Cycle and GABA Shunt in Arabidopsis thaliana Col-0. Sci Rep

6:26515.

31.   Chao D-Y, Dilkes B, Luo H, Douglas A, Yakubova E, Lahner B, Salt DE. 2013.

Polyploids Exhibit Higher Potassium Uptake and Salinity Tolerance in Arabidopsis.

Science (80-) 341:658–659.

32.   Hollister JD, Arnold BJ, Svedin E, Xue KS, Dilkes BP, Bomblies K. 2012. Genetic

Adaptation Associated with Genome-Doubling in Autotetraploid Arabidopsis arenosa.

PLoS Genet 8:e1003093.

33.   Kerwin R, Feusier J, Corwin J, Rubin M, Lin C, Muok A, Larson B, Li B, Joseph B,

Francisco M, Copeland D, Weinig C, Kliebenstein DJ. 2015. Natural genetic variation in

Arabidopsis thaliana defense metabolism genes modulates field fitness. Elife 2015:1–28.

34.   van de Voorde TFJ, van der Putten WH, Bezemer TM. 2012. Soil inoculation method

determines the strength of plant-soil interactions. Soil Biol Biochem 55:1–6.

35.   Mauricio R. 1998. Costs of Resistance to Natural Enemies in Field Populations of the

Annual Plant Arabidopsis thaliana. Am Nat 151:20–28.

36.   Fierer N, Jackson RB. 2006. The diversity and biogeography of soil bacterial

communities. Proc Natl Acad Sci 103:626–631.

37.   Lauber CL, Hamady M, Knight R, Fierer N. 2009. Pyrosequencing-based assessment of

soil pH as a predictor of soil bacterial community structure at the continental scale. Appl

Environ Microbiol 75:5111–5120.

38. Rubin MJ, Brock MT, Weinig C, Baker RL, Wilcox S, Anderson K, Seth J, Rubin MJ.

2018. Circadian rhythms are associated with shoot architecture in natural settings. New

Phytol 219.

39. Bulgarelli D, Rott M, Schlaeppi K, Ver Loren van Themaat E, Ahmadinejad N, Assenza

F, Rauf P, Huettel B, Reinhardt R, Schmelzer E, Peplies J, Gloeckner FO, Amann R,

Eickhorst T, Schulze-Lefert P. 2012. Revealing structure and assembly cues for

Arabidopsis root-inhabiting bacterial microbiota. Nature 488:91–5.

40. Huse SM, Young VB, Morrison HG, Antonopoulos DA, Kwon J, Dalal S, Arrieta R,

Hubert NA, Shen L, Vineis JH, Koval JC, Sogin ML, Chang EB, Raffals LE. 2014.

Comparison of brush and biopsy sampling methods of the ileal pouch for assessment of

mucosa-associated microbiota of human subjects. Microbiome 2:5.

41. Callahan BJ, McMurdie PJ, Rosen MJ, Han AW, Johnson AJA, Holmes SP. 2016.

DADA2: High-resolution sample inference from Illumina amplicon data. Nat Methods

13:581–583.

42. Quast C, Pruesse E, Yilmaz P, Gerken J, Schweer T, Yarza P, Peplies J, Glöckner FO.

2013. The SILVA ribosomal RNA gene database project: Improved data processing and

web-based tools. Nucleic Acids Res 41:D590–D596.

43. Shetty SA, Hugenholtz F, Lahti L, Smidt H, de Vos WM. 2017. Intestinal microbiome

landscaping: Insight in community assemblage and implications for microbial modulation

strategies. FEMS Microbiol Rev 41:182–199.

44. Harrison JG, Calder WJ, Shastry V, Buerkle CA. 2020. Dirichlet-multinomial modelling

outperforms alternatives for analysis of microbiome and other ecological count data. Mol

Ecol Resour 20:481–497.

45.   Holmes I, Harris K, Quince C. 2012. Dirichlet multinomial mixtures: Generative models

for microbial metagenomics. PLoS One 7.

46.   Fordyce JA, Gompert Z, Forister ML, Nice CC. 2011. A Hierarchical Bayesian Approach

to Ecological Count Data: A Flexible Tool for Ecologists. PLoS One 6:e26785.

47.   Love MI, Huber W, Anders S. 2014. Moderated estimation of fold change and dispersion

for RNA-seq data with DESeq2. Genome Biol 15:550.

48.   Carpenter B, Gelman A, Hoffman MD, Lee D, Goodrich B, Betancourt M, Brubaker M,

Guo J, Li P, Riddell A. 2017. Stan: A Probabilistic Programming Language. J Stat Softw

76:1–32.

49.   Stan Development Team. 2018. rstan: the R interface to Stan v. 2.18.2.

50.   Besag J, Green P, Higdon D, Mengersen K. 1989. Institute of Mathematical Statistics is

collaborating with JSTOR to digitize, preserve, and extend access to Statistical Science. ®

www.jstor.org. Statistics (Ber) 10:409–435.

51.   Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. 1990. Basic local alignment

search tool. J Mol Biol 215:403–410.

52.   Johnson M, Zaretskaya I, Raytselis Y, Merezhuk Y, McGinnis S, Madden TL. 2008.

NCBI BLAST: a better web interface. Nucleic Acids Res 36.

53.   Agarwala R, Barrett T, Beck J, Benson DA, Bollin C, Bolton E, Bourexis D, Brister JR,

Bryant SH, Canese K, Cavanaugh M, Charowhas C, Clark K, Dondoshansky I, Feolo M,

Fitzpatrick L, Funk K, Geer LY, Gorelenkov V, Graeff A, Hlavina W, Holmes B, Johnson

M, Kattman B, Khotomlianski V, Kimchi A, Kimelman M, Kimura M, Kitts P, Klimke

W, Kotliarov A, Krasnov S, Kuznetsov A, Landrum MJ, Landsman D, Lathrop S, Lee JM,

Leubsdorf C, Lu Z, Madden TL, Marchler-Bauer A, Malheiro A, Meric P, Karsch-

Mizrachi I, Mnev A, Murphy T, Orris R, Ostell J, O'Sullivan C, Palanigobu V, Panchenko

AR, Phan L, Pierov B, Pruitt KD, Rodarmer K, Sayers EW, Schneider V, Schoch CL,

Schuler GD, Sherry ST, Siyan K, Soboleva A, Soussov V, Starchenko G, Tatusova TA,

Thibaud-Nissen F, Todorov K, Trawick BW, Vakatov D, Ward M, Yaschenko E,

Zasypkin A, Zbicz K. 2018. Database resources of the National Center for Biotechnology

Information. Nucleic Acids Res 46:D8–D13.

54.   Alix K, Gérard PR, Schwarzacher T, Heslop-Harrison JSP. 2017. Polyploidy and

interspecific hybridization: Partners for adaptation, speciation and evolution in plants. Ann

Bot. Oxford University Press.

55.   Bai Y, Müller DB, Srinivas G, Garrido-Oter R, Potthoff E, Rott M, Dombrowski N,

Münch PC, Spaepen S, Remus-Emsermann M, Hüttel B, McHardy AC, Vorholt JA,

Schulze-Lefert P. 2015. Functional overlap of the Arabidopsis leaf and root microbiota.

Nature 528:364–369.

56.   Staley C, Ferrieri AP, Tfaily MM, Cui Y, Chu RK, Wang P, Shaw JB, Ansong CK,

Brewer H, Norbeck AD, Markillie M, do Amaral F, Tuleski T, Pellizzaro T, Agtuca B,

Ferrieri R, Tringe SG, Paša-Tolić L, Stacey G, Sadowsky MJ. 2017. Diurnal cycling of

rhizosphere bacterial communities is associated with shifts in carbon metabolism.

Microbiome 5:65.

57.   Edger PP, Heidel-Fischer HM, Bekaert M, Rota J, Glöckner G, Platts AE, Heckel DG,

Der JP, Wafula EK, Tang M, Hofberger JA, Smithson A, Hall JC, Blanchette M, Bureau

TE, Wright SI, dePamphilis CW, Eric Schranz M, Barker MS, Conant GC, Wahlberg N,

Vogel H, Pires JC, Wheat CW. 2015. The butterfly plant arms-race escalated by gene and

genome duplications. Proc Natl Acad Sci 112:8362–8366.

58.   Lebeis SL, Paredes SH, Lundberg DS, Breakfield N, Gehring J, McDonald M, Malfatti S,

del Rio TG, Jones CD, Tringe SG, Dangl JL, Glavina del Rio T, Jones CD, Tringe SG,

Dangl JL. 2015. Salicylic acid modulates colonization of the root microbiome by specific

bacterial taxa. Science (80-) 349:860–864.

59.   Badri D V., Zolla G, Bakker MG, Manter DK, Vivanco JM. 2013. Potential impact of soil

microbiomes on the leaf metabolome and on herbivore feeding behavior. New Phytol

198:264–273.

60.   Fitzpatrick CR, Copeland J, Wang PW, Guttman DS, Kotanen PM, Johnson MTJ. 2018.

Assembly and ecological function of the root microbiome across angiosperm plant

species. Proc Natl Acad Sci U S A 115:E1157–E1165.

61.   Ishii S, Ashida N, Ohno H, Segawa T, Yabe S, Otsuka S, Yokota A, Senoo K. 2017.

Noviherbaspirillum denitrificans sp. nov., a denitrifying bacterium isolated from rice

paddy soil and Noviherbaspirillum autotrophicum sp. nov., a denitrifying, facultatively

autotrophic bacterium isolated from rice paddy soil and proposal to reclassify H. Int J Syst

Evol Microbiol 67:1841–1848.

62.   Segraves KA. 2017. The effects of genome duplications in a community context. New

511   Phytol 215:57–69.

512   63.   Bomblies K, Madlung A. 2014. Polyploidy in the Arabidopsis genus. Chromosom Res.

513   Springer Netherlands.

514   64.   Van de Peer Y, Mizrachi E, Marchal K. 2017. The evolutionary significance of

515   polyploidy. Nat Rev Genet 18:411–424.

517   **Figures**

Figure 1: Genotype and ploidy differentially affect rhizosphere bacterial community abundances between the Col and Ler genotypes. a)
Families identified as more abundant in the diploid rhizosphere are shown above grey line, and those more abundant in the tetraploid
rhizosphere appear below grey line. b) Families identified as more abundant in the Landsberg rhizosphere are shown above grey line,
and those more abundant in the rhizosphere of tetraploids appear below grey line. Log10 fold changes were calculated from relative
abundance estimates obtained through hierarchical Bayesian modeling of read counts (see main text). Points represent individual ASVs
within families.

Figure 2: a,b) Residuals of observed growth differences in above and belowground biomass for plants grown in soils inoculated with microbiomes shaped by each genotype, with effects of block removed. All genotypes inoculated with Columbia tetraploid microbiome had significantly reduced above and below ground biomass. c) Bacterial families identified as more abundant in the Columbia tetraploid rhizosphere are shown above grey line. Those more abundant in the rhizosphere of all other inoculums appear below grey line. Log10 fold changes were calculated from relative abundance estimates obtained through hierarchical Bayesian modeling of read counts (see main text). Points represent individual ASVs within families

521   **Supplemental Tables and Figures**

523   **Supplemental Table 1**: Number of reads at each step of sequence data processing

Processing Step	Number of Reads
Initial	3 282 410
Filtering and Trimming, Sample Inference, Chimera Removal	1 985 490
Removal of Chloroplast, Mitochondrial, Eukaryotic DNA and Plant-less blanks	1 647 741

525   **Supplemental Table 2**: Effect of inoculum shaped by plant genotype on plant performance

Trait	P - Value
Aboveground biomass	< 0.001
Belowground biomass	< 0.001
Root : Shoot	**ns**
Days to germination	**ns**
Days to flowering	**ns**
Seed Mass	**ns**

528