Title: The intrinsic neonatal hippocampal network: rsfMRI findings

Abbreviated Title: The intrinsic neonatal hippocampal network

Authors:

Athena L. Howell (contribution: running registration and fMRI analyses, statistical analyses and writing the manuscript),

David E. Osher (preprocessing fMRI data and editing the manuscript),

Jin Li (preprocessing fMRI data and editing the manuscript),

and Zeynep M. Saygin (PI/advisor, guiding the research and analyses, checking results and editing the manuscript)

The Ohio State University, 43210

Please address correspondence to:

Athena L. Howell, The Ohio State University, 225 Psychology Building, 1835 Neil Avenue, Columbus, OH, 43210. E-mail: howell.551@osu.edu

# of pages: 19 (in body of manuscript)

# of Figures: 5 Figures

# of words in abstract: 201

24 # of words in introduction: 575

25 # of words in discussion: 1077

26 Acknowledgements; The authors declare no competing financial interests

# **Abstract**

Many adults cannot voluntarily recall memories before the ages of 3-5, a phenomenon referred to as "infantile amnesia". The development of the hippocampal network likely plays a significant part in the emergence of the ability to form long-lasting memories. In adults, the hippocampus has specialized and privileged connections with certain cortical networks, which presumably facilitate its involvement in memory encoding, consolidation, and retrieval. Is the hippocampus already specialized in these cortical connections at birth? And are the topographical principles of connectivity (e.g. long-axis specialization) present at birth? We analyzed resting-state hippocampal connectivity in neonates scanned within one week of birth (Developmental Human Connectome Project) and compared them to adults (Human Connectome Project). We explored the connections of the whole hippocampus and its long-axis specialization to seven canonical cortical networks. We found that the neonatal hippocampal networks show clear immaturity at birth: adults showed hippocampal connectivity that was unique for each cortical network, whereas neonates showed no differentiation in hippocampal connectivity across these networks. Further, neonates lacked long-axis specialization (i.e., along anterior-posterior axis) of the hippocampus in its differential connectivity patterns to the cortical networks. This immaturity in connectivity may contribute to immaturity in memory formation in the first years of life.

"New and Noteworthy":

While animal data, and anatomical and behavioral human data from young children

suggest that the hippocampus is immature at birth, to date, there are no direct assessments of

human hippocampal functional connectivity (FC) very early in life. Our study explores the FC of

the hippocampus to the cortex at birth, allowing insight into the development of human memory

systems.

## **Introduction**

Many adults cannot voluntarily recall memories before the ages of 3-5, a phenomenon

referred to as "infantile amnesia" (Alberini & Travaglia, 2017). One potential reason for this is

that the hippocampus (the primary brain structure responsible for episodic memory formation in

adults) and its connections with the rest of the brain may be immature at birth. Indeed, the

hippocampus does appear to be immature at birth; evidence in macaques suggests it continues to

mature after one year of age (roughly age 3-5 in humans) (Jabés et al., 2011) and human data

indicates that volumetric and structural changes in the hippocampus continue through childhood

(DeMaster et al., 2014; Gilmore et al., 2012; Seress 2007). Further, episodic memory performance

may be influenced by changes in the patterns of hippocampal connectivity from middle childhood

to adulthood, including along the long-axis of the hippocampus (Blankenship et al., 2017;

DeMaster et al., 2014; Ghetti et al., 2010; Gogtay et al., 2006; Poppenk & Moscovitch, 2011;

Riggins et al., 2016). At younger ages, hippocampal gray matter volume has been linked to early

language ability (Can et al., 2013) and one recent study showed potential hippocampal activation

for learned items in 2-year old toddlers (Prabhakar et al., 2018). However, the intrinsic

connectivity of hippocampus very early in life is less well understood. Therefore, an understanding

of the hippocampal network at birth and its development may lead to greater understanding of

memory development.

Recently, Wael and colleagues (2018) showed the hippocampus has a clear intrinsic pattern

of functional connectivity (FC) to a set of cortical networks in adults. Specifically, they showed

higher (i.e. most positive) connectivity from the hippocampus to the Default Mode and Limbic

networks and lowest (i.e. least positive) connectivity to the Frontoparietal and Ventral Attention

networks (from Yeo et al., 2011), Further, this connectivity pattern differed between the anterior

and posterior portions of the hippocampus, with anterior hippocampus showing larger differences

in connectivity to the networks than posterior hippocampus. This so-called long-axis specialization

of the hippocampus is consistent with previous research showing that the anterior and posterior

hippocampus display different patterns of structural and functional connectivity and may be

uniquely activated in response to cognitive, memory and spatial demands (for reviews see Poppenk

et al., 2013, Strange et al., 2014). The development of the hippocampal network and the long-axis

gradient likely plays a significant part in the emergence of the ability to form long-lasting

memories. For instance, the work of Riggins et al. (2016) examines the relationship of

anterior/posterior connectivity and episodic memory in 4- and 6-year old children and finds

developmental differences even between these two ages. Although this work in young children is

notable, the fact remains that we know very little about the hippocampus, its connections, and its

relationship to memory-formation during the earliest stages of life.

To this end, we compared the resting-state hippocampal connectivity patterns to a set of

cortical networks in neonates and adults. Resting state connectivity, determined by spontaneously

correlated activity of disparate brain regions, is used as a reliable marker of intrinsic functional

connectivity (FC) between brain regions (Biswal et al., 1995; Raichle, 2009; Smith, 2013; Sporns,

2013); further, FC at rest is predictive of task-based activity (Cole et al., 2014; Osher et al., 2019;

Smith et al., 2009; Tobyne et al., 2018).

More recently, developmental studies using FC have shown the FC of some networks is

mature at birth while others take months or longer to become adultlike (for reviews see Gao et al.,

2017 and Grayson & Fair, 2017). In particular, multiple studies indicate the connectivity of visual

and somatomotor networks is not only functional but highly adult-like at birth (Gao et al., 2015b;

Lin et al., 2008; Liu et al., 2008). Other areas, such as the default mode network, dorsal attention

network, frontoparietal network, and some perceptual regions show relatively immature functional

and structural characteristics at birth and experience large modifications postnatally (Gao et al.,

2015b; Natu et al., 2019), although the frontoparietal network may have important functional roles

even within the first year of life (e.g. Linke et al., 2018).

To assess hippocampal maturity at birth, we analyzed FC between seven intrinsic networks

and the hippocampus as a whole as well as along the hippocampal long-axis in both neonates and

adults. We also compared neonatal vs. adult hippocampal connectivity to the cortex at a finer,

voxelwise scale. Based on previous literature suggesting the immaturity of the hippocampus at

birth, we hypothesized that neonates would differ from adults in their hippocampal connectivity

to the cortex, particularly to the more immature networks (e.g. default mode and frontoparietal).

**Materials and Methods**

**Participants**

<u>Neonates:</u>

Neonatal data comes from the initial release of the Developing Human Connectome Project

(dHCP) (http://www.developingconnectome.org, Markopoulos et al., 2018). Neonates were

recruited and imaged in London at the Evelina Neonatal Imaging Centre after gathering informed

parental consent to image and release the data. The study was approved by the UK Health Research

Authority. 40 neonates were included in our analyses (15 female, 36-44 weeks old at scan).

<u>Adults:</u>

Adult data comes from the Human Connectome Project (HCP), WU-Minn HCP 1200

Subject Data Release (https://www.humanconnectome.org/study/hcp-young-adult, Van Essen et

al., 2013). Participants were scanned at Washington University in St. Louis (WashU). We included

40 participants in our analyses (15 female; 20-35 years old). These adult participants were motion-

matched to the neonates. Specifically, we matched each neonatal participant with an adult from

the HCP dataset with the same gender who showed the most similar motion parameter (i.e.,

framewise displacement, FD). While the resulting group of adults were motion-matched to the

neonates, we found that the groups were significantly different in the average gray-matter tSNR,

with neonates exhibiting higher tSNR values ($t(78)$=-6.8774, $p$=1.3469x10$^{-9}$). To ensure that any

results were not driven by tSNR differences between groups, we identified an additional group of

HCP adults whose tSNR was matched to the tSNR of the neonates $(t(78)$=-1.5237, $p$=.132) and

replicated our results (see Extended Data, Figure 1-1).

**MRI Acquisition**

<u>Neonates:</u>

All acquisition information comes from the dHCP data release documentation. Imaging

was carried out on a 3T Philips Achieva (running modified R3.2.2 software) using an imaging

system specifically designed for neonates with a 32 channel phased array head coil (Hughes, E.J.,

et al.). Neonates were scanned during natural sleep; resting-state FC patterns have been shown to

stay largely consistent while awake, asleep, or under anesthesia (Liu et al., 2015; Larson-Prior et

al., 2009).

*Resting-state fMRI*

High temporal resolution fMRI developed specifically for neonates was collected using

multiband (MB) 9x accelerated echo=planar imaging (TE/TR=38/392ms, voxel size = 2.15 x 2.15

x 2.15mm$^3$). The resting state scan lasted approximately 15 minutes and consisted of 2300 volumes

for each run. No in-plane acceleration or partial Fourier transform was used. Single-band reference

scans with bandwidth matched readout and additional spin-echo acquisitions were also acquired

with both AP/PA fold-over encoded directions.

*Anatomical MRI*

High-resolution T2-weighted and inversion recovery T1-weighted multi-slice fast spin-

echo images were acquired with in-plane resolution 0.8 x $0.8mm^2$ and 1.6mm slices overlapped

by 0.8mm (T2-weighted: TE/TR= 156/12000ms; T1 weighted: TE/TR/TI = 8.7/4795/1740ms)

<u>Adults:</u>

All acquisition information comes from the HCP data release documentation. Scanning for

the 1200 WU-Minn HCP subject was carried out on a customized 3T Connectome Scanner adapted

from a Siemens Skyra (Siemens AG, Erlanger, Germany), equipped with a 32-channel Siemens

receiver head coil and a "body" transmission coil specifically designed by Siemens to

accommodate the smaller space (due to special gradients) of the WU-Minn and MGH-UCLA

Connectome scanners.

*Resting-State fMRI*

Participants were scanned using the Gradient-echo EPI sequence (TE/TR = 33.1/720ms,

flip angle = $52^o$, 72 slices, voxel size = 2 x 2 x $2mm^3$). Scanning lasted approximately 15 minutes

consisting of 1200 volumes for each run. Each participant finished two resting-state fMRI sessions.

For each session, two phases were encoded: one right-to-left (RL) and the other left-to-right (LR).

For our analyses, we used the LR phase encoding from the first session. Participants were

instructed to relax and keep their eyes open and fixated on a bright, projected cross-hair against a

dark background.

*Anatomical MRI*

High-resolution T2-weighted and T1-weighted images were acquired with an isotropic

voxel resolution of 0.7mm$^3$ (T2-weighted 3D T2-SPACE scan: TE/TR=565/3200ms; T1-weighted

3D MPRAGE: TE/TR/TI = 2.14/2400/1000ms).

**MRI Preprocessing**

<u>Neonates:</u>

The dHCP data was preprocessed using the dHCP minimal preprocessing pipelines

(Makropoulos et al., 2018). Anatomical MRI preprocessing included bias correction, brain

extraction using BET from FSL (FMRIB Software Library) and segmentation of the T2w volume

using their DRAW-EM algorithm (Makropoulos et al., 2014). The resulted gray and white matter

segmentations were used as anatomical masks in further analyses; these masks were manually

checked for accuracy.

Minimal preprocessing for the resting-state fMRI included (Fitzgibbon et al., 2016)

distortion correction, motion correction, 2-stage registration of the MB-EPI functional image to

the T2 structural image, temporal high-pass filtering (150s high-pass cutoff), and ICA denoising

using FSL's FIX (Salimi-Khorshidi, et al., 2014).  In addition to this minimal preprocessing, we

smoothed the data (Gaussian filter, FWHM = 3mm) across the gray matter, and applied a band-

pass filter at 0.009-0.08 Hz. To further denoise the data, we used aCompCor (Behzadi et al., 2007)

to regress out physiological noise (heartbeat, respiration, etc.) from the white matter and

cerebrospinal fluid (CSF).

<u>Adults:</u>

HCP data was preprocessed using the HCP minimal preprocessing pipelines (Glasser et al.,

2013). For the anatomical data, a Pre-FreeSurfer pipeline was applied to correct gradient distortion,

produce an undistorted "native" structural volume space for each adult participant by ACPC

registration (hereafter referred to as "acpc space"), extract the brain, perform a bias field

correction, and register the T2-weighted image to the T1-weighted image. Additionally, each

participant's brain was aligned to a common MNI152 template brain (with 0.7mm isotropic

resolution). Then, the FreeSurfer pipeline (based on FreeSurfer 5.3.0-HCP) was performed with a

number of enhancements specifically designed to capitalize on HCP data (Glasser et al., 2013).

The goal of this pipeline was to segment the volume into predefined structures, to reconstruct the

white and pial cortical surfaces, and to perform FreeSurfer's standard folding-based surface

registration to their surface atlas (fsaverage).

For the resting-state fMRI data, minimal functional analysis pipelines included: removing

spatial distortions, motion correction, registering the fMRI data to structural and MNI152

templates, reducing the bias field, normalizing the 4D image to a global mean, and masking the

data with the final brain mask. After completing these steps, the data were further denoised using

the ICA-FIX method (Salimi-Khorshidi, et al., 2014). To mirror the adult and neonatal

preprocessing pipelines, we unwarped the data from MNI152 to acpc space, allowing both groups

to be analyzed in "native" space. We then applied spatial smoothing (Gaussian filter, FWHM =

3mm) within the gray matter, band-pass filtered at 0.009-0.08 Hz and implemented aCompCor to

regress out physiological noise, just as we did with the neonates.

All subsequent analyses in neonates and adults were performed in each subject's native

space, except for the whole-brain voxelwise analysis.

**Connectivity analyses**

We used the 7-network cortical parcellation identified by Yeo et al. (2011). For the

whole-hippocampus and long-axis analyses, the hippocampal label was binarized from

FreeSurfer's (surfer.nmr.mgh.harvard.edu) aparc+aseg parcellation and visually inspected for

accuracy in each subject. For the first long-axis gradient analysis this label was further sectioned

into anterior and posterior portions via manual segmentation using FreeSurfer, with the uncal

apex as the dividing marker (Poppenk & Moscovitch., 2011). All labels (cortical networks,

hippocampal labels) were originally in CVS average-35 MNI152 space and then registered to

each individual subject's anatomical data using ANTs (Advanced Normalization Tool)

3dWarpMultiTransform (ANTs version 2.1.0; http://stnava.github.io/ANTs; Avants et al., 2011).

ANTs is routinely used for developmental dataset registrations (Alexander et al., 2019; Dean et

al., 2018). The resulting registrations were checked for accuracy. Similarly, for the long-axis

gradient analysis, the hippocampal label in CVS was split into nine equally spaced "slices" along

the anterior-posterior axis. Using the same ANTs registration technique for all ROIs provided an

extra measure of consistency between groups and between analyses; however, as an added

quality check we ran our whole-hippocampus to network analysis using the binarized

hippocampal label provided by the dHCP and HCP for each individual. These second results are

nearly identical (see Extended Data, Figure 2-1) to the first (Figure 2) thus increasing confidence

that our results are not due to registration error.

After registration to the anatomical data, we registered the labels onto the functional data

in neonates using an inverse warp of the func2anat matrix provided by the dHCP. In adults, the

labels in acpc space after ANTs registration were then resampled to 2mm cubic voxels to align

with the functional data. We manually checked individuals from each sample to ensure the

accuracy and fit of the labels to the individual functional data.  We extracted the BOLD activation

in each label over the time course, averaged within each label, and correlated the hippocampal

activity—first whole hippocampus, then along the long-axis (for both anterior-posterior and

gradient slices)—with activity in each of the 7 networks to create a Fisher's Z-scored correlation

matrix using Matlab 2018b (The MathWorks, Inc., Natick, Massachusetts, United States).

We also explored differences in the hippocampal connectivity to the whole cortex at a voxelwise

scale between adults and neonates to determine whether specific regions within the networks were

driving adult-neonate differences. Hippocampal connectivity to the cortex was calculated by

correlating the average hippocampal signal and the signal of each voxel within the cortical gray

matter mask during the time course for each individual in functional space. To compare the

connectivity between adults and neonates, images from both groups were registered to the template

space (i.e., CVS average-35 MNI152) before running a between-group analysis. Although this is

the only template-space analysis we performed, template-space analyses have been routinely

performed to compare infants to adults using similar registration methods (e.g. Gao et al., 2009;

Gao et al., 2015a).

**Experimental Design and Statistical Analyses**

Where t-tests were performed between regions we corrected for multiple comparisons

using the Holm-Bonferroni correction (Holm, 1979); all connectivity values were Fisher's Z

transformed (Fisher, 1915) to normalize the data.

Before doing any of the planned analyses, we first performed data quality checks.  To make

sure there was no significant motion difference between groups, we calculated the framewise

displacement (FD) (Power et al., 2012) based on the six motion parameters estimated from a rigid-

body transformation provided by dHCP and HCP. We manually checked the registration of the

gray and white matter masks as well as the network and hippocampal labels in the adults and

neonates to the registration was accurate. Because we are performing comparisons of correlations

between groups, we next wanted to ensure that the correlation distributions were similar and were

normally distributed in both neonates and adults; we did this by assessing the correlation of each

voxel to every other voxel in the brain and plotting the distribution of those correlations. We also

performed between-subject reliability of correlation matrices within and across the adult and

neonate groups. We calculated the connectivity of each region (i.e. each of the seven networks and

the hippocampus) to every other region for each subject. This connectivity matrix was then

correlated with every other subject's value either between- or within-groups to assess inter-subject

reliability; in other words, we correlated the connectivity of every adult to every other adult

(within-group) and neonate to neonate, as well as comparing every adult to every neonate

(between-group).

Our first analysis examined the relationship of the whole hippocampus to the seven cortical

networks. After running a one-way ANOVA with network as the independent variable and

connectivity as the dependent variable for both groups, we computed pairwise comparisons

between each unique combination of connectivity values to the networks (e.g. Hipp-Lim vs Hipp-

VA) to determine networks with significantly different FC to the hippocampus (Snedecor and

Cochran, 1989). Rose plots comparing the connectivity pattern of adults and neonates were created

by subtracting the mean connectivity across all networks from each individual network (for adults

and neonates separately) and plotting the resulting magnitude to show the relative connectivity

patterns of the hippocampus to the networks for each group and to compare these patterns between

groups.

For our hippocampal-cortical voxelwise analysis, we used FSL's randomise function to

compare between groups and perform permutation testing (to correct for multiple comparisons) in

order to determine areas of greater connectivity in adults vs neonates and visa versa. After mapping

the individual correlation matrices from subject space into a common template space, we used

randomise with default 5000 permutations and clustered the results using FSL's threshold-free

cluster enhancement (TFCE), which corrects for family-wise error (FWE). This produced a list of

potential clusters with each cluster's associated p-value; the p-values were then thresholded at a

$p < 0.0005$, and only those clusters that remained significant after that point are reported in this

paper.

For the first long-axis hippocampus analysis, we first computed a two-way ANOVA in

each group (separately) using location (i.e. anterior or posterior hippocampus) and network as

independent variables and FC as the dependent variable. Pairwise comparisons were then made

between the anterior and posterior FC values to each network for each group (e.g. adult antHipp-

Lim vs adult postHipp-Lim). For the second long-axis analysis, we conducted a two-way ANOVA

at each slice using group and network as independent variables and connectivity as the dependent

variable. We also computed a one-way ANOVA at each slice for each group with network as the

independent variable. As in the whole-hippocampal analysis, rose plots were created by

subtracting out the mean connectivity to all networks (e.g. mean connectivity of adult anterior

hippocampus to all networks) from each network and group in the anterior and posterior labels

individually to demonstrate comparative connectivity differences between the anterior and

posterior regions in each group.

**Results**

(FIGURE 1 HERE)

**Preliminary data-checks**

Comparison of the framewise displacement in adults and neonates showed no significant

difference of FD between adults and neonates (t(78)=-0.48, p=0.63).  Visual inspection of the

gray and white matter masks (which are critical for resting-state preprocessing) in Figure 1a

shows they are accurately delineating gray/white matter in both neonates and adults; the cortical

networks and hippocampal labels also appear to be correctly localized, suggesting that the

regions are accurately identified in both neonates and adults (Figure 1a). Figure 1b demonstrates

that both neonates and adults have normally-distributed correlation values that are centered

around 0. Between-subject reliability of correlation matrices within and across the adult and

neonate groups showed the connectivity matrices (i.e. region-to-region connectivity of each of

the seven networks and the hippocampus to each other) of each adult subject to each other adult

subject were highly correlated, as were the matrices of each neonate subject to each other

neonate subject, and a pairwise comparison of subject variability within groups (e.g. adult-adult

correlations compared to neonate-neonate correlations) was not significant (t(78)=0.76, p=0.45).

But subject-to-subject correlations across the two groups were significantly lower than the

within-group correlations (adult-adult vs adult-neo t(78) = 14.09, p=$3.87 \times 10^{-23}$, neonate-neonate

vs adult-neonate (t(78)=11.95, p=$2.63 \times 10^{-19}$)  suggesting that while the connectivity data are

reliable, neonates have different connectivity patterns than adults.

**Whole Hippocampus**

We first explored the connectivity of the whole hippocampus to the cortical networks. In

adults, there was a main effect of network suggesting that some networks are more strongly

connected with the hippocampus than others (Figure 2; one-way ANOVA, $F(6,273)=47.11$,

$p=1.84 \times 10^{-39}$). Subsequent pairwise comparisons showed a clear hierarchy of connectivity, such

that hippocampal connectivity was highest to the Limbic (Lim) network (vs hippocampal

connectivity to: Ventral Attention or VA ($t(78)=12.95$, $p_{HB}=8.42 \times 10^{-20}$); FrontoParietal or FP

($t(78)=11.76$, $p_{HB}=1.20 \times 10^{-17}$), Dorsal Attention or DA ($t(78)=10.09$, $p_{HB}=1.50 \times 10^{-14}$);Visual or

Vis ($t(78)=7.20$, $p_{HB}=4.52 \times 10^{-9}$); and SomatoMotor or SM ($t(78)=5.97$, $p_{HB}=7.91 \times 10^{-7}$)).

Hippocampal connectivity to the Default Mode Network (DM) was higher than hippocampal

connectivity to: VA ($t(78)=10.32$, $p_{HB}=5.80 \times 10^{-15}$); FP ($t(78)=9.07$, $p_{HB}=1.33 \times 10^{-12}$); DA

($t(78)=7.24$, $p_{HB}=4.04 \times 10^{-9}$); Vis ($t(78)=4.63$, $p_{HB}=1.45 \times 10^{-4}$); and SM ($t(78)=3.16$, $p_{HB}=0.014$)).

Hippocampal-SM connectivity was 3rd highest, and higher than hippocampal connectivity to: VA

($t(78)=7.83$, $p_{HB}=3.19 \times 10^{-10}$); FP ($t(78)=6.46$, $p_{HB}=1.09 \times 10^{-7}$); and DA ($t(78)=4.35$,

$p_{HB}=3.61 \times 10^{-4}$)). Hippocampal-Vis connectivity was the next highest (vs VA ($t(78)=5.49$,

$p_{HB}=5.31 \times 10^{-6}$); FP ($t(78)=4.16$, $p_{HB}=6.38 \times 10^{-4}$), and connectivity with DA was higher than with

VA ($t(78)=3.89$, $p_{HB}=1.47 \times 10^{-3}$). In summary, hippocampal connectivity was highest to Lim,

followed by DM, then SM, Vis, and DA; hippocampal connectivity was lowest (i.e. negatively

correlated) with the FP and VA networks. In previous literature, the hippocampus is occasionally

included as a part of the DM network; our finding of high Hippocampal-DM correlation and

anti-correlation between the hippocampus and attention (i.e. FP and VA) networks falls in line

with earlier work on the connectivity of the DM network (e.g. Buckner, Andrews-Hanna &

Schacter, 2008) and is a good sign of the reliability of our results.

In contrast to the adult pattern, although neonates did show a main effect of network

($F(6,273)=5.12$, $p=2.27 \times 10^{-5}$), pairwise comparisons indicated that only the Lim and SM

networks significantly differ from the rest, with significantly greater connectivity from the

hippocampus to Lim vs DA ((t(78)=5.31, $p_{HB}$=2.15x10$^{-5}$) and Lim vs FP (t(78)=4.22,

$p_{HB}$=1.33x10$^{-3}$) and significantly greater connectivity to SM vs DA(t(78)=3.35, $p_{HB}$=0.023).

Pairwise comparisons between adults and neonates showed significant differences

between the groups, with significantly less connectivity in adults to Vis (t(78)=-2.64,

$p_{HB}$=0.040), DA(t(78) =-2.77, $p_{HB}$=0.035), FP(t(78)=-5.33, $p_{HB}$=5.49x10$^{-6}$) and VA (t(78)=-8.62,

$p_{HB}$=5.86x10$^{-13}$) networks compared to neonates.

(FIGURE 2 HERE)**Hippocampus to Cortex voxelwise analysis**

We next explored the connectivity of the hippocampus to the entire cortex at a voxelwise

scale; because our previous analysis only focused on 7 canonical networks, we may have missed

differences between neonates and adults at a finer grain than that seen on a network level.

Thresholding the unpaired t-test results of the whole-brain clusters at p<0.0005 produced 26

significant FWE-corrected (Smith & Nichols, 2009) clusters in the neonates > adults comparison

(i.e. 26 clusters where neonatal hippocampal FC significantly exceeds adult hippocampal FC)

and 14 significant clusters in the adults > neonates comparison (Figure 3). Specifically,  neonates

show greater hippocampal FC to frontal and parietal areas, bilateral lingual and pericalcarine

cortex and cuneus when compared to adults; frontoparietal differences were particularly

prevalent within the right hemisphere. Adults, on the other hand, displayed greater hippocampal

FC than the neonates primarily to bilateral isthmus cingulate and precuneus. Cluster sizes and

indices for clusters greater than 200 voxels along with peak voxel location and associated brain

regions are reported in Figures 3-1 and 3-2  and largely follow the results from the 7-network

analysis—the neonatal hippocampus shows greater FC to frontoparietal and attention-relevant

areas, whereas the adult hippocampus shows greater FC with regions associated with the default

mode and limbic networks.

(FIGURE 3 HERE)

**Anterior-Posterior Hippocampus**

We next explored the anterior vs. posterior hippocampal connectivity patterns in neonates

and adults; previous literature in both humans and other animals suggest functional

differentiation of the anterior and posterior hippocampal segments, and thus we may expect these

segments to have differences in FC to the 7 cortical networks. In adults, a two-way ANOVA

indicated a main effect of network ($F(6,546)=60.04., p=5.04 \times 10^{-57}$) and an interaction between

network and anterior/poster hippocampus ($F(6,546)=14.31, p=3.54 \times 10^{-15}$) (Figure 4).

In neonates, the two-way ANOVA showed only a significant main effect for network

($f(6,546)=7.67, p=6.30 \times 10^{-8}$) (Figure 4).  Pairwise comparisons between the anterior and

posterior portions of the hippocampus in adults show greater anterior vs posterior connectivity to

the Lim ($t(78)=3.53, p_{HB}=0.0035$), DMN ($t(78)=2.38, p_{HB}=0.03$) and SM ($t(78)=3.19,$

$p_{HB}=0.0082$) networks, and decreased anterior vs posterior connectivity to the DA ($t(78)=-3.07,$

$p_{HB}=0.0087$), Frontoparietal ($t(78)=-5.79, p_{HB}=9.99 \times 10^{-7}$) and VA ($t(78)=-3.92, p_{HB}=0.0011$)

networks. These results suggest the anterior hippocampus was primarily driving the negative

correlations with VA & FP seen at the level of the whole hippocampus in adults. Neonates,

however, show no significant differences between the anterior and posterior portions of the

hippocampus to any of the networks, suggesting no differentiation/specialization of the

hippocampal segments in their connections to the rest of the brain.

(FIGURE 4 HERE)

**Long-Axis Gradient**

Finally, we investigated the long-axis gradient, which has been demonstrated to map onto

a differential functional gradient of the hippocampus. We broke up the hippocampus in each

subject into 9 different segments along the anterior-posterior axis and compared the 7-network

connectivity to these segments in neonates and adults. Adults showed clear differentiation of

network connectivity along the long-axis while neonates showed no clear differentiation (Figure

5). The Lim and DM in adults appeared to have an initial rise and fall of FC along the anterior-

posterior gradient of the hippocampus which differentiated them from the Vis, SM, and DA, and

the FP and VA showed a similar rise and fall of negative FC along the gradient. One-way

ANOVAs for adults and neonates at each slice indicated a main effect of network in adults in all

but the most posterior slice (Slice 1 ($F_{(6,273)}$=23.61, p=$1.89 \times 10^{-22}$), Slice 2 ($F_{(6,273)}$=40.45,

p=$4.19 \times 10^{-35}$), Slice 3 ($F_{(6,273)}$=50.09, p=$2.61 \times 10^{-41}$), Slice 4 ($F_{(6,273)}$=49.56, p=$5.48 \times 10^{-41}$),

Slice 5 ($F_{(6,273)}$=49.07, p=$1.11 \times 10^{-40}$), Slice 6 ($F_{(6,273)}$=25.10, p=$1.12 \times 10^{-23}$), Slice 7

($F_{(6,273)}$=13.40, p=$2.57 \times 10^{-13}$), Slice 8 ($F_{(6,273)}$=5.51, p=$2.12 \times 10^{-5}$)). In the neonates, there

was no main effect of network in any of the slices (at p<0.001). To compare between the two

groups, we performed two-way ANOVAs (with network and group as independent variables and

FC value as the dependent variable) for each of the 9 slices. There was a significant interaction

between network and group for the anterior 7 slices (Slice 1 ($F_{(6,546)}$=7.30, p=$1.64 \times 10^{-7}$), Slice

2 ($F_{(6,546)}$=16.25, p=$2.95 \times 10^{-17}$), Slice 3 ($F_{(6,546)}$=18.98, p=$3.99 \times 10^{-20}$), Slice 4

($F_{(6,546)}$=17.22, p=$2.83 \times 10^{-18}$), Slice 5 ($F_{(6,546)}$=17.93, p=$5.05 \times 10^{-19}$), Slice 6 ($F_{(6,546)}$=5.79,

p=$7.26 \times 10^{-6}$), Slice 7 ($F_{(6,546)}$=4.87, p=$7.27 \times 10^{-5}$) and Slice 8($F_{(6,546)}$=3.89, p=$8.26 \times 10^{-4}$), but

no group differences for the most posterior slice. These results show that the biggest

differentiation of hippocampal connectivity to the 7 networks occurs in the anterior 2/3s of the

hippocampus in adults and that neonates do not show this differentiation.

(FIGURE 5 HERE)

**Discussion**

Our results show that the intrinsic connectivity of the hippocampal network is not fully

mature at birth. Previous functional and volumetric evidence in both non-human primates and

humans suggests that the hippocampus continues to develop beyond one year of age, even into

middle childhood (Blankenship et al., 2017; Jabes et al., 2011; Keresztes et al., 2018; Lavenex

and Banta Lavenex, 2013; Riggins et al., 2016). Although there is some evidence to suggest that

the hippocampus is playing a key role in memory formation even early on in rodents (Alberini &

Travaglia, 2017; Travaglia et al., 2018), it has been suggested that the long-lasting memories of

very young children may be created in a fundamentally different way from adult long-term

memories and may rely on cortical mechanisms rather than the traditional hippocampal method

(Ellis & Turke-Browne, 2018; Gómez & Edgkin, 2016). Interestingly, multiple studies

comparing preterm to term infants show no differences in gray matter volume in the

hippocampus with decreased gestational age, implying the better part of hippocampal growth is

accomplished prior to birth (Alexander et al., 2019; Ge et al., 2015; Thompson et al., 2008); our

work suggests that although the physical bulk of the hippocampus may exist at birth, its

connections do not. Specifically, the hippocampus does not have preferential connectivity to any

particular network at birth and lacks any long-axis gradient of connectivity, suggesting that the

hippocampus, the cortical networks it interacts with, or some combination of both, are immature

at birth and may therefore be unable to form long-term memories using adult-like mechanisms.

Indeed, the cortex itself is still maturing early on (e.g. Gao et al., 2015b; Ofen et al., 2007;

Salzwedel et al., 2019) and it is likely this cortical immaturity, in addition to hippocampal

immaturity, is contributing to the differences in memory formation between adults and neonates.

Adults showed a clear hierarchy of FC to the seven networks (consistent with Wael et al.

(2018)), whereas neonates lacked this hierarchy. Further, the comparison between adults and

neonates shows significant differences between groups to all networks except the SM network,

and only marginally significant differences between groups in the Vis and DA networks.  The

similarity between adults and neonates in connectivity to the SM and Vis networks may be due

to the relative maturity of these areas at birth (Arcaro & Livingstone, 2017; Deen et al., 2017;

Hurk et al., 2017; Gao et al. 2017, Dall'Orso et al., 2018).

To more specifically determine which regions in the networks were responsible for the

differences seen between adults and neonates, we conducted a voxel-wise cortical analysis. Our

results indicate that neonates have higher connectivity to much of the cortex as compared to

adults with the exception of areas of bilateral medial orbitofrontal, isthmus cingulate and

precuneus. This is consistent with Riggins et al.'s (2016) conclusion that 4-year old children rely

more on regions "outside" the canonical hippocampal network to complete episodic memory

tasks, and other research suggesting the infant cortex is more broadly tuned than in adults (Ellis

&Turk-Browne, 2018).  The few regions where adults display higher FC than neonates reside

mainly within DM network and highlight the immaturity of this network: adults show

significantly greater DM-Hippocampal connectivity than neonates, consistent with Gao et al.'s

(2015) finding that this network is one of the last to develop in the first year of life.

Our anterior-posterior analysis and long-axis gradient analyses again suggest that the FC

differentiation of the hippocampus is lacking at birth. Consistent with previous literature, adults

display changes along the long-axis such that the anterior hippocampus shows greater

connectivity to the Lim network than the posterior hippocampus but greater posterior vs anterior

FC to the attention (i.e. FP and VA) networks; in fact, the anterior hippocampus is especially

anti-correlated with these networks, as is consistent with previous literature (Buckner et al.,

2009; Wael et al., 2018).  The greatest differentiation in FC to the networks in adults occurred

within the anterior two-thirds of the hippocampus. In contrast, neonates showed no specificity to

any of the networks along the long-axis or the anterior-posterior analysis. Blankenship et al.

(2017), Langnes et al. (2018), and Riggins et al. (2016) show evidence of specialization along

the longitudinal axis in 4- and 6-year old children but no such evidence is seen in our results,

suggesting that maturational changes within the hippocampus may occur before age 4 to produce

the preferential connectivity seen in children and adults. Future studies of infants and toddlers

can better elucidate when after birth this change in specialization of the long-axis occurs.

Several limitations warrant discussion. A major problem in imaging children is motion

artifact. We used the motion-corrected data that were released by the dHCP, took steps in

preprocessing to ensure that physiological artifacts were removed from the data in both neonates

and adults (Power et al., 2014; Yan et al., 2013), and further motion-matched the neonatal and

adult groups. Given that motion-related artifacts are a major confound in FC analyses (Power et

al., 2012; Satterthwaite et al., 2013), our approach should minimize the risk of spurious

correlations. Other steps we took to minimize potential confounds included visual inspection of

spatial registration results (and using established registration procedures that have been

previously performed on infants (Alexander et al., 2019; Dean et al., 2018; Gao et al., 2009; Gao

et al., 2015a)), performing the analyses in the native-space of each individual, and checking the

reliability of the correlation values across participants in each group to ensure they were not

particularly noisy in the neonatal group.  A result of particular note is that neonates showed

primarily positive FC from the hippocampus to the networks, while adults showed slightly

negative FC for some networks. Blankenship et al., (2017) similarly fail to show any negative

hippocampal FC in their sample of 4- and 6-year old children (but this may be due to their

preprocessing steps, see Murphy & Fox, 2017 for discussion). Here, we used the same

preprocessing steps in both neonates and adults and used aCompCor and other preprocessing

steps that should not necessarily remove negative correlations if they were there. Indeed, we

found a normal distribution of correlation values in both neonates and adults (Figure 1)

suggesting that negative correlations do exist in neonates, but not between the hippocampus and

the cortex.  Further, regardless of the negative vs. positive correlation differences we observe a

difference in the pattern of FC in adults (demonstrated in the rose plots) primarily in the anterior

portion of the hippocampus; this is missing in neonates.

Differences in arousal states between the groups present another challenge. Mitra et al.,

(2017) showed differences in resting-state connectivity between sleeping infants and waking

adults. However, observation of Mitra et al.'s data suggests although the magnitude of

connectivity may differ between arousal states, the overall pattern of connectivity remains

similar (i.e. the same clusters of connectivity are observed in sleep and in rest and their relative

comparison to other clusters remains similar across sleep states and age groups). Further,

although notable differences are seen between the 24-mo sleeping infants and waking adults in

Mitra et al., this difference is far less pronounced in the younger 6-mo infants. Based on previous

EEG studies (e.g. Roffwarg, 1966), it is possible that younger infants experience less slow-wave

sleep and more REM sleep and thus, younger infants (vs. older infants) during sleep would be

expected to look more like awake adults due to the high similarity of REM and wakefulness

activity patterns in EEG, particularly in infants. Because we would expect more awake-like REM

sleep and less slow-wave sleep in young infants, we believe that the neonates in the current study

are unlikely to show major wake/sleep confounds in their connectivity patterns. Finally, analysis

of the same dataset but specifically of visual network connectivity showed striking *similarities* in

connectivity patterns (https://www.biorxiv.org/content/10.1101/712455v1) and therefore further

suggest that any differences in data acquisition and/or sleep states between adults and neonates

are unlikely to systematically lead to the differences in network connectivity that we find here.

Finally, we found the motion- and gender-matched HCP adults used this manuscript

tended to have lower tSNR than their respective dHCP counterparts. To ascertain this

discrepancy was not the cause of observed group differences, we identified a separate group of

40 HCP adults whose tSNR matched that of the 40 neonates used here and performed our whole

hippocampus to network analysis on this group. The resulting pattern matches the pattern

observed from the previous analyses (i.e. using the original 40 HCP adults), reiterating that

identified differences in connectivity patterns between adults and neonates are likely not spurious

byproducts of discrepant data quality (Extended Data, Figure 1-1).

In conclusion, our results suggest that the resting-state FC patterns of the human

hippocampus are immature at birth. This immaturity may play a key role in infantile amnesia and

the vast differences between adults and neonates shown here suggests a fundamentally different

memory and learning system from that of adults may be present at this point in development.

**<u>References</u>**

Alberini, C. M., & Travaglia, A. (2017). Infantile Amnesia: A Critical Period of Learning to

Learn and Remember. *Journal of Neuroscience*, *37*(24), 5783–5795.

Alexander, B., Kelly, C. E., Adamson, C., Beare, R., Zannino, D., Chen, J., … Thompson, D. K.

(2019). Changes in neonatal regional brain volume associated with preterm birth and

perinatal factors. *NeuroImage*, *185*, 654–663.

Arcaro, M. J., & Livingstone, M. S. (2017). A hierarchical, retinotopic proto-organization of the

primate visual system at birth. *ELife*, *6*.

Avants, B. B., Tustison, N. J., Song, G., Cook, P. A., Klein, A., & Gee, J. C. (2011). A

Reproducible Evaluation of ANTs Similarity Metric Performance in Brain Image

Registration. *NeuroImage*, *54*(3), 2033–2044.

Behzadi, Y., Restom, K., Liau, J. & Liu, T.T.J.N. A component based noise correction method

(CompCor) for BOLD and perfusion based fMRI. *Neuroimage, 37*, 90-101 (2007)

Biswal, B., Yetkin, F. Z., Haughton, V. M., & Hyde, J. S. (1995). Functional connectivity in the

motor cortex of resting human brain using echo-planar mri. *Magnetic Resonance in*

*Medicine*, *34*(4), 537–541.

Blankenship, S. L., Redcay, E., Dougherty, L. R., & Riggins, T. (2017). Development of

hippocampal functional connectivity during childhood. *Human Brain Mapping*, *38*(1),

182–201.

Buckner, R. L., Andrews‐Hanna, J. R., & Schacter, D. L. (2008). The Brain's Default

Network. *Annals of the New York Academy of Sciences*, 1124(1), 1–38.

Buckner, R. L., Sepulcre, J., Talukdar, T., Krienen, F., Liu, H., Hedden, T., … Johnson, K. A.

(2009). Cortical Hubs Revealed by Intrinsic Functional Connectivity: Mapping, Assessment

of Stability, and Relation to Alzheimer's Disease. *The Journal of Neuroscience : The*

*Official Journal of the Society for Neuroscience*, *29*(6), 1860–1873.

Can, D. D., Richards, T., & Kuhl, P. (2013). Early gray-matter and white-matter concentration in

infancy predict later language skills: A whole brain voxel-based morphometry study.

*Brain and Language*, 124(1), 34–44. https://doi.org/10.1016/j.bandl.2012.10.007

Cole, M. W., Bassett, D. S., Power, J. D., Braver, T. S., & Petersen, S. E. (2014). Intrinsic and

task-evoked network architectures of the human brain. *Neuron*, *83*(1), 238–251.

Dall'Orso, S., Steinweg, J., Allievi, A. G., Edwards, A. D., Burdet, E., & Arichi, T. (2018).

Somatotopic Mapping of the Developing Sensorimotor Cortex in the Preterm Human Brain.

*Cerebral Cortex*, *28*(7), 2507–2515.

Dean, D. C., Planalp, E. M., Wooten, W., Schmidt, C. K., Kecskemeti, S. R., Frye, C., …

Davidson, R. J. (2018). Investigation of brain structure in the 1-month infant. *Brain*

*Structure and Function*, *223*(4), 1953–1970.

Deen, B., Richardson, H., Dilks, D. D., Takahashi, A., Keil, B., Wald, L. L., … Saxe, R. (2017).

Organization of high-level visual cortex in human infants. *Nature Communications*, *8*,

13995.

DeMaster, D., Pathman, T., Lee, J. K., & Ghetti, S. (2014). Structural development of the

hippocampus and episodic memory: Developmental differences along the

anterior/posterior axis. *Cerebral Cortex (New York, N.Y.: 1991)*, *24*(11), 3036–3045.

Ellis, C. T., & Turk-Browne, N. B. (2018). Infant fMRI: A Model System for Cognitive

Neuroscience. *Trends in Cognitive Sciences*, *22*(5), 375–387.

Fisher, R. A. (1915). Frequency Distribution of the Values of the Correlation Coefficient in

Samples from an Indefinitely Large Population. *Biometrika*, *10*(4), 507.

Fitzgibbon, S.P., et al. The developing Human Connectome Project (dHCP): minimal 406

functional pre-processing pipeline for neonates. *in Fifth Biennial Conference on Resting*

*State and Brain Connectivity* (2016)

Gao, W., Alcauter, S., Elton, A., Hernandez-Castillo, C. R., Smith, J. K., Ramirez, J., & Lin, W.

(2015). Functional Network Development During the First Year: Relative Sequence and

Socioeconomic Correlations. *Cerebral Cortex (New York, NY)*, *25*(9), 2919–2928.

Gao, W., Alcauter, S., Smith, J. K., Gilmore, J., & Lin, W. (2015). Development of Human Brain

Cortical Network Architecture during Infancy. *Brain Structure & Function*, *220*(2),

1173–1186.

Gao, W., Lin, W., Grewen, K., & Gilmore, J. H. (2017). Functional Connectivity of the Infant

Human Brain: Plastic and Modifiable. *The Neuroscientist*, *23*(2), 169–184.

Gao, W., Zhu, H., Giovanello, K. S., Smith, J. K., Shen, D., Gilmore, J. H., & Lin, W. (2009).

Evidence on the emergence of the brain's default network from 2-week-old to 2-year-old

healthy pediatric subjects. *Proceedings of the National Academy of Sciences*, *106*(16),

6790–6795.

Ge, X., Shi, Y., Li, J., Zhang, Z., Lin, X., Zhan, J., Ge, H., Xu, J., Yu, Q., Leng, Y., Teng, G.,

Feng, L., Meng, H., Tang, Y., Zang, F., Toga, A. W., & Liu, S. (2015). Development of

the human fetal hippocampal formation during early second trimester. *NeuroImage*, *119*,

33–43.

Ghetti, S., DeMaster, D. M., Yonelinas, A. P., & Bunge, S. A. (2010). Developmental

Differences in Medial Temporal Lobe Function during Memory Encoding. *Journal of

Neuroscience*, *30*(28), 9548–9556.

Gilmore, J. H., Shi, F., Woolson, S. L., Knickmeyer, R. C., Short, S. J., Lin, W., … Shen, D.

(2012). Longitudinal Development of Cortical and Subcortical Gray Matter from Birth to

2 Years. *Cerebral Cortex*, *22*(11), 2478–2485.

Glasser, M.F., et al. The minimal preprocessing pipelines for the Human Connectome Project.

*Neuroimage*, *80*, 105-124 (2013)

Gómez, R. L., & Edgin, J. O. (2016). The extended trajectory of hippocampal development:

Implications for early memory development and disorder. *Developmental Cognitive*

*Neuroscience*, *18*, 57–69.

Gogtay, N., Nugent, T. F., Herman, D. H., Ordonez, A., Greenstein, D., Hayashi, K. M., …

Thompson, P. M. (2006). Dynamic mapping of normal human hippocampal

development. *Hippocampus*, *16*(8), 664–672.

Grayson, D. S., & Fair, D. A. (2017). Development of large-scale functional networks from birth

to adulthood: A guide to the neuroimaging literature. *NeuroImage*, *160*, 15–31.

Holm, S. (1979). A simple sequentially rejective multiple test procedure. Scandinavian Journal

of Statistics, 6, 65–70

Hughes, E.J., et al. A dedicated neonatal brain imaging system. *Magn Reson Med*. *78*, 794-804

(2017).

Hurk, J. van den, Baelen, M. V., & Beeck, H. P. O. de. (2017). Development of visual category

selectivity in ventral visual cortex does not require visual experience. *Proceedings of the*

*National Academy of Sciences*, *114*(22), E4501–E4510.

Jabès, A., Lavenex, P. B., Amaral, D. G., & Lavenex, P. (2011). Postnatal development of the

hippocampal formation: A stereological study in macaque monkeys. *The Journal of*

*Comparative Neurology*, *519*(6), 1051–1070.

Jenkinson, M., Beckmann, C. F., Behrens, T. E. J., Woolrich, M. W., & Smith, S. M. (2012).

FSL. *NeuroImage*, *62(2)*, 782-790.

Keresztes, A., Ngo, C. T., Lindenberger, U., Werkle-Bergner, M., & Newcombe, N. S. (2018).

Hippocampal Maturation Drives Memory from Generalization to Specificity. *Trends in*

*Cognitive Sciences*, *22*(8), 676–686.

Langnes, E., Vidal-Piñeiro, D., Sneve, M. H., Amlien, I. K., Walhovd, K. B., & Fjell, A. M.

(2018). Development and Decline of the Hippocampal Long-Axis Specialization and

Differentiation During Encoding and Retrieval of Episodic Memories. *Cerebral Cortex*,

*29*(8), 3398–3414.

Larson-Prior, L. J., Zempel, J. M., Nolan, T. S., Prior, F. W., Snyder, A. Z., & Raichle, M. E.

(2009). Cortical network functional connectivity in the descent to sleep. *Proceedings of the*

*National Academy of Sciences*, *106*(11), 4489–4494.

Lavenex, P., & Banta Lavenex, P. (2013). Building hippocampal circuits to learn and remember:

Insights into the development of human memory. *Behavioural Brain Research*, *254*, 8–21.

Lin, W., Zhu, Q., Gao, W., Chen, Y., Toh, C.-H., Styner, M., … Gilmore, J. H. (2008).

Functional Connectivity MR Imaging Reveals Cortical Functional Connectivity in the

Developing Brain. *American Journal of Neuroradiology*, *29*(10), 1883–1889.

Linke, A. C., Wild, C., Zubiaurre-Elorza, L., Herzmann, C., Duffy, H., Han, V. K., Lee, D. S. C.,

& Cusack, R. (2018). Disruption to functional networks in neonates with perinatal brain

injury predicts motor skills at 8 months. *NeuroImage. Clinical, 18,* 399–406.

Liu, W.-C., Flax, J. F., Guise, K. G., Sukul, V., & Benasich, A. A. (2008). Functional

connectivity of the sensorimotor area in naturally sleeping infants. *Brain Research*, *1223*,

42–49.

Liu, X., Yanagawa, T., Leopold, D. A., Fujii, N., & Duyn, J. H. (2015). Robust Long-Range

Coordination of Spontaneous Neural Activity in Waking, Sleep and Anesthesia. *Cerebral*

*Cortex*, *25*(9), 2929–2938.

Makropoulos, A., et al. Automatic whole brain MRI segmentation of the developing neonatal

brain. *IEEE Transactions on Medical Imaging, 33*, 1818-1831 (2014).

Makropoulos, A., et al. The developing human connectome project: A minimal processing

pipeline for neonatal cortical surface reconstruction. *Neuroimage*, *173*, 88-112 (2018).

Murphy, K., & Fox, M. D. (2017). Towards a consensus regarding global signal regression for

resting state functional connectivity MRI. *NeuroImage*, *154*, 169–173.

Mitra, A., Snyder, A. Z., Tagliazucchi, E., Laufs, H., Elison, J., Emerson, R. W., Shen, M. D.,

Wolff, J. J., Botteron, K. N., Dager, S., Estes, A. M., Evans, A., Gerig, G., Hazlett, H. C.,

Paterson, S. J., Schultz, R. T., Styner, M. A., Zwaigenbaum, L., IBIS Network, …

Raichle, M. (2017). Resting-state fMRI in sleeping infants more closely resembles adult

sleep than adult wakefulness. *PloS One*, *12*(11),

Natu, V. S., Gomez, J., Barnett, M., Jeska, B., Kirilina, E., Jaeger, C., Zhen, Z., Cox, S., Weiner,

K. S., Weiskopf, N., & Grill-Spector, K. (2019). Apparent thinning of human visual

cortex during childhood is associated with myelination. *Proceedings of the National*

*Academy of Sciences of the United States of America*, *116*(41), 20750–20759.

Ofen, N., Kao, Y.-C., Sokol-Hessner, P., Kim, H., Whitfield-Gabrieli, S., & Gabrieli, J. D. E.

(2007). Development of the declarative memory system in the human brain. *Nature*

*Neuroscience*, *10*, 1198.

Osher, D. E., Brissenden, J. A., & Somers, D. C. (2019). Predicting an individual's dorsal

attention network activity from functional connectivity fingerprints. *Journal of*

*Neurophysiology*, *122*(1), 232–240.

Poppenk, J., Evensmoen, H. R., Moscovitch, M., & Nadel, L. (2013). Long-axis specialization of

the human hippocampus. *Trends in Cognitive Sciences*, *17*(5), 230–240.

Poppenk, J., & Moscovitch, M. (2011). A Hippocampal Marker of Recollection Memory Ability

among Healthy Young Adults: Contributions of Posterior and Anterior Segments.

*Neuron*, *72*(6), 931–937.

Power, J. D., Barnes, K. A., Snyder, A. Z., Schlaggar, B. L., & Petersen, S. E. (2012). Spurious

but systematic correlations in functional connectivity MRI networks arise from subject

motion. *NeuroImage*, *59*(3), 2142–2154.

Power, J. D., Mitra, A., Laumann, T. O., Snyder, A. Z., Schlaggar, B. L., & Petersen, S. E.

(2014). Methods to detect, characterize, and remove motion artifact in resting state fMRI.

*NeuroImage*, *84*, 320–341.

Prabhakar, J., Johnson, E. G., Nordahl, C. W., & Ghetti, S. (2018). Memory-related hippocampal

activation in the sleeping toddler. *Proceedings of the National Academy of Sciences of the*

*United States of America*, 115(25), 6500–6505. https://doi.org/10.1073/pnas.1805572115

Raichle, M. E. (2009). A Paradigm Shift in Functional Brain Imaging. *Journal of Neuroscience*,

*29*(41), 12729–12734.

Riggins, T., Geng, F., Blankenship, S. L., & Redcay, E. (2016). Hippocampal functional

connectivity and episodic memory in early childhood. *Developmental Cognitive*

*Neuroscience*, *19*, 58–69

Roffwarg, H. P., Muzio, J. N., & Dement, W. C. (1966). Ontogenetic development of the human

sleep-dream cycle. *Science (New York, N.Y.)*, *152*(3722), 604–619.

Salimi-Khorshidi, G., et al. Automatic denoising of functional MRI data: combining 409

independent component analysis and hierarchical fusion of classifiers. *Neuroimage*, *90,*

449-468  (2014)

Salzwedel, A. P., Stephens, R. L., Goldman, B. D., Lin, W., Gilmore, J. H., & Gao, W. (2019).

Development of Amygdala Functional Connectivity During Infancy and Its Relationship

With 4-Year Behavioral Outcomes. *Biological Psychiatry: Cognitive Neuroscience and*

*Neuroimaging*, *4*(1), 62–71.

Satterthwaite, T. D., Elliott, M. A., Gerraty, R. T., Ruparel, K., Loughead, J., Calkins, M. E., …

Wolf, D. H. (2013). An improved framework for confound regression and filtering for

control of motion artifact in the preprocessing of resting-state functional connectivity data.

*NeuroImage*, *64*, 240–256.

Seress, L. (2007). Comparative anatomy of the hippocampal dentate gyrus in adult and

developing rodents, non-human primates and humans. In H. E. Scharfman (Ed.),

*Progress in Brain Research* (pp. 23–798).

Smith, S. M., Fox, P. T., Miller, K. L., Glahn, D. C., Fox, P. M., Mackay, C. E., … Beckmann,

C. F. (2009). Correspondence of the brain's functional architecture during activation and

rest. *Proceedings of the National Academy of Sciences of the United States of America*,

*106*(31), 13040–13045.

Smith, S. M., & Nichols, T. E. (2009). Threshold-free cluster enhancement: Addressing

problems of smoothing, threshold dependence and localisation in cluster inference.

*NeuroImage*, *44*(1), 83–98.

Smith, S. M., Vidaurre, D., Beckmann, C. F., Glasser, M. F., Jenkinson, M., Miller, K. L., …

Van Essen, D. C. (2013). Functional connectomics from resting-state fMRI. *Trends in*

*Cognitive Sciences*, *17*(12), 666–682.

Snedecor, GW, Cochran, WG - Ames: Iowa State Univ. Press Iowa, 1989

Sporns, O. (2013). Structure and function of complex brain networks. *Dialogues in Clinical*

*Neuroscience*, *15*(3), 247–262.

Strange, B. A., Witter, M. P., Lein, E. S., & Moser, E. I. (2014). Functional organization of the

hippocampal longitudinal axis. *Nature Reviews Neuroscience*, *15*(10), 655–669.

Thompson, D. K., Wood, S. J., Doyle, L. W., Warfield, S. K., Lodygensky, G. A., Anderson, P.

J., Egan, G. F., & Inder, T. E. (2008). Neonate hippocampal volumes: Prematurity,

perinatal predictors, and 2-year outcome. *Annals of Neurology*, *63*(5), 642–651.

Tobyne, S. M., Somers, D. C., Brissenden, J. A., Michalka, S. W., Noyce, A. L., & Osher, D. E.

(2018). Prediction of individualized task activation in sensory modality-selective frontal

cortex with 'connectome fingerprinting.' *NeuroImage*, *183*, 173–185.

Travaglia, A., Steinmetz, A. B., Miranda, J. M., & Alberini, C. M. (2018). Mechanisms of

critical period in the hippocampus underlie object location learning and memory in infant

rats. *Learning & Memory*, *25*(4), 176–182.

Wael, R. V. de, Larivière, S., Caldairou, B., Hong, S.-J., Margulies, D. S., Jefferies, E., …

Bernhardt, B. C. (2018). Anatomical and microstructural determinants of hippocampal

subfield functional connectome embedding. *Proceedings of the National Academy of*

*Sciences*, *115*(40), 10154–10159.

Wheelock, M. D., Hect, J. L., Hernandez-Andrade, E., Hassan, S. S., Romero, R., Eggebrecht, A.

T., & Thomason, M. E. (2019). Sex differences in functional connectivity during fetal brain

development. *Developmental Cognitive Neuroscience*, *36*, 100632.

Yan, C.-G., Cheung, B., Kelly, C., Colcombe, S., Craddock, R. C., Di Martino, A., … Milham,

M. P. (2013). A comprehensive assessment of regional variation in the impact of head

micromovements on functional connectomics. *NeuroImage*, *76*, 183–201.

Yeo, B. T., Krienen, F. M., Sepulcre, J., Sabuncu, M. R., Lashkari, D., Hollinshead, M., …

Buckner, R. L. (2011). The organization of the human cerebral cortex estimated by

intrinsic functional connectivity. *Journal of Neurophysiology*, *106*(3), 1125–1165.

**Legends**

**Figure 1: Preliminary Data Checks** a) Gray matter (red), white matter (yellow) and network

registrations on the anatomical images of a representative adult and neonate subject; registration

image: blue=hippocampus, white = Vis, red = SM, purple (dark) = DA, pink = VA, green = Lim,

yellow =  FP, orange = DM b) voxelwise correlations distributions of a representative adult and

neonate c) between-subject and between-group correlations demonstrate high within-group

reliability of connectivity but low between-group reliability between adults and neonates. (*)

indicates significance at p<0.05; ns denotes non-significance

**Figure 1-1: tSNR-Matched Adult Hippocampus to Networks.** Hippocampal-network

connectivity of 40 tSNR-matched HCP adults again shows very similar results to the motion-

matched and binarized-hippocampal analyses. Hippocampal connectivity in adults shows a clear

hierarchy, with strong positive connectivity to Lim and DMN and negative connectivity to FP

and VA (*) indicates significance at pHB<0.05; (***) indicates significance at pHB<0.005.

**Figure 2: Hippocampal Connectivity to Cortical Networks** a) Comparison of hippocampal

connectivity to the seven cortical networks in adults showed a hierarchy of hippocampal

connectivity, whereby the highest FC was with Lim, followed by DM, SM, and Vis, almost no

FC with DA, and negative FC with FP and VA. b) In contrast, neonates show the same level of

FC to almost all of the 7 networks. Rose plot to the right shows adult connectivity compared to

neonates to highlight the differences between groups in the pattern of hippocampal FC to these

networks. Brain images on the top right depict connectivity between the hippocampus (left) and

the seven cortical networks (right). (*) indicates significance at $p_{HB}<0.05$; (***) indicates

significance at $p_{HB}<0.005$. Lim=Limbic; DM=Default Mode; SM=Somatomotor; Vis=visual;

DA=Dorsal Attention; FP=FrontoParietal; VA=Ventral Attention

**Figure 2-1: Binarized Whole Hippocampus to Networks.** Hippocampal connectivity to the

networks using a binarized HCP/dHCP hippocampal ROI yields very similar results to the ANTs

registered hippocampal ROI (see figure 2). As with the initial analysis, hippocampal connectivity

in adults shows a clear hierarchy whereas neonates display very few differences in hippocampal

connectivity strength to the networks. (*) indicates significance at pHB<0.05; (***) indicates

significance at pHB<0.005.

770
771     **Figure 3:  Hippocampal Connectivity to Cortex.** Comparison of adult and neonate

772     hippocampal connectivity to the cortex at a voxelwise grain. FWE-corrected results for the

773     contrast of neonate > adult connectivity is shown in warm colors and the contrast of adult >

774     neonate is denoted by cool colors.

775
**Figure 3-1: Hippocampal Connectivity to Cortex, Adult>Neo** Clusters are listed from largest

to smallest.  Peak coordinates (MAX) are listed in MNI space as well as center of gravity (COG)

for each cluster

**Figure 3-2:  Hippocampal Connectivity to Cortex, Neo>Adult** Clusters are listed from largest

to smallest. Peak coordinates (MAX) are listed in MNI space as well as center of gravity (COG)

for each cluster

**Figure 4: Anterior/Posterior Hippocampal Connectivity to Networks** a) Anterior vs

posterior hippocampal-network connectivity in adults b) Anterior vs posterior hippocampal-

network connectivity in neonates c) Rose plot comparing anterior vs posterior hippocampal-

network connectivity pattern in adults d) Rose plot comparing anterior vs posterior hippocampal-

network connectivity pattern in neonates. Brain image to the right shows the anterior (red) and

posterior (yellow) hippocampal labels (*) indicates significance at $p_{HB}<0.05$; (***) indicates

significance at $p_{HB}<0.005$.

**Figure 5: Connectivity along the Long-Axis Gradient to Networks** Comparison of the

connectivity of the long axis gradient of the hippocampus to the 7 networks in a) adults and b)

neonates. The slices are arranged anterior-to-posterior. Lighter coloring surrounding each line

represents the standard error. Brain image on the right demonstrates the hippocampus (blue)

segmented into slices (white lines). (*) are slices where the ANOVA shows an interaction

between network and group at $p<0.001$.

**<u>Figures</u>**

(Figure 1)

(Extended Data for Figure 1, Figure 1-1)

(Figure 2)

(Extended data for Figure 2; Figure 2-1)

(Figure 3)

| Cluster | Regions | Voxels | MAX | MAX (X) | MAX (Y) | MAX (Z) | COG (X) | COG (Y) | COG (Z) |
|---|---|---|---|---|---|---|---|---|---|
| 1 | (L) Posterior Cingulate; Isthmus Cingulate, Precuneus | 2370 | 8.08 | -10 | -57 | 17 | -6.23 | -55.2 | 20.8 |
| 2 | (R) Isthmus Cingulate; Precuneus | 1255 | 8.27 | 15 | -54 | 19 | 9.3 | -56.1 | 19.1 |
| 3 | (L) Inferior Parietal | 574 | 6.85 | -42 | -77 | 43 | -44.3 | -74.8 | 39.2 |
| 4 | (L) Middle Temporal Cortex | 403 | 5.72 | -62 | -1 | -20 | -63.2 | -7.54 | -18.2 |
| 5 | (L) Medial Orbital Frontal | 303 | 7.45 | -10 | 39 | -11 | -7.75 | 41.9 | -11.7 |
| 6 | (L) Middle Temporal Cortex; Superior Temporal Cortex | 235 | 6.88 | -52 | -13 | -14 | -53.1 | -11.6 | -13.7 |

813    (Extended Data for Figure 3; Figure 3-1)

| Cluster | Regions | Voxels | MAX | MAX (X) | MAX (Y) | MAX (Z) | COG (X) | COG (Y) | COG (Z) |
|---|---|---|---|---|---|---|---|---|---|
| 1 | (R) Rostral Middle Frontal; Pars Triangularis; Pars Orbitalis; Lateral Orbitofrontal; Pars Opercularis; Insula; Caudal Middle Frontal; Precentral; Postcentral | 16290 | 8.72 | 57 | 14 | 4 | 44.9 | 26.2 | 21.8 |
| 2 | (R) Superior Frontal; Paracentral | 4702 | 6.88 | 4 | 26 | 61 | 9.85 | 12.7 | 59.4 |
| 3 | (L) Supramarginal | 3278 | 8.54 | -65 | -42 | 34 | -59.6 | -38.9 | 30.7 |
| 4 | (R) Supramarginal; Inferior Parietal | 3226 | 7.93 | 62 | -36 | 48 | 61.8 | -35.9 | 37.1 |
| 5 | (R) Lingual; Pericalcarine (L) Lingual; Pericalcarine | 2794 | 6.22 | -19 | -66 | 2 | 1.81 | -77.1 | 5.9 |
| 6 | (L) Rostral Middle Frontal | 796 | 6.85 | -34 | 51 | 29 | -36.3 | 46.6 | 28.9 |
| 7 | (R) Lateral Orbitofrontal; Pars Orbitalis | 458 | 5.59 | 46 | 22 | -7 | 39.4 | 24.6 | -7.38 |
| 8 | (L) Superior Frontal | 350 | 5.74 | -17 | 7 | 66 | -13.6 | 8.52 | 69.4 |
| 9 | (R) Insula | 238 | 5.25 | 42 | 3 | -6 | 40.4 | 6.44 | -3.65 |

821    (Extended Data for Figure 3; Figure 3-2)

824     (Figure 4)

(Figure 5)

828
829