TITLE

 Molecular principles of Piezo1 activation by increased membrane tension.

4 **AUTHOR NAME AND AFFILIATIONS**

 Dario De Vecchis[1], David J Beech[1] and Antreas C Kalli[1,2*]

 [1]School of Medicine and [2]Astbury Centre for Structural Molecular Biology, University of Leeds,

 Leeds LS2 9JT, United Kingdom.

 *Corresponding author

 SUMMARY PARAGRAPH

14 **Piezo1 is a mechanosensitive channel involved in many cellular functions and responsible for**
 sensing shear-stress and pressure forces in cells1–3. Piezo1 plays a critical role in the circulatory
 system and tissue development. Mutations on Piezo1 are linked to human diseases such as
 lymphedema2,4 or hematological disorders such as hemolytic anaemia5 and resistance to
 malaria6. Hypotheses for Piezo1 gating include the "force-from-lipids" principle7,8 that
 suggests that Piezo1 senses mechanical forces through the bilayer1,9 and a direct involvement
 of the cytoskeleton as well as the extracellular matrix in Piezo1 activation10,11. However, the
 molecular and structural changes underpinning the Piezo1 gating mechanism and how the
 channel senses forces in the membrane remain unknown. Here we reveal the activation
 mechanism of Piezo1 and the structural rearrangements that occur when Piezo1 moves from a
 closed to an open state when mechanical tension is applied to the cell membrane. Our results
 show that Piezo1's curved shape is stable in a native-like model membrane without tension
 creating a membrane indentation with a trilobed topology. Upon stretching Piezo1 adapts to
 the stretched bilayer by flattening and expansion of its blade region. In our simulations Piezo1
 expands up to a planar circular area of ~680 nm^2 comparable with previous structural data and
 hypotheses12–14. Piezo1 flattening and expansion results in changes in the beam helix tilt angle.
 These movements result in the tilting and lateral movement of the pore lining TM37 and TM38
 helices. This leads to the opening of the channel and to the movement of lipids that occupy
 Piezo1 pore region outside of this region, revealing for the first time the structural changes that
 happen during Piezo1 mechanical activation. The changes in the blade region are transmitted
 to helices TM37 and 38 via hydrophobic interactions and by interactions of neighbouring

subunits via the elbow region. The flat structure of Piezo1 identified in this study exposes the C-terminal extracellular domain (CED) that in the closed state is hidden in the membrane and presumably from shear stress. Our results provide new structural data for different states of Piezo1 and suggest the molecular principles by which mechanical force opens the Piezo1 channel, thus coupling force to physiological effect via ion permeation.

Figure 1. Piezo1 channel flattens and its CED becomes exposed in response to increased membrane tension. For each system, the applied negative pressure in each simulation is indicated with different colours: black, 1 bar; red, -5 bar; green, -20 bar; blue, -30 bar; orange, -40 bar. (**a**) Snapshots of the last frame from simulations of Piezo1 embedded in a model membrane bilayer. (**b**) Correlation between the area, i.e. x-side × y-side, of the simulated box (A_{box}) and the root mean-square deviation (RMSD) calculated using the protein Cα of the Piezo1 blades. (**c**) RMSD calculated using the protein Cα of the C-terminal extracellular domain (CED), dashed lines (lower values), and of the rest of the protein (i.e. the Piezo1 blades), straight lines (higher values). (**d**) Projected area of the Piezo1 membrane indentation on the membrane plane (A_{proj}, indicated by a dashed blue circle in **a**) as a function of time. For the -40 system the timeline is the average between the two repeats. The standard deviation is indicated.

To investigate how mechanical forces act on the Piezo1 channel, we simulated the recent Piezo1

structure solved by cryo-electron microscopy (cryo-EM) in a model membrane using a serial multi-

scale molecular dynamics (MD) simulation approach. Missing residues were added to the structure

prior to the simulations (see Methods). An initial coarse-grained (CG) simulation of 500 ns was

carried out to equilibrate the bilayer around Piezo1 and create the unique Piezo1 footprint suggested

previously[12–14]. Although the CG approach was critical for creating the Piezo1/membrane system,

it does not allow us to study the Piezo1 dynamics. For this reason, the system was backmapped to an

atomistic resolution and further simulated in independent repeat simulations applying pressure to the

bilayer (i.e. along x and y axes) ranging from the native-like pressure of +1 bar to -40 bar as described

in Methods. Under these conditions, we were able to calculate the lateral pressure (P_L) applied on the

bilayer which is 14.2 mN/m for the -5 bar, 45.8 mN/m for the -20 bar, 59 mN/m for the -30 bar and

67.8 ± 0.35 mN/m for the -40 bar system. The asymmetric bilayer used in our simulations mimics the

native endothelial membrane in which Piezo1 functions[15] (see Methods).

Tension causes Piezo1 to flatten and expand

The different pressures cause expansion of the simulated box to a different degree with the membrane

in the system adapting to each pressure. This results in an increase of the area per lipid (APL) in both

leaflets for all systems; the APL for the system with the highest tension in the bilayer was $90.09 \pm$

0.84 Å^2 and 89.46 ± 0.96 Å^2 the upper and the lower leaflets, respectively (see Supplemental Table

S1 for all APLs). A similar approach revealed that an increase of the APL resulted in the transition of

the TREK-2 mechanosensitive channel for a down to an up conformation[16].

Our analysis above shows that the APL reaches a plateau for both leaflets within the first 10 ns of the

simulation. Therefore, each system was simulated for at least 50 ns to allow sufficient time for Piezo1

to equilibrate in the bilayer. Indeed, our analysis shows that the opening of the channel occurs after

the first 10 ns of simulation in conjunction with the increase of the APL (Fig. 4a, b and Supplemental

Table S1). However, to further understand if more extended simulation time alters the APL, two of

our systems the -30 bar and the +1 bar systems were extended to 100 ns. In both cases the analysis

confirms than no change in the APL is observed in the longer simulations (Supplemental Table S1).

Moreover, a further repeat simulation at -40 bar was carried out obtaining a comparable result, thus

indicating that the systems are equilibrated, and the simulated time is sufficient to study the Piezo1

structural transitions under tension. The Piezo1 structure shows a remarkable adaptation to a stretched

bilayer with its N-terminal blades, which remain embedded within the bilayer during all simulations,

to extend towards more flatten conformations (Fig. 1a). The degree of flattening was different for the

different tensions with the system with the highest tension resulting in an almost flat Piezo1 structure

(Fig. 1a). Fig. 1b shows the correlation between the area of the simulated box and the root mean-

square deviation (RMSD) calculated using the Cα atoms of the blades, demonstrating that they both

increase until the bilayer stops expanding. Note that in the bilayer without tension (+1 bar) Piezo1

maintains its curved shape even though this simulation is twice as long (i.e. 100 ns) as some of the

simulations with a tension applied to the bilayer. This shows that the curved shape of Piezo1 blades

forces the bilayer to adopt a curved trilobed shape. Interestingly, whilst Piezo1 reaches a flat

conformation in both the -30 bar and -40 bar systems, larger conformational changes with more

opening of the pore are observed in the -40 bar simulations. Even when the -30 bar systems were

extended up to 100 ns (compared to 50 ns of the -40), the structural changes in this system were

smaller (Fig. 1a, b), suggesting that there is a cross-talk between Piezo1 and bilayer with Piezo1

sensing forces from the bilayer after it reaches the flat conformation. Our RMSD analysis shows the

Piezo1 blades as the major contributors to the RMSD drift, whereas the CED mostly retained its initial

conformation with comparable values between the systems (Fig. 1c).

Figure 2. Coordinated tilting of Piezo1 channel beams and ion pore helices in response to

increased membrane tension.(**a**) Partial density profile along the z direction (that is perpendicular

to the bilayer x,y plane) for the protein (black line), the membrane (magenta line) and the solvent (i.e.

water and ions, cyan line). Values from the first repeat at -40 bar are shown here. Insets are the last

frames from simulations as shown in Fig. 1a. (**b**) Tilt angle relative to the bilayer normal for the beam

helix. (**c**) Correlation between the RMSD of the Piezo1 blades and the tilt angle relative to the bilayer

normal of the TM38 helix. (**d**) Correlation between the distances from the centre of mass of the TM38

or TM37 helices and the Piezo1 pore (**e**) Correlation between the RMSD of the Piezo1 blades and the

tilt angle relative to the bilayer normal of the TM37 helix (**f**) Snapshots of the last frame from the +1

bar and -40 bar systems with the Piezo1 triskelion shown from the extracellular side. The centre of

the triskelion is highlighted. The TM38 (cartoon) and the TM37 (ribbon) from each Piezo1 chain are

also shown. Membrane and solvent are not shown for clarity. The colour code for the applied negative

pressure in each simulation is indicated and is the same as in Fig. 1.

In addition, our results suggest that upon flattening Piezo1 also expands thus increasing the in-plane

projection of the Piezo1 membrane indentation (A_{proj}; Fig 1a). This is in good agreement with recent

studies that also suggested an increase of the A_{proj} involved in the Piezo1, and recently Piezo2,

activation[12–14]. The A_{proj} can be approximated by a circle. By approximating the Piezo1 triskelion to

a regular triangle, we calculated the side of this triangle to further determine the circumscribed circle

(i.e. the A_{proj}) as described in the Methods. Our approximation is justified by the moderate standard

deviation values of the side of the triskelion (i.e. the side of the triangle) which is ± 0.23 nm for the

+1 bar, ± 0.53 nm for the -5 bar, ± 0.95 nm for the -20 bar, ± 0.88 nm for the -30 bar and ± 1.07 nm

and ± 1.23 nm for the two repeats at -40 bar. This result is also indicative of a concurrent movement

of the Piezo1 blades that flatten at the same time. Fig. 1d shows how the increase of applied

mechanical force during simulations (e.g. more tension), causes an increase of the A_{proj} values which

reach ~680 nm^2 at -40 bar (Fig. 1d, orange line). Recent cryo-EM structural data of the full-length

Piezo2, estimated a A_{proj} of the dome to 700 nm^2 [14]. Although the current value for the full-length

Piezo1 is still unknown, this agreement shows that upon flattening Piezo1 covers the area that is

within the dome in the closed state[14]. This is in good agreement with the suggestion that the dome

provides an energy store for Piezo1 activation[12]. We also note that in our atomistic simulations, even

the +1 bar system shows a mild increase of the A_{proj} with respect to the initial coordinates from the

cryo-EM (Fig. 1d, black line). Overall, our approach provides novel molecular insights into the

activation and dynamics of Piezo1 in a native-like lipid environment in which this channel functions.

Moreover, we dynamically quantify the different A_{proj} expansions under different mechanical stress

and the area covered by a flat activated Piezo1 channel. Finally, the extended structure obtained at -

40 bar is corroborated by recent structural data which positively correlate the size of the vesicle with

a flattened Piezo1 structure[13]. Flattening of Piezo1 may also be needed for Yoda1 binding[17].

Figure 3. Expansion of the Piezo1 channel pore in response to increased membrane tension. (**a**) Volume of the cavity of the Piezo1 pore region as a function of time for each simulated system. The applied negative pressure in each simulation is indicated with different colours: black, 1 bar; red, -5 bar; green, -20 bar; blue, -30 bar; orange, -40 bar. (**b**) Number of water molecules overlapping with

the calculated cavity in **a** as a function of time for each simulated system. (**c**) Volume of the cavity

within the core of the Piezo1 structure, for the 1 bar and the -40 bar systems. The structures are the

last frames from each simulation and show the Piezo1 pore region. For the -40 bar system the second

repeat simulation is shown. The cavity (sliced) is shown as spheres and coloured according to the

occupancy during the trajectory, from less populated (blue) to more populated (red). Residues

selected for the calculation of the cavity are shown in ribbon (including the TM38 helix, see Methods).

TM37, elbow and base helix are represented as cartoon cylinders. The Piezo1 chains are shown in the

same colours as in Fig. 1a. (**d**) Water molecules within 8 Å from the residues selected for the

calculation of the cavity for the 1 bar and the -40 bar systems. Water molecules are depicted in light

cyan transparent spheres. The structures are the last frames from simulations and show the Piezo1

pore region. For the -40 bar system the second repeat is shown. The protein orientation and colour

code are the same as in **c**. (**e**) Detail of the Piezo1 channel mouth for the 1 bar and the -40 systems

shown from a side (above) and a extracellular (below) view. The structures are the same shown in **c**

and **d** and show the hydrophobic interactions between the pore-lining helices TM38 (yellow residues)

and the TM37 (dark grey residues). The salt bridge between the elbow and the TM38 from a

neighbouring subunit is also shown (red and blue residues). Residues from the Piezo1 hydrophobic

gate (residues L2475, V2476) are in light grey.

Structural rearrangements in Piezo1 pore

Piezo1 flattening caused by membrane tension results in the displacement of the CED relative to the

membrane plane (Fig 2a). Although our simulations did not show any changes in the position of the

CED domain relative to the transmembrane (TM) domain, CED is hidden within the dome at the

beginning of the simulations or in the system without tension applied in the bilayer. Flattening of

Piezo1 and of the bilayer results in the exposure of CED as it emerges outside the dome. The degree

of CED exposure above the dome increases as the tension in the bilayer increases; in the -30 bar and

-40 bar systems in which a flat Piezo1 was reached CED is fully exposed. The density analysis also

revealed that a considerable portion of the Piezo1 structure remained consistently exposed to the

cytoplasm even after the expansion (Fig 2a). It is possible that these cytoplasmic regions remain

exposed during activation providing a platform for Piezo1 interactions with the cytoskeleton, which

is in agreement with its mechanoprotective role in Piezo1 function[9].

Another Piezo1 component which undergoes noticeable structural rearrangement is the beam helix

(residues 1300-1365; Fig. 2b). Our data reveal that during the simulations with tension in the bilayer

there is a change in the tilting angle of each Piezo1 beam (going from ~65° to ~85°), until reaching a

position that is almost parallel to the bilayer surface when Piezo1 is in a flat conformation. Interestingly, the beam tilt movement is correlated with the change in the RMSD (Supplemental Fig. S1a) suggesting that this movement contributes to Piezo1 flattening. The beam is a long α-helix exposed to the cytoplasm which underpasses a great portion of each blade, precisely the three C-terminal proximal bundles. Therefore, our data suggest that each Piezo1 beam may act as a set of long levers and their concerted movement during mechanical sensing may possibly contribute to gating.

The precise molecular details of Piezo1 gating are still largely unknown. The protein is also characterised by a fast inactivation phase after gating[18,19]. Recent observations proposed a functional gate possibly regulating Piezo1 inactivation[20]. This gate is composed by the two conserved hydrophobic residues Leu2475 and Val2476 in the pore-lining TM helix 38[20]. Calculation of the tilt angle of the TM38 relative to the bilayer normal shows that higher negative pressure, and thus larger tension, corresponds to a higher tilting of the pore-lining TM38 (Supplemental Fig. S1b). The highest tilt angle is observed in the -40 bar system (Fig. 2c). Calculation of the distance of TM38 relative to the pore centre of mass also shows that larger tension, corresponds to a larger displacement (Supplemental Fig. S1c). Therefore, during flattening and expansion TM38 both increases its tilt angle and its distance from the pore centre of mass. The change in the TM38 tilting and distance from the channel pore is correlated with the changes in the RMSD calculated for the Piezo1 blades (Fig. 2c and Supplemental Fig. S1c). This shows that TM38 displacement/tilting occurs whilst Piezo1 flattens and expands (Fig. 2f). Interestingly, we found that the distance from the pore and tilting of the TM38 also correlates with the distance from the pore and tilting of the TM37 (Fig. 2d, e and Supplemental Fig. S1d, e), which is located anti-parallel (Fig. 2f). In conclusion, our simulations suggest a simultaneous lateral movement and tilting of the TM38 and TM37 that results in the opening of the pore.

Tension leads to Piezo1 channel opening

Calculation of the water-filled cavity in Piezo1 pore region shows that the volume of this cavity increases with the applied negative pressures in our simulations until it reaches ~75000 Å^3 in the -40 bar (Fig. 3a). In our calculation the cavity is defined by the three TM38 (one from each subunit) and also some regions of the CED (see Methods), because the loops that connect this domain with the TM37/38 form a structure similar to a "cage" just above the channel mouth which is characterised by a funnel-shape (Fig. 3c). Fig. 3a also shows that tension of -30 bar on the membrane, although it contributes to flatting of Piezo1 structure and almost maximise the beam orientation, is not sufficient to open the channel pore and maximise the volume of the cavity; the cavity in the -30 system is not

hydrated as much as in the -40 bar (Fig. 3b, d). Note that in the simulations of the system with no tension to the bilayer (1 bar) the cavity did not increased and the channel was not hydrated. We also found that lipid and cholesterol molecules initially overlap with this volume and therefore they are within the Piezo1 pore, but they increasingly leave the cavity as higher tension is applied to the bilayer (Fig. 4a). Three independent system set-ups followed by three independent CG equilibrations demonstrate that lipids get within the channel mouth during the equilibration phase. This demonstrates that there is enough space in the current closed structure of Piezo1 for lipids to occupy the channel mouth. This may also suggest that the presence of lipids in the pore may be required in order to maintain the channel mouth closed. Our results show that a higher tension to the membrane (that increases the APL in the proximity of the channel mouth), results in higher volume in the pore cavity as it enables lipids and cholesterol that occupy the channel mouth to move outside of it in addition to a larger movement of the pore lining helices. This hypothesis may explain biochemical data which suggest a functional role of lipids in Piezo1 regulation[21–23]. Therefore, our simulations suggest that the tension applied to the bilayer during functional activation, results not only in the displacement of TM38 and TM37 but also in the movement of lipids outside the Piezo1 pore. These two events occur simultaneously and may both be required for Piezo1 full opening.

An open question is how the mechanical stimuli are transferred from the blades to the channel pore. Our results suggest that the flattening and expansion of the blades is transmitted to the TM helices 37 and 38 guiding the pore opening. Noticeably, the pore region of the homologous Piezo2, has a remarkably similar structure14 which lead us to postulate a similar mechanism. In Piezo1 the pore-lining TM38, is located parallel, although with an opposite orientation, to the TM37. TM37 together with the elbow region (residues 2116-2142), a base helix (residues 2149 to 2175) and hairpin helices (residues 2501 to 2534) from all three subunits, create a cuff that encloses the pore-lining TM3812,13 (Fig. 3e). Noticeably, in all our simulations we noticed two main pivotal interactions: the first is the interaction between the TM37 and TM38 via the hydrophobic residues Ile2466, Leu2469, Tyr2470, Ile2473 and Val2477, that are adjacent to the aforementioned hydrophobic functional gate in TM38 (residues Leu2475 and Val2476[20]) and the residue Phe2198, Ile2202, Phe2205 and Phe2209 from the TM37 (Fig. 3e). Therefore, extension and expansion of the blades cause the displacement of TM37 that in turn "pulls" the pore-lining helix TM38 due their hydrophobic interactions. This results in the opening of the channel and might explain why in the -40 bar simulation, in which more expansion was observed, the pore was more open (Fig. 3e).

The second pivotal interactions occur in a region towards the C-terminal end of TM37 and TM38. Previous work suggested the importance of the negative charge at position Glu2133 (Glu2416 in Piezo2) for conductance and ion selectivity[24]. In particular, the single mutant E2133A showed half the unitary conductance with respect to the wildtype phenotype and the negative charge from the glutamate has been suggested to allosterically modulate the selectivity of the Piezo1 filter[24]. The Piezo1 cryo-EM structure allowed to precisely map the position of the Glu2133, which is located in a helix named elbow. Interestingly, the tip of each elbow points towards the TM38 of another Piezo1 subunit and in our model we found that Glu2133 on the elbows forms a salt bridge with the Arg2482 in a neighbouring subunit in agreement with previous suggestions[25] (Fig. 3e). This salt bridge is retained in all our simulations between at least two Piezo1 chains with above 70% of occurrence, even after Piezo1 conformational rearrangement due to the applied negative pressure (Supplemental Table S2). This indicates that this interface is possibly maintained during the conformational transition we observed.

In agreement with this hypothesis, functional data[24] demonstrated that the mutant G2133D increases the conductance with respect to the wildtype, which is probably due to the short chain of the aspartate that would pull backward (i.e towards the elbow) the TM38, thus reducing the pore constriction. Furthermore, this hypothesis rationalises the effect of the less efficient mutant E2133K that was shown to decrease Piezo1 conductance[24]. That is, the mutation to lysine opposite to the Arg2482 would push forward the TM38 due to repulsion, contributing in the increase of the channel constriction. In the E2133K mutant, the salt bridge is broken and so is the connection between the elbow and the TM38 which results in an impairment of the functional displacement connected to the blade expansion. In this condition, although less conducting, the channel is still functional[24], which may be due to the aforementioned hydrophobic interactions between TM37 and TM38 in synergy with the proposed of lipid displacement which may act in concert. In conclusion, the predicted salt bridge between Glu2133 with Arg2482 in a neighbouring subunit would provide an interaction interface between the Piezo1 chains so that each blade extension would control the gating of a nearby Piezo1 chain (Fig. 4b). This may also provide a mechanism that ensures a prompt and mutual propagation of sensing between Piezo1 subunits.

Figure 4. Proposed mechanism for the Piezo1 channel opening in response to membrane tension. (**a**) Snapshots from the second repeat simulation at -40 bar with the associated simulation time are shown. Membrane bilayer and solvent (i.e. water and ions) are not shown for clarity. The three Piezo1 subunits are indicated with different colours as in Fig. 1a. In the extracellular view at the top, the CED domain has been removed for clarity. Phosphorous and oxigen atoms from lipid headgroups and cholesterol respectively are shown (only lipids that are within 20 Å from the residues selected for the calculation of the Piezo1 pore region cavity; see Methods). The colour code is: POPC, tan; POPE, pink; POPS, red; PIP_2, green; cholesterol, magenta. The Cα atoms from interacting residues between the helices TM37, TM38 and the elbow are indicated as spheres as in Fig. 3e. In the Piezo1 structures in the middle panel, water molecules within 8 Å from the residues selected for the calculation of the cavity (see Methods) are shown as light cyan transparent spheres. (**b**) Cartoon representation explaining the Piezo1 activation mechanism coupled to membrane tension. The colour code is the same as in **a**. The hydrophobic interactions between the TM37 and TM38 and the salt bridge between TM38 and the elbow from a neighbouring subunit, are indicated in yellow/grey and in blue/red, respectively. Lipids that initially occupy the Piezo1 channel mouth and progressively move away from it with the increase tension, are schematically depicted in green. Membrane tension and Piezo1 blade expansion are indicated by grey arrows.

In conclusion, this study reveals the molecular principles by which Piezo1 channel opens in response to increase membrane tension. The steps during Piezo1 opening are shown in Fig. 4b and are: 1) In a

membrane without tension, Piezo1's curved shape forces the bilayer to curve creating a characteristic

indentation with a trilobed topology. This indentation is stable in native condition and the CED is

hidden within it. In this inactive resting conformation, Piezo1 channel mouth is occupied by lipids.

2) Membrane tension on the bilayer results in the flattening and extension of its blades and tilting of

its beams helices. 3) Piezo1 expansion and flattening allows the blades to "pull" helices 37 and 38,

via hydrophobic interactions, to tilt and progressively move away from the channel pore opening

Piezo1. 4) In the open conformation Piezo1 blades are fully extended and flat, CED is exposed, beam

helices are almost parallel to the lipid bilayer. The elbow regions from neighbouring subunits, also

pull the pore-lining helices 38. At the same time the increase in the area per lipid due to tension allows

displacement of the lipids that occupy the channel mouth resulting in an open. Therefore, both

conformational changes and the increase in the APL that allows movement of lipids outside the

channel pore region are required for Piezo1 opening.

This work provides molecular understanding of the structural rearrangements within Piezo1 that lead

to gating and highlight the critical role of membrane lipids in Piezo1 opening. Our study demonstrates

how the fascinating fold and shape of Piezo channels allow this protein to sense mechanical force

thus playing a central role in transduction of mechanical signals throughout the membrane. Our

findings have important implications for understanding mechanosensing in general, which is critical

in stem cell niche commitment, vascular biology and development and are instrumental for

understanding Piezo1 channel-related disease such as lymphoedema, anaemia and malaria.

METHODS

Modeling the Piezo1 trimer. Structural data were obtained from the cryo-EM structure PDB:

6B3R[12]. Missing residues were added with MODELLER (v 9.19)[26] and the loop refinement tool[27]

was used to remove a knot in one chain between residues 2066-2074. The best loop was selected out

of 10 candidates according to the discrete optimized protein energy method[28]. The final Piezo1 model

does not comprises the first 576 residues because they are not present in the template and residues

718-781, 1366-1492, 1579-1654, 1808-1951 which are exposed to the cytosol and predicted as

unstructured. Therefore, each chain is composed by five non-overlapping fragments: residues 577-

717, 782-1365, 1493-1578, 1655-1807 and 1952-2547. The Piezo1 model obtained was further

energy minimised in vacuum with GROMACS 2016[29] prior simulations.

Coarse-grained simulations. Prior atomistic simulations the Piezo1 model obtained as described above was converted to a CG resolution and energy minimised. The coarse-grained molecular dynamics (CG-MD) simulation was essential to equilibrate the lipid bilayer around the protein and reconstitute the Piezo1 membrane indentation. The CG-MD simulations were performed using the Martini 2.2 force field[30] and GROMACS 2016[29]. To model the protein secondary and tertiary structure an elastic network model with a cut-off distance of 7 Å was used. The elastic network restricts any major conformational change within the protein during the CG-MD simulations. For the equilibration simulation Piezo1 model was inserted in a complex asymmetric bilayer using the INSert membrANE tool[31]. Three independent system assembly steps and CG equilibrations were carried out, the first from both were used in this study. The composition of the model bilayer is the following: For the outer leaflet, 1-palmitoyl-2-oleyl-phosphtidylcholine (POPC) 55%, sphingomyelin (SM) 5%, 1-palmitoyl-2-oleyl-phosphtidylethanolamine (POPE) 20% and cholesterol 20%. For the inner leaflet, POPC 50%, POPE 20%, 1-palmitoyl-2-oleyl-phosphtidylserine (POPS) 5%, cholesterol 20% and phosphatidylinositol 4,5-bisphosphate (PIP_2) 5%. The system was neutralised with a 150 mM concentration of NaCl. The model was further energy minimised and subsequently equilibrated for 500 ns with the protein particles restrained (1000 kJ $\cdot$ mol^{-1} $\cdot$ nm^{-2}) to allow the membrane bilayer to equilibrate around the model. The equilibration was performed at 323 K, with protein, lipids and solvent separately coupled to an external bath using the v-rescale thermostat [32] (coupling constant of 1.0). The temperature of 323 K is above the transition temperatures of all lipid species in the systems, therefore avoiding the lipids to undergo phase transitions to the gel phase. Pressure was maintained at 1 bar (coupling constant of 1.0) with semi-isotropic conditions and compressibility of 3×10^{-6} using the Berendsen barostat [33]. Lennard-Jones and Coulombic interactions were shifted to zero between 9 and 12 Å, and between 0 and 12 Å, respectively.

Atomistic simulations. After the CG-MD equilibration the system was carefully checked. Two molecules of POPC were removed because the head-group was trapped between Piezo1 transmembrane bundles. Moreover, in order to restore membrane asymmetry, two molecules of PIP_2 were removed because they flipped to the outer leaflet. The system obtained was minimised and further converted to atomistic resolution as described in [34]. A concentration of 3.0 mM of Ca^{2+} (i.e. 50 ions) was added to the box and neutrality restored with counterions. The obtained system was energy minimised and subsequently equilibrated in four NPT ensemble runs of 20000 steps each with an increasing time-step from 0.2 fs to 2 fs and the protein particles restrained (1000 kJ $\cdot$ mol^{-1} $\cdot$ nm^{-2}). A further equilibration step of 5 ns with a time-step of 1 fs and Cα atoms restrained (1000 kJ $\cdot$ mol^{-1} $\cdot$ nm^{-2}) was performed in order to relax the Piezo1 model embedded in a highly curved bilayer (i.e.

the membrane indentation). Stretch-induced conformational changes in the Piezo1 model were
investigated by NPT ensemble unrestrained simulations of 50 ns were the bilayer plane (xy-plane)
pressure was varied semiisotropically between -40, -30, -20, -5 and +1 bar, whereas the pressure in
bilayer normal (z) direction kept at +1 bar. The +1 bar and the -30 bar systems were further extended
to 100 ns. A similar approach revealed insightful for investigate structural transitions for the TREK-
2 mechanosensitive channel[16]. All the atomistic systems were simulated using GROMACS 2016[29]
with CHARMM36 force field[34] and a 2 fs time step. A Berendsen semi-isotropic pressure coupling[33]
at 1 bar was used during all the equilibration phases. The Parrinello-Rahman barostat[36] was further
used for the stretch-induced simulations. All simulations were performed at 323 K with protein, lipids
and solvent coupled to an external bath using the v-rescale thermostat. Long-range electrostatics were
managed using the particle-mesh Ewald method[37]. The LINCS algorithm was used to constrain bond
lengths[38].
378
Analysis and computer graphics
To calculate the area of the box the tool gmx energy from GROMACS 2016[29] was used to extract,
for each frame, the x-side and the y-side of the simulated box. The area was subsequently calculated
by multiplying these values as x-side × y-side. The root mean-square deviation was calculated on the
protein Cα using the gmx rms tool from GROMACS 2016[29]. For each system the initial coordinates
were considered as a reference structure. For the system at -40 bar, the error is the standard deviation.
The A_{proj} was calculated as follow: The program visual molecular dynamics 1.9.3 (VMD)[39]
(http://www.ks.uiuc.edu/Research/vmd/) was used to calculate distances between the residue Ala641,
located within the first bundle embedded in the bilayer, from each Piezo1 chain and for each
trajectory. Values were then averaged by chain with the standard deviation as error. The average
distance calculated was then used to calculate the radius r of the circumscribed circle using an in-
house script dividing the average distance (i.e. the side of a regular triangle) by square root 3. The
area of the circumference was then calculated as $\pi \times r^2$. In every step the standard deviation was
calculated. The average partial density was calculated using the GROMACS 2016[29] tool gmx density.
The tilting angle relative to the bilayer normal and distance from the Piezo1 pore of the Piezo1 beams,
pore-lining helices 38 and 37, were calculated using gmx bundle and gmx distance tools from
GROMACS 2016[29]. For all calculations, the Piezo1 pore region comprised the same residues that
were used for the calculation of the cavity (see below). The tool trj_cavity[40]
(https://sourceforge.net/projects/trjcavity/) was used for the calculation of the volume of the Piezo1
pore cavity as well as for the water molecules that overlap to it. The considered residues were 2450-
2547, 2214-2224 and 2326-2334. The trajectories were fitted on the Cα of the considered residues

before the calculation and options -dim 4 (i.e. the degree of how buried they were) and -spacing 1.4

(i.e. the size of the grid voxel in Å) were used. Salt bridges were calculated with VMD 1.9.3[39] using

a 6 Å cut-off and, when indicated (see Supplemental Table S2), without considering the first 30 ns of

simulation. For the calculation of the APL the tool GridMAT-MD[41], version 2.0

http://www.bevanlab.biochem.vt.edu/GridMAT-MD/) was used. The lateral pressure P_L was

calculated as explained for the TREK-2 mechanosensitive channel[16]. Molecular graphics were done

using VMD 1.9.3[39]. Data were plotted using Grace (http://plasma-gate.weiz-mann.ac.il/Grace/).

REFERENCES

1.   Wu, J., Lewis, A. H. & Grandl, J. Touch, Tension, and Transduction – The Function and

Regulation of Piezo Ion Channels. *Trends Biochem. Sci.* **42**, 57–71 (2017).

2.   Beech, D. J. & Kalli, A. C. Force Sensing by Piezo Channels in Cardiovascular Health and

Disease. *Arterioscler. Thromb. Vasc. Biol.* ATVBAHA119313348 (2019).

doi:10.1161/ATVBAHA.119.313348

3.   Li, J. *et al.* Piezo1 integration of vascular architecture with physiological force. *Nature* **515**,

279–282 (2014).

4.   Fotiou, E. *et al.* Novel mutations in PIEZO1 cause an autosomal recessive generalized

lymphatic dysplasia with non-immune hydrops fetalis. *Nat. Commun.* **6**, 8085 (2015).

5.   Andolfo, I. *et al.* Multiple clinical forms of dehydrated hereditary stomatocytosis arise from

mutations in PIEZO1. *Blood* **121**, 3925–3935 (2013).

6.   Ma, S. *et al.* Common PIEZO1 Allele in African Populations Causes RBC Dehydration and

Attenuates Plasmodium Infection. *Cell* **173**, 443-455.e12 (2018).

7.   Teng, J., Loukin, S., Anishkin, A. & Kung, C. The force-from-lipid (FFL) principle of

mechanosensitivity, at large and in elements. *Pflugers Arch.* **467**, 27–37 (2015).

8.   Martinac, B., Adler, J. & Kung, C. Mechanosensitive ion channels of E. coli activated by

amphipaths. *Nature* **348**, 261–3 (1990).

9.   Cox, C. D. *et al.* Removal of the mechanoprotective influence of the cytoskeleton reveals

PIEZO1 is gated by bilayer tension. *Nat. Commun.* **7**, 10366 (2016).

10. Poole, K., Moroni, M. & Lewin, G. R. Sensory mechanotransduction at membrane-matrix

interfaces. *Pflügers Arch. - Eur. J. Physiol.* **467**, 121–132 (2015).

11. Prager-Khoutorsky, M., Khoutorsky, A. & Bourque, C. W. Unique Interweaved Microtubule

Scaffold Mediates Osmosensory Transduction via Physical Interaction with TRPV1. *Neuron*

83, 866–878 (2014).

12. Guo, Y. R. & MacKinnon, R. Structure-based membrane dome mechanism for Piezo

mechanosensitivity. *Elife* **6**, e33660 (2017).

13.   Lin, Y.-C. *et al.* Force-induced conformational changes in PIEZO1. *Nature* 1–5 (2019).

doi:10.1038/s41586-019-1499-2

14.   Wang, L. *et al.* Structure and mechanogating of the mammalian tactile channel PIEZO2.

Nature 1–5 (2019). doi:10.1038/s41586-019-1505-8

15.   Murphy, E. J., Joseph, L., Stephens, R. & Horrocks, L. A. Phospholipid composition of

cultured human endothelial cells. *Lipids* **27**, 150–153 (1992).

16.   Aryal, P. *et al.* Bilayer-Mediated Structural Transitions Control Mechanosensitivity of the

TREK-2 K2P Channel. *Structure* **25**, 708-718.e2 (2017).

17.   Botello-Smith, W. M. *et al.* A mechanism for the activation of the mechanosensitive Piezo1

channel by the small molecule Yoda1. *Nat. Commun.* **10**, 4503 (2019).

18.   Wu, J. *et al.* Inactivation of Mechanically Activated Piezo1 Ion Channels Is Determined by

the C-Terminal Extracellular Domain and the Inner Pore Helix. *Cell Rep.* **21**, 2357–2366

(2017).

19.   Coste, B. *et al.* Piezo1 and Piezo2 are essential components of distinct mechanically

activated cation channels. *Science* **330**, 55–60 (2010).

20.   Zheng, W., Gracheva, E. O. & Bagriantsev, S. N. A hydrophobic gate in the inner pore helix

is the major determinant of inactivation in mechanosensitive Piezo channels. *Elife* **8**, e44003

(2019).

21.   Borbiro, I., Badheka, D. & Rohacs, T. Activation of TRPV1 channels inhibits

mechanosensitive piezo channel activity by depleting membrane phosphoinositides. *Sci.*

Signal. **8**, ra15 (2015).

22.   Romero, L. O. *et al.* Dietary fatty acids fine-tune Piezo1 mechanical response. *Nat. Commun.*

10, 1200 (2019).

23.   Pliotas, C. & Naismith, J. H. Spectator no more, the role of the membrane in regulating ion

channel function. *Curr. Opin. Struct. Biol.* **45**, 59–66 (2017).

24.   Coste, B. *et al.* Piezo1 ion channel pore properties are dictated by C-terminal region. *Nat.*

Commun. **6**, 7223 (2015).

25.   Kefauver, J. M. *et al.* Structure of the mechanically activated ion channel Piezo1. *Nature* **554**,

481–486 (2017).

METHODS ONLY REFERENCES:

26.   Šali, A. & Blundell, T. L. Comparative protein modelling by satisfaction of spatial restraints.

J. Mol. Biol. **234**, 779–815 (1993).

27.   Fiser, A., Do, R. K. & Sali, A. Modeling of loops in protein structures. *Protein Sci.* **9**, 1753–
469   73 (2000).

470   28.   Shen, M.-Y. & Sali, A. Statistical potential for assessment and prediction of protein
471   structures. *Protein Sci.* **15**, 2507–24 (2006).

472   29.   Abraham, M. J. *et al.* GROMACS: High performance molecular simulations through multi-
473   level parallelism from laptops to supercomputers. *SoftwareX* **1–2**, 19–25 (2015).

474   30.   De Jong, D. H. *et al.* Improved parameters for the martini coarse-grained protein force field.
475   *J. Chem. Theory Comput.* **9**, 687–697 (2013).

476   31.   Wassenaar, T. A., Ingólfsson, H. I., Böckmann, R. A., Tieleman, D. P. & Marrink, S. J.
477   Computational Lipidomics with *insane* : A Versatile Tool for Generating Custom Membranes
478   for Molecular Simulations. *J. Chem. Theory Comput.* **11**, 2144–2155 (2015).

479   32.   Bussi, G., Donadio, D. & Parrinello, M. Canonical sampling through velocity rescaling. *J.*
480   *Chem. Phys.* **126**, 14101 (2007).

481   33.   Berendsen, H. J. C., Postma, J. P. M., van Gunsteren, W. F., DiNola, A. & Haak, J. R.
482   Molecular dynamics with coupling to an external bath. *J. Chem. Phys* **81**, 3684–3690 (1984).

483   34.   Wassenaar, T. A., Pluhackova, K., Böckmann, R. A., Marrink, S. J. & Tieleman, D. P. Going
484   backward: A flexible geometric approach to reverse transformation from coarse grained to
485   atomistic models. *J. Chem. Theory Comput.* **10**, 676–690 (2014).

486   35.   Lee, S. *et al.* CHARMM36 united atom chain model for lipids and surfactants. *J. Phys.*
487   *Chem. B* **118**, 547–556 (2014).

488   36.   Parrinello, M. & Rahman, A. Polymorphic transitions in single crystals: A new molecular
489   dynamics method. *J. Appl. Phys.* **52**, 7182–7190 (1981).

490   37.   Darden, T., York, D. & Pedersen, L. Particle mesh Ewald: An N·log(N) method for Ewald
491   sums in large systems. *J. Chem. Phys.* **98**, 10089–10092 (1993).

492   38.   Hess, B., Bekker, H., Berendsen, H. J. C. & Fraaije, J. G. E. M. LINCS: A linear constraint
493   solver for molecular simulations. *J. Comput. Chem.* **18**, 1463–1472 (1997).

494   39.   Humphrey, W., Dalke, A. & Schulten, K. VMD: visual molecular dynamics. *J. Mol. Graph.*
495   **14**, 33–8, 27–8 (1996).

496   40.   Paramo, T., East, A., Garzón, D., Ulmschneider, M. B. & Bond, P. J. Efficient
497   characterization of protein cavities within molecular simulation trajectories: trj_cavity. *J.*
498   *Chem. Theory Comput.* **10**, 2151–2164 (2014).

499   41.   Allen, W. J., Lemkul, J. A. & Bevan, D. R. GridMAT-MD: A grid-based membrane analysis
500   tool for use with molecular dynamics. *J. Comput. Chem.* **30**, 1952–1958 (2009).

ACKNOWLEDGEMENTS

504 ACK and DDV are funded by an Academy of Medical Sciences and Wellcome Trust Springboard
505 Award. DJB is funded by a Wellcome Trust Investigator Award.

COMPETING INTERESTS

508 We have no competing interests to report.

MATERIAL AND CORRESPONDANCE

511 Any requests can be sent to Antreas C Kalli.

AUTHOR CONTRIBUTION

514 D.DV. Built the model, performed simulations and analysed the data. The research was
515 conceptualised by A.C.K. and D.J.B., A.C.K and D.J.B. acquired funding and supervised the project.
516 D.DV. Wrote the first draft. All authors revised the first draft.

ADDITIONAL INFORMATION

519 A supplemental figure (Supplemental Fig. S1) and two tables of additional data (Supplemental Table
520 S1 and Supplemental Table S2) accompany this paper.

DATA AVAILABILITY

523 Data will be available a doi at the University of Leeds

Supplemental Figure S1. Correlation of Piezo1 pore-lining helices with changes in Piezo1 conformation. For each system, the applied negative pressure in each simulation is indicated with different colours: black, 1 bar; red, -5 bar; green, -20 bar; blue, -30 bar; orange, -40 bar. (**a**) Correlation between the root mean-square deviation (RMSD) of the Piezo1 blades and the tilt angle relative to the bilayer normal of the beam helix.(**b**) Tilt angle relative to the bilayer normal for the TM38 helix. (**c**) Correlation between the RMSD of the Piezo1 blades and the distance from the centre of mass of the TM38 helix and the Piezo1 pore. (**d**) Correlation between the tilt angle relative to the bilayer normal of the TM38 and TM37 helices. (**e**) Correlation between the RMSD of the Piezo1 blades and the distance from the respective centre of mass of the TM37 helix and the Piezo1 pore.

Supplemental Table S1. Area per lipid during simulations.

	+1 bar		-5 bar		-20 bar		-30 bar		-40 bar	
	up	*lw*	*up*	*lw*	*up*	*lw*	*up*	*lw*	*up ± sd*	*lw ± sd*
0 ns	40.19	64.92	40.19	64.92	40.19	64.92	40.19	64.92	40.19 ± 0	64.92 ± 0
10 ns	40.21	63.16	43.72	68.32	63.06	68.57	77.10	78.13	86.83 ± 2.58	92.8 ± 1.87
20 ns	37.21	74.51	45.26	64.95	66.61	65.83	77.66	76.03	90.09 ± 0.28	88.88 ± 0.26
30 ns	38.30	69.36	48.39	59.68	67.62	64.85	78.80	76.02	90.18 ± 0.66	89 ± 0.94
40 ns	42.52	57.29	51.91	55.94	66.60	64.88	78.30	76.19	90.19 ± 0.14	89.16 ± 0.08
50 ns	40.04	65.08	53.24	53.57	66.86	64.18	78.52	76.09	90.09 ± 0.84	89.46 ± 0.96
60 ns	40.60	63.62					78.05	75.78		
70 ns	38.29	72.12					77.72	75.99		
80 ns	39.45	63.92					78.30	76.39		
90 ns	41.68	61.33					78.38	76.22		
100 ns	40.77	62.91					77.75	76.31		

Values are in Å^2. *up*, upper leaflet; *lw*, lower leaflet; *sd*, standard deviation.

Supplemental Table S2. Predicted salt bridge between the elbow and the pore-lining helix 38

	+1 bar	-5 bar	-20 bar	-30 bar	-40 bar (rep1)	-40 bar (rep 2)
E2133.a - R2482.c	100	100	100	100	100	100
E2133.a - R2482.b	98	98	100	96	70 (98)	100 (100)
E2133.b - R2482.c	100	100	100	7 (62)	8 (89)	7 (83)

When occurrence was lower than 70%, numbers between parentheses indicate occurrence within the first 30 ns of simulations.