Non-selective response inhibition in Go/NoGo task: Bayesian analysis of fMRI data

Ruslan Masharipov, Maxim Kireev*, Alexander Korotkov, and Svyatoslav Medvedev

1 - N.P. Bechtereva Institute of the Human Brain, Russian Academy of Sciences, Academika Pavlova Street 9, St. Petersburg, 197376, Russia

Corresponding author: Maxim Kireev;
Address: Academika Pavlova Street 9, St. Petersburg, 197376, Russia
E-mail: kireev@ihb.spb.ru
Phone number: +7 812 670-09-51
Fax number: +7 812 2343247

Abstract

Response inhibition is typically considered a brain mechanism selectively triggered by particular "inhibitory" stimuli or events. Based on recent research, an alternative non-selective mechanism was proposed by several authors. Presumably, the inhibitory brain activity may be triggered not only by the presentation of "inhibitory" stimuli but also by any imperative stimuli, including Go stimuli, when the context is uncertain. Earlier support for this notion was mainly based on the absence of a significant difference between neural activity evoked by equiprobable Go and NoGo stimuli. Previous neuroimaging studies utilized classical null hypothesis significance testing, making it impossible to accept the null hypothesis. Therefore, the current research aimed to provide evidence for practical equivalence of neuronal activity in Go and NoGo trials using Bayesian analysis of functional magnetic resonance imaging (fMRI) data. Thirty-four healthy participants performed a cued Go/NoGo task with an equiprobable presentation of Go and NoGo stimuli. To reveal brain areas previously associated with response inhibition in similar experimental conditions, we performed a meta-analysis of fMRI studies using equal probability Go/NoGo tasks. As a result, we observed overlap between response inhibition areas and areas demonstrating the practical equivalence of neuronal activity located in the right dorsolateral prefrontal cortex, parietal cortex, premotor cortex, and left inferior frontal gyrus. Thus, obtained results favour the existence of non-selective response inhibition, which can act in settings of context uncertainty, modelled by the equal probability of Go and NoGo stimuli.

Keywords: non-selective response inhibition, functional MRI, Bayesian inference, accepting the null hypothesis, practical equivalence, Go/NoGo task

Highlights

- Non-selective response inhibition was assessed by equiprobable Go/NoGo task
- Bayesian analysis of fMRI data was combined with a meta-analysis of fMRI studies
- Nodes of response inhibition system were equally involved in Go and NoGo trials

•   Evidence for non-selective response inhibition in uncertainty context was found

1. Introduction

Response inhibition is the ability to suppress inappropriate, automatic, reflexive, or

habitual prepotent responses to produce a controlled goal-directed response (Friedman et al.,

2008, Isoda & Hikosaka, 2011). It is generally accepted in the literature that response inhibition

works in close relation to other processes associated with cognitive control, such as working

memory, voluntary attention, conflict monitoring, and action selection (Buchsbaum et al., 2005,

Nee, Wager & Jonides, 2007, Simmonds, Pekar & Mostofsky, 2008, Criaud & Boulinguez,

2013, Cieslik et al., 2015). Moreover, response inhibition is thought to represent a multifaceted

phenomenon rather than a unitary brain mechanism. A distinction is made between action

cancellation and action restraint (Swick, Ashley & Turken, 2011, Cieslik et al., 2015, Zhang,

Geng & Lee, 2017) as well as between reactive and proactive response inhibition mechanisms

(Jaffard et al., 2007, 2008, Aron, 2011, Braver, 2012, Erika-Florence, Leech & Hampshire,

2013).

According to the conventional view, the response inhibition process is selectively triggered

by "inhibitory" stimuli that result in increased neuronal activity in brain structures responsible

for inhibitory control (Verbruggen & Logan, 2008, Verbruggen et al., 2014, Logan et al., 2014,

Criaud et al., 2017). However, in several cases, the concept of selective response inhibition fails

to explain observed behavioural and neurophysiological phenomena. Manipulating the

probability of the occurrence of "inhibitory" stimuli and the subjects' awareness of the

probability of the appearance may slow down the motor response (Chevrier, Noseworthy &

Schachar, 2007, Jaffard et al., 2007, Boulinguez et al., 2008, Vink et al., 2014, Vink et al., 2015,

Dunovan et al., 2015, Meffert et al., 2016, Hsieh, Wu & Tang, 2016). Moreover, when necessary

to rapidly suppress a specific action, the action is inhibited along with all other potential actions.

Therefore such inhibition can affect the entire motor system (Coxon, Stinear & Byblow, 2007,

Aron & Verbruggen, 2008, Badry et al., 2009, Duque & Ivry, 2009, Duque et al., 2010, Majid et

al., 2012, MacDonald et al., 2017). In attempting to explain the effects mentioned above, several

authors proposed the concept of non-selective ("global") response inhibition (Frank, 2006, Aron,

2011, Albares et al., 2014, Criaud et al., 2017). It is thought that those non-selective mechanisms

serve to prevent inappropriate or premature responses at the expense of the speed of execution of

correct actions. First, response inhibition mechanisms may non-selectively inhibit all potential

responses to further selectively execute an appropriate response (non-selective inhibition of

multiple concurrent motor responses). Second, inhibition may be triggered not only by the

presentation of "inhibitory" stimuli but also by the occurrence of any imperative stimuli

instructing on the necessity to suppress or execute a prepared action (non-selectivity of inhibitory stimulus perception). In the present work, we consider the latter mechanism.

A tentative neurophysiological model of the non-selective or "global" response inhibition involved in the resolution of interference between several competing response options was proposed by Frank (2006) and included the cortico-subthalamic "hyper-direct" pathway (Nambu et al., 1996, Haynes & Haber, 2013) which is capable of rapidly and non-selectively suppress all potential response options. As the research area developed, it was hypothesized that the model might be applied not only to tasks with multiple concurrent response options but also to simpler tasks where the subject has to choose between executing and refraining from an action (Albares et al., 2014, Criaud et al., 2017). The authors used a cued Go/NoGo task, wherein a preparatory cue stimulus indicated the probability of a NoGo stimulus occurrence. Within the framework of the selective inhibition model, the inhibition process would only be triggered by the identification of the NoGo stimulus. According to the hypothesis of non-selective inhibition, when the context is uncertain (equal probability of NoGo and Go stimuli), the need for response inhibition arises for both NoGo and Go trials. Experimental assessment of the hypotheses revealed no statistically significant differences between NoGo and Go trials both in the amplitude of early components of event-related potentials (ERP) (Albares et al., 2014) and in the level of neuronal activity measured by functional magnetic resonance imaging (fMRI) (Criaud et al., 2017). The authors considered this fact as evidence favouring the presence of a "non-selective" inhibitory mechanism that may be engaged when the context is uncertain. The analysis of early ERP components suggests that non-selective inhibition may work as a gating mechanism acting on the initiation of the prepared action.

However, a critical limitation of the abovementioned studies was that the authors could not accept the null hypothesis, so they did not provide direct proof of the practical equivalence of neuronal activity level between Go and NoGo trials in the inhibition-related brain structures. Indeed, within the framework of classical (frequentist) null hypothesis significance testing (NHST), we cannot accept the null hypothesis based on the absence of a significant difference. We can only reject it (Kruschke & Liddell, 2015). Thus, the question of experimental support of non-selective response inhibition remains unanswered, and answering this question requires overcoming the methodological limitation of NHST, which is possible using Bayesian statistics (Kruschke & Liddell, 2015).

Therefore, the present study aimed to verify the non-selective response inhibition hypothesis by using the fMRI data obtained for an equiprobable Go/NoGo task. Based on the results of prior studies, it may be suggested that if the hypothesis on the non-selectivity of inhibition is correct, then the brain structures responsible for response inhibition will

demonstrate practically equivalent levels of neuronal activity in equiprobable Go and NoGo trials. To assess this prediction, the Bayesian parameter inference was applied. Instead, if the hypothesis on the selectivity of response inhibition is correct, then activation of the inhibition-related brain structures will be observed in the NoGo conditions compared with Go conditions.

2. Material and methods

2.1. A meta-analysis of fMRI studies using equal probability Go/NoGo tasks.

Given that a practically equivalent level of neuronal activity may be identified not only for response inhibition-related structures but, for example, those related to sensory processing of visual stimuli, working memory, and attention, we conducted a meta-analysis of fMRI studies aimed at revealing brain structures associated with response inhibition. Studies using equal probability Go/NoGo tasks comparable to the present study were selected for the meta-analysis. We searched for studies that, similar to the investigation by Criaud et al. (2017), compared the neuronal activity in the condition of equiprobable Go and NoGo stimuli presentation with the control Go condition, wherein the subject did not need to inhibit the prepared action.

We searched for these studies in the PubMed database in the period of 01/01/2000 to 15/07/2019 using the following keywords: "((fmri) OR (functional magnetic resonance)) AND ((nogo) OR (no-go))." Four additional studies were identified through manual searches. As a result, 726 papers were identified. At the first stage of selection, we excluded reviews, meta-analyses, and papers repeatedly reporting the results of fMRI studies (see the flow chart of analysis at Fig. 1).

Figure 1. Flow chart of study selection in the meta-analysis.

Studies dealing with Flanker task modifications of Go/NoGo task or based on unequal

probability of appearance of Go and NoGo trials were also excluded from the analysis. As a

result, 593 papers were excluded. Task designs of the remaining 113 papers provided for equal

probability Go and NoGo stimuli presentation. At the next step, 82 of the papers were excluded

based on the following criteria: auditory and sensorimotor Go/NoGo tasks were used; only

emotion-laden task conditions were used (emotionally neutral conditions were either absent or

not considered separately); the necessary control Go-condition (100% probability of the Go-

stimulus presentation) was not used; the coordinates for the contrast of interest "50/50%

Go/NoGo > 100% Go-control" within a group of healthy subjects were not reported. In 11 out of

the 31 remaining studies, healthy volunteer subjects under the age of 18 (children and

adolescents) were studied. These papers were also excluded from analysis because this study

focused on the brain activity of healthy, adult subjects. In all remaining studies except one, only

block designs were used. To make our sample more homogenous, we excluded the only eligible study with an event-related design (Criaud et al., 2017). The final meta-analysis included 20 studies (452 healthy subjects, mean age of 29 years, 210 foci) (for more details see supplementary materials, Table S1).

The meta-analysis was conducted using GingerALE 3.0.2 software (http://brainmap.org/ale) with a cluster-forming threshold of 0.001 (uncorrected) and a cluster-level threshold of 0.05 corrected for family-wise error (FWE) (Eickhoff et al., 2016). All the coordinates were converted into Montreal Neurological Institute (MNI) space. The median full width at half maximum (FWHM) value of Gaussian function was set at a level of 13.5 mm.

2.2. Power analysis.

To justify the sample size, we estimated the effect sizes typical of selective inhibition and performed an a priori power analysis. We used Consortium for Neuropsychiatric Phenomic LA5c dataset (Poldrack et al., 2016, Gorgolewski et al., 2017) as it contains fMRI data from a relatively large cohort (N = 115) of healthy subjects performing Stop-Signal Task (see supplementary materials for details). This classical inhibitory paradigm models a situation where the inhibitory brain activity selectively triggered by infrequent inhibitory stimuli (stop signal) but not by Go stimuli (Criaud et al., 2017). The mean effect sizes for the selective inhibition contrast ("Correct-Stop > Go" contrast, see Fig. S1) were estimated within the brain regions revealed by the current meta-analysis. They ranged from 0.57 to 1.12 Cohen's d. A priori power analysis using GPower 3.1.9.7 (Faul et al., 2007) indicated that the sample size of 34 subjects would be sufficient to detect selective inhibition effect (d = 0.57, two-tailed one-sample test, alpha = 0.05) with the power of 0.9 (see Fig. S2). The power of 0.8 or 0.9 is commonly used for sample size calculations (Dell, Holleran & Ramakrishnan, 2002). However, obtaining non-significant results with high power would not provide direct evidence for the null hypothesis of non-selective inhibition in settings of context uncertainty (Amrhein, Korner-Nievergelt & Roth, 2017). Bayesian inference or frequentist equivalence testing should be used to provide evidence for the null hypothesis (Kruschke & Liddel, 2015, Lakens, 2017, Kruschke, 2018).

2.3. Subjects.

Thirty-four healthy, right-handed volunteer subjects (24 women, aged (mean ± SD): 25.9 ± 5.2) participated in the present fMRI study. An Oldfield test was used to determine the dominant arm (Oldfield, 1971). The subject volunteers signed written informed consent to participate in the study and were paid for their participation. All the procedures were performed in accordance with the Declaration of Helsinki and were approved by the Ethics Committee of the N.P. Bechtereva Institute of the Human brain of the Russian Academy of Sciences.

2.4. Experimental task and study procedure.

We used a paired stimulus modification of the Go/NoGo task (see Fig. 2) (Kropotov,
2016). This task was originally developed to dissociate cognitive processes, such as response
inhibition, conflict monitoring, sensory mismatch and category discrimination (Kropotov et al.,
2011, 2015). Given that imperative Go and NoGo stimuli were presented in an equiprobable
manner, previously collected fMRI data were used for the purposes of the current study. Each
trial consisted of the two consequently presented stimuli. The first preparatory cue stimulus
warned the subjects on the presentation of the second, imperative stimulus, or indicated no need
for any response to the second stimulus. The study included two variants of the experiment's
instructions. According to the first instruction (see Fig. 2A), the subject should press the
response button upon presentation of the pair of images "animal-animal" ("A-A Go" trials) and
refrain from acting upon presentation of the pair "animal-plant" ("A-P NoGo" trials). According
to the second experiment's instruction (see Fig. 2B), the subject acts after the presentation of the
pair "animal-plant" ("A-P Go" trials) and suppresses an action upon presentation of the pair
"animal-animal" ("A-A NoGo" trials).

Figure 2. Experimental design of the Go/NoGo task. S1 – first stimulus (preparatory cue). S2
– second stimulus (imperative stimulus). A – images of animals, P – images of plants. A) First
experiment: "A-A Go", "A-P NoGo". B) Second experiment: "A-P Go", "A-A NoGo". Red
boxes highlight trials that were compared to test the hypothesis of selective and non-selective
response inhibition.

In both experiments, if the first presented stimulus was an image of a "plant," subjects do
not need to perform any actions in response to the presentation of any second stimuli of a trial. In
such a trial, the subject should ignore the second stimulus and wait for the next pair of stimuli
("P-A Ignore" and "P-P Ignore" trials). Accordingly, it was assumed that there is no inhibition of
prepared action in "Ignore" conditions. The "Ignore" trials were included in the adopted

Go/NoGo task to evaluate preparatory brain activity, such as preparing to receive a relevant stimulus (attentional set) and preparing to make a movement (motor set), by comparing them to Go and NoGo trials. However, the issue related to preparatory processes is beyond the scope of the present study. Fifty pairs of each type of stimuli were randomly presented in each experiment. The order of following the instructions was counterbalanced among the subjects. Two variants of the present Go/NoGo task made it possible to control the differences in the load on working memory between A-P/A-A - Go/NoGo stimuli (Kropotov & Ponomarev, 2015).

The fMRI data obtained using this experimental task allowed us to test the hypothesis on the *non-selectivity* of response inhibition in the current study given that a similar task design was previously used for the same goal by Criaud et al. (2017). The imperative Go and NoGo stimuli were presented after the preparatory stimulus with equal probability, as reported in event-related studies by (Albares et al., 2014, Di Russo et al., 2016, Kropotov, 2016, Criaud et al., 2017). Such an equally probable presentation of imperative stimuli provides several advantages. First, it minimizes the difference in cognitive load between Go and NoGo conditions arising from task complexity (Criaud & Boulinguez, 2013, Criaud et al., 2017). Second, such a design enables the exclusion of effects associated with a low frequency of the NoGo stimuli presentation ("oddball" effects) confounding the effect of inhibition (Di Russo et al., 2016). Third, it creates maximum uncertainty regarding the probability of the presentation of an imperative stimulus and thus minimizes the conflict between two response types, which makes it possible to distinguish between error monitoring and conflict resolution processes on the one hand and response inhibition processes on the other (Lavric, Pizzagalli & Forstmeier, 2004).

In total, 100 NoGo trials, 100 Go trials, 100 P-A Ignore trials, and 100 P-P Ignore trials were presented over two experimental sessions. In the absence of stimulation, a fixation cross was displayed in the centre of the screen. The stimuli were presented for 100 ms, and the interstimulus interval was 1000 ms. The intertrial interval jittered from 2800 to 3200 ms with an increment step of 100 ms. Additionally, to improve design efficiency, 100 zero-events (fixation crosses) were randomly inserted between the stimuli pairs (trials), and their duration jittered from 3000 to 5000 ms with an increment size of 500 ms. The action to be performed consisted of pressing a button with the right thumb. The duration of one task session was 17.5 minutes. Before starting the fMRI study, the subjects performed a training task. The Invivo's Eloquence fMRI System (Invivo, Orlando, FL, USA) was used to deliver the stimuli, synchronize with fMRI acquisition, and record reaction times of subjects bottom pressing. The task presentation sequence and all temporal parameters of the stimuli presentation were programmed using the E-prime 2.0 software package (Psychology Software Tools Inc., Pittsburgh, PA, USA).

2.5. Image acquisition.

A Philips Achieva 3.0 Tesla scanner (Philips Medical Systems, Best, Netherlands) was
used for the study. The structural T1-images were registered with the following parameters: field
of view (FOV) – 240×240 mm, repetition time (TR) – 25 ms, echo time (TE) – 2.2 ms, 130 axial
slices with a thickness of 1 mm and pixel size of 1×1 mm, flip angle – 30°. For the registration of
T2*-images, a single-pulse echo planar imaging (EPI) sequence was used. The period of data
registration from 31 axial slices was 2 seconds (TR = 2 s, TE = 35 ms). The following
parameters were employed: FOV – 200×186 mm, flip angle – 90°, voxel size – 3×3×3 mm. Two
dummy scans were performed prior to each session. To minimize head motions, we used an MR-
compatible soft cervical collar.

2.6. Preprocessing of fMRI-images.

Image preprocessing included the following: realignment to the first image of the session,
slice time correction, co-registration, segmentation, normalization to an MNI template, and
spatial smoothing (8 mm FWHM). Preprocessing and statistical analysis of the images were
performed using an SPM12 (Statistical parametric mapping) software package
(http://www.fil.ion.ucl.ac.uk/spm).

2.7. Statistical analysis of fMRI data.

The first level of analysis was conducted using frequentist parameter estimation. The
second level of analysis was conducted using both the frequentist and Bayesian parameter
estimation (Friston et al., 2002a, Friston & Penny, 2003, Neumann & Lohmann, 2003). Onset
times of second stimuli presentation (separately for "A-A Go", "A-P NoGo", "P-A Ignore
Exp1", "P-P Ignore Exp1", and also "A-P Go", "A-A NoGo", "P-A Ignore Exp2", "P-P Ignore
Exp2"), erroneous button pressing, and missing the responding in Go trials were used to create
regressors of the general linear model (GLM) for each subject. Events were impulse responses
with a duration of zero convolved with the canonical haemodynamic response function (HRF).
Low-frequency drift was removed by temporal high pass filtering with a cut-off frequency of
1/128 Hz. Six head motion parameters were introduced in the GLM as nuisance regressors to
account for the movement artefacts (Johnstone et al., 2006). Beta-coefficients reflecting an
increase in the blood oxygenation level-dependent (BOLD) signal in the experimental condition
relative to the implicit baseline were converted into percent signal change (PSC) following the
procedure recommended in (Poldrack, Mumford, Nichols, 2011, p.186). To that end, beta-
coefficients for each condition were divided by the mean value of beta-coefficients for the
constant term and multiplied by 100 and a scaling factor (SF) that is needed so that the peak of
an isolated BOLD response equals to the 1 (SF = 0.21). Linear contrast, c, with a sum of positive
terms equal to 1 was used to calculate the difference of the BOLD-signal percent changes in the
conditions under study: 0.5 · ["A-P NoGo" + "A-A NoGo"] – 0.5 · ["A-P Go" + "A-A Go"] (the

"NoGo vs. Go" comparison). The contrasts from thirty-four subjects were used as variables to verify the hypotheses on selective and non-selective response inhibition on the second level of analysis. Only grey-matter voxels were included in the second-level analysis. To that end, a mask was created based on the segmentation of each subject's structural T1-images.

2.8. Verification of the hypotheses on selective and non-selective response inhibition.

The classical NHST approach estimates the probability (p-value) of obtaining actual data's best-fitting parameter value, β, or something more extreme under the null hypothesis that the experimental effect is $\theta = c\beta = 0$. One can never accept the null hypothesis using NHST because the p-value does not represent the probability of the null hypothesis and the probability an effect equals exactly zero is itself zero (Friston et al., 2002a). A non-significant result can be caused by two conditions (Dienes, 2014): 1) there is no effect, and our data are against the alternative hypothesis; or 2) our data are insufficient to distinguish alternative hypothesis from the null hypothesis, and we cannot confidently make any inference (low statistical power). When the obtained result is not significant, it is recommended to use frequentist equivalence testing (Lakens, 2017) or Bayesian inference (Penny & Ridgway, 2013; Dienes, 2014; Kruschke & Liddel, 2015, Kruschke, 2018). In the present study, the Bayesian parameter inference was utilized (Friston & Penny, 2003; Penny & Ridgway, 2013; Kruschke, 2018).

The Bayesian approach estimates the posterior probability (PP) distribution of effect θ given the data using the likelihood and the prior knowledge. For the second-level Bayesian analysis, SPM12 implements the global shrinkage prior (Friston & Penny, 2003). It represents a prior belief that, on average, in the whole brain, there is no global experimental effect. If the posterior probability of the effect exceeding the effect size threshold, γ, is greater than the predefined probability threshold, $\alpha = 95\%$, then the hypothesis on the presence of "NoGo > Go" effect will be accepted (see Fig. S3A):

$$PP_{NoGo>Go} = p(\theta > \gamma|Data) \geq \alpha.$$

If the effect value falls within the interval [-γ; γ] with a probability of $\alpha = 0.95$, then the hypothesis of null "NoGo = Go" effect will be accepted, supporting the practical equivalence (Kruschke, 2018) of the BOLD signal between compared conditions (see Fig. S3B):

$$PP_{NoGo=Go} = p(-\gamma < \theta < \gamma|Data) \geq \alpha.$$

The hypothesis on the presence of "Go > NoGo" effect will be accepted if (see Fig. S3C):

$$PP_{Go>NoGo} = p(\theta < -\gamma|Data) \geq \alpha.$$

If none of the above criteria are satisfied, the data in particular voxel are insufficient to distinguish the null hypothesis from the alternative hypothesis ("low-confidence" voxels, see Fig. S3D) (Magerkurth et al., 2015). For such voxels, it is impossible to make any inference confidently, so the power should be increased. In the current study, our conclusions were based

only on voxels in which the effect size values (PSC) fell within or outside the interval [-γ; γ] with

the PP > 0.95. Thus, for these voxels, the obtained data are sufficient to make an inference

confidently.

Bayesian parameter inference with a zero effect size threshold corresponds to a false

discovery rate correction for multiple comparisons within the NHST framework (Genovese,

Lazar & Nichols, 2002, Friston & Penny, 2003). In the SPM12, the group-level effect size

threshold is set at one standard deviation of the prior variance of the contrast (prior SD_θ) by

default (Friston & Penny, 2003). For the contrast under consideration, prior $SD_\theta = 0.103\%$ of

BOLD-signal. The interval [-γ, γ] can be thought as the neuronal ''background noise level''

(Eickhoff et al., 2008) or as a region of practical equivalence (ROPE) that expresses which effect

size values (PSC) are equivalent to the null value for current practical purposes (Kruschke, 2018,

the "ROPE-only" decision rule, see Fig. S3). Additionally, the results of Bayesian parameter

inference were compared with the results of classical inference using a FWE-corrected voxel-

wise threshold of $p < 0.05$.

The hypothesis on the *selectivity* of response inhibition implies a selective increase in the

neuronal activity in response to the presentation of NoGo stimuli compared to Go stimuli

("NoGo > Go", PP($\theta > 0.103\%$) > 0.95). Regarding the hypothesis on the *non-selectivity* of

response inhibition, Go and NoGo trials are expected to result in practically equivalent neuronal

activity ("NoGo = Go", PP($-0.103\% < \theta < 0.103\%$) > 0.95) in the brain structures responsible for

response inhibition (which were revealed by the current meta-analysis). For visualization

purposes, posterior probabilities were converted to LogOdds = log(PP/(1-PP)) (LogOdds > 3

correspond to PP > 0.95). Anatomical localization of clusters was identified using xjView

toolbox (http://www.alivelearn.net/xjview).

3. Results

As a result of a meta-analysis of 20 fMRI studies, we identified the brain structures that

were characterized by increased neuronal activity demonstrated in the settings of equal

probability Go and NoGo stimuli presentation compared to the control Go conditions ("50/50%

Go/NoGo > 100% Go-control" contrast). In such control conditions, the response inhibition

process was not involved because the subjects were instructed always to press the button. The set

of revealed brain structures includes: (1) right dorsolateral prefrontal cortex (DLPFC), (2) right

inferior parietal lobule (IPL), (3) right temporoparietal junction (TPJ), (4) bilateral inferior

frontal gyrus (IFG) and anterior insula (also known as anterior insula/frontal operculum (AIFO)),

(5) right premotor cortex (PMC) and frontal eye field (FEF), (6) bilateral anterior cingulate

cortex (ACC) and supplementary motor area (SMA), and (7) bilateral thalamus (see Fig. 3, Table

1).

Meta-analysis

Figure 3. Results of the meta-analysis. The result of the meta-analysis of 20 fMRI studies using equal probability Go/NoGo tasks ("50/50% Go/NoGo > 100% Go-control" contrast).

Table 1. The result of the meta-analysis of 20 fMRI studies using equal probability Go/NoGo tasks ("50/50% Go/NoGo > 100% Go-control" contrast).

№	Cluster size, mm^3	Peak MNI-coordinate, mm	Peak ALE-score	Anatomical localization (L – left, R – right hemisphere; BA – Brodmann area)
1	10160	42 34 32	0.0114	R: DLPFC (middle frontal gyrus), BA 8,9
2	8368	46 18 -10	0.0138	R: IFG/Anterior insula (AIFO), BA 13, 45, 47
3	4896	-34 24 0	0.0088	L: IFG/Anterior insula (AIFO), BA 13, 44, 45, 47
4	4176	60 -44 26	0.0090	R: TPJ (supramarginal gyrus and superior temporal gyrus), BA 22, 39, 40
5	3408	5 18 40	0.0103	R/L: SMA, ACC, BA 6, 8, 24, 32
6	1664	12 -8 10	0.0089	R: Thalamus
7	1584	42 6 40	0.0076	R: PMC, FEF (precentral gyrus, middle frontal gyrus), BA 6, 8, 9
8	1456	-12 -10 10	0.0084	L: Thalamus
9	1424	30 -62 38	0.0071	R: IPL (angular gyrus), BA 7, 39

Abbreviations: DLPFC – dorsolateral prefrontal cortex, IFG – inferior frontal gyrus, AIFO – anterior insula/frontal operculum, TPJ – temporoparietal junction, SMA – supplementary motor area, ACC – anterior cingulate cortex, PMC – premotor cortex, FEF – frontal eye field, IPL – inferior parietal lobule.

Both classical NHST and Bayesian inference applied in the present fMRI study do not reveal any significant increase in neuronal activity in NoGo trials compared to Go trials. Thus, the "NoGo > Go" effect predicted by the hypothesis of selective response inhibition was not

observed. In addition, both statistical methods revealed similar activation patterns for the "Go > NoGo" effect (Dice coefficient = 0.8), demonstrating the consistency between utilized Bayesian parameter inference with $\gamma = 1$ prior SD_θ and LogOdds > 3 (PP > 0.95) thresholding approach and NHST with pFWE<0.05 (see Fig. 4, Fig. S4A and Tables S2 and S3 in the supplementary materials).

Figure 4. Results of classical voxel-wise and Bayesian parameter inference on the obtained fMRI data. A) Classical NHST inference using a FWE-corrected voxel-wise threshold of p < 0.05. B) Bayesian parameter inference using the effect size threshold $\gamma = 1$ prior $SD_\theta = 0.103\%$, LogOdds > 3 (PP > 0.95). The "NoGo > Go" effect was not detected by any of the methods. Red

colour depicts the "Go > NoGo" effect. Green colour depicts the "NoGo = Go" effect (practical equivalence of neuronal activity in Go and NoGo trials). White colour depicts A) non-significant voxels; B) "low-confidence" voxels.

Meanwhile, Bayesian analysis has made it possible to define the brain structures, wherein practical equivalence of neuronal activity was observed in Go and NoGo trials. The null "NoGo = Go" effect was revealed for a widely distributed set of regions throughout the whole brain surrounding clusters of activations revealed in the "Go > NoGo" comparison (see Fig. 4B). The "NoGo = Go" regions were separated from activation clusters by regions that consisted of "low confidence" voxels (Magerkurth et al., 2015). For "low confidence" voxels, the data obtained are insufficient to accept or reject the null hypothesis (the number of "low confidence" decreases with increasing sample size, see Fig. S4B). The "NoGo = Go" area cannot be reasonably split into several parts such as activation clusters. Therefore, we limited ourselves to specifying the anatomic localization of brain areas where the "NoGo = Go" area overlaps with the clusters revealed by the current meta-analysis.

As it can be inferred from the revealed overlap between the results of the meta-analysis and the results of abovementioned Bayesian analysis, only a few of the brain structures demonstrated both increased activity in the "50/50% Go/NoGo > 100% Go-control" meta-analytical comparison and the practical equivalence of the BOLD signal in the "NoGo vs. Go" comparison of the present fMRI data set. Overlap between both analyses was observed in (1) right DLPFC, (2) right IPL, (3) right TPJ, (4) left IFG and anterior insula (AIFO), (5) right PMC, and FEF (see Fig. 5, Table 2).

Figure 5. Overlap between meta-analysis and Bayesian meta-analysis. Overlap between null "NoGo = Go" effect regions and response inhibition brain structures revealed by the current

meta-analysis ("50/50% Go/NoGo > 100% Go-control" contrast). Abbreviations: IFG – inferior frontal gyrus, Ins – insula, DLPFC – dorsolateral prefrontal cortex, FEF – frontal eye field, PMC – premotor cortex, IPL – inferior parietal lobule, BA – Brodmann area, L – left, R – right.

Table 2. Overlap between null "NoGo = Go" effect regions revealed by Bayesian analysis of the obtained fMRI data and the results of the current meta-analysis.

№	Cluster size, mm^3	Centroid MNI-coordinates	Anatomical localization (L – left, R – right hemisphere; BA – Brodmann area)
1	1323	39 6 42	R: PMC, FEF, BA 6, 8
2	1188	30 -60 45	R: IPL
3	918	39 33 33	R: DLPFC, BA 9
4	432	-39 30 3	L: IFG
5	162	-27 27 0	L: Anterior insula
6	162	-39 15 1	R: TPJ

Abbreviations: DLPFC – dorsolateral prefrontal cortex, PMC – premotor cortex, FEF – frontal eye field, IPL – inferior parietal lobule, IFG – inferior frontal gyrus, TPJ – temporoparietal junction.

4. Discussion

The present study results confirmed the hypothesis of non-selective response inhibition in the context uncertainty associated with the equiprobable presentation of Go and NoGo stimuli. Bayesian analysis of the obtained fMRI data provides evidence of the practical equivalence of neuronal activity evoked by Go and NoGo stimuli in several brain structures. According to the results of our meta-analysis, some of these brain structures, including right DLPFC, IPL, and TPJ, left IFG (AIFO), right PMC, and FEF, were associated with response inhibition. These structures were repeatedly found activated whenever the condition of equal probability Go and NoGo stimuli presentation was compared with the control Go condition ("50/50% Go/NoGo > 100% Go control" contrast). In addition, we did not observe the "NoGo > Go" effect in the current study. This finding is consistent with the data of several previous event-related fMRI studies that did not find a significant activity increase in NoGo trials compared to Go trials under similar experimental conditions (Lee et al., 2015, Di Russo et al., 2016, Criaud et al., 2017, see also random-effects analysis within visual and audio modalities in Laurens, Kiehl & Liddle, 2004).

However, the question is how precisely observed activity can be associated with response inhibition per se in those brain regions in which accepted null hypothesis overlapped with response inhibition brain regions resulted from the meta-analysis. On the one hand, the frontoparietal structures (DLPFC, IPL, TPJ) are engaged not only in Go/NoGo tasks but also in experiments examining task switching, resolution of cognitive conflicts, working memory, and

attention focusing, i.e., external-directed activities ("multiple demand system" (Duncan, 2010); "task-general network" (Cole et al., 2014); "extrinsic mode network" (Hugdahl et al., 2015)). On the other hand, the most prominent feature of brain activity during Go/NoGo tasks compared to diverse non-inhibitory tasks is a right-dominant activity in DLPFC (meta-analyses: Buchsbaum et al., 2005, Nee, Wager & Jonides, 2007, Levy & Wagner, 2011, Cieslik et al., 2015, Zhang, Geng & Lee, 2017). Among different response inhibition tasks, several meta-analyses showed that an action-restraint (Go/NoGo task) elicits stronger activation in the right DLPFC and parietal cortex compared to another inhibition process, namely, action cancellation (stop-signal task) (Swick, Ashley & Turken, 2011, Cieslik et al., 2015, Zhang, Geng & Lee, 2017). Furthermore, "response uncertainty" tasks modelled by equal probability Go/NoGo stimuli evoked more activity in the right DLPFC (near the cluster revealed in the current study with coordinates [x=39 y=33 z=33], see Table 2) compared to Go/NoGo and stop-signal tasks with a low probability of inhibitory stimuli ("response override" tasks) (Levy & Wagner, 2011).

In meta-analyses of Simmonds, Pekar & Mostofsky (2008) and Criaud & Boulinguez (2013), it was supposed that the increased activity in the right DLPFC observed during Go/NoGo tasks may be due to increased demands caused by working-memory and attentional load rather than the inhibition processes per se. The authors explained it by the fact that most of the fMRI studies used complex designs. Thus, it would be relatively more difficult for the subject to identify a NoGo stimulus, which was believed to increase the need for inhibitory control and consequently increase the brain inhibitory activity in NoGo trials. However, this confounding effect was controlled in the current fMRI study because we utilized equally probable presentation of Go and NoGo stimuli, and the performed meta-analysis included only studies using condition with equiprobable Go and NoGo stimuli presentation.

Another prefrontal structure related to non-selective response inhibition in the current study was the left IFG (opercular part, AIFO). At the same time, the right IFG is commonly associated with response inhibition (Aron, Robbins & Poldrack, 2014). Based on the study of patients with brain damage (Aron et al., 2003) and other neuroimaging studies using primarily the stop-signal tasks (Aron, Robbins & Poldrack, 2014), it was claimed that the right IFG represents a key node of the response inhibition brain system. Currently, this opinion is under active discussion (Mostofsky & Simmonds, 2008, Hampshire et al., 2010, Hampshire, 2015a, Hampshire & Sharp, 2015b, Swick, Ashley & Turken, 2008, Swick & Chatham, 2014). It is noted that Aron et al. predominantly relied upon studies using the stop-signal task and did not consider studies using Go/NoGo tasks (Swick & Chatham, 2014). The studies by Swick, Ashley & Turken (2008) and Kramer et al. (2013) revealed worse performance in the Go/NoGo task when left (not right) IFG was damaged. According to these findings, the left IFG can also be

involved in inhibitory control during action restraint, which is additionally confirmed by our results.

Regarding the practically equivalent involvement of PMC in both Go and NoGo trials, it was previously demonstrated to participate in the planning and coordination of actions because its electrical stimulation results in involuntary motor actions (Desmurget et al., 2009). According to Duque et al. (2012), premotor cortex function is associated with inhibition of any premature actions and control of the time of action execution. The authors refer to this brain mechanism as "impulse control", a concept that is similar to non-selective response inhibition (Criaud et al., 2017).

In summary, it can be noted that the selectivity of response inhibition has been more extensively studied to date than its non-selectivity. A considerable amount of research on inhibitory control has used tasks with a low probability of inhibitory stimuli. However, over time, it has become apparent that under several conditions, non-selectivity of response inhibition can in principle occur. Studies on non-selectivity of response inhibition typically focus on the possible brain mechanisms for non-selective ("global") inhibition of motor responses in the settings of interference among multiple concurrent response options (De Jong, Coles & Logan, 1995, Coxon, Stinear & Byblow, 2007, Aron & Verbruggen, 2008, Frank, 2006, Duque & Ivry, 2009, Duque et al., 2010, MacDonald et al., 2017). Recently, the possibility of involvement of non-selective response inhibition not only for NoGo stimuli but also for Go stimuli was supposed by Criaud et al. (2017). In the current study, the practical equivalence of BOLD signals evoked by NoGo and Go stimuli was demonstrated for a number of brain areas associated with response inhibition as identified by the meta-analysis of fMRI studies. The present study results revealed in the settings of context uncertainty modelled by the equal probability of Go and NoGo trials proved the idea that response inhibition can act as a non-selective mechanism of action inhibition. Thus, one promising area of further research of brain mechanisms of response inhibition (and inhibitory control in general) is the study of the interplay between selective and non-selective inhibition as a function of the context uncertainty degree.

5. Conclusion

For the first time, combining a meta-analysis approach and second-level Bayesian analysis yielded results favouring the existence of non-selective response inhibition. In the present work, selectivity means that inhibition is triggered only by an inhibitory stimulus. The overlap between brain areas previously associated with response inhibition and brain areas demonstrating the practical equivalence of neuronal activity in equiprobable Go and NoGo trials was observed in the right DLPFC, IPL, TPJ, PMC and left IFG (AIFO). When a subject was waiting for an equiprobable Go or NoGo stimulus, a non-selective inhibitory control process occurred in both

Go trials and NoGo trials in opposition to the model of selective response inhibition. This type of response inhibition prevents the performance of any premature motor actions and operates in non-selective, "global" mode. Its involvement is favoured by context uncertainty caused by the equally probable presentation of Go and NoGo stimuli. Presumably, upon the identification of a Go stimulus, the inhibition is released, and the process of action execution is initiated, i.e., it acts as a gating mechanism for accessing a prepared motor program. At the same time, we do not discard the opportunity of involvement of selective inhibitory mechanisms in a less uncertain context. Therefore, future research should address the issue of how brain mechanisms of selective and non-selective response inhibition are inter-related.

6. Acknowledgments

Current study in a part for Bayesian analysis of fMRI data was supported by the Russian Science Foundation grant #19-18-00454. The meta-analysis of Go/NoGo fMRI studies was performed within the state assignment of the Ministry of Education and Science of Russian Federation (theme number AAAA-A19-119101890066-2).

7. References

Albares, M., Lio, G., Criaud, M., Anton, J.-L., Desmurget, M., & Boulinguez, P. (2014). The dorsal medial frontal cortex mediates automatic motor inhibition in uncertain contexts: Evidence from combined fMRI and EEG studies. *Human Brain Mapping*, *35*(11), 5517–5531. https://doi.org/10.1002/hbm.22567

Amrhein, V., Korner-Nievergelt, F., & Roth, T. (2017). The earth is flat (p > 0.05): significance thresholds and the crisis of unreplicable research. *PeerJ*, 5, e3544. https://doi.org/10.7717/peerj.3544

Aron, Adam R. (2011). From Reactive to Proactive and Selective Control: Developing a Richer Model for Stopping Inappropriate Responses. *Biological Psychiatry*, *69*(12), e55–e68. https://doi.org/10.1016/j.biopsych.2010.07.024

Aron, Adam R., Fletcher, P. C., Bullmore, E. T., Sahakian, B. J., & Robbins, T. W. (2003). Stop-signal inhibition disrupted by damage to right inferior frontal gyrus in humans. *Nature Neuroscience*, *6*(2), 115–116. https://doi.org/10.1038/nn1003

Aron, Adam R., Robbins, T. W., & Poldrack, R. A. (2014). Inhibition and the right inferior frontal cortex: one decade on. *Trends in Cognitive Sciences*, *18*(4), 177–185. https://doi.org/10.1016/j.tics.2013.12.003

Aron, Adam R., & Verbruggen, F. (2008). Stop the Presses. *Psychological Science*, *19*(11), 1146–1153. https://doi.org/10.1111/j.1467-9280.2008.02216.x

Badry, R., Mima, T., Aso, T., Nakatsuka, M., Abe, M., Fathi, D., Foly, N., Nagiub, H., Nagamine, T., & Fukuyama, H. (2009). Suppression of human cortico-motoneuronal excitability

during the Stop-signal task. *Clinical Neurophysiology*, 120(9), 1717–1723. https://doi.org/10.1016/j.clinph.2009.06.027

Boulinguez, P., Jaffard, M., Granjon, L., & Benraiss, A. (2008). Warning Signals Induce Automatic EMG Activations and Proactive Volitional Inhibition: Evidence From Analysis of Error Distribution in Simple RT. *Journal of Neurophysiology*, *99*(3), 1572–1578. https://doi.org/10.1152/jn.01198.2007

Braver, T. S. (2012). The variable nature of cognitive control: a dual mechanisms framework. *Trends in Cognitive Sciences*, *16*(2), 106–113. https://doi.org/10.1016/j.tics.2011.12.010

Buchsbaum, B. R., Greer, S., Chang, W.-L., & Berman, K. F. (2005). Meta-analysis of neuroimaging studies of the Wisconsin Card-Sorting task and component processes. *Human Brain Mapping*, *25*(1), 35–45. https://doi.org/10.1002/hbm.20128

Chevrier, A. D., Noseworthy, M. D., & Schachar, R. (2007). Dissociation of response inhibition and performance monitoring in the stop signal task using event-related fMRI. *Human Brain Mapping*, *28*(12), 1347–1358. https://doi.org/10.1002/hbm.20355

Cieslik, E. C., Mueller, V. I., Eickhoff, C. R., Langner, R., & Eickhoff, S. B. (2015). Three key regions for supervisory attentional control: Evidence from neuroimaging meta-analyses. *Neuroscience & Biobehavioral Reviews*, *48*, 22–34. https://doi.org/10.1016/j.neubiorev.2014.11.003

Cole, M. W., Bassett, D. S., Power, J. D., Braver, T. S., & Petersen, S. E. (2014). Intrinsic and Task-Evoked Network Architectures of the Human Brain. *Neuron*, *83*(1), 238–251. https://doi.org/10.1016/j.neuron.2014.05.014

Coxon, J. P., Stinear, C. M., & Byblow, W. D. (2007). Selective Inhibition of Movement. *Journal of Neurophysiology*, *97*(3), 2480–2489. https://doi.org/10.1152/jn.01284.2006

Criaud, M., & Boulinguez, P. (2013). Have we been asking the right questions when assessing response inhibition in go/no-go tasks with fMRI? A meta-analysis and critical review. *Neuroscience & Biobehavioral Reviews*, *37*(1), 11–23. https://doi.org/10.1016/j.neubiorev.2012.11.003

Criaud, M., Longcamp, M., Anton, J.-L., Nazarian, B., Roth, M., Sescousse, G., … Boulinguez, P. (2017). Testing the physiological plausibility of conflicting psychological models of response inhibition: A forward inference fMRI study. *Behavioural Brain Research*, *333*, 192–202. https://doi.org/10.1016/j.bbr.2017.06.030

De Jong, R., Coles, M. G. H., & Logan, G. D. (1995). Strategies and mechanisms in nonselective and selective inhibitory motor control. *Journal of Experimental Psychology: Human Perception and Performance*, *21*(3), 498–511. https://doi.org/10.1037//0096-1523.21.3.498

Dienes, Z. (2014). Using Bayes to get the most out of non-significant results. *Frontiers in Psychology*, *5*. https://doi.org/10.3389/fpsyg.2014.00781

Dell, R. B., Holleran, S., & Ramakrishnan, R. (2002). Sample Size Determination. *ILAR Journal*, 43(4), 207–213. https://doi.org/10.1093/ilar.43.4.207

Desmurget, M., Reilly, K. T., Richard, N., Szathmari, A., Mottolese, C., & Sirigu, A. (2009). Movement Intention After Parietal Cortex Stimulation in Humans. *Science*, *324*(5928), 811–813. https://doi.org/10.1126/science.1169896

Di Russo, F., Lucci, G., Sulpizio, V., Berchicci, M., Spinelli, D., Pitzalis, S., & Galati, G. (2016). Spatiotemporal brain mapping during preparation, perception, and action. *NeuroImage*, *126*, 1–14. https://doi.org/10.1016/j.neuroimage.2015.11.036

Duncan, J. (2010). The multiple-demand (MD) system of the primate brain: mental programs for intelligent behaviour. *Trends in Cognitive Sciences*, *14*(4), 172–179. https://doi.org/10.1016/j.tics.2010.01.004

Dunovan, K., Lynch, B., Molesworth, T., & Verstynen, T. (2015). Competing basal ganglia pathways determine the difference between stopping and deciding not to go. *ELife*, *4*. https://doi.org/10.7554/eLife.08723

Duque, J., & Ivry, R. B. (2009). Role of Corticospinal Suppression during Motor Preparation. *Cerebral Cortex*, *19*(9), 2013–2024. https://doi.org/10.1093/cercor/bhn230

Duque, J., Labruna, L., Verset, S., Olivier, E., & Ivry, R. B. (2012). Dissociating the Role of Prefrontal and Premotor Cortices in Controlling Inhibitory Mechanisms during Motor Preparation. *Journal of Neuroscience*, *32*(3), 806–816. https://doi.org/10.1523/JNEUROSCI.4299-12.2012

Duque, J., Lew, D., Mazzocchio, R., Olivier, E., & Ivry, R. B. (2010). Evidence for Two Concurrent Inhibitory Mechanisms during Response Preparation. *Journal of Neuroscience*, *30*(10), 3793–3802. https://doi.org/10.1523/JNEUROSCI.5722-09.2010

Eickhoff, S. B., Grefkes, C., Fink, G. R., & Zilles, K. (2008). Functional Lateralization of Face, Hand, and Trunk Representation in Anatomically Defined Human Somatosensory Areas. *Cerebral Cortex*, *18*(12), 2820–2830. https://doi.org/10.1093/cercor/bhn039

Eickhoff, Simon B., Nichols, T. E., Laird, A. R., Hoffstaedter, F., Amunts, K., Fox, P. T., … Eickhoff, C. R. (2016). Behavior, sensitivity, and power of activation likelihood estimation characterized by massive empirical simulation. *NeuroImage*, *137*, 70–85. https://doi.org/10.1016/j.neuroimage.2016.04.072

Erika-Florence, M., Leech, R., & Hampshire, A. (2014). A functional network perspective on response inhibition and attentional control. *Nature Communications*, *5*(1). https://doi.org/10.1038/ncomms5073

Faul, F., Erdfelder, E., Lang, A.-G., & Buchner, A. (2007). G*Power 3: A flexible statistical power analysis program for the social, behavioral, and biomedical sciences. *Behavior Research Methods*, 39(2), 175–191. https://doi.org/10.3758/bf03193146

Frank, M. J. (2006). Hold your horses: A dynamic computational role for the subthalamic nucleus in decision making. *Neural Networks*, *19*(8), 1120–1136. https://doi.org/10.1016/j.neunet.2006.03.006

Friedman, N. P., Miyake, A., Young, S. E., DeFries, J. C., Corley, R. P., & Hewitt, J. K. (2008). Individual differences in executive functions are almost entirely genetic in origin. *Journal of Experimental Psychology: General*, *137*(2), 201–225. https://doi.org/10.1037/0096-3445.137.2.201

Friston, K. J., Glaser, D. E., Henson, R. N. A., Kiebel, S., Phillips, C., & Ashburner, J. (2002b). Classical and Bayesian Inference in Neuroimaging: Applications. *NeuroImage*, *16*(2), 484–512. https://doi.org/10.1006/nimg.2002.1091

Friston, K. J., & Penny, W. (2003). Posterior probability maps and SPMs. *NeuroImage*, *19*(3), 1240–1249. https://doi.org/10.1016/S1053-8119(03)00144-7

Friston, K. J., Penny, W., Phillips, C., Kiebel, S., Hinton, G., & Ashburner, J. (2002a). Classical and Bayesian Inference in Neuroimaging: Theory. *NeuroImage*, *16*(2), 465–483. https://doi.org/10.1006/nimg.2002.1090

Ganos, C., Rothwell, J., & Haggard, P. (2018). Voluntary inhibitory motor control over involuntary tic movements. *Movement Disorders*, *33*(6), 937–946. https://doi.org/10.1002/mds.27346

Genovese, C. R., Lazar, N. A., & Nichols, T. (2002). Thresholding of Statistical Maps in Functional Neuroimaging Using the False Discovery Rate. *NeuroImage*, *15*(4), 870–878. https://doi.org/10.1006/nimg.2001.1037

Gorgolewski, K. J., Durnez, J., & Poldrack, R. A. (2017). Preprocessed Consortium for Neuropsychiatric Phenomics dataset. *F1000Research*, 6, 1262. https://doi.org/10.12688/f1000research.11964.2

Hampshire, A. (2015a). Putting the brakes on inhibitory models of frontal lobe function. *NeuroImage*, *113*, 340–355. https://doi.org/10.1016/j.neuroimage.2015.03.053

Hampshire, A., Chamberlain, S. R., Monti, M. M., Duncan, J., & Owen, A. M. (2010). The role of the right inferior frontal gyrus: inhibition and attentional control. *NeuroImage*, *50*(3), 1313–1319. https://doi.org/10.1016/j.neuroimage.2009.12.109

Hampshire, A., & Sharp, D. J. (2015b). Contrasting network and modular perspectives on inhibitory control. *Trends in Cognitive Sciences*, *19*(8), 445–452. https://doi.org/10.1016/j.tics.2015.06.006

Haynes, W. I. A., & Haber, S. N. (2013). The Organization of Prefrontal-Subthalamic Inputs in Primates Provides an Anatomical Substrate for Both Functional Specificity and Integration: Implications for Basal Ganglia Models and Deep Brain Stimulation. *Journal of Neuroscience*, 33(11), 4804–4814. https://doi.org/10.1523/jneurosci.4674-12.2013

Hsieh, S., Wu, M., & Tang, C.-H. (2016). Adaptive Strategies for the Elderly in Inhibiting Irrelevant and Conflict No-Go Trials while Performing the Go/No-Go Task. *Frontiers in Aging Neuroscience*, *7*. https://doi.org/10.3389/fnagi.2015.00243

Hugdahl, K., Raichle, M. E., Mitra, A., & Specht, K. (2015). On the existence of a generalized non-specific task-dependent network. *Frontiers in Human Neuroscience*, *9*. https://doi.org/10.3389/fnhum.2015.00430

Isoda, M., & Hikosaka, O. (2011). Cortico-basal ganglia mechanisms for overcoming innate, habitual and motivational behaviors. *European Journal of Neuroscience*, *33*(11), 2058–2069. https://doi.org/10.1111/j.1460-9568.2011.07698.x

Jaffard, M., Benraiss, A., Longcamp, M., Velay, J.-L., & Boulinguez, P. (2007). Cueing method biases in visual detection studies. *Brain Research*, *1179*, 106–118. https://doi.org/10.1016/j.brainres.2007.08.032

Jaffard, M., Longcamp, M., Velay, J.-L., Anton, J.-L., Roth, M., Nazarian, B., & Boulinguez, P. (2008). Proactive inhibitory control of movement assessed by event-related fMRI. NeuroImage, 42(3), 1196–1206. https://doi.org/10.1016/j.neuroimage.2008.05.041

Johnstone, T., Ores Walsh, K. S., Greischar, L. L., Alexander, A. L., Fox, A. S., Davidson, R. J., & Oakes, T. R. (2006). Motion correction and the use of motion covariates in multiple-subject fMRI analysis. *Human Brain Mapping*, *27*(10), 779–788. https://doi.org/10.1002/hbm.20219

Krämer, U. M., Solbakk, A.-K., Funderud, I., Løvstad, M., Endestad, T., & Knight, R. T. (2013). The role of the lateral prefrontal cortex in inhibitory motor control. *Cortex*, *49*(3), 837–849. https://doi.org/10.1016/j.cortex.2012.05.003

Kropotov, J. D. (2016). *Functional Neuromarkers for Psychiatry: Applications for Diagnosis and Treatment* (1st ed.). Cambridge, MA: Academic Press.

Kropotov, J. D., Ponomarev, V. A., Hollup, S., & Mueller, A. (2011). Dissociating action inhibition, conflict monitoring and sensory mismatch into independent components of event related potentials in GO/NOGO task. NeuroImage, 57(2), 565–575. https://doi.org/10.1016/j.neuroimage.2011.04.060

Kropotov, J. D., & Ponomarev, V. A. (2015). Differentiation of neuronal operations in latent components of event-related potentials in delayed match-to-sample tasks. *Psychophysiology*, *52*(6), 826–838. https://doi.org/10.1111/psyp.12410

Kruschke, J. K. (2018). Rejecting or Accepting Parameter Values in Bayesian Estimation. Advances in Methods and Practices in Psychological Science, 1(2), 270–280. https://doi.org/10.1177/2515245918771304

Kruschke, J. K., & Liddell, T. M. (2015). The Bayesian New Statistics: Two Historical Trends Converge. *SSRN Electronic Journal*. https://doi.org/10.2139/ssrn.2606016

Lakens, D. (2017). Equivalence Tests. *Social Psychological and Personality Science*, *8*(4), 355–362. https://doi.org/10.1177/1948550617697177

Laurens, K. R., Kiehl, K. A., & Liddle, P. F. (2004). A supramodal limbic-paralimbic-neocortical network supports goal-directed stimulus processing. *Human Brain Mapping*, *24*(1), 35–49. https://doi.org/10.1002/hbm.20062

Lavric, A., Pizzagalli, D. A., & Forstmeier, S. (2004). When 'go' and 'nogo' are equally
frequent: ERP components and cortical tomography. *European Journal of Neuroscience*, *20*(9),
2483–2488. https://doi.org/10.1111/j.1460-9568.2004.03683.x

Lee, K.-H., Pluck, G., Lekka, N., Horton, A., Wilkinson, I. D., & Woodruff, P. W. R. (2015).
Self-harm in schizophrenia is associated with dorsolateral prefrontal and posterior cingulate
activity. *Progress in Neuro-Psychopharmacology and Biological Psychiatry*, *61*, 18–23.
https://doi.org/10.1016/j.pnpbp.2015.03.005

Levy, B. J., & Wagner, A. D. (2011). Cognitive control and right ventrolateral prefrontal cortex:
reflexive reorienting, motor inhibition, and action updating. *Annals of the New York Academy of*
Sciences, *1224*(1), 40–62. https://doi.org/10.1111/j.1749-6632.2011.05958.x

Logan, G. D., Van Zandt, T., Verbruggen, F., & Wagenmakers, E.-J. (2014). On the ability to
inhibit thought and action: General and special theories of an act of control. *Psychological*
Review, *121*(1), 66–95. https://doi.org/10.1037/a0035230

MacDonald, H. J., McMorland, A. J. C., Stinear, C. M., Coxon, J. P., & Byblow, W. D. (2017).
An Activation Threshold Model for Response Inhibition. *PLOS ONE*, *12*(1), e0169320.
https://doi.org/10.1371/journal.pone.0169320

Magerkurth, J., Mancini, L., Penny, W., Flandin, G., Ashburner, J., Micallef, C., … Weiskopf,
N. (2015). Objective Bayesian fMRI analysis - a pilot study in different clinical environments.
Frontiers in Neuroscience, *9*. https://doi.org/10.3389/fnins.2015.00168

Majid, D. S. A., Cai, W., George, J. S., Verbruggen, F., & Aron, A. R. (2011). Transcranial
Magnetic Stimulation Reveals Dissociable Mechanisms for Global Versus Selective
Corticomotor Suppression Underlying the Stopping of Action. *Cerebral Cortex*, 22(2), 363–371.
https://doi.org/10.1093/cercor/bhr112

Meffert, H., Hwang, S., Nolan, Z. T., Chen, G., & Blair, J. R. (2016). Segregating attention from
response control when performing a motor inhibition task. *NeuroImage*, *126*, 27–38.
https://doi.org/10.1016/j.neuroimage.2015.11.029

Mostofsky, S. H., & Simmonds, D. J. (2008). Response Inhibition and Response Selection: Two
Sides of the Same Coin. *Journal of Cognitive Neuroscience*, *20*(5), 751–761.
https://doi.org/10.1162/jocn.2008.20500

Nambu, A., Takada, M., Inase, M., & Tokuno, H. (1996). Dual somatotopical representations in
the primate subthalamic nucleus: evidence for ordered but reversed body-map transformations
from the primary motor cortex and the supplementary motor area. *The Journal of Neuroscience*,
16(8), 2671–2683. https://doi.org/10.1523/jneurosci.16-08-02671.1996

Nee, D. E., Wager, T. D., & Jonides, J. (2007). Interference resolution: Insights from a meta-
analysis of neuroimaging tasks. *Cognitive, Affective, & Behavioral Neuroscience*, *7*(1), 1–17.
https://doi.org/10.3758/CABN.7.1.1

Neumann, J., & Lohmann, G. (2003). Bayesian second-level analysis of functional magnetic
resonance images. *NeuroImage*, *20*(2), 1346–1355. https://doi.org/10.1016/S1053-
8119(03)00443-9

Oldfield, R. C. (1971). The assessment and analysis of handedness: The Edinburgh inventory.
Neuropsychologia, *9*(1), 97–113. https://doi.org/10.1016/0028-3932(71)90067-4

Poldrack, R. A., Mumford, J. A., & Nichols, T. E. (2011). Handbook of Functional MRI Data
Analysis. Cambridge, United Kingdom: Cambridge University Press.

Poldrack, R. A., Congdon, E., Triplett, W., Gorgolewski, K. J., Karlsgodt, K. H., Mumford, J.
A., Sabb, F. W., Freimer, N. B., London, E. D., Cannon, T. D., & Bilder, R. M. (2016). A
phenome-wide examination of neural and cognitive function. *Scientific Data*, 3(1), 1–12.
https://doi.org/10.1038/sdata.2016.110

Simmonds, D. J., Pekar, J. J., & Mostofsky, S. H. (2008). Meta-analysis of Go/No-go tasks
demonstrating that fMRI activation associated with response inhibition is task-dependent.
Neuropsychologia, *46*(1), 224–232. https://doi.org/10.1016/j.neuropsychologia.2007.07.015

Swick, D., Ashley, V., & Turken, A. U. (2008). Left inferior frontal gyrus is critical for response
inhibition. *BMC Neuroscience*, *9*(1), 102. https://doi.org/10.1186/1471-2202-9-102

Swick, D., Ashley, V., & Turken, A. U. (2011). Are the neural correlates of stopping and not
going identical? Quantitative meta-analysis of two response inhibition tasks. *NeuroImage*, *56*(3),
1655–1665. https://doi.org/10.1016/j.neuroimage.2011.02.070

Swick, D., & Chatham, C. H. (2014). Ten years of inhibition revisited. *Frontiers in Human*
Neuroscience, *8*. https://doi.org/10.3389/fnhum.2014.00329

Verbruggen, F., Best, M., Bowditch, W. A., Stevens, T., & McLaren, I. P. L. (2014). The
inhibitory control reflex. *Neuropsychologia*, *65*, 263–278.
https://doi.org/10.1016/j.neuropsychologia.2014.08.014

Verbruggen, F., & Logan, G. D. (2008). Response inhibition in the stop-signal paradigm. *Trends*
in Cognitive Sciences, *12*(11), 418–424. https://doi.org/10.1016/j.tics.2008.07.005

Vink, M., Kaldewaij, R., Zandbelt, B. B., Pas, P., & du Plessis, S. (2015). The role of stop-signal
probability and expectation in proactive inhibition. *European Journal of Neuroscience*, *41*(8),
1086–1094. https://doi.org/10.1111/ejn.12879

Vink, M., Zandbelt, B. B., Gladwin, T., Hillegers, M., Hoogendam, J. M., van den Wildenberg,
W. P. M., … Kahn, R. S. (2014). Frontostriatal activity and connectivity increase during
proactive inhibition across adolescence and early adulthood. *Human Brain Mapping*, *35*(9),
4415–4427. https://doi.org/10.1002/hbm.22483

Woolrich, M. W., Behrens, T. E. J., Beckmann, C. F., & Smith, S. M. (2005). Mixture models
with adaptive spatial regularization for segmentation with an application to FMRI data. *IEEE*
Transactions on Medical Imaging, *24*(1), 1–11. https://doi.org/10.1109/TMI.2004.836545

Zhang, R., Geng, X., & Lee, T. M. C. (2017). Large-scale functional neural network correlates of
response inhibition: an fMRI meta-analysis. *Brain Structure and Function*, *222*(9), 3973–3990.
https://doi.org/10.1007/s00429-017-1443-x

Meta-analysis

ALE-score: 0.005 — 0.014

A) Classical voxel-wise inference

T-value: 5.59 ▮ 17.55
Go>NoGo   Non-significant

B) Bayesian parameter inference

Log Odds: 3 ▮ 27   3 ▮ 27
Go>NoGo   NoGo = Go   Low-confidence

Overlap between meta-analysis and Bayesian analysis