1   **Non-selective response inhibition in Go/NoGo task: Bayesian analysis of fMRI data**

2   Ruslan Masharipov, Maxim Kireev*, Alexander Korotkov, and Svyatoslav Medvedev
3
4   1 - N.P. Bechtereva Institute of the Human Brain, Russian Academy of Sciences, Academika Pavlova Street 9, St.
5   Petersburg, 197376, Russia
6
7   Corresponding author: Maxim Kireev;
8   Address: Academika Pavlova Street 9, St. Petersburg, 197376, Russia
9   E-mail: kireev@ihb.spb.ru
10   Phone number: +7 812 670-09-51
11   Fax number: +7 812 2343247
12

Abstract

Response inhibition is typically considered a selective mechanism triggered by particular

"inhibitory" stimuli or events. Based on recent research, an alternative non-selective mechanism

was proposed by several authors. Presumably, the mechanism may involve non-selective

inhibition of all potential responses to allow execution of an appropriate response, and such

inhibition may be triggered not only by the presentation of "inhibitory" stimuli but also by any

imperative stimuli, including Go stimuli. Earlier support for this notion was mainly based on the

absence of a significant difference between neural activity evoked by equiprobable Go and

NoGo stimuli. Previous neuroimaging studies utilized classical null hypothesis significance

testing, which makes it impossible to accept the null hypothesis. Therefore, the current research

aimed to provide evidence for practical equivalence of neuronal activity in Go and NoGo trials

using Bayesian analysis of functional magnetic resonance imaging (fMRI) data. Twenty healthy

participants performed a cued Go/NoGo task with an equiprobable presentation of Go and NoGo

stimuli. To reveal brain areas previously associated with response inhibition in similar

experimental conditions, we performed a meta-analysis of fMRI studies using equal probability

Go/NoGo tasks. As a result, we observed overlap between response inhibition areas and areas

demonstrating practical equivalence of neuronal activity located in the right dorsolateral

prefrontal cortex, inferior parietal lobule, premotor cortex, and left inferior frontal gyrus. Thus,

obtained results favour the existence of non-selective response inhibition, which can act in

settings of context uncertainty, modelled by the equal probability of Go and NoGo stimuli.

Key words: non-selective response inhibition, functional MRI, Bayesian inference, accepting

the null hypothesis, practical equivalence, Go/NoGo task

Highlights

• Non-selective response inhibition was assessed by equiprobable Go/NoGo task

• Bayesian analysis of fMRI data was combined with a meta-analysis of fMRI studies

• Nodes of response-inhibition system were equally involved in Go and NoGo trials

• Evidence for non-selective response inhibition in uncertainty context was found

1. Introduction

Response inhibition is the ability to suppress inappropriate, automatic, reflexive, or

habitual prepotent responses to produce a controlled goal-directed response (Friedman et al.,

2008, Isoda & Hikosaka, 2011). It is generally accepted in the literature that response inhibition

works in close relation to other processes associated with cognitive control, such as working

memory, voluntary attention, conflict monitoring and action selection (Buchsbaum et al., 2005,

Nee, Wager & Jonides, 2007, Simmonds, Pekar & Mostofsky, 2008, Criaud & Boulinguez,

2013, Cieslik et al., 2015). Moreover, response inhibition is thought to represent a multifaceted

phenomenon rather than a unitary brain mechanism. A distinction is made between action

cancellation and action restraint (Swick, Ashley & Turken, 2011, Cieslik et al., 2015, Zhang,

Geng & Lee, 2017) as well as between reactive and proactive response inhibition mechanisms

(Jaffard et al., 2007, 2008, Aron, 2011, Braver, 2012, Erika-Florence, Leech & Hampshire,

2013).

According to the conventional view, the response inhibition process is selectively triggered

by "inhibitory" stimuli that result in increased neuronal activity in brain structures responsible

for inhibitory control (Verbruggen & Logan, 2008, Verbruggen et al., 2014, Logan et al., 2014,

Criaud et al., 2017). However, in several cases, the concept of selective response inhibition fails

to explain observed behavioural and neurophysiological phenomena. Manipulating the

probability of the occurrence of "inhibitory" stimuli and the subjects' awareness of the

probability of the appearance may slow down the motor response (Chevrier, Noseworthy &

Schachar, 2007, Jaffard et al., 2007, Boulinguez et al., 2008, Vink et al., 2014, Vink et al., 2015,

Dunovan et al., 2015, Meffert et al., 2016, Hsieh, Wu & Tang, 2016). Moreover, when it is

necessary to suppress a specific action selectively, that action is inhibited as well as all other

potential actions (Coxon, Stinear & Byblow, 2007, Aron & Verbruggen, 2008, Duque & Ivry,

2009, Duque et al., 2010, MacDonald et al., 2017). In attempting to explain the effects

mentioned above, several authors proposed the concept of non-selective ("global") response

inhibition (Frank, 2006, Aron, 2011, Criaud et al., 2017). First, response inhibition mechanisms

may non-selectively inhibit all potential responses to further selectively execute an appropriate

response, i.e., non-selective inhibition of multiple concurrent motor responses. Second,

inhibition may be triggered not only by the presentation of "inhibitory" stimuli but also by the

occurrence of any imperative stimuli instructing on the necessity to suppress or execute a

prepared action (non-selectivity of inhibitory stimulus perception). It is thought that those non-

selective mechanisms serve to prevent inappropriate or premature responses at the expense of the speed of execution of correct actions.

A tentative neurophysiological model of the non-selective or "global" response inhibition involved in the resolution of interference between several competing response options was proposed by Frank (2006), including the cortico-subthalamic "hyper-direct" pathway, which is capable of rapidly and simultaneously non-selectively suppressing all potential response options. As the research area developed, it was hypothesized that the model might be applied not only to tasks with multiple concurrent response options but also to simpler tasks where the subject has to choose between executing and refraining from an action (Albares et al., 2014, Criaud et al., 2017). The authors used a cued Go/NoGo task, wherein a preparatory cue stimulus indicated the probability of the occurrence of a NoGo stimulus. Within the framework of the selective inhibition model, the inhibition process would only be triggered by the identification of the NoGo stimulus. According to the hypothesis of non-selective inhibition, when the context is uncertain (equal probability of NoGo and Go stimuli), the need for response inhibition arises for both NoGo and Go trials. Experimental assessment of the hypotheses revealed no statistically significant differences between NoGo and Go trials both in the amplitude of early components of event-related potentials (ERP) (Albares et al., 2014) and in the level of neuronal activity measured by functional magnetic resonance imaging (fMRI) (Criaud et al., 2017). The authors considered this fact as evidence in favour of the presence of a "non-selective" inhibitory mechanism that may be engaged when the context is uncertain. The results of the analysis of early ERP components suggest that non-selective inhibition may act as a gating mechanism acting on the initiation of the motor program of the prepared action.

However, a critical limitation of the abovementioned studies was that the authors cannot accept the null hypothesis, so they did not provide direct proof of the practical equivalence of neuronal activity level between Go and NoGo trials in the inhibition-related brain structures. Indeed, within the framework of classical (frequentist) null hypothesis significance testing (NHST), we cannot accept the null hypothesis based on the absence of a significant difference. We can only reject it (Kruschke & Liddell, 2015). Thus, the question of experimental support of non-selective response inhibition remains unanswered, and answering this question requires overcoming the methodological limitation of NHST, which is possible using Bayesian statistics (Kruschke & Liddell, 2015).

Therefore, the present study aimed to verify the non-selective response inhibition hypothesis by using the fMRI data obtained for an equiprobable Go/NoGo task. Based on the results of prior studies, it may be suggested that if the hypothesis on the non-selectivity of inhibition is correct, then the brain structures responsible for response inhibition will

demonstrate practically equivalent levels of neuronal activity in equiprobable Go and NoGo
trials. To assess this prediction, the Bayesian parameter inference was applied. Instead, if the
hypothesis on the selectivity of response inhibition is correct, then activation of the inhibition-
related brain structures will be observed in the NoGo-conditions compared with Go-conditions.

2. Material and methods

2.1. A meta-analysis of fMRI studies using equal probability Go/NoGo tasks.

Given that a practically equivalent level of neuronal activity may be identified not only for
response inhibition-related structures but, for example, those related to sensory processing of
visual stimuli, working memory, and attention, we conducted a meta-analysis of fMRI studies
aimed at revealing brain structures associated with response inhibition. Studies using equal
probability Go/NoGo tasks comparable to the present study were selected for the meta-analysis.
We searched for studies that, similar to the investigation by Criaud et al. (2017), compared the
neuronal activity in the condition of equiprobable Go and NoGo stimuli presentation with the
control Go condition, wherein the subject did not need to inhibit the prepared action.

We searched for these studies in the PubMed database in the period of 01/01/2000 to
15/07/2019 using the following keywords: "((fmri) OR (functional magnetic resonance)) AND
((nogo) OR (no-go))." Four additional studies were identified through manual searches. As a
result, 726 papers were identified. At the first stage of selection, we excluded reviews, meta-
analyses, and papers repeatedly reporting the results of fMRI studies (see the flow chart of
analysis at Fig. 1).

Figure 1. Flow chart of study selection in the meta-analysis.

Studies dealing with Flanker task modifications of Go/NoGo task or based on unequal
probability of appearance of Go and NoGo trials were also excluded from the analysis. As a
result, 593 papers were excluded. Task designs of the remaining 113 papers provided for equal
probability Go and NoGo stimuli presentation. At the next step, 82 of the papers were excluded
based on the following criteria: auditory and sensorimotor Go/NoGo tasks were used; only
emotion-laden task conditions were used (emotionally neutral conditions were either absent or
not considered separately); the necessary control Go-condition (100% probability of the Go-
stimulus presentation) was not used; or the coordinates for the contrast of interest "Go/NoGo >
Go-control" within a group of healthy subjects were not reported. In 11 out of the 31 remaining
studies, healthy volunteer subjects under the age of 18 (children and adolescents) were studied.
These papers were also excluded from analysis because this study focused on the brain activity
of healthy, adult subjects. In all remaining studies except one, only block designs were used. To
make our sample more homogenous, we excluded the only eligible study with an event-related
design (Criaud et al., 2017). The final meta-analysis included 20 studies (452 healthy subjects,
mean age of 29 years, 210 foci) (for more details see supplementary materials, Table S1).

The meta-analysis was conducted using GingerALE 3.0.2 software (http://brainmap.org/ale) with a cluster-forming threshold of 0.001 (uncorrected) and a cluster-level threshold of 0.05 corrected for family-wise error (FWE) (Eickhoff et al., 2016). All the coordinates were converted into Montreal Neurological Institute (MNI) space. The median full width at half maximum (FWHM) value of Gaussian function was set at a level of 13.5 mm.

2.2. Subjects.

Twenty healthy, right-handed volunteer subjects (16 women, aged (mean ± SD): 23.9 ± 4.6) participated in the present fMRI study. An Oldfield test was used to determine the dominant arm (Oldfield, 1971). The subject volunteers signed written informed consent to participate in the study and were paid for their participation. All the procedures were performed in accordance with the Declaration of Helsinki and were approved by the Ethics Committee of the N.P. Bechtereva Institute of the Human brain of the Russian Academy of Sciences.

2.3. Experimental task and study procedure.

We used a paired stimulus modification of the Go/NoGo task (see Fig. 2) (Kropotov, 2016). This task was originally developed to dissociate cognitive processes, such as response inhibition, conflict monitoring, sensory mismatch and category discrimination (Kropotov et al., 2011, 2015). Given that imperative Go and NoGo stimuli were presented in an equiprobable manner, previously collected fMRI data were used for the purposes of the current study. Each trial consisted of the two consequently presented stimuli. The first preparatory cue stimulus warned the subjects on the presentation of the second, imperative stimulus, or indicated that there was no need for any response to the second stimulus. The study included two variants of the experiment's instructions. According to the first instruction (see Fig. 2A), the subject should press the response button upon presentation of the pair of images "animal-animal" ("A-A Go" trials) and refrain from acting upon presentation of the pair "animal-plant" ("A-P NoGo" trials). According to the second experiment's instruction (see Fig. 2B), the subject acts after the presentation of the pair "animal-plant" ("A-P Go" trials) and suppresses an action upon presentation of the pair "animal-animal" ("A-A NoGo" trials).

Figure 2. Experimental design of the Go/NoGo task. S1 – first stimulus (preparatory cue). S2 – second stimulus (imperative stimulus). A – images of animals, P – images of plants. A) First experiment: "A-A Go", "A-P NoGo". B) Second experiment: "A-P Go", "A-A NoGo". Trials that were compared to test the hypothesis of selective and non-selective response inhibition are highlighted by red boxes.

In both experiments, if the first presented stimulus was an image of a "plant," subjects do not need to perform any actions in response to the presentation of any second stimuli of a trial. In such a trial, the subject should ignore the second stimulus and wait for the next pair of stimuli ("P-A Ignore" and "P-P Ignore" trials). Accordingly, it was assumed that there is no inhibition of prepared action in "Ignore" conditions. The "Ignore" trials were included in the adopted Go/NoGo task to evaluate preparatory brain activity, such as preparing to receive a relevant stimulus (attentional set) and preparing to make a movement (motor set), by comparing them to Go and NoGo trials. However, the issue related to preparatory processes is beyond the scope of the present study. Fifty pairs of each type of stimuli were randomly presented in each experiment. The order of following the instructions was counterbalanced among the subjects. Two variants of the present Go/NoGo task made it possible to control the differences in the load on working memory between A-P/A-A - Go/NoGo stimuli (Kropotov & Ponomarev, 2015).

The fMRI data obtained using this experimental task allowed us to test the hypothesis on the *non-selectivity* of response inhibition in the current study given that a similar task design was previously used for the same goal by Criaud et al. (2017). The imperative Go and NoGo stimuli were presented after the preparatory stimulus with equal probability, as reported in event-related studies by (Albares et al., 2014, Di Russo et al., 2016, Kropotov, 2016, Criaud et al., 2017). Such an equally probable presentation of imperative stimuli provides several advantages. First, it minimizes the difference in cognitive load between Go and NoGo conditions arising from task complexity (Criaud & Boulinguez, 2013, Criaud et al., 2017). Second, such a design enables the exclusion of effects associated with a low frequency of the NoGo stimuli presentation ("oddball" effects) confounding the effect of inhibition (Di Russo et al., 2016). Third, it creates maximum uncertainty regarding the probability of the presentation of an imperative stimulus and thus minimizes the conflict between two response types, which makes it possible to distinguish between error monitoring and conflict resolution processes on the one hand and response inhibition processes on the other (Lavric, Pizzagalli & Forstmeier, 2004).

In total, 100 NoGo trials, 100 Go trials, 100 P-A Ignore trials, and 100 P-P Ignore trials were presented over two experimental sessions. In the absence of stimulation, a fixation cross was displayed in the centre of the screen. The stimuli were presented for 100 ms, and the

interstimulus interval was 1000 ms. The intertrial interval jittered from 2800 to 3200 ms with an increment step of 100 ms. Additionally, to improve design efficiency, 100 zero-events (fixation crosses) were randomly inserted between the stimuli pairs (trials), and their duration jittered from 3000 to 5000 ms with an increment size of 500 ms. The action to be performed consisted of pressing a button with the right thumb. The duration of one task session was 17.5 minutes. Before starting the fMRI study, the subjects performed a training task. The Invivo's Eloquence fMRI System (Invivo, Orlando, FL, USA) was used to deliver the stimuli, synchronize with fMRI acquisition, and record reaction times of subjects bottom pressing. The task presentation sequence and all temporal parameters of the stimuli presentation were programmed using the E-prime 2.0 software package (Psychology Software Tools Inc., Pittsburgh, PA, USA).

2.4. Image acquisition.

A Philips Achieva 3.0 Tesla scanner (Philips Medical Systems, Best, Netherlands) was used for the study. The structural T1-images were registered with the following parameters: field of view (FOV) – 240×240 mm, repetition time (TR) – 25 ms, echo time (TE) – 2.2 ms, 130 axial slices with a thickness of 1 mm and pixel size of 1×1 mm, flip angle – 30°. For the registration of T2*-images, a single-pulse echo planar imaging (EPI) sequence was used. The period of data registration from 31 axial slices was 2 seconds (TR = 2 s, TE = 35 ms). The following parameters were employed: FOV – 200×186 mm, flip angle – 90°, voxel size – 3×3×3 mm. Two dummy scans were performed prior to each session. To minimize head motions, we used an MR-compatible soft cervical collar.

2.5. Preprocessing of fMRI-images.

Image preprocessing included the following: realignment to the first image of the session, slice time correction, co-registration, segmentation, normalization to an MNI template, and spatial smoothing (8 mm FWHM). Preprocessing and statistical analysis of the images were performed using an SPM12 (Statistical parametric mapping) software package (http://www.fil.ion.ucl.ac.uk/spm).

2.6. Statistical analysis of fMRI data.

The first level of analysis was conducted using frequentist parameter estimation. The second level of analysis was conducted using both the frequentist and Bayesian parameter estimation (Friston et al., 2002a, Friston & Penny, 2003, Neumann & Lohmann, 2003). Onset times of second stimuli presentation (separately for "A-A Go", "A-P NoGo", "P-A Ignore Exp1", "P-P Ignore Exp1", and also "A-P Go", "A-A NoGo", "P-A Ignore Exp2", "P-P Ignore Exp2") and erroneous button pressing and missing the responding in Go trials were used to create regressors of the general linear model (GLM) for each subject. Events were impulse responses with a duration of zero that were convolved with the canonical haemodynamic

response function (HRF). Low-frequency drift was removed by temporal high pass filtering with a cut-off frequency of 1/128 Hz. Six head motion parameters were introduced in the GLM as nuisance regressors to account for the movement artefacts (Johnstone et al., 2006). Beta-coefficients reflecting an increase in the blood oxygenation level dependent (BOLD) signal in the experimental condition relative to the implicit baseline were converted into percent signal change (PSC) following the procedure recommended in (Poldrack, Mumford, Nichols, 2011, p.186). To that end, beta-coefficients for each condition were divided by the mean value of beta-coefficients for the constant term and multiplied by 100 and a scaling factor (SF) that is needed so that the peak of an isolated BOLD response equals to the 1 (SF = 0.21). Linear contrast, c, with a sum of positive terms equal to 1 was used to calculate the difference of the BOLD-signal percent changes in the conditions under study: 0.5 · ["A-P NoGo" + "A-A NoGo"] – 0.5 · ["A-P Go" + "A-A Go"] (the "NoGo vs. Go" comparison). The contrasts from 20 subjects were used as variables to verify the hypotheses on selective and non-selective response inhibition on the second level fMRI analysis. Only grey-matter voxels were included in the second level analysis. To that end, a mask was created based on the data of the segmentation of each subject's structural T1-images.

2.7. Verification of the hypotheses on selective and non-selective response inhibition.

The classical NHST approach estimates the probability (p-value) of obtaining actual data's best-fitting parameter value, β, or something more extreme under the null hypothesis that the experimental effect is $\theta = c\beta = 0$. One can never accept the null hypothesis using NHST because the p-value does not represent the probability of the null hypothesis and the probability an effect equals exactly zero is itself zero (Friston et al., 2002a). A non-significant result can be caused by two conditions (Dienes, 2014): 1) there is no effect, and our data are against the alternative hypothesis; or 2) our data are insufficient to distinguish alternative hypothesis from the null hypothesis, and we cannot confidently make any inference (low statistical power). When the obtained result is not significant, it is recommended to supply interval estimates, e.g., confidence intervals or highest density intervals (Mumford, 2012; Dienes, 2014; Kruschke, 2018), and use frequentist equivalence testing (Lakens, 2017) or Bayesian inference (Penny & Ridgway, 2013; Dienes, 2014; Kruschke, 2018). In present study, the Bayesian parameter inference was utilized (Friston & Penny, 2003; Penny & Ridgway, 2013; Kruschke, 2018).

The Bayesian approach estimates the posterior probability (PP) distribution of effect θ given the data using the likelihood and the prior knowledge. For the second level Bayesian analysis, SPM12 implements the global shrinkage prior (Friston & Penny, 2003). It represents a prior belief that, on average, in the whole brain, there is no global experimental effect. If the posterior probability of the effect exceeding the effect size threshold, γ, is greater than the

predefined probability threshold, α = 95%, then the hypothesis on the presence of "NoGo > Go" effect will be accepted:

$$PP_{NoGo>Go} = p(\theta > \gamma | Data) \geq \alpha.$$

If the effect value falls within the interval $[-\gamma; \gamma]$ with a probability of α = 0.95, then the hypothesis of null "NoGo = Go" effect will be accepted, supporting the practical equivalence (Kruschke, 2018) of the BOLD signal between compared conditions:

$$PP_{NoGo=Go} = p(-\gamma < \theta < \gamma | Data) \geq \alpha.$$

The hypothesis on the presence of "Go > NoGo" effect will be accepted if

$$PP_{Go>NoGo} = p(\theta < -\gamma | Data) \geq \alpha.$$

If none of the above criteria are satisfied, the data in particular voxel are insufficient to distinguish the null hypothesis from the alternative hypothesis ("low-confidence" voxels) (Magerkurth et al., 2015). For such voxels, it is impossible to make any inference confidently, so the power should be increased. In the current study, our conclusions were based only on voxels in which the effect size values (PSC) fell within or outside the ROPE with the PP > 0.95. Thus, for these voxels, the obtained data are sufficient to make an inference confidently.

Bayesian parameter inference with a zero effect size threshold corresponds to a false discovery rate correction for multiple comparisons within NHST framework (Genovese, Lazar & Nichols, 2002, Friston & Penny, 2003). In the SPM12, the group-level effect size threshold is set at one standard deviation of the prior variance of the contrast (prior SD_θ) by default (Friston & Penny, 2003). For the contrast under consideration, prior SD_θ = 0.11% of BOLD-signal. The interval $[-\gamma, \gamma]$ can be thought as the neuronal ''background noise level'' (Eickhoff et al., 2008) or as a region of practical equivalence (ROPE) that expresses which effect size values (PSC) are equivalent to the null value for current practical purposes (Kruschke, 2018, the "ROPE-only" decision rule). Additionally, the results of Bayesian parameter inference were compared with the results of classical inference using a FWE-corrected voxel-wise threshold of p < 0.05.

The hypothesis on the *selectivity* of response inhibition implies a selective increase in the neuronal activity in response to the presentation of NoGo stimuli compared to Go stimuli ("NoGo > Go", PP(θ > 0.11%) > 0.95). Regarding the hypothesis on the *non-selectivity* of response inhibition, Go and NoGo trials are expected to result in practically equivalent neuronal activity ("NoGo = Go", PP(-0.11% < θ < 0.11%) > 0.95) in the brain structures responsible for response inhibition (which were revealed by the current meta-analysis). For visualization purposes, posterior probabilities were converted to LogOdds = log(PP/(1-PP)) (LogOdds > 3 correspond to PP > 0.95). Anatomical localization of clusters was identified using xjView toolbox (http://www.alivelearn.net/xjview).

3. Results

As a result of a meta-analysis of 20 fMRI studies, we identified the brain structures that were characterized by increased neuronal activity demonstrated in the settings of equal probability Go and NoGo stimuli presentation compared to the control Go conditions. In such control conditions, the response inhibition process was not involved because the subjects were instructed always to press the button ("Go/NoGo > Go-control"). The set of revealed brain structures includes: (1) left dorsolateral prefrontal cortex (DLPFC), (2) left inferior parietal lobule (IPL), (3) left temporoparietal junction (TPJ), (4) bilateral inferior frontal gyrus (IFG) and anterior insula (also known as anterior insula/frontal operculum (AIFO)), (5) left premotor cortex (PMC) and frontal eye field (FEF), (6) bilateral anterior cingulate cortex (ACC) and supplementary motor area (SMA), and (7) bilateral thalamus (see Fig. 3, Table 1).

Figure 3. Results of the meta-analysis. The result of the meta-analysis of 20 fMRI studies using equal probability Go/NoGo tasks (contrast "Go/NoGo > Go-control").

Table 1. The result of the meta-analysis of 20 fMRI studies using equal probability Go/NoGo tasks (contrast "Go/NoGo > Go-control").

№	Cluster size, mm³	Peak MNI-coordinate, mm	Peak ALE-score	Anatomical localization (L – left, R – right hemisphere; BA – Brodmann area)
1	10160	42 34 32	0.0114	R: DLPFC (middle frontal gyrus), BA 8,9
2	8368	46 18 -10	0.0138	R: IFG/Anterior insula (AIFO), BA 13, 45, 47
3	4896	-34 24 0	0.0088	L: IFG/Anterior insula (AIFO), BA 13, 44, 45, 47
4	4176	60 -44 26	0.0090	R: TPJ (supramarginal gyrus and superior temporal gyrus), BA 22, 39, 40
5	3408	5 18 40	0.0103	R/L: SMA, ACC, BA 6, 8, 24, 32
6	1664	12 -8 10	0.0089	R: Thalamus
7	1584	42 6 40	0.0076	R: PMC, FEF (precentral gyrus, middle

				frontal gyrus), BA 6, 8, 9
8	1456	-12 -10 10	0.0084	L: Thalamus
9	1424	30 -62 38	0.0071	R: IPL (angular gyrus), BA 7, 39

Abbreviations: DLPFC – dorsolateral prefrontal cortex, IFG – inferior frontal gyrus, AIFO –

anterior insula/frontal operculum, TPJ – temporoparietal junction, SMA – supplementary motor

area, ACC – anterior cingulate cortex, PMC – premotor cortex, FEF – frontal eye field, IPL –

inferior parietal lobule.

Both classical NHST and Bayesian inference applied in the present fMRI study do not

reveal any significant increase in neuronal activity in NoGo trials compared to Go trials. Thus,

the "NoGo > Go" effect predicted by the hypothesis of selective response inhibition was not

observed. In addition, both statistical methods revealed similar activation patterns for the "Go >

NoGo" effect (Dice coefficient = 0.8), demonstrating the consistency between utilized Bayesian

parameter inference with $\gamma = 1$ prior SD_θ and LogOdds > 3 (PP > 0.95) thresholding approach

and NHST with pFWE<0.05 (see Fig. 4, and Tables S2 and S3 in the supplementary materials).

A) Classical voxel-wise inference

B) Bayesian parameter inference

Figure 4. Results of classical voxel-wise and Bayesian parameter inference on the obtained fMRI data. A) Classical NHST inference using a FWE-corrected voxel-wise threshold of p < 0.05. B) Bayesian parameter inference using the effect size threshold $\gamma = 1$ prior $SD_\theta = 0.11\%$, LogOdds > 3 (PP > 0.95). The "NoGo > Go" effect was not detected by any of the methods. Red colour depicts the "Go > NoGo" effect. Green colour depicts the "Go = NoGo" effect (practical equivalence of neuronal activity in Go and NoGo trials). White colour depicts A) non-significant voxels; B) "low-confidence" voxels.

Meanwhile, Bayesian analysis has made it possible to define the brain structures, wherein practical equivalence of neuronal activity was observed in Go and NoGo trials. The null "NoGo = Go" effect was revealed for a widely distributed set of regions throughout the whole brain surrounding clusters of activations revealed in the "Go > NoGo" comparison (see Fig. 4B). The "NoGo = Go" regions were separated from activation clusters by regions that consisted of "low confidence" voxels (Magerkurth et al., 2015). For "low confidence" voxels, the data obtained are insufficient to accept or reject the null hypothesis. The "NoGo = Go" area cannot be reasonably split into several parts such as activation clusters. Therefore, we limited ourselves to specifying the anatomic localization of brain areas where the "NoGo = Go" area overlaps with the clusters revealed by the current meta-analysis.

As it can be inferred from the revealed overlap between the results of the meta-analysis and the results of abovementioned Bayesian analysis, only a few of the brain structures demonstrated both increased activity in the "Go/NoGo > Go-control" meta-analytical comparison and the practical equivalence of the BOLD signal in the "NoGo vs. Go" comparison of the present fMRI data set. Overlap between both analyses was observed in (1) right DLPFC, (2) right IPL, (3) right TPJ, (4) left IFG and anterior insula (AIFO), (5) right PMC, and FEF (see Fig. 5, Table 2).

Figure 5. Overlap between meta-analysis and Bayesian meta-analysis. Overlap between null "NoGo = Go" effect regions and response inhibition brain structures revealed by the current meta-analysis. Abbreviations: IFG – inferior frontal gyrus, Ins – insula, DLPFC – dorsolateral prefrontal cortex, FEF – frontal eye field, PMC – premotor cortex, IPL – inferior parietal lobule, BA – Brodmann area, L – left, R – right.

Table 2. Overlap between null "NoGo = Go" effect regions revealed by Bayesian analysis of the obtained fMRI data and the results of the current meta-analysis.

№	Cluster size, mm^3	Centroid MNI-coordinates	Anatomical localization (L – left, R – right hemisphere; BA – Brodmann area)
1	1836	39 33 33	R: DLPFC, BA 9
2	1377	39 6 42	R: PMC, FEF, BA 6, 8
3	1215	30 -60 45	R: IPL
4	702	-33 21 -12	L: IFG
5	513	-39 27 0	L: Anterior insula
6	216	54 -42 15	R: TPJ
7	162	-27 27 0	L: Anterior insula

Abbreviations: DLPFC – dorsolateral prefrontal cortex, PMC – premotor cortex, FEF – frontal eye field, IPL – inferior parietal lobule, IFG – inferior frontal gyrus, TPJ – temporoparietal junction.

4. Discussion

The results of the present study confirmed the hypothesis of non-selective response inhibition at least in the context uncertainty associated with the equiprobable presentation of Go and NoGo stimuli. Bayesian analysis of the obtained fMRI data provides evidence of the practical equivalence of neuronal activity evoked by Go and NoGo stimuli in several brain structures. According to the results of our meta-analysis, some of these brain structures, including right DLPFC, IPL, and TPJ, left IFG (AIFO), right PMC, and FEF, were associated

with response inhibition. These structures were repeatedly found activated whenever the condition of equal probability Go and NoGo stimuli presentation was compared with the control Go condition ("Go/NoGo > Go control" contrast). In addition, we did not observe the "NoGo > Go" effect in the current study. This finding is consistent with the data of a number of previous event-related fMRI studies that did not find significant activity increase in NoGo trials compared to Go trials under similar comparisons (Lee et al., 2015, Di Russo et al., 2016, Criaud et al., 2017, see also random-effects analysis within visual and audio modalities in Laurens, Kiehl & Liddle, 2004).

However, the question is how precisely observed activity can be associated with response inhibition per se in those brain regions in which accepted null hypothesis overlapped with response inhibition brain regions resulted from the meta-analysis. The frontoparietal structures (DLPFC, IPL, TPJ) are engaged not only in Go/NoGo tasks but also in experiments examining task switching, resolution of cognitive conflicts, working memory, and attention focusing, i.e., external-directed activities ("multiple demand system" (Duncan, 2010); "task-general network" (Cole et al., 2014); "extrinsic mode network" (Hugdahl et al., 2015)). In addition, the most prominent feature of brain activity during Go/No Go tasks compared to diverse non-inhibitory tasks is a right-dominant activity in DLPFC (meta-analyses: Buchsbaum et al., 2005, Nee, Wager & Jonides, 2007, Levy & Wagner, 2011, Cieslik et al., 2015, Zhang, Geng & Lee, 2017). Among different response inhibition tasks, several meta-analyses showed that an action-restraint (Go/NoGo task) elicits stronger activation in the right DLPFC and parietal cortex compared to another inhibition process, namely, action cancellation (stop-signal task) (Swick, Ashley & Turken, 2011, Cieslik et al., 2015, Zhang, Geng & Lee, 2017). Furthermore, "response uncertainty" tasks modelled by equal probability Go/NoGo stimuli evoked more activity in the right DLPFC (near the cluster revealed in current study with coordinates [x=39 y=33 z=33], see Table 2) compared to Go/NoGo and stop-signal tasks with low probability of NoGo stimuli ("response override" tasks) (Levy & Wagner, 2011).

Similarly, in meta-analyses of Simmonds, Pekar & Mostofsky (2008) and Criaud & Boulinguez (2013), it was supposed that the increased activity in the right DLPFC observed during Go/NoGo tasks may be due to increased demands caused by working-memory and attentional load rather than the inhibition processes per se. The authors explained it by the fact that most of the fMRI studies used complex designs. Thus, it would be relatively more difficult for the subject to identify a NoGo stimulus, which was believed to increase the need for inhibitory control and consequently increase the brain inhibitory activity in NoGo trials. However, this confounding effect was controlled in the current fMRI study because we utilized

equally probable presentation of Go and NoGo stimuli and the performed meta-analysis included only studies using condition with equiprobable Go and NoGo stimuli presentation.

Another prefrontal structure, the right IFG (its opercular part, AIFO), which was revealed in the present study as related to non-selective response inhibition, is commonly associated with response inhibition (Aron, Robbins & Poldrack, 2014). Based on the study of patients with brain damage (Aron et al., 2003) and other neuroimaging studies using primarily the stop-signal tasks (Aron, Robbins & Poldrack, 2014), it was claimed that the right IFG represents a key node of the response inhibition brain system. Currently, this opinion is under active discussion (Mostofsky & Simmonds, 2008, Hampshire et al., 2010, Hampshire, 2015a, Hampshire & Sharp, 2015b, Swick, Ashley & Turken, 2008, Swick & Chatham, 2014). It is noted that Aron et al. predominantly relied upon studies using the stop-signal task and did not consider those using Go/NoGo tasks (Swick & Chatham, 2014). The studies by Swick, Ashley & Turken (2008) and Kramer et al. (2013) revealed worse performance in the Go/NoGo task when left (not right) IFG was damaged. According to these findings, the left IFG can also be involved in inhibition control during action restraint, which is additionally confirmed by our results.

Regarding the practically equivalent involvement of PMC in both Go and NoGo trials, it was previously demonstrated to participate in the planning and coordination of actions because its electrical stimulation results in involuntary motor actions (Desmurget et al., 2009). According to Duque et al. (2012), premotor cortex function is associated with inhibition of any premature actions and control of the time of action execution. The authors refer to this brain mechanism as "impulse control", i.e., a concept that is similar to non-selective response inhibition (Criaud et al., 2017).

In summary, it can be noted that the selectivity of response inhibition has been more extensively studied to date than its non-selectivity. A substantial body of literature was devoted to action cancelation studied with the use of stop-signal tasks with a low probability of the presentation of NoGo stimuli. However, over time, it has become apparent that under several conditions, non-selectivity of response inhibition can in principle occur. Studies on non-selectivity of response inhibition typically focus on the possible brain mechanisms for non-selective inhibition of motor responses in the settings of interference among multiple concurrent response options associated exclusively with NoGo stimuli (De Jong, Coles & Logan, 1995, Coxon, Stinear & Byblow, 2007, Aron & Verbruggen, 2008, Frank, 2006, Duque & Ivry, 2009, Duque et al., 2010, MacDonald et al., 2017). Moreover, the possibility of involvement of non-selective response inhibition not only for NoGo stimuli but also for Go stimuli was previously supposed by Criaud et al. (2017). In the current study, the practical equivalence of BOLD signals for NoGo and Go stimuli was demonstrated for a number of brain areas associated with response

inhibition as identified by the meta-analysis of fMRI studies. The results of the present study revealed in the settings of context uncertainty modelled by the equal probability of Go and NoGo trials proved the idea that response inhibition can act as a non-selective mechanism of action inhibition. Thus, one promising area of further research of brain mechanisms of response inhibition (and inhibitory control in general) is the study of the interplay between selective and non-selective inhibition as a function of the context uncertainty degree.

5. Conclusion

For the first time, combining a meta-analysis approach and second-level Bayesian analysis yielded results favouring the existence of non-selective response inhibition. The overlap between brain areas previously associated with response inhibition and brain areas demonstrating practical equivalence of neuronal activity in equiprobable Go and NoGo trials was observed in the right DLPFC, IPL, PMC and left IFG (AIFO). When a subject was waiting for an equiprobable Go or NoGo stimulus, a non-selective inhibitory control process was involved with any imperative stimulus, i.e., it occurred in Go trials and NoGo trials. In opposition to the model of selective response inhibition, this type of response inhibition prevented the performance of any premature motor actions and operated in non-selective, "global" mode. Its involvement is favoured by context uncertainty caused by the equally probable presentation of Go and NoGo stimuli. Upon the identification of a Go stimulus, inhibition is released, and the process of action execution is initiated, i.e., it acts as a gating mechanism for accessing a prepared motor program. We do not discard the opportunity of involvement of selective inhibitory mechanisms in a less uncertain context. Therefore, future research should address the issue of how brain mechanisms of selective and non-selective response inhibition are inter-related.

6. Acknowledgments

Current study in a part for Bayesian analysis of fMRI data was supported by the Russian Science Foundation grant #19-18-00454. The meta-analysis of Go/NoGo fMRI studies was performed within the state assignment of the Ministry of Education and Science of Russian Federation (theme number AAAA-A19-119101890066-2).

7. References

Albares, M., Lio, G., Criaud, M., Anton, J.-L., Desmurget, M., & Boulinguez, P. (2014). The dorsal medial frontal cortex mediates automatic motor inhibition in uncertain contexts: Evidence from combined fMRI and EEG studies. *Human Brain Mapping*, *35*(11), 5517–5531. https://doi.org/10.1002/hbm.22567

Aron, Adam R. (2011). From Reactive to Proactive and Selective Control: Developing a Richer Model for Stopping Inappropriate Responses. *Biological Psychiatry*, *69*(12), e55–e68. https://doi.org/10.1016/j.biopsych.2010.07.024

Aron, Adam R., Fletcher, P. C., Bullmore, E. T., Sahakian, B. J., & Robbins, T. W. (2003). Stop-
signal inhibition disrupted by damage to right inferior frontal gyrus in humans. *Nature*
Neuroscience, *6*(2), 115–116. https://doi.org/10.1038/nn1003

Aron, Adam R., Robbins, T. W., & Poldrack, R. A. (2014). Inhibition and the right inferior
frontal cortex: one decade on. *Trends in Cognitive Sciences*, *18*(4), 177–185.
https://doi.org/10.1016/j.tics.2013.12.003

Aron, Adam R., & Verbruggen, F. (2008). Stop the Presses. *Psychological Science*, *19*(11),
1146–1153. https://doi.org/10.1111/j.1467-9280.2008.02216.x

Boulinguez, P., Jaffard, M., Granjon, L., & Benraiss, A. (2008). Warning Signals Induce
Automatic EMG Activations and Proactive Volitional Inhibition: Evidence From Analysis of
Error Distribution in Simple RT. *Journal of Neurophysiology*, *99*(3), 1572–1578.
https://doi.org/10.1152/jn.01198.2007

Braver, T. S. (2012). The variable nature of cognitive control: a dual mechanisms framework.
Trends in Cognitive Sciences, *16*(2), 106–113. https://doi.org/10.1016/j.tics.2011.12.010

Buchsbaum, B. R., Greer, S., Chang, W.-L., & Berman, K. F. (2005). Meta-analysis of
neuroimaging studies of the Wisconsin Card-Sorting task and component processes. *Human*
Brain Mapping, *25*(1), 35–45. https://doi.org/10.1002/hbm.20128

Chevrier, A. D., Noseworthy, M. D., & Schachar, R. (2007). Dissociation of response inhibition
and performance monitoring in the stop signal task using event-related fMRI. *Human Brain*
Mapping, *28*(12), 1347–1358. https://doi.org/10.1002/hbm.20355

Cieslik, E. C., Mueller, V. I., Eickhoff, C. R., Langner, R., & Eickhoff, S. B. (2015). Three key
regions for supervisory attentional control: Evidence from neuroimaging meta-analyses.
Neuroscience & Biobehavioral Reviews, *48*, 22–34.
https://doi.org/10.1016/j.neubiorev.2014.11.003

Cole, M. W., Bassett, D. S., Power, J. D., Braver, T. S., & Petersen, S. E. (2014). Intrinsic and
Task-Evoked Network Architectures of the Human Brain. *Neuron*, *83*(1), 238–251.
https://doi.org/10.1016/j.neuron.2014.05.014

Coxon, J. P., Stinear, C. M., & Byblow, W. D. (2007). Selective Inhibition of Movement.
Journal of Neurophysiology, *97*(3), 2480–2489. https://doi.org/10.1152/jn.01284.2006

Criaud, M., & Boulinguez, P. (2013). Have we been asking the right questions when assessing
response inhibition in go/no-go tasks with fMRI? A meta-analysis and critical review.
Neuroscience & Biobehavioral Reviews, *37*(1), 11–23.
https://doi.org/10.1016/j.neubiorev.2012.11.003

Criaud, M., Longcamp, M., Anton, J.-L., Nazarian, B., Roth, M., Sescousse, G., … Boulinguez,
P. (2017). Testing the physiological plausibility of conflicting psychological models of response
inhibition: A forward inference fMRI study. *Behavioural Brain Research*, *333*, 192–202.
https://doi.org/10.1016/j.bbr.2017.06.030

De Jong, R., Coles, M. G. H., & Logan, G. D. (1995). Strategies and mechanisms in nonselective
and selective inhibitory motor control. *Journal of Experimental Psychology: Human Perception*
and Performance, *21*(3), 498–511. https://doi.org/10.1037//0096-1523.21.3.498

Dienes, Z. (2014). Using Bayes to get the most out of non-significant results. *Frontiers in*
Psychology, *5*. https://doi.org/10.3389/fpsyg.2014.00781

Desmurget, M., Reilly, K. T., Richard, N., Szathmari, A., Mottolese, C., & Sirigu, A. (2009).
Movement Intention After Parietal Cortex Stimulation in Humans. *Science*, *324*(5928), 811–813.
https://doi.org/10.1126/science.1169896

Di Russo, F., Lucci, G., Sulpizio, V., Berchicci, M., Spinelli, D., Pitzalis, S., & Galati, G. (2016).
Spatiotemporal brain mapping during preparation, perception, and action. *NeuroImage*, *126*, 1–
14. https://doi.org/10.1016/j.neuroimage.2015.11.036

Duncan, J. (2010). The multiple-demand (MD) system of the primate brain: mental programs for
intelligent behaviour. *Trends in Cognitive Sciences*, *14*(4), 172–179.
https://doi.org/10.1016/j.tics.2010.01.004

Dunovan, K., Lynch, B., Molesworth, T., & Verstynen, T. (2015). Competing basal ganglia
pathways determine the difference between stopping and deciding not to go. *ELife*, *4*.
https://doi.org/10.7554/eLife.08723

Duque, J., & Ivry, R. B. (2009). Role of Corticospinal Suppression during Motor Preparation.
Cerebral Cortex, *19*(9), 2013–2024. https://doi.org/10.1093/cercor/bhn230

Duque, J., Labruna, L., Verset, S., Olivier, E., & Ivry, R. B. (2012). Dissociating the Role of
Prefrontal and Premotor Cortices in Controlling Inhibitory Mechanisms during Motor
Preparation. *Journal of Neuroscience*, *32*(3), 806–816.
https://doi.org/10.1523/JNEUROSCI.4299-12.2012

Duque, J., Lew, D., Mazzocchio, R., Olivier, E., & Ivry, R. B. (2010). Evidence for Two
Concurrent Inhibitory Mechanisms during Response Preparation. *Journal of Neuroscience*,
30(10), 3793–3802. https://doi.org/10.1523/JNEUROSCI.5722-09.2010

Eickhoff, S. B., Grefkes, C., Fink, G. R., & Zilles, K. (2008). Functional Lateralization of Face,
Hand, and Trunk Representation in Anatomically Defined Human Somatosensory Areas.
Cerebral Cortex, *18*(12), 2820–2830. https://doi.org/10.1093/cercor/bhn039

Eickhoff, Simon B., Nichols, T. E., Laird, A. R., Hoffstaedter, F., Amunts, K., Fox, P. T., …
Eickhoff, C. R. (2016). Behavior, sensitivity, and power of activation likelihood estimation
characterized by massive empirical simulation. *NeuroImage*, *137*, 70–85.
https://doi.org/10.1016/j.neuroimage.2016.04.072

Erika-Florence, M., Leech, R., & Hampshire, A. (2014). A functional network perspective on
response inhibition and attentional control. *Nature Communications*, *5*(1).
https://doi.org/10.1038/ncomms5073

Frank, M. J. (2006). Hold your horses: A dynamic computational role for the subthalamic
nucleus in decision making. *Neural Networks*, *19*(8), 1120–1136.
https://doi.org/10.1016/j.neunet.2006.03.006

Friedman, N. P., Miyake, A., Young, S. E., DeFries, J. C., Corley, R. P., & Hewitt, J. K. (2008). Individual differences in executive functions are almost entirely genetic in origin. *Journal of Experimental Psychology: General*, *137*(2), 201–225. https://doi.org/10.1037/0096-3445.137.2.201

Friston, K. J., Glaser, D. E., Henson, R. N. A., Kiebel, S., Phillips, C., & Ashburner, J. (2002b). Classical and Bayesian Inference in Neuroimaging: Applications. *NeuroImage*, *16*(2), 484–512. https://doi.org/10.1006/nimg.2002.1091

Friston, K. J., & Penny, W. (2003). Posterior probability maps and SPMs. *NeuroImage*, *19*(3), 1240–1249. https://doi.org/10.1016/S1053-8119(03)00144-7

Friston, K. J., Penny, W., Phillips, C., Kiebel, S., Hinton, G., & Ashburner, J. (2002a). Classical and Bayesian Inference in Neuroimaging: Theory. *NeuroImage*, *16*(2), 465–483. https://doi.org/10.1006/nimg.2002.1090

Ganos, C., Rothwell, J., & Haggard, P. (2018). Voluntary inhibitory motor control over involuntary tic movements. *Movement Disorders*, *33*(6), 937–946. https://doi.org/10.1002/mds.27346

Genovese, C. R., Lazar, N. A., & Nichols, T. (2002). Thresholding of Statistical Maps in Functional Neuroimaging Using the False Discovery Rate. *NeuroImage*, *15*(4), 870–878. https://doi.org/10.1006/nimg.2001.1037

Hampshire, A. (2015a). Putting the brakes on inhibitory models of frontal lobe function. *NeuroImage*, *113*, 340–355. https://doi.org/10.1016/j.neuroimage.2015.03.053

Hampshire, A., Chamberlain, S. R., Monti, M. M., Duncan, J., & Owen, A. M. (2010). The role of the right inferior frontal gyrus: inhibition and attentional control. *NeuroImage*, *50*(3), 1313–1319. https://doi.org/10.1016/j.neuroimage.2009.12.109

Hampshire, A., & Sharp, D. J. (2015b). Contrasting network and modular perspectives on inhibitory control. *Trends in Cognitive Sciences*, *19*(8), 445–452. https://doi.org/10.1016/j.tics.2015.06.006

Hsieh, S., Wu, M., & Tang, C.-H. (2016). Adaptive Strategies for the Elderly in Inhibiting Irrelevant and Conflict No-Go Trials while Performing the Go/No-Go Task. *Frontiers in Aging Neuroscience*, *7*. https://doi.org/10.3389/fnagi.2015.00243

Hugdahl, K., Raichle, M. E., Mitra, A., & Specht, K. (2015). On the existence of a generalized non-specific task-dependent network. *Frontiers in Human Neuroscience*, *9*. https://doi.org/10.3389/fnhum.2015.00430

Isoda, M., & Hikosaka, O. (2011). Cortico-basal ganglia mechanisms for overcoming innate, habitual and motivational behaviors. *European Journal of Neuroscience*, *33*(11), 2058–2069. https://doi.org/10.1111/j.1460-9568.2011.07698.x

Jaffard, M., Benraiss, A., Longcamp, M., Velay, J.-L., & Boulinguez, P. (2007). Cueing method biases in visual detection studies. *Brain Research*, *1179*, 106–118. https://doi.org/10.1016/j.brainres.2007.08.032

Jaffard, M., Longcamp, M., Velay, J.-L., Anton, J.-L., Roth, M., Nazarian, B., & Boulinguez, P.
(2008). Proactive inhibitory control of movement assessed by event-related fMRI. NeuroImage,
42(3), 1196–1206. https://doi.org/10.1016/j.neuroimage.2008.05.041

Johnstone, T., Ores Walsh, K. S., Greischar, L. L., Alexander, A. L., Fox, A. S., Davidson, R. J.,
& Oakes, T. R. (2006). Motion correction and the use of motion covariates in multiple-subject
fMRI analysis. *Human Brain Mapping*, 27(10), 779–788. https://doi.org/10.1002/hbm.20219

Krämer, U. M., Solbakk, A.-K., Funderud, I., Løvstad, M., Endestad, T., & Knight, R. T. (2013).
The role of the lateral prefrontal cortex in inhibitory motor control. *Cortex*, 49(3), 837–849.
https://doi.org/10.1016/j.cortex.2012.05.003

Kropotov, J. D. (2016). *Functional Neuromarkers for Psychiatry: Applications for Diagnosis
and Treatment* (1st ed.). Cambridge, MA: Academic Press.

Kropotov, J. D., Ponomarev, V. A., Hollup, S., & Mueller, A. (2011). Dissociating action
inhibition, conflict monitoring and sensory mismatch into independent components of event
related   potentials   in   GO/NOGO   task.   NeuroImage,   57(2),   565–575.
https://doi.org/10.1016/j.neuroimage.2011.04.060

Kropotov, J. D., & Ponomarev, V. A. (2015). Differentiation of neuronal operations in latent
components of event-related potentials in delayed match-to-sample tasks. *Psychophysiology*,
52(6), 826–838. https://doi.org/10.1111/psyp.12410

Kruschke, J. K. (2018). Rejecting or Accepting Parameter Values in Bayesian Estimation.
Advances   in   Methods   and   Practices   in   Psychological   Science,   1(2),   270–280.
https://doi.org/10.1177/2515245918771304

Kruschke, J. K., & Liddell, T. M. (2015). The Bayesian New Statistics: Two Historical Trends
Converge. *SSRN Electronic Journal*. https://doi.org/10.2139/ssrn.2606016

Lakens, D. (2017). Equivalence Tests. *Social Psychological and Personality Science*, 8(4), 355–
362. https://doi.org/10.1177/1948550617697177

Laurens, K. R., Kiehl, K. A., & Liddle, P. F. (2004). A supramodal limbic-paralimbic-
neocortical network supports goal-directed stimulus processing. *Human Brain Mapping*, 24(1),
35–49. https://doi.org/10.1002/hbm.20062

Lavric, A., Pizzagalli, D. A., & Forstmeier, S. (2004). When 'go' and 'nogo' are equally
frequent: ERP components and cortical tomography. *European Journal of Neuroscience*, 20(9),
2483–2488. https://doi.org/10.1111/j.1460-9568.2004.03683.x

Lee, K.-H., Pluck, G., Lekka, N., Horton, A., Wilkinson, I. D., & Woodruff, P. W. R. (2015).
Self-harm in schizophrenia is associated with dorsolateral prefrontal and posterior cingulate
activity. *Progress in Neuro-Psychopharmacology and Biological Psychiatry*, 61, 18–23.
https://doi.org/10.1016/j.pnpbp.2015.03.005

Levy, B. J., & Wagner, A. D. (2011). Cognitive control and right ventrolateral prefrontal cortex:
reflexive reorienting, motor inhibition, and action updating. *Annals of the New York Academy of
Sciences*, 1224(1), 40–62. https://doi.org/10.1111/j.1749-6632.2011.05958.x

Logan, G. D., Van Zandt, T., Verbruggen, F., & Wagenmakers, E.-J. (2014). On the ability to
inhibit thought and action: General and special theories of an act of control. *Psychological*
Review, *121*(1), 66–95. https://doi.org/10.1037/a0035230

MacDonald, H. J., McMorland, A. J. C., Stinear, C. M., Coxon, J. P., & Byblow, W. D. (2017).
An Activation Threshold Model for Response Inhibition. *PLOS ONE*, *12*(1), e0169320.
https://doi.org/10.1371/journal.pone.0169320

Magerkurth, J., Mancini, L., Penny, W., Flandin, G., Ashburner, J., Micallef, C., … Weiskopf,
N. (2015). Objective Bayesian fMRI analysis - a pilot study in different clinical environments.
Frontiers in Neuroscience, *9*. https://doi.org/10.3389/fnins.2015.00168

Meffert, H., Hwang, S., Nolan, Z. T., Chen, G., & Blair, J. R. (2016). Segregating attention from
response control when performing a motor inhibition task. *NeuroImage*, *126*, 27–38.
https://doi.org/10.1016/j.neuroimage.2015.11.029

Mostofsky, S. H., & Simmonds, D. J. (2008). Response Inhibition and Response Selection: Two
Sides of the Same Coin. *Journal of Cognitive Neuroscience*, *20*(5), 751–761.
https://doi.org/10.1162/jocn.2008.20500

Mumford, J. A. (2012). A power calculation guide for fMRI studies. *Social Cognitive and*
Affective Neuroscience, *7*(6), 738–742. https://doi.org/10.1093/scan/nss059

Nee, D. E., Wager, T. D., & Jonides, J. (2007). Interference resolution: Insights from a meta-
analysis of neuroimaging tasks. *Cognitive, Affective, & Behavioral Neuroscience*, *7*(1), 1–17.
https://doi.org/10.3758/CABN.7.1.1

Neumann, J., & Lohmann, G. (2003). Bayesian second-level analysis of functional magnetic
resonance images. *NeuroImage*, *20*(2), 1346–1355. https://doi.org/10.1016/S1053-
8119(03)00443-9

Oldfield, R. C. (1971). The assessment and analysis of handedness: The Edinburgh inventory.
Neuropsychologia, *9*(1), 97–113. https://doi.org/10.1016/0028-3932(71)90067-4

Poldrack, R. A., Mumford, J. A., & Nichols, T. E. (2011). Handbook of Functional MRI Data
Analysis. Cambridge, United Kingdom: Cambridge University Press.

Simmonds, D. J., Pekar, J. J., & Mostofsky, S. H. (2008). Meta-analysis of Go/No-go tasks
demonstrating that fMRI activation associated with response inhibition is task-dependent.
Neuropsychologia, *46*(1), 224–232. https://doi.org/10.1016/j.neuropsychologia.2007.07.015

Swick, D., Ashley, V., & Turken, A. U. (2008). Left inferior frontal gyrus is critical for response
inhibition. *BMC Neuroscience*, *9*(1), 102. https://doi.org/10.1186/1471-2202-9-102

Swick, D., Ashley, V., & Turken, A. U. (2011). Are the neural correlates of stopping and not
going identical? Quantitative meta-analysis of two response inhibition tasks. *NeuroImage*, *56*(3),
1655–1665. https://doi.org/10.1016/j.neuroimage.2011.02.070

Swick, D., & Chatham, C. H. (2014). Ten years of inhibition revisited. *Frontiers in Human*
Neuroscience, *8*. https://doi.org/10.3389/fnhum.2014.00329

Verbruggen, F., Best, M., Bowditch, W. A., Stevens, T., & McLaren, I. P. L. (2014). The
inhibitory   control   reflex.   *Neuropsychologia*,   *65*,   263–278.
https://doi.org/10.1016/j.neuropsychologia.2014.08.014

Verbruggen, F., & Logan, G. D. (2008). Response inhibition in the stop-signal paradigm. *Trends*
in Cognitive Sciences, *12*(11), 418–424. https://doi.org/10.1016/j.tics.2008.07.005

Vink, M., Kaldewaij, R., Zandbelt, B. B., Pas, P., & du Plessis, S. (2015). The role of stop-signal
probability and expectation in proactive inhibition. *European Journal of Neuroscience*, *41*(8),
1086–1094. https://doi.org/10.1111/ejn.12879

Vink, M., Zandbelt, B. B., Gladwin, T., Hillegers, M., Hoogendam, J. M., van den Wildenberg,
W. P. M., … Kahn, R. S. (2014). Frontostriatal activity and connectivity increase during
proactive inhibition across adolescence and early adulthood. *Human Brain Mapping*, *35*(9),
4415–4427. https://doi.org/10.1002/hbm.22483

Woolrich, M. W., Behrens, T. E. J., Beckmann, C. F., & Smith, S. M. (2005). Mixture models
with adaptive spatial regularization for segmentation with an application to FMRI data. *IEEE*
Transactions on Medical Imaging, *24*(1), 1–11. https://doi.org/10.1109/TMI.2004.836545

Zhang, R., Geng, X., & Lee, T. M. C. (2017). Large-scale functional neural network correlates of
response inhibition: an fMRI meta-analysis. *Brain Structure and Function*, *222*(9), 3973–3990.
https://doi.org/10.1007/s00429-017-1443-x