1

Activation of G-protein coupled estradiol receptor 1 in the dorsolateral striatum attenuates

preference for cocaine and saccharin in male but not female rats

Authors:

Jacqueline A. Quigley, M.S.[1] and Jill B. Becker, Ph.D.[1,2]

[1]Psychology Department, [2]Molecular and Behavioral Neuroscience Institute,

University of Michigan, Ann Arbor MI, 48109 USA

Corresponding Author:

Jill B. Becker

205 Zina Pitcher Place, Ann Arbor, MI 48109

734-763-4363

jbbecker@umich.edu

Abstract

There are sex differences in the response to psychomotor stimulants, where females exhibit a greater response than males, due to the presence of the gonadal hormone estradiol (E2). Extensive research has shown that E2 enhances drug-seeking and the rewarding properties of cocaine for females. The role of E2 in male drug-seeking, however, is not well understood. The current study investigated pharmacological manipulation of E2 receptors in the dorsolateral striatum (DLS) on preference for cocaine in gonad-intact male and female rats. In males, activation of G-protein coupled E2 receptor 1 (GPER1), via administration of ICI 182,780 or G1, attenuated conditioned place preference for 10mg/kg cocaine, while inhibition of GPER1, via G15, enhanced preference at a 5mg/kg cocaine dose. Similarly, GPER1 activation, via G1, prevented males from forming a preference for 0.1% saccharin (SACC) versus plain water. Surprisingly, activation of GPER1 did not alter preference for cocaine or SACC in females. These studies also examined the quantity of E2 receptor mRNA in the dorsal striatum, using qPCR. No sex differences in relative mRNA expression of ERα, ERβ, and GPER1 were observed. However, there was greater GPER1 mRNA, relative to ERα and ERβ, in both males and females. The results presented here indicate that E2, acting via GPER1, may be protective against drug preference in male rats.

Introduction

More men report abusing cocaine and qualify for cocaine use disorder, however,
cocaine use in women is gradually increasing and there is growing evidence to suggest that
women are more vulnerable to addiction [1]. The time of first drug use to admission for addiction
treatment is typically shorter for women than men [2]. When entering treatment, women also
present with greater social, behavioral, and psychological symptoms related to substance use
disorder, despite having abused drugs for a shorter period of time [3,4]. These women are also
taking greater amount of drug when entering treatment and report experiencing enhanced
cravings compared to their male counterparts [5,6]. Historically, we have attributed sex-specific
behaviors related to addiction as being mediated by cultural influences combined with
differences in neurobiological function between males and females, including a sex-specific role
of the gonadal hormone estradiol (E2) that enhances drug-taking in females [7,8].

Research into the biological bases for sex differences in cocaine addiction has focused
on how E2 alters the rewarding properties of cocaine in females. Clinical models have found
that when E2 levels are high, women report an enhanced euphoria or "high" when abusing
smoked cocaine [9]. Similarly, female rats show enhanced cocaine craving and motivation for
cocaine when E2 levels are highest [10–13]. While these data suggest that E2 plays a major role in
facilitating motivation and other behaviors in females, the role of E2 in modulating males'
addiction-like behaviors should also be understood. Testosterone produced by the male testes
is aromatized to E2 in the brain and periphery to act on E2 receptors in the male brain.

E2 acts by binding to receptors alpha (ERα), beta (ERβ), and G-protein coupled E2
receptor 1 (GPER1). The cellular location of traditional E2 receptors α and β are either in the
cell nucleus, or after palmitoylation, on the extracellular membrane in association with caveolin
proteins [14–16]. GPER1 activates intracellular signaling cascades mediated by cAMP, ERK, and
PI3K and localized on the plasma membrane and in the endoplasmic reticulum [17]. In females,
ERα and GPER1 in the dorsal striatum are localized to GABAergic medium spiny neurons with

recurrent collaterals onto dopamine (DA) terminals, as well as cholinergic interneurons and glia

[18,19]. The localization of ERα and GPER1 in the dorsal striatum of males remains unknown.

For males and females alike, the rewarding effects of cocaine are attributed to the drug's

direct effects on DA reuptake in the striatum [20,21]. However, there are sex differences in the

effects of cocaine on increases in extracellular DA and receptor binding that are driven by E2 in

the dorsal striatum. In females, E2 binding to GABAergic neurons decreases GABA release and

this disinhibits DA terminals and ultimately increases stimulated DA release locally [16,22]. This E2-

induced increase in DA release is associated with an enhancement of the effects of cocaine and

other psychostimulants on drug-induced behaviors, such as behavioral sensitization [23–26].

Previous research has not found that E2 has similar effects on cocaine-induced increases in DA

in males [23]. Finally, the effects of E2 in dorsolateral striatum (DLS) are mediated by mGluR

signaling in females [24]

There is extensive support for E2 in mediating drug abuse liability in females but a lack

of attention to understanding the role of E2 in males. There is recent evidence from gene

knockout studies, however, that GPER1 could be playing a modulatory role in the preference for

morphine in males [27]. Given that previous research has highly implicated the DLS as an

important region for studying the effects of E2 on addiction-like behaviors in females, the current

set of studies were deigned to investigate how intra-DLS GPER1 modulates preference for

rewarding stimuli in males as well as females. A cocaine conditioned place preference (CPP)

and saccharin (SACC) two-bottle choice behavioral paradigms were used to assess how

GPER1 activation or inhibition alters preference for rewarding stimuli. Finally, the current study

also used qPCR to determine relative mRNA levels of ERs in the dorsal striatum of both males

and females.

Materials and Methods

Animals

A total of 62 male and 46 female gonad-intact Sprague-Dawley rats were used in the

current set of experiments, as detailed in Figure 1A. Animals were ordered from Charles River

Breeding Laboratory (Portage, MI, USA) and were approximately 75 days old on arrival.

Animals were maintained on a 14:10 light/dark cycle in a temperature-controlled climate of 72°F

± 2°F, in ventilated laboratory cages. Rats had ad libitum access to water and

phytoestrogen-free rat chow (2017 Teklad Global, 14% protein rodent maintenance diet, Harlan

rat chow; Harlan Teklad, Madison, WI, USA). Animals were initially housed in same-sex pairs

until undergoing surgery, after which they were subsequently housed individually. All animals

were weighed daily to determine good health and at this time, females were also vaginally

lavaged to determine stage of estrus. All animal care and experimental procedures were

carried out in accordance with the National Institutes of Health guidelines on laboratory animal

use and care, using a protocol approved by University of Michigan Institutional Use and Care of

Animals Committee.

Stereotaxic Surgery and Drug Preparation

One week after arriving in the laboratory, rats underwent surgery for the implantation of

bi-lateral guide cannula aimed at the DLS (AP: +0.4 ML: +/-3.6 DV: -4.0). On the day of surgery,

rats were injected with carprofen (5mg/kg s.c.) and 30 minutes later were anesthetized with

ketamine (50mg/kg i.p.) and dexmedetomidine (0.25mg/kg i.p.), then prepared in a stereotaxic

frame. At the conclusion of the surgery, rats were given atipamezole hydrochloride (0.5mg/kg

i.p.) and 3ml 0.9% saline (s.c.). Every 24 hours for three days post-surgery, rats were given

carprofen (5mg/kg s.c.) prophylactically for post-operative pain. No animal underwent behavioral

testing for at least 7 days after surgery.

During surgery, 33-gauge solid stylets were inserted into the 26-gauge hollow guide

cannula that were fixed on animals' skull. These stylets were flush with the bottom of the guide

cannula and did not protrude into the brain. Treatment conditions were randomly assigned to

animals prior to behavioral testing. Control animals received 100% cholesterol (CHOL) and

experimental animals received either 10% ICI (ERα/ERβ antagonist; GPER1 agonist), G1

(agonist targeting GPER1) or G15 (antagonist targeting GPER1) dissolved in Cholesterol

(Control) via stylets, which protruded from the guide cannula by 1mm and delivered treatment

directly into the DLS. Treatment stylets were prepared as previously described [28]. In order to

insert stylets, rats were briefly anesthetized with 5% isoflurane.

Hollow guides and interlocking treatment stylets were manufactured by and purchase

from P1 Technologies. Drugs were obtained from the following sources: ICI182,780 (ICI) (Santa

Cruz Biotechnology, purity ≥ 98%); G1 (Cayman Chemical, purity ≥ 98%); G15 (Cayman

Chemical, purity ≥ 95%); Cholesterol (Santa Cruz Biotechnology, purity ≥ 92%)

Conditioned Place Preference (CPP)

Animals were tested on a CPP paradigm that took place over 10 consecutive days, as

illustrated in Figure 2 A. The CPP apparatus consisted of two side chambers (15.5 inches x 12

inches) and a center neutral chamber (15.5 inches x 7.5 inches). On day 1 (pre-test), rats were

placed in the novel chamber and were allowed to move freely between all compartments for 30

minutes. Immediately following pre-test session, treatment stylets were inserted. For eight days

thereafter, animals were trained to associate each of the three chambers with another stimulus

(drug paired; neutral; vehicle paired). Each conditioning session began with a 10-minute

habituation in the center neutral chamber. Animals were then removed from the chamber and

received an intra-peritoneal (i.p.) injection of either cocaine or vehicle and immediately placed in

one of the side chambers (drug paired; vehicle paired) for 30 minutes. Animals were conditioned

to each stimulus (drug or vehicle) 4 times each, every other day. Which treatment animals

received first was counterbalanced. On day 10 (Test), rats were placed in the three-

compartment chamber and were again allowed to move freely between all compartments for 30

minutes.

The side in which animals spent the most time during pre-test was treated as their

"preferred chamber", which differed for each animal. A biased design was utilized, so that each

individual animal's initially preferred chamber was paired with saline, and their initially un-

preferred chamber was paired with cocaine for conditioning. ANYMAZE tracking software

(Stoelting Co., Wood Dale, IL 60191) was utilized to track the amount of time spent in each

chamber.

Either a 5mg/kg or 10mg/kg cocaine dose was utilized during conditioning in order to be

able to test both an increase and decrease in cocaine CPP after ER manipulation. For example,

both males and females acquire cocaine CPP at a 10mg/kg conditioning dose therefore we

utilized that dose when investigating a potential decrease in CPP (i.e., ICI and G1). When

investigating whether there was an increase cocaine CPP in males, we used a dose that did not

generally produce a CPP, 5mg/kg, which allowed us to identify a potential increase in CPP (i.e.,

G15) without a ceiling effect. This dose of cocaine did produce CPP for cocaine in females (data

not shown), so we did not test the effects of G15 in females with 5mg/kg cocaine.

Two Bottle Choice Experiment (SACC versus H_2O)

Animals were tested on a two-bottle choice paradigm that took place over 12

consecutive days, as illustrated in Figure 1 B. Briefly, on days 1-4, two bottles both containing

water were available on each animal's home cage to determine that were was no significant

difference between total liquid intake between animals. During days 5-12, one bottle contained

water and the second bottle contained 0.1% saccharin (Sigma-Aldrich; purity ≥ 92%) dissolved

in water. Placement of the bottle (left versus right) was switched daily to account for a potential

side of cage preference. On day 5, four hours before the SACC bottle was introduced, stylets

containing either 10% G1 in cholesterol or cholesterol alone were inserted and remained in

place for the duration of the experiment. Once daily, one hour prior to the start of dark cycle,

bottles were removed from home cages, weighed, and refilled.

Euthanasia and Tissue Preparation

Animals received 0.5 ml of Sodium Pentobarbital (i.p). Once the animal was fully

sedated, it was perfused transcardially with 0.1M phosphate buffered saline followed by 4%

paraformaldehyde. The brain of each rat was also dissected and post-fixed in 4%

paraformaldehyde for 24 hours and afterwards stored in 10% sucrose. Brains were sliced on

either a microtome or cryostat in 60-micron sections then were mounted on slides, stained with

cresyl violate and cover slipped. Sections were analyzed for accurate guide cannulae

placements, depicted in Table 1, by an observer blind to experimental conditions. Only animals

that had accurate cannulae placements are shown and were included in the final analysis. For

animals in the H_2O versus SACC experiment, females' ovaries and uterus and males' testes

and vas deferens were dissected and weighed after the animal was perfused as a proxy to

determine if G1 treatment in the brain affected peripheral gonadal tissues.

Quantitative Polymerase Chain Reaction (qPCR)

Naïve gonad-intact male and female rats were given 0.5 ml of FatalPlus (i.p.; 195 mg

sodium pentobarbital; Vortech Pharmaceuticals, Ltd; Dearborn, MI). Once the animal was fully

anesthetized the brain was rapidly removed and placed into ice-cold saline. The dorsal striatum

from both hemispheres was micro-dissected from each animal and stored at -80°C for later

processing.

Tissue was extracted using a phenol-chloroform reaction using Trizol (Cat. No. 97064-

950, Amresco) as the lysis reagent. Next, a QuantiTect® Reverse Transcription Kit was used for

genomic DNA wipeout and cDNA synthesis (Cat. No. 205314, Qiagen). Relative gene

expression was measured using a RealMasterMix™ Fast SYBR Kit (Cat. No. A25742, Applied

Biosystems). Primers (not designed to determine specific splice variants, but to detect all

variants) were purchased from Qiagen for: ERα (Cat. No. QT00386925, Qiagen Primer;

reference sequence NM_012689), ERβ (Cat. No. QT00190113, Qiagen Primer; reference

sequence NM_012754), and GPER1 (Cat. No. QT00376943, Qiagen Primer; reference

sequence NM_133573). These genes were compared against the housekeeping gene HPRT1

(Cat No. QT00199640, Qiagen Primer) and relative gene expression was quantified using the

2^ddCT method. Samples were run in triplicates at the Biomedical Research Core Facilities at

the University of Michigan (https://brcf.medicine.umich.edu/cores/advanced-

genomics/technologies/real-time-pcr/).

Statistical Analysis

CPP data were analyzed using 2-way ANOVAs. In the case of a significant interaction, a

Bonferroni correction was used for multiple comparisons. For males and females independently,

CPP data were analyzed by time spent in the drug-paired chamber (pre-test versus test)

between treatment conditions, within each sex. Behavioral testing for males and females was

not done simultaneously therefore, we did not compare them statistically.

Daily preference of 0.1% SACC versus water was calculated as a percentage: (0.1%

SACC consumed/(0.1%SACC + Water consumed))*100. An unpaired t-test was performed to

identify treatment group differences in preference score. Unpaired t-tests were performed to

determine if gonad weights were different between G1 or CHOL treated males or females.

Relative mRNA expression of ERα, ERβ, and GPER1 was analyzed between sexes.

Unpaired t tests were performed to identify sex differences in ERβ and GPER1 expression

within the dorsal striatum. Nonparametric testing was conducted for ERα because data were not

normally distributed for males.

All statistical analyses were performed using GraphPad Prism v8.0 and IBM SPSS

Statistics v27.0. All data sets were tested for normality and equal variance between groups.

Effect sizes are reported as partial eta ($n^{2)}$ or partial eta-squared (n^2p) for F-tests and Cohen's d

(d) for t-tests. The threshold for significance in all experiments was set to $p < 0.05$ and sample

sizes per experiment were determined based on pilot studies. Animals were excluded from

statistical analyses if the placements of the guide cannula were off target in both the CPP and

SACC versus H_2O experiments (< 5% of total animals were excluded).

Results

Effects of ER manipulation on cocaine CPP in males

The effects of ER manipulation on cocaine CPP in males, as measured by time spent in

the drug-paired chamber, is shown in Figure 3 A-C. Administration of ICI, a nonselective ERα/β

antagonist and GPER1 agonist, attenuated males' preference for 10mg/kg cocaine (Figure 3A).

A two-way repeated measures ANOVA revealed a treatment x test session interaction ($F_{(1,15)}$ =

5.758; p = 0.0299, n^2p = 0.277). A Bonferroni multiple comparisons test revealed that CHOL

treated animals increased time spent in the cocaine-paired chamber (4.745 ± 0.974, p = 0.0004)

but ICI treated males did not (0.482 ± 1.033, p > 0.9999).

The goal of our follow up study was to differentiate between the effects of ICI on

inhibition of ERα and ERβ versus activation of GPER1. We did this by using G1, a selective

GPER1 agonist. Figure 3 B represents males treated with G1 compared to CHOL. A two-way

repeated measures ANOVA revealed that G1 treatment replicated the behavioral effect of ICI

treatment on cocaine CPP; there was a significant interaction between test session and

treatment ($F_{(1,15)}$ = 7.429; p = 0.016, n^2p = 0.331) and multiple comparisons revealed that at

10mg/kg cocaine, G1 treated males did not acquire a CPP for cocaine (1.272 ± 0.972, p =

0.3804), but CHOL animals did (4.745 ± 0.8741, p = 0.0001).

We further investigated whether administration of the GPER1 antagonist, G15, could

cause enhanced preference for cocaine in males. Illustrated in Figure 3 C, at a 5mg/kg

conditioning dose, CHOL treated males did not show a CPP for cocaine however, G15 treated

males did. A two-way repeated measures ANOVA revealed a significant interaction between

test session and treatment ($F_{(1,15)}$ = 8.194; p = 0.0119, n^2p = 0.353). A Bonferroni multiple

comparisons test revealed that males treated with G15 spent more time in the drug-paired

chamber after conditioning while CHOL treated males did not (1.591 ± 0.6143, p = 0.0406;

0.826 ± 0.579, p = 0.3182; respectively).

Effects of ER manipulation on cocaine CPP in females

The effects of ER manipulation on cocaine CPP in females are shown in Figure 3 D-E.

When cocaine CPP was determined in females treated with CHOL, ICI, or G1, all groups formed

a preference for 10mg/kg cocaine. A two-way repeated measures ANOVA was performed to

compare test sessions by treatment. Figure 3 D represents CHOL and ICI treated females,

where there was a main effect of test session ($F_{(1,16)}$ = 13.99; p = 0.018, n^2p = 0.467). There

was no main effect of treatment however, and no interaction between test session and

treatment. Similar results are shown in Figure 3 E, which represents CHOL and G1 treated

females, where there was only a main effect of test session ($F_{(1,16)}$ = 20.05; p = 0.0004, n^2p =

0.556).

Effects of intra DLS ER manipulation on saccharin preference in males and females

The effects of the GPER1 agonist, G1, on preference for 0.1% SACC versus water is

illustrated in Figure 3 D. An unpaired t-test comparing preference scores between CHOL and

G1 treated males revealed significant group differences ($t_{(10)}$ = 2.589; p = 0.0270, d = 1.495) in

preference scores. There were no differences in total water intake on days 1-4, prior to

treatment. There were also no significant differences in total liquid consumed on days 5-12

between treatment groups.

Consistent with the effects of G1 on cocaine CPP, there was no effect of G1 on females'

preference for 0.1% SACC over water, shown in Figure 4 C. An unpaired t-test was used to

reveal no significant group differences ($t_{(10)}$ = 0.7998; p = 0.4424, d = 0.462). Both G1 and

CHOL groups had equal variance and data were normally distributed. There were no

differences in total liquid consumption on days 1-4, prior to treatment or during treatment on

days 5-12.

Effects of intra DLS ER manipulation on gonad weights in males and females

There was no effect of G1 treatment intra-DLS on gonad weight in either males or

females. In females, this included uterus weight ($t_{(10)}$ = 0.4571; p = 0.6574, d = 0.263) and

ovary weights ($t_{(10)}$ = 1.892; p = 0.0878. d = 1.09). For males, this included testes ($t_{(10)}$ =

0.1270; p = 0.9014, d = 0.073) and vas deferens weights ($t_{(10)}$ = 1.625; p = 0.1352, d = 0.938).

Relative mRNA expression of ERα, Erβ, and GPER1 in the dorsal striatum of males and

females

As shown in Figure 5 A-C, there were no sex differences in relative mRNA expression of

ERs in the dorsal striatum. For ERα, male data points violated normality testing, therefore an

unpaired nonparametric Mann-Whitney U test was performed to identify group differences. No

sex differences in ERα expression were found (U = 19; p = 0.1949, n = 0.142). Unpaired t tests

revealed no group differences for ERβ ($t_{(14)}$ = 0.3334; p = 0.7438, d = 0.166), or GPER1($t_{(14)}$ =

0.7504; p = 0.4654, d = 0.374).

The expression of GPER1 mRNA was greater than ERα and Erβ mRNA relative to the

HRPT housekeeping gene, as illustrated in Figure 5D. A significant one-way ANOVA

determined differences between ER subtypes ($F_{(2,45)}$ = 180.3; p < 0.0001, n^2 = 0.889). A

Bonferroni multiple comparisons test determined a significant difference in relative expression of

GPER1 versus ERα (p < 0.0001) and GPER1 versus ERβ (p < 0.0001) but no difference

between ERα and ERβ (p > 0.9999).

Discussion

Our study is the first to report that GPER1 activation in the DLS modulates preference

formation for cocaine in male rats. We have shown that activation of GPER1 in the DLS is

sufficient to attenuate cocaine CPP, while inhibition of GPER1 receptors produces a significant

preference for cocaine at a dose that does not usually induce CPP in males. We also identified

the relative mRNA expression of ERs in the dorsal striatum and found no sex differences,

indicating that the sex-specific effect of GPER1 activation/inhibition is not due to an overall sex

difference in the expression of receptors within this brain region.

Previous work in male rats found that animals that formed a strong taste aversion to

amphetamine, relative to animals that formed a low taste aversion, also formed a stronger place

preference for amphetamine, suggesting a common mechanism mediates both effects [29]. Our

findings that GPER1 activation inhibits preference for cocaine in males was replicable with an

alternative reward, 0.1% SACC, which was preferred in control males as well as both

experimental and control females. These animals still drank comparable amounts of liquid and

consumed equitable amounts of chow compared to controls, which suggests that GPER1

activation does not cause an overall malaise in male animals. Instead, we hypothesize that

GPER1 activation could be altering the learned rewarding effects of cocaine and SACC. In both

experiments, the rewarding stimuli were introduced after G1 was administered into the DLS.

Future experiments are needed to determine whether GPER1 activation would decrease the

rewarding properties of cocaine or SACC after a preference has been established, or if the

effect of GPER1 is limited to the initial establishment of a preference.

We initially hypothesized that the pharmacological manipulation of ERs in the DLS of

females would alter their cocaine CPP, based on previous findings that E2 regulates the

rewarding properties of cocaine in females [30]. We predicted that administering ICI, an ERα and

ERβ antagonist would inhibit CPP formation, but our results do not support this. Both ICI and G1

attenuated cocaine CPP in males, but not females. Since ICI is an ERα and ERβ antagonist,

and a GPER1 agonist, this suggests that it was the GPER1 agonist action that attenuated CPP

in males. We postulate that the effects of ICI in females were seen because it is also an agonist

for GPER1 and that we would have needed to give an antagonist all three receptors to inhibit

cocaine preference formation in females. We also recognize that a significant limitation of this

study is the lack of varied doses, as we only administered 10% drug:vehicle in both sexes.

Although this was a sufficient dose in males, it could be that females need a higher or lower

dose to cause a change in behavior.

Females show enhanced rotational behavior and sensitization after cocaine exposure

compared to males and this effect is mediated by ERα [25,31]. Work in dissociated medium spiny

neurons from dorsal striatum of females has shown E2 decreases L-type calcium current, which

implicates ERβ [32,33]. In neurons from males, the response to E2 was significantly less [34].

Finally, E2 treatment reduces GABA release, and overexpression of ERα in the dorsal striatum

also enhances the inhibitory effects of E2 on GABA release [31,35]. Thus, ERα and ERβ are

implicated in the effects of E2 on striatal function in females.

Our study used qPCR to explore relative RNA expression of ERα, ERβ, and GPER1 in

the dorsal striatum and we did not find any sex differences. These findings are consistent with

recent evidence supporting no sex differences in protein levels of ERα or GPER1 in the dorsal

striatum of adult males and females [36]. Our study did determine that GPER1 mRNA levels are

greater than ERα and ERβ, which do not differ from one another.

We did not differentiate the medial versus lateral subsections of the dorsal striatum, and

distribution of ERs within the dorsal striatum could differ by sex. It is known that in females,

membrane receptors that are coupled to mGluR mediate the rapid effects of E2 in the DLS, but

this mechanism has not been investigated in males [22,24]. Whether there are sex differences in

the signaling pathways mediating the effects of E2 in the DLS needs to be explored. Together,

these data suggest that the sex-specific behavioral outcomes of GPER1 manipulation are likely

due to differences in the downstream effects of receptor activation, rather than sex differences

in overall expression.

In conclusion, we report that we have identified a novel role for GPER1 in males. To our

knowledge, this is the first set of studies to show that activation or inhibition of GPER1 in the

DLS is sufficient to alter cocaine conditioned place preference in males. Based on the role of E2

seen in female drug-seeking, we historically hypothesized an increase in motivated behaviors

behaviors after E2 treatment, and therefore designed studies to detect an increase in drug-

seeking or drug preference, rather than a decrease. This could be one reason that a role for E2

in drug-seeking in males has been missed until now. Given these results, we postulate that

GPER1 is a potential target for decreasing motivation to attain cocaine in males, which is

currently under investigation in our laboratory.

Funding and Disclosure

Funding was contributed by NIH R01-DA-039952. The authors have no conflicts of interest to

disclose.

Acknowledgements

In alphabetical order, we thank Kendra Beaudoin, Hannah Epstein, Brianna Graham, Lahin

Lalani and Benjamin Lipkin for assisting in craniotomies, day-to-day animal handling as well as

behavioral testing. We would also like to acknowledge Caitlin Posillico for her integral role in

tissue processing and qPCR data collection. We thank Molly Logsdon for helping with creating

the figures for this manuscript and Brandon Luma for the construction of the chambers used in

the CPP behavioral Paradigm.

Author Contributions

All authors had full access to all the data in the study and take responsibility for the integrity of

the data and accuracy of the data analysis. *Conceptualization:* Quigley, J.A. and Becker, J.B.;

Methodology: Quigley, J.A. and Becker, J.B.; *Investigation:* Quigley, J.A.; *Formal Analysis:*

Quigley, J.A.; *Resources:* Becker, J.B.; *Writing – Original Draft:* Quigley, J.A.; *Writing – Revision*

and Editing: Quigley, J.A. and Becker, J.B.; *Visualization:* Quigley, J.A.; *Supervision:* Becker,

J.B.; *Funding:* Becker, J.B.

References

1.   John WS, Wu L-T. Trends and correlates of cocaine use and cocaine use disorder in the
United States from 2011 to 2015. *Drug Alcohol Depend.* 2017;180:376-384.
doi:10.1016/j.drugalcdep.2017.08.031.

2.   Hernandez-Avila CA, Rounsaville BJ, Kranzler HR. Opioid-, cannabis- and alcohol-
dependent women show more rapid progression to substance abuse treatment. *Drug
Alcohol Depend.* 2004;74(3):265-272. doi:10.1016/j.drugalcdep.2004.02.001.

3.   Greenfield SF, Back SE, Lawson K, Brady KT. Substance abuse in women. *Psychiatr Clin
North Am* 2010;33(2):339-355. doi:10.1016/j.psc.2010.01.004.

4.   Sanvicente-Vieira B, Rovaris DL, Ornell F, et al. Sex-based differences in
multidimensional clinical assessments of early-abstinence crack cocaine users. *PLoS One*
2019;14(6):e0218334. doi:10.1371/journal.pone.0218334.

5.   Kosten TA, Gawin FH, Kosten TR, Rounsaville BJ. Gender differences in cocaine use and
treatment response. *J Subst Abuse Treat* 1993;10(1):63-66.

6.   Back SE, Payne RL, Wahlquist AH, et al. Comparative profiles of men and women with
opioid dependence: results from a national multisite effectiveness trial. *Am J Drug Alcohol
Abuse* 2011;37(5):313-323. doi:10.3109/00952990.2011.596982.

7.   Carroll ME, Lynch WJ, Roth ME, Morgan AD, Cosgrove KP. Sex and estrogen influence
drug abuse. *Trends Pharmacol. Sci.* 2004;25(5):273-279. doi:10.1016/j.tips.2004.03.011.

8.   Becker JB. Sex differences in addiction. *Dialogues Clin Neurosci* 2016;18(4):395-402.

9.   Sofuoglu M, Dudish-Poulsen S, Nelson D, Pentel PR, Hatsukami DK. Sex and menstrual
cycle differences in the subjective effects from smoked cocaine in humans. *Exp Clin
Psychopharmacol* 1999;7(3):274-283.

10.   Becker JB, Hu M. Sex differences in drug abuse. *Front Neuroendocrinol* 2008;29(1):36-
47. doi:10.1016/j.yfrne.2007.07.003.

11.   Hu M, Crombag HS, Robinson TE, Becker JB. Biological basis of sex differences in the
propensity to self-administer cocaine. *Neuropsychopharmacology* 2004;29(1):81-85.
doi:10.1038/sj.npp.1300301.

12.   Roberts DC, Bennett SA, Vickers GJ. The estrous cycle affects cocaine self-administration
on a progressive ratio schedule in rats. *Psychopharmacology* 1989;98(3):408-411.
doi:10.1007/BF00451696.

13.   Nicolas C, Russell TI, Pierce AF, et al. Incubation of Cocaine Craving After Intermittent-
Access Self-administration: Sex Differences and Estrous Cycle. *Biol. Psychiatry*
2019;85(11):915-924. doi:10.1016/j.biopsych.2019.01.015.

14.   Razandi M, Pedram A, Greene GL, Levin ER. Cell membrane and nuclear estrogen
receptors (ERs) originate from a single transcript: studies of ERalpha and ERbeta
expressed in Chinese hamster ovary cells. *Mol. Endocrinol.* 1999;13(2):307-319.

451   doi:10.1210/mend.13.2.0239.

452 15. Luoma JI, Boulware MI, Mermelstein PG. Caveolin proteins and estrogen signaling in the
453   brain. *Mol. Cell. Endocrinol.* 2008;290(1-2):8-13. doi:10.1016/j.mce.2008.04.005.

454 16. Fuentes N, Silveyra P. Estrogen receptor signaling mechanisms. *Adv. Protein Chem.*
455   *Struct. Biol.* 2019;116:135-170. doi:10.1016/bs.apcsb.2019.01.001.

456 17. Otto C, Rohde-Schulz B, Schwarz G, et al. G protein-coupled receptor 30 localizes to the
457   endoplasmic reticulum and is not activated by estradiol. *Endocrinology*
458   2008;149(10):4846-4856. doi:10.1210/en.2008-0269.

459 18. Almey A, Milner TA, Brake WG. Estrogen receptor α and G-protein coupled estrogen
460   receptor 1 are localized to GABAergic neurons in the dorsal striatum. *Neurosci. Lett.*
461   2016;622:118-123. doi:10.1016/j.neulet.2016.04.023.

462 19. Almey A, Filardo EJ, Milner TA, Brake WG. Estrogen receptors are found in glia and at
463   extranuclear neuronal sites in the dorsal striatum of female rats: evidence for cholinergic
464   but not dopaminergic colocalization. *Endocrinology* 2012;153(11):5373-5383.
465   doi:10.1210/en.2012-1458.

466 20. Kiyatkin EA. Cocaine enhances the changes in extracellular dopamine in nucleus
467   accumbens associated with reinforcing stimuli: a high-speed chronoamperometric study in
468   freely moving rats. *Eur. J. Neurosci.* 1993;5(3):284-291.

469 21. Pettit HO, Justice JB. Dopamine in the nucleus accumbens during cocaine self-
470   administration as studied by in vivo microdialysis. *Pharmacol. Biochem. Behav.*
471   1989;34(4):899-904. doi:10.1016/0091-3057(89)90291-8.

472 22. Yoest KE, Quigley JA, Becker JB. Rapid effects of ovarian hormones in dorsal striatum
473   and nucleus accumbens. *Horm. Behav.* 2018;104:119-129.
474   doi:10.1016/j.yhbeh.2018.04.002.

475 23. Cummings JA, Jagannathan L, Jackson LR, Becker JB. Sex differences in the effects of
476   estradiol in the nucleus accumbens and striatum on the response to cocaine:
477   neurochemistry and behavior. *Drug Alcohol Depend.* 2014;135:22-28.
478   doi:10.1016/j.drugalcdep.2013.09.009.

479 24. Song Z, Yang H, Peckham EM, Becker JB. Estradiol-Induced Potentiation of Dopamine
480   Release in Dorsal Striatum Following Amphetamine Administration Requires Estradiol
481   Receptors and mGlu5. *Eneuro* 2019;6(1). doi:10.1523/ENEURO.0446-18.2019.

482 25. Hu M, Becker JB. Effects of sex and estrogen on behavioral sensitization to cocaine in
483   rats. *J. Neurosci.* 2003;23(2):693-699.

484 26. Robinson TE, Becker JB, Presty SK. Long-term facilitation of amphetamine-induced
485   rotational behavior and striatal dopamine release produced by a single exposure to
486   amphetamine: sex differences. *Brain Res.* 1982;253(1-2):231-241.

487 27. Sun K, Wang F, Ma L, et al. Genetic knockout of the G protein-coupled estrogen receptor
488   1 facilitates the acquisition of morphine-induced conditioned place preference and
489   aversion in mice. *Biochem. Biophys. Res. Commun.* 2020.

490   doi:10.1016/j.bbrc.2020.03.003.

28. Becker JB, Snyder PJ, Miller MM, Westgate SA, Jenuwine MJ. The influence of estrous cycle and intrastriatal estradiol on sensorimotor performance in the female rat. *Pharmacol. Biochem. Behav.* 1987;27(1):53-59. doi:10.1016/0091-3057(87)90476-X.

29. Turenne SD, Miles C, Parker LA, Siegel S. Individual differences in reactivity to the rewarding/aversive properties of drugs: Assessment by taste and place conditioning. *Pharmacology Biochemistry and Behavior* 1996;53(3):511-516. doi:10.1016/0091-3057(95)02042-X.

30. Hu M, Becker JB. Acquisition of cocaine self-administration in ovariectomized female rats: effect of estradiol dose or chronic estradiol administration. *Drug Alcohol Depend.* 2008;94(1-3):56-62. doi:10.1016/j.drugalcdep.2007.10.005.

31. Schultz KN, von Esenwein SA, Hu M, et al. Viral vector-mediated overexpression of estrogen receptor-alpha in striatum enhances the estradiol-induced motor activity in female rats and estradiol-modulated GABA release. *J. Neurosci.* 2009;29(6):1897-1903. doi:10.1523/JNEUROSCI.4647-08.2009.

32. Boulware MI, Mermelstein PG. Membrane estrogen receptors activate metabotropic glutamate receptors to influence nervous system physiology. *Steroids* 2009;74(7):608-613. doi:10.1016/j.steroids.2008.11.013.

33. Mermelstein PG, Becker JB, Surmeier DJ. Estradiol reduces calcium currents in rat neostriatal neurons via a membrane receptor. *J. Neurosci.* 1996;16(2):595-604.

34. Becker JB, James D, Mermelstein PG. Estradiol reduces calcium currents in rat neostriatal neurons via a membrane receptor. *J. Neurosci.* 1996.

35. Hu M, Watson CJ, Kennedy RT, Becker JB. Estradiol attenuates the K+-induced increase in extracellular GABA in rat striatum. *Synapse* 2006;59(2):122-124. doi:10.1002/syn.20221.

36. Krentzel AA, Willett JA, Johnson AG, Meitzen J. Estrogen receptor alpha, G-protein coupled estrogen receptor 1, and aromatase: Developmental, sex, and region-specific differences across the rat caudate-putamen, nucleus accumbens core and shell. *J. Comp. Neurol.* 2020. doi:10.1002/cne.24978.

Figure 1.
(A) Number of male and female animals per treatment condition for each experiment. **(B)** Representative cannula placement into the DLS.

Figure 2.
(A) The apparatus used for conditioned place preference training and testing was a three-compartment chamber with three areas, differentiated by distinct tactile and visual cues. As described in the timeline, the pre-test took place on day 1, followed by 8 conditioning sessions, ending with a final test session. Version 1 (V1) and version 2 (V2) refer to conditioning either beginning with saline or cocaine on day 2, which was counterbalanced across animals. Treatment stylets were inserted after pre-test on day 1 and remained inserted for continuous treatment through the final test session. **(B)** A two-bottle choice paradigm was utilized to determine preference for 0.1% SACC versus plain water. Two bottles were accessible to individually housed animals in their home cages. On days 1-4, both bottles contained H_2O only. During days 5-12, one bottle contained H_2O and the other contained 0.1% SACC dissolved in H_2O. Treatment stylets were inserted before SACC introduction on day 5 and remained for the duration of the experiment.

Figure 3.
GPER1 activation/inhibition intra-DLS alters cocaine CPP for males but not females. A cocaine CPP is inferred if animals spend more time in drug-paired chamber during test versus pre-test. **(A)** At a conditioning dose of 10mg/kg cocaine, CHOL (control) treated males form a CPP however, treatment of ICI (nonselective ERα/ERβ antagonist; GPER1 agonist) or **(B)** G1 (selective GPER1 agonist) attenuates males CPP for cocaine. **(C)** Treatment of G15 (selective GPER1 antagonist) causes a CPP for a 5mg/kg dose of cocaine; CHOL treated males do not form a CPP. **(D-E)** Females treated with CHOL, ICI, and G1 form a CPP for 10mg/kg cocaine. CPP data are shown as mean +/- SEM and each point corresponds to an individual animal. *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$.

Figure 4.
GPER1 activation intra-DLS attenuates preference for 0.1% SACC versus H_2O for males but not females. **(A)** G1 treatment causes a conditioned avoidance of 0.1% SACC, as indicated by a preference score < 50%. G1 treated males have a significantly lower preference score, averaged across days, than CHOL (control) males. **(B)** Both G1 and CHOL (control) treated females similarly formed a preference for 0.1% SACC over H_2O, as indicated by group preference scores exceeding 50%.
Preference score data are shown as mean +/- SEM with data points representing individual days. *$p < 0.05$, **$p < 0.01$.

Figure 5.
(A-C) There are no significant differences in relative mRNA expression of estradiol receptors in the dorsal striatum between females and males. Data are shown as males relative to females. **(D)** Relative mRNA expression of GPER1 is greater than ERα and ERβ, which do not differ from one another. Data are shown as GPER1 and ERβ relative to ERα.
Data are shown as mean +/- SEM with data points representing individual animals. ****$p < 0.0001$

A

	Treatment	Male	Female
CPP	ICI (ERα/β antagonist; GPER1 agonist) (10mg/kg cocaine)	8	8
	G1 (GPER1 agonist) (10mg/kg cocaine)	8	8
	CHOL (10mg/kg cocaine)	9	10
	G15 (GPER1 antagonist) (5mg/kg cocaine)	8	--
	CHOL (5mg/kg cocaine)	9	--
SACC vs. H₂O	G1 (GPER1 agonist)	6	6
	CHOL	6	6
qPCR	*naïve*	8	8

B

Males   **Females**

Males: 0.1% SACC vs H$_2$O

Females: 0.1% SACC vs H$_2$O