Dopamine modulates prediction error forwarding in the nonlemniscal inferior colliculus

Catalina Valdés-Baizabal[1,2†], Guillermo V. Carbajal[1,2†],

David Pérez-González[1,2] & Manuel S. Malmierca[1,2,3*]*

[1] Auditory Neuroscience Laboratory, Institute of Neuroscience of Castilla y León, Calle Pintor Fernando Gallego 1, 37007 Salamanca, Spain

[2] The Salamanca Institute for Biomedical Research (IBSAL), 37007 Salamanca, Spain.

[3] Department of Biology and Pathology, Faculty of Medicine, Campus Miguel de Unamuno, University of Salamanca, 37007 Salamanca, Spain

[†] Equal contribution

[*] Corresponding authors

Short title: Dopamine modulates prediction error forwarding in the nonlemniscal IC

Keywords: inferior colliculus (IC), stimulus-specific adaptation (SSA), predictive coding, predictive processing, perceptual learning, perceptual inference, prediction error, precision, dopamine, eticlopride

Abstract

The predictive processing framework describes perception as a hierarchical predictive model of sensation. Higher-level neural structures constrain the processing at lower-level structures by suppressing synaptic activity induced by predictable sensory input. But when predictions fail, deviant input is forwarded bottom-up as 'prediction error' to update the perceptual model. The earliest prediction error signals identified in the auditory pathway emerge from the nonlemniscal inferior colliculus (IC). The drive that these feedback signals exert on the perceptual model depends on their 'expected precision', which determines the postsynaptic gain applied in prediction error forwarding. Expected precision is theoretically encoded by the neuromodulatory (e.g., dopaminergic) systems. To test this empirically, we recorded extracellular responses from the rat nonlemniscal IC to oddball and cascade sequences before, during and after the microiontophoretic application of dopamine or eticlopride (a D_2-like receptor antagonist). Hence, we studied dopaminergic modulation on the subcortical processing of unpredictable and predictable auditory changes. Results demonstrate that dopamine reduces the net neuronal responsiveness exclusively to unexpected input, without significantly altering the processing of expected auditory events at population level. We propose that, in natural conditions, dopaminergic projections from the thalamic subparafascicular nucleus to the nonlemniscal IC could serve as a precision-weighting mechanism mediated by D_2-like receptors. Thereby, the levels of dopamine release in the nonlemniscal IC could modulate the early bottom-up flow of prediction error signals in the auditory system by encoding their expected precision.

List of abbreviatures

CAS: cascade condition.

CSI: common SSA index.

dB SPL: decibels sound pressure level.

DEV: deviant condition.

FR: firing rate (spikes/s).

FRA: frequency response area.

IC: inferior colliculus.

iMM: index of neuronal mismatch.

PE: prediction error.

SFR: spontaneous firing rate (spikes/s).

SPF: subparafascicular nucleus of the thalamus.

SSA: stimulus-specific adaptation.

STD: standard condition.

TDT: Tucker-Davis Technologies.

Introduction

Perceptual systems prune redundant sensory input as a means of sparing processing resources while providing saliency to input that is rare, unique, and therefore potentially more informative. This perceptual function has been classically studied in humans using the auditory oddball paradigm [1], where the successive repetition of a tone ('standard condition', henceforth 'STD') is randomly interrupted by an 'oddball' tone ('deviant condition', henceforth 'DEV'). When applied to animal models, the oddball paradigm unveils a phenomenon of neuronal short-term plasticity called stimulus-specific adaptation (SSA), measured as the difference between DEV and STD responses [2].

SSA first emerges in the auditory system at the level of the inferior colliculus (IC), mainly in its nonlemniscal portion (i.e., the IC cortices) [3]. As a site of convergence of both ascending and descending auditory pathways, the IC plays a key role in processing deviant sounds over redundant ones [4] and shaping the auditory context [5]. The complex computational network of the IC integrates excitatory, inhibitory and rich neuromodulatory input [6,7]. This includes dopaminergic innervation from the subparafascicular nucleus (SPF) of the thalamus to the nonlemniscal IC [8–12]. Previous reports have detected mRNA coding for dopaminergic D_2-like receptors in the IC [9,13] and proved its functional expression as protein [14], while other studies demonstrated that dopamine modulates the auditory responses of IC neurons in heterogeneous manners [10,14]. However, the involvement of dopaminergic modulation of SSA in the IC is yet to be confirmed.

In recent years, subcortical SSA in the auditory system has been reinterpreted in the context of the predictive processing [15]. This conceptual

framework posits that hierarchically coupled neuronal populations infer the

hidden causes of sensation and predict upcoming sensory regularities using a

generative model of the world [16–25]. In such hierarchical model, higher neural

populations try to explain away or inhibit the sensory input prompted by the

hidden states of the world. As a result, lower-level neural populations receiving

those top-down predictions decrease their responsiveness to expected sensory

inputs, which during an oddball paradigm manifest functionally as SSA of the STD

response. But when encountering a DEV, the generative model fails to predict

that 'oddball', forwarding a prediction error (PE) signal which reports the

unexpected portions of sensory input to the higher-level neural population. That

bottom-up flow of PE signals serves to provide feedback and update the inferred

representations about the states of the world along each level of the processing

hierarchy. In a previous study from our lab performed in awake and anaesthetized

rodents, we demonstrated that DEV responses in the nonlemniscal IC were better

explained as PE signaling activity [26].

In the predictive processing framework, there are only two sorts of things

that need to be inferred about the world: the state of the world, and the uncertainty

about that state [27]. On the one hand, representations about the states of the

world emerge from the hierarchical exchange of top-down predictions and

bottom-up PEs, which is embodied in the synaptic activity of the nervous system.

On the other hand, this inferential process entails a certain degree of uncertainty,

which is encoded in terms of *expected precision* or confidence by means of the

postsynaptic gain [27–29]. Thereby, synaptic messages are weighted according

to their expected precision as they are passed along the processing hierarchy.

When expected precision is high, PE signals receive postsynaptic amplification

to strengthen its updating power. Conversely, when imprecision is expected, PE

signals undergo negative gain to prevent misrepresentations. Neuromodulators

(such as dopamine) cannot directly excite or inhibit postsynaptic responses, but

only modulate the postsynaptic responses to other neurotransmitters. Therefore,

according to some predictive processing implementations [27,30–32], the only

possible function of neuromodulatory systems is to encode the expected

precision.

In this study, we perform microiontophoretic injections of dopamine and

eticlopride (a D_2-like receptor antagonist) while recording single- and multi-unit

responses under oddball and regular sequences to determine whether

dopaminergic input to the nonlemniscal IC modulates response properties and

predictive processing. Our results demonstrate that dopamine has a profound

effect on how unexpected sounds are processed in the nonlemniscal IC. We

argue this could be compatible with a dopaminergic encoding of the expected

precision of PE signals, acting as a regulatory mechanism at the level of the

auditory brainstem.

138 Materials and Methods

Surgical procedures

We conducted experiments on 31 female Long-Evans rats aged 9–17

weeks with body weights between 150–250 gr. All methodological procedures

were approved by the Bioethics Committee for Animal Care of the University of

Salamanca (USAL-ID-195), and performed in compliance with the standards of

the European Convention ETS 123, the European Union Directive 2010/63/EU

and the Spanish Royal Decree 53/2013 for the use of animals in scientific

research.

We first induced surgical anesthesia with a mixture of ketamine/xylazine

(100 and 20 mg/Kg respectively, intramuscular) and then maintained it with

urethane (1.9 g/Kg, intraperitoneal). To ensure a stable deep anesthetic level, we

administered supplementary doses of urethane (~0.5 g/Kg, intraperitoneal) when

the corneal or pedal withdrawal reflexes were present. We selected urethane over

other anesthetic agents because it better preserves normal neural activity, having

a modest, balanced effect on inhibitory and excitatory synapses [33–36].

Prior to the surgery, we recorded auditory brainstem responses (ABR) with

subcutaneous needle electrodes to verify the normal hearing of the rat. We

acquired the ABR using a RZ6 Multi I/O Processor (Tucker-Davis Technologies,

TDT) and BioSig software (TDT) before beginning each experiment. ABR stimuli

consisted of 0.1 ms clicks at a rate of 21 clicks/s, delivered monaurally to the right

ear in 10 dB steps, from 10 to 90 decibels of sound pressure level (dB SPL), in a

closed system through a speaker coupled to a small tube sealed in the ear.

After normal hearing was confirmed, we placed the rat in a stereotaxic

frame where the ear bars were replaced by hollow specula that accommodated

the sound delivery system. We performed a craniotomy in the left parietal bone

to expose the cerebral cortex overlying the left IC. We removed the dura overlying

the left IC and covered the exposed cortex with 2% agar to prevent desiccation.

Data acquisition procedures

Experiments were performed inside a sound-insulated and electrically

shielded chamber. All sound stimuli were generated using a RZ6 Multi I/O

Processor (TDT) and custom software programmed with OpenEx Suite (TDT) and MATLAB (MathWorks). In search of evoked auditory neuronal responses from the IC, we presented white noise bursts and sinusoidal pure tones of 75 ms duration with 5 ms rise-fall ramps. Once the activity of an auditory unit was clearly identified, we only used pure tones to record the experimental stimulation protocols. All protocols ran at 4 stimuli per second and were delivered monaurally in a closed-field condition to the ear contralateral to the left IC through a speaker. We calibrated the speaker using a ¼-inch condenser microphone (model 4136, Brüel&Kjær) and a dynamic signal analyzer (Photon+, Brüel&Kjær) to ensure a flat spectrum up to ~73 dB SPL between 0.5 and 44 kHz, and that the second and third signal harmonics were at least 40 dB lower than the fundamental at the loudest output level.

To record extracellular activity while carrying out microiontophoretic injections, we attached a 5-barrel glass pipette to a hand-manufactured, glass-coated tungsten microelectrode (1.4–3.5 MΩ impedance at 1 kHz), with the tip of the electrode protruding 15–25 µm from the pipette tip [37]. We place the electrode over the exposed cortex, forming an angle of 20° with the horizontal plane towards the rostral direction. Using a piezoelectric micromanipulator (Sensapex), we advanced the electrode while measuring the penetration depth until we could observe a strong spiking activity synchronized with the train of searching stimuli.

Analog signals were digitized with a RZ6 Multi I/O Processor, a RA16PA Medusa Preamplifier and a ZC16 headstage (TDT) at 12 kHz sampling rate and amplified 251×. Neurophysiological signals for multiunit activity were band-pass filtered between 0.5 and 4.5 kHz. Stimulus generation and neuronal response

processing and visualization were controlled online with custom software created

with the OpenEx suite (TDT) and MATLAB. A unilateral threshold for automatic

action potential detection was manually set at about 2–3 standard deviations of

the background noise. Spike waveforms were displayed on the screen and

overlapped on each other in a pile-plot to facilitate isolation of units. Recorded

spikes were considered to belong to a single unit when the SNR of the average

waveform was larger than 5 (51% of the recorded units).

Stimulation protocols

For all recorded neurons, we first computed the frequency-response area

(FRA), which is the map of response magnitude for each frequency/intensity

combination (Fig. 1). The stimulation protocol to obtain the FRA consisted of a

randomized sequence of sinusoidal pure tones ranging between 0.7–44 kHz, 75

ms of duration with 5 ms rise-fall ramps, presented at a 4 Hz rate, randomly

varying frequency and intensity of the presented tones (3–5 repetitions of all

tones).

Protocol 1: Oddball paradigm (DEV and STD)

In a first round of experiments, we used the oddball paradigm (Fig. 4A) to

study SSA. We presented trains of 400 stimuli containing two different

frequencies (f_1 and f_2) in a pseudo-random order at a 4Hz repetition rate and at

a level of 10–40 dB above threshold. Both frequencies were within the excitatory

FRA previously determined for the neuron (Fig. 1), and evoked similar firing rates

(FR). One frequency (f_1) appeared with high probability within the sequence

(STD; P=0.9). The succession of STD tones was randomly interspersed with the

second frequency (f_2), presented with low probability within the sequence (DEV;

P=0.1). After obtaining one data set, the relative probabilities of the two stimuli

were reversed, with f_2 becoming the STD and f_1 becoming the DEV (Fig. 4A).

This allows to control for the physical characteristics of the sound in the evoked

response, such that the differential response between DEV and STD of a given

tone can only be due to their differential probability of appearance. The separation

between f_1 and f_2 was 0.28 (49 units) or 0.5 (45 units; Protocol 2 was also applied

to these units) octaves, which is within the range of frequency separations used

in other previous studies [3,26,38–43]. The units from those two groups were

pooled together, since their responses did not differ significantly. Deviant and

standard responses were averaged from all stimulus presentations from both

tested frequencies..

The Common SSA Index (CSI) was calculated as:

233
$$CSI = \frac{DEV_{f1} + DEV_{f2} - STD_{f1} - STD_{f2}}{DEV_{f1} + DEV_{f2} + STD_{f1} + STD_{f2}}$$

where DEV_{fi} and STD_{fi} are FRs in response to a frequency f_i when it was

presented in deviant and standard conditions, respectively. The CSI ranges

between -1 to +1, being positive if the DEV response was greater than the STD

response. The firing rates in response to DEV or STD stimuli were calculated

using 100 ms windows starting at the beginning of each stimulus. Spontaneous

firing rates were calculated using 75 ms windows previous to each individual

stimulus.

Protocol 2: Oddball paradigm + Cascade sequence (CAS)

In light of the recent discovery of PE signals recorded in the nonlemniscal

IC [26], we decided to adapt our stimulation protocol to that of Parras and

colleagues for a second round of experiments which incorporated the cascade

sequence (CAS) [44]. By arranging a set of 10 tones in a regular succession of

ascending or descending frequency, no tone is ever immediately repeated.

Consequently, whereas CAS does not induce SSA—as opposed to STD—, its

pattern remains predictable, so the next tone in the sequence can be expected—

as opposed to DEV. Thus, this design contains 3 conditions of auditory transit:

(1) *no change* or *predictable repetition* (i.e., STD), which is the most susceptible

to SSA or repetition suppression (Fig. 4A, bottom); (2) *predictable change* (i.e.,

CAS; Fig. 4B); and (3) *unpredictable change* (i.e., DEV), which should allegedly

elicit the strongest PE signaling when it surprisingly interrupts the uniform train of

STDs (Fig. 4A, top).

Therefore, after computing the FRA (Fig. 1), we selected 10 evenly-spaced

tones at a fixed sound intensity 10–40 dB above minimal response threshold, so

that at least two tones fell within the FRA limits. Those 10 frequencies were

separated from each other by 0.5 octaves, in order to make the results

comparable to those of [26]. We used the 10 tones to build the ascending and

descending versions of CAS (Fig. 4B). We selected 2 tones within that lot to

generate the ascending and descending versions of the oddball paradigm (Fig.

4A), comparing the resultant DEV with their corresponding CAS versions (Fig.

4B). All sequences were 400 tones in length, at the same, constant presentation

rate of 4 Hz. Thus, each frequency could be compared with itself in DEV, STD

and CAS conditions (Fig. 4A-B), obtaining 40 trials per condition. To allow

comparison between responses from different neurons, we normalized the spike

count evoked by each tone in DEV, STD and CAS as follows:

269
$$DEV_N = \frac{DEV}{N} \quad ; \quad STD_N = \frac{STD}{N} \quad ; \quad CAS_N = \frac{CAS}{N}$$

Where,

$$N = \sqrt{DEV^2 + STD^2 + CAS^2}$$

From these normalized responses, we computed the index of neuronal

mismatch (iMM) as:

$$iMM = DEV_N - STD_N$$

These indices range between -1 and 1. The iMM is largely equivalent to

the classic CSI as an index of SSA, as previously demonstrated by Parras and

colleagues (see their supplementary figure 2 in [26]). Nevertheless, please

beware the CSI provides one index for each pair of tones in the oddball paradigm,

whereas the iMM provides one index for each tone tested.

Dopaminergic manipulation procedures

After recording the chosen stimulation protocol in a *'control condition'*, i.e.,

before any dopaminergic manipulation, we applied either dopamine or the D_2-like

receptor antagonist eticlopride (Sigma-Aldrich Spain) iontophoretically through

multi-barreled pipettes attached to the recording electrode. The glass pipette

consisted of 5 barrels in an H configuration (World Precision Instruments,

catalogue no. 5B120F-4) with the tip broken to a diameter of 30–40 µm [37]. The

center barrel was filled with saline for current compensation (165 mM NaCl). The

others were filled with dopamine (500 mM) or eticlopride (25 mM). Each drug was

dissolved in distilled water and the acidity of the solution was adjusted with HCl

to pH 3.5 for dopamine and pH 5 for eticlopride. The drugs were retained in the

pipette with a -20 nA current and ejected using 90 nA currents (Neurophore BH-

2 system, Harvard Apparatus). Thus, we released dopamine or eticlopride into

the microdomain of the recorded neuron at concentrations that have been

previously demonstrated effective in *in vivo* studies [10]. About 5 minutes after

the drug injection, we repeated the FRA and the chosen stimulation protocol

continuously until the drug was washed away, leaving roughly 2–3 minutes

between the end of one recording set and the beginning of the next one. The

recording set showing a maximal SSA alteration relative to the control values was

considered the *'drug condition'* of that neuron. We established the *'recovery*

condition' when the CSI returned to levels that did not significantly differ from

control values, never before 40 minutes post-injection. We used either dopamine

or eticlopride during protocol 1, while only dopamine was tested during protocol

2.

Histological verification procedures

At the end of each experiment, we inflicted electrolytic lesions (5 µA, 5 s)

through the recording electrode. Animals were sacrificed with a lethal dose of

pentobarbital, after which they were decapitated, and the brains immediately

immersed in a mixture of 1% paraformaldehyde and 1% glutaraldehyde in 1 M

PBS. After fixation, tissue was cryoprotected in 30% sucrose and sectioned in the

coronal plane at 40 µm thickness on a freezing microtome. We stained slices with

0.1% cresyl violet to facilitate identification of cytoarchitectural boundaries.

Finally, we assigned the recorded units to one of the main subdivisions of the IC

using the standard sections from a rat brain atlas as reference [45].

Data analysis

The peristimulus histograms representing the time-course of the

responses (Fig. 2E-F, 3E-F) were calculated using 1 ms bins an then smoothed

with a 6 ms gaussian kernel ("ksdensity" function in MATLAB) to estimate the spike-density function over time.

All the data analyses were performed with SigmaPlot (Systat Software) and MATLAB software, using the built-in functions, the Statistics and Machine Learning toolbox for MATLAB, as well as custom scripts and functions developed in our laboratory. Unless stated otherwise, all average values for trials and neurons in the present study are expressed as *'median [interquartile range]'*, since the data did not follow a normal distribution (one-sample Kolmogorov-Smirnov test).

We performed a bootstrap procedure to analyze dopaminergic effects on each individual neuron. SSA indices, both CSI [2] and iMM [26], are calculated from the averages of the single-trial responses to DEV, STD and CAS. Consequently, only one value of such indexes can be obtained for each unit and condition. Therefore, to test the drug effects on each unit, we calculated the 95% bootstrap confidence intervals for the SSA index in the control condition. The bootstrap procedure draws random samples (with replacement) from the spike counts evoked on each trial, separately for DEV and STD stimuli, and then applies either the CSI or the iMM formula. This procedure is repeated 10000 times, thus obtaining a distribution of expected CSI values based on the actual responses from a single recording. We applied this procedure using the *bootci* MATLAB function, as in previous studies of SSA neuromodulation in the IC [38,39,42,46], which returned the 95% confidence interval for the CSI in the control condition. We considered drug effects to be significant when SSA index in the drug condition did not overlap with the confidence interval in the control condition.

We used the Wilcoxon Signed Rank test (*signrank* function in MATLAB) to

check for differences at the population level between the control and drug CSI

and FRs.

Results

In order to study the role of dopamine in shaping SSA in the nonlemniscal

IC, we recorded responses from a total of 142 single- and multi-units in 31 young

adult Long Evans rats. In a first series of experiments (protocol 1, see *Methods*)

we presented the oddball paradigm before, during and after microiontophoretic

application of dopamine (n=94) or eticlopride (n=43). In an additional series of

experiments (n=43, from the former pool of units), we also presented two cascade

sequences (ascending and descending) in addition to the oddball paradigm [26]

(protocol 2, see *Methods)*. Histological verification located all recorded neurons

in the rostral cortex of the IC, where SSA indexes tend to be higher [3,4,43].

Dopamine effects on the CSI

The microiontophoretic application of dopamine caused an average 15%

decrease of SSA in our sample, falling from a median CSI of 0.51 [0.25–0.78] in

the control condition to 0.43 [0.15–0.73] after dopamine application (p=0.014;

Figure 2A). Such average reduction was caused by a 26% drop in the median

DEV response (control FR: 5.63 [3.25–10.00] spikes/s; dopamine FR: 4.19 [1.63–

7.63]; p<0.001; red in Figure 2B,C), while the STD response did not show a

significant change (control FR: 1.71 [0.64–3.83]; dopamine FR: 1.63 [0.36–3.65];

p=0.284; blue in Figure 2B,D). As observed in previous reports [3], the

spontaneous firing rate (SFR) found in the nonlemniscal IC tended to be very

scarce (for individual examples, see Figs. 2E-F and 3E-F) and did not change significantly with dopamine application (control SFR: 0.18 [0.05–0.77]; dopamine SFR: 0.15 [0.03–0.90]; p=0.525; gray in Figure 2B).

Previous studies had reported heterogeneous dopaminergic effects on the response of IC neurons [10,14], so we performed a bootstrap analysis to evaluate the statistical significance of CSI changes unit by unit. We confirmed such heterogeneity across our sample, with 42 units following the population trend by decreasing their CSI, whereas 33 units showed CSI increments under dopamine; 19 units remained unaltered (Fig. 2A). Figure 2E shows the response of a unit to STD (blue) and DEV (red) in the control condition (left panel), during dopamine application (middle panel) and after recovery (right panel). The application of dopamine caused an increment of the STD response and a decrement of the DEV response, leading to a decrease of the CSI. In contrast, the unit in Figure 2F showed a decrement of the response to both STD and DEV during the application of dopamine, thus resulting in an increase of the CSI. The effects of dopamine peaked around 8-10 minutes after microiontophoretic application, followed by a progressive recovery to baseline values that could take beyond 90 minutes (Fig. 2E-F, right panels).

Eticlopride effects on the CSI

We aimed to determine whether dopaminergic effects on the CSI were mediated by D_2-like receptors, as suggested by previous reports [9,13]. To test endogenous dopaminergic modulation on SSA mediated by D_2-like receptors, we applied eticlopride, a D_2-like receptor antagonist, to 43 units. We observed no significant response changes at sample level (DEV FR: p=0.609; STD FR:

p=0.769; SFR: p=0.405; CSI change: p=0.170; Figure 3A-D). However, we

performed a bootstrap analysis to evaluate the statistical significance of CSI

changes in each unit under eticlopride influence, which revealed that only 11 units

remained unaffected. The CSI had significantly increased in 19 units and

decreased in 13 units (Fig. 3A), implying that eticlopride was indeed antagonizing

endogenous dopaminergic modulation mediated by D_2-like receptors on those

units.

Dopamine effects on the iMM and CAS

To test whether dopamine modulates PE signaling in the nonlemniscal IC,

we performed an additional set of experiments following stimulation protocol 2

(see *Methods*), which was based on the methodology of a previous study [26].

Alongside the oddball paradigm (Fig. 4A), we recorded responses of 43 units to

two cascade sequences, which consisted of 10 tones presented in a predictable

succession of increasing or decreasing frequencies (Fig. 4B).

We used a bootstrap analysis to evaluate the statistical significance of the

effect of dopamine on the iMM of each recording, which confirmed that 23 units

underwent heterogeneous iMM changes (Fig. 4C, colored dots, each

representing one tested frequency) whereas another 18 remained stable (Fig.

4C, gray dots). Results agreed with those obtained using the CSI, as the iMM of

the sample fell by 22%, from a median of 0.57 [0.41-0.69] in the control condition

to a median of 0.45 [0.27–0.65] under dopaminergic influence (p=0.002; Fig. 4C).

This was caused by a significant reduction in the median DEV response (control

normalized FR: 0.70 [0.58–0.80]; dopamine normalized FR: 0.64 [0.47–0.78];

p=0.002), while the STD response was not affected (control normalized FR: 0.11

[0.02–0.23]; dopamine normalized FR: 0.10 [0.03–0.31]; p=0.188). Most

interestingly, CAS response also remained unaffected by dopamine application

(control normalized FR: 0.68 [0.52–0.77]; dopamine normalized FR: 0.71 [0.57–

0.83]; p=0.115; Fig. 5D).

Discussion

We recorded single- and multi-unit activity in the nonlemniscal IC under an

auditory oddball paradigm while performing microiontophoretic applications of

dopamine and eticlopride (D_2-like receptor antagonist). Following the discovery

of PE signaling activity in the nonlemniscal IC [26], we included cascade

sequences [44] in a subset of experiments to address dopamine role from a

predictive processing standpoint. This resulted in 3 stimulation conditions: (1)

STD or *expected repetition* (Fig. 4A, bottom), susceptible of generating intense

SSA; (2) DEV or *unexpected change* (Fig. 4A, top), which should elicit the

strongest PE signaling; and (3) CAS or *expected change* (Fig. 4B), a condition

featuring the same STD-to-DEV step, but which neither undergo SSA (unlike

STD) nor should entail a PE (or, at least, not as strong as DEV). Our results

revealed that dopamine modulates SSA and PE signaling in the nonlemniscal IC.

Dopamine reduces DEV responses and thereby SSA indices at population level

Dopamine application caused a 15% reduction of SSA in the nonlemniscal

IC (Figs. 2A, 4C) due to general drop in DEV responses of 25% (Fig. 2B, C).

Neither STD (Fig. 2B, D) nor CAS responses were significantly affected at

population level (Fig. 4D). The differential effect of dopamine on DEV and STD

cannot be explained by the differences in their control FR, since CAS yielded FRs

as high as DEV that were not similarly reduced by dopamine (Fig. 4D). In other

words, dopamine application decreased the responsiveness to *unexpected*

stimuli while the responsiveness to the expected stimuli remained stable.

Therefore, dopamine likely modulates net PE signaling from the nonlemniscal IC.

Eticlopride effects did not describe significant tendencies at population

level (Fig. 3A-D). Nevertheless, 75% of our sample manifested significant SSA

changes under eticlopride (Fig. 3A, colored dots). This confirms the release of

endogenous dopamine, as well as the functional expression of D_2-like receptors

in the nonlemniscal IC. Taken together with previous findings regarding

dopaminergic modulation of the IC [9,10,13,14], the reduction of SSA and PE

signaling is most likely mediated by D_2-like receptors.

The net reduction of DEV responses with dopamine is unique as compared

with the effects of other neurotransmitters and neuromodulators on IC neurons.

GABAergic and glutamatergic manipulations alter the general excitability of IC

neurons, thereby exerting symmetrical effects on STD and DEV responses which

result in a gain control of SSA [42,46,47]. Conversely, cholinergic and

cannabinoid manipulation yield asymmetrical effects that mostly affect STD

responses [38,39]. Activation of M1 muscarinic receptors and CB1 cannabinoid

receptors tended to reduce average SSA by increasing responsiveness to

repetitive stimuli (i.e., STD). Dopamine also deliver asymmetrical effects, but in

contrast with the aforementioned cases, the activation of D_2-like receptors tends

to reduce population SSA by decreasing responsiveness to surprising stimuli

(i.e., DEV).

Intrinsic and synaptic properties generate heterogeneous dopaminergic effects

In line with previous reports [10,14], dopaminergic effects were

heterogeneous across units. Complex dopaminergic interactions altering the

excitation-inhibition balance cannot be accurately tracked, because the exact

location and neuronal types expressing D_2-like receptors in the IC are yet to be

determined. Notwithstanding, the heterogeneity of dopaminergic effects must

result from distinctive intrinsic and synaptic properties.

D_2-like receptors are coupled to G proteins which regulate the activity of

manifold voltage-gated ion channels, adjusting excitability depending on the

repertoire expressed in each neuron [48]. D_2-like receptors coupled to Gi/o

proteins can both increase potassium currents and decrease calcium currents via

βγ subunit complex, thereby reducing excitability [48]. The opening probability of

calcium channels can also diminish by the activation of D_2-like receptors coupled

to Gq proteins [48]. D_2-like receptor activation can augment or reduce sodium

currents depending on the receptor subtypes expressed on the neuronal

membrane [48]. Furthermore, D_2-like receptor activation can also reduce NMDA

synaptic transmission, decreasing the FR [48]. In addition, nonlemniscal IC

neurons express hyperpolarization-activated cyclic nucleotide-gated (HCN)

channels [49,50], which can be modulated by dopamine and yield mixed effects

on neuronal excitability [51].

In addition, dopamine and eticlopride can interact with D_2-like receptors

expressed in a presynaptic neuron. Both glutamatergic and GABAergic

projections converge onto single IC neurons [6,52,53], which may also receive

dopaminergic inputs from the SPF [8–12]. Dopamine could potentially activate

presynaptic D_2-like receptors expressed in a glutamatergic neuron, as described

in striatal medium spiny neurons [54–56], or conversely in a GABAergic neuron,

as demonstrated in the ventral tegmental area [57].

SPF dopaminergic projections to the IC cortices: An early precision-weighting

mechanism in the auditory system?

Dopaminergic function has been traditionally studied in the context of

reinforcement learning, where dopamine is thought to encode the discrepancy

between expected and observed reward in a 'reward PE' [58,59]. Dopaminergic

neurons report positive PE values by increasing their firing and negative PE

values by reducing their tonic discharge rates, thereby guiding the learning

process [60]. Hence, these signed 'reward PEs' encoded by dopaminergic

neurons are substantially different from the unsigned 'sensory PEs' encoded by

auditory neurons in the nonlemniscal IC [21,23]. According to this interpretation,

our dopamine ejections in the nonlemniscal IC could mimic reinforcing signals,

which in natural conditions would come from the SPF [8–12]. Speculatively, such

dopaminergic input might aim to induce long-term potentiation on IC neurons to

build lasting associations between acoustic cues and rewarding outcomes,

thereby contributing to establish reward expectations. However, we fail to see

why these positive reward PEs would mitigate the transmission of sensory PEs

from the nonlemniscal IC, as evidenced by the reduced DEV responses we have

observed after dopamine application.

An alternative interpretation from the predictive processing framework

argues that information (i.e., PE) about the hidden states of the world cannot be

encoded by dopamine release, since dopamine cannot directly excite the

postsynaptic responses which would be needed to mediate the influence of that

information [27]. Dopamine can only modulate the postsynaptic responses to

other neurotransmitters, a function more compatible with expected precision

encoding and PE weighting. This approach spares the need of two distinct types

of PE signaling, while better accommodating some findings that were not easily

explained as reward PEs [27]. A significant portion of dopaminergic neurons

increase their firing in response to aversive stimuli and cues which predict them,

contrary to how reward PEs should work [61–63]. Most relevant to the present

study, dopaminergic neurons also respond to conditions where the reward PE

should theoretically be zero [64,65], including novel or unexpected stimuli [66–

69].

Everything considered, we propose that the most suited way of interpreting

our data is through the lens of a precision-weighting mechanism. A tentative

predictive processing explanation is that dopamine release in the nonlemniscal

IC encodes expected imprecision. In other words, the role of dopaminergic input

to the IC cortices could be to apply a net negative gain which dampens the

forward propagation of PE signals. Such net inhibitory effect over sensory

transmission would especially affect DEV responses, as the unexpected

interruptions of an otherwise very regular train of stimuli will elicit the strongest

PE signaling activity of all conditions tested in this study.

In natural conditions, such midbrain-level modulation of the bottom-up PE

flow could be shaped by SPF dopaminergic neurons projecting to the IC cortices

[11,13,70]. Many auditory nuclei project to the SPF, including auditory cortex,

auditory thalamus, the superior olivary complex and the IC itself, providing the

SPF with rich auditory information [70–74]. Even other centers, which perform

higher-order functions in the sensory processing and integration of auditory

information, send projections to the SPF, such as the medial prefrontal cortex

and the deep layers of the superior colliculus [73]. Hence, the dopaminergic

activity of the SPF must be interwoven to a great extent with the general

functioning of the auditory system. The reciprocal connectivity of the SPF with

many nuclei at multiple levels of the auditory pathway could provide the structural

basis for an optimal encoding of expected precision, modulating via dopaminergic

input the weight of PEs forwarded from the nonlemnical IC.

Limitations

Our proposal might seem at odds with some previous works regarding

neuromodulation in the predictive processing framework. Whereas cholinergic

and NMDA manipulation are often reported to yield precision-weighting effects

[20,30,75], dopaminergic effects on PE are less common in the literature [76],

and they are usually linked to processes of active inference [27,77–80], rather

than perceptual inference and learning. Besides, classic neuromodulators are

often thought to increase the expected precision of PE signaling [30], contrary to

the inhibitory net effects of dopamine that we found in the nonlemniscal IC.

Notwithstanding, it is important to keep in mind that the current view on the

relationship between neuromodulation and expected precision derives from, and

mainly refers to, cortical data. Cortical intrinsic circuitry and its neuromodulatory

sources are vastly different to those of the nonlemniscal IC. Cortical predictive

processing implementations have proposed specific hypotheses about the

neuronal types encoding precision-weighted PEs in a defined canonical

microcircuit [16,21,23]. Unfortunately, to the best of our knowledge [4,6], current

understanding on the intrinsic circuitry of the nonlemniscal IC does not allow to

directly import such hypotheses into auditory midbrain processing. The observed heterogeneity of dopaminergic effects in our sample is indeed compatible with distinct neuronal types fulfilling differentiated processing roles in the intrinsic circuitry of the nonlemniscal IC. However, it is not possible to distinguish between neuronal types and assign them putative roles solely based on their functional data [81]. In any case, our results make room for distinct roles of neuromodulation along the multiple stages of predictive processing along the auditory hierarchy. A possibility that could be more adequately addressed in future studies.

Conclusions

Our study demonstrates that dopamine modulates auditory midbrain processing of unexpected input. We propose that dopamine release in the nonlemniscal IC could encode expected imprecision, consequently reducing the postsynaptic gain of PE signals and thereby dampening their drive over higher-level processing stages. The dopaminergic projections from the thalamic SPF to the IC cortices could be the biological substrate of this early precision-weight mechanism. Thus, despite being usually neglected by most corticocentric approaches, our results confirm subcortical structures as a key element of predictive processing, at least in the auditory system.

Acknowledgments

CVB and GVC contributed equally to this work. We thank Drs Edward L Bartlett, Nell Cant, Adrian Rees and Richard Rosch for their useful comments on previous versions of the manuscript. We also thank Mr Antonio Rivas Cornejo for taking care of histological processing.

Financial disclosure

Financial support provided by Spanish MINECO (SAF2016-75803-P) to MSM.

CVB held a grant from Mexican CONACYT (216652). GVC held a fellowship from

the Spanish MICINN (BES-2017-080030).

FIGURES

Figure 1

Figure 1. Effect of dopamine on the FRA. **A)** FRA of a neuron in control condition

(left panel) and after dopamine application (right panel). **B)** The subtraction of the

control FRA from that during application of dopamine in A reveals that dopamine

increased the excitability of this neuron. **C)** FRA of another neuron in control

condition (left panel) and after dopamine application (right panel). **D)** The

subtraction of the control FRA from that during application of dopamine in C

reveals that dopamine decreased the excitability of this neuron.

Figure 2

Figure 2. Effects of dopamine on the CSI. **A)** Scatter plot of the CSI in control condition versus dopamine application. Units that underwent significant CSI changes are represented in purple, whereas the rest are marked as gray dots. The purple cross on the abscissa axis represents one CSI measurement which ordinate value falls out of scale (y = −0.59). **B)** Violin plots of the SFR (gray), DEV response (red) and STD response (blue). Control conditions are represented in the left half of each violin (no color) while dopamine effects are on display in the right half (colored). Horizontal thick black lines mark the median of each distribution, while vertical bars cover the interquartile range. Regarding statistical significance, **n.s.** indicates that p > 0.05 and ****** indicates that p < 0.001. **C)** Scatter plot of DEV responses in control condition versus dopamine application. **D)** Scatter plot of STD responses in control condition versus dopamine application. **E)** Peristimulus histogram of a unit before (left panel), during (middle panel) and after (right panel) dopamine application. In this case, dopamine reduced the CSI. **F)** Another example showing the opposite effects.

640   *Figure 3*

Figure 3. Effects of eticlopride on the CSI. **A)** Scatter plot of the CSI in control condition versus eticlopride application. Units that underwent significant CSI changes are represented in green, whereas the rest are marked as gray dots. **B)** Violin plots of the SFR (gray), DEV response (red) and STD response (blue). Control conditions are represented in the left half of each violin (no color) while eticlopride effects are on display in the right half (colored). Horizontal thick black lines mark the median of each distribution, while vertical bars cover the interquartile range. Regarding statistical significance, **n.s.** indicates that $p > 0.05$. **C)** Scatter plot of DEV responses in control condition versus eticlopride application. **D)** Scatter plot of STD responses in control condition versus eticlopride application. **E)** Peristimulus histogram of a unit before (left panel), during (middle panel) and after (right panel) eticlopride application. In this case, eticlopride reduced the CSI. **F)** Another unit example showing the opposite effects.

	Figure 4

Figure 4. Dopamine effects on predictable versus unpredictable auditory events.

A) Oddball paradigm, displaying two experimental conditions for a given f_i target

tone. **B)** Cascade sequences highlighting the f_i target tone. **C)** Scatter plot of the

iMM in control condition versus dopamine application. Frequencies that

underwent significant iMM changes are represented in purple, whereas the rest

are marked as gray dots. **D)** Violin plots of the CAS (green), DEV (red) and STD

(blue) normalized responses. Control conditions are represented in the left half

of each violin (no color) while dopamine effects are on display in the right half

(colored). Horizontal thick black lines mark the median of each distribution, while

the boxplots inside each distribution indicate the interquartile range, with the

confidence interval for the median indicated by the notches. Regarding statistical

significance, **n.s.** indicates that p > 0.05 and ** indicates that p < 0.01 (repeated

measures ANOVA, Dunn-Sidak correction).

REFERENCES

1. Näätänen R, Gaillard AWK, Mäntysalo S. Early selective-attention effect

on evoked potential reinterpreted. Acta Psychol (Amst). 1978;42: 313–

329. doi:10.1016/0001-6918(78)90006-9

2. Ulanovsky N, Las L, Nelken I. Processing of low-probability sounds by

cortical neurons. Nat Neurosci. 2003;6: 391–398. doi:10.1038/nn1032

3. Malmierca MS, Cristaudo S, Pérez-González D, Covey E. Stimulus-

Specific Adaptation in the Inferior Colliculus of the Anesthetized Rat. J

Neurosci. 2009;29: 5483–5493. doi:10.1523/JNEUROSCI.4153-08.2009

4. Malmierca MS, Carbajal GV, Escera C. Deviance Detection and Encoding

Acoustic Regularity in the Auditory Midbrain. In: Kandler K, editor. The

Oxford Handbook of the Auditory Brainstem. Oxford University Press;

2019. pp. 706–740. doi:10.1093/oxfordhb/9780190849061.013.19

5. Petersen CL, Hurley LM. Putting it in context: Linking auditory processing

with social behavior circuits in the vertebrate brain. Integr Comp Biol.

2017;57: 865–877. doi:10.1093/icb/icx055

6. Ito T, Malmierca MS. Neurons, Connections, and Microcircuits of the

Inferior Colliculus. In: Oliver D, Cant N, Fay R, Popper A, editors. The

Mammalian Auditory Pathways. Springer, Cham; 2018. pp. 127–167.

doi:10.1007/978-3-319-71798-2_6

7. Hurley LM. Neuromodulatory Feedback to the Inferior Colliculus. In:

Kandler K, editor. The Oxford Handbook of the Auditory Brainstem.

Oxford University Press; 2018.

doi:10.1093/oxfordhb/9780190849061.013.15

8. Hurd YL, Suzuki M, Sedvall GC. D1 and D2 dopamine receptor mRNA

expression in whole hemisphere sections of the human brain. J Chem

Neuroanat. 2001;22: 127–137. doi:10.1016/S0891-0618(01)00122-3

9. Nevue AA, Felix RA, Portfors C V. Dopaminergic projections of the

subparafascicular thalamic nucleus to the auditory brainstem. Hear Res.

2016;341: 202–209. doi:10.1016/j.heares.2016.09.001

10. Gittelman JX, Perkel DJ, Portfors C V. Dopamine modulates auditory

responses in the inferior colliculus in a heterogeneous manner. JARO - J

Assoc Res Otolaryngol. 2013;14: 719–729. doi:10.1007/s10162-013-

0405-0

11. Batton AD, Blaha CD, Bieber A, Lee KH, Boschen SL. Stimulation of the

subparafascicular thalamic nucleus modulates dopamine release in the

inferior colliculus of rats. Synapse. 2018: 0–2. doi:10.1002/syn.22073

12. Weiner DM, Levey AI, Sunahara RK, Niznik HB, O'Dowd BF, Seeman P,

et al. D1 and D2 dopamine receptor mRNA in rat brain. Proc Natl Acad

Sci U S A. 1991;88: 1859–63. doi:10.1073/PNAS.88.5.1859

13. Nevue AA, Elde CJ, Perkel DJ, Portfors C V. Dopaminergic Input to the

Inferior Colliculus in Mice. Front Neuroanat. 2016;9: 168.

doi:10.3389/fnana.2015.00168

14. Hoyt JM, Perkel DJ, Portfors C V. Dopamine acts via D2-like receptors to

modulate auditory responses in the inferior colliculus. eNeuro. 2019;6:

ENEURO.0350-19.2019. doi:10.1523/ENEURO.0350-19.2019

15. Carbajal GV, Malmierca MS. The Neuronal Basis of Predictive Coding

Along the Auditory Pathway: From the Subcortical Roots to Cortical

Deviance Detection. Trends Hear. 2018;22: 233121651878482.

doi:10.1177/2331216518784822

16. Friston KJ. A theory of cortical responses. Philos Trans R Soc B Biol Sci.

2005;360: 815–836. doi:10.1098/rstb.2005.1622

17. Friston KJ. The free-energy principle: a rough guide to the brain? Trends

Cogn Sci. 2009;13: 293–301. doi:10.1016/j.tics.2009.04.005

18. Garrido MI, Kilner JM, Stephan KE, Friston KJ. The mismatch negativity:

a review of underlying mechanisms. Clin Neurophysiol. 2009;120: 453–

463. doi:10.1016/j.clinph.2008.11.029

19. Auksztulewicz R, Friston KJ. Repetition suppression and its contextual

determinants in predictive coding. Cortex. 2016;80: 125–140.

doi:10.1016/j.cortex.2015.11.024

20. Auksztulewicz R, Schwiedrzik CM, Thesen T, Doyle W, Devinsky O,

Nobre AC, et al. Not All Predictions Are Equal: "What" and "When"

Predictions Modulate Activity in Auditory Cortex through Different

Mechanisms. J Neurosci. 2018;38: 8680–8693.

doi:10.1523/JNEUROSCI.0369-18.2018

21. Bastos AM, Usrey WM, Adams RA, Mangun GR, Fries P, Friston KJ.

Canonical Microcircuits for Predictive Coding. Neuron. 2012;76: 695–711.

doi:10.1016/j.neuron.2012.10.038

22. Garrido MI, Kilner JM, Kiebel SJ, Stephan KE, Baldeweg T, Friston KJ.

Repetition suppression and plasticity in the human brain. Neuroimage.

2009;48: 269–279. doi:10.1016/j.neuroimage.2009.06.034

23. Keller GB, Mrsic-Flogel TD. Predictive processing: a canonical cortical

computation. Neuron. 2018;100: 424–435.

doi:10.1016/j.neuron.2018.10.003

24. Clark A. Surfing uncertainty: prediction, action, and the embodied mind.

Oxford University Press; 2016.

25. Hohwy J. The predictive mind. Oxford, England: Oxford University Press;

2013. doi:10.5860/choice.169393

26. Parras GG, Nieto-Diego J, Carbajal GV, Valdés-Baizabal C, Escera C,

Malmierca MS. Neurons along the auditory pathway exhibit a hierarchical

organization of prediction error. Nat Commun. 2017;8: 2148.

doi:10.1038/s41467-017-02038-6

27. Friston KJ, Shiner T, FitzGerald THB, Galea JM, Adams RA, Brown HR,

et al. Dopamine, affordance and active inference. Sporns O, editor. PLoS

Comput Biol. 2012;8: e1002327. doi:10.1371/journal.pcbi.1002327

28. Feldman H, Friston KJ. Attention, Uncertainty, and Free-Energy. Front

Hum Neurosci. 2010;4: 215. doi:10.3389/fnhum.2010.00215

29. Spratling MW. Reconciling predictive coding and biased competition

models of cortical function. Front Comput Neurosci. 2008;2: 4.

doi:10.3389/neuro.10.004.2008

30. Moran RJ, Campo P, Symmonds M, Stephan KE, Dolan RJ, Friston KJ.

Free Energy, Precision and Learning: The Role of Cholinergic

Neuromodulation. J Neurosci. 2013;33: 8227–8236.

doi:10.1523/JNEUROSCI.4255-12.2013

31. Parr T, Friston KJ. Uncertainty, epistemics and active Inference. J R Soc

Interface. 2017;14. doi:10.1098/rsif.2017.0376

32. Parr T, Friston KJ. The Anatomy of Inference: Generative Models and

Brain Structure. Front Comput Neurosci. 2018;12: 90.

doi:10.3389/fncom.2018.00090

33. Hara K, Harris RA. The anesthetic mechanism of urethane: The effects on

neurotransmitter-gated ion channels. Anesth Analg. 2002;94: 313–318.

doi:10.1213/00000539-200202000-00015

34. Maggi CA, Meli A. Suitability of urethane anesthesia for

physiopharmacological investigations in various systems. Part 1: General

considerations. Experientia. 1986;42: 109–114. doi:10.1007/BF01952426

35. Sceniak MP, MacIver MB. Cellular Actions of Urethane on Rat Visual

Cortical Neurons In Vitro. J Neurophysiol. 2006;95: 3865–3874.

doi:10.1152/jn.01196.2005

36. Duque D, Malmierca MS. Stimulus-specific adaptation in the inferior

colliculus of the mouse: anesthesia and spontaneous activity effects.

Brain Struct Funct. 2015;220: 3385–3398. doi:10.1007/s00429-014-0862-

1

37. Ayala YA, Pérez-González D, Duque D, Palmer AR, Malmierca MS.

Extracellular Recording of Neuronal Activity Combined with

Microiontophoretic Application of Neuroactive Substances in Awake Mice.

J Vis Exp. 2016; e53914. doi:10.3791/53914

38. Ayala YA, Malmierca MS. Cholinergic Modulation of Stimulus-Specific

Adaptation in the Inferior Colliculus. J Neurosci. 2015;35: 12261–12272.

doi:10.1523/JNEUROSCI.0909-15.2015

39. Valdés-Baizabal C, Parras GG, Ayala YA, Malmierca MS.

Endocannabinoid Modulation of Stimulus-Specific Adaptation in Inferior

Colliculus Neurons of the Rat. Sci Rep. 2017;7: 6997.

doi:10.1038/s41598-017-07460-w

40. Ayala YA, Pérez-González D, Duque D, Nelken I, Malmierca MS.

Frequency discrimination and stimulus deviance in the inferior colliculus

and cochlear nucleus. Front Neural Circuits. 2013;6: 119.

doi:10.3389/fncir.2012.00119

41. Pérez-González D, Malmierca MS. Variability of the time course of

stimulus-specific adaptation in the inferior colliculus. Front Neural Circuits.

2012;6: 107. doi:10.3389/fncir.2012.00107

42. Ayala YA, Malmierca MS. The effect of inhibition on stimulus-specific

adaptation in the inferior colliculus. Brain Struct Funct. 2018;223: 1391–

1407. doi:10.1007/s00429-017-1546-4

43. Duque D, Pérez-González D, Ayala YA, Palmer AR, Malmierca MS.

Topographic Distribution, Frequency, and Intensity Dependence of

Stimulus-Specific Adaptation in the Inferior Colliculus of the Rat. J

Neurosci. 2012;32: 17762–17774. doi:10.1523/JNEUROSCI.3190-

12.2012

44. Ruhnau P, Herrmann B, Schröger E. Finding the right control: The

mismatch negativity under investigation. Clin Neurophysiol. 2012;123:

507–512. doi:10.1016/j.clinph.2011.07.035

45. Paxinos G, Watson C. The rat brain in stereotaxic coordinates. Elsevier;

2007.

46. Pérez-González D, Hernández O, Covey E, Malmierca MS. GABA A-

mediated inhibition modulates stimulus-specific adaptation in the inferior

colliculus. Coleman MJ, editor. PLoS One. 2012;7: e34297.

doi:10.1371/journal.pone.0034297

47. Ayala YA, Pérez-González D, Malmierca MS. Stimulus-specific

adaptation in the inferior colliculus: The role of excitatory, inhibitory and

modulatory inputs. Biol Psychol. 2016;116: 10–22.

doi:10.1016/j.biopsycho.2015.06.016

48. Neve KA, Seamans JK, Trantham-Davidson H. Dopamine receptor

signaling. Journal of Receptors and Signal Transduction. Taylor &

Francis; 2004. pp. 165–205. doi:10.1081/RRS-200029981

49. Ahuja TK, Wu SH. Intrinsic membrane properties and synaptic response

characteristics of neurons in the rat's external cortex of the inferior

colliculus. Neuroscience. 2007;145: 851–865.

doi:10.1016/j.neuroscience.2006.12.031

50. Koch U, Braun M, Kapfer C, Grothe B. Distribution of HCN1 and HCN2 in

rat auditory brainstem nuclei. Eur J Neurosci. 2004;20: 79–91.

doi:10.1111/j.0953-816X.2004.03456.x

51.   He C, Chen F, Li B, Hu Z. Neurophysiology of HCN channels: From

cellular functions to multiple regulations. Prog Neurobiol. 2014;112: 1–23.

doi:10.1016/J.PNEUROBIO.2013.10.001

52.   LeBeau FE, Malmierca MS, Rees A. Iontophoresis in vivo demonstrates a

key role for GABA(A) and glycinergic inhibition in shaping frequency

response areas in the inferior colliculus of guinea pig. J Neurosci.

2001;21: 7303–12. doi:21/18/7303 [pii]

53.   Malmierca MS. Chapter 29 – Auditory System. In: Paxinos G, editor. The

Rat Nervous System. Academic Press; 2015. pp. 865–946.

doi:10.1016/B978-0-12-374245-2.00029-2

54.   Calabresi P, Mercuri NB, Sancesario G, Bernardi G. Electrophysiology of

dopamine-denervated striatal neurons. Implications for Parkinson's

disease. Brain. 1993;116 (Pt 2: 433–52. Available:

http://www.ncbi.nlm.nih.gov/pubmed/8096420

55.   Bamford NS, Robinson S, Palmiter RD, Joyce JA, Moore C, Meshul CK.

Dopamine modulates release from corticostriatal terminals. J Neurosci.

2004;24: 9541–52. doi:10.1523/JNEUROSCI.2891-04.2004

56.   Higley MJ, Sabatini BL. Competitive regulation of synaptic Ca2+ influx by

D2 dopamine and A2A adenosine receptors. Nat Neurosci. 2010;13: 958–

966. doi:10.1038/nn.2592

57.   Michaeli A, Yaka R. Dopamine inhibits GABA(A) currents in ventral

tegmental area dopamine neurons via activation of presynaptic G-protein

coupled inwardly-rectifying potassium channels. Neuroscience. 2010;165:

1159–69. doi:10.1016/j.neuroscience.2009.11.045

58. Sutton RS, Barto AG. Reinforcement Learning: An Introduction. 2nd

editio. Cambridge,MA: The MIT Press; 1998.

59. Schultz W, Dayan P, Montague PR. A neural substrate of prediction and

reward. Science (80-). 1997;275: 1593–1599.

doi:10.1126/science.275.5306.1593

60. Diederen KMJ, Ziauddeen H, Vestergaard MD, Spencer T, Schultz W,

Fletcher PC. Dopamine Modulates Adaptive Prediction Error Coding in

the Human Midbrain and Striatum. J Neurosci. 2017;37: 1708–1720.

doi:10.1523/JNEUROSCI.1979-16.2016

61. Zweifel LS, Fadok JP, Argilli E, Garelick MG, Jones GL, Dickerson TMK,

et al. Activation of dopamine neurons is critical for aversive conditioning

and prevention of generalized anxiety. Nat Neurosci. 2011;14: 620–628.

doi:10.1038/nn.2808

62. Matsumoto M, Hikosaka O. Two types of dopamine neuron distinctly

convey positive and negative motivational signals. Nature. 2009;459:

837–841. doi:10.1038/nature08028

63. Bromberg-Martin ES, Matsumoto M, Hikosaka O. Dopamine in

Motivational Control: Rewarding, Aversive, and Alerting. Neuron. Elsevier;

2010. pp. 815–834. doi:10.1016/j.neuron.2010.11.022

64. Sadacca BF, Jones JL, Schoenbaum G. Midbrain dopamine neurons

889  compute inferred and cached value prediction errors in a common

890  framework. Elife. 2016;5. doi:10.7554/eLife.13665

891 65. Young AMJ, Joseph MH, Gray JA. Latent inhibition of conditioned

892  dopamine release in rat nucleus accumbens. Neuroscience. 1993;54: 5–

893  9. doi:10.1016/0306-4522(93)90378-S

894 66. Menegas W, Babayan BM, Uchida N, Watabe-Uchida M. Opposite

895  initialization to novel cues in dopamine signaling in ventral and posterior

896  striatum in mice. Elife. 2017;6. doi:10.7554/eLife.21886

897 67. Strecker RE, Jacobs BL. Substantia nigra dopaminergic unit activity in

898  behaving cats: Effect of arousal on spontaneous discharge and sensory

899  evoked activity. Brain Res. 1985;361: 339–350. doi:10.1016/0006-

900  8993(85)91304-6

901 68. Ljungberg T, Apicella P, Schultz W. Responses of monkey dopamine

902  neurons during learning of behavioral reactions. J Neurophysiol. 1992;67:

903  145–163. doi:10.1152/jn.1992.67.1.145

904 69. Horvitz JC. Mesolimbocortical and nigrostriatal dopamine responses to

905  salient non- reward events. Neuroscience. 2000;96: 651–656.

906  doi:10.1016/S0306-4522(00)00019-1

907 70. Yasui Y, Nakano K, Mizuno N. Descending projections from the

908  subparafascicular thalamic nucleus to the lower brain stem in the rat. Exp

909  Brain Res. 1992;90: 508–518. doi:10.1007/BF00230933

910 71. LeDoux JE, Ruggiero DA, Reis DJ. Projections to the subcortical

911  forebrain from anatomically defined regions of the medial geniculate body

in the rat. J Comp Neurol. 1985;242: 182–213.

doi:10.1002/cne.902420204

72. Yasui Y, Kayahara T, Nakano K, Mizuno N. The subparafascicular

thalamic nucleus of the rat receives projection fibers from the inferior

colliculus and auditory cortex. Brain Res. 1990;537: 323–327.

doi:10.1016/0006-8993(90)90378-O

73. Wang J, Palkovits M, Usdin TB, Dobolyi A. Afferent connections of the

subparafascicular area in rat. Neuroscience. 2006;138: 197–220.

doi:10.1016/j.neuroscience.2005.11.010

74. Arnault P, Roger M. Ventral temporal cortex in the rat: Connections of

secondary auditory areas Te2 and Te3. J Comp Neurol. 1990;302: 110–

123. doi:10.1002/cne.903020109

75. Rosch RE, Auksztulewicz R, Leung PD, Friston KJ, Baldeweg T.

Selective Prefrontal Disinhibition in a Roving Auditory Oddball Paradigm

Under N-Methyl-D-Aspartate Receptor Blockade. Biol Psychiatry Cogn

Neurosci Neuroimaging. 2019;4: 140–150.

doi:10.1016/j.bpsc.2018.07.003

76. Todd J, Harms L, Schall U. Mismatch negativity: Translating the potential.

Front Psychiatry. 2013;4: 171. doi:10.3389/fpsyt.2013.00171

77. Marshall L, Mathys CD, Ruge D, de Berker AO, Dayan P, Stephan KE, et

al. Pharmacological Fingerprints of Contextual Uncertainty. Rushworth

MFS, editor. PLoS Biol. 2016;14: e1002575.

doi:10.1371/journal.pbio.1002575

78.   FitzGerald THB, Dolan RJ, Friston KJ. Dopamine, reward learning, and

active inference. Front Comput Neurosci. 2015;9: 136.

doi:10.3389/fncom.2015.00136

79.   Friston KJ, Schwartenbeck P, FitzGerald THB, Moutoussis M, Behrens T,

Dolan RJ. The anatomy of choice: Dopamine and decision-making. Philos

Trans R Soc B Biol Sci. 2014;369. doi:10.1098/rstb.2013.0481

80.   Schwartenbeck P, FitzGerald THB, Mathys CD, Dolan R, Friston KJ. The

dopaminergic midbrain encodes the expected certainty about desired

outcomes. Cereb Cortex. 2015;25: 3434–3445.

doi:10.1093/cercor/bhu159

81.   Ito T. Different coding strategy of sound information between GABAergic

and glutamatergic neurons in the auditory midbrain. J Physiol. 2020;598:

1039–1072. doi:10.1113/JP279296

948