1 **Metagenomic analysis of a blood stain from the French revolutionary Jean-Paul Marat (1743-**
2 **1793)**
3

4 Toni de-Dios[1*], Lucy van Dorp[2*+], Philippe Charlier[3,4*], Sofia Morfopoulou[2,5], Esther Lizano[1],
5 Celine Bon[6], Corinne Le Bitouzé[7], Marina Alvarez-Estape[1], Tomas Marquès-Bonet[1,8,9,10],
6 François Balloux[2+], Carles Lalueza-Fox[1+]
7
[1]Institute of Evolutionary Biology (CSIC-Universitat Pompeu Fabra), 08003 Barcelona, Spain
[2]UCL Genetics Institute, University College London, London WC1E 6BT, UK
[3]Département de la Recherche et de l'Enseignement, Musée du Quai Branly - Jacques Chirac,
75007 Paris, France
[4]Université Paris-Saclay (UVSQ), Laboratory Anthropology, Archaeology, Biology (LAAB),
78180 Montigny-le-bretonneux, France
[5]Division of Infection and Immunity, University College London, London WC1E 6BT, UK
[6]Département Hommes, Natures, Sociétés, Muséum National d'Histoire Naturelle, 75116 Paris,
France
[7]Archives Nationales, 75004 Paris, France
[8]Catalan Institution of Research and Advanced Studies (ICREA), 08010 Barcelona, Spain
[9]CNAG-CRG, Centre for Genomic Regulation, Barcelona Institute of Science and Technology
(BIST), 08036 Barcelona, Spain
[10]Institut Català de Paleontologia Miquel Crusafont, Universitat Autònoma de Barcelona,
08193 Cerdanyola del Vallès, Barcelona, Spain
23
24 *These authors equally contributed to this work
25 + Co-corresponding authors

26
27 **Keywords**
28 Ancient DNA, metagenomics, infection
29
30 **Abstract**
31

The French revolutionary Jean-Paul Marat (1743-1793) was assassinated in 1793 in his
bathtub, where he was trying to find relief from the debilitating skin disease he was suffering
from. At the time of his death, Marat was annotating newspapers, which got stained with his
blood and were subsequently preserved by his sister. We extracted and sequenced DNA from
the blood stain and also from another section of the newspaper, which we used for
comparison. Results from the human DNA sequence analyses were compatible with a
heterogeneous ancestry of Marat, with his mother being of French origin and his father born in
Sardinia. Metagenomic analyses of the non-human reads uncovered the presence of fungal,
bacterial and low levels of viral DNA. Relying on the presence/absence of microbial species in
the samples, we could cast doubt on several putative infectious agents that have been
previously hypothesised as the cause of his condition but for which we detect not a single
sequencing read. Conversely, some of the species we detect are uncommon as environmental
contaminants and may represent plausible infective agents. Based on all the available
evidence, we hypothesize that Marat may have suffered from a fungal infection (seborrheic
dermatitis), possibly superinfected with bacterial opportunistic pathogens.

1 Introduction

Jean-Paul Marat (1743-1793) was a famous French physician, scientist and journalist, best known for his role as Jacobin leader during the French Revolution. Marat's parents were Giovanni Mara, born in Cagliari, Sardinia, who later added a "t" to his family name to give it a French feel and Louise Cabrol, a French Huguenot from Castres. Marat was stabbed to death in his bathtub by the Girondist' supporter Charlotte Corday on July 13th, 1793 (Figure 1a). Upon his death, his sister Charlotte Albertine kept two issues of Marat's newspaper *l'Ami du Peuple* (nº506 and nº678, published on June 30th, 1791 and August 13th, 1792, respectively), which he was annotating the day of his assassination and that got stained with his blood (Figure 1b). Albertine gave the issues to the collector François-Nicolas Maurin (1765-1848) in 1837. After his death, as explained by a handwritten note by writer Anatole France dated from October 10th, 1864, the two issues ended up in the possession of baron Carl De Vinck who in 1906 donated them to the Département des Estampes, Bibliothèque National de France, in Paris (see notice in the Catalogue Général: https://catalogue.bnf.fr/ark:/12148/cb40261215w).

Marat's health during the last years before his assassination is shrouded in mystery. He suffered from a severe itching skin disease from which he found some relief by spending most of his time in a medicinal bathtub over which he placed a board to use as a writing desk. His condition, which he attributed to his stay in the sewers of Paris while hiding from his political enemies, has been the subject of numerous medical debates and has been alternatively attributed to scabies, syphilis, atopic eczema, seborrheic dermatitis or dermatitis herpetiformis (1–5), the latter as a potential manifestation of celiac disease (6). It has been suggested that his condition affected his character and turned it more violent (1).

With the intention of shedding light on these issues, we retrieved two samples from one of the newspapers stained with Marat's blood, one sample from the blood stain and a second one from a non-stained area in the upper corner of the paper, to be used as a comparison. A principal concern was to use a non-destructive approach to explore Marat's genomic footprint; therefore, the samples were taken with forensic swabs. The DNA extracted from both samples was used to build genomic libraries that were subjected to second-generation sequencing using the Illumina platform. DNA reads where subsequently classified, separating the human reads – most likely deriving from Marat's blood – from those assigned to microbial species. The analysis of both sets of DNA sequences allowed characterisation of Marat's ancestry as well as identification of the potential pathogens responsible for his debilitating skin condition.

2 Material and methods

2.1 *DNA extraction and sequencing*
Forensic swabs were obtained from one of the newspapers Marat was annotating at the time of his assassination (Figure 1). One swab was taken from the blood stain and another from an area of the newspaper without visual evidence of blood. The blood swab was extracted with a buffer composed of 10 mM TrisHCl, 10 mM EDTA, 2 mM SDS, 50 mM DTT; proteinase K was added after one hour incubation. The extract was subsequently concentrated and purified using a Qiagen column kit. DNA extraction from both swabs was performed together with extraction blanks (no sample). A total of 35 ul of each sample was used for library preparation following the BEST protocol (7). Libraries were quantified using BioAnalyzer and sequenced by HiSeq 4000 (Illumina). Library blanks were also performed for each library batch. We generated 568,623,176 DNA reads from the blood stain, of which 74,244,610 reads mapped to the human reference genome (Table S1). From these, we retrieved a complete human mitochondrial (mtDNA) genome at a mean depth of coverage of 18.156X and a nuclear genome at 0.125X.

2.2 Mapping and variant calling

Raw sequences adapters were removed using *Cutadapt* (8). Reads were then aligned against the Human Reference genome (GRCh37/hg19) and the mitochondrial reference genome (rCRS) as well as for a set of microbial candidates using *BWA* v.07.3 (9) and Bowtie2 (10). We employed two aligners as mapping sensitivity of different aligners can vary between different samples when working with aDNA (11). Duplicate reads were discarded using *Picard* tools (12). Unique mapped reads were filtered for a mapping quality equal of above 30 (Table S1). All mapped sequences (human nuclear, human mitochondrial and microbial) were assessed for post-mortem damage patterns at the ends of reads using *MapDamage* v.2 (13), which can be used as a sign of historic authenticity over modern contamination (Fig. S1) . Post-mortem damage signals were also obtained for each read using *pmdtools* (14) (Fig. S2). Mapping statistics including the depth of coverage were recorded using *Qualimap* (15). Due to the low coverage of the human sample, we performed a pseudo-haploid calling approach, common to the processing of aDNA, using the *SAMtools Pileup* tool (16). This data was then merged with the Human Origins dataset for its use in population genetics analyses (17,18).

2.3 Modern DNA Contamination

Schmutzi was used to estimate the amount of modern DNA contamination in the mitochondrial (mtDNA) genome (19) likely deriving from the DNA of those who have handled the newspaper in the years following Marat's death. We identified mitochondrial contamination based on the inferred deamination patterns as 52.5% +/- 4.5% with the full haplogroup profiles provided in Table S2. This allowed the modern DNA sequences to be delineated from the ancient DNA sequences using *Jvarkit* and a custom script (20) by selecting the human reads with mismatches in their first or last three nucleotides. This reduced the amount of modern mtDNA contamination to 0-0.1%. The depth of coverage was recorded using *Qualimap*.

We also independently estimated the amount of contamination based on the heterozygous sites in the X chromosome using *angsd v0.925-21* (21,22). We obtained an estimate of 3.2% modern contamination. As with the mtDNA genome we filtered reads with mismatches in the first or last three nucleotides, taking forward only those reads for additional population genetics analyses. After applying both filters, the resultant mean depth of coverage for the mitochondrial genome was 4.038x and 0.029x for the nuclear genome.

2.4 Uniparental Markers and sex Determination analyses

The mtDNA haplogroup was determined using *SAMtools* pileup tool calling the positions defined in the *Phylotree* database (23). We used a genome browser (*IGV*.v2.4.14) to study the genomic context of each possible SNP (24). Only those SNPs that were present in two or more reads, and those which were not located at the ends of the reads, were considered. The contamination was estimated by calculating the ratio of discordant reads at haplogroup-diagnostic positions. Molecular sex was assigned with Ry_compute (22), a script designed for the sex identification of low coverage individuals (Fig. S3).

2.5 Population Genetics Analysis

Principal Component Analysis (PCA) was performed using *SmartPCA* in *EIG* v6.0.1 with a subset of modern individuals from the Human Origins dataset (25). This subset contained 434 present-day Europeans and 616,938 autosomal SNPs, plus our sample (Figure 2). The Marat sample was projected using the option *lsqproject*. We also considered the Marat sample projected into an expanded dataset of West Eurasian populations (Fig. S4). As projected individuals' components tend to 0, we also carried out a control analysis using Han Chinese,

151 French and Marat (Fig. S5). The results were visualised using the R package *GGplot2* (26). This
152 dataset confirmed that Marat is not artefactually placed at the centre of the plot.
153
To formally test the relationship of the Marat sample to relevant geographic regions we
calculated *f4* statistics of the form *f4*(Mbuti, Marat; X. Y) where X and Y are tested for
combinations of possible ancestral sources: Sardinian, French, English, Italian_North, Basque,
Spanish. *f4* values were calculated in qpDstat of AdmixTools v.5.0 (27) with statistical
significance assessed through Z-scores following jack-knife resampling (Table S3). This statistic
tests the covariance in allele frequency differences between an African out-group (Mbuti) and
Marat relative to the clade formed by X and Y. Positive values of *f4* indicate a closer affinity of
Marat to Y relative to X, with negative values indicating a closer relationship of Marat to X
relative to Y.
163
We additionally ran an unsupervised clustering analysis using *ADMIXTURE* v1.3 and another
subset of the Human Origins dataset (28). This subset included 881 individuals from Europe,
West Asia and North Africa typed over 616,938 shared autosomal SNPs. We filtered the
dataset by removing SNPs in high linkage disequilibrium using PLINK.v1.9 (29), removing all
SNPs with a r^2 threshold of 0.4 within a 200 SNP sliding window, advancing by 50 SNPs each
time. We performed the clustering analysis using K values ranging from 1 to 10, with 10
replicates for each value of K. We selected K according to the lowest cross-validation error
value (K=4). The *ADMIXTURE* results at K=4 were visualised using Pong (30) (Figure 2).
172
2.6 Metagenomic analysis
We first removed adapters and merged the paired-end reads into longer single-end sequences
using AdapterRemoval v2 (31). We removed PCR duplicates with exact sequence identity using
dedupe from the BBMap suite of tools (https://sourceforge.net/projects/bbmap/). We
subsequently used the default preprocessing pipeline designed for metaMix which consists of
removing human and rRNA sequences using bowtie2 followed by megaBLAST, as well as low
quality and low complexity reads using prinseq (32) (-lc_method dust -lc_threshold 7 -
min_qual_mean 15). The number of reads filtered at each step are provided in Table S4. We
screened the remaining high quality DNA reads for the presence of possible pathogens using
both KrakenUniq (33) against the Kraken database compiled in Lassalle et al 2018 (34) and
metaMix (35) using megaBLAST and a local custom database consisting of the RefSeq
sequences of bacteria, viruses, parasites, fungi and human, as of July 2019. KrakenUniq was
run with default parameters. The metaMix-nucleotide mode was run with the default read
support parameter of 10 reads was used (Table S5) and the default number of 12 MCMC
chains. The number of the MCMC iterations is automatically calculated by metaMix based on
the number of species to explore for each dataset, resulting in 10,000 iterations for the blood
sample and 3,230 iterations for the paper swab.
190
The relative proportion of reads assigned to different species by KrakenUniq and metaMix was
highly correlated; R^2=0.94 and R^2=0.82, for the blood stain and the unstained paper,
respectively (Fig. S6). However, metaMix tended to assign a higher number of reads to
individual species, closer to the number found by mapping directly to the microbial genomes
and we observed important discrepancies for the number of reads assigned to some of the
species (Table S6). Additionally, metaMix results for both the blood stain and the unstained
paper consisted of fewer species compared to KrakenUniq, even when the same read support
threshold was applied to KrakenUniq, indicating increased specificity due to the MCMC
exploration of the species space, that comes at an increased computational cost.
200
In order to compare the accuracy of the two assignment tools, we further explored the
presence of clinically relevant species by mapping the quality-filtered subset of reads (Table

S2) used in metagenomic assignment against the reference genomes of different candidate
genera of fungi and bacteria using *bowtie2* (10) and *BWA* v.07.3 (Fig. S7-S13). For all reads
mapping to individual reference genomes, mapDamage v2 (13) was also run to assess evidence
of nucleotide mis-incorporation characteristic of post-mortem damage. These mapping results
were systematically supporting the metaMix assignments over those obtained with
KrakenUniq (Table S6). This led us to rely on metaMix for all metagenomic assignments
presented in the paper.
210
211 Besides testing for the presence and absence of species, we tested whether some
212 microorganisms were overrepresented in the blood stain compared to the unstained section of
213 the paper using a one-sided binomial test and a significance threshold of 0.95 (Table S5).
214
As an additional control, we also conducted metagenomic analysis of two publicly available
ancient metagenomes obtained from parchment of comparable age to the Marat newspaper
(36). We followed the same pre-processing pipeline described for the Marat samples, first
removing adapters and PCR duplicates before employing the default metaMix pre-processing
pipeline, this time removing reads that mapped to either the human or sheep, cow and goat
reference genomes. As before, metaMix-nucleotide mode was run with with a read support
parameter of 10 reads and with 12 MCMC chains x 2,325 and 6,130 iterations respectively for
ERR466100 and ERR466101. We provide the breakdown of read filtering steps in Table S7 and
our raw metaMix results in Table S8.
224
2.7 Phylogenetic analysis
In the case of *Malassezia*, a phylogenetic analysis of the mitochondrial DNA genome with
available modern strains on the Short Read Archive (SRA) was performed. We called variant
positions using *GATK UnifiedGenotyper* (37) and generated a Maximum Likelihood tree using
RAxML-NG specifying a GTR substitution model and 100 bootstrap resamples (38). The tree
was rooted with *M. globosa* (Fig. S10)*.
231
We also conducted a phylogenetic analysis for *C. acnes*, combining our historical strain with all
C. acnes genomes deposited in the SRA covering the reference at an average depth >10x, and
with *C. namnetense* as an outgroup (SRR9222443). The only *C. acnes* genomes sequenced at
medium to high depth are those reported by Gomes et al 2017 (39). A Maximum Likelihood
tree was generated over the 21,751SNP alignment using RAxML-NG (Fig. S13) and clonal
complexes and phylotypes were assigned based on the PubMLST *C. acnes* definitions database
(https://pubmlst.org/bigsdb?db=pubmlst_pacnes_seqdef).

3 Results

3.1 Human ancestry analysis

We generated 568,623,176 DNA reads from the blood stain, of which 74,244,610 reads mapped to the human reference genome (Table S1). From these, we retrieved a complete human mitochondrial (mtDNA) genome at a mean depth of coverage of 4.038x and the nuclear genome at 0.029x (Table S1). The predominant mtDNA haplotype was H2a2a1f, although we found evidence of some additional mtDNA sequences, notably a K1a15 haplotype. The ratio of sexual chromosome to autosomal DNA reads indicated that the sample donor was male (Fig. S3).

The human DNA reads showed evidence of post-mortem deamination occurring in 1% of the ends of sequencing reads, indicating authentic ancient DNA damage (Fig. S1-S2). This is similar to the degree of damage that has been observed in aDNA obtained from other human specimens of a similar age (40). For further analyses we selected only those reads that displayed C to T or G to A substitutions at the 5' or 3' end, respectively. After this procedure, the degree of mitochondrial contamination was reduced to 0-0.01%.

To explore the ancestry of Marat in the context of modern European populations, we performed Principal Component Analysis (PCA) (Figure 2a and Fig. S4-5) and unsupervised clustering in ADMIXTURE (Figure 2b). Our sample projected among modern French individuals sampled from France in the population genetic analyses. This result is broadly compatible with proposed hypotheses relating to the ancestry of Marat (2). *f4* statistics suggest a closer affinity of Marat to modern Italian, English, Sardinian, Basque and French populations relative to those from Spain (Table S3). However, these trends are subtle and we note that mixed ancestries are difficult to discern, especially when only limited genetic data is available.

3.2 Metagenomic analysis

We conducted metagenomic species assignments with the 9,788,947 deduplicated, quality controlled and low complexity filtered DNA reads (combined merged and non-merged) that did not map to the human genome (see Methods and Table S4). We used metaMix (35), a Bayesian mixture model framework developed to resolve complex metagenomic mixtures, which classified ~9% of the non-human reads into 1,328 microbial species (Table S5). The species assignments were replicated with KrakenUniq (33), which led to largely consistent, if less accurate, results (~7% classified into 3,213 species, Fig. S6, Table S6). Thus, we relied on the metaMix species assignments throughout the paper, unless stated otherwise.

We detected the presence of a wide range of microorganisms, including some expected to develop on decaying cellulose and/or dried blood, but also others recognized as opportunistic human pathogens from the following bacterial genera: *Acidovorax, Acinetobacter, Burkholderia, Chryseobacterium, Corynebacterium, Cutibacterium, Micrococcus, Moraxella, Paraburkholderia, Paracoccus, Pseudomonas, Rothia, Staphylococcus, Streptococcus* and the fungal genera *Aspergillus, Penicillium, Talaromyces* and *Malassezia* as well as HPV (type 179 and type 5) and HHV6B viruses, albeit the latter supported by a very low numbers of reads (Table S5-S6). Some of the DNA reads, notably from *Aspergillus glaucus*, *Cutibacterium acnes*, *Malassezia restricta* and *Staphylococcus epidermis* showed typical misincorporation patterns that are considered indicative of these sequences being authentically old (Fig. S7).

We additionally sequenced the swab taken from the unstained paper sample. In this case, only 96,252 pairs of reads were obtained (56,616 merged, 25,712 non-merged, 35,216

deduplicated and filtered combined merged and non-merged), with 52% of the reads that could be classified with metaMix into 66 species and 36% with KrakenUniq into 374 species, respectively (see Methods and Table S4). Although very little DNA could be retrieved from the section of the document that had not been blood-stained, we tried to identify microorganisms that were statistically significantly over-represented in the blood stain relative to the unstained paper. Amongst these and besides, as expected, *Homo sapiens*, different species of *Aspergillus* and *Acinetobacter* were significantly overrepresented in the blood stain (Table S5). It remains questionable however whether the unstained paper represents a suitable negative control given that the newspaper had been extensively manipulated by Marat. Significant over-representation of *Aspergillus spp.* and *Acinetobacter spp.* in the blood stain relative to the rest of the document could also be due to the blood providing better conditions for the growth of iron-limited microbes. Indeed, *Aspergillus spp.* and *Acinetobacter spp.* are commonly found in the environment but are also grown in blood agar. As such, it is plausible that these represent *post-mortem* contaminants. Indeed, for *Acinetobacter spp.* we identified no post-mortem damage pattern.

Metagenomic analysis of historical samples can be challenging as the resulting microbial communities typically comprise an unknown mixture of endogenous species as well as contaminants, both contemporary and modern. To mitigate this problem, we relied on a 'differential diagnostics' approach (Table 1), where we specifically tested for the presence of reads from pathogens that could plausibly have led to Marat's symptoms, most of which have been previously hypothesised in the literature (1–5). Such a differential approach is more stringent than the standard approach in clinical diagnostics aiming to identify the full list of microbes present in the samples after enforcement of a read-number threshold (41,42). Our approach allows limiting the number of species to be tested to a small list of plausible candidates. Second, the lack of detection of even one read from a focal microbial species by direct mapping falsifies the null hypothesis that it was not involved in the disease.

We did not identify a single sequencing read in either the blood stain or the unstained paper for the agents of syphilis, leprosy, scrofula (tuberculosis) and diabetic candidiasis (thrush) (Table 1, Table S5). We additionally tested for scabies, which is caused by burrowing of the mite *Sarcoptes scabiei* under the skin. Since the metagenomic reference database did not include arthropod genomes, this was tested separately by blasting all the non-human reads against the *Sarcoptes scabiei* genome (GCA_000828355.1). Again, we detected not a single read matching to *Sarcoptes scabiei*, which makes scabies an implausible cause for Marat's skin disease (Table 1, Table S5).

Conversely, metaMix recovered 15,926 and 83 filtered DNA reads from the blood stain and the unstained paper respectively, assigned to *Malassezia restricta* a fungal pathogen causing seborrheic dermatitis, which has been previously hypothesized as one of the most plausible causes for Marat's condition (1–4). Direct mapping of all reads to *M. restricta* (GCA_003290485.1) resulted in 19,194 reads from the blood stain dataset mapping over 17.17% of the reference genome. KrakenUniq failed to identify *M. restricta*, instead assigning 627 reads sequenced from the blood stain to *M. sympodialis*. However, further analysis of the *Malassezia* reads based on genome mapping pointed to most (80.3%) being uniquely assigned to *M. restricta* rather than *M. sympodialis* (Fig. S8). This allowed us to reconstruct a complete *M. restricta* mtDNA genome at 0.84X coverage. The *Malassezia* reads were evenly distributed along the full genome assembly supporting no mixing or misclassification of the species (Fig. S9).

We placed our Marat *M. restricta* mitochondrial genome in phylogenetic context by building a maximum likelihood phylogeny including our historical strain and available present-day mtDNA

343   *M. restricta* genomes. Although the total number of samples is small, the fact that the *M.*
344   *restricta* mtDNA molecule recovered from Marat's blood is placed basal to modern strains (Fig.
345   S10) and exhibits some post-mortem damage (Fig. S5) further support its authenticity.
346
We also recovered 587 filtered reads assigned by metaMix to *Staphylococcus aureus* in the
blood stain but none in the reads obtained from the unstained paper. The differential
representation in the two samples is not significantly different due to the far lower number of
reads in the unstained sample (Table S5). Although a common commensal, *S. aureus* is also a
frequent human pathogen and the leading cause of atopic eczema. In order to confirm the
metagenomic assignments to *S. aureus*, we mapped the raw microbial reads to a series of
reference genomes from various species in the *Staphylococcus* genus. This allowed us to
identify 888 reads mapping against the *S. aureus* reference genome, out of which 758 uniquely
mapped to *S. aureus* (Fig. S11). The presence of *S. aureus*, but with a relatively low number of
reads, may be compatible with a secondary infection by *S. aureus* rather than *S. aureus* being
the initial cause of Marat's condition. Alternatively, Marat, or someone who also handled the
newspaper, could have carried *S. aureus* as a skin commensal.
359
The most prevalent microbial species in the blood stain was *Cutibacterium acnes*
(formerly *Propionibacterium acnes* (43)), which was also present in the unstained paper (Table
S5). *C. acnes* is largely a commensal and part of the normal skin biota present on most healthy
adult humans' skin, including in association with *S. epidermis* which we also observe in our
sample (Table S5-S6, Fig. S11) (44). *C. acnes* is also a frequent contaminant in metagenomic
samples (45,46). However, *C. acnes* can also be involved in severe acneiform eruptions (47)
and we cannot exclude the possibility that it could have contributed to Marat's condition.
86,019 reads mapped to the *C. acnes* reference genome (GCF_000008345.1), yielding an
alignment of 3.4X average coverage (Fig. S12) and exhibiting modest post-mortem damage
(Fig. S7).
370
A phylogeny of Marat *C. acnes* with a collection of publicly available modern strains (39,46)
places our historic genome on a short branch falling basal to Type I strains, supporting its age
and authenticity (Fig. S13). This phylogenetic placement suggests our Marat strain falls into *C.*
acnes phylotype I (*C. acnes subsp. acnes*) rather than II (*C. acnes subsp. defendens*). Whilst our
Marat strain does not cluster with phylotype Ia, the type more commonly associated with skin
surface associated acne vulgaris (48), its position, basal to Type Ib strains cannot exclude its
involvement in soft or deep tissue infections (49).
378
Delineating contaminants and commensals from plausible pathogens remains challenging from
this type of data source, in particular due to the absence of a suitable control. To alleviate this
issue, we conducted full taxonomic assignments of two ancient metagenomes generated from
historical parchment samples dating to the 17th and 18th centuries (PA1 and PA2 respectively)
(36). Although these samples were obtained from livestock (ruminant) skins whereas we are
working with cellulose paper, we anticipate that they may have been used and handled in a
comparable way to the newspaper Marat was annotating. In this way they represent what can
be considered as the most biologically comparable ancient metagenomes available to date. An
equivalent metaMix analysis applied to these filtered sequencing reads (Table S7) identified
not a single read assigned to *M. restrica*, *S. aureus* or *C. acnes* (Table S8). We therefore do not
systematically expect a significant number of reads for the three species we suggest as most
plausible candidates for Marat's condition.
391

4 Discussion

Over the last decade, ancient-pathogen genomics has made great progress by borrowing technological advances originally developed for the study of human ancient DNA (50,51). Although most microbial data has been secondarily generated from the sequencing of ancient human bones or teeth (51–54) other, rare samples, such as preserved tissues (55,56) or microscope slides from antique medical collections have been analysed (50,57). We are aware of no previous attempt to leverage ancient DNA technology to diagnose infections in historical characters, despite previous sequencing of remains from other prominent historical figures such as King Richard III and the putative blood of Louis XVI (58,59).

In this work we analysed both human and 'off-target' microbial reads to shed light on an important historical figure of the French Revolution and his skin condition. Due to the loss of Marat's remains after their removal from the Panthéon in February 1795, the paper stained with his blood likely represents the only available biological material to study both his ancestry and the cause of his skin condition. Although second-generation sequencing techniques have been applied to the analysis of ancient parchments (60) our work represents the first instance where this methodological approach has been applied to old cellulose paper.

The presence and relative abundance of different microorganisms in the documents Marat was annotating is affected by their endogenous presence as well as contemporary and modern contamination both for the blood and unstained sample. Some microorganisms present in the samples might reflect skin microbiome signatures. Whilst some other microorganisms represent environmental contaminants and are likely unrelated to Marat's condition. In order to identify the most likely candidates for Marat's condition we tested a set of proposed diagnoses, which we considered as plausible if we detected at least one read assigned to the causative infectious agent (Table 1).

Potential conditions for which we detected not a single supportive read included syphilis, tuberculosis (scrofula), leprosy, diabetic candidiasis or scabies. We appreciate that absence of evidence for an infectious agent does not constitute incontrovertible evidence of its absence. Moreover, it is not uncommon for metagenomic diagnostics applied to clinical samples to fail to identify reads from the likely infectious agent above the predefined diagnostic threshold, or even fail to detect any read at all (42,43). As such, the absence of reads from a putative pathogen makes it less plausible as the agent of Marat's condition but does not definitely rule them out.

Conversely, we detected and validated microbial reads for two of the conditions we tested, seborrheic dermatitis (*Malassezia spp.*), atopic dermatitis (*Staphylococcus aureus*) and cannot exclude severe acneiform eruptions (*Cutibacterium acnes*) as a third, given the age and phylogenetic position of the *C. acnes* genome we obtained. For all three cases, the number of reads would have exceeded the threshold suggested for detection in clinical metagenomic diagnostic (42,43)(41), even when considering the swab from the unstained paper as a control.

The presence of *Malassezia restricta* is of particular interest because this fungus is specialized to live on the skin (61). Although also a common commensal and contaminant in metagenomic studies, *Malassezia* has been described in various skin conditions, including dandruff, atopic eczema, folliculitis and seborrheic dermatitis (62,63). Interestingly, the latter symptoms would fit those described in Marat (5). The *M. restricta* reads we identified were not statistically significantly overrepresented in the blood's stain relative to the unstained paper, although they could be expected to be present in both samples if someone heavily infected was holding the newspaper. Although we cannot confidently claim the reads in Marat's blood are directly

444 associated with Marat himself, we do identify post-mortem damage in these reads and a
445 phylogenetic placement in a modern mitochondrial DNA phylogeny consistent with these
446 reads being indeed old (Fig. S7, Fig. S10). We also do not systematically expect the presence of
447 *M. restricta* on parchment of a similar age (Table S8).
448
Also of possible interest is the widespread presence of *Cutibacterium acnes subsp. acnes*,
which although a common commensal or contaminant can also be implicated in severe
acneiform eruptions, which constitutes the top hit in the blood sample and falls basal to
phylotype I strains currently in circulation. As with *M. restricta*, we do not observe a single *C.*
acnes read in two biologically equivalent historic parchment metagenomes (Table S8).
Staphylococcus aureus, which is frequently detected in cases of atopic dermatitis, is also
present in reads obtained from the blood's stain, although in fairly low number.
456
Whilst our results do not allow us to reach a definite diagnosis of Marat's condition, they
allowed us to cast doubt on several previous hypotheses and provide, using all the available
evidence, some plausible aetiologies. We suggest that Marat could have been suffering from
an advanced fungal or polymicrobial infection, either primary or secondary to another
condition. Future metagenomic analysis of additional documents in Marat's possession during
his assassination could help confirm the microbial composition found in this study and
strengthen these observations.
464
465 Our work further illustrates the potential of sequencing technologies for the generation of
466 (meta-)genomic information from difficult, singular samples and opens new avenues to
467 address medical hypotheses of major historical interest.
468

469 **Acknowledgments and Funding**
470
This work was supported by Obra Social ''La Caixa'' and Secretaria d'Universitats i Recerca
(GRC2017-SGR880) (T.M.-B. and C.L.-F.), BFU2017-86471-P and PGC2018-101927-B-I00
(MINECO/FEDER, UE) (T.M-B) and PGC2018-095931-B-100 (MINECO/FEDER, UE) (C.L.-F.). T.M-
B. is also supported by a U01 MH106874 grant and Howard Hughes International Early Career
and CERCA Programme del Departament d'Economia i Coneixement de la Generalitat de
Catalunya. S.M is funded by a Welcome Trust post-doctoral fellowship (206478/Z/17/Z). L.v.D
and F.B. acknowledge financial support from the Newton Fund UK-China NSFC initiative (grant
MR/P007597/1) and the BBSRC (equipment grant BB/R01356X/1). Full metagenomic reads are
available at NCBI under BioProject ID XXXXX.

Author Contributions
482
483 P.C., L.v.D., F.B. and C.L.-F. conceived and designed the study; P.C. and C.L.B sourced the
484 newspaper; C.B., M.A-E. and E.L. developed and performed laboratory analysis; T.d.-D., L.v.D.,
485 S.M. analysed data and performed computational analyses; T.d.-D., L.v.D., S.M., F.B., and C.L.-
486 F. wrote the paper with inputs from all co-authors.
487

488 **Competing Interests**
489
490 The authors have no competing interests to declare.

References

1. Bayon HP. The Medical Career of Jean-Paul Marat. Proc R Soc Med. 1945;39(1):39–44.
2. Cohen JHL, Cohen EL. Doctor Marat and His Skin. Med Hist. 1958;2(04):281–6.
3. Dotz W. Jean Paul Marat. His life, cutaneous disease, death, and depiction by Jacques Louis David. Am J Dermatopathol. 1979;1(3):247–50.
4. Jelinek JE. Jean-Paul Marat. The differential diagnosis of his skin disease. Am J Dermatopathol. 1979;1(3):251–2.
5. Dale PM. Medical Biographies. The Ailments of Thirty-Three Famous Persons. Norman, editor. Press, University of Oklahoma; 1952.
6. Coto-Segura C, Coto-Segura P, Santos-Juanes J. The Skin of a Revolutionary. JAMA Dermatology. 2011;147(5):539.
7. Carøe C, Gopalakrishnan S, Vinner L, Mak SST, Sinding MHS, Samaniego JA, et al. Single-tube library preparation for degraded DNA. Johnston S, editor. Methods Ecol Evol. 2018;9(2):410–9.
8. Martin M. Cutadapt removes adapter sequences from high-throughput sequencing reads. EMBnet.journal. 2011;17(1):10.
9. Li H, Durbin R. Fast and accurate short read alignment with Burrows-Wheeler transform. Bioinformatics. 2009;25(14):1754–60.
10. Langmead B, Salzberg SL. Fast gapped-read alignment with Bowtie 2. Nat Methods. 2012;9(4):357–9.
11. Taron UH, Lell M, Barlow A, Paijmans JLA. Testing of Alignment Parameters for Ancient Samples: Evaluating and Optimizing Mapping Parameters for Ancient Samples Using the TAPAS Tool. Genes (Basel). 2018;9(3).
12. Broad Institute. Picard.
13. Jónsson H, Ginolhac AA, Schubert M, Johnson PLFF, Orlando L, Jonsson H, et al. mapDamage2.0: fast approximate Bayesian estimates of ancient DNA damage parameters. In: Bioinformatics. England, England; 2013. p. 1682–4.
14. Skoglund P, Northoff BH, Shunkov M V, Derevianko AP, Pääbo S, Krause J, et al. Separating endogenous ancient DNA from modern day contamination in a Siberian Neandertal. Proc Natl Acad Sci U S A. 2014;111(6):2229–34.
15. García-Alcalde F, Okonechnikov K, Carbonell J, Cruz LM, Götz S, Tarazona S, et al. Qualimap: Evaluating next-generation sequencing alignment data. Bioinformatics. 2012;28(20):2678–9.
16. Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, et al. The Sequence Alignment/Map format and SAMtools. Bioinformatics. 2009;25(16):2078–9.
17. Lazaridis I, Patterson N, Mittnik A, Renaud G, Mallick S, Kirsanow K, et al. Ancient human genomes suggest three ancestral populations for present-day Europeans. Nature. 2014;513(7518):409–13.
18. Lazaridis I, Nadel D, Rollefson G, Merrett DC, Rohland N, Mallick S, et al. Genomic insights into the origin of farming in the ancient Near East. Nature. 2016;536(7617):419–24.
19. Renaud G, Slon V, Duggan AT, Kelso J. Schmutzi: Estimation of contamination and endogenous mitochondrial consensus calling for ancient DNA. Genome Biol. 2015;16(1):1–18.
20. Pierre L. JVarkit: java-based utilities for Bioinformatics. Figshare. 2015.
21. Korneliussen TS, Albrechtsen A, Nielsen R. ANGSD: Analysis of Next Generation Sequencing Data. BMC Bioinformatics. 2014;15(1):356.
22. Skoglund P, Storå J, Götherström A, Jakobsson M. Accurate sex identification of ancient human remains using DNA shotgun sequencing. J Archaeol Sci. 2013;40(12):4477–82.
23. van Oven M. PhyloTree Build 17: Growing the human mitochondrial DNA tree. Forensic Sci Int Genet Suppl Ser. 2015;5:e392--e394.

24. Robinson JT, Thorvaldsdóttir H, Winckler W, Guttman M, Lander ES, Getz G, et al.
Integrative genomics viewer. Nat Biotechnol. 2011;29(1):24–6.

25. Patterson N, Price AL, Reich D. Population structure and eigenanalysis. PLoS Genet.
2006;2(12):e190.

26. Wickham H. ggplot2: Elegant Graphics for Data Analysis. Springer-Verlag New York;
2016.

27. Patterson N, Moorjani P, Luo Y, Mallick S, Rohland N, Zhan Y, et al. Ancient Admixture in
Human History. Genetics. 2012;192(3):1065–93.

28. Alexander DH, Novembre J, Lange K. Fast model-based estimation of ancestry in
unrelated individuals. Genome Res. 2009;19(9):1655–64.

29. Purcell S, Neale B, Todd-Brown K, Thomas L, Ferreira MAR, Bender D, et al. PLINK: A
Tool Set for Whole-Genome Association and Population-Based Linkage Analyses. Am J
Hum Genet. 2007;81(3):559–75.

30. Behr AA, Liu KZ, Liu-Fang G, Nakka P, Ramachandran S. pong: fast analysis and
visualization of latent clusters in population genetic data. Bioinformatics.
2016;32(18):2817–23.

31. Schubert M, Lindgreen S, Orlando L. AdapterRemoval v2: rapid adapter trimming,
identification, and read merging. BMC Res Notes. 2016;9:88.

32. Schmieder R, Edwards R. Quality control and preprocessing of metagenomic datasets.
Bioinformatics. 2011;27(6):863–4.

33. Breitwieser FP, Baker DN, Salzberg SL. KrakenUniq: confident and fast metagenomics
classification using unique k-mer counts. Genome Biol. 2018;19(1):198.

34. Lassalle F, Spagnoletti M, Fumagalli M, Shaw L, Dyble M, Walker C, et al. Oral
microbiomes from hunter-gatherers and traditional farmers reveal shifts in commensal
balance and pathogen load linked to diet. Mol Ecol. 2018;27(1):182–95.

35. Morfopoulou S, Plagnol V. Bayesian mixture analysis for metagenomic community
profiling. Bioinformatics. 2015;31(18):2930–8.

36. Teasdale MD, van Doorn NL, Fiddyment S, Webb CC, O'Connor T, Hofreiter M, et al.
Paging through history: parchment as a reservoir of ancient DNA for next generation
sequencing. Philos Trans R Soc B Biol Sci. 2015;370(1660):20130379.

37. McKenna A, Hanna M, Banks E, Sivachenko A, Cibulskis K, Kernytsky A, et al. The
Genome Analysis Toolkit: A MapReduce framework for analyzing next-generation DNA
sequencing data. Genome Res. 2010;20(9):254–60.

38. Kozlov AM, Darriba D, Flouri T, Morel B, Stamatakis A. RAxML-NG: a fast, scalable and
user-friendly tool for maximum likelihood phylogenetic inference. Bioinformatics. 2019;

39. Gomes A, van Oosten M, Bijker KLB, Boiten KE, Salomon EN, Rosema S, et al. Sonication
of heart valves detects more bacteria in infective endocarditis. Sci Rep.
2018;8(1):12967.

40. Rasmussen M, Guo X, Wang Y, Lohmueller KE, Rasmussen S, Albrechtsen A, et al. An
Aboriginal Australian genome reveals separate human dispersions into Asia. Science
(80-). 2011;334:94–8.

41. Wilson MR, Sample HA, Zorn KC, Arevalo S, Yu G, Neuhaus J, et al. Clinical Metagenomic
Sequencing for Diagnosis of Meningitis and Encephalitis. N Engl J Med.
2019;380(24):2327–40.

42. Miller S, Naccache SN, Samayoa E, Messacar K, Arevalo S, Federman S, et al. Laboratory
validation of a clinical metagenomic sequencing assay for pathogen detection in
cerebrospinal fluid. Genome Res. 2019;29(5):831–42.

43. Scholz CFP, Kilian M. The natural history of cutaneous propionibacteria, and
reclassification of selected species within the genus Propionibacterium to the proposed
novel genera Acidipropionibacterium gen. nov., Cutibacterium gen. nov. and
Pseudopropionibacterium gen. nov. Int J Syst Evol Microbiol. 2016;66(11):4422–32.

44. Dreno B, Martin R, Moyal D, Henley JB, Khammari A, Seite S. Skin microbiome and acne

vulgaris: Staphylococcus, a new actor in acne. Exp Dermatol. 2017;26(9):798–803.

45.   Salter SJ, Cox MJ, Turek EM, Calus ST, Cookson WO, Moffatt MF, et al. Reagent and
597   laboratory contamination can critically impact sequence-based microbiome analyses.
598   BMC Biol. 2014;12(1):87.

599   46.   Mollerup S, Friis-Nielsen J, Vinner L, Hansen TA, Richter SR, Fridholm H, et al.
600   Propionibacterium acnes: Disease-causing agent or common contaminant? detection in
601   diverse patient samples by next- generation sequencing. Burnham C-AD, editor. J Clin
602   Microbiol. 2016;54(4):980–7.

603   47.   Platsidaki E, Dessinioti C. Recent advances in understanding Propionibacterium acnes (
604   Cutibacterium acnes) in acne. F1000Research. 2018;7:F1000 Faculty Rev-1953.

605   48.   Lomholt HB, Kilian M. Population Genetic Analysis of Propionibacterium acnes Identifies
606   a Subpopulation and Epidemic Clones Associated with Acne. PLoS One.
607   2010;5(8):e12277.

608   49.   Nazipi S, Stodkilde-Jorgensen K, Scavenius C, Bruggemann H. The Skin Bacterium
609   Propionibacterium acnes Employs Two Variants of Hyaluronate Lyase with Distinct
610   Properties. Microorganisms. 2017;5(3):57.

611   50.   Gelabert P, Sandoval-Velasco M, Olalde I, Fregel R, Rieux A, Escosa R, et al.
612   Mitochondrial DNA from the eradicated European Plasmodium vivax and P. falciparum
613   from 70-year-old slides from the Ebro Delta in Spain. Proc Natl Acad Sci U S A.
614   2016;113(41):11495–500.

615   51.   Rasmussen S, Allentoft ME, Nielsen K, Orlando L, Sikora M, Sjögren K-G, et al. Early
616   Divergent Strains of Yersinia pestis in Eurasia 5,000 Years Ago. Cell. 2015;163(3):571–
617   82.

618   52.   Rascovan N, Sjögren K-G, Kristiansen K, Nielsen R, Willerslev E, Desnues C, et al.
619   Emergence and Spread of Basal Lineages of Yersinia pestis during the Neolithic Decline.
620   Cell. 2019;176(1):295-305.e10.

621   53.   Margaryan A, Hansen HB, Rasmussen S, Sikora M, Moiseyev V, Khoklov A, et al. Ancient
622   pathogen DNA in human teeth and petrous bones. Ecol Evol. 2018;8(6):3534–42.

623   54.   Mühlemann B, Jones TC, Damgaard P de B, Allentoft ME, Shevnina I, Logvin A, et al.
624   Ancient hepatitis B viruses from the Bronze Age to the Medieval period. Nature.
625   2018;557(7705):418–23.

626   55.   Marciniak S, Prowse TL, Herring DA, Klunk J, Kuch M, Duggan AT, et al. Plasmodium
627   falciparum malaria in 1st–2nd century CE southern Italy. Curr Biol. 2016;26(23):R1220–
628   2.

629   56.   Devault AM, Golding GB, Waglechner N, Enk JM, Kuch M, Tien JH, et al. Second-
630   Pandemic Strain of Vibrio cholerae from the Philadelphia Cholera Outbreak of 1849 . N
631   Engl J Med. 2014;370(4):334–40.

632   57.   de-Dios T, van Dorp L, Gelabert P, Carøe C, Sandoval-Velasco M, Fregel R, et al. Genetic
633   affinities of an eradicated European Plasmodium falciparum strain. Microb Genomics.
634   2019;mgen000289.

635   58.   King TE, Fortes GG, Balaresque P, Thomas MG, Balding D, Delser PM, et al. Identification
636   of the remains of King Richard III. Nat Commun. 2014;5:5631.

637   59.   Olalde I, Sánchez-Quinto F, Datta D, Marigorta UM, Chiang CWK, Rodríguez JA, et al.
638   Genomic analysis of the blood attributed to Louis XVI (1754–1793), king of France. Sci
639   Rep. 2014;4:4666.

640   60.   Teasdale M, Doorn N, Fiddyment S, Webb C, O'Connor T, Hofreiter M, et al. Paging
641   through history: Parchment as a reservoir of ancient DNA for next generation
642   sequencing. Philos Trans R Soc B Biol Sci. 2015;370.

643   61.   Saunders CW, Scheynius A, Heitman J. Malassezia Fungi Are Specialized to Live on Skin
644   and Associated with Dandruff, Eczema, and Other Skin Diseases. PLOS Pathog.
645   2012;8(6):e1002701.

646   62.   Ashbee R, Bignell E, editors. Pathogenic yeasts. In: The yeast handbook. Berlin

647   Heidelberg: Springer-Verlag; 2010. p. 209–30.

648   63.   Gupta AK, Batra R, Bluhm R, Boekhout T, Dawson TL. Skin diseases associated with

649   Malassezia species. J Am Acad Dermatol. 2004;51(5):785–98.

651　**Figures and Tables**
652

653
654
655　**Figure 1:** a) "La mort de Marat"; portrait of Jean-Paul Marat after his assassination, by Jacques-
656　Louis David (1793). Preserved at Musées Royaux des Beaux-Arts de Belgique, Brussels. b)
657　Sampling the page of l'*Ami du Peuple* stained with Marat's blood that has been analysed.
658

659
660　**Figure 2:** a) Principal Component Analysis (PCA) of modern human European populations with
661　Marat's ancient DNA reads projected. Symbols provide the country and region, where
662　provided, as given in the legend at right. b) Admixture analysis with modern European samples
663　and Marat. Both analyses are coherent with Marat's suggested French and Italian combined
664　ancestry.

Table 1: List of diseases tested for associated agents and presence in the blood stain and the unstained paper samples. The following symbols denote the abundance of reads for each infectious agent tested. ✓: present; ✓✓: top ten; ✓✓✓: top hit; ✗: absent.

Disease	Pathogen	Blood	Unstained paper
Syphilis	*Treponema pallidum*	✗	✗
Scrofula (tuberculosis)	*Mycobacterium tuberculosis*[1]	✗	✗
Leprosy	*Mycobacterium leprae*	✗	✗
Diabetic candidiasis (thrush)	*Candida* sp.	✗	✗
Scabies	*Sarcoptes scabiei*	✗	✗
Seborrheic dermatitis	*Malassezia* sp.	✓✓	✓
Atopic dermatitis	*Staphylococcus aureus*	✓	✗
Severe acneiform eruptions	*Cutibacterium acnes*	✓✓✓	✓✓

[1] Scrofula can also be caused by other Mycobacteria in particular *M. scrofulaceum* and *M. avium intracellulare*, which are also absent from both samples.