GLP-1 Notch - LAG-1 CSL control of the germline stem cell fate is mediated by

transcriptional targets *lst-1* and *sygl-1*

Jian Chen[1], Ariz Mohammad[1], Nanette Pazdernik[1,5], Huiyan Huang[2], Beth Bowman[3,6],

Eric Tycksen[4], Tim Schedl[1*]

[1] Department of Genetics, Washington University School of Medicine

[2] Department of Pediatrics, Washington University School of Medicine

[3] Department of Biology, Emory University

[4] Genome Technology Access Center, McDonnell Genome Institute, Washington

University School of Medicine

[5] Current address, Integrated DNA Technologies

[6] Current address, Vanderbilt University

*Corresponding author: Tim Schedl

Department of Genetics

Washington University School of Medicine

Campus Box 8232

4523 Clayton Avenue,

St Louis, MO, 63110

ts@wustl.edu

Running title: Germline GLP-1 signaling transcriptional targets

28 **Key words**: germline stem cell, transcription, GLP-1 signaling, *lag-1*, *lst-1*, *sygl-1*

29 **Abstract**

Stem cell systems are essential for development and maintenance of polarized tissues.

Intercellular signaling pathways control stem cell systems, where niche cells signal stem

cells to maintain the stem cell fate/self renewal and inhibit differentiation. In the *C.*

elegans germline stem cell system, GLP-1 Notch signaling specifies the stem cell fate.

However, the downstream transcriptional targets of GLP-1 signaling that mediate the

stem cell fate have not been fully enumerated. We employed a genome-wide approach

to uncover transcriptional targets of GLP-1 signaling - the intersection of genes

identified as directly bound by LAG-1, the *C. elegans* Notch pathway sequence-specific

DNA binding protein, from ChIP-seq experiments, with genes identified as requiring

GLP-1 signaling for RNA accumulation, from RNA-seq analysis. *lst-1* and *sygl-1*, genes

previously identified as transcriptional targets from a bioinformatic candidate gene

approach, were bound by germline LAG-1 and their expression dependent on *glp-1* and

germline *lag-1* activity. No additional genes were identified as both bound by LAG-1 and

whose mRNA level was dependent on *glp-1* and *lag-1*. Genes were identified as likely

secondary effects of GLP-1 signaling with the properties that their *glp-1* dependent

mRNA accumulation could be explained by a requirement for *lst-1* and *sygl-1* activity

and their lack of LAG-1 binding. Furthermore, *glp-1* dependent peak accumulation of

FBF-2, which promotes the stem cell fate, is explained by a requirement for *lst-1* and

sygl-1 activity. Finally, we showed that *lag-1* is germline autonomously required for the

stem cell fate and that elevated LAG-1 accumulation is spatially limited to the stem cell

region by posttranscriptional regulation that, in part, requires *lst-1* and *sygl-1*. These

findings are consistent with the possibility that *lst-1* and *sygl-1* are the sole germline

GLP-1 signaling mRNA transcriptional targets, which largely or completely mediate the

stem cell fate.

Introduction

Stem cell systems are required for the development and maintenance of polarized tissues, controlling the position, number and timing of differentiated cell type production. Stem cell systems have non-stem niche cells that signal nearby cells to promote the stem cell fate/self renewal and to inhibit differentiation. Niche – stem cell signaling pathways include Notch, BMP, Wnt and JAK/Stat (Morrison and Kimble 2006; Morrison and Spradling 2008; Simons and Clevers 2011). A deep understanding of how a stem cell system works requires knowledge of the full repertoire of gene products that are immediately downstream of the signaling pathway, which then mediate stem cell fate/self renewal and inhibit differentiation.

Here we focus on the *C. elegans* germline stem cell system, which shares a number of features with other stem cell systems (Hubbard and Schedl, 2019). Niche - germline stem cell signaling employs the Notch pathway, which has been extensively studied in *C. elegans* (Greenwald and Kovall 2013). The worm germline is a polarized tube-shaped cellular assembly line designed for the rapid production of gametes. The germline is capped by the distal tip cell (DTC), which is a large somatic niche cell that polarizes germline cellular organization. Germ cells adjacent to the DTC are in a region called the progenitor zone, which distally contains stem cells, then cells completing a terminal mitotic cell cycle and cells undergoing meiotic S-phase, followed proximally by cells undergoing the earliest stages of meiotic prophase, leptotene and zygotene (**Fig 1**; Kimble and Crittenden 2005; Kimble and Seidel 2008; Hubbard and Schedl, 2019). The

DTC expresses two Notch pathway DSL (for Delta, Serate, LAG-2) ligands, LAG-2 and

APX-1 (Henderson *et al.* 1994; Tax and Thomas 1994; Nadarajan *et al.* 2009). *C.*

elegans has two Notch receptors, GLP-1 and LIN-12 (Lambie and Kimble 1991;

Greenwald and Kovall 2013). GLP-1 is expressed in progenitor zone germ cells and

continuously required to promote the germline stem cell fate. Genetic loss of *glp-1* in

larval or adult stages results in loss of all germline stem cells because of their

premature entry into meiotic prophase (Austin and Kimble 1987; Crittenden *et al.* 1994).

Conversely, g̲ain of f̲unction (gf) mutations in *glp-1* result in a tumorous germline, with a

vast excess of stem cells and reduced or no meiotic prophase cells (Berry *et al.* 1997;

Pepper *et al.* 2003). *lin-12* functions only in somatic cell fate specification, including

mediating the anchor cell – ventral uterine (AC/VU) decision and 2^o vulval cell fate

specification. *lin-12* and *glp-1* function redundantly in late embryonic fates, and their

simultaneous loss results in an early larval lethal phenotype called Lag (for l̲in-12 a̲nd

glp-1; (Lambie and Kimble 1991)). The current model is that when the DTC presenting

LAG-2 and APX-1 interacts with GLP-1 expressed germ cells, ligand dependent

cleavage of the receptor generates the GLP-1 i̲ntrac̲ellular d̲omain, GLP-1(ICD), which

translocates to the nucleus and associates with the sequence specific DNA binding

protein LAG-1. LAG-1 is a founding member of the CSL family of DNA binding proteins

[for C̲BF1 (or RBPJ) in mammals, S̲u(H) in Drosophila and L̲AG-1](Christensen *et al.*

1996). Complete loss of *lag-1* results in L1 larval lethality, reflecting LAG-1 functioning

as the DNA binding co-factor for both GLP-1(ICD) and LIN-12(ICD), while a *lag-1*

hypomorphic mutant results in incompletely penetrant loss of germline stem cells due to

premature entry into meiotic prophase (Lambie and Kimble 1991; Qiao *et al.* 1995).

LAG-1 and other CSL proteins share the same *in vitro* DNA binding site (GTGGGAA,

LAG-1/CSL binding site hereafter) (Brou *et al.* 1994; Tun *et al.* 1994; Christensen *et al.*

1996). GLP-1(ICD), bound to LAG-1, and associated with SEL-8 (also called LAG-3)

that functions similarly to Drosophila Mastermind, forms an activation complex that

transcribes GLP-1 signaling targets. LAG-1 is thus central to GLP-1 signaling as its

sequence specific DNA binding determines which genes are transcriptional targets.

Through its transcriptional targets, GLP-1 signaling pathway promotes the stem cell

fate, at least in part, by inhibiting three parallel pathways that promote meiotic

development, the GLD-1 pathway, the GLD-2 pathway and SCF^{PROM-1} (**Fig 1**).

Two germline GLP-1 signaling transcriptional targets, *lst-1* and *sygl-1*, have been

identified that are redundantly required to promote the stem cell fate (Kershner *et al.*

2014; Lee *et al.* 2016). A number of lines of evidence support this identification. First,

genetic manipulation of *lst-1* and *sygl-1* give the same phenotypes as manipulation of

glp-1: the *lst-1 sygl-1* double null mutant has the identical premature meiotic entry of all

cells in the L2 stage as *glp-1* null; loss of *lst-1* and *syg-1* later in larval and early adult

stages results in all stem cells entering meiotic prophase as is observed with loss of *glp-*

1; and ubiquitous overexpression of *lst-1* or *sygl-1* in the germline results in an

tumorous phenotype similar to *glp-1(gf)* (Kershner *et al.* 2014; Shin *et al.* 2017).

Second, epistasis analysis place *lst-1* and *sygl-1*, like *glp-1*, upstream of the meiotic

entry pathway genes. Loss of any two of the meiotic entry pathway genes (e.g., *gld-2*

gld-1 double null mutant) results in a tumorous germline due to a failure of germ cells to

enter meiotic prophase. The *gld-2 gld-1; glp-1* triple null mutant and the *gld-2 gld-1 lst-1*

sygl-1 quadruple null mutant have a tumorous germline phenotype, because the

premature meiotic entry of stem cells in the absence of *glp-1* and *lst-1* & *sygl-1* requires

the activity of the meiotic entry pathways (Kershner *et al.* 2014). Third, *lst-1* and *sygl-1*

nascent transcripts are restricted to the distal most germ cells, contacting the DTC, and

this expression requires *glp-1* activity (**Fig 1**; (Kershner *et al.* 2014; Lee *et al.* 2016)).

Fourth, LAG-1 binding sites in a *sygl-1* promoter reporter are required for distal germline

specific expression (Kershner *et al.* 2014). Finally, *lst-1* was previously identified as a

LIN-12 transcriptional target in vulval development (Yoo *et al.* 2004). The Pumilio family

RNA binding proteins FBF-1 and FBF-2 also promote the stem cell fate, functioning in

mRNA degradation and translational repression of meiotic entry pathway gene products

(Crittenden *et al.* 2002; Suh *et al.* 2009; Voronina *et al.* 2012). LST-1 and SYGL-1 act,

at least in part, through binding FBF-1 and FBF-2 and post-transcriptionally repressing

GLD-1 accumulation (Shin *et al.* 2017; Haupt *et al.* 2019).

lst-1 and *sygl-1* were identified in a candidate RNAi screen of 15 genes that meet the

following criteria – (i) the gene contained a cluster of at least four LAG-1 binding sites

and (ii) its mRNA was a target of FBF-1 regulation based on FBF-1 immunoprecipitation

followed by microarray analysis (Ketshner *et al.* 2010; Kershner *et al.* 2014). The

presence of LAG-1 binding sites in genes regulated by Notch signaling transcription is

logically predicted. However, as FBF-1 promotes the stem cell fate and functions in

mRNA degradation and translational repression, it is counterintuitive that GLP-1

signaling transcriptional targets would be FBF-1 posttranscriptional repression targets.

Therefore, it is possible that there are additional GLP-1 signaling transcriptional targets

which could be identified in an unbiased approach. Three other GLP-1 signaling

germline transcriptional targets have been reported, *fbf-2*, *utx-1* and *lip-1* (Lamont *et al.*

2004; Lee *et al.* 2006; Seelk *et al.* 2016), but with significantly less experimental support

than *lst-1* and *sygl-1*.

We have taken a genome-wide approach to identify transcriptional targets of GLP-1

signaling, through intersection of genes identified as directly bound by LAG-1, from

ChIP-seq experiments, with genes identified as requiring GLP-1 signaling for RNA

accumulation, from RNA-seq transcriptomics analysis. An important part of our

approach was identifying genes whose RNA level, or protein level, was dependent on

glp-1 signaling. However, in *glp-1* null mutants, germ cells enter meiosis in the L2 stage

complicating comparative RNA and protein accumulation studies because of very

different germ cell number and type. Therefore, we have taken advantage of the

epistasis of *gld-2 gld-1* double null over the *glp-1* null mutant, comparing meiotic entry

defective tumorous germlines, with and without *glp-1* activity, similar to other studies

(Hansen *et al.* 2004; Lamont *et al.* 2004; Kershner *et al.* 2014; Lee *et al.* 2016; Seelk *et*

al. 2016; Shin *et al.* 2017). We identified *lst-1* and *sygl-1* as the only genes that had

both germline LAG-1 binding and whose expression was dependent on *glp-1* and

germline *lag-1* activity. We used a time course following auxin-induced degradation of

LAG-1 to distinguish between primary versus secondary changes in RNA level in GLP-1

signaling. *lst-1* and *sygl-1* were the first genes (2hrs treatment) whose mRNA level were

dependent on LAG-1 activity, representing a primary effect. Five additional genes were

then identified at a later time point (4hrs treatment) whose RNA accumulation was

dependent of LAG-1 activity. Consistent with these changes being a secondary effect,

the five genes were not bound by LAG-1 and for three of the genes, RNA accumulation

was dependent on *lst-1* and *sygl-1* activity. We additionally showed that *glp-1*

dependent peak FBF-2 accumulation (Lamont *et al.* 2004) is fully explained by a

requirement for downstream *lst-1* and *sygl-1* activity. Finally, we expand our

understanding of *lag-1* function and expression. We showed that *lag-1* is germline

autonomously required for the stem cell fate and found that LAG-1 is spatially restricted,

with peak accumulation in the stem cell region of the progenitor zone. Peak

accumulation is regulated posttranscriptionally, with about 50% of *glp-1* dependent

LAG-1 accumulation contributed by *lst-1* and *sygl-1* activity. Together, our results are

consistent with the possibility that *lst-1* and *sygl-1* are the only mRNA transcriptional

targets and that GLP-1 signaling is mediated largely or completely by *lst-1* and *sygl-1*.

Results

LAG-1 accumulation is spatially restricted and germline autonomously required

for the stem cell fate

As a first step in examining GLP-1 Notch – LAG-1 CSL transcriptional control of the

germline stem cell fate, we determined the LAG-1 protein accumulation pattern, which

has not previously been reported. The endogenous *lag-1* locus was tagged with 3xHA

at the C-terminus of the LAG-1, using CRISPR/Cas9 engineering, to generate *lag-*

1(oz530[lag-1::3xHA]), hereafter called *lag-1::HA* (**Fig 2A; Materials and Methods**).

The *lag-1:: HA* strain appears phenotypically wild type; we did not observe phenotypes

associated with *lag-1* loss of function (Lambie and Kimble 1991; Qiao *et al.* 1995),

including loss of *glp-1* - embryonic lethality or a smaller progenitor zone due to stem

cells undergoing spatially premature meiotic entry, loss of *lin-12* - egg-laying/vulva

defects, or loss of both *lin-12* and *glp-1* - Lag larval arrest (**S1 Fig**). We examined

germline LAG-1::HA accumulation by anti-HA antibody staining in dissected

hermaphrodite gonad preparations (**Fig 2B-C; Materials and Methods**). In young

adults (1 day past mid-L4 larval stage) LAG-1::HA was found in germ cell nuclei in the

distal most ~10 cell diameters of the progenitor zone (PZ), during late pachytene,

diplotene and diakinesis of oogenesis. The late oogenic accumulation is consistent with

maternal loading of LAG-1 for early embryonic GLP-1 signaling in specification of

certain blastomere cell fates (Priess 2005). In mid/late L4, LAG-1::HA was also found in

germ cell nuclei in the distal most ~10 cell diameters of the PZ, but not observed in late

pachytene cells in the proximal gonad, which correspond to germ cells undergoing

spermatogenesis (Ellis and Schedl 2007). Distal germline LAG-1::HA staining was

variable in 1-day adults, and the staining intensity was weaker than in the L4 stage. We

also observed strong LAG-1:::HA staining in nuclei of all somatic gonad cells, the DTC,

and all sheath and spermathecal cells, as well as in polyploidy intestinal cells, in both

the L4 and adult stage dissection preparations.

LAG-1 has been assumed to function in the germline to promote the stem cell fate,

however the observation of strong LAG-1 accumulation in the DTC raises the possibility

that it may have a non-autonomous function in promoting the stem cell fate. We used

the auxin-inducible degradation (AID) system that destabilizes degron tagged proteins

(Zhang *et al.* 2015), to test if LAG-1 is require in the germline to promote the stem cell

fate. The degron tag was placed at the C-terminus of endogenous LAG-1 locus,

followed by 3xHA, using CRISPR/Cas9 engineering, to generate *lag-1(oz536oz537),*

hereafter called *lag-1::degron::HA* (**Fig 2A; Materials and Methods**). Germline specific

degradation was achieved by germline specific expression of the TIR1 F-box substrate

specificity protein, using the *gld-1* promoter (*ieSi64[gld-1p::TIR1::mRuby]*; (Zhang *et al.*

2015); **Materials and Methods**). In the absence of auxin treatment, the *lag-*

1::degron::HA strain is phenotypically wild type, with LAG-1::degron::HA accumulation

indistinguishable from LAG-1::HA (**Fig 2B, D; S1 Fig**). Auxin mediated degradation was

initiated at mid-L4. After 4 hrs of auxin treatment, LAG-1::degron::HA distal germ cell

nuclear staining was no longer detected and this was also true at 24hrs of auxin

treatment (**Fig 2D**). In contrast, strong LAG-1::degron::HA staining was detected in the

DTC, sheath and spermathecal cells and the intestine at both 4 and 24 hrs of auxin

treatment, consistent with germline specific degradation of LAG-1 (**Fig 2D**).

To assess inappropriate entry of stem cells into meiotic prophase following loss of LAG-

1::degron::HA, we stained dissected gonads for progenitor zone marker CYE-1 cyclin E

(Fox *et al.* 2011) and meiotic chromosome axis protein HIM-3 (Zetka *et al.* 1999;

Hansen *et al.* 2004b)(**Materials and Methods**). After 4 hrs of auxin treatment, when

LAG-1::degron::HA is no longer detected, the position of the progenitor zone – leptotene

boundary [in cell diameters (CD) from the distal tip] was not significantly different from

wild type and from animals not treated with auxin (**Fig 2E; S1 Fig**). Following 24hrs of

auxin treatment all progenitor zone cells were CYE-1 negative and HIM-3 positive,

indicating that all the stem cells had entered meiotic prophase. The kinetics of stem cell

meiotic entry following germline loss of LAG-1::degron::HA (also see below) are

consistent with the kinetics observed following loss of GLP-1 signaling through shift of

glp-1 temperature sensitive mutants to the restrictive temperature; the progenitor zone

is maintained at 4 hrs and absent by 10 hrs (Fox and Schedl 2015). Together, the

above results indicate that LAG-1 is germline autonomously required and expressed in

the appropriate distal germ cells to promote the stem cell fate.

LAG-1 accumulation is positively regulated by GLP-1 signaling and negatively

regulated by the GLD-1 and GLD-2 meiotic entry pathways

We next investigated how LAG-1 accumulation is regulated in the distal germline. The

lag-1 gene contains multiple consensus LAG-1 binding sites (Christensen *et al.* 1996),

and along with the distal germline restricted accumulation pattern, suggests the

possibility that LAG-1 accumulation occurs through a positive transcriptional loop via

GLP-1 signaling. If the GLP-1(ICD) - LAG-1 complex is regulating *lag-1* transcription, we

predicted that the *lag-1* mRNA would be spatially restricted to the first ~5 - 10 cell

diameters (CD) from the distal tip in wild type, where GLP-1 dependent transcription

and cytoplasmic mRNA of known targets *sygl-1* and *lst-1* are observed, and that *lag-1*

mRNA accumulation would depend on GLP-1 activity (Lee *et al.* 2016, 2019). We

performed s̲ingle m̲olecule f̲luorescent *i̲n s̲itu* h̲ybridization (smFISH) on dissected

gonads with *lag-1* mRNA complementary probes, analyzing expression quantitatively by

counting foci along the distal-proximal axis of the germline (Spike *et al.* 2014, **Materials**

and Methods). In wild type, we found that *lag-1* mRNA was uniformly distributed

throughout the distal 25 CD; abundant *lag-1* mRNA was found from 10 – 25 CD from

the distal tip, a region where *sygl-1* and *lst-1* mRNAs are not observed (**S2 Fig**). To

examine GLP-1 dependence, we used the *gld-2 gld-1* null double mutant meiotic entry

defective background (**Introduction**) and compared foci distribution relative to wild type

in *gld-2 gld-1* with either *glp-1(+)* or the *glp-1* null allele *q175*. As an internal control, wild

type gonads were co-dissected and smFISH performed together with *gld-2 gld-1* or *gld-*

2 gld-1; glp-1(q175). The *gld-2 gld-1* mutant background had no significant effect on the

uniform distribution of *lag-1* mRNA foci relative to wild type. Furthermore, loss of *glp-1*

activity had no significant effect on the uniform distribution of *lag-1* mRNA foci in *gld-2*

gld-1 (**S2 Fig**). The uniform distal *lag-1* mRNA accumulation, with or without *glp-1*

activity, is not consistent with the model that GLP-1 signaling promotes germline *lag-1*

transcription.

The above results indicate that distal restricted LAG-1 accumulation occurs through a

post-transcriptional mechanism. To investigate this mechanism, we first quantified the

distal – proximal accumulation of LAG-1::HA from anti-HA antibody staining in an

otherwise wild type background, and subtracted background signal from staining of *lag-*

1(+) germlines, lacking 3xHA (**Materials & Methods**). Because germline staining for

LAG-1::HA in 24hr adults was variable, we employed mid/late L4 hermaphrodites where

staining is more consistent from gonad to gonad. High LAG-1::HA was observed from 1

– 5 CD from the distal tip (peak) followed by a significant fall to a basal level from ~17 -

25 CD (**Fig 3A-B**). To allow comparisons, the mean peak intensity at 4 CD was set to

100. From peak to proximal base at 25 CD, there was an ~7-fold drop in LAG-1::HA

levels, with accumulation at base significantly above background, ~15% of peak.

We then investigated the possibility that GLP-1 signaling indirectly regulates LAG-1

peak levels, possibly through targets *sygl-1* and *lst-1*. The pattern and level of LAG-

1::HA was first assessed in the *gld-2 gld-1* background, and then in *gld-2 gld-1; glp-1*

null triple mutants and the *gld-2 gld-1 sygl-1 lst-1* quadruple null mutants, each co-

dissected and stained with wild type, with and without *lag-1::HA* to allow background

subtraction and normalization (**Fig 3A, C-D; Materials and Methods**). In *gld-2 gld-1*,

LAG-1::HA displayed a similar overall expression pattern as in wild type, but with

elevated accumulation throughout the progenitor zone, with peak levels increased by

~25% (also see below). In the absence of *glp-1*, LAG-1::HA level was reduced

throughout the progenitor zone, with a gradual rise from distal to proximal. Compared to

the peak level at 4 CD, LAG-1::HA is 39% of *glp-1(+)* level, indicating that ~39% of

peak LAG-1 accumulation is independent of GLP-1 signaling, with a similar level

continuing into the proximal progenitor zone. Thus, ~61% of peak LAG-1 accumulation

depends on GLP-1 signaling. In the absence of *sygl-1* and *lst-1*, we observed a peak of

LAG-1 accumulation that was intermediate between the presence and absence of *glp-1*,

~70% of the LAG-1::HA peak in *gld-2 gld-1*. The difference in accumulation indicates

that *lst-1* and *sygl-1* activity account for ~50% of *glp-1* dependent LAG-1 accumulation

(**Fig 3C-D**). These results don't provide an explanation for the remaining 50% of *glp-1*

dependent control of LAG-1 accumulation, which is not reliant on control of *lag-1* mRNA

level (**S2 Fig**). Figure 3E summarizes the GLP-1 signaling dependent and independent

control of LAG-1 peak accumulation, indicating that about one third of LAG-1

accumulation is promoted by GLP-1 transcriptional targets *lst-1* and *sygl-1*.

GLD-1 and GLD-2 repress LAG-1 accumulation in the proximal part of the progenitor

zone (**Fig 3B; S3 Fig**). We found that in the *gld-2 gld-1* double null mutant, LAG-1 level

was elevated ~4.5-fold, as assessed at 25 CD from the distal tip. GLD-1 alone accounts

for more that half of this repression, as LAG-1 level is elevated almost 3-fold in the *gld-1*

single null mutant, which is consistent with *lag-1* mRNA being identified as a GLD-1

target in RNA pull-down experiments (Jungkamp *et al.* 2011; Wright *et al.* 2011). As loss

of *gld-2* function alone does not affect LAG-1 level, the remaining repression is

apparently through the combined action of GLD-1 and GLD-2. The above results

indicate that the fall in LAG-1 level in the proximal part of the progenitor zone is largely

through translational repression by GLD-1 and GLD-2, as well as the absence of *lst-1*

and *sygl-1* promoting LAG-1 accumulation, which are spatially restricted to the distal

most ~5 – 10 CD. Peak LAG-1 accumulation is also repressed by the combined

activities of GLD-1 and GLD-2 (**Fig 3; S3 Fig**). This modest repression is presumably

because of lower *gld-1* and *gld-2* activity in the distal most 10-cell diameters (Hubbard &

Schedl, 2019).

Genome-wide identification of GLP-1 Notch – LAG-1 CSL transcriptional targets:

ChIP-seq

Multiple lines of evidence indicate that *lst-1* and *sygl-1* are transcriptional targets of

GLP-1 signaling to promote the germline stem cell fate (**Introduction**; (Kershner *et al.*

2014; Lee *et al.* 2016, 2019)). *lst-1* and *sygl-1* were identified in an RNAi candidate

screen of 15 genes that meet the following criteria – the gene contained a cluster of at

least four LAG-1 binding sites and its mRNA was a target of the Pumilio family RNA

binding protein FBF-1 (Kershner *et al.* 2014). FBF-1 promotes the stem cell fate through

acting in translational repression/mRNA destabilization; thus it is counterintuitive that

GLP-1 signaling transcriptional targets would be FBF-1 posttranscriptional targets,

which presumably would lead to downregulation of their expression. Therefore, it is

possible that there are additional GLP-1 signaling transcriptional targets that remain to

be identified. We have taken a genome-wide approach to identify transcriptional targets

of GLP-1 signaling, through intersection of genes identified as directly bound by LAG-1,

from ChIP-seq experiments, with genes that are identified as requiring GLP-1 signaling

for RNA accumulation, from RNA-seq transcriptomics analysis.

LAG-1 CSL is the DNA binding protein that mediates transcription of GLP-1 and LIN-12

signaling targets, and was used in the ChIP-seq experiments. We first performed

conventional ChIP-seq on whole mid-L4 worms, using the endogenous *lag-1* gene

tagged with GFP and 3xFLAG, *lag-1(ar611[lag-1::GFP::3xFLAG])* (**S4 Fig**; gift from Iva

Greenwald). We separately performed anti-FLAG or anti-GFP Chromatin

immunoprecipitation (ChIP) experiments, both antibodies can efficiently pulled-down

tagged LAG-1 and gave a significant fold enrichment of *lst-1* and *sygl-1* DNA fragments

compared to control genes in ChIP-qPCR (**S4C Fig, Materials and Methods**).

Following high throughput sequencing, peaks were identified with greater than a two-

fold enrichment compared to input control, and a false discovery rate (FDR <0.05). The

Homer suite (Heinz *et al.* 2010) was used to annotate the peaks to their nearest

transcription start site (TSS). Seventy-six genes were identified as binding to LAG-1

from the intersection of the FLAG-IP (one biological replicate) and GFP-IP (two

biological replicates) ChIP-seq (**S4 Fig; S3 Table**). Known somatic LIN-12 signaling

targets in vulval development, *lst-1* and *mir-61/250* (Yoo et al. 2004; Yoo and

Greenwald 2005), and germline target *sygl-1* (Kershner *et al.* 2014) were identified with

each having a single major ChIP-seq peak that covered multiple canonical LAG-1/CSL

binding sites. The *lag-1* gene contained multiple peaks, particularly in the large first

intron that contains multiple canonical LAG-1 binding sites. The *lag-1* first intron is also

bound by a large number of transcription factors (~50% of those analyzed to date,

http://www.modencode.org/), but this is less than the definition of a Highly Occupied

Target (HOT), where >65% of transcription factors bind to a region (Gerstein *et al.* 2010;

Van Nostrand and Kim 2013). Thus, the *lag-1* gene may function in a LIN-12 (and

possibly GLP-1) signaling dependent positive autoregulatory feedback loop with LAG-1

in the soma, consistent with *lag-1* reporter gene analysis in the anchor cell – ventral

uterine cell decision (K. Luo and I. Greenwald, personal communication; Choi *et al.*

2013), although this does not appear to be the case in the germline (see above). We did

not identify peaks for other reported LIN-12 or GLP-1 signaling somatic targets, such as

ref-1 in the embryo or *lip-1* in vulval development ((Berset et al. 2001; Neves and Priess

2005); **S3 Table**). This maybe a result of our analysis being from only a single stage

(mid-L4) or because only a small number of cells are expressing the target gene under

LIN-12 or GLP-1 control, resulting in only a small amount of total LAG-1 bound DNA,

which is below the limit of detection in the ChIP-seq assay. We performed *de novo*

discovery of over-represented DNA sequence motifs among the 75 genes and found the

highest hit to be an 9-mer that contains the canonical CBF/LAG-1 binding site (*p-value:*

e-42)(**S4H Fig**). Thus, we believe that the bulk of the genes identified are bound by LAG-

1 *in vivo*, although this data does not address the issue of whether the genes are

regulated by GLP-1 or LIN-12 signaling.

To identify genes whose transcription promotes the stem cell fate through direct LAG-1

binding, we performed germline specific LAG-1 ChIP-seq, where the approach was

guided by our expression and germline autonomy analysis above. LAG-1 is modestly

expressed in germline stem cells compared to much higher expression in late stage

pachytene and diplotene oogenic germ cells and somatic cells (**Fig 2, 3**). This

necessitated performing germline LAG-1 ChIP-seq on L4 stage hermaphrodites that

lack oogenic germ cells in pachytene and diplotene. We generated a transgenic strain

with the following components: (1) A fosmid transgene where the BioTag, a 23-amino

acid peptide that is recognized and biotinylated by the *E. coli* enzyme BirA biotin ligase

(Cronan 1990), was placed at the C-terminus of LAG-1 (*ozls43[lag-1::3xFLAG::BioTag*,

hereafter called *lag-1::BioTag*) (**S5A Fig; Materials and Methods**); (2) A transgene

with germline specific expression of *E. coli BirA* biotin ligase from the *pie-1* promoter; (3)

The *lag-1* deletion allele, *tm3052*, which demonstrated that *lag-1::BioTag* in *ozls43*

produced functional, rescuing, LAG-1. Germline specific expression of BirA should

result in germline restricted biotinylation of LAG-1::BioTag, which can be pulled down by

streptavidin beads (**Fig 4A**). However, because of high levels of endogenous *C.*

elegans biotinylated proteins (Ooi *et al.* 2009; Watts *et al.* 2018) we were not able to

directly pull-down the low levels of biotinylated LAG-1::BioTag. To overcome this issue,

we performed sequential ChIP, first with anti-FLAG and then with streptavidin beads,

followed by library construction directly on the beads due to ultra-low quantities of DNA

following sequential ChIP (**Fig 4B**). Germline specific LAG-1 ChIP-seq data was

analyzed as described for whole worm ChIP-seq.

One hundred and thirty seven genes with germline specific LAG-1 peaks were identified

in common between three biological replicates, including *lst-1*, *sygl-1* and *mir-61/250*

(**Fig 4C-D**). The *lag-1* gene was also found to contain multiple peaks. However, other

linked genes on the fosmid (e.g., *zen-4*) and the selectable marker used (*unc-119*) also

contained multiple peaks, which were absent in the whole worm ChIP-seq. Thus, it is

likely that there are multiple integrated copies of *ozIs43*, leading to an elevation of

background sequences being pulled down and detected in the germline ChIP-seq

experiments, therefore these genes were not considered to be specifically bound by

germline LAG-1 . *De novo* motif identification recovered the same 9-mer that contains

the canonical CBF/LAG-1 binding site (*p-value: e-^{34}*) as found in the whole worm

experiment, consistent with LAG-1 binding many of these genes *in vivo*, in germ cells.

Thirty-six genes were identified in common between germline specific and whole worm

that show LAG-1 ChIP-seq peaks, supporting that LAG-1 binds to these genes *in vivo*

(**S5E Fig**). Together, these results provide biochemical support that *lst-1* and *sygl-1* are

germline GLP-1 transcriptional targets.

Genome-wide identification of GLP-1 Notch – LAG-1 CSL transcriptional targets:

RNA-seq

To identify genes that require GLP-1 signaling for expression we performed RNA-seq

comparing two strains, one with GLP-1 signaling ON, using *glp-1* gain of function (gf)

allele *ar202* that produces a large number of proliferating germ cells undergoing GLP-1

signaling at the restrictive temperature (Pepper *et al.* 2003), and the other with GLP-1

signaling OFF, using the null allele *q175*. As described above, we employed the meiotic

entry defective *gld-2 gld-1* double null mutant to allow examination of the effect of *glp-1*

null in proliferating germ cells; *glp-1* gf was also placed in this background so that

genotype was identical, except for *glp-1* activity status (**Fig 5A; Materials and**

Methods). Further, we used dissected gonad preparations to significantly enrich for

expression changes that occur in the germline. *In situ* hybridization and qRT-PCR was

used to confirm that in GLP-1 ON, transcriptional target *sygl-1* and *lst-1* were expressed

throughout the germline at significantly elevated levels, compared to GLP-1 OFF where

expression was similar to background (**Fig 5B-C**, **S6A Fig**). RNA-seq was performed on

5 biological replicates, following Ribo-zero treatment, from GLP-1 ON and GLP-1 OFF

dissected gonad RNA preparations. Heatmap and principle component analysis (PCA)

demonstrated significant differences between the GLP-1 ON and GLP-1 OFF RNA-seq

results (**S6B-C Fig**). We identified 94 genes whose RNA accumulation was dependent

of GLP-1 signaling, with greater than 2-fold elevation of reads in GLP-1 ON versus

GLP-1 OFF, FDR<0.05, and requiring greater than two counts per million reads (CPM)

(**Materials and Methods; S5 Table**).

Intersection of the 137 genes identified as bound by LAG-1 with the 94 genes whose

RNA accumulation was dependent on GLP-1 signaling yielded only two genes, *lst-1* and

sygl-1 (**Fig 4, 5**). Three additional genes have been reported as germline GLP-1

signaling targets, *fbf-2*, *utx-1*, and *lip-1* (Lamont *et al.* 2004; Lee *et al.* 2006; Seelk *et al.*

2016). We did not observe LAG-1 germline ChIP-seq peaks (**S5B Fig**) for *fbf-2*, *utx-1* or

lip-1. For GLP-1 dependent RNA accumulation in GLP-1 ON versus OFF, *fbf-2* mRNA

level was unchanged, *utx-1* mRNA level was below the 2 CPM cutoff, while *lip-1* mRNA

increased 2.3 fold in GLP-1 ON. Consistent with unchanged mRNA level, an FBF-2::fbf-

2 3'UTR reporter driven by the heterologous *pie-1* promoter gives a qualitatively similar

FBF-2 progenitor zone protein accumulation as observed in wild type ((Lamont *et al.*

2004; Wang *et al.* 2016); see below), supporting GLP-1 signaling independent control of

fbf-2 mRNA accumulation. Prior work indicated that *lip-1* mRNA accumulation in the

progenitor zone was primarily controlled by FBF-1 mediated degradation (Lee *et al.*

2006). Additionally, we note that dissected gonad preparations, while largely containing

proliferating germ cells, also contain the DTC, all sheath cells, and variable amounts of

spermathecal and uterine cells, depending on the efficacy of the dissection and carcass

removal. The extent to which GLP-1 signaling is occurring in the somatic gonad in the

young adult is unknown. The RNA-seq method we employed does not recover RNAs

less than ~100nt. Thus, it remains possible that small RNAs, including *mir-61/250*, are

transcriptional targets of GLP-1 signaling (**Fig 4C-D**). Together, the above experiments

provide robust *in vivo* biochemical support for *lst-1* and *sygl-1* being mRNA

transcriptional targets of germline GLP-1 signaling, and potentially being the only GLP-1

signaling transcriptional targets that promote the stem cell fate.

FBF-2 accumulation is controlled by GLP-1 signaling transcriptional targets *lst-1*

and *sygl-1*

Previous work showed that FBF-2 accumulation is enriched in the progenitor zone, with

peak FBF-2 accumulation dependent on GLP-1 signaling (Lamont *et al.* 2004). We used

CRISPR/Cas9 tagged *fbf-2(q932[3xV5::fbf-2])* (Shin et al. 2017), hereafter called *fbf-*

2::V5, to quantitatively examine FBF-2 accumulation, following anti-V5 antibody staining

in dissected gonads from young adult hermaphrodites (**Materials and Methods**). We

found that FBF-2::V5 displayed a peak of accumulation at 8 – 13 CD from the distal tip,

followed by a somewhat gradual fall to low proximal levels by ~35 CD (**Fig 6A-B**),

similar to previously reported (Lamont *et al.* 2004), with peak accumulation ~4 fold

higher than base. In *gld-2 gld-1* double null mutant germlines, the FBF-2::V5 peak was

~80% of wild type, with a the fall more rapid than in wild type and a flat base from ~18

CD through 35 CD, with the peak also ~4 fold higher than base (**Fig 6A, C**). In the *gld-2*

gld-1; glp-1 triple null mutant germlines, FBF-2::xV5 levels are low throughout the

progenitor zone (**Fig 6A, C-D**), with the level ~4 fold lower than in the *gld-2 gld-1* double

mutant. Thus, peak FBF-2 accumulation requires GLP-1 signaling, with ~15% of FBF-2

accumulation being GLP-1 signaling independent, as previously reported (Lamont *et al.*

2004). Given that FBF-2 accumulation appears to be controlled post-transcriptionally

(Wang *et al.* 2016), and the *lst-1* and *sygl-1* promote peak LAG-1 accumulation in the

progenitor zone (**Fig 3**), we next examined whether LST-1 and SYGL-1 promoted peak

FBF-2 accumulation. Analysis of *gld-2 gld-1 lst-1 sygl-1* quadruple null mutant germlines

showed low FBF-2 accumulation throughout the progenitor zone (**Fig 6**), consistent with

LST-1 and SYGL-1 being required for peak FBF-2 accumulation. FBF-2 levels at the

peak position in wild type are almost 2 fold lower in the absence of *lst-1* and *sygl-1*,

compared to the absence of *glp-1*; the reason for this difference is unknown. In the

proximal progenitor zone, GLD-1 appears to promote FBF-2 accumulation as the level

is lower in the *gld-1* null mutant (**S7 Fig**). Since GLD-1 acts in translational repression,

the effect on FBF-2 accumulation is presumably indirect. In summary, GLP-1 dependent

peak FBF-2 accumulation in the progenitor zone appears to be completely explained

through post-transcriptional regulation by *lst-1* and *sygl-1* activity (**Fig 6E**).

Germline LAG-1 functions in transcriptional activation of *lst-1* and *sygl-1*

Only two of 137 germline LAG-1 ChIP-seq peaks are associated with GLP-1 dependent

activation of target transcription. The remaining peaks may reflect LAG-1 acting

independent of GLP-1 signaling, either as a transcriptional activator or repressor. In

Drosophila, it is known that in the absence of Notch ICD, *Su(H)* CSL functions in

actively repressing Notch transcriptional target genes (Bray 2006). To test if LAG-1 has

a GLP-1 signaling independent transcriptional function in the germline, we performed

LAG-1 AID followed by gonad dissection and RNA-seq (**Fig 7**). We used a strain

containing *lag-1::degron*, *glp-1(ar202)* gf that at the restrictive temperature will have the

bulk of germ cells undergoing GLP-1 signaling, and the meiotic entry defective double

mutant *gld-2 gld-1*, to allow analysis of proliferating germ cells, with or without auxin

treatment. Germline specific degradation of LAG-1 was achieved with *gld-*

1p::tir1::mRudy as described above (**Fig 2D, Materials and Methods**).

We performed RNA-seq following 48hrs of auxin treatment from the L1 stage at 25°C, a

time where LAG-1 is undetectable, and the same time point where GLP-1 ON and OFF

RNA-seq was performed (**Fig 5**). Heatmap and principal component analysis

demonstrated that changes in RNAs from LAG-1 ON (minus auxin) and LAG-1 OFF

(plus auxin) were significantly different, while biological replicates were similar (**S8A-B;**

S6 Table). Ninety-four genes were identified whose expression was dependent on LAG-

1 (activated genes), where ~70% (64) of these were also GLP-1 dependent genes (**S8C**

Fig; S5-6 Tables). Only two of the LAG-1 activated genes overlapped with LAG-1

germline specific ChIP-seq peak containing genes, *lst-1* and *sygl-1* (**S8E Fig**). Forty

eight genes were identified that were LAG-1 repressed, where ~80% of these were also

GLP-1 repressed genes (**S8D Fig**). There was no overlap between LAG-1 repressed

genes and germline LAG-1 ChIP-seq peak genes (**S8F Fig**), which is not consistent

with LAG-1 functioning in transcriptional repression in the *C. elegans* germline. We note

that the LAG-1 activated/repressed genes list and the GLP-1 signaling

activated/repressed gene list derive from strains that functionally differ in at least two

ways that may have resulted in incomplete overlap of gene lists. First, LAG-1 AID may

not completely eliminate LAG-1 protein, while the *glp-1(q175)* allele is null. Second,

while both use dissected gonads for RNA-seq, LAG-1 AID degrades LAG-1 specifically

in the germline, while *glp-1(q175)* lacks GLP-1 signaling in both the germline and the

soma. The above results indicate that while LAG-1 has an essential function in

transcriptional activation of *lst-1* and *sygl-1* for germline GLP-1 signaling, it does not

have an essential GLP-1 signaling independent function in either transcriptional

activation or repression. The disconnect we observed between the large number of

genes containing a germline LAG-1 ChIP-seq peak(s) but whose transcription appears

not to be regulated by GLP-1 signaling or LAG-1 is similarly observed in yeast and

mammalian cells where transcription factors bind many genes, but regulate significantly

fewer (Lenstra and Holstege 2012; Cusanovich *et al.* 2014). The biological significance

of this disconnect is currently unclear.

Primary versus secondary GLP-1 signaling mRNA targets

The LAG-1 AID system described above provides a route to distinguish between genes

whose expression is a primary or a secondary effect of GLP-1 signaling/LAG-1 activity.

By performing a time course of LAG-1 degradation, using RNA-seq as the readout,

genes whose RNA levels are transcriptionally controlled by GLP-1 signaling should

change expression earlier, while genes whose RNA level is indirectly controlled/a

secondary effect of GLP-1 signaling, should change expression later.

We performed a LAG-1 AID time course, harvesting RNA following 0.5, 1, 2, 4 and

48hrs of auxin treatment (**Fig 7**). *sygl-1* and *lst-1* mRNA levels, as assessed by qRT-

PCR, dropped significantly by 1hr of auxin treatment and by 2hrs were not significantly

different from 48hrs of auxin treatment or GLP-1 OFF. Staining for LAG-

1::degron::3xHA indicated that at 2 and 4hrs of auxin treatment, LAG-1 was at a

low/undetectable level, while staining for WAPL-1 show no change in the size of the

progenitor zone (**S9 Fig**), indicating that the switch of all the stem cell to meiotic

development, following loss of LAG-1, was only just beginning. RNA-seq was therefore

performed at 2 and 4hrs, time points, where there was minimal pleiotropy from loss of

germline stem cells. Multidimensional scaling (MDS) plots revealed that the 0, 2, and

4hrs auxin treatment clustered together, but separately from 48hrs auxin treatment,

indicating that there are few changes in expression at early times following degradation

of LAG-1 compared to strong loss of LAG-1 (**S8G Fig**). At 2hr of auxin treatment,

among the GLP-1 dependent genes, only *lst-1* and *sygl-1* were identified as LAG-1

mRNAs targets (**Fig 7B-D**). By 4hrs of auxin treatment, 5 additional genes were

identified whose RNAs displayed LAG-1, as well as GLP-1, dependent accumulation

(**Fig 7D-E**). The new 5 genes from the 4hr time point could represent additional GLP-1 -

LAG-1 transcriptional targets, which have longer RNA half-lives than *lst-1* and *sygl-1*, or

they may represent changes in RNA level that are secondary effects of GLP-1 signaling.

We did not detect germline LAG-1 ChIP-seq peaks or canonical LAG-1/CSL binding

sites in these 5 genes, indicating that they are not GLP-1 transcriptional targets. From

these results, and our previous finding that *lst-1* and *sygl-1* promote LAG-1 and FBF-2

protein accumulation, we reasoned that the accumulation of RNA for these 5 genes

maybe dependent on *lst-1* and *sygl-1* activity.

We first examined the kinetics of loss of LST-1 and SYGL-1 following LAG-1 AID. Peak

LST-1 falls ~5 fold at 2hrs and ~10 fold at 4hrs, while SYGL-1 falls ~3 fold at 2hrs and

10-fold at 4hrs auxin treatment (**S10 Fig**). Given their rapid loss, the significant

reduction of LST-1 and SYGL-1 could be responsible for the reduction in RNA level of

the 5 genes at the 4hrs auxin treatment time point. To test this, we measure RNA level

by qRT-PCR for the 5 genes (*epg-5, tbx-2*, C17H12.36, *ucr-2.1*, F40D4.13) from

dissected gonads with GLP-1 ON, GLP-1 ON but lacking *lst-1* and *sygl-1* activity (LST-1

SYGL-1 OFF) and GLP-1 OFF (see **Fig 7F** and legend for full genotype). For *epg-5,*

tbx-2 and C17H12.36, higher RNA accumulation in GLP-1 ON was dependent on *lst-1*

and *sygl-1* activity. Thus, *epg-5*, *tbx-2* and C17H12.36 are indirect or secondary targets

of GLP-1 signaling, with their RNA accumulation dependent on the downstream *lst-1*

and *sygl-1* gene activity. It is likely that other genes whose RNA level are dependent on

both LAG-1 at 48hrs and GLP-1 signaling are secondary targets that also rely on *lst-1*

and *sygl-1* activity. The LAG-1 AID time course provides an estimate of the maximum

half-life of the *lst-1* and *sygl-1* mRNAs, 1hr, and the LST-1 and SYGL-1 proteins, 2hrs,

at 25°C.

Discussion

A central question for the *C. elegans* germline stem cell system is the identity of the

GLP-1 Notch dependent transcriptional targets that promote the stem cell fate. We

performed a genome-wide search for transcriptional targets through germline specific

ChIP-seq analysis with LAG-1, the Notch signaling CSL DNA binding protein homolog,

and intersected these results with transcriptomics analysis under conditions of the

presence or absence of GLP-1 signaling and the presence or absence of LAG-1. We

provide molecular support for two previously identified transcriptional targets, *lst-1* and

sygl-1. Our results are consistent with the possibility of no additional direct mRNA

transcriptional targets of germline GLP-1 signaling. We further report examples of GLP-

1 dependent control of RNA level and protein accumulation that are indirect, through

LST-1 and SYGL-1. We elaborate on these and other findings below.

611   GLP-1 - LAG-1 transcriptional control of the germline stem cell fate

We present three lines of genome-wide molecular support that *lst-1* and *sygl-1* are

direct GLP-1 signaling transcriptional targets. First, we identified robust germline LAG-1

ChIP-seq peaks in the promoters of *lst-1* and *sygl-1*, overlapping the position of

consensus LAG-1/CSL binding sites. Second, transcriptomics analysis demonstrated

GLP-1 signaling and germline LAG-1 dependent *lst-1* and *sygl-1* mRNA accumulation.

Third, in a LAG-1 AID time course, *lst-1* and *sygl-1* mRNAs were the first RNAs to fall,

as early as 0.5 to 1hr after auxin treatment, indicating that their mRNA loss was a

direct/primary effect of reduced LAG-1 activity. Combined with prior work that *lst-1* and

sygl-1 transcription in the distal progenitor zone is dependent on GLP-1 signaling and

that distal *sygl-1* reporter gene expression requires canonical LAG-1 binding sites

(Kershner *et al.* 2014; Lee *et al.* 2016), our results strengthen the conclusion that *lst-1*

and *sygl-1* are direct transcriptional targets.

This work supports the possibility that there may not be any additional direct mRNA

transcriptional targets of GLP-1 signaling. Our results are not consistent with three

reported genes, *fbf-2*, *lip-1* and *utx-1*, being GLP-1 transcriptional targets (Lamont *et al.*

2004; Lee *et al.* 2006; Seelk *et al.* 2016). ChIP-seq peaks were not observed for the

three genes. *fbf-2* mRNA level was unchanged with or without GLP-1 signaling or LAG-

1 protein. For *lip-1*, while GLP-1 signaling and LAG-1 promoted mRNA accumulation, in

the LAG-1 time course *lip-1* mRNA level was unchanged at 2 and 4hrs of auxin

treatment, but was reduced at 48hrs (**S5-7 Tables**). The delayed fall in *lip-1* mRNA level

suggests an indirect/secondary effect of GLP-1 signaling and LAG-1 activity. For *utx-1*,

its mRNA read count was below the 2 CPM cut-off for reliable RNA level assessment.

From transcriptomics analysis, while we identified genes whose RNA accumulation was

GLP-1 signaling dependent (94) or germline LAG-1 dependent (94), other than *lst-1* and

sygl-1, they lacked germline or whole worm LAG-1 ChIP-seq peaks. From the LAG-1

AID time course, five genes were identified where their RNA level was dependent of

LAG-1 at 4hrs of auxin treatment, raising the possibility that they were direct/primary

targets of LAG-1 activity. However, in addition to the five genes lacking LAG-1 ChIP-

seq peaks, RNA accumulation for three of the genes was dependent on *lst-1* and *sygl-1*.

Thus, our genome-wide studies did not identify any new candidate GLP-1 signaling

mRNA transcriptional target genes. However, the RNA-seq approach employed cannot

assess the level of small RNAs (less than ~100nt), and a number of small RNA genes

were identified as containing germline LAG-1 ChIP-seq peaks (e.g, *mir-61/250*, five 21

U RNA genes). Thus, it is possible that there are small RNA genes that are germline

GLP-1 signaling transcriptional targets. We note that because of the low amount of DNA

that was obtained for the germline LAG-1 ChIP-seq experiments, it is possible that we

missed some weak ChIP-seq peaks. We also note that all of the transcriptomics

analyses were performed using dissected gonads from the *gld-2 gld-1* meiotic entry

double mutant background, to allow germline RNA level analysis in the absence of *glp-1*

or *lag-1* activity. We cannot rule out the possibility that the absence of *gld-1* and *gld-2*

activity may mask GLP-1 signaling dependent changes in RNA levels. Nevertheless, the

observation that the premature meiotic entry phenotype of the *lst-1 sygl-1* null double

mutant is the same as *glp-1* null (Kershner *et al.* 2014) indicates that no additional

transcriptional targets genes are necessary to account for germline GLP-1 signaling.

We explored aspects of *lag-1* function and expression that had not previously been

reported. We found that *lag-1* is germline autonomously required for the stem cells fate,

as predicted from our current understanding of Notch signaling and notwithstanding

strong LAG-1 accumulation in all somatic gonad cells. In the soma, LAG-1 appears to

function in a positive autoregulatory feedback loop to promote accumulation in cells

undergoing the LIN-12 dependent AC/VU decision (K. Luo and I. Greenwald, personal

communication; (Choi *et al.* 2013)). Consistent with this possibility, we found multiple

LAG-1 ChIP-seq peaks in the *lag-1* gene in the whole worm ChIP-seq experiments. In

the germline, however, *lag-1* mRNA was found at a constant level in the progenitor

zone, in the presence or absence of *glp-1* activity. This finding is consistent with

germline *lag-1* transcription being independent of GLP-1 signaling, with a *lag-1* positive

transcriptional feedback loop not being active in the germline. We found that LAG-1

protein accumulation is spatially restricted, high in the distal most 5 CD (peak) from the

tip and then falls 7-fold to a base level ~17 CD from the tip. Importantly, the germ cells

with peak LAG-1 accumulation correspond to those where GLP-1 dependent nascent

lst-1 and *sygl-1* transcripts are observed (Lee *et al.* 2016, 2019). While GLP-1 signaling

is not controlling *lag-1* mRNA level, we nevertheless found that 61% of peak LAG-1

accumulation was GLP-1 signaling dependent, suggesting a posttranscriptional

mechanism (**Fig 3E**). This is consistent with the observation that many genes in the

distal germline are regulated posttranscriptionally through the 3'UTR (Merritt et al. 2008;

Merritt and Seydoux 2010). We found that ~50% of the GLP-1 dependent LAG-1

accumulation required *lst-1* and *sygl-1* activity. The basis for the remaining ~50% of

GLP-1 dependent LAG-1 accumulation is not known. Perhaps GLP-1(ICD) is stabilizing

LAG-1 protein. The fall in LAG-1 levels in the proximal progenitor zone can be

attributed, at least in part, to GLD-1 and GLD-2 activity. We speculate that the spatial

restriction of LAG-1 contributes to controlling the size of the stem cells pool, with peak

levels required for efficient *lst-1* and *sygl-1* transcription in the first 5 CD, and that lower

levels in the proximal progenitor zone decrease the probability of stochastic GLP-1

signaling triggering transcription of *lst-1* and *sygl-1* in cells as they progress toward

meiotic development.

Mammalian CBF1 and Drosophila Su(H), orthologs of LAG-1, can act as transcriptional

repressors in the absence of Notch signaling (Bray 2006, 2016). Following germline

specific loss of LAG-1, we found that *lst-1* and *sygl-1* RNA levels dropped significantly,

equivalent to the absence of GLP-1 signaling. Thus, LAG-1 is required for GLP-1

dependent expression and, correspondingly, does not appear to function in repression

of transcriptional targets *lst-1* and *sygl-1*. In the genome-wide studies, none of the

germline LAG-1 repressed genes (RNA level elevated following LAG-1 AID) contain

germline LAG-1 ChIP-seq peaks, indicating that transcriptional repression is not a

general property of LAG-1 in the germline. We note that the Notch signaling dynamics in

the *C. elegans* germline differs from well-known lateral signaling examples, which may

then affect mechanistic aspects of transcriptional control. GLP-1 signaling and *lst-1* and

sygl-1 activity are continuously required in the *C. elegans* germline to promote the stem

cell fate, from the L1 larval stage through at least mid-adulthood, under optimal growth

conditions (Hubbard and Schedl, 2019). Thus, with the exception of the early L1 stage,

germline stem cells are born undergoing GLP-1 signaling, and only lose GLP-1

signaling when cells are displaced away from the DTC niche (ON > OFF). In contrast,

cells undergoing lateral signaling initially lack Notch signaling, then undergo Notch

signaling, and then may continue or downregulate Notch signaling (OFF>ON>OFF).

These differences in Notch signaling dynamics are likely mirrored in the transcriptional

states of respective cells and may provide an explanation for differences observed

between the *C. elegans* germline and other cells and organisms in Notch mediated

transcriptional control.

GLP-1 signaling indirectly mediates control of RNA and protein levels through

LST-1 and SYGL-1

We have found a number of examples of GLP-1 dependent RNA and protein

accumulation that are through the activity of *lst-1* and *sygl-1*. Lamont et al. (2004)

reported that distal peak FBF-2 accumulation requires GLP-1 signaling. We found that

the *glp-1* dependent peak FBF-2 accumulation can be fully explained by the activity of

lst-1 and *sygl-1* (**Fig 6**). FBF-2 activity functions in repression of the GLD-1 and GLD-2

meiotic entry pathways. As described above, ~60% of peak LAG-1 accumulation is

GLP-1 dependent, about 50% of which can be attributed to *lst-1* and *sygl-1* activity.

From the LAG-1 AID time course, we identified 5 genes whose RNA accumulation

depended on LAG-1, as well as *glp-1* activity. For *epg-5*, *tbx-2* and C17H12.36 we

found that their GLP-1 and LAG-1 dependent expression can be explained by the

requirement for *lst-1* and *sygl-1* activity. The remaining two genes, *ucr-2.1* & F40D4.13,

show a trend to *lst-1* and *sygl-1* dependence, although it is not significant with the

number of replicates examined. Previous work reported that GLP-1 signaling inhibits

GLD-1 accumulation in the distal progenitor zone (Hansen *et al.* 2004b) and this

posttranscriptional repression required *lst-1* and *sygl-1* activity (Shin *et al.* 2017). Thus,

the bulk of germline gene expression changes ascribed to GLP-1 signaling, and *lag-1*

function, can be attributed to the activity of transcriptional targets *lst-1* and *sygl-1*. These

findings are consistent with the possibility that there are no additional mRNA targets of

GLP-1 signaling to promote the stem cell fate.

LST-1 and SYGL-1 have been reported to function in conjunction with FBF-1 and FBF-2

in direct translational repression/mRNA destabilization in the posttranscriptional

inhibition of GLD-1 accumulation (Shin *et al.* 2017; Haupt *et al.* 2019). FBF-1 has also

been shown to function in translational repression/mRNA destabilization in the

posttranscriptional inhibition of FBF-2 accumulation (Lamont *et al.* 2004). In contrast, we

find that *lst-1* and *sygl-1* function in posttranscriptional activation of FBF-2 and LAG-1

accumulation. Future work will be necessary to determine if *lst-1* and *sygl-1* act with or

separately from FBF-1 and FBF-2 in posttranscriptional activation of FBF-2 and LAG-1

accumulation. Similarly, *lst-1* and *sygl-1* promote *epg-5*, *tbx-2* and C17H12.36 RNA

accumulation, and the mechanism by which this occurs remains to be determined.

Materials and Methods

Strain maintenance

749 Unless otherwise noted, *C. elegans* strains were maintained at 20°C through

750 conventional methods (Brenner 1974). The animals were grown on NGM plates seeded

751 with OP50 bacteria. *glp-1(ar202)* is a <u>t</u>emperature <u>s</u>ensitive (*ts*) allele and strains with

752 this allele were maintained at 15°C. A complete list of strains used in this study is

753 provided in **S1 Table**.

Generation of CRIPSR alleles and transgenes for *lag-1*

lag-1(oz530)[lag-1::3xHA] allele (Fig 2A) was crossed into relevant genetic backgrounds

for quantification of LAG-1 accumulation. This allele was generated through in vitro

assembled RNA protein complex (RNP) CRISPR (Arribere *et al.* 2014; Paix *et al.* 2015).

Briefly, each component was pooled in volume of 20µl with final concentration as

follows before injection: 0.25 ng/µl HiFi Cas-9 (IDT, #1081061), 30 µM tracrRNA (IDT,

#1072534), 15 µM *lag-1* specific crRNA(atacagtaatcccgcgagag<u>NGG</u>) (IDT, Alt-R™),

0.02 µM *lag-1* specific single strand DNA repair templates (3xHA sequences are

underlined)

(cctacaaatcattggaacgacatggaccgtgcagaattgtgtccaattac<u>TACCCTTACGACGTGCCAGAT</u>

<u>TACGCTTACCCCTACGACGTACCAGACTACGCCTACCCATACGACGTCCCAGACTA</u>

CGCTTAGattAAactctcgcgggattactgtatctttatattgtctcctaatttctcccaattcgt) (IDT), 15 µM *pha-*

1 crRNA and 0.02 µM *pha-1(e2123)* specific repair template (Ward 2014). The injected

animals were raised at 25°C to identify *pha-1(e2123)* rescued animals. PCR was used

to screen for edits (see **S2 Table** for oligonucleotide information). Generated alleles

were then examined by Sanger sequencing to ensure there were no extraneous

mutations. One allele, *lag-1(oz530),* outcrossed twice with wild type, was used for

further analysis.

lag-1(oz536oz537)[lag-1::degron::3xHA] CRISPR allele was used to assess LAG-1

protein function in the germ cells (**Fig 2A**). This allele was generated through the Self

Excising Cassette (SEC) method (Dickinson *et al.* 2015), due to the insert size. The

SEC contains *sqt-1* roller marker and hygromycin antibiotic selection marker to facilitate

the screening (Dickinson *et al.* 2015). Briefly, each component was mixed in a final

volume of 20 µl with the following concentration prior injection of wild type animals: 50

ng/µl Cas-9 plasmid (pDD121, *cas-9* driven by *eft-3* promoter), 50 ng/µl *lag-1* small

guide RNA (sgRNA) plasmid (*lag-1* sgRNA sequence atacagtaatcccgcgagag was

cloned into plasmid DR274 U6 through *BsaI* site), 10 ng/µl *lag-1* specific repair template

and 2.5 ng/µl *myo-2p::gfp* co-injection marker. The *lag-1* specific repair template was

constructed through Golden Gate cloning method (Schwartz and Jorgensen 2016). The

injected animals were raised at 20°C for three days prior of adding 500 µl of 5 mg/µl

hygromycin. Six days later, animals that survived the antibiotic treatment and without

myo-2p::gfp injection marker were selected for PCR to screen for inserts (see **S2 Table**

for oligonucleotide information). *lag-1(oz536)*(rollers, with SEC in) was verified through

Sanger sequencing, followed by two times outcrosses with wild type prior to heat shock

to remove the SEC (Dickinson *et al.* 2015) to generate *lag-1(oz536oz537)*

lag-1(ozIs43)[lag-1::3xFLAG::BioTag + unc-119] transgene fosmid (**S5A Fig**) was

generated through insertion of short 3xFLAG and BioTag sequence through

Recombinering (Sharan *et al.* 2009) at the C-termimal of LAG-1 in pCC1FOS vector

(Source BioScience , CBGtg9050C1288D). BioTag is a 23 amino acid peptide that can

be recognized specifically by biotin ligase BirA in a sequence dependent manner (Ooi *et*

al. 2009). All transgenic alleles were generated through bombardment and *unc-119*

rescue; three alleles were found to rescue the phenotype of *lag-1(tm3052)*[loss of

function allele of *lag-1*]. One of the alleles, *lag-1(ozIs43)* was chosen to generate strain

BS1193 (**S1 Table**) and used for germline LAG-1 ChIP-seq analysis (**Fig 4**).

Auxin treatment

The auxin treatment was used to degrade endogenous LAG-1 protein in the germline.

Indole 3-acetic acid (IAA), the native plant hormone used in this study was ordered from

Alfa Aesar (#A10556) (Zhang *et al.* 2015). Like Zhang et al., 2015, we noticed that high

concentration of auxin (4 mM) inhibited OP50 bacteria growth. Therefore, all treatments

were done in NGM plates supplemented with 1 mM auxin (auxin plates). The un-seeded

auxin plates were stored at 4°C in the dark, and used within two weeks. The auxin

plates were seeded with OP50 bacteria and kept at 20°C for 24 hours before use. The

auxin treatment was conducted either at 20°C (**Fig 2E**) or at 25°C (**Fig 7**, **S8-10 Fig**)

with seeded plates preincubated at appropriate temperature for 2 hours. To achieve

germline specific removal of LAG-1 protein, three different germline-expressed TIR1

transgenes were tested for degradation of LAG-1 protein. Efficacy of LAG-1 protein

degradation was assayed by checking mRNA abundance of GLP-1/LAG-1

transcriptional targets, *lst-1* & *sygl-1*, via qPCR. The TIR1 transgenes were driven by

either *sun-1* promoter (Zhang *et al.* 2015), *pie-1* promoter (Kasimatis *et al.* 2018) or *gld-*

1 promoter (Zhang *et al.* 2015). We found that *gld-1* promoter driven TIR1 transgene

gave the largest reduction of *lst-1* and *sygl-1* mRNA, and was used in this study.

Immunostaining and Progenitor Zone (PZ) size

The germline extrusion and staining procedure is as described (Brenner & Schedl

2016). The primary antibodies used are: WAPL-1 at 1:2000 (Rabbit, Novus, #49300002)

(Mohammad *et al.* 2018), HIM-3 at 1:600 (Rabbit, Novus, 53470002), CYE-1 at 1:100

(mouse) (Brodigan *et al.* 2003), and HA at 1:100 (Rat, Roche, #11867423001), V5 at

1:1000 (mouse, Bio-rad, #MCA1360)(Shin et al., 2017), OLLAS at 1:500 (Rat, Novus,

#NBP1-96713)(Shin et al., 2017). Both CYE-1 and WAPL-1 antibodies staining gives

similar estimate of PZ size compared to REC-8 in *C. elegans* germline (Mohammad *et*

al. 2018). Hyperstack images were captured and the surface nuclei slice usually have

the best signal for both antibodies. Therefore this slice was used to count rows of cells

are positive for either antibody as described previously (Mohammad *et al.* 2018).

Protein quantification

In this work, detailed protein expression quantification method was developed to study

LAG-1, FBF-2, LST-1 and SYGL-1 protein expression regulation in relevant genetic

backgrounds. Quantification of protein in the distal end of antibody stained germlines

was done as described (Brenner and Schedl 2016) with noted differences.

Since ChIP-seq experiments were performed at L4 stage, and also LAG-1 protein

expression seems to be more robustly present in the progenitor zone of L4 stage

animals as compared to 1-day old adults, we carried out quantification of LAG-1

accumulation in dissected gonads of L4 stage worms. FBF-2 quantification is carried out

in young-adult germlines (~8 hr post-L4/adult molt). At this stage, the wild type worms

have started laying eggs and in tumorous animals like *gld-1* null germlines, the proximal

tumor is still small enough to have not invaded the entire gonad; thus, the polarity of the

germline is still intact (Mohammad *et al.* 2018). Auxin treatment of animals carrying

tagged LST-1 and SYGL-1 was started at mid-L4 stage and dissection and staining,

thus, quantification of protein was carried out at 2 or 4 hours later at 25°C.

Apart from the protein to be quantified, gonads were co-stained with DAPI and antibody

against WAPL-1. DAPI is used to count and mark cell diameter (cd) and WAPL-1 was

used to measure progenitor zone length. WAPL-1 staining was also used to distinguish

between different genotypes that were dissected and stained together. Hyperstack

images were captured using a 63X objective lens on a spinning disk confocal

microscope (PerkinElmer-Cetus, Norwalk, CT). Exposure time, which is kept constant

for an individual experiment, is set by using auto exposure in Volocity software (Perkin-

Elmer) for each experiment using an epitope-tagged strain in wild type background. To

capture distal the end of germline, two overlapping hyperstack images were captured to

give a coverage length of ~120-150 microns (equivalent to ~40-50 cd). The images

were exported, stitched in pairs, using Bio-Formats and Stitching plugin in Fiji (Preibisch

et al. 2009; Linkert *et al.* 2010; Schindelin *et al.* 2012). In Fiji, DAPI images were used to

draw a line, starting at the distal end to the desired cd (Brenner and Schedl 2016).

FBF-2 protein was quantified 35 cd from the distal end of the germlines, whereas

quantification of LAG-1, SYGL-1 and LST-1 was done using 25 cd. A width of 75 pixel

was used to collect protein levels for each line. Some of the gonads become out of

focus across their length, thus intensity values were collected for two or more slices of

the hyperstacks for every pixel and a maximum value is selected for every pixel of the

line. After manually drawing the line, ImageJ API

(https://imagej.nih.gov/ij/developer/api/index.html) was used with custom python scripts

to collect and store intensity data. The pixel intensities thus obtained, were processed

similar to *Brenner & Schedl 2016* to produce protein level graphs. *Brenner & Schedl*

2016, have quantified GLD-1 levels using antibody against GLD-1, they have used *gld-*

1(-) germlines to remove "background noise". Since our quantification involved antibody

staining against epitope-tagged animals, we instead, used N2 to remove "background

noise". Also instead of *spo-11(-)* germlines as internal normalization controls, we have

used strains carrying epitope-tagged protein, without any other mutations, as internal

controls and have termed them as wild type germlines.

mRNA detection and quantification

The procedure for *In situ* hybridization is adopted from Jones et al., 1996. Briefly, young

adult animals were dissected in a glass dish. The dissected gonads were fixed with 3%

paraformaldehyde / 0.25% glutaraldehyde / 0.1 M K2HPO4 (pH7.2) for 2 hours at RT,

followed by post-fixed with 100% methanol at -20°C. After three washes in PBST to

remove residual methanol, the gonads were incubated with 50 µg/ml protease K in

PBST for 30 mins, followed by 15 mins re-fixation in 3% paraformaldehyde / 0.25%

glutaraldehyde / 0.1 M K2HPO4 (pH7.2), three washes in PBST, 15 mins incubation in

PBST with 2 mg/ml glycine to remove residue aldehyde, then another three washes in

PBST prior to hybridization. The hybridization buffer contains: 5xSSC, 50% deionized

formamide, 100 µg/ml autoclaved Herring sperm DNA, 50 µg/ml Heparin and 0.1 %

Tween-20. The gonads were pre-hybridized in PBST/hybridization buffer (1:1) for 5

mins at 48°C, followed by 1 hour of incubation in hybridization buffer. PCR primers (see

S2 Table for oligonucleotide information) were used to amplify *sygl-1* and *lst-1* cDNA

from total RNA preparation and 100 ng of DNA was used to perform 35 rounds of single

oligo PCR to incorporate digoxigenin labelled nucleotide into single stranded DNA

probes that's complementary to *sygl-1* and *lst-1* mRNA. Each probe was diluted in 1 ml

hybridization buffer (for ten 100 µl hybridization). After O/N incubation in either probe,

gonads were washed with hybridization buffer three times to remove the excess probe,

followed by three washes in PBST. The gonads were then incubated with anti-

digoxigenin antibody (Sigma, #11333089001) O/N at 4°C, followed by three washes in

PBST, then developed in AP substrate BCIP/NBT (Sigma, #B5655). The detection of

sygl-1 takes 30mins and *lst-1* takes 1 hour.

Single molecule fluorescent *In situ* hybridization (smFISH) experiments were similarly

performed as described (Spike et al., 2014). *lag-1* Stellaris smFISH probes was

designed by and ordered from Biosearch Technologies. The fixed dissected gonads

were incubated with probe at final concentration of 5 μM, at 37°C O/N. All the

subsequent washes were done at 37°C: three washes in 2xSSC with 10% formamide,

three washes in PBST. DAPI was introduced in the last wash to stain DNA. Gonads

were mounted and imaged similar to immunostaining.

To quantify smFISH foci, hyperstack images of the distal end of the gonads were

acquired using a 63X objective lens on a spinning disk confocal microscope

(PerkinElmer-Cetus, Norwalk, CT). A suitable z-stack distance (0.4 micron) was used in

order to capture all smFISH foci. Image acquisition along the entire thickness of the

gonad results in progressive bleaching of the smFISH foci, hampering its accurate

quantification. To circumvent this issue, instead of covering the entire thickness of the

germline, we only took images of the germline covering one germ cell thickness from

the surface. Two overlapping hyperstack images were acquired for each gonadal arm to

get at least 25 cell diameters from the distal end. Images were stitched and further

processed in Fiji. For every gonadal image, all the other surrounding artifacts were

cleared and gonads were rotated and cropped to get an optimum size to minimize the

computational cost of further processing. 3D Objects Counter plugin (Bolte and

Cordelières 2006) was used to get smFISH foci and their positions. A threshold value

was selected, which would include most of the foci. For every germline, the first 25 cell

diameters were marked, forming an imaginary line passing through the center of the

gonadal tube. These markers and imaginary lines with a line-width of 100 pixels,

roughly corresponding to about 10 microns, were used to assign smFISH foci to a

particular cell diameter.

Chromatin Immunoprecipitation (ChIP)

The overall ChIP procedure was adapted from Berkseth et al., 2013.

L4 stages animals raised at 20°C were used for both whole worm and germline specific

LAG-1 ChIP experiments in large quantities. Typically, freshly starved plates with lots of

L1 stage animals were chunked onto NA22 bacteria plates (Berkseth *et al.* 2013), and

grown to adult stage at 20°C. Gravid adult hermaphrodites were bleached to get

synchronized L1 population, which were plated onto NA22 plates (NA22 bacteria plates

were seeded three days prior). Each NA22 plate (100 mm x 15 mm) housed around

5×10^4 animals till L4 stages or 2×10^4 animals till adult stage. If required, repeated

growing/bleaching were used to obtain even larger quantities of animals. The L4

animals were washed off the plate using PBST and were further washed three times

with PBST to remove NA22 bacteria, and frozen into small worm balls in liquid nitrogen.

The frozen worm balls were ground up in liquid nitrogen with mortar and pestles. The

worm powder was fixed in 1% paraformaldehyde in PBS for 15 mins, and then post-

fixed with 125 mM glycine for 5 mins. After centrifugation, the worm pellet was washed

in PBS three times before re-suspending in 1% SDS buffer. The worm suspension

containing fixed DNA-protein complexes was sheared through Bioruptor (Diagenode,

Denville, NJ) on the high setting for 20 cycles (30 seconds ON/30 seconds OFF), to

generate ~ 500 base pair length fragmented DNA. After centrifugation, the supernatant

was pre-cleared by protein-G Dynabeads (ThermoFisher, Waltham, MA) before the

immunoprecipitation experiment.

For immunoprecipitation of LAG-1 from the whole worm lysate, different GFP and FLAG

antibodies were first tested for their ability to immunoprecipitate LAG-1 (**S4B Fig**). Two

antibodies were chosen for further analysis: FLAG M2, which has have been

successfully used in other ChIP experiments (Gerace and Moazed 2015; Chen *et al.*

2018), and anti-GFP antibody, which immunoprecipitated the highest amount of LAG-1

(**S4B Fig**). Two μg of either FLAG (mouse, Sigma, #F1804) or GFP (goat, Rockland,

#600-101-215) antibodies was used to immunoprecipitate LAG-1 protein from 2 grams

of worm powder. After O/N incubation at 4°C, the precipitated DNA-protein complexes

were washed and eluted in 1% SDS for 10 mins. DNA was then reverse crosslinked to

prepare sequencing libraries following the manufacturer's instructions (Kapa Biosystem,

#KK8500). To immunoprecipitate germline LAG-1, L4 stage animals were used as it

lacks LAG-1 from late stage of oogenesis that is not involved in GLP-1 signaling (**Fig 2**).

However, direct ChIP with streptavidin beads was not successful, likely due to

interference from large amounts of endogenous biotinylated protein (Ooi *et al.* 2009;

Watts *et al.* 2018); hence, a sequential ChIP method was developed to overcome this

issue. First, 300 μl FLAG M2 beads/6 grams of worm powder per replicate (Sigma,

#M8823) was first used to precipitate all LAG-1 proteins from both germline and somatic

tissues, following the same procedure from whole worm ChIP. After eluting in 1% SDS,

the DNA-protein complexes was diluted five times and re-immunoprecipitated with

streptavidin beads to specifically pull-down biotinylated LAG-1. Since the binding

between biotin and streptavidin is very stable, the precipitated DNA-protein complex

was washed in 2% SDS, 1% SDS for 10 mins each to remove any none specific

binding. Unlike the whole worm ChIP, the DNA amount after sequential ChIP was ultra

low, so the NGS libraries were prepared directly with DNA on the streptavidin beads

following manufacturer's instructions (Kapa Biosystem, #KK8500).

Sequencing and Bioinformatic analysis was performed by Genome Technology Access

Center at the McDonnell Genome Institute (GTAC @ MGI, Washington University in St.

Louis). Short sequencing reads (50 nt) were aligned back to *C. elegans* reference

genome and ChIP-seq peaks were identified using MACS2.0 (Feng *et al.* 2012) by

comparing ChIP library to input DNA library, for both whole worm and germline ChIP

experiments. The association of peak to closest gene annotation was done with

HOMER suite (Heinz *et al.* 2010). HOMER suite was also used to identify overly

represented motifs in ChIP-seq data. Gene list comparisons and Venn diagrams

preparation were both conducted in R.

RNA-seq analysis

All RNA-seq experiments conducted in this study used isolated gonads from dissected

worms with a tumorous germline. All strains carried *gld-1 gld-2* double mutant

background. For GLP-1 ON and OFF RNA-seq, gain of function (gf) allele *glp-1(ar202)*

or null allele *glp-1(q175)* were introduced into the *gld-1(q485) gld-2(q497)* mutant

background. *glp-1(ar202)* is a temperature sensitive gain of function allele, used to

produce cells throughout the germline that are undergoing GLP-1 signaling. This

increased our ability to detect gene expression differences. For the auxin induced LAG-

1 degradation time course RNA-seq experiments, *lag-1(oz536oz537)* and *ieSi64[gld-*

1p::tir1] were introduced into *gld-1(q485) gld-2(q497); glp-1(ar202)*. Animals were raised

at 25°C for 48 hours from synchronized L1s population before dissection. For shorter

periods of auxin treatment (e.g, 2 hours and 4 hours), animals grown on NGM plates

were transferred to seeded auxin plates (pre-conditioned to 25°C) and treated for the

required time, prior to gonad extrusion.

1000

Fifty young adults were dissected in a glass dish and the gonads were dissected away

from rest of the animal's body at the spermatheca. The isolated gonads were

centrifuged for 15 sec and re-suspended in 500 µl Trizol (Invitrogen, #15596026). The

RNA isolation was done according to manufacturer's instructions. RNA was then treated

with DNase (Thermo Fisher Scientific, #EN0525) to remove the contaminating genomic

DNA before proceeding to the removal of ribosome RNA (rRNA). The rRNA were

removed with the Ribo-Zero kit (Illumina, #RZE1206) according to manufacturer's

instructions. The conversion of RNA to dsDNA was done with NEBNext RNA first strand

(NEB, #E7525S) and second strand (NEB, E6111S) kits. The converted dsDNA was

used to prepare NGS libraries similar to ChIP-seq (Kapa Biosystem, #KK8500).

1011

The sequencing and bioinformatic analysis was performed by the Genome Technology

Access Center at the McDonnell Genome Institute. The samples were sequenced on an

Illumina HiSeq 3000 with 50 nt reads to a mean depth of 34.5 million reads and then

aligned back to the Ensembl Release 76 *C. elegans* genome with STAR version 2.0.4b

(Dobin *et al.* 2013). Gene counts were then quantitated with Subread:featureCount

version 1.4.5 (Liao *et al.* 2014). The gene counts were then TMM scaled with EdgeR

(Robinson *et al.* 2009) and voomWithQualityWeights (Liu *et al.* 2015) transformed with

Limma (Ritchie *et al.* 2015). A Limma generalized linear model without intercept was

then fitted to the data and tested for statistical significance via Limma's empirical Bayes

moderated t-statistics. The genes were considered to be activated by GLP-1 if the

following three criteria were met: (1) its expression in GLP-1 ON gonads was at least

2.0 fold compared to GLP-1 OFF gonads. (2) the Benjamini-Hochberg false discovery

rate (FDR) is less than 0.05 and (3) the gene expression level as determined by Counts

Per Million (CPM) must be greater than 2.0. The same criteria were applied to obtain

the gene list for either activated or repressed by LAG-1 in the time course RNA-seq

analysis. Gene list comparisons and Venn diagrams preparation were both done in R.

1028

Immunoblot analysis

The immunobloting was performed as described (Jones et al., 1996). Immunoblot was

used to determine the efficiency of different commercial antibodies in pull-down of LAG-

1 in ChIP conditions (**S4 Fig**). After ChIP experiments, the protein DNA complexes were

eluted in 1x SDS sample buffer by heating at 95°C for 5 mins. The eluted samples were

run on 10 % SDS-PAGE gel, then transferred to PVDF membrane. To detect LAG-1

after ChIP, FLAG antibody (Sigma, F1804) was used at 1:1250 dilution. The detection

of signal was achieved with HRP-conjugated anti-mouse secondary antibody (Jackson

Immunoresearch, #115-035-146).

1038

qPCR analysis

For analysis of mRNA abundance in tumorous germline of relevant genotypes (**Fig 5,**

7), total mRNA was isolated from dissected gonads with Trizol (Invitrogen, #15596026)

following manufacturer's protocol. The RNA was then DNase treated before cDNA

synthesis with Superscript III (Invitrogen, #18080051). The cDNAs were usually diluted

five fold prior to qPCR analysis. For analysis of DNA enrichment after ChIP

experiments, the DNA concentration was determined and the same amount of DNA was

used for each qPCR reaction. iTaq Universal SYBR Green Supermix (Biorad,

#1725121) and CFX96 Real time System (Biorad) were used for qPCR analysis. The

delta Ct method was used to calculate the relative mRNA abundance compare to the

ama-1 control (Kumar *et al.* 2019). The primers used for qPCR analysis are listed in **S2**

Table.

1051

Data processing and statistical analysis

Data stored as either text or csv files, were imported into a R programming

envrionment, and graphs were plotted using the ggplot2 package (http://ggplot2.org).

The significance bars were generated using the ggsignif package (https://CRAN.R-

project.org/package=ggsignif) in R. Statistical significance was determined using either

two-tailed Student's t-test (for n < 30) or Z-test (for n ≥ 30).

1058

Acknowledgments

We thank our colleagues in worm community for unpublished alleles: Iva Greenwald,

Columbia University, *lag-1(ar611);* William Kelly, Emory University, *ckSi11.* We also

thank Julie Ni and Sam Gu for their help in the initial phases of ChIP-seq analysis and

our colleagues Mike Nonet, Jane Hubbard, Iva Greenwald, Dave Hansen, David

Greenstein, Jordan Ward, and Dan Dickenson for helpful advice and discussions. We

thank WormBase, funded by the National Human Genome Research Institute; the

Caenorhabditis Genetics Center, funded by National Institutes of Health Office of

Research Infrastructure Programs; and Shohei Mitani and the Japanese National

Bioresource Project for strains. This work was supported by National Institutes of Health

R01 GM-100756 to T.S.

1070

1071 **References**

1072

Arribere J. A., R. T. Bell, B. X. H. Fu, K. L. Artiles, P. S. Hartman, *et al.*, 2014 Efficient

marker-free recovery of custom genetic modifications with CRISPR/Cas9 in

caenorhabditis elegans. Genetics. https://doi.org/10.1534/genetics.114.169730

Berkseth M., K. Ikegami, S. Arur, J. D. Lieb, and D. Zarkower, 2013 TRA-1 ChIP-seq

reveals regulators of sexual differentiation and multilevel feedback in nematode sex

determination. Proc. Natl. Acad. Sci. 110: 16033–16038.

https://doi.org/10.1073/pnas.1312087110

Berset T., E. F. Hoier, G. Battu, S. Canevascini, and A. Hajnal, 2001 Notch inhibition of

RAS signaling through MAP kinase phosphatase LIP-1 during C. elegans vulval

development. Science (80-.). https://doi.org/10.1126/science.1055642

Bolte S., and F. P. Cordelières, 2006 A guided tour into subcellular colocalization

analysis in light microscopy. J. Microsc. 224: 213–232.

Bray S. J., 2006 Notch signalling: A simple pathway becomes complex. Nat. Rev. Mol.

Cell Biol.

Bray S. J., 2016 Notch signalling in context. Nat. Rev. Mol. Cell Biol.

Brenner S., 1974 The genetics of Caenorhabditis elegans. Genetics.

Brenner J. L., and T. Schedl, 2016 Germline stem cell differentiation entails regional

control of cell fate regulator GLD-1 in Caenorhabditis elegans. Genetics.

https://doi.org/10.1534/genetics.115.185678

Brodigan T. M., J. i. Liu, M. Park, E. T. Kipreos, and M. Krause, 2003 Cyclin E

expression during development in Caenorhabditis elegans. Dev. Biol.

https://doi.org/10.1016/S0012-1606(02)00032-5

Chen X., L. Lu, S. Qian, M. Scalf, L. M. Smith, *et al.*, 2018 Canonical and noncanonical

actions of arabidopsis histone deacetylases in ribosomal RNA processing. Plant

Cell. https://doi.org/10.1105/tpc.17.00626

Choi V. N., S. K. Park, and B. J. Hwang, 2013 Clustered LAG-1 binding sites in lag-

1/CSL are involved in regulating lag-1 expression during lin-12/Notch-dependent

cell-fate specification. BMB Rep. https://doi.org/10.5483/BMBRep.2013.46.4.269

Christensen S., V. Kodoyianni, M. Bosenberg, L. Friedman, and J. Kimble, 1996 lag-1, a

gene required for lin-12 and glp-1 signaling in Caenorhabditis elegans, is

homologous to human CBF1 and Drosophila Su(H). Development.

Cronan J. E., 1990 Biotination of proteins in vivo. A post-translational modification to

label, purify, and study proteins. J. Biol. Chem.

Cusanovich D. A., B. Pavlovic, J. K. Pritchard, and Y. Gilad, 2014 The Functional

Consequences of Variation in Transcription Factor Binding. PLoS Genet.

https://doi.org/10.1371/journal.pgen.1004226

Dickinson D. J., A. M. Pani, J. K. Heppert, C. D. Higgins, and B. Goldstein, 2015

Streamlined genome engineering with a self-excising drug selection cassette.

Genetics. https://doi.org/10.1534/genetics.115.178335

Dobin A., C. A. Davis, F. Schlesinger, J. Drenkow, C. Zaleski, *et al.*, 2013 STAR:

Ultrafast universal RNA-seq aligner. Bioinformatics.

https://doi.org/10.1093/bioinformatics/bts635

E. J. A. Hubbard, and T. Schedl, 2019 Biology of the Caenorhabditis elegans Germline Stem Cell

System. Genetics. (In Press)

Feng J., T. Liu, B. Qin, Y. Zhang, and X. S. Liu, 2012 Identifying ChIP-seq enrichment

using MACS. Nat. Protoc. https://doi.org/10.1038/nprot.2012.101

Fox P. M., V. E. Vought, M. Hanazawa, M. H. Lee, E. M. Maine, *et al.*, 2011 Cyclin e

and CDK-2 regulate proliferative cell fate and cell cycle progression in the C.

elegans germline. Development. https://doi.org/10.1242/dev.059535

Fox P. M., and T. Schedl, 2015 Analysis of germline stem cell differentiation following

loss of GLP-1 notch activity in Caenorhabditis elegans. Genetics.

https://doi.org/10.1534/genetics.115.178061

Gerace E., and D. Moazed, 2015 Affinity Pull-Down of Proteins Using Anti-FLAG M2

Agarose Beads, in *Methods in Enzymology*,.

Gerstein M. B., Z. J. Lu, E. L. Van Nostrand, C. Cheng, B. I. Arshinoff, *et al.*, 2010

Integrative analysis of the Caenorhabditis elegans genome by the modENCODE

project. Science (80-.). https://doi.org/10.1126/science.1196914

Hansen D., E. J. A. Hubbard, and T. Schedl, 2004a Multi-pathway control of the

proliferation versus meiotic development decision in the Caenorhabditis elegans

germline. Dev. Biol. https://doi.org/10.1016/j.ydbio.2003.12.023

Hansen D., L. Wilson-Berry, T. Dang, and T. Schedl, 2004b Control of the proliferation

versus meiotic development decision in the C. elegans germline through regulation

of GLD-1 protein accumulation. Development. https://doi.org/10.1242/dev.00916

Haupt K. A., A. L. Enright, A. S. Ferdous, A. M. Kershner, H. Shin, *et al.*, 2019 The

molecular basis of LST-1 self-renewal activity and its control of stem cell pool size.

Development. https://doi.org/10.1242/dev.181644

Heinz S., C. Benner, N. Spann, E. Bertolino, Y. C. Lin, *et al.*, 2010 Simple Combinations

of Lineage-Determining Transcription Factors Prime cis-Regulatory Elements

Required for Macrophage and B Cell Identities. Mol. Cell.

https://doi.org/10.1016/j.molcel.2010.05.004

Jungkamp A. C., M. Stoeckius, D. Mecenas, D. Grün, G. Mastrobuoni, *et al.*, 2011 In

vivo and transcriptome-wide identification of RNA binding protein target sites. Mol.

Cell. https://doi.org/10.1016/j.molcel.2011.11.009

Kasimatis K. R., M. J. Moerdyk-Schauwecker, and P. C. Phillips, 2018 Auxin-mediated

sterility induction system for longevity and mating studies in Caenorhabditis

elegans. G3 Genes, Genomes, Genet. https://doi.org/10.1534/g3.118.200278

Kershner A. M., H. Shin, T. J. Hansen, and J. Kimble, 2014 Discovery of two GLP-

1/Notch target genes that account for the role of GLP-1/Notch signaling in stem cell

maintenance. Proc. Natl. Acad. Sci. https://doi.org/10.1073/pnas.1401861111

Kodoyianni V., E. M. Maine, and J. Kimble, 1992 Molecular basis of loss-of-function

mutations in the glp-1 gene of Caenorhabditis elegans. Mol. Biol. Cell.

https://doi.org/10.1091/mbc.3.11.1199

Kumar S., B. M. Egan, Z. Kocsisova, D. L. Schneider, J. T. Murphy, *et al.*, 2019

Lifespan Extension in C. elegans Caused by Bacterial Colonization of the Intestine

and Subsequent Activation of an Innate Immune Response. Dev. Cell.

https://doi.org/10.1016/j.devcel.2019.03.010

Lamont L. B., S. L. Crittenden, D. Bernstein, M. Wickens, and J. Kimble, 2004 FBF-1

and FBF-2 regulate the size of the mitotic region in the C. elegans germline. Dev.

Cell. https://doi.org/10.1016/j.devcel.2004.09.013

Lee M. H., and T. Schedl, 2001 Identification of in vivo mRNA targets of GLD-1, a maxi-

KH motif containing protein required for C. elegans germ cell development. Genes

Dev. https://doi.org/10.1101/gad.915901

Lee M. H., B. Hook, L. B. Lamont, M. Wickens, and J. Kimble, 2006 LIP-1 phosphatase

controls the extent of germline proliferation in Caenorhabditis elegans. EMBO J.

https://doi.org/10.1038/sj.emboj.7600901

Lee C. H., E. B. Sorensen, T. R. Lynch, and J. Kimble, 2016 C. elegans GLP-1/Notch

activates transcription in a probability gradient across the germline stem cell pool.

Elife. https://doi.org/10.7554/eLife.18370

Lee C., H. Shin, and J. Kimble, 2019 Dynamics of Notch-Dependent Transcriptional

Bursting in Its Native Context. Dev. Cell.

https://doi.org/10.1016/j.devcel.2019.07.001

Lenstra T. L., and F. C. P. Holstege, 2012 The discrepancy between chromatin factor

location and effect. Nucl. (United States). https://doi.org/10.4161/nucl.19513

Liao Y., G. K. Smyth, and W. Shi, 2014 FeatureCounts: An efficient general purpose

program for assigning sequence reads to genomic features. Bioinformatics.

https://doi.org/10.1093/bioinformatics/btt656

Linkert M., C. T. Rueden, C. Allan, J. M. Burel, W. Moore, *et al.*, 2010 Metadata

matters: Access to image data in the real world. J. Cell Biol. 189: 777–782.

Liu R., A. Z. Holik, S. Su, N. Jansz, K. Chen, *et al.*, 2015 Why weight? Modelling sample

and observational level variability improves power in RNA-seq analyses. Nucleic

Acids Res. https://doi.org/10.1093/nar/gkv412

Mohammad A., K. Vanden Broek, C. Wang, A. Daryabeigi, V. Jantsch, *et al.*, 2018

Initiation of meiotic development is controlled by three post-transcriptional

pathways in caenorhabditis elegans. Genetics.

https://doi.org/10.1534/genetics.118.300985

Neves A., and J. R. Priess, 2005 The REF-1 family of bHLH transcription factors pattern

C. elegans embryos through Notch-dependent and Notch-independent pathways.

Dev. Cell. https://doi.org/10.1016/j.devcel.2005.03.012

Nostrand E. L. Van, and S. K. Kim, 2013 Integrative analysis of C. elegans

modENCODE ChIP-seq data sets to infer gene regulatory interactions. Genome

Res. https://doi.org/10.1101/gr.152876.112

Ooi S. L., J. G. Henikoff, and S. Henikoff, 2009 A native chromatin purification system

for epigenomic profiling in Caenorhabditis elegans. Nucleic Acids Res.

https://doi.org/10.1093/nar/gkp1090

Paix A., A. Folkmann, D. Rasoloson, and G. Seydoux, 2015 High efficiency, homology-

directed genome editing in Caenorhabditis elegans using CRISPR-

Cas9ribonucleoprotein complexes. Genetics.

https://doi.org/10.1534/genetics.115.179382

Pepper A. S. R., D. J. Killian, and E. J. A. Hubbard, 2003 Genetic analysis of

Caenorhabditis elegans glp-1 mutants suggests receptor interaction or competition.

Genetics.

Preibisch S., S. Saalfeld, and P. Tomancak, 2009 Globally optimal stitching of tiled 3D

microscopic image acquisitions. Bioinformatics 25: 1463–1465.

https://doi.org/10.1093/bioinformatics/btp184

Ritchie M. E., B. Phipson, D. Wu, Y. Hu, C. W. Law, *et al.*, 2015 Limma powers

differential expression analyses for RNA-sequencing and microarray studies.

Nucleic Acids Res. https://doi.org/10.1093/nar/gkv007

Robinson M. D., D. J. McCarthy, and G. K. Smyth, 2009 edgeR: A Bioconductor

package for differential expression analysis of digital gene expression data.

Bioinformatics. https://doi.org/10.1093/bioinformatics/btp616

Schindelin J., I. Arganda-Carreras, E. Frise, V. Kaynig, M. Longair, *et al.*, 2012 Fiji: An

open-source platform for biological-image analysis. Nat. Methods.

Schwartz M. L., and E. M. Jorgensen, 2016 SapTrap, a toolkit for high-throughput

CRISPR/Cas9 gene modification in Caenorhabditis elegans. Genetics.

https://doi.org/10.1534/genetics.115.184275

Seelk S., I. Adrian-Kalchhauser, B. Hargitai, M. Hajduskova, S. Gutnik, *et al.*, 2016

Increasing notch signaling antagonizes PRC2-mediated silencing to promote

reprograming of germ cells into neurons. Elife.

https://doi.org/10.7554/eLife.15477.001

Sharan S. K., L. C. Thomason, S. G. Kuznetsov, and D. L. Court, 2009

Recombineering: A homologous recombination-based method of genetic

engineering. Nat. Protoc. https://doi.org/10.1038/nprot.2008.227

Shin H., K. A. Haupt, A. M. Kershner, P. Kroll-Conner, M. Wickens, *et al.*, 2017 SYGL-1

and LST-1 link niche signaling to PUF RNA repression for stem cell maintenance in

Caenorhabditis elegans. PLoS Genet. https://doi.org/10.1371/journal.pgen.1007121

Spike C. A., D. Coetzee, C. Eichten, X. Wang, D. Hansen, *et al.*, 2014 The TRIM-NHL

protein LIN-41 and the OMA RNA-binding proteins antagonistically control the

prophase-to-metaphase transition and growth of caenorhabditis elegans oocytes.

Genetics. https://doi.org/10.1534/genetics.114.168831

Wang X., J. R. Olson, D. Rasoloson, M. Ellenbecker, J. Bailey, *et al.*, 2016 Dynein light

chain DLC-1 promotes localization and function of the PUF protein FBF-2 in

germline progenitor cells. Dev. https://doi.org/10.1242/dev.140921

Ward J. D., 2014 Rapid and precise engineering of the caenorhabditis elegans genome

with lethal mutation co-conversion and inactivation of NHEJ repair. Genetics.

https://doi.org/10.1534/genetics.114.172361

Watts J. S., D. G. Morton, K. J. Kemphues, and J. L. Watts, 2018 The biotin-ligating

protein BPL-1 is critical for lipid biosynthesis and polarization of the Caenorhabditis

elegans embryo. J. Biol. Chem. https://doi.org/10.1074/jbc.M117.798553

Yoo A. S., C. Bais, and I. Greenwald, 2004 Crosstalk between the EGFR and LIN-

12/Notch Pathways in C. elegans Vulval Development. Science (80-.).

https://doi.org/10.1126/science.1091639

Yoo A. S., and I. Greenwald, 2005 Developmental biology: LIN-12/notch activation

leads to microRNA-mediated down-regulation of Vav in C. elegans. Science (80-.).

https://doi.org/10.1126/science.1119481

Zhang L., J. D. Ward, Z. Cheng, and A. F. Dernburg, 2015 The auxin-inducible

degradation (AID) system enables versatile conditional protein depletion in C.

elegans. Development 142: 4374–4384. https://doi.org/10.1242/dev.129635

1250

1251 **Figure legends**

1252

Figure 1. Overview of GLP-1 signaling in the distal germline of *C. elegans*. (A)

Schematic of distal end of germline in adult hermaphrodite. Distal end of the germline is

capped by a somatic distal tip cell (DTC). Progenitor zone cells are shown in green;

meiotic nuclei are red. Dashed line indicate meiotic entry. GLP-1 signaling maintains

germline stem cell fate and GLP-1 mediated transcription occupy first 6~7 cell diameter

(cd) in the distal germline (Lee et al., 2016, Lee et al., 2019). GLP-1 transcriptional

targets, *lst-1* and *sygl-1, are* normally expressed in first 5 cd and 10 cd, respectively

(Shin et al., 2017). (B) Genetic pathway controlling stem cell vs meiotic entry balance in

the distal end of *C. elegans* germline. GLP-1 signaling employs its targets, LST-1 and

SYGL-1, to repress meiotic entry pathway genes, GLD-1, GLD-2 and SCF^{PROM-1}. GLP-

1(ICD), GLP-1 intracellular domain.

1264

Figure 2. LAG-1 is cell-autonomously required for the germline stem cell fate. (A)

Diagrams of two alleles of *lag-1* at endogenous locus*, lag-1(oz530)[lag-1::3xHA]* (top)

and *lag-1(oz536oz537)[lag-1::degron::3xHA]* (bottom). Purple boxes, exons; lines,

introns; pink boxes, untranslated region; yellow boxes, 3xHA; green box, degron. (B &

C) Images of HA-stained (LAG-1, yellow) germlines from dissected hermaphrodites, co-

stained with DAPI (cyan) for (B) adult and (C) L4 stage. Asterisk, distal end; dashed

lines, meiotic entry; solid white lines, positions of LAG-1 accumulation; red arrowhead,

sheath cell nuclei; white arrows, spermatheca. (D & E) Images of (D) HA-stained (LAG-

1, yellow) and (E) CYE-1 (green), HIM-3-stained (red) germlines from dissected

hermaphrodites. L4 stage animals with the following genotype *lag-1(oz536oz537)[lag-*

1::degron::3xHA]; ieSi64[gld-1p::TIR1::mRuby::gld-1 3'UTR] were treated with or without

auxin for 4 hours (top two panels) or 24 hours (bottom two panels). Asterisk, distal end;

dashed lines, meiotic entry; solid white lines, positions of LAG accumulation; red

arrowhead, sheath cell nuclei. Scale bar is 10 μm.

1279

Figure 3. Post-transcriptional regulation of LAG-1 by GLP-1 signaling. (A) Images

of HA-stained (LAG-1, yellow) germlines from dissected L4 hermaphrodites of the

indicated genotype. Asterisk, distal end; dashed white lines, meiotic entry. Scale bar is

10 μm. (B & C) Plot of LAG-1 levels for indicated genotype. *lag-1(oz530)[lag-1::3xHA]* is

used for quantification, see materials and methods for alleles used for other genotype.

Numbers in bracket shows the sample size. Dots, mean; Error bars, mean ± SD. (D)

Graph showing LAG-1 peak levels (see B) in L4 hermaphrodites of indicated genotype.

Numbers indicate mean values of LAG-1 level for each genotype and numbers in

bracket shows the sample size. Dots, mean; Error bars, mean ± SD. P-value ≤ 0.01 (*);

≤ 0.001 (**); ≤ 0.0001 (***); > 0.01 non-significant (NS.). (E) Model depicting genetic

control of peak accumulation for LAG-1. Sixty one percent of LAG-1 peak level is

attributed to GLP-1 signaling, in which GLP-1 transcriptional targets *lst-1* and *sygl-1*

account for 31% of LAG-1 peak level.

1293

Figure 4. Genome-wide identification of germline-specific LAG-1 targets by

sequential ChIP-seq. (A) Diagram showing LAG-1 was biotinylated specifically in

germline. A sequence dependent, germline-expressed biotin ligase, BirA, was used to

biotinylate LAG-1 via BioTag. The strain harbored two transgenes, i) *ckSi11[pie-*

1p::birA::gfp], where biotin ligase birA was driven by germline specific promoter *pie-1*

and ii) *ozIs43[lag-1p::lag-1::3xFLAG::BioTag]*. The complete genotype for this strain is

lag-1(tm3052); glp-1(ar202); ozIs43[lag-1p::lag-1::3xFLAG::BioTag]; ckSi11[pie-

1p::birA::gfp]. (B) The workflow of sequential ChIP-seq. Direct ChIP was not feasible

due to the interference of endo-geneous biotinylated proteins (Ooi et al., 2009; Watts et

al., 2018). Sequential ChIP-seq was developed to overcome this issue and to facilitate

DNA library preparation from ultra-low amount of DNA. (C) Genome browser tracks for

sygl-1, lst-1 and *mir-61/250* after sequential ChIP-seq. The raw reads were normalized

to the control, and the signals were presented as log2-fold change after normalization.

Black arrows, canonical LAG-1/CSL binding motif GTGGGAA (Tun et al., 1994; Brou et

al., 1994; Christensen et al., 1996). (D) Venn diagram showing 137 genes were

overlapped from three biological replicates of germline LAG-1 ChIP-seq analyses. Rep,

replicate. (E) The overly represented motif discovered by HOMER suite with germline

specific ChIP-seq data (top) and the reported canonical LAG-1/CSL binding motif

(bottom) (Tun et al., 1994; Brou et al., 1994; Christensen et al., 1996).

1313

Figure 5. Genome-wide identification of GLP-1-dependent genes by

transcriptomic analysis. (A) Schematics showing the harvesting of tumorous

germlines for either qPCR analysis or transcriptomic analysis. The genotype for GLP-1

ON animal (green) is *gld-2(q497) gld-1(q485); glp-1(ar202)* and GLP-1 OFF (red)

animal is *gld-2(q497) gld-1(q485); glp-1(q175).* (B) *In situ* hybridization (ISH) used to

detect *sygl-1* mRNA expression in GLP-1 ON and GLP-1 OFF young adult animals.

Dotted lines showing boundary of the germline. Black arrow indicates distal end of the

germline. (C) *sygl-1* and *lst-1* transcript levels analysis via qPCR with dissected

germline tumor. The expression level for either genes in GLP-1 OFF background were

set as one. Three biological replicates were conducted and two tailed t-test was used

for statistical analysis. P-value ≤ 0.01 (*); ≤ 0.001 (**); ≤ 0.0001 (***); > 0.01 non-

significant (NS.). (D) Venn diagram identifying GLP-1 transcriptional targets through

integrative genome-wide approach. Five biological replicates were used to conduct

transcriptomic analysis and identified 94 GLP-1-dependent genes (blue circle). The

GLP-1 transcriptional targets were defined as genes whose mRNA expression was

controlled by GLP-1 and also had LAG-1 occupied their promoter regions (red circle,

data from Fig 4D).

1331

Figure 6. FBF-2 expression is post-transcriptionally controlled by GLP-1 signaling

through transcriptional targets LST-1 and SYGL-1. (A) Images of V5-stained (FBF-2,

green) germlines from dissected young adult hermaphrodites of the indicated genotype.

Asterisk, distal end; dashed white line, meiotic entry. Scale bar is 10 µm. (B & C) Plot of

FBF-2 levels for indicated genotype. *fbf-2(q932)* is used for quantification, see materials

and methods for alleles used for other genotype. Numbers in bracket shows the sample

size. Dots, mean; error bars, mean ± SD. (D) Graph showing FBF-2 peak levels (see B)

in L4 hermaphrodites of indicated genotype. Dots, mean; Error bars, mean ± SD. P-

value ≤ 0.01 (*); ≤ 0.001 (**); ≤ 0.0001 (***); > 0.01 non-significant (NS.). Numbers

indicate mean values for each genotype. (E) Model depicting genetic control of FBF-2

peak levels. Total of 85% of FBF-2 peak level can be attributed to GLP-1 signaling

through GLP-1 transcriptional targets *lst-1* and *sygl-1*.

1344

Figure 7. Time-course transcriptomic analysis to identify LAG-1-dependent genes

in the germline. (A) Diagram showing time-course qPCR and transcriptomic analysis

following germline-specific degradation of LAG-1. This strain harbors germline-

expressed *tir1*, an ubqiutin E3 ligase that drive target degradation through proteolysis in

the presence of co-factor auxin (Zhang et al., 2015). The CRISPR allele *lag-*

1(oz536oz537) has degron fused C-terminally to LAG-1, which is recognized by *tir1*,

resulting in germline-specific degradation of LAG-1 protein. The complete genotype for

this strain is *gld-2(q497) gld-1(q485); glp-1(ar202); lag-1(oz536oz537)[lag-*

1::degron::3xHA]; ieSi64[gld-1p::TIR1::mRuby::gld-1 3'UTR]. Isolated gonads, after

dissection are used to quantify the abundance of RNAs, either through qPCR or

transciptomic analysis. (B & C) Quantification of two GLP-1/LAG-1 targets *sygl-1* and

lst-1 mRNA abundance after auxin treatment at different time points. Three biological

replicates were conducted and two-tailed t-test was used for statistical analysis. P-value

≤ 0.01 (*); ≤ 0.001 (**); ≤ 0.0001 (***); > 0.01 non-significant (NS.). (D) Venn diagram

showing genes' RNA expression dependent on LAG-1 at various time points through

transcriptomic analysis. The presented genes were also controlled by GLP-1. Four

biological replicates were conducted for each time point. (E) Expression of 5 more

genes were reduced after 4-hour auxin treatment, in addition to *lst-1* and *sgyl-1*

(identified at 2 hour auxin treatment). These genes' relative fold reduction in both GLP-1

OFF and LAG-1 OFF (via 4 hours auxin treatment) from transcriptomic analysis were

presented. (F) qPCR analysis for genes expression level from (E) in different mutant

background. The genotypes are, **GLP-1 ON**: *gld-2(q497) gld-1(q485); glp-1(ar202).*

GLP-1 OFF: *gld-2(q497) gld-1(q485); glp-1(q175).* **GLP-1 ON LST-1 SYGL-1 OFF**: *gld-*

2(q497) gld-1(q485) lst-1(ok814) sygl-1(tm5040); glp-1(ar202). Five biological replicates

were conducted for each genotype and two-tailed t-test was used for statistical analysis.

P-value ≤ 0.01 (*); ≤ 0.001 (**); ≤ 0.0001 (***); > 0.01 non-significant (NS.). (G) Model

depicting genetic control of RNA accumulation for genes from (E).

1372

Supplemental figure 1. Characterization of epitope-tagged *lag-1* alleles. (A & B)

The dead eggs and L1 arrest frequency (A) and progenitor zone length (B) in WT, *lag-*

1(oz530)[lag-1::3xHA] and *lag-1(oz536oz537)[lag-1::degron::3xHA].* (B) Graph showing

distance, in cell diameters, between the distal tip of the germline and the row of cells at

proximal end of the continuous zone of WAPL-1 staining for L4 hermaphrodites of

indicated genotype. Data are plotted as horizontal dot plots with each dot representing

length in cell diameter to zone end for one gonad. Numbers in bracket shows the

sample size. Thick vertical lines represent mean and horizontal lines represent mean ±

SD. P-value ≤ 0.01 (*); ≤ 0.001 (**); ≤ 0.0001 (***); > 0.01 non-significant (NS.). (C)

Progenitor zone length and Germline proliferation defective (Glp) phenotype after L4

stage animals were treated with or without auxin for either 4 hours or 24 hours (in Fig

2D & E). The genotype for auxin treatment was *lag-1(oz536oz537)[lag-*

1::degron::3xHA]; ieSi64[gld-1p::TIR1::mRuby::gld-1 3'UTR].

1386

1387 **Supplemental figure 2. *lag-1* mRNA expression analysis by smFISH.** (A) Z project

1388 of one nucleus thick (see materials and methods) distal germlines of L4

1389 hermaphrodites, probed for LAG-1 transcripts using smFISH (white) and DAPI (blue).

1390 Asterisk, distal end; Scale bar, 10 µm. (B & C) Density plot of *lag-1* mRNA foci for

1391 indicated genotype. Numbers in bracket shows the sample size.

1392

Supplemental figure 3. LAG-1 levels analysis. (A) Plot of LAG-1 levels for indicated

genotype. *lag-1(oz530)[lag-1::3xHA]* is used for quantification, see materials and

methods for alleles used for other genotype. Numbers in bracket shows the sample

size. Dots, mean; Error bars, mean ± SD. (B) Graph showing LAG-1 base levels (see

Fig 3B) in L4 hermaphrodites of indicated genotype. Numbers in bracket shows the

sample size. Dots, mean; Error bars, mean ± SD. P-value ≤ 0.01 (*); ≤ 0.001 (**); ≤

0.0001 (***); > 0.01 non-significant (NS.). Numbers indicate mean values for each

genotype.

1401

Supplemental figure 4. Genome-wide identification for LAG-1 targets in whole

animal by ChIP-seq analysis. (A) Schematics of *lag-1* allele at endogenous locus, *lag-*

1(ar611)[lag-1::gfp::3xFLAG] (unpublished strain from Iva Greenwald), used for whole

worm ChIP-seq analysis. (B) Different antibodies were tested to determine if they are

competent for ChIP. Same amount of extracts were used each ChIP, followed by

western blot analysis with FLAG antibody. Bottom band detects the heavy chain of

antibodies. FLAG antibody from Sigma and GFP antibody from Rockland were selected

for rest analysis below. (C) ChIP-qPCR analysis for *sygl-1* and *lst-1* promoter regions

bound by LAG-1. A non-peak region in the *xol-1* promoter was used as a negative

control (ctr). *** for p<0.0001. Error bars, mean ± SD. (D) Genome browser tracks

showing 10 kb genomic region for *sygl-1*, *lst-1 and mir-61/250,* 40 kb genomic region for

lag-1 and 20 kb genomic region for *glp-1 and lin-12* after ChIP-seq. Raw reads were

normalized to control, and signal intensity were presented as log2 fold change. Black

arrow heads, canonical LAG-1/CSL binding motif GTGGGAA (Tun et al., 1994; Brou et

al., 1994; Christensen et al., 1996). (E) Venn diagram showing the overlapping genes

identified through FLAG antibody and GFP antibody ChIP-seq analysis. Both data lists

were filtered for more than 2-fold change of signal (ChIP/control) with a moderate False

Discovery Rate (FDR<0.05). (F & G) Protein coding vs. non-coding distribution (F) and

chromosome position (G) of 75 genes from E. The overly represented motif discovered

by HOMER suite with the ChIP-seq data (top) and the canonical LAG-1/CSL binding

sequence (Brou et al., 1994, Tun et al., 1994 and Christensen et al., 1996)(bottom).

1423

Supplemental figure 5. Supplemental information for genome-wide identification

of germline LAG-1 targets. (A) The structure of *lag-1(ozIs43)[lag-1p::lag-*

1::3xFLAG::biotag::lag-1 3'UTR] fosmid transgene. 3xFLAG and Biotag sequence were

inserted into a fosmid that contains native regulatory sequence for *lag-1* gene, *lag-*

1(ozIs43) was able to rescue *lag-1* null allele *tm3052*. (B) Genome browser tracks

showing *sygl-1* and other four putative germline GLP-1/LAG-1 transcriptional targets

from literature: 10 kb genomic region for *sygl-1* (this study), *fbf-2 (Lamont et al., 2004)*

and lip-1 (Lee et al., 2006), 20 kb genomic region for *utx-1 (Seelk et a., 2016)* and *glp-1*

(Christensen et al., 1996). Raw reads were normalized to control, and the signal

intensity were presented as log2 fold change. Black arrow heads, canonical LAG-1/CSL

binding motif GTGGGAA (Tun et al., 1994; Brou et al., 1994; Christensen et al., 1996).

Red arrow heads, LAG-1 binding site (LBS) from original references (see B) where

LAG-1 was suggested to bind. (C & D) Protein coding/ non-coding distribution (C) and

chromosome position (D) of 137 genes from Fig 4D. (E) Venn diagram showing the

overlapping genes from germline and whole worm ChIP-seq analysis of LAG-1.

1439

Supplemental figure 6. Supplemental information for transcriptomic analysis to

identify GLP-1-dependent genes. (A) *In situ* hybridization used to determine *lst-1*

mRNA expression in young adult animals. The genotype are, GLP-1 ON: *gld-2(q497)*

gld-1(q485) glp-1(ar202) and GLP-1 OFF: *gld-2(q497) gld-1(q485) glp-1(q175).* Dotted

lines showing the boundary of the germline. Black arrow indicate distal gonadal end. (B

& C) Heatmap (B) and Principle Component Analysis (PCA) (C) were generated using

top 500 genes with most significant p-value from differential gene expression analysis.

1447

Supplemental figure 7. FBF-2 protein accumulation analysis. (A) Plot of FBF-2

levels for indicated genotype. *fbf-2(q932)* is used for quantification, see materials and

methods for alleles used for other genotype. Numbers in bracket shows the sample

size. Dots, mean; Error bars, mean ± SD. (B) Graph showing FBF-2 base levels (see

Fig 6B) in early adult hermaphrodites of indicated genotype. Numbers indicate mean

values for each genotype and numbers in bracket shows the sample size. Dots, mean;

Error bars, mean ± SD. P-value ≤ 0.01 (*); ≤ 0.001 (**); ≤ 0.0001 (***); > 0.01 non-

significant (NS.).

1456

Supplemental figure 8. Time-course transcriptomic analysis upon LAG-1

degradation in germline. (A & B) Heatmap (A) and Principle Component Analysis

(PCA) (B) for top 500 genes with most significant p-values, the differential gene

expression analysis was done between animals treated with or without auxin for 48

hours. (C & D) The differentially-expressed genes upon auxin treatment for 48 hours

were compared to genes affected by GLP-1 to discover: the overlapping genes

activated by both LAG-1 and GLP-1 (C) or those genes repressed by both LAG-1 and

GLP-1 (D). (E & F) The differentially-expressed genes upon auxin treatment for 48

hours were compared to putative LAG-1 targets through LAG-1 germline ChIP-seq

analysis to determine: LAG-1 transcriptional targets (E) and if LAG-1 can repress gene

expression (F). (G) Multiple dimension scaling analysis showing the similarities of the

RNA-seq samples conducted in this study. Five biological replicates each were

conducted for GLP-1 ON (in blues circle) and GLP-1 OFF (in pink circle). Four biological

replicates were conducted for time course RNA-seq analysis following LAG-1

degradation by auxin treatment. The 48-hour auxin treated samples were grouped in

yellow circle. With shorter period of treatment, i.e, 2 hours and 4 hours, the

transcriptomic profile are not very distinct from un-treated samples and were grouped in

the black circle.

1475

Supplemental figure 9. Time-course auxin treatment to degrade LAG-1. (A & B)

Representative images of anti-HA-stained (LAG-1, yellow, left panels) and WAPL-1-

stained (pink, right panels) after L4 stage animals were treated with or without auxin for

2 hours (A) or 4 hours (B) at 25°C. Asterisks, distal end of germline. Red arrow heads,

LAG-1 expression in the sheath cells. (C & D) Graph showing distance, in cell

diameters, between the distal end of the germline and the row of cells at proximal end of

the continuous zone of WAPL-1 staining for auxin-treated hermaphrodites. Data are

plotted as horizontal dot plots with each dot representing length in cell diameter to zone

end for one gonad. Numbers in bracket shows the sample size. Thick vertical lines

represent mean and horizontal lines represent mean ± SD. P-value ≤ 0.01 (*); ≤ 0.001

(**); ≤ 0.0001 (***); > 0.01 non-significant (NS.).

1487

Supplemental figure 10. SYGL-1 and LST-1 protein expression analysis following

2-hour and 4-hour auxin treatment to degrade LAG-1 protein. (A & B)

Representative images of SYGL-1 (A) and LST-1 (B) protein expression after animals

were treated with or without auxin for 4 hours. The genotype for auxin treatment are

sygl-1(q983)[3xOLLAS::sygl-1]; lag-1(oz536oz537)[lag-1::degron::3xHA]; ieSi64[gld-

1p::TIR1::mRuby::gld-1 3'UTR] (for quantification of SYGL-1 expression) and *lst-*

1(q1003)[lst-1::3xOLLAS]; lag-1(oz536oz537)[lag-1::degron::3xHA]; ieSi64[gld-

*1p::TIR1::mRuby::gld-1 3'UTR] (*for quantification of LST-1 expression*).* (C & D) Plots of

SYGL-1 (C) and LST-1 (D) levels in auxin-treated germlines. Numbers in bracket shows

the sample size. Dots, mean; Error bars, mean ± SD. (E & F) Comparison of SYGL-1

(E) and LST-1 (F) peak levels (see C & D) in auxin-treated germlines. Numbers indicate

1499 mean values for each genotype and numbers in bracket shows the sample size. Error

1500 bars, mean ± SD. P-value ≤ 0.01 (*); ≤ 0.001 (**); ≤ 0.0001 (***); > 0.01 non-significant

1501 (NS.). Numbers indicate mean values for each genotype.

1502

1503 **S1 Table.** Strains used in this study.

1504 **S2 Table.** Oligonucleotides used in this study.

1505 **S3 Table.** List of putative LAG-1 peak regions identified through whole worm ChIP-seq analysis.

1506 **S4 Table.** List of putative LAG-1 peak region identified through germline ChIP-seq analysis.

1507 **S5 Table.** List of genes activated or repressed by GLP-1.

1508 **S6 Table.** List of genes activated or repressed by LAG-1, after 48 hours of auxin treatment.

1509 **S7 Table.** List of genes activated or repressed by LAG-1, after 2 and 4 hours of auxin treatment.

1510 **S8 Table.** List of putative LAG-1 targets from literature.

Figure 1

A

Progenitor Zone (PZ)

Leptotene

DTC

GLP-1 transcription

LST-1

SYGL-1

Meiotic Entry

B

LAG-2
APX-1 → GLP-1 → GLP-1(ICD)
LAG-1

lst-1

sygl-1

?

GLD-1

GLD-2

SCF PROM-1

Meiotic
Development

Stem
Cell Fate

niche

germ cells

Figure 2

Figure 3

Figure 4

A

ckSi11[pie-1p::birA::gfp]

LAG-1 Biotag BirA biotin

Sequence specific biotinylation of LAG-1

Biotinylated LAG-1 in germline

ozIs43[lag-1p::lag-1::3xflag::biotag]

B Sequential ChIP-seq workflow:

1. ChIP with FLAG antibody

wash | elute

2. Re-ChIP with streptavidin beads

wash

3. DNA libraries prepared from DNA on the beads

C

10 kb

Rep-1
Rep-2
Rep-3

sygl-1

Rep-1
Rep-2
Rep-3

lst-1

Rep-1
Rep-2
Rep-3

mir-61/250

D

Rep-3
71

35   31

Rep-1   **137**   Rep-2
311   395

E

ChIP-seq motif   p-value

GCGTGGGAAA   1^{e-34}

Canonical LAG-1/CSL binding site

GTGGGAA

Figure 5

A

GLP-1 ON

GLP-1 OFF

qPCR or
RNA-seq

B

GLP-1 ON

GLP-1 OFF

C

Relative mRNA level

**

GLP-1 ON
GLP-1 OFF

*

sygl-1   *lst-1*

D

GLP-1
dependent
genes

Germline LAG-1
ChIP-seq peak
annotated genes

lst-1
sygl-1

Figure 6

Figure 7

Supplemental Figure 1 (S1 Fig)

A

Genotype	Dead eggs (%)	L1 arrest (%)	n
WT	0.5	0.0	803
LAG-1::3xHA	0.5	0.0	2139
LAG-1::degron: 3xHA	0.5	0.1	1953

B

C

Auxin (Hour)	- Auxin (4 h)	+ Auxin (4 h)	- Auxin (24 h)	+ Auxin (24 h)
WAPL-1 CD	21.9 ± 1.6	21.5 ± 1.4	19.4 ± 2.5	0
Glp (%)	0	0	0	100
n	35	26	35	30

Supplemental Figure 2 (S2 Fig)

A

B

C

Supplemental Figure 3 (S3 Fig)

A

B

Supplemental Figure 4 (S4 Fig)

Supplemental Figure 5 (S5 Fig)

A

Fosmid transgene structure

lag-1p *lag-1* 3xFLAG Biotag 3'UTR

lag-1 (ozIs43)
[lag-1::3xFLAG::Biotag]

B

10 kb

Rep-1
Rep-2
Rep-3

sygl-1

Rep-1
Rep-2
Rep-3

fbf-2

Rep-1
Rep-2
Rep-3

lip-1

20 kb

Rep-1
Rep-2
Rep-3

utx-1

Rep-1
Rep-2
Rep-3

glp-1

C

Protein-coding_77%

non-coding 23%

D

Chr II 12.4%
Chr III 17.5%
Chr I 14.6%
Chr IV 16.1%
Chr X 14.6%
Chr V 24.8%

E

Germline LAG-1 ChIP-seq peak annotated genes 101

Whole worm LAG-1 ChIP-seq peak annotated genes 39

Supplemental Figure 6 (S6 Fig)

Supplemental Figure 7 (S7 Fig)

A

B Base levels

Supplemental Figure 8 (S8 Fig)

Supplemental Figure 9 (S9 Fig)

Supplemental Figure 10 (S10 Fig)