Hat1-Dependent Lysine Acetylation Targets Diverse Cellular Functions

Paula A. Agudelo Garcia and Mark R. Parthun

Department of Biological Chemistry and Pharmacology, The Ohio State University, Columbus, OH 43210

Corresponding author:
Mark R. Parthun, PhD.
206 Rightmire Hall
1060 Carmack Road
Columbus, OH 43210
Tel: 614-292-6215
Email: parthun.1@osu.edu

Running title: Identification of the mammalian Hat1-dependent acetylome

ABSTRACT

Lysine acetylation has emerged as one of the most important post-translational modifications, regulating different biological processes. However, its regulation by lysine acetyltransferases is still unclear in most cases. Hat1 is a lysine acetyltransferase originally identified based on its ability to acetylate histones. Using an unbiased proteomics approach, we have determined how loss of Hat1 affects the mammalian acetylome. $Hat1^{+/+}$ and $Hat1^{-/-}$ mouse embryonic fibroblast (MEF) cells lines were grown in both glucose- and galactose-containing media, as Hat1 is required for growth on galactose due to defects in mitochondrial function. Following trypsin digestion of whole cell extracts, acetylated peptides were enriched by acetyllysine affinity purification and acetylated peptides were identified and analyzed by label-free quantitation. Comparison of the acetylome from $Hat1^{+/+}$ cells grown on galactose and glucose demonstrated that there are large carbon source-dependent changes in the mammalian acetylome where the acetylation of enzymes involved in glycolysis was the most affected. Comparisons of the acetylomes from $Hat1^{+/+}$ and $Hat1^{-/-}$ cells identified 65 proteins whose acetylation decreased by at least 2.5-fold in cells lacking Hat1. In $Hat1^{-/-}$ cells, acetylation of the auto regulatory loop of CBP was the most highly affected, decreasing by up to 20-fold. In addition to proteins involved in chromatin structure, Hat1-dependent acetylation was also found in a number of transcriptional regulators, including p53, and mitochondrial proteins. Hat1 mitochondrial localization suggests that it may be directly involved in the acetylation of mitochondrial proteins.

INTRODUCTION

The acetylation of the ε-amino group of lysine residues in proteins is a widely

utilized post-translational modification that regulates a variety of cellular

processes, including chromatin structure, transcriptional regulation, DNA damage

repair, cellular metabolism, cytoskeletal dynamics and apoptosis(1,2). The critical

role played by protein acetylation is reflected in the number of human disease

states, such as cancers and neurological disorders, which result from alterations

in this modification(3). Proteomic studies have identified several thousand

acetylated proteins in mammalian cells (known as the acetylome) (4-7). Given its

abundance and relevance, acetylation may rival phosphorylation in importance

as a regulatory and signaling modification. An intriguing difference between

acetylation and phosphorylation regards the repertoire of enzymes responsible

for these modifications. While there are more than 500 kinases encoded in

mammalian genomes, there are fewer than 20 lysine acetyltransferases(1).

Hence, it is likely that each known protein acetyltransferase will have multiple

substrates and will be involved in multiple cellular processes(8,9).

Histone acetyltransferase 1 (Hat1) was the first protein ε-amino group lysine

acetyltransferase to be identified and is the most widely conserved among

eukaryotes (10-12). Like most lysine acetyltransferases, Hat1 was identified

based on its ability to acetylate histones. Hat1 was originally purified from yeast

cytoplasmic extracts in a complex with Hat2 (Rbap46/Rbbp7 in mammalian

cells). Biochemically, the Hat1 complex acetylates histone H4 lysine residues 5

and 12 and only acetylates histones free in solution (the Hat1 reaction is blocked

by the presence of nucleosomal DNA)(11,13). This observation suggested that Hat1 was responsible for the evolutionarily conserved diacetylation of newly synthesized histone H4 (on lysines 5 and 12) that is known to accompany the process of replication-coupled chromatin assembly, a hypothesis that was subsequently confirmed(14).

In vivo studies of Hat1 have identified several phenotypes associated with loss of this enzyme. These phenotypes, which include defects in heterochromatin gene silencing, DNA damage sensitivity, genome instability and alterations in gene expression, are likely due to the histone modifying function of Hat1. However, recent studies have identified mitochondrial defects associated with Hat1 loss, suggesting that this enzyme may influence key cellular pathways through the acetylation of non-histone substrates. To determine whether Hat1 influences non-histone protein acetylation, we have used a proteomic strategy to identify Hat1-dependent sites of acetylation. The abundance of acetylated peptides was compared between Hat1$^{+/+}$ and Hat1$^{-/-}$ mouse embryonic fibroblasts (MEFs) grown on both glucose and galactose. As previously reported in liver from mice fed a high fat died or starved; these experiments indicated that the shift in carbon source from glucose to galactose had a dramatic effect on the acetylome(15,16). In particular, there were changes in the acetylation of many of the enzymes required for glycolysis. In addition, loss of Hat1 resulted in a significant decrease in the acetylation of >70 proteins. These proteins are involved in a wide range of cellular functions, including chromatin structure, transcriptional regulation and mitochondrial function. Interestingly, acetylation levels of the protein

acetyltransferase CBP were the most highly affected in Hat1$^{-/-}$ MEFs suggesting that Hat1 may be a regulator of CBP function. In addition, we demonstrate that Hat1 exhibits mitochondrial localization consistent with the possibility that this enzyme may be directly involved in mitochondrial protein acetylation.

MATERIALS AND METHODS

MEF cell culture. E14.5 embryos were dissected from uterus; head and internal organs were removed. Tissue was disaggregated using an 18-gauge syringe and brought to single cell suspension with trypsin incubation at 37°C. Cells were then plated onto 100 mm tissue culture plates, passaged upon confluency and maintained in Dulbecco's modified Eagle medium (DMEM-Sigma) with 10% fetal bovine serum (FBS-Gibco) and 1X Pen/Strep antibiotics (Sigma). SV40 T immortalized MEFs (iMEFs) were derived from primary Hat1$^{+/+}$ and Hat1$^{-/-}$ embryonic day 13.5 embryos. To establish iMEFs, early passage cells were transformed with SV-40 T antigen containing plasmid pBSSVD2005 (ADDGENE, Cambridge, MA). Early passage cells were seeded at 25% confluency in 6 well plates and transfected with 2 ug of expression vector using Fugene reagent (Roche). Cells were harvested and seeded into 100 mm dishes after 48 hrs of transfection. The cells were split at 1 in 10 dilutions until passage 5.

Mitochondrial Isolation. Cells were resuspended in mitochondrial isolation buffer (0.8M sucrose, 20mM tris-HCl ph7.2, 40mM KCl, 2mM EGTA, 1mg/mL BSA, 0.2mM PMSF), kept on ice for 2 min and lysed with Dounce homogenizer. Homogenate was centrifuged at 1500g for 10min at 4°C. Supernatant was further

centrifuged at 17.000g for 30min at 4°C. Pellet was resuspended in mitochondrial extraction buffer without BSA.

Subcellular Fractionation. Protein fractions from nucleus, cytosol and mitochondria were obtained according to Dimauro et al.(17). Cells were resuspended in STM buffer (250mM sucrose, 50mM Tris-HCl pH 7.5, 5mM MgCl2) and homogenized at low speed for 1 min, vortexed and kept on ice for 30min. Samples were then centrifuged at 800g for 15min, supernatant and pellet were separated; pellet was washed twice with STM buffer and resuspended in NET buffer (20mM hepes pH7.9, 1.5mM MgCL2, 0.5NaCl, 0.2mM EDTA, 20% glycerol and 1%triton x-100) this was saved as the nuclear fraction. Supernatant was centrifuged one more time at 11.000g for 10min. Supernatant was treated with 100% acetone and centrifuged at 12.000g for 5min to obtain the cytosolic fraction. Pellet containing the mitochondrial fraction was washed twice with STM buffer and resuspended in SOL buffer (50mM tris-HCl pH 6.8, 1mM EDTA, 0.5% Triton X-100).

Immunofluorescence. Hek293 cells were seeded on 6 well plates containing coverslips. Cells were fixed with 4% paraformaldehyde, blocked with 5% BSA and primary antibodies Hat1 (Abcam) was incubated overnight, washed and replaced with goat-anti-rabbit secondary antibody. Cells were mounted using Vectashield mounting media containing DAPI and imaged using a wide field fluorescence microscope. Mitolight was used according to manufacturer's instructions.

PTM scan lysine acetylation. 2x10^8 cells for each experimental condition were collected. Cells were washed with PBS and resuspended in 10mL of urea lysis buffer. Cell lysates were frozen in dry ice/ethanol for at least 30 minutes, or longer if necessary, for cell lysate to freeze completely. The acetylome of each sample was analyzed by PTMScan (cell signaling) using their standard procedure.

RESULTS

Hat1 plays a role in multiple cellular processes through the acetylation of histone H4 and the regulation of chromatin structure(18-20).   In addition, it has been reported that Hat1 is required for mitochondrial function(21,22). Given the large number of orphan acetylated proteins observed in mitochondria and the fact that Hat1 acetylates histone H4 in the cytosol, we decided to identify potential non-histone substrates of Hat1 and its additional functions. To do so, we took a proteomic approach to determine how loss of Hat1 influenced the mammalian acetylome.

The mammalian acetylome is altered by a shift from glucose to galactose.
Since it has been previously reported that metabolic stress can induce changes in acetylation and consistent with a role for Hat1 in mitochondrial function, we used two different growth conditions: glucose and galactose, to characterize the cellular acetylome from immortalized Hat1$^{+/+}$ and Hat1$^{-/-}$ MEFs(15,16,21).

Immortalized Hat1$^{+/+}$ and Hat1$^{-/-}$ MEFs were grown in media containing either glucose or galactose as the primary carbon source. Whole cell extracts were prepared, and proteins were digested with trypsin. Acetylated peptides were

enriched by affinity purification using an anti-acetyl lysine affinity resin. The resulting peptides were analyzed by LC-MS/MS and the abundance of the acetylated peptides in each sample was determined using label free quantitation.

We first compared Hat1$^{+/+}$ cells grown in glucose and galactose to determine the effect of the carbon source on the mammalian acetylome. 470 peptides, derived from 364 proteins were detected, whose level of acetylation changed by +/- 2.5-fold in cells grown in galactose relative to glucose (Figure 1A). Acetylation levels increased on 173 of these peptides (37%) and decreased on 297 (63%) (Supporting Table 1). The list of proteins whose acetylation changes by +/- 2.5-fold was analyzed by Ingenuity Pathway Analysis (Figure 1B). Strikingly, the most significantly enriched pathway was glycolysis. Gluconeogenesis was also highly enriched as it shares many of the same enzymes. The acetylation state of the enzymes needed to convert galactose to glucose-6-phosphate was not altered. Of the 10 enzymes required to convert glucose to pyruvate, 7 showed a significant change in acetylation. It has been shown that most of the enzymes involved in glycolysis, as well as enzymes in the majority of metabolic pathways, are acetylated(2,23). In fact, some of these sites of acetylation are known to regulate enzymatic activity. For example, acetylation of lysine 305 of pyruvate kinase promotes the enzyme's degradation(24). However, none of the sites of acetylation that were found to change in response to growth on galactose have been the subject of functional characterization.

We also analyzed the acetylomes between Hat1$^{-/-}$ MEFs grown in glucose and galactose. Table 1 lists the 50 proteins that had the largest increases and

decreases in acetylation in Hat1$^{+/+}$ MEFs and compares them to the changes observed in Hat1$^{-/-}$ counterparts under the same growth conditions. It is apparent that loss of Hat1 had very little impact on the acetylation status of proteins whose acetylation decreased when grown on galactose. However, there were marked changes in the proteins whose acetylation increased in wild type cells grown in galactose. For 40% of these proteins, the increased acetylation seen when the cells were switched to galactose decreased by at least 5-fold in the Hat1$^{-/-}$ cells. For example, PDZK1 showed a 108-fold increase in acetylation when Hat1$^{+/+}$ MEFs were grown on galactose compared to only a 2.9-fold increase in Hat1$^{-/-}$ MEFs. This set of proteins was not enriched for any specific cellular pathways but it includes 2 proteins that are critical for mitochondrial function, AGK and ATP6V1E1. Interestingly, Agk$^{-/-}$ cells have been reported to have very similar phenotypes to Hat1$^{-/-}$ cells, including poor growth on galactose media, defective mitochondrial structure and poor mitochondrial metabolism(25). One possibility is that Hat1 mitochondrial function is executed through the acetylation of Agk regulating its activity.

Identification of Hat1-dependent sites of acetylation. Hat1-dependent sites of acetylation were identified by comparing the acetylome of Hat1$^{+/+}$ MEFs grown on glucose to that of Hat1$^{-/-}$ MEFS grown on glucose and the acetylome of Hat1$^{+/+}$ MEFs grown on galactose to that of Hat1$^{-/-}$ MEFs grown on galactose. This analysis identified 84 sites in 65 proteins whose acetylation decreased by at least 2.5-fold in Hat1$^{-/-}$ cells (Table 2). Of these 84 sites of acetylation, 29 have not been previously reported (~35%, Table 2). It was previously proposed that

Hat1 acetylates lysine residues in the context of the sequence GXG**K**XG based on a comparison of the ability of Hat1 to acetylate histone H4 and H2A from a variety of eukaryotes(11). The sequence context of the Hat1-dependent sites of acetylation was analyzed to determine if there was an associated motif. Figure 2A shows the motifs identified for the total pool of acetylated peptides analyzed and for those peptides that changed by more than 2.5-fold in the Hat1$^{-/-}$ samples. It is clear that glycine residues were enriched at the -3, -1 and +1 positions, as originally predicted. This motif also indicated that there was a strong preference for alanine at the +5 position and for glycine at the -7 position in the peptides whose acetylation was Hat1-dependent.

The list of proteins whose acetylation is Hat1-dependent was subjected to Gene Ontology (GO) analysis. Hat1-dependent protein acetylation was observed on proteins involved in a variety of cellular pathways, including ones not directly related to chromatin transactions (Figure 2B). These pathways included transcriptional regulation, translation, mitochondria and RNA processing.

Hat1 and the acetylation of chromatin related proteins. As expected, the most highly represented group of proteins whose acetylation was Hat1-dependent was those involved in chromatin and DNA binding (Figure 3A). Hat1-dependent acetylation was observed for all of the core histones, as well as for histone H1.4. These sites of acetylation included histone H2A lysine 5 (and the corresponding residues in histone H2A.Z and H2A.J) that was previously identified as a Hat1 substrate(11,26). Histone H4 lysine residues 5 and 12, the most well characterized Hat1 substrates, were not identified in these data sets.

This is likely due to the abundance of trypsin cleavage sites in the NH_2-terminal tail of H4 that results in the generation of fragments containing these residues that are too small to be reliably observed.

Interestingly, the acetylation of two histone acetyltransferases (CBP and Hbo1) was Hat1-dependent (Table 2). The protein whose acetylation was most highly dependent on Hat1 was CBP, decreasing up to 20-fold in Hat1[-/-] cells (Figure 3A). Importantly, the Hat1-dependent sites of acetylation on CBP are all located in the auto-regulatory loop (AL) in the CBP HAT domain (Figure 3B). The acetylation state of the AL of CBP is critical for its acetyltransferase activity(27). The dependence of the CBP AL on Hat1 suggests that Hat1 may influence the acetylation of some proteins indirectly through the activation of CBP. In fact, comparison of the Hat1-dependent sites of acetylation with an acetylomics analysis of CBP[-/-] MEFs shows that approximately 45% of these sites are CBP substrates (Table 2)(8). The potential regulation of CBP by Hat1 is particularly intriguing given the role of CBP in the acetylation of newly synthesized histone H3(28). Several reports have suggested that Hat1 is involved in the transcriptional regulation of several genes in different models(29-33). However, this evidence is conflicted by Hat1's incapacity to acetylate histones in a nucleosome context. One scenario is that Hat1 influences gene expression indirectly through the acetylation of CBP regulating the Hat activity of this transcription factor.

Hat1 dependent protein acetylation of transcriptional regulators.

Transcriptional regulatory proteins were also highly represented in the Hat1

acetylome (Figure 4A). An antibody specific for this site of p53 acetylation is available allowing us to validate the results of the proteomic analysis. We found that mouse p53 lysine 381 acetylation is induced by growth of MEFs in galactose (Supporting Table 1). Figure 4B shows immunofluorescence images of Hat1$^{+/+}$ and Hat1$^{-/-}$ MEFs grown in galactose. Staining for total p53 indicated that Hat1 loss did not affect p53 protein levels. However, there was a significant decrease of p53 lysine 381 acetylation in the Hat1$^{-/-}$ MEFs, confirming that Hat1 regulates p53 acetylation.

Hat1 and the acetylation of mitochondrial proteins. Proteomic approaches have identified a large number of mitochondrial proteins as being acetylated. Mitochondrial protein acetylation increases in mouse liver during fasting, caloric restriction or a long-term high fat diet (12, 75, 76), indicating that the acetylation patterns in this organelle can respond to dietary changes. Interestingly, mitochondrial proteins are also enriched in the Hat1-dependent acetylome (Figure 5A). Recent reports have clearly identified defects in mitochondrial function in the absence of Hat1(21,22,34). The observation that the acetylation of a subset of mitochondrial proteins is Hat1-dependent suggests the possibility that Hat1 may influence mitochondrial function through mitochondrial protein acetylation.

Even though several mitochondrial proteins are acetylated, there is a debate as to whether mitochondrial protein acetylation results from chemical or enzymatic reactions(35-39). To better understand if Hat1 directly acetylates mitochondrial proteins, we first examined whether Hat1 localizes in this compartment. We took

two biochemical approaches, performing sub-cellular fractionation and mitochondrial purification (Figure 5B). In both cases, Hat1 was detected in the mitochondrial compartment. To further confirm this result, we used confocal microscopy to visualize Hat1 and the mitochondrial marker MitoLight. In addition to nuclear and cytoplasmic localization, Hat1 also co-localized with MitoLight in confocal sections (Figure 5C, yellow staining), suggesting that Hat1 may function as a mitochondrial lysine acetyltransferase.

DISCUSSION

Hat1 was the first lysine acetyltransferase identified but its confirmed substrates are limited to histones. We have used an unbiased proteomics approach to determine how loss of Hat1 influences the mammalian acetylome. This analysis identified 65 proteins whose acetylation is dependent on Hat1. Detailed biochemical experiments will need to be performed to determine whether these proteins are direct substrates of Hat1 or whether Hat1 regulates their acetylation through the regulation of the enzymatic activity of other HATs like CBP. The list of targets that we have identified is likely to be an underestimate of the influence of Hat1 on the mammalian acetylome. In our study, trypsin was used to digest proteins. Trypsin cleaves proteins after lysine or arginine but its activity is blocked by acetyllysine. Hence, loss of Hat1-dependent acetyllysine residues could actually result in an increase in the abundance of some peptides in our study. In addition, trypsin digestion of proteins with high lysine and arginine content can result in peptides that are too small for accurate identification. This

is the case with some histones and is likely to be the reason why the validated Hat1 targets, H4 lysine residues 5 and 12, were not detected.

Hat1 was originally isolated from yeast based on its ability to acetylate histone H4 on lysine residues 5 and 12(11). Interestingly, the sequence contexts of H4 lysines 5 and 12 are very similar (GRGK$_5$GG and GLGK$_{12}$GG). As the NH$_2$-tail of histone H4 is nearly invariant throughout evolution, Hat1 from yeast was capable of acetylating histone H4 from a variety of eukaryotes. Yeast Hat1 was also found to acetylate lysine residue 5 of histone H2A but only from specific species (11,13,26). Examination of the sequence context of histone H2A lysine 5 from a variety of species indicated that Hat1 only acetylated this residue when it was in the context GXG**K**XG, which was hypothesized to be a recognition site for Hat1(11). The subsequent crystal structure of the yeast enzyme demonstrated that there was a structural basis for this recognition site(40). Analysis of the peptides that showed Hat1-dependent acetylation in the current study supports the importance of the glycine residues at positions -3. -1 and +2 but also suggests that these residues are not strictly required. In addition, the Hat1-dependent peptides show a marked preference for an alanine at the +5 position and a glycine at the -7 position. Interestingly, neither of these residues is present at lysines 5 and 12 of H4. However, one or both of the -7 glycine or the +5 alanine are found in canonical mammalian histone H2A and the H2A variants H2A.Z and H2A.J, all of which have decreased acetylation in the absence of Hat1 in our dataset.

A significant fraction of the Hat1-dependent acetylome may be sites that are indirectly regulated by Hat1 through the HATs CBP and Hbo1. Indeed, nearly half of the sites of acetylation that decrease by at least 2.5-fold in Hat1$^{-/-}$ cells were also decreased in CBP$^{-/-}$ cells. There are both direct and indirect mechanisms by which Hat1 may regulate acetylation of the AL of CBP. Hat1 may directly acetylate the lysine residues in the AL of CBP. Alternatively, Hat1 could directly interact with CBP and alter CBP structure or activity to facilitate the autoacetylation of the AL by CBP(27). Hat1 could also interact with other factors that regulate the autoacetylation of the AL, such as RNAs that bind CBP and activate its acetyltransferase activity(41).

Lysine 381 acetylation on p53 (mouse) is both Hat1- and CBP-dependent. Interestingly, Hat1 and p53 share some phenotypes. It was recently shown that mice that are heterozygous for Hat1 (Hat1$^{+/-}$) display premature aging and a significantly shortened lifespan(21). Mice heterozygous for p53 have a similarly shortened lifespan. In addition, heterozygosity of p53 rescues the extreme lifespan decrease of Sirt6$^{-/-}$ mice(42). It will be interesting to determine whether the role of Hat1 in mammalian aging is linked to the acetylation of p53.

While many mitochondrial proteins are acetylated, there is debate as to whether this acetylation is the result of non-specific chemical reactions or is enzyme-catalyzed(35-39). The high levels of acetyl-CoA in the mitochondria certainly suggests that chemical acetylation of mitochondrial is likely to occur. However, there are a number of reports demonstrating the localization of other lysine acetyltransferases to mitochondria(36-38,43). Our results, combined with the

recent demonstration that Hat1 is required for proper mitochondrial function, raise the possibility that Hat1 may also be involved in mitochondrial protein acetylation(21). Hat1 doesn't contain a mitochondrial localization signal, but its translocation to this organelle could occur through the interaction with mitochondrial proteins or mitochondrial transporters. Future studies will focus on determining whether the Hat1-dependent sites of acetylation on mitochondrial proteins are a result of the direct action of Hat1 in the mitochondria. Together the data provided in our study shows that Hat1 influences a wide range of biological process and that its function extends beyond the nuclear compartment.

ACKNOWLEDGEMENTS

This work was support by a grant form the National Institutes of Health (R01 GM062970).

CONFLICT OF INTEREST

The authors declare that they have no conflict of interest.

AUTHOR CONTRIBUTIONS

P.A.A.G. designed experiments, performed experiments, analyzed data and wrote the manuscript. M.R.P. designed experiments, analyzed data and wrote the manuscript.

REFERENCES

1.	Drazic, A., Myklebust, L. M., Ree, R., and Arnesen, T. (2016) The world of protein acetylation. *Biochim Biophys Acta* **1864**, 1372-1401
2.	Narita, T., Weinert, B. T., and Choudhary, C. (2019) Functions and mechanisms of non-histone protein acetylation. *Nat Rev Mol Cell Biol* **20**, 156-174
3.	Arif, M., Senapati, P., Shandilya, J., and Kundu, T. K. (2010) Protein lysine acetylation in cellular function and its role in cancer manifestation. *Biochim Biophys Acta* **1799**, 702-716
4.	Kim, G. W., and Yang, X. J. (2010) Comprehensive lysine acetylomes emerging from bacteria to humans. *Trends Biochem Sci* **36**, 211-220
5.	Kim, S. C., Sprung, R., Chen, Y., Xu, Y., Ball, H., Pei, J., Cheng, T., Kho, Y., Xiao, H., Xiao, L., Grishin, N. V., White, M., Yang, X. J., and Zhao, Y. (2006) Substrate and functional diversity of lysine acetylation revealed by a proteomics survey. *Mol Cell* **23**, 607-618
6.	Zhao, S., Xu, W., Jiang, W., Yu, W., Lin, Y., Zhang, T., Yao, J., Zhou, L., Zeng, Y., Li, H., Li, Y., Shi, J., An, W., Hancock, S. M., He, F., Qin, L., Chin, J., Yang, P., Chen, X., Lei, Q., Xiong, Y., and Guan, K. L. (2010) Regulation of cellular metabolism by protein lysine acetylation. *Science* **327**, 1000-1004
7.	Ke, M., Shen, H., Wang, L., Luo, S., Lin, L., Yang, J., and Tian, R. (2016) Identification, Quantification, and Site Localization of Protein Posttranslational Modifications via Mass Spectrometry-Based Proteomics. *Adv Exp Med Biol* **919**, 345-382
8.	Weinert, B. T., Narita, T., Satpathy, S., Srinivasan, B., Hansen, B. K., Scholz, C., Hamilton, W. B., Zucconi, B. E., Wang, W. W., Liu, W. R., Brickman, J. M., Kesicki, E. A., Lai, A., Bromberg, K. D., Cole, P. A., and Choudhary, C. (2018) Time-Resolved Analysis Reveals Rapid Dynamics and Broad Scope of the CBP/p300 Acetylome. *Cell* **174**, 231-244 e212
9.	Fournier, M., Orpinell, M., Grauffel, C., Scheer, E., Garnier, J. M., Ye, T., Chavant, V., Joint, M., Esashi, F., Dejaegere, A., Gonczy, P., and Tora, L. (2016) KAT2A/KAT2B-targeted acetylome reveals a role for PLK4 acetylation in preventing centrosome amplification. *Nat Commun* **7**, 13227
10.	Kleff, S., Andrulis, E. D., Anderson, C. W., and Sternglanz, R. (1995) Identification of a gene encoding a yeast histone H4 acetyltransferase. *J Biol Chem* **270**, 24674-24677
11.	Parthun, M. R., Widom, J., and Gottschling, D. E. (1996) The major cytoplasmic histone acetyltransferase in yeast: links to chromatin replication and histone metabolism. *Cell* **87**, 85-94
12.	Deng, W., Wang, C., Zhang, Y., Xu, Y., Zhang, S., Liu, Z., and Xue, Y. (2016) GPS-PAIL: prediction of lysine acetyltransferase-specific modification sites from protein sequences. *Sci Rep* **6**, 39787
13.	Verreault, A., Kaufman, P. D., Kobayashi, R., and Stillman, B. (1998) Nucleosomal DNA regulates the core-histone-binding subunit of the human Hat1 acetyltransferase. *Curr Biol* **8**, 96-108
14.	Nagarajan, P., Ge, Z., Sirbu, B., Doughty, C., Agudelo Garcia, P. A., Schlederer, M., Annunziato, A. T., Cortez, D., Kenner, L., and Parthun, M. R. (2013) Histone acetyl transferase 1 is essential for mammalian development, genome

stability, and the processing of newly synthesized histones H3 and H4. *PLoS Genet* **9**, e1003518

15.   Hebert, A. S., Dittenhafer-Reed, K. E., Yu, W., Bailey, D. J., Selen, E. S., Boersma, M. D., Carson, J. J., Tonelli, M., Balloon, A. J., Higbee, A. J., Westphall, M. S., Pagliarini, D. J., Prolla, T. A., Assadi-Porter, F., Roy, S., Denu, J. M., and Coon, J. J. (2013) Calorie restriction and SIRT3 trigger global reprogramming of the mitochondrial protein acetylome. *Mol Cell* **49**, 186-199

16.   Yang, L., Vaitheesvaran, B., Hartil, K., Robinson, A. J., Hoopmann, M. R., Eng, J. K., Kurland, I. J., and Bruce, J. E. (2011) The fasted/fed mouse metabolic acetylome: N6-acetylation differences suggest acetylation coordinates organ-specific fuel switching. *J Proteome Res* **10**, 4134-4149

17.   Dimauro, I., Pearson, T., Caporossi, D., and Jackson, M. J. (2012) A simple protocol for the subcellular fractionation of skeletal muscle cells and tissue. *BMC Res Notes* **5**, 513

18.   Agudelo Garcia, P. A., Hoover, M. E., Zhang, P., Nagarajan, P., Freitas, M. A., and Parthun, M. R. (2017) Identification of multiple roles for histone acetyltransferase 1 in replication-coupled chromatin assembly. *Nucleic Acids Res* **45**, 9319-9335

19.   Parthun, M. R. (2007) Hat1: the emerging cellular roles of a type B histone acetyltransferase. *Oncogene* **26**, 5319-5328

20.   Parthun, M. R. (2012) Histone acetyltransferase 1: More than just an enzyme? *Biochim Biophys Acta*

21.   Nagarajan, P., Agudelo Garcia, P. A., Iyer, C. C., Popova, L. V., Arnold, W. D., and Parthun, M. R. (2019) Early-onset aging and mitochondrial defects associated with loss of histone acetyltransferase 1 (Hat1). *Aging cell*, e12992

22.   Marin, T. L., Gongol, B., Zhang, F., Martin, M., Johnson, D. A., Xiao, H., Wang, Y., Subramaniam, S., Chien, S., and Shyy, J. Y. (2017) AMPK promotes mitochondrial biogenesis and function by phosphorylating the epigenetic factors DNMT1, RBBP7, and HAT1. *Sci Signal* **10**

23.   Guan, K. L., and Xiong, Y. (2011) Regulation of intermediary metabolism by protein acetylation. *Trends Biochem Sci* **36**, 108-116

24.   Xiong, Y., Lei, Q. Y., Zhao, S., and Guan, K. L. (2011) Regulation of glycolysis and gluconeogenesis by acetylation of PKM and PEPCK. *Cold Spring Harb Symp Quant Biol* **76**, 285-289

25.   Vukotic, M., Nolte, H., Konig, T., Saita, S., Ananjew, M., Kruger, M., Tatsuta, T., and Langer, T. (2017) Acylglycerol Kinase Mutated in Sengers Syndrome Is a Subunit of the TIM22 Protein Translocase in Mitochondria. *Mol Cell* **67**, 471-483 e477

26.   Tafrova, J. I., and Tafrov, S. T. (2014) Human histone acetyltransferase 1 (Hat1) acetylates lysine 5 of histone H2A in vivo. *Mol Cell Biochem* **392**, 259-272

27.   Park, S., Stanfield, R. L., Martinez-Yamout, M. A., Dyson, H. J., Wilson, I. A., and Wright, P. E. (2017) Role of the CBP catalytic core in intramolecular SUMOylation and control of histone H3 acetylation. *P Natl Acad Sci USA* **114**, E5335-E5342

28. Das, C., Roy, S., Namjoshi, S., Malarkey, C. S., Jones, D. N., Kutateladze, T. G., Churchill, M. E., and Tyler, J. K. (2014) Binding of the histone chaperone ASF1 to the CBP bromodomain promotes histone acetylation. *Proc Natl Acad Sci U S A* **111**, E1072-1081

29. Vlad, M. L., Manea, S. A., Lazar, A. G., Raicu, M., Muresian, H., Simionescu, M., and Manea, A. (2019) Histone Acetyltransferase-Dependent Pathways Mediate Upregulation of NADPH Oxidase 5 in Human Macrophages under Inflammatory Conditions: A Potential Mechanism of Reactive Oxygen Species Overproduction in Atherosclerosis. *Oxidative medicine and cellular longevity* **2019**, 3201062

30. Fan, P., Zhao, J., Meng, Z., Wu, H., Wang, B., Wu, H., and Jin, X. (2019) Overexpressed histone acetyltransferase 1 regulates cancer immunity by increasing programmed death-ligand 1 expression in pancreatic cancer. *J Exp Clin Cancer Res* **38**, 47

31. Gruber, J. J., Geller, B., Lipchik, A. M., Chen, J., Salahudeen, A. A., Ram, A. N., Ford, J. M., Kuo, C. J., and Snyder, M. P. (2019) HAT1 Coordinates Histone Production and Acetylation via H4 Promoter Binding. *Mol Cell* **75**, 711-724 e715

32. Han, N., Shi, L., Guo, Q., Sun, W., Yu, Y., Yang, L., Zhang, X., and Zhang, M. (2017) HAT1 induces lung cancer cell apoptosis via up regulating Fas. *Oncotarget* **8**, 89970-89977

33. Sadler, A. J., Suliman, B. A., Yu, L., Yuan, X., Wang, D., Irving, A. T., Sarvestani, S. T., Banerjee, A., Mansell, A. S., Liu, J. P., Gerondakis, S., Williams, B. R., and Xu, D. (2015) The acetyltransferase HAT1 moderates the NF-kappaB response by regulating the transcription factor PLZF. *Nat Commun* **6**, 6795

34. Miao, B. P., Zhang, R. S., Yang, G., Sun, J. J., Tang, Y. Y., Liang, W. F., Liu, T., Wen, Z., Yang, P. C., and Nie, G. H. (2018) Histone acetyltransferase 1 up regulates Bcl2L12 expression in nasopharyngeal cancer cells. *Arch Biochem Biophys* **646**, 72-79

35. Ghanta, S., Grossmann, R. E., and Brenner, C. (2013) Mitochondrial protein acetylation as a cell-intrinsic, evolutionary driver of fat storage: chemical and metabolic logic of acetyl-lysine modifications. *Crit Rev Biochem Mol Biol* **48**, 561-574

36. Di Domenico, A., Hofer, A., Tundo, F., and Wenz, T. (2014) Mitochondrial protein acetylation mediates nutrient sensing of mitochondrial protein synthesis and mitonuclear protein balance. *IUBMB Life* **66**, 793-802

37. Scott, I., Webster, B. R., Chan, C. K., Okonkwo, J. U., Han, K., and Sack, M. N. (2014) GCN5-like protein 1 (GCN5L1) controls mitochondrial content through coordinated regulation of mitochondrial biogenesis and mitophagy. *J Biol Chem* **289**, 2864-2872

38. Scott, I., Webster, B. R., Li, J. H., and Sack, M. N. (2012) Identification of a molecular component of the mitochondrial acetyltransferase programme: a novel role for GCN5L1. *Biochem J* **443**, 655-661

39. Webster, B. R., Scott, I., Han, K., Li, J. H., Lu, Z., Stevens, M. V., Malide, D., Chen, Y., Samsel, L., Connelly, P. S., Daniels, M. P., McCoy, J. P., Jr., Combs, C. A.,

Gucek, M., and Sack, M. N. (2013) Restricted mitochondrial protein acetylation initiates mitochondrial autophagy. *J Cell Sci* **126**, 4843-4849

40. Dutnall, R. N., Tafrov, S. T., Sternglanz, R., and Ramakrishnan, V. (1998) Structure of the histone acetyltransferase Hat1: a paradigm for the GCN5-related N-acetyltransferase superfamily. *Cell* **94**, 427-438

41. Bose, D. A., Donahue, G., Reinberg, D., Shiekhattar, R., Bonasio, R., and Berger, S. L. (2017) RNA Binding to CBP Stimulates Histone Acetylation and Transcription. *Cell* **168**, 135-149 e122

42. Ghosh, S., Wong, S. K., Jiang, Z., Liu, B., Wang, Y., Hao, Q., Gorbunova, V., Liu, X., and Zhou, Z. (2018) Haploinsufficiency of Trp53 dramatically extends the lifespan of Sirt6-deficient mice. *Elife* **7**

43. Stilger, K. L., and Sullivan, W. J., Jr. (2013) Elongator protein 3 (Elp3) lysine acetyltransferase is a tail-anchored mitochondrial protein in Toxoplasma gondii. *J Biol Chem* **288**, 25318-25329

FIGURE LEGENDS

Figure 1. The effect of carbon source on the mouse acetylome. A) Pie chart showing the number of sites of lysine acetylation that increase or decrease when mouse embryonic fibroblasts are switch from growth in glucose-containing media to galactose-containing media. B) Results Ingenuity Pathway Analysis of the proteins whose acetylation changed by at least 2.5-fold between growth on glucose and galactose. C) Schematic diagram of the pathway of glycolysis. Enzymes in green increased at least 2.5-fold when cells on galactose, enzymes in red decreased at least 2.5-fold on galactose and enzymes in black were unchanged.

Figure 2. Characterization of the Hat1-dependent acetylome. A) Top, motif derived from the bulk acetylated peptides. Bottom, Motif derived from comparison of pepetides whose acetylation is effected by Hat1 deletion. B) Proteins whose acetylation decreased at least 2.5-fold in Hat1 KO cells were analyzed by Gene Ontology. The number of proteins involved in the indicated pathways are presented in a pie chart.

Figure 3. Hat1-dependent acetylation of chromatin proteins. A) A subset of chromatin-related proteins whose acetylation decreased at least 2.5-fold in Hat1 KO cells. B) Schematic diagram showing the domains of CBP. Hat1-dependent sites of acetylation cluster in the autoregulatory loop.

Figure 4. . Hat1-dependent acetylation of transcriptional regulators. A) Subset of transcriptional regulators whose acetylation decreased at least 2.5 fold in Hat1 KO cells. B) Hat1 WT and Hat1 KO were grown in galactose and stained with antibodies recognizing unmodified p53 and p53 acetylated at lysine 381 (386 in human). Fluorescent signals were quantitated (right).

Figure 5. Hat1is required for normal mitochondrial function. A) Subset of mitochondrial proteins whose acetylation decreased at least 2.5 fold in Hat1 KO cells. B) Cells were fractionted int o neuclear, cytoplasmic and mitochondrial extracts. Western blots of the extracts were probed with the indicated markers. VDAC is mitochondrial, GAPDH is cytoplasmic nd Lamin is nuclear. C) Mitochondrial were purified by a percoll gradient. Western blot was probed with the indicated antibodies. D)HEK293 cells were transfected with MitoLight expressing vector. Cells were visualized with a confocal microscope and an individual Z-plane is shown.

Table 1: Top 50 proteins whose acetylation increases (green) or decreases (red) during growth on galactose relative to growth on glucose

Protein Name	HAT1+/+ Galactose : Glucose	HAT1-/- Galactose : Glucose
PDZK1	108.1	2.9
SULT1A1	66.0	49.4
EHD3	30.0	3.3
DDIT3	26.8	3.4
ADAM1A	18.4	46.7
CBP	13.6	-1.7
CXorf57	13.1	1.7
OXGR1	11.5	17.4
KIF18B	11.1	-3.9
PKM	8.9	9.4
MTHFD2	8.8	6.9
Titin	8.7	1.6
EIF5A	8.4	1.8
HMHA1	7.9	5.3
H2A	7.8	-3.9
COL27A1	7.7	1.2
H1	7.4	1.0
DOCK4	7.3	8.9
RPL8	7.2	1.9
DCXR	6.9	15.4
PKM	6.9	6.9
GAPDH	6.8	3.8
CDK1	6.6	2.5
VPS16	6.5	4.2
C19orf47	6.5	2.6
ATP6V1E1	6.3	-1.1
TSR2	6.3	1.6
ANXA2	6.2	2.3
H2AZ	6.0	-2.2
NCL	6.0	-3.1
AGK	5.9	-1.4
INTS8	5.6	3.7
H2AZ	5.6	-1.8
SYNJ2	5.6	-1.4
H2AFJ	5.4	-1.5
APOB	5.4	2.6
EPG5	5.3	8.2
CTU2	4.9	-1.2
MYH9	4.8	-1.1
GDI2	4.8	4.1
FASN	4.8	4.3
SLC25A4	4.8	2.8
TPR	4.8	-2.5
XPG	4.7	3.6
CCDC45	4.7	1.0
RAVER1	4.7	3.7
SLC25A5	4.6	2.3
PRPF38B	4.6	3.5
KIAA2022	4.5	1.2
ATP5A1	4.5	2.8

Protein Name	HAT1+/+ Galactose : Glucose	HAT1-/- Galactose : Glucose
DATF1	-42.3	-13.0
CXXC1	-39.0	-4.6
GTF3B	-32.5	-90.1
CCNT1	-30.1	-55.0
MgcRacGAP	-27.0	-5.2
Za20d3	-17.5	-14.4
WDHD1	-17.4	-47.5
GPATCH8	-17.4	-10.5
CSRP1	-17.2	-4.3
PCNP	-15.0	-11.3
SMARCC1	-14.3	-11.7
CDK13	-12.5	-26.0
SRC-2	-12.0	-5.5
hnRNP M	-11.4	-6.6
Bcl-9	-10.9	-3.5
BRD2	-10.5	-3.3
C19orf43	-10.4	-14.4
RTF1	-9.9	-6.4
CHD-9	-9.9	-10.4
CCL5	-9.5	-5.0
PNUTS	-9.4	-17.4
Bcl-9	-9.2	-8.0
HELLS	-9.1	-4.1
MEF2D	-9.1	-9.3
FIR	-9.0	-3.5
AKAP8L	-9.0	-5.8
ZC3H14	-8.8	-10.1
TIF1G	-8.5	-7.8
RBM17	-8.4	-7.9
RDBP	-8.0	-8.5
HCFC1	-8.0	-3.8
BAZ1A	-8.0	-2.5
NFAT5	-8.0	-4.0
BRD3	-7.9	-3.0
ARPP-19	-7.8	-7.8
TCF20	-7.7	-10.1
CBP	-7.6	-5.0
MED8	-7.5	-8.8
anillin	-7.5	-2.8
HMG20B	-7.5	-6.8
CBLL1	-7.5	-5.4
RPS20	-7.4	-3.8
Mena	-7.4	-3.0
CCNT1	-7.4	-8.8
ZMAT2	-7.3	-11.3
JunB	-7.3	-8.1
PNUTS	-7.2	-11.9
MED6	-7.2	-7.4
ZNF292	-7.1	-1.6
BRD2	-7.1	-3.6

Table 2: Acetylation sites that decrease by at least 2.5-forld in Hat1$^{-/-}$ cells relative to Hat1$^{+/+}$ cells.

PROTEIN	ACETYLATION SITE(s)	HAT1-/-/HAT1+/+	CBP SUBSTRATE
CBP	1588, 1592, 1593	-20.2	YES
EHD3	13	-13	NO
HSP105	850***	-7.1	NO
KIF18B	775***, 776***	-6.3	NO
H2B	17	-6.1	YES
THUMPD1	284	-5.6	YES
TUSC3	310***	-5.4	NO
Titin	10074***	-5.4	NO
H2A	6	-5.0	YES
RPL8	42***	-4.9	NO
H2AFJ	6, 10	-4.9	YES
PDZK1	248	-4.8	NO
H2AZ	8, 12	-4.7	YES
CBP	1584, 1587	-4.3	YES
CBP	1592, 1593	-4.2	YES
NCL	132	-4.1	YES
RPL14	201***	-4.1	NO
NCL	135***, 138***	-4.0	NO
ID1	19***	-4.0	NO
C/EBP-zeta	503***	-3.9	NO
RRP15	228	-3.8	YES
MTA2	592	-3.8	YES
H2AZ;	5, 8	-3.8	YES
COL27A1	540***	-3.6	NO
hnRNP U	223***	-3.6	NO
TPR	531	-3.6	NO
CCDC27	224***	-3.5	NO
RPL6	5	-3.5	YES
calpastatin	15	-3.5	YES
H4	9	-3.4	NO
TIA1	145***	-3.4	NO
VEGF iso6	93***	-3.4	NO
NIP30	168	-3.3	YES
ATP6V1E1	68	-3.3	NO
C19orf47	16***8, 171***	-3.2	NO
NUCKS	35	-3.1	YES
PDLIM2	279***	-3.1	NO
HSD17B4	46	-3.1	NO
TWIST1	33***	-3.0	NO
H3	19	-3.0	YES
RPS30	11***	-3.0	NO
PPP2R2A	402	-2.9	NO
RAI1	1076, 1079	-2.9	NO
AGK;ATP5B;DECR1	231;159;244	-2.9	YES
VDAC3	90	-2.9	NO
SET	38	-2.9	NO
p53	376, 381	-2.8	YES
DAP	20	-2.8	YES
MLLT10	451	-2.8	YES
SYNJ2	1034***	-2.8	NO
EWS	643	-2.7	YES
U5-200kD	55	-2.7	YES
Lasp-1	207	-2.7	YES
H1E	168	-2.7	YES
GALNT5	306***	-2.7	NO
NPM1	154	-2.7	YES
HDGF	105	-2.6	YES
DMAP1	463	-2.6	YES
KIAA2022	672	-2.6	NO
DPY30	40	-2.6	YES
Bcl-9L	36	-2.6	NO
Hbo1	434***	-2.6	NO
UBF	559	-2.6	YES
Myb	428***	-2.6	NO
anillin	642***	-2.6	NO
DOCK7	1952***	-2.5	NO
HMGA1	15	-2.5	YES
DDX42	807	-2.5	YES
CCDC45	586	-2.5	YES
NCL	62	-2.5	NO
NCL	225***, 226***	-2.5	NO

***Not Previously Reported

A

470 PEPTIDES

297   173

■ UP
■ DOWN

B

-log₁₀P VALUE

C

GLUCOSE

HEXOKINASE

GLUCOSE-6-PHOSPHATE

PHOSPHOGLUCOSE ISOMERASE

FRUCTOSE-6-PHOSPHATE

PHOSPHOFRUCTOKINASE

FRUCTOSE-1,6-DIPHOSPHATE

ALDOLASE

DIHYDROXYACETONE PHOSPHATE ⟷ GLYCERALDEHYDE-3-PHOSPHATE

TRIOSEPHOSPHATE ISOMERASE   GLYCERALDEHYDE-3-PHOSPHATE DEHYDROGENASE

GLYCERALDEHYDE-1,3-DIPHOSPHATE

PHOSPHOGLYCERATE KINASE

GLYCERATE-3-PHOSPHATE

PHOSPHOGLYCEROMUTASE

GLYCERATE-2-PHOSPHATE

ENOLASE

PHOSPHOENOLPYRUVATE

PYRUVATE KINASE

PYRUVATE

Figure 1

Figure 2

A

PROTEIN	FOLD DECREASE	SEQUENCE*
CBP	20.2 4.3 4.2	$_{1588}$**K**KNN**KK**TNK$_{1596}$ $_{1581}$GDS**K**NA**K**K$_{1588}$ $_{1590}$NN**KK**TNK$_{1596}$
Histone H2B	6.1	$_{15}$**KK**AVTK$_{20}$
Histone H2A	5.0	$_{4}$G**K**QGG**K**AR$_{11}$
HISTONE H2A.J	4.9	$_{5}$G**K**QGG**K**VR$_{12}$
HISTONE H2A.Z	4.7 3.8	$_{5}$AG**K**DSG**K**AK$_{13}$ $_{1}$AGG**K**AG**K**DSGK$_{11}$
HISTONE H4	3.4	$_{4}$GKGG**K**GLGK$_{12}$
HISTONE H3	3.0	$_{18}$**K**QLATK$_{23}$
HISTONE H1.4	2.7	$_{160}$KPAAAAGA**K**K$_{169}$
MTA2	3.8	$_{585}$SSSQPAA**K**R$_{593}$
HBO1	2.6	$_{429}$LFLDH**K**TLYYDVEPF LFYVMTEADNTGCHLI GYFSK$_{4648}$

*Acetylated lysine denoted by **K**

B

CBP

🔵=Hat1-DEPENDENT ACETYL LYSINE

Figure 3

A

PROTEIN	FOLD DECREASE	SEQUENCE*
ID1	6.0	$_3$VASGSAAAAAGPSCSL**K**AGR$_{22}$
C/EBP-z	3.9	$_{498}$QIGDD**K**VR$_{505}$
TWIST1	3.0	$_{27}$QQPASG**K**R$_{34}$
p53	2.8	$_{376}$**K**TMVK**K**$_{381}$
AF-10	2.8	$_{432}$SFDNSPGELGSSSLPTAGY**K**R$_{452}$
UBF	2.6	$_{550}$APPAATNSS**K**K$_{560}$
MYB	2.6	$_{214}$LDAPTLPSTPLIGH**K**$_{428}$

*Acetylated lysine denoted by **K**

B

Figure 4

A

PROTEIN	FOLD DECREASE	SEQUENCE*
AGK**	2.9	$_{231}$GPL**K**TK$_{236}$
ATP5B**	2.9	$_{156}$GPI**K**TK$_{158}$
DECR1**	2.9	$_{2417}$GPI**K**TK$_{243}$
VDAC3	2.9	$_{75}$WNTDNTLGTEISWEN**K**L AEGLK$_{96}$

*Acetylated lysine denoted by **K**

**Isobaric peptides

B

C

D

DAPI   MitoLight

HAT1   MERGE

Figure 5