1 **Meiosis and Kinetochore genes are used by cancer cells as genome destabilizers and**

2 **transformation catalysts**

4 Roshina Thapa[1], Swetha Vasudevan[1], Mimi Abo-Ayoub Ashqar [1], Eli Reich[1], Nataly Kravchenko-

5 Balasha[1*] and Michael Klutstein[1]*

6
7
8 [1] Institute of Dental Sciences, Faculty of Dental Medicine, The Hebrew University of Jerusalem,
9 P.O.B. 12272, Ein Kerem, Jerusalem, ISRAEL 91120
10
11 Corresponding Authors: Michael Klutstein and Nataly Kravchenko-Balasha
12 Email: michaelk@ekmd.huji.ac.il ;   natalyk@ekmd.huji.ac.il

18   * M.K and N.K.B contributed equally to this work

20 **Keywords:** Cancer, Chromosomal instability, Meiosis genes, Kinetochore genes, overexpression,

21 CNV, surprisal analysis.

Abstract:

Cancer cells have an altered transcriptome which contributes to their altered behaviors compared to normal cells. Indeed, many tumors express high levels of genes participating in meiosis or kinetochore biology, but the role of this high expression has not been fully elucidated. In this study we explore the relationship between this overexpression and genome instability and transformation capabilities of cancer cells. For this, we obtained expression data from 5 different cancer types which were analyzed using computational information-theoretic analysis. We were able to show that highly expressed meiotic/kinetochore genes were enriched in the altered gene networks characterizing unstable cancer types with high chromosome instability (CIN). However, altered networks characterizing the cancers with low CIN did not include meiotic and kinetochore genes. Representative gene candidates, found by the analysis to be correlated with a CIN phenotype, were further explored by transfecting genomically-stable (HCT116) and unstable (MCF7) cancer cell lines with vectors overexpressing those genes. This overexpression resulted in abnormal cell divisions, defective spindle formation and increased transformation properties in stable cancer HCT116 cells. Interestingly the genome stability properties of the unstable MCF7 cancer cells were less affected by gene overexpression. Our results indicate that overexpression of both meiosis and kinetochore genes drives genomic instability and cancer progression.

Introduction:

Cancer is a complex disease, characterized by numerous genomic aberrations and by dysregulation of gene expression. Along with overexpression of oncogenes and repression of tumor suppressors, tumors often express various tissue specific genes, not necessarily related to their primary tissue of origin (1-3). In particular, cancer cells have been found to frequently express genes that are normally restricted to the testis. These genes can be referred to as cancer/testis antigens (CTA) (4). Many CTA have been found to be involved in meiotic divisions, which occur in the testis and include processes with inherent genome instability (see below). This property of the meiotic genes has led to the intriguing idea that the expression of CTA in tumors drives genome and chromosome instability in those tumors (5, 6).

Chromosome instability (CIN) is considered to be one of the most important cancer hallmarks. CIN tumors, which have the propensity to constantly change their genome, have worse prognosis than non-CIN tumors (7, 8). Previous work shows that CIN tumors use several molecular mechanisms to achieve their instability, such as replication stress and modulation of the spindle assembly checkpoint (9-11). Due to inherent functions of CTA genes involved in meiosis, that include mono-orientation of sister kinetochores and DNA double-strand break formation and repair, CTA have become prime candidates initiating an additional mechanism to achieve CIN (12-15).

Several small scale studies have already shown that a cohort of meiotic genes is expressed in different tumors (6, 16-19). Of special note is the fact that overexpression of meiotic cohesin Rec8 in fission yeast cells causes uniparental disomy of chromosomes in mitotic cells (20).

Another emerging player in the generation of CIN in cancer is the kinetochore (21). Kinetochores are protein complexes built on centromeres, the specialized loci on eukaryotic chromosomes, playing a key role in mediating chromosome segregation (21). This is mainly achieved through the physical connection between microtubules and the centromeric DNA (22). The balance between all the different kinetochore components is crucial for maintaining genome stability and correct ploidy. Under- or overexpression of different kinetochore components may lead to the formation of chromosomes with very little microtubule attachment, or on the contrary, too many microtubules binding to a chromosome (22-25). Eventually this may lead to non-disjunction and aneuploidy (23-25). Overexpression of specific kinetochore components such as the inner-

centromere protein CENP-A (centromere specific ortholog of histone H3 serving as the structural

basis of the kinetochore) leads to deposition of kinetochore components on additional loci in a

chromosome already containing a centromere, and the formation of di-centric chromosomes,

resulting in a breakage-fusion-bridge cycle of chromosomes and CIN (26). On the other hand,

insufficient CENP-A can result in senescence of cells and apoptosis (27, 28). Misregulation of

kinetochore components has been observed in many tumors (29). Alterations in the expression

levels of kinetochore genes may also cause CIN in tumors, as well as affect the prognosis of

specific patients and their response to therapy (29). Despite all these studies the role that the

kinetochores play as drivers of CIN during tumorigenesis is not fully understood.

To explore further the relationship between meiosis and kinetochore genes and genome instability

we performed a large scale computational analysis of normal and cancer tissues which were

obtained from breast, bladder, stomach, colorectal and cervical cancer and normal tissues, all from

TCGA data (https:\\xena.ucsc.edu\). We have demonstrated that tumors with high CIN

overexpressed meiosis and kinetochore genes in cancer specific gene correlation networks.

Although tumors were heterogeneous and could be characterized by different altered gene

expression networks, meiosis and kinetochore altered transcripts could be found in different

compositions in high CIN tumors but not in low CIN patients within the same type of cancer (see

Fig. 1).

To further validate our hypothesis that meiosis and kinetochore genes drive CIN we performed

experimental studies in genomically stable and unstable cancer cell lines (CIN+ and CIN-,(8)). We

have demonstrated that overexpression of meiosis and kinetochore genes in cancer cell-lines

increases genome instability in this setting. Moreover, we show that this over-expression elevates

significantly genome instability in genomically stable cancer cell lines, but less in unstable cell

lines. Overexpression of these genes also led to the enhanced transformation and invasiveness

properties of the cancer cell lines, providing an experimental evidence for the involvement of

meiosis and kinetochore genes in genome instability and cellular transformation. An overview of

the study is summarized in Figure 1.

Materials and Methods:

1. Data analysis

1.1 Thermodynamic-based information theoretical approach (Surprisal Analysis)

Matrix of gene expression data was obtained from TCGA database for each cancer type. Every dataset is profiled for thousands of transcripts (total 20,530). The matrix was used as an input file for the information-theoretic surprisal analysis using MATLAB software (30). The analysis was used in several studies, for example (30-32). Briefly, we identify in normal and tumor subsets, the expected gene expression levels at the steady state (a state in which the biological processes are balanced) and deviations thereof for each transcript i. The deviations occur due to environmental/genomic constraints. Any biochemical/genetic perturbation can be considered as a constraint and elicit a coordinated change in a group of transcripts. These groups of transcripts are named **unbalanced processes** and identified through calculations of $G_{i\alpha}$ values (=weights of participation) for each transcript i in each process α (α=1,2..3). Each transcript can participate in more than one unbalanced process due to non-linearity of biological networks. Only the transcripts located on the tails of the distributions of $G_{i\alpha}$ values (See plots in the Tables S3-S7) are analyzed further for biological meaning. Additionally, the analysis identifies an amplitude, $\lambda_{\alpha}(k)$, or an importance of each process α in each tissue k (**Fig. S1**). Sign of $G_{i\alpha}$ and, $\lambda_{\alpha}(k)$ means correlation or anti-correlation between the transcripts in the same process α (in case of $G_{i\alpha}$) or α between the same processes in different tumors (in case of $\lambda_{\alpha}(k)$). For example, if the process α is assigned the values: $\lambda_{\alpha}(1) = 37$, $\lambda_{\alpha}(20) = 0$, $\lambda_{\alpha}(33) = -39$, it means that this process influences the tumors of the patients indexed 1 and 33 in the opposite directions, while it is inactive in patient 20. In order to calculate whether a particular transcript was induced or reduced due to a process α, the product $G_{i\alpha}*\lambda_{\alpha}(k)$ is calculated for each transcript.

 In summary we identify for each transcript a set of unbalanced processes in which it is involved and quantify how important each process is in each normal/cancer sample. Thus, a comprehensive map of unbalanced processes is obtained for each cancer or normal sample that allows to characterize each tissue in heterogeneous datasets in detail. Detailed description on how surprisal

154 analysis is implemented in biology and how $G_{i\alpha}$ and, $\lambda_\alpha (k)$ are computed is provided in detail in

155 (30-32)

1.2 Computation of copy number variation

We obtained data of copy number variation (CNVs) in cancer population from the TCGA genomic

database. Copy number variation (CNVs) are a type of structural variant involving alterations in

the number of copies of specific regions of DNA, which can be either deleted or duplicated. These

chromosomal deletions and duplications involve large stretches of DNA (that is, thousands of

nucleotides, which may span many different genes) but can range considerably in size as well as

prevalence. Only CNVs larger than 1 Mb (large CNVs usually correlate with the genomic stability)

were considered for further analysis and are thus termed lCNVs (large chromosome number

variations). The number of lCNVs for each sample were summed and determined using R tools.

The distribution for each cancer type is shown in **Fig. 2**.

2. Experimental methods

2.1 Cell lines and culture

HCT116 human colon adenocarcinoma cell lines were maintained in Dulbecco's Modified Eagle's

medium (DMEM, Sigma), supplemented with 10% fetal bovine serum (FBS), 1% PenStrep (100

U/mL Penicillin and 100 μg/mL Streptomycin) and 4 mM L-glutamine in a 37 °C incubator (5%

CO_2). MCF-7 breast cancer cell lines were maintained in RPM-1640, (Sigma), supplemented with

10% fetal bovine serum (FBS), 1% PenStrep (100 U/mL Penicillin and 100 μg/mL Streptomycin)

and 4 mM L-glutamine in a 37 °C incubator (5% CO_2).

2.2 DNA preparation and Transfection.

Meiotic and kinetochore genes tagged with Enhanced Green Fluorescent Protein (EGFP) (from

BD Biosciences) were used. The GenElute, HP Plasmid Midiprep Kits (Bio Basic Inc, Canada) .

was used to isolate the plasmid. All cells were transfected 24h after initial plating. Transfections

were performed using the Mirus transfection reagent (cat-81094967, Zotal, USA) with a 4:1

(transfection reagent: DNA) ratio. Transfection efficiency for HCT116 and MCF-7 cells was

evaluated by counting the number of GFP positive cells by immunoflorescent microscope and

calculating the percentage based on the total number of cells.

2.3 Cell synchronization and immunostaining.

HCT116 and MCF-7 cells were synchronized by double thymidine block (33). Cells were treated

with 2mM thymidine for 18 h in medium supplemented with 10% FBS. After washing twice

with PBS, cells were cultured in fresh medium for 9h and again treated for 15h with media

containing 2mM thymidine (10% FBS). After washing cells with PBS, the block was released by

the incubation of cells in fresh medium and cells were harvested at 9h (HCT116) and 11h (MCF-

7) and fixed with methanol. After that immunostaining was performed. Cells were washed 3

times with PBS and blocked with 5% BSA diluted with PBS. After that, cells were incubated

with first Ab (α-tubulin antibodies, T5168, Sigma, USA, 1:400 dilution) for 2 hr at room temp.

After washing twice with PBS, cells were incubated with a secondary Ab (donkey anti-mouse

IgG antibodies, life science,USA, 1:200 dilution) for 1 hr at room temp followed by washing the

cells twice with PBS. Again, the cells were incubated with Hoechst 33342 (cat: PIR-62249

Thermo scientific, Germany) diluted 1:10,000 for 5-30min for DNA visualization.

2.4 Soft agar assay

Colony formation on soft agar was assayed in triplicate by plating 5000 cells in a layer of 0.3%

(w/v) agar in assay DMEM (HCT116) and RPMI medium (MCF-7) medium, on top of a 0.6%

(w/v) agar layer. Plates were incubated at 37 °C and 5% (v/v) CO_2 for 3 weeks, and the medium

was replaced every 4 d. Colonies were stained using 0.005% (w/v) Crystal Violet solution, and an

image of the whole well was acquired using an Olympus SZ61 stereomicroscope. Colonies were

counted using ImageJ software (http://imagej.nih.gov/ ij/). The area and number of colonies was

calculated.

3. Statistics

Significance was determined using a two-tailed Student's *t* test. The $p < .001$ was considered as

extremely significant (***), $p < .01$ as highly significant (**), and $p < .05$ as statistically

significant (*).

Results:

Degree of genome instability varies between different cancer types

We hypothesized that the extent of alterations in gene expression levels of meiotic and kinetochore genes may be related to the degree of genome instability. To explore this, we determined the distribution of large DNA tracts which exhibited copy number variations in five different cancer types, namely breast (n=1104), bladder (n=407), stomach (n=415), colorectal (n=383) and cervical (n=305) cancer patients and compared them to normal tissues from breast (n=114), bladder (n=19) stomach (n=20) colorectal (n=51) and cervix (n=3), all from TCGA database. Copy number variation (**CNV**) is a type of chromosomal structural variation that involves alterations in DNA copy number of specific genome regions. Those regions can be either deleted or duplicated. The chromosomal deletions and duplications can involve large stretches of DNA, e.g. thousands of nucleotides, which may span many different genes. In our analysis we included only CNVs that were larger than 1 Mb (lCNV= large CNVs of more than $1*10^6$ Base pairs, 1 Mb). We found that Bladder tumors had large number of DNA regions with lCNVs (<u>high lCNV value</u>) with a maximum value of 850 1CNVs and an average value of 112 1CNVs. Breast and stomach cancers also show high values of lCNVs (with a maximum of 821 1CNVs, and average of 109 1CNVs for breast and a maximum of 650 lCNVs and average of 108 1CNVs for stomach cancer) (**Fig 2,B**). On the other hand, colorectal and cervical cancer types have less DNA regions with lCNV (max 402, avg lCNVs=79 and max 278, avg 1CNVs= 80 respectively) (**Fig. 2A, B**). (**p<0.001**)

Altered gene expression networks, characterizing cancer samples with genome instability, are enriched with overexpressed kinetochore and meiosis transcripts

To explore the relationship between the altered expression of meiosis and kinetochore genes and genome instability we performed a large scale unbiased computational analysis of over 2800 normal and cancer tissues from 5 cancer types mentioned above. We utilized a computational information-theoretic surprisal analysis (<u>SA</u>) (31, 32) in order to identify altered gene co-expressed networks in cancer tissues, named *unbalanced processes*. Several different unbalanced processes (=networks) may occur in a particular cancer type due to inter-patient heterogeneity. SA deciphers

a number of unbalanced processes in each cancer type, by decoding the expected expression levels

of the tested molecules at the steady state (i.e. the balanced, unconstrained state), and the deviations

thereof due to environmental or genomic constraints. These constraints elicit unbalanced processes

in the system (see (31, 32) for more details). Co-varying altered transcripts that deviate from the

steady state significantly and in a coordinated manner are grouped and each group represents an

unbalanced process α. The analysis determines those transcripts through calculation of a "weight

of participation", $G_{i\alpha}$, of each transcript i in a process α (**Table S1**). Transcripts with significant

$G_{i\alpha}$ values are analyzed further using the David database (https://david.ncifcrf.gov/) in order to

assign a biological meaning for each process α. Every altered transcript can be involved in several

unbalanced processes.

Next the analysis assigns an amplitude, $\lambda_\alpha(k)$, an importance of each process α in each tissue k.

Table S2 lists amplitudes of all processes in every cancer type and in every tissue. Several distinct

unbalanced processes can be active in each cancer type/cancer tissue(32, 34). Detailed description

of the analysis can be found in (32).

Using SA we identified 16 distinct unbalanced processes in breast cancer **(Fig. 3A**, rigorous error

analysis, as described in (32, 34, 35) and Methods, is applied in order to determine a number of

unbalanced processes beyond the noise in each dataset). Processes with lower indices, such as

processes 1 and 2, were the most dominant and appeared in high a percentage of the patients **(Fig.**

S1). Each process included different biological categories. For example, the most dominant

process, process 1, which was found in 21% of breast cancer patients (239 of 1095 breast cancer

samples, **Fig. S1**). This process included induced transcripts involved in biological categories such

as cell division, sister chromatid cohesion and chromosome segregation **(Table S3)**. Process 2

appeared in 20% of breast cancer patients **(Table S3)** and included induced transcripts involved in

cell division, mitotic nuclear division and sister chromatid cohesion **(Table S3)**. Interestingly, less

common unbalanced processes (with higher indices) included less kinetochore and meiosis genes

and consequently less kinetochore/meiosis related enriched biological categories **(Fig. 3A)**.

Similar results were found for the two other types of cancer with high values of lCNVs: bladder

cancer **(Fig. 3B)** and stomach cancer **(Fig. 3C) (Tables S4, S5)**.

In contrast, the most dominant unbalanced processes in cancers with lower values of lCNVs

(colorectal and cervical cancers) were not particularly enriched for meiosis and/or kinetochore

genes (**Fig. 3D, E, Table S6-S7).** These results show a correlation between high lCNV values and

overexpression of meiosis/ kinetochore genes **(Fig. 3A-C)** and evince a possibility that highly

expressed meiosis and kinetochore genes in cancer might be involved in genome instability of

those cancers.

To further investigate the correlation between the altered expression levels of meiosis/kinetochore

genes and genome instability, we examined every cancer type individually. We compared cancer

samples with highly unstable genomes (10% of the samples with highest lCNVs values) to the

cancer samples with relatively stable genomes (10% of the samples with lowest lCNVs values)

within <u>the same cancer type</u>. The results demonstrate that in breast cancer, the most dominant

processes harboring meiosis/kinetochore genes appeared mainly in cancer tissues with highly

unstable genomes (high lCNVs, **Fig. 4A**). Patients having more stable genomes (lower lCNVs)

did not harbor those networks (**Fig. 4B**). Similar results were found in bladder and stomach cancers

(**Fig. 4C, D and Fig. 4E, F** respectively**).** In contrast, the correlation between gene expression

levels of meiosis/kinetochore genes and genome instability was not found in the cancer types with

relatively stable genomes, (colorectal and cervical cancers, **Fig. S2)**.

These results point to a high correlation between participation of meiosis/kinetochore in dominant

unbalanced cancer-specific processes and genome instability, showing that meiosis and

kinetochore genes may have an active role in driving CIN. This strong correlation prompted us to

examine our hypothesis experimentally.

Over expression of representative meiosis and kinetochore genes in genomically stable and

unstable cancer cell lines

In order to test our hypothesis, we have overexpressed representative kinetochore and meiosis

related genes, which were found in the analysis (see for example Table S2), in cancer cell lines.

The two kinetochore genes we overexpressed, CENP-A and HJURP, represent the structural basis

of the kinetochore structure (19, 36). CENP-A is the histone H3 homolog that forms a platform

upon which all other kinetochore components assemble (37). HJURP is the CENP-A chaperone,

responsible for recruiting it to the centromere (38). We also overexpressed four meiosis-related

genes: DMC1, SPO11, SMC1B and REC8. DMC1 and SPO11 are involved in homologous

recombination and DNA double strand break repair during meiosis (39, 40). SMC1B and REC8

are meiosis specific components of cohesin, and participate in homologous chromosome pairing

and in sister chromatid mono-orientation (41-43). We have overexpressed these genes in two

cancer cell-lines- HCT116, a colon cancer cell line which is CIN negative and has a relatively

stable genome, and MCF7, a breast cancer cell line which shows high chromosome instability (8,

44). The genes were fused to GFP to monitor their expression (**Fig. S3**).

Over expression of meiosis and kinetochore genes promotes genome instability in cancer cell

lines

In order to check whether the overexpressed meiosis and kinetochore genes promote genome

instability we evaluated the number of cells with lagging chromosomes, anaphase bridges, uneven

segregation of chromosomes and deviation from a bipolar spindle configuration as a means to

estimate genome stability (25, 45-47). **Figure 5 (B, C and G)** shows that overexpression of both

kinetochore genes, CENP-A and HJURP, and one of the meiosis genes, DMC1 in HCT116, caused

a significant elevation in anaphase bridges and lagging chromosomes (more examples in **Fig. S4**).

Other meiosis-related genes did not affect the chromosome segregation phenotype when over-

expressed. **Figure 5 (E, F and I)** also shows that overexpression of all meiosis and kinetochore

genes in HCT116 cells caused a significant elevation in mono-polar and multi-polar spindle

formation compared to an empty plasmid (G, H, I and J) (more examples in **Fig. S5**).

In contrast, overexpression of meiosis and kinetochore genes in the genomically unstable MCF7

cells caused a significantly less severe phenotype **Figure 5** (**H and J**). Only CENP-A

overexpression caused a small elevation in the occurrence of mono-polar and multi-polar spindles

(**Fig. 5J**). However, all other phenotypes, related to the spindle (**Fig. 5J**) and chromosome

segregation (**Fig. 5H**), were not affected in MCF7 cells in response to induced expression of

meiosis/kinetochore genes.

These results demonstrate that overexpression of kinetochore and meiosis genes in genomically

stable cells has the ability to promote genome instability. The same overexpression has a

significantly smaller effect in a cell line that has already acquired a high degree of genome

instability before the gene transfection.

Over expression of meiosis and kinetochore genes promotes invasiveness of cancer cells

Cancers with unstable genomes are often more invasive than cancers with stable genomes (8, 48,

49). Therefore, we hypothesized that overexpression of meiosis and kinetochore genes and

promotion of genome instability could induce invasiveness and transformation properties of cancer
cells.
To examine a change in the transformation properties of the cells we tested an ability of HCT116
and MCF7 cells to generate colonies in soft agar following overexpression of the
meiosis/kinetochore genes (50, 51). **Figure 6 and S6** show that overexpression of all meiosis and
kinetochore genes in HCT116 cells enhanced both the number of colonies generated and the size
of the colonies, demonstrating enhanced cancer transformation properties. Surprisingly, although
genome instability parameters were not increased in the MCF7 cells upon the gene overexpression,
an elevation in the number and size of the colonies in soft agar was detected in this cell line (**Fig.**
S7). These results show that induced invasiveness and cellular transformation of the tested cell
lines may correspond to the induced expression of meiosis and kinetochore genes, although not
necessarily linked to the ability of those genes to induce genome instability parameters.
350
351
Discussion
Genome instability and the mechanisms behind it are among central questions in cancer biology
in recent decades (52). Here we investigate a new possible route to achieve genome instability in
cancer by overexpression of genes which participate in meiosis or in kinetochore formation. The
process of meiosis includes inherent genome instability which occurs through meiotic homologous
recombination and sister chromatid mono-orientation. The perturbed expression of kinetochore
proteins has also potential to affect the processes involved in proper chromosome segregation and
genome stability.
We have taken a multi-pronged approach to support our hypothesis and have used computational
analysis of large cancer datasets and an experimental approach utilizing cancer cell line models.
Using information-theoretic computational analysis of five different cancer types obtained from
TCGA database, we have shown that the most abundant altered gene expression networks,
characterizing unstable cancers, were enriched with meiosis and kinetochore genes. Altered gene
expression networks characterizing cancers with low lCNVs were not enriched with those genes.
Moreover, in unstable cancers patients with the highest lCNVs were characterized by the
unbalanced processes enriched with meiosis and kinetochore genes in contrast to the patients with
low lCNVs within the same cancer type which did harbor those processes.

Although these analyses were merely restricted to a correlation, the finding of this correlation in
five major cancer types, and the extension of the correlation to the specific patient groups within
each cancer hints to a strong link between the overexpression of meiosis and kinetochore genes
and genomic instability in tumors.

In order to go beyond this correlation and demonstrate a causative effect, we performed
experiments in which several representative meiosis and kinetochore genes, as identified by the
computational analysis, were expressed in two cancer cell lines. The overexpression caused an
elevation of genome instability parameters in the stable HCT116 cell lines, but not in the less stable
MCF7 cell line. However, overexpression caused both cell lines to increase their invasiveness and
transformation properties as measured by colony formation ability in soft agar.

These results could suggest that induced invasiveness (observed in both cell lines) is not directly
related to the induced genome instability (observed in HCT116) although they both result from the
overexpression of the same genes. Another possibility is that even a slight and undetectable
increase in genome instability (as in MCF-7), can cause a large effect on invasiveness of the cells.
Further experiments are needed to distinguish between these possibilities.

Our results also demonstrate that meiosis and kinetochore genes can serve as markers for genome
instability. Future work should assess the accuracy and sensitivity of those markers and whether
downregulating meiosis and kinetochore genes could be used as a therapeutic approach.

In conclusion, we have shown that genome instability in tumors could be driven by overexpression
of specific classes of genes, namely meiosis and kinetochore genes, which are involved in genome
organization and maintenance of undifferentiated cells. This finding may have medical
implications regarding the identification of genome instability in tumors and the treatment of
unstable tumors through manipulation of these gene networks.

Figure Legends:

Figure 1. Schematic overview of how mis-expressed meiosis and kinetochore genes induce
chromosome instability. 2821 tumors tissues of five cancer types obtained from TCGA datasets
were analyzed. Cancer types were categorized into two subtypes: cancer with unstable genomes
(unstable cancers) and cancer with a more stable genome (stable cancers). Information-theoretic
analysis is utilized to study the altered gene expression networks in the entire population. We find

that the most dominant cancer-specific altered networks in unstable cancers were enriched with

meiotic and kinetochore but not in stable cancers. The experimental overexpression of meiosis and

kinetochore genes in cancer cell lines induced genomic instability phenotypes: anaphase bridges

(right) and spindle defects (multipolar spindle, left).

Figure 2. Distribution of Large Copy number variation (lCNVs) values in different cancer

types. Bladder cancer has a high number of DNA regions with lCNVs (max 850, avg lCNVs=

112) followed by breast cancer and stomach cancers (max 821, avg lCNVs =109 and max 650 avg

CNVs = 108 respectively). These three cancer types are categorized as unstable cancer types

(p<0.001). On the other hand, colorectal, and cervical cancer types have a low number of DNA

regions with lCNVs (max 402, avg lCNVs=79 and max 278, avg 1CNVs= 80 respectively), and

are categorized as more stable cancer types. Fig. 2A shows the full distribution of lCNVs values

in all samples, while Fig. 2B shows a zoom–in on cases having up to 200 DNA regions with lCNVs

(low CNV values).

Figure 3. Involvement of kinetochore and meiosis genes in unbalanced processes as

identified using surprisal analysis in both unstable and stable cancers. Every dataset is

profiled for thousands of transcripts (total 20,530), which are resolved into altered networks

(unbalanced processes) characterizing each tumor and normal tissue. All transcripts, which

deviate from the balance state in the same (coordinated) way, are organized in groups,

unbalanced processes. Meiosis and kinetochore genes were found to participate in the most

dominant processes in unstable cancers: processes 1 and 2 in breast cancer (Bc, **A)**, processes 2,

3 and 4 in bladder cancer (Bl, **B)** and process 1and 3 in stomach cancer (Sc, **C)**. There is a

significantly lower percentage of meiosis and kinetochore genes in the dominant processes

characterizing colorectal (Co, **D**) and cervical cancers (Ce, **E).**

Figure 4. The correlation between over-expression of meiosis/kinetochore genes and genomic

instability within each cancer type. Unstable breast, bladder and stomach cancers were analyzed

in this section. Each cancer type was analyzed separately as following: Distribution of lCNVs

values was generated for each cancer type. 10% of the samples with highest lCNVs values

(unstable group) and 10% of the samples with lowest lCNVs values (stable group) were selected

in each cancer type. Cancer specific unbalanced processes of 10% of the samples with highest
lCNVs values **(A, C, E)** were compared to the processes appeared in 10% of the samples with
lowest lCNVs values **(B, D, F)**. The most dominant processes harboring meiosis/kinetochore genes
appeared mainly in unstable group **(A, C, E))** in all three cancer types. Patients having more stable
genomes (stable group) harbored the processes which did not include meiosis and kinetochore
genes **(B, D, F)**. Sign "-" means anticorrelation: -1 in **B, D, F** means that all induced transcripts,
including meiosis and kinetochore genes, in process 1 are reduced in process -1 and vice versa.
Tables S3-S7include the information regarding biological categories induced in processes -1, -2
and -3 **(B, D, F)**, which characterize stable groups within breast, bladder and stomach cancers.
440
Figure 5. **Overexpression of meiosis (DMC1, REC8, SPO11 and SMC1B) and kinetochore**
genes (CENP-A and HJURP) causes a genomic instability phenotype. The effect of
overexpressed meiosis and kinetochore genes in stable (HCT116) and unstable (MCF-7) cancer
cell lines was measured by staining cells with DAPI and anti-tubulin IF (see Methods). Scoring
the cells for genome instability was performed by counting defects during anaphase (anaphase
bridges, lagging chromosomes) and apparent spindle formation defects (multipolar and unipolar).
Presented are immunofluorescence images showing the effects of overexpression of meiosis
(DMC1) and kinetochore (CENP-A) genes. Showing abnormal cell divisions i.e. Anaphase Bridge
(**B**) and lagging chromosomes (**C**) and apparent spindle formation defects (unipolar **E** and tripolar
F). Quantification of chromosome segregation and spindle defects in meiosis and kinetochore
transfected stable cell line HCT116 shows more significant defects compared to the MCF 7 cell
line (**G-J**). Statistical significance is shown by asterisks (*$P < 0.05$; **$P < 0.01$; ***$P < 0.001$).
Scale bar equals 10μm.
454
Figure 6. Overexpression of meiosis (DMC1, REC8, SPO11 and SMC1B) and kinetochore
genes (CENP-A and HJURP) causes elevated invasiveness Overexpressed meiotic and
kinetochore genes in a stable cell line (HCT 116) have the ability to promote invasiveness in a soft
agar assay. A soft agar colony formation assay was applied for the detection of transformed cells
when overexpressed with meiotic and kinetochore genes in a stable cell line. The number of
colonies **(A)** and size of the colonies (**B** for full scale or **C** for a zoom in on colonies up to 3000
mm^2) were compared (see Methods). Statistically significant differences compared with empty

vector (No genes) were determined using a Student's 2-tailed t test (*P < 0.05; **P < 0.01; ***P

< 0.001)

Acknowledgments:

We thank the Klutstein and Kravshenko-Balasha labs for fruitful discussions. This work was

supported by a research grant from the Israeli Cancer Association (grant number 20190056) to

MK and NKB and a special donation from the ICA USA section.

Availability of data and reagents:

The authors affirm that all data necessary for confirming the conclusions of this article are

represented fully within the article and its tables and figures. All the necessary reagents from this

study can be shared.

References:

1.　　Nakade S, Yamamoto T, Sakuma T. Cancer induction and suppression with transcriptional
control and epigenome editing technologies. Journal of human genetics. 2018;63(2):187-94.
2.　　Ko JY, Oh S, Yoo KH. Functional Enhancers As Master Regulators of Tissue-Specific Gene
Regulation and Cancer Development. Molecules and cells. 2017;40(3):169-77.
3.　　Klutstein M, Nejman D, Greenfield R, Cedar H. DNA Methylation in Cancer and Aging.
Cancer Res. 2016;76(12):3446-50.
4.　　Bruggeman JW, Koster J, Lodder P, Repping S, Hamer G. Massive expression of germ cell-
specific genes is a hallmark of cancer and a potential target for novel treatment development.
Oncogene. 2018;37(42):5694-700.
5.　　Whitehurst AW. Cause and consequence of cancer/testis antigen activation in cancer.
Annual review of pharmacology and toxicology. 2014;54:251-72.
6.　　Lindsey SF, Byrnes DM, Eller MS, Rosa AM, Dabas N, Escandon J, et al. Potential role of
meiosis proteins in melanoma chromosomal instability. Journal of skin cancer.
2013;2013:190109.
7.　　Krem MM, Press OW, Horwitz MS, Tidwell T. Mechanisms and clinical applications of
chromosomal instability in lymphoid malignancy. British journal of haematology. 2015;171(1):13-
28.
8.　　Burrell RA, McClelland SE, Endesfelder D, Groth P, Weller MC, Shaikh N, et al. Replication
stress links structural and numerical cancer chromosomal instability. Nature.
2013;494(7438):492-6.
9.　　Musacchio A. The Molecular Biology of Spindle Assembly Checkpoint Signaling Dynamics.
Current biology : CB. 2015;25(20):R1002-18.

10.   Foijer F, Xie SZ, Simon JE, Bakker PL, Conte N, Davis SH, et al. Chromosome instability
induced by Mps1 and p53 mutation generates aggressive lymphomas exhibiting aneuploidy-
induced stress. Proceedings of the National Academy of Sciences of the United States of America.
2014;111(37):13427-32.
11.   Iwaizumi M, Shinmura K, Mori H, Yamada H, Suzuki M, Kitayama Y, et al. Human Sgo1
downregulation leads to chromosomal instability in colorectal cancer. Gut. 2009;58(2):249-60.
12.   Sakuno T, Tada K, Watanabe Y. Kinetochore geometry defined by cohesion within the
centromere. Nature. 2009;458(7240):852-8.
13.   Watanabe Y, Nurse P. Cohesin Rec8 is required for reductional chromosome segregation
at meiosis. Nature. 1999;400(6743):461-4.
14.   Ishiguro K, Kim J, Fujiyama-Nakamura S, Kato S, Watanabe Y. A new meiosis-specific
cohesin complex implicated in the cohesin code for homologous pairing. EMBO reports.
2011;12(3):267-75.
15.   Hunter N. Meiotic Recombination: The Essence of Heredity. Cold Spring Harbor
perspectives in biology. 2015;7(12).
16.   Feichtinger J, Larcombe L, McFarlane RJ. Meta-analysis of expression of l(3)mbt tumor-
associated germline genes supports the model that a soma-to-germline transition is a hallmark
of human cancers. International journal of cancer. 2014;134(10):2359-65.
17.   Erenpreisa J, Cragg MS, Salmina K, Hausmann M, Scherthan H. The role of meiotic cohesin
REC8 in chromosome segregation in gamma irradiation-induced endopolyploid tumour cells.
Experimental cell research. 2009;315(15):2593-603.
18.   Kalejs M, Ivanov A, Plakhins G, Cragg MS, Emzinsh D, Illidge TM, et al. Upregulation of
meiosis-specific genes in lymphoma cell lines following genotoxic insult and induction of mitotic
catastrophe. BMC cancer. 2006;6:6.
19.   Gong C, Li T, Li Q, Yan L, Wang T. Rice OsRAD21-2 is expressed in actively dividing tissues
and its ectopic expression in yeast results in aberrant cell division and growth. Journal of
integrative plant biology. 2011;53(1):14-24.
20.   Folco HD, Chalamcharla VR, Sugiyama T, Thillainadesan G, Zofall M, Balachandran V, et al.
Untimely expression of gametogenic genes in vegetative cells causes uniparental disomy. Nature.
2017;543(7643):126-30.
21.   Pesenti ME, Weir JR, Musacchio A. Progress in the structural and functional
characterization of kinetochores. Current opinion in structural biology. 2016;37:152-63.
22.   Pavin N, Tolic IM. Self-Organization and Forces in the Mitotic Spindle. Annual review of
biophysics. 2016;45:279-98.
23.   Thangavelu PU, Lin CY, Vaidyanathan S, Nguyen THM, Dray E, Duijf PHG. Overexpression
of the E2F target gene CENPI promotes chromosome instability and predicts poor prognosis in
estrogen receptor-positive breast cancer. Oncotarget. 2017;8(37):62167-82.
24.   Qi ST, Wang ZB, Ouyang YC, Zhang QH, Hu MW, Huang X, et al. Overexpression of SETbeta,
a protein localizing to centromeres, causes precocious separation of chromatids during the first
meiosis of mouse oocytes. Journal of cell science. 2013;126(Pt 7):1595-603.
25.   Klutstein M, Fennell A, Fernandez-Alvarez A, Cooper JP. The telomere bouquet regulates
meiotic centromere assembly. Nat Cell Biol. 2015;17(4):458-69.

26.   Tomonaga T, Matsushita K, Yamaguchi S, Oohashi T, Shimada H, Ochiai T, et al.
Overexpression and mistargeting of centromere protein-A in human primary colorectal cancer.
Cancer research. 2003;63(13):3511-6.
27.   McGregor M, Hariharan N, Joyo AY, Margolis RL, Sussman MA. CENP-A is essential for
cardiac progenitor cell proliferation. Cell cycle (Georgetown, Tex). 2014;13(5):739-48.
28.   Lermontova I, Koroleva O, Rutten T, Fuchs J, Schubert V, Moraes I, et al. Knockdown of
CENH3 in Arabidopsis reduces mitotic divisions and causes sterility by disturbed meiotic
chromosome segregation. The Plant journal : for cell and molecular biology. 2011;68(1):40-50.
29.   Zhang W, Mao JH, Zhu W, Jain AK, Liu K, Brown JB, et al. Centromere and kinetochore
gene misexpression predicts cancer patient survival and response to radiotherapy and
chemotherapy. Nature communications. 2016;7:12619.
30.   Flashner-Abramson E, Abramson J, White FM, Kravchenko-Balasha N. A thermodynamic-
based approach for the resolution and prediction of protein network structures. Chemical
Physics. 2018;514:20-30.
31.   Remacle F, Kravchenko-Balasha N, Levitzki A, Levine RD. Information-theoretic analysis of
phenotype changes in early stages of carcinogenesis. Proceedings of the National Academy of
Sciences of the United States of America. 2010;107(22):10324-9.
32.   Vasudevan S, Flashner-Abramson E, Remacle F, Levine RD, Kravchenko-Balasha N.
Personalized disease signatures through information-theoretic compaction of big cancer data.
Proceedings of the National Academy of Sciences of the United States of America.
2018;115(30):7694-9.
33.   Kim JA, Tan Y, Wang X, Cao X, Veeraraghavan J, Liang Y, et al. Comprehensive functional
analysis of the tousled-like kinase 2 frequently amplified in aggressive luminal breast cancers.
Nature communications. 2016;7:12991.
34.   Flashner-Abramson E VS, Adejumobi IA, Sonnenblick A, Kravchenko-Balasha N. . Decoding
cancer heterogeneity: studying patient-specific signaling signatures towards personalized cancer
therapy. Theranostics. 2019(in press).
35.   Flashner-Abramson E VS, Adejumobi IA, Sonnenblick A, Kravchenko-Balasha N. . Decoding
cancer heterogeneity: studying patient-specific signaling signatures towards personalized cancer
therapy. . Theranostics2019.
36.   Stellfox ME, Bailey AO, Foltz DR. Putting CENP-A in its place. Cellular and molecular life
sciences : CMLS. 2013;70(3):387-406.
37.   Fukagawa T, Earnshaw WC. The centromere: chromatin foundation for the kinetochore
machinery. Developmental cell. 2014;30(5):496-508.
38.   Zasadzinska E, Foltz DR. Orchestrating the Specific Assembly of Centromeric
Nucleosomes. Progress in molecular and subcellular biology. 2017;56:165-92.
39.   Bannister LA, Schimenti JC. Homologous recombinational repair proteins in mouse
meiosis. Cytogenetic and genome research. 2004;107(3-4):191-200.
40.   Li W, Ma H. Double-stranded DNA breaks and gene functions in recombination and
meiosis. Cell research. 2006;16(5):402-12.
41.   Takabayashi S, Yamauchi Y, Tsume M, Noguchi M, Katoh H. A spontaneous smc1b
mutation causes cohesin protein dysfunction and sterility in mice. Experimental biology and
medicine (Maywood, NJ). 2009;234(8):994-1001.

42. Murdoch B, Owen N, Stevense M, Smith H, Nagaoka S, Hassold T, et al. Altered cohesin
gene dosage affects Mammalian meiotic chromosome structure and behavior. PLoS genetics.
2013;9(2):e1003241.

43. Rankin S. Complex elaboration: making sense of meiotic cohesin dynamics. The FEBS
journal. 2015;282(13):2426-43.

44. Ben-David U, Siranosian B, Ha G, Tang H, Oren Y, Hinohara K, et al. Genetic and
transcriptional evolution alters cancer cell line drug response. Nature. 2018;560(7718):325-30.

45. Case CM, Sackett DL, Wangsa D, Karpova T, McNally JG, Ried T, et al. CKAP2 ensures
chromosomal stability by maintaining the integrity of microtubule nucleation sites. PloS one.
2013;8(5):e64575.

46. Hudson DF, Amor DJ, Boys A, Butler K, Williams L, Zhang T, et al. Loss of RMI2 Increases
Genome Instability and Causes a Bloom-Like Syndrome. PLoS genetics. 2016;12(12):e1006483.

47. Staples CJ, Myers KN, Beveridge RD, Patil AA, Lee AJ, Swanton C, et al. The centriolar
satellite protein Cep131 is important for genome stability. Journal of cell science. 2012;125(Pt
20):4770-9.

48. Wangsa D, Quintanilla I, Torabi K, Vila-Casadesus M, Ercilla A, Klus G, et al. Near-tetraploid
cancer cells show chromosome instability triggered by replication stress and exhibit enhanced
invasiveness. FASEB journal : official publication of the Federation of American Societies for
Experimental Biology. 2018;32(7):3502-17.

49. Chua MLK, Lo W, Pintilie M, Murgic J, Lalonde E, Bhandari V, et al. A Prostate Cancer
"Nimbosus": Genomic Instability and SChLAP1 Dysregulation Underpin Aggression of Intraductal
and Cribriform Subpathologies. European urology. 2017;72(5):665-74.

50. Pan Y, Tong JHM, Lung RWM, Kang W, Kwan JSH, Chak WP, et al. RASAL2 promotes tumor
progression through LATS2/YAP1 axis of hippo signaling pathway in colorectal cancer. Mol
Cancer. 2018;17(1):102.

51. Geng CH, Zhang CL, Zhang JY, Gao P, He M, Li YL. Overexpression of Sirt6 is a novel
biomarker of malignant human colon carcinoma. Journal of cellular biochemistry.
2018;119(5):3957-67.

52. Negrini S, Gorgoulis VG, Halazonetis TD. Genomic instability--an evolving hallmark of
cancer. Nature reviews Molecular cell biology. 2010;11(3):220-8.

2821 tumor and normal tissues

Stomach Bladder and Breast

Colorectal and cervical

Unstable genomes

Stable genomes

Information-theoretic analysis to identify perturbed gene network in the entire population

networks enriched with meoisis and kinetochore genes

Unstable cancers

Gene expression (%) of meoisis/kinetochore

Stable cancers

Gene expression (%) of meoisis/kinetochore

Experimental overexpression of meoisis and kinetochore genes

Multipolar spindle

10 um

Anaphase bridge

10 um

Meoisis and kinetechore genes induce chromosomal instability

A

B

Breast cancer ■ Bladder cancer ■ Stomach cancer
Cervical cancer ■ Colorectal cancer

10% of Highest CNVs

10% of lowest CNVs

# of unbalanced process

# of unbalanced process

Meiosis genes   Kinetochore genes

Chromosome seggregation

Normal Division

A — Tubulin | DAPI | Merge

Lagging chromosome

B — HCT116 CENPA, HCT116 DMC1, MCF 7 CENPA, MCF 7 DMC1

Anaphase bridge

C — HCT116 CENPA, HCT116 DMC1, MCF 7 CENPA, MCF 7 DMC1

Spindle defects

Bipolar

D — Tubulin | DAPI | Merge

Unipolar

E — HCT116 CENPA, HCT116 DMC1, MCF 7 CENPA, MCF 7 DMC1

Tripolar

F — HCT116 CENPA, HCT116 DMC1, MCF 7 CENPA, MCF 7 DMC1

Chromosome seggregation

G — HCT 116 — Seggregation(%)

H — MCF 7 — Seggregation(%)

Normal division
Anaphase bridge
Lagging chromosomes

Seggregation defects

I — HCT 116 — Seggregation(%)

J — MCF 7 — Seggregation(%)

Equal seggregation
Unequal seggregation

Spindle defects

K — HCT 116 — Spindle (%)

L — MCF 7 — Spindle (%)

Unipolar
Tripolar
Bipolar

NO GENES, CENPA, HJURP, DMC1, SPO11, SMC1B, REC8

HCT116

Misexpression of genes with roles in kinetochore structure or in meiosis contributes to genome instability in cancer

Roshina Thapa[1], Swetha Vasudevan[1], Mimi Abo-Ayoub Ashqar [1], Eli Reich[1], Nataly Kravchenko-Balasha[1*] and Michael Klutstein[1]*

[1] Institute of Dental Sciences, Faculty of Dental Medicine, The Hebrew University of Jerusalem, P.O.B. 12272, Ein Kerem, Jerusalem, ISRAEL 91120

Corresponding Authors: Michael Klutstein and Nataly Kravchenko-Balasha
Email: michaelk@ekmd.huji.ac.il ; natalyk@ekmd.huji.ac.il

* M.K and N.K.B contributed equally to this work

Supplementary files include:

Fig. S1. Amplitudes ($\lambda_1 (k)$) of unbalanced process 1 in each tumor and normal tissues.

Fig. S2. The correlation between over-expression of meiosis/kinetochore genes and genomic instability within stable cancer types

Fig. S3. Overexpression of meiosis and kinetochore genes in HCT116 and MCF-7.

Fig. S4. Immunofluorescent staining showing the cell division defects.

Fig. S5. Immunofluorescent staining showing the spindle defects.

Fig. S6. Meiosis/kinetochore genes induce number and size if the colonies in soft agar

Fig. S7. Soft agar assay analysis showing the number of colonies and area of MCF-7 cell line when overexpressed with meiotic and kinetochore genes.

Figure S1

λ_1 of Breast cancer

Figure S1. Amplitudes ($\lambda_1(k)$) of unbalanced process 1 in each breast tumor (blue dots) and breast normal tissues (orange dots). The most dominant (characterized the biggest number of the patients in the dataset) process 1 distinguishes between cancer (blue dots) and non- cancer (orange dots) tissues. The blue box marks threshold limits. Plots of amplitudes for all unbalanced processes in breast and other cancer types can be found in **Table S2**.

Figure S2

Figure S2. The correlation between over-expression of meiosis/kinetochore genes and genomic instability within stable cancer types. Stable cancer types, cervical cancer (Ce) and colorectal cancer (CC), were analyzed in this section. Each cancer type was analyzed separately as following: Distribution of lCNVs values was generated for each cancer type. 10% of the samples with highest lCNVs values (unstable group) and 10% of the samples with lowest lCNVs values (stable group) were selected in each cancer type. Cancer specific unbalanced processes of 10% of the samples with highest lCNVs values **(A, C)** were compared to the processes appeared in 10% of the samples with lowest lCNVs values (**B, D**). The most dominant processes characterizing two groups were not enriched with meiosis/kinetochore genes.

Supplementary Figure 3 (S3): Meiosis/kinetochore genes were fused to GFP and the over-expression pattern in the cell lines shows that the overexpression was successful. Overexpression of meiotic genes (DMC1, REC8, SMC1B, SPO11) and kinetochore genes (CENP-A and HJURP) in the HCT 116 cell line (**A**) and MCF-7 cell line (**B**) is presented.

Figure S4

Chromosome seggregation

Supplementary Figure 4 (S4): Immunofluorescent staining showing the effect of overexpressed meiosis genes (DMC1, REC8, SMC1B and SPO11) and kinetochore genes (CENP-A and HJURP). The figure shows normal division (**A**) and anaphase bridges (white arrow) in HJURP and SPO11 overexpression (**B**). Scale bar is 10μm.

Supplementary Figure 5 (S5): Immunofluorescent staining showing effect of Overexpressed meiosis genes (DMC1, REC8, SMC1B and SPO11) and kinetochore genes (CENP-A and HJURP) in HCT116 and MCF-7 cell lines. **A&D** represent normal division. Overexpressed genes caused abnormal spindle defects: unipolar spindle formation (**B&E** in HCT116 and MCF 7 cells) and tripolar spindle formation (**C &F** in HCT116 and MCF-7 cells). Scale bar in B is 10 μm.

Supplementary Figure 6 (S6): Meiosis/kinetochore genes induce number and size of colonies in soft agar. HCT116 and MCF7 cells overexpressing the meiosis/kinetochore genes mentioned above were seeded in soft agar. Representative images of soft agar assay for HCT116 (**A**) and MCF-7 cell lines (**B**) are shown. Scale bar 10µm.

Figure S7

Supplementary Figure 7 (S7): A soft agar colony formation assay was applied for the detection of transformed cells when overexpressed with meiotic and kinetochore genes in an unstable cell line (MCF-7). The number of colonies **(Fig A)** and size of the colonies **(Fig B** for full scale or **C** for a zoom in on colonies up to 4000 mm^2**)** were measured using imageJ . Statistically significant differences compared with empty vector (No genes) were determined using a Student's 2-tailed t test (*P < 0.05; **P < 0.01; ***P < 0.001).