Productive lifespan and resilience rank can be predicted from on-farm first parity sensor time series but not using a common equation across farms

I. Adriaens[1,2,3*], N.C. Friggens[4], W. Ouweltjes[5], H. Scott[3], B. Aernouts[1], J. Statham[3]

[1] KU Leuven, Department of Biosystems, Biosystems Technology Cluster, Campus Geel, Kleinhoefstraat 4, 2440 Geel, Belgium

[2] KU Leuven, Department of Biosystems, Division Mechatronics, Biostatistics and Sensors, Kasteelpark Arenberg 30, 3001 Leuven, Belgium

[3] RAFT Solutions Ltd., Mill Farm, Studley Road, Ripon HG4 2QR, United Kingdom

[4] UMR Modélisation Systémique Appliquée aux Ruminants, INRAE, AgroParisTech, Université Paris-Saclay, 16 Rue Claude Bernard, 75005, Paris, France

[5] Wageningen Research, Breeding & Genomics, Droevendaalsesteeg 1, 6708PB Wageningen, The Netherlands

* Corresponding author: Ines Adriaens, ines.adriaens@kuleuven.be, Kasteelpark Arenberg 30 box 2456, 3001 Leuven, Belgium, +32 (0) 16 32 88 73

ABSTRACT

A dairy cow's lifetime resilience and her ability to re-calve gain importance on dairy farms as they affect all aspects of the sustainability of the dairy industry. Many modern farms today have milk meters and activity sensors that accurately measure yield and activity at a high frequency for monitoring purposes. We hypothesized that these same sensors can be used for precision phenotyping of complex traits such as lifetime resilience or productive lifespan. The objective of this study was to investigate if lifetime resilience and productive lifespan of dairy cows can be predicted using sensor-derived proxies of first parity sensor data. We used a data set from 27 Belgian and British dairy farms with an automated milking system containing at least 5 years of successive measurements. All of

these farms had milk meter data available, and 13 of these farms were also equipped with activity

sensors. This subset was used to investigate the added value of activity meters to improve the model's

prediction accuracy. To rank cows for lifetime resilience, a score was attributed to each cow based

on her number of calvings, her 305-day milk yield, her age at first calving, her calving intervals and

the days in milk at the moment of culling, taking her entire lifetime into account. Next, this lifetime

resilience score was used to rank the cows within their herd resulting in a lifetime resilience ranking.

Based on this ranking, the cows were classified in a low (last third), moderate (middle third) or high

(first third) resilience category. In total 45 biologically-sound sensor features were defined from the

time-series data, including measures of variability, lactation curve shape, milk yield perturbations,

activity spikes indicating estrous events and activity dynamics representing health events. These

features, calculated on first lactation data, were used to predict the lifetime resilience rank and thus,

the classification within the herd (low/moderate/high). Using a specific linear regression model

progressively including features stepwise selected at farm level (cut-off P-value of 0.2), classification

performances were between 35.9% and 70.0% (46.7 ± 8.0, mean ± standard deviation) for milk yield

features only and between 46.7% and 84.0% (55.5 ± 12.1, mean ± standard deviation) for lactation

and activity features together. This is respectively 13.7 and 22.2% higher than what random

classification would give. Moreover, using these individual farm models, only 3.5% and 2.3% of the

cows were classified high while being low and vice versa, while respectively 91.8% and 94.1% of

the wrongly classified animals were predicted in an adjacent category. A common equation across

farms to predict this rank could not be found, which demonstrates the variability in culling and

management strategies across farms and within farms over time. The lack of a common model

structure across farms suggests the need to consider local (and evidence based) culling management

rules when developing decision support tools for dairy farms. With this study we showed the potential

of precision phenotyping of complex traits based on biologically meaningful features derived from

readily available sensor data. We conclude that first lactation milk and activity sensor data have the

potential to predict cows' lifetime resilience rankings within farms but that consistency over farms is

currently lacking.

Key Words: resilience; precision phenotyping; prediction model; longevity; precision livestock

farming.

56   **INTRODUCTION**

Increasing the longevity of dairy cows is key for the dairy sector's sustainability. Cows with a

long productive lifespan typically have a good reproductive performance, few health problems and

produce milk in an efficient and consistent way. A dairy cow typically only starts to make profit for

the farmer during her second lactation and she reaches her full production potential as late as in her

third lactation (Cabrera, 2018). Early culling and short longevity thus clearly has a negative impact

on the economic efficiency of the herd. More importantly, longevity and optimizing the ratio between

the productive and non-productive life is also crucial for the fulfillment of societal demands and to

reduce environmental impacts of the sector (van Knegsel et al., 2014).

One step towards the optimization of the farm management with respect to longevity would be

the identification of animals that have a high probability of completing several lactations, or more

specifically, that are 'resilient'. Resilient animals can be considered as animals that avoid early culling

by coping well with the farm's management conditions. These animals reproduce easily, produce

consistently and react well on imposed challenges and (physiological) stress (Ahlman et al., 2011).

Correct and timely identification of these resilient animals would not only allow for optimization of

culling and (advanced) breeding decisions, but also for the long-term selection of animals that thrive

in their specific farm environments (e.g., infection pressure, feed quality, milking conditions, …).

Optimized efficiency on farm would require that the lifetime resilience of an animal can be

predicted as soon as possible. Genetic indicators for lifetime resilience are not yet available as

phenotypic information on this complex trait is lacking. Nevertheless, recent technological

developments led to an increased implementation of sensor systems and automation to improve the

herd management and reduce labor requirements (Steeneveld and Hogeveen, 2015). Besides for the

detection of health problem and fertility events, many of these sensor systems also have the potential

to provide targeted information about other, more complex traits (Friggens and Thorup, 2015). In this

study, we hypothesized that common sensor data such as milk production and activity time series can

be used to predict a complex trait such as lifetime resilience. Simultaneously, additional benefits of

these technologies will be generated from the calculation of precision phenotypes and their use for

the characterization of overall and relative performance of the animals within the farm context and

compared to herd mates (Royal et al., 2000; Tenghe et al., 2015; Sorg et al., 2017). Accordingly,

when sensor data can be used to this purpose, selection of animals on these more complex traits

becomes possible, which, when combined with the genetic merit of each animal, can boost future

breeding efforts at farm and population level (Köning and May, 2019).

In order to use sensor data for the prediction of lifetime resilience such that it can be used for both

decision support and precision phenotyping, we propose to derive biologically meaningful proxies

for the cow's physiological status from the high-frequency milk yield and activity dynamics available

from commercially available sensor systems. It was shown previously that each change in feed intake

or energy allocation (for e.g. an immune response) may result in yield perturbations (Ben Abdelkrim

et al., 2019) and thus, the milk yield dynamics mirror the animal's physiological status. Similarly,

activity dynamics reflect potential estrus and the more general behavioral responses of the cows to

physiological and environmental stress (Rutten et al., 2013). Because of the link between health and

fertility performance and longevity and culling, the proposed concept is that through characterization

of these dynamics, it will be possible to predict their lifetime resilience.

This study aimed at developing meaningful milk yield and activity features from first lactation

sensor time series and combining these features into a model capable of predicting the lifetime

resilience of the cows on each farm. This model could be used to help farmers to identify animals that

cope well with their specific farm contexts, for example to aid breeding (e.g., dam selection for sexed

semen, embryo transfer or the use of a beef sire) or culling decisions as early as after the first lactation.

As such, there is still time to take decisions that directly contribute to the farm's efficiency by

selecting animals which perform well on that particular farm.

MATERIALS AND METHODS

Data Collection and Selection

Available Data. Software back-ups of the farm management system were collected on

respectively 34 and 42 Belgian and British farms with an automated milking system (**AMS**). From

this database, 27 farms were selected based on (1) the accessibility and reliability of at least 5 years

of contiguous data, and (2) the availability of daily milk yield at individual cow level. The time period

covered by these data varied between 2005 and 2019. All these 27 farms had either an AMS of Lely

(Lely Industries N.V., Maasluis, the Netherlands; No. = 16) or of DeLaval (DeLaval International,

Tumba, Sweden; No. = 11). On average 2.4 lactations were recorded per cow's life. All farms had

intensive production systems with cows kept indoors and fed with both forage and concentrates. Other

management practices differed among herds but were not further documented in the software back-

up files of the farm management system.

All data tables were extracted from the restored back-up files of the AMS software system using

SQL Server Management Studio (Microsoft, Redmond, WA, United States). The further data mining,

pre-processing and merging of these data tables and the rest of the analyses described below was done

in Matlab R2017a (The MathWorks Inc., Natick, MA, United States). Both the full data set of 27

farms all having daily milk records (data set 1, **DS1**) and a subset of 13 farms also having daily

activity data available (data set 2, **DS2**, all milked by a Lely AMS) were used for this study. An

overview of the characteristics of both data sets is given in Table 1.

Cow Selection. After extraction of data tables, individual cows on each farm were selected based

on the availability of sensor data for their entire lifetime production. Because exact culling dates were

not always available, we elected to apply a criterion to discriminate between cows that likely had

been dried off towards the end of the time span covered by the dataset available for that farm (not to

be included in the analysis) or removed from the herd (to be included in the analysis). For each farm,

the 95% confidence interval on the average dry period length was calculated. If the last milk record

was before the end of the data set minus the upper 95% confidence interval boundary, that cow had

97.5% chance that she was removed from the herd and she was included. Accordingly, only cows

that met this criterion and for which the date of her first calving was within the timespan of the

available data for that farm were selected. An overview of the characteristics of this selection is

provided in Table 1. Data set 1 consisted of 3754 unique cows and 9395 unique lactations, while DS2

included 2075 cows with 5286 lactations. Per farm, respectively 24 to 308 cows (139 ± 82, mean ±

SD) with in total 44 to 799 lactations (348 ± 229) and 57 to 308 cows (160 ± 84) with 113 to 799

lactations (407 ± 264) were selected for DS1 and DS2.

Sensor data. The milk yield sensor data were recorded by the AMS using ICAR-approved milk

meters as integrated in the Lely and DeLaval robots. The available activity sensor data were recorded

by Lely (Lely Industries N.V., Maasluis, the Netherlands) neck-mounted activity sensors and

consisted of raw 2-hourly measures of acceleration in x, y and z planes that were aggregated in single

daily activity records, but no further details of the individual sensor systems were available. Although

in this study the raw data were all similar with 2-hourly values varying between 0 and 300, the

presented methodology is independent of the actual raw values and can be applied on all measures of

activity for which daily aggregates are available as long as they represent the behavioral changes of

the cows linked with estrous and health events.

Calculation of Lifetime Resilience Ranking

For this study, the lifetime resilience of a cow was considered primarily as the cumulative result

of her ability to recalve (and thus, to extend her productive lifespan) supplemented with secondary

corrections for age at first calving, calving intervals, 305-day milk yield, health events and number

of inseminations (Friggens and De Haas, 2019). With a high weight given to each newly started

lactation (i.e. parity number), the secondary corrections mainly allow discrimination between all cows

reaching a certain parity. This definition was agreed upon within the EU Horizon 2020 GenTORE

consortium consisting of researchers, animal experts, veterinarians, technology suppliers and

geneticists and is further detailed in Friggens and De Haas, (2019). Because the number of

inseminations and health events were not consistently available for all herds over the entire time

period, the final equation for calculating lifetime resilience scores (**RS**) excluded these variables and

was (Eq. 1):

$RS_i = \overline{CI}$ (Eq. 1)

$+300 * L_i$

$+(730 - AFC_i)$

$+\sum_{j=1}^{L_i-1} \left(\overline{CI}_j - CI_{i,j} \right)$

$+\sum_{j=1}^{L_i} \left(\frac{\sum_{k=1}^{\max(305, DIM_{i,j})} MY_{i,j,k}}{\sum_{k=1}^{\max(305, DIM_{i,j})} \overline{MY_{i,J,k}}} - 1 \right) * 100$

$+ \min(0, (DIM_{i,L_i} - 100))$

With:

RS_i Lifetime resilience score for cow *i*

$\overline{CI}$ Average calving interval of the herd

L_i Lactation number in which cow *i* exited the herd (last lactation number of a

172   cow)

173   AFC_i    Age at first calving of cow i in days

174   $CI_{i,j}$    Calving interval of cow i between the start of lactation j and $(j+1)$

175   $\overline{CI}_j$    Average calving interval between the start of lactation j and $(j+1)$ of all cows

176      in the herd

177   $MY_{i,j,k}$    Milk production (in kg) of cow i at day k of lactation j

178   $\overline{MY_{i,j,k}}$    Average milk production (in kg) at day k of all cows in the herd in lactation j

179   $DIM_{i,j}$    Days in milk (**DIM**) of cow i at the end of her lactation j

180   DIM_{i,L_i}    Days in milk of cow i at the end of her last lactation Li

182   This way, each RS consists of (1) a baseline equal to the average calving interval of that herd to avoid

183   negative lifetime resilience scores (this does not contribute to the ranking); (2) a bonus of 300 points

184   given for each recalving (newly started lactation); (3) a penalty or bonus score given to cows older

185   respectively younger than 24 months at their first calving equal to 1 point per day longer or shorter

186   than 730 days (i.e. 24 months); (4) a penalty or bonus score equal to the number of days the calving

187   interval is respectively shorter or longer than the average calving interval of the same parity in the

188   herd; (5) a penalty or bonus score equal to the percentage the 305-day milk production is respectively

189   lower or higher compared to the average 305-day production of the corresponding parity for all

190   lactations in the herd, reflecting production performance in the most relevant part of the lactation;

191   and (6) a penalty score equal to 100-DIM_{exit} for cows exiting the herd before day 100 in lactation

192   assuming that these cows are involuntarily removed from the herd.

193    Based on this score, which assumes that these factors reflect the accumulated effects of the cows'

194   resilience over their lifetime, cows were ranked within farms. For example, if the average CI of a

195   herd is 400 days, the average CI between the start of first and second parity is 380 days and the

196   average 305-day milk production in first lactation is 8000 kg, then a cow that calved twice (the first

197   time aged 775 days), had a calving interval of 420 days between the start of the first and second

lactation, produced 5% more milk than average in the first 305 days of the first lactation (8400 kg)

and 20% less than her herd peers in the first 90 days of the second lactation, and that was culled day

90 in the second lactation will receive a lifetime resilience score of $RS = 400 + 300 * 2 + (730 - 775)$

$+ (380 - 420) + 5 - 20 + (90 - 100) = 890$ points. Because the weights in the lifetime RS were

arbitrarily chosen and based on expert knowledge, and the main interest is to distinct the lowest from

the highest resilient animals, the RS was converted into an on-farm resilience rank (**RR**) to rank the

cows for their lifetime resilience performance. This way, the exact weights or points assigned to each

variable in Eq. 1 are of less importance. In the final lifetime RR, high ranked cows ('highly resilient

animals') represent animals recalving many times, having the (theoretically) optimal age at first

calving, having short calving intervals (and thus good reproductive performance), and producing

proportionally more milk compared to their herd mates. The number of lactations affects this ranking

the most because of the high weight for each new lactation started. Before entering the lifetime RR

in the models, it was scaled for each farm using $RR_{scaled} = (RR\text{-}RR_{min})/(RR_{max} - RR_{min})$ with RR_{min}

equals 1 and RR_{max} equals the maximum rank (equivalent to the number of cows included in the

ranking for that farm). The resulting RR_{scaled} varied between 0 (i.e. the highest ranked cow) and 1 (i.e.

the lowest ranked cow), and thus, no scale effects caused by the varying number of animals included

per farm would influence the prediction models. In the rest of this manuscript, the RR refers to the

scaled RR.

Sensor Features (SF)

Milk Yield SF. The time-series data of two sensors were included in this study: (1) milk meter

sensors from which daily milk yields were calculated and (2) activity sensors from which the two-

hourly raw data were aggregated into daily activity records. Sensor features were calculated for each

of the cows for which the first lactation was longer than 200 days, because 200 days is enough to

grasp a good image of the time-series dynamics as the second part of the lactation curve after the peak

can be estimated by a linear function (Wood, 1967). Only the data of the first 305 days of the first

lactation were included for the calculations.

As explained more in detail in Appendix A, in total 30 milk yield SF were calculated based on

the daily milk yield dynamics. Moreover, a methodology was developed to calculate SF from the

dynamics of the lactation curves using both the theoretical shape and the deviations from this

theoretical shape as proxies for the cow's physiological status. Accordingly, SF were defined in the

following categories: (1) lactation shape characteristics including peak yield, consistency, days in

milk of peak, etc.; (2) goodness-of-fit and variability measures including the characteristics of

lactation model residuals; and (3) perturbation features characterizing the disturbances in the lactation

dynamics and including the development and recovery rates, the number of perturbations, etc. To

determine the theoretical shape of the lactation curve (i.e., potential production when no perturbations

are present), a simple lactation model was iteratively fitted such that perturbations were excluded and

thus, did not influence the lactation model's coefficients (Adriaens et al., 2018). The chosen model

was the nonlinear Wood model: $\boldsymbol{TMY = A*e^{-B*DIM}*DIM^C}$ with *A, B* and *C* being the model's

coefficients, *TMY* the total daily milk yield in kg and *DIM* the days in milk expressed in days (Wood,

1967). The Wood model (gamma function) describes the lactation curve with an increasing phase, a

peak and an almost linear decreasing phase, describing the overall lactation dynamics with only three

coefficients. Its simplicity reduces the computational power needed when repeatedly fitting the

nonlinear model, and therefor this equation was preferred over more complex lactation models. In

each iteration, the residuals were calculated by subtracting the fitted Wood model from the milk yield

data. Next, all the residuals smaller than 85% of the theoretical curve (i.e. Wood's model) were

removed and the model was refitted in a next iteration. This procedure was repeated for each lactation

curve individually until the difference of the average root mean squared error (**RMSE**) between two

iterations was smaller than 0.10 kg for that curve, or for at most 20 iterations. The final model

coefficients represent the lactation shape when no perturbations would have been present, and the

model's residuals reflect the perturbations and 'unexpected' milk yield dynamics. An example of the

daily milk yield data, the iterated Wood model and the corresponding residuals is shown in Figure 1.

Next, lactation SF were calculated from the final coefficients of Wood's model (A, B and C), the

residuals of all daily milk yield records and the periods identified as perturbations. For the latter,

major events (i.e. periods of at least 10 days of successively negative residuals with at least one day

of milk production lower than 80% of the theoretical production) were discriminated from minor

events (i.e. periods of at least 5 days of successively negative residuals with at least one day of milk

production between 90% and 80% of the expected production). It was assumed that large

perturbations (major events) may represent severe health problems which might influence culling and

rebreeding decisions, while smaller (minor) perturbations are more probably linked to chronical or

subclinical infections, with a different effect on culling or longevity. A detailed description of the

milk yield SF and how they were calculated can be found in Appendix A.

Activity Sensor Features. In addition to the lactation SF, for DS2 also activity SF were calculated

from the daily aggregated raw activity measures. Fifteen different SF were defined in the following

categories: (1) features related to the absolute (within-herd) levels, i.e. variability and autocorrelation;

(2) fertility-related characteristics based on short spikes representing estrous behavior; and (3) overall

activity-related characteristics based on changes in average activity during longer periods of time

which possibly relates to e.g. health events. To identify the short spikes of the second category, a

median smoother using a window of 4 days was used and subtracted from the raw daily activity data

to obtain residual activity levels. A short spike was identified as an increase above 40% of the

maximal residual. For the identification of the longer-term patterns in the data, a 20 day-window

median smoother was applied and subtracted from the daily activity data to obtain the residuals. A

threshold of 20% of these minimal and maximal activity residuals was set to identify changes in

activity of several days compared to the previous period. The details for the activity SF calculations

are given in Appendix B. Although all the raw values in DS2 had similar variability and magnitude

(i.e., with 2-hourly measures between 0 and 300 and similar longitudinal patterns), the developed

procedure can be applied on all sorts of activity data, also those originating from other types of sensors

independently of the exact activity levels, as long as daily measurements are available.

Standardization of the SF. The mean and SD differed across SF. For example, days in milk of

peak varied between 10 and 150 days, while the lactation persistency comprised values between 0.001

and 1. When using the SF as variables in a prediction model, this can cause an imbalance in the fitting

and selection procedure and it limits the interpretation of the model regression coefficients. Therefore,

standardization of the SF was needed. Before entering the SF in the models, each SF was standardized

within herd using mean centering (i.e., subtracting the within-herd mean of that SF) and by dividing

them by the within-herd SD. Accordingly, the standardized SF within a herd have a mean of zero and

a SD of one, which thus corrects for differences in their order of magnitude and solves interpretability

issues. This standardization step ensures that a higher absolute value of a model coefficient indicates

a larger effect on the model outcome (i.e., the lifetime RR). Standardized SF (both milk yield and

activity) with values smaller than -3 or higher than 3 (mean $\pm$ 3*SD) were considered as outliers and

replaced by zeros (i.e., the average value) to avoid missing and unbalanced data. The objective was

to develop a tool to evaluate and forecast the (phenotypic) performance of an animal in the herd early

in her productive life to still have time to take breeding decisions that would directly contribute to the

farms' performance, and so that 'high risk' animals could be monitored closer. Therefore, in this

study only the first parity SF were taken into account as proxies for performance, health and fertility

to predict their lifetime resilience and recalving ability on farm.

Exploratory Analysis

In this study, a model was sought to predict the lifetime RR of all the animals on a specific farm.

Ideally, a common model structure that is valid for all farms would be obtained, as this would allow

the calculation of a limited and universal number of SF indicative for the animals' lifetime resilience.

As a first step to evaluate the consistency between the SF and the lifetime RR across farms, the

Pearson linear correlation coefficient between each SF and the lifetime RR at individual farm level

was calculated. High positive and negative correlations would indicate a strong effect of that SF on

the lifetime RR, and thus a potential candidate for inclusion in further prediction models.

In a second step, mutual correlations between the SF were explored for all farms together. This

initial data exploration using data of all farms together pointed out some significant (but small) linear

correlations between the SF. However, at individual farm level, these correlations were often

inconsistent and the sign of the correlations differed between farms. To investigate whether an

underlying latent structure existed in the SF and avoid future multi-collinearity in the prediction

models, a principal component analysis was carried out on the SF of both DS1 and DS2. These

principal component analyses showed that respectively 8 and 24 principal components with

eigenvalues higher than 1 (Kaiser criterion) explained only 71% and 74% of the variance, suggesting

that a latent structure for data reduction over all farms did not exist.

Individual Farm Model Development

Several multivariate modelling techniques including partial least squares and general linear

mixed models were tested, but all had poor prediction performance (with classification results equal

or worse than what random classification would have given, i.e., less than a third correctly classified)

or showed significant overfitting of the data. Ultimately, a separate multivariate linear regression

model relating the SF to the lifetime RR within farm was constructed as follows (Eq. 2):

$$RR_{scaled} = \beta X + \varepsilon \qquad \text{(Eq. 2)}$$

With RR_{scaled} being the scaled lifetime RR between 0 and 1 as defined above. The β vector contains

the regression coefficients for the standardized SF in the design matrix X, and ε are the residual errors.

A backward stepwise regression procedure was used to identify redundant SF in X applying a P-value

of 0.2 as the inclusion threshold. This means that if there was a probability of more than 20% that

removing the SF had a significant deteriorating effect on the prediction model, it was included. The

chosen threshold might seem uncommonly high but given the high variability in the SF both between

and within farms, we deemed relevant to include any feature having a tendency towards significance.

Ten-fold cross-validation (**CV**) was performed to evaluate the prediction performance of the

obtained models and identify overfitting. To this end, ten times all the cows of each farm were

assigned to either the calibration (90% of the animals) or validation (10% of the animals) set using

random sampling from a uniform distribution, but applying the additional criterion that both the

calibration and the validation set contained at least one animal ranked in the highest third, one in the

middle third and one in the lowest third of the ranked cows. In each CV-cycle, the cows in the

calibration set were used to estimate the regression coefficients β and the obtained model was used

to predict the RR_{scaled} of the cows in the validation set. The average prediction results over all ten CV-

cycles were considered to represent the final model performance.

Model Evaluation

The initial model fit at farm level was evaluated using the RMSE (i.e., the $RMSE_{TR}$) and the

R^2_{adj}, calculated as shown in equation 3 (Eq. 3):

339
$$R^2_{adjusted} = 1 - \frac{\left(1-(1-\frac{SSE}{SSTO})\right)(n-1)}{1-(k-1)}$$
(Eq. 3)

With SSE = residual sum of squares of the regression, SSTO the total sum of squares (i.e. the mean

value of the outcome RR_{scaled}), n the number of data points of each farm and k the number of SF

retained in the final model for that farm.

To evaluate classification performance and discriminate between high and low resilient cows

(which is of practical relevance), the cows of each farm were divided into three different categories

based on their ranking: high (**H**; top third), moderate (**M**; middle third) and low (**L**, bottom third)

resilient animals. When the predicted RR_{scaled} did not cover the full range of 0 to 1, and to be able to

calculate these high, medium and low ranked categories for each farm, a farm-individual correction

factor was applied on the predicted RR_{scaled} scores as follows (Eq. 4):

$$\widehat{RR}_{scaled,corr,i} = \frac{\widehat{RR}_{scaled,i} - A}{B - A} \qquad \text{(Eq. 4)}$$

With $\widehat{RR}_{scaled,i}$ being the predicted RR_{scaled} of the i^{th} cow and A and B are farm-specific coefficients

representing respectively the minimum and maximum of all the predicted RR_{scaled} for that farm in the

calibration set of each CV-cycle. The RR_{scaled} of the cows in the validation set of each CV-cycle were

predicted using each individual farm model (Eq. 2) and their category (H, M, L) was determined after

applying the correction using the farm-specific coefficients (Eq. 4). Both the root mean squared error

of cross-validation (**RMSECV,** Eq. 5) and the classification accuracy were evaluated in this CV to

assess the models' prediction performance.

$$RMSECV = \sqrt{\frac{1}{N}\sum_1^N\left(RR_{scaled,i} - \widehat{RR_{scaled,i}}\right)^2} \qquad \textbf{(Eq. 5)}$$

To evaluate whether a common model structure across farms could be identified or whether

specific features are highly correlated to lifetime resilience in all farms, we evaluated the overlap in

retained features for each farm, both in terms of their inclusion or exclusion in each farm-specific

model and the sign of their regression coefficients.

Prediction performance improvement of the models including and excluding activity features

was assessed using a one-sided paired t-test on the percentage correctly classified using the null

hypothesis "*activity features do not improve (i.e. increase) the percentage correctly classified*

animals" and on the proportion oppositely classified using testing the null hypothesis "*activity*

features do not improve (i.e. decrease) the percentage oppositely classified animals".

RESULTS AND DISCUSSION

Lifetime Resilience

This study investigates the possibility to predict 'lifetime resilience' from SF of the first lactation.

Currently, there is no consensus on what 'resilience' exactly is, and definitions found in literature

include the 'ability to maintain performance regardless of pathogen burden' and 'the adaptation

ability to a broad range of environmental conditions' (Mulder and Rashidi, 2017; Köning and May,

2018). Our approach differs from these definitions and considers lifetime resilience as the cumulative

effect of good health and fertility and a high adaptability to challenges, resulting in long productive

life spans.

Figure 2 shows the lifetime resilience score plotted against the lactation in which the cows were

culled for one farm as an example. Each circle represents one animal on the farm, and a higher

resilience score also means a higher lifetime resilience ranking. The animals on this example farm

exited the farm in between their first and seventh lactation. This figure confirms that in general for

our definition of resilience, the total number of lactations each cow has started has the biggest impact

on the final lifetime RR. The cows ranked lower than their herd mates with higher last lactation

numbers are seen as spikes. These are mainly cows that are removed from the herd immediately after

calving, and thus for which the penalty given for exiting before day 100 in lactation has a large effect

(DIM_{i,L_i} in Eq. 1).

The inclusion of the 305-day yield originates from the idea that to differentiate two animals

exiting the herd after the same number of lactations, the one with the highest production in the first

305 days (and thus, probably not encountering severe health events) would probably be the more

resilient one. Ideally penalties or bonus points for health events and the number of inseminations

would also be included, but the health and insemination records were not sufficiently complete over

the whole time period for all farms. Consistent and correct registration, collection, mining and storage

of data remains a challenge in the development of on-farm applications (Hudson et al., 2018).

Sensor Feature Definition and Overview

The present study shows an example of how real-farm high-frequency sensor data can be used

beyond monitoring and detection applications (Boichard and Brochard, 2012). We used the

longitudinal, high-granularity milk yield and activity data of respectively 27 and 13 commercial dairy

operations with an AMS to calculate biologically meaningful features of the cows. This is a unique

data set, not only because of its commercial nature (as opposed to research farm data), but also

because these high-frequency time series allowed for the inclusion of the *dynamics* of milk yield and

daily activity. The availability of at least 5 years of successive measurements per farm uniquely

permitted us to study the accumulated effect of health and fertility traits of many animals over their

entire lifetime.

Today, cows exit the herd for many different reasons, for which the most common are poor

reproduction performance, udder health problems, metabolic disorders in early lactation and claw

health and locomotion disorders (Ahlman et al., 2011; Santos et al., 2016). The SF were calculated

starting from expert knowledge and biological hypotheses on the supposed effect of these culling

reasons on the sensor time series, and included features characterizing the perturbations and dynamics

of the lactation and activity curves. For example Elgersma et al., (2018) showed that fluctuations in

milk yield can reflect the cow's health status. Table 2 gives a summary of the most important SF over

all farms for DS1 and DS2. As can be seen from the minimum and maximum value of each SF, some

of these SF have extreme values which upon further investigation appear to result from erroneous

calculations rather than from real deviating curves. These outliers were set to zero for the analysis.

Both the milk yield and the activity features were defined using biological knowledge of how

these time series typically respond to changes in health and nutritional status (Højsgaard and Friggens,

2010; Codrea et al., 2011; Bjerre-Harpoth et al., 2012). We thereby always started from the data-own

baseline, which was identified using a median smoother that is insensitive to sudden changes or

differences in absolute value or baselines. Through characterization of the residuals from this median

smoother and the correction and standardization at herd level, group changes and differences in the

intensity of the responses between herds were corrected for.

Predicting Lifetime Resilience Ranking from Milk Yield Features

Pearson Linear Correlations. The Pearson linear correlation coefficients between the lactation

SF and RR are shown in Figure 3. A high (positive or negative) correlation between a SF and the RR

suggests that there is a large effect of the SF on the RR. The average correlations per SF over all

farms varied between ρ = -0.134 and ρ = +0.1492, and for some of the farms individual features

showed correlations of more than +/- 0.4. Visual exploration of the correlation scatter plots (results

not shown) did not show non-linear relationships either. Although consistency would have been

expected, Figure 3 already suggests that there is only little consistency across farms in which SF can

be predictive for the lifetime RR of the cows of a particular farm. For example, some of the features

represent the effect of health events on the milk yield data through the characterization of

perturbations. Because one could imagine that severe health events and the associated losses

negatively affect longevity on all the farms, we supposed that there would be a consistently high

correlation between perturbation-related SF and the RR. However, this consistency is lacking,

especially in terms of the correlations' sign (negative vs. positive). The highest and most consistent

correlations are obtained for SF representing model fit and size of the residuals (RMSE of the Wood

model (SF2), number of residuals below 85% of the predicted value (SF23) and average size of the 3

largest negative residuals (SF24)).

Model calibration. In the stepwise procedure for selecting the SF that are included in the

multilinear regression model, the best possible combination of SF to fit the RR on each farm is

determined, and SF are included or excluded dependent on whether they improve the model fit. The

cross-validation step reveals whether the final model structure overfitted the training data and whether

the selected SF are indeed meaningful for predicting the RR. If a SF is included in the multilinear

regression model, the absolute value of its regression coefficient is directly related to its effect because

of the standardization procedure of the SF (mean centering and equalizing the variance).

The R^2_{adj} of the individual multivariate linear regression models of each farm varied between

0.03 and 0.61 (0.22 ± 0.16, mean ± SD) and the RMSE$_{TR}$ was between 0.17 and 0.27 (0.23 ± 0.03,

mean ± SD). Between 2 to 12 milk yield features were retained, and all SF were included at least once

in one of the models of the individual farms (Table 3). The SF most often retained in the models were

associated with the goodness-of-fit of the estimated Wood curves (SF21, 22, 25, 27, 28, 29), the size

and number of perturbations (SF13, 14, 15), and their associated milk losses (SF11, 12, 18). This

suggests that the SF that are proxies for (subclinical or chronic) health events are most informative

over the RR, and thus these health events influence the cow's longevity. It appears that farmers do

take the health of the cows into account when making culling and re-insemination decisions, although

not consistently and not on all farms. Moreover, the effects found in this study are rather weak, and

in some cases it can be assumed that only the combined result of different features or only the extreme

values might influence productive lifespan and RR. For example, a cow with severe clinical mastitis

is likely to show a large and sudden drop in milk yield (Rajala-Schultz et al., 1999; Gröhn et al., 2004;

Andersen et al., 2011) while a cow with subclinical mastitis may have a lactation curve which can be

well modelled with a lactation model and which appears to be normal. Both can have impact on

reproduction performance and the probability of culling (Lavon et al., 2010; Wathes, 2012;

Wolfenson et al., 2015), but mathematically capturing the differences without additional health or

treatment information is not possible. General lactation curve characteristics such as peak height and

peak DIM, slopes, rate of the increasing phase of the lactation and persistency of the lactation after

the peak) were only included in the models of only 12 out of 27 farms, suggesting that for example

having a high milk production in the first lactation compared to herd mates barely affects the RR (and

thus the ability to recalve and longevity). The variability between farms is demonstrated again by the

fact that the regression coefficients were consistently above or below zero for only 2 out of the 30

SF, and so only 2 SF had a consistently positive or negative effect on the RR.

Cross validation. With the CV, the repeatability, generality and overfitting of the models was

tested. The within-farm CV showed similar performances (RMSEC $= 0.24 \pm 0.03$) compared to the

$RMSE_{TR}$ of the initial models with all animals included, indicating that models were not overfitted.

On average $46.7 \pm 8.0\%$ of the animals were classified in the correct H, M, L category, and on average

$4.4 \pm 3.5\%$ of the cows were classified 'high' where they should have been 'low' and vice versa. Also

here the large differences in model performance between the different farms stands out, with a range

of correctly classified cows between35.8 and 70.0%, and the range of oppositely classified cows

between 0.0 and 16.5%. When looking deeper into which cows are predicted in the right category, it

was found that the models do not solely predict animals exiting the herd after the first lactation

correctly but also cows exiting the herd in a later lactation. For cows culled already in the first

lactation, health issues are expected to be the major reason (Pinedo et al., 2010). For cows culled at a

later stage, it might not be expected that first lactation features are predictive for the RR unless these

characteristics are highly repeatable over time or represent chronic or repeating conditions that fail to

cure. However, information on the exact reason and timing of culling of the cows could not be taken

into account in the model as this information was not available in the data sets. As it is expected that

this information can contribute to the model, future research should focus on solving this

multidimensionality issue and developing new ways to take these complex interactions into account.

Predicting Lifetime Resilience Ranking from both Yield and Activity Features

Pearson Linear Correlations. Data set 2 consisted of 13 farms for which, besides milk yield

features (No. = 30 features), activity features (No. = 15 features) could also be calculated. These

features included both general characteristics of the daily activity (skewness, variability, absolute

daily level) and specific features associated with short term and longer period activity changes. Farm-

individual Pearson correlation coefficients (Figure 4) between the activity features and the RR varied

between $\rho = -0.41$ and $\rho = +0.44$ and only the number of short activity peaks in (SF34) was

consistently associated with a higher ranking (lower number of peaks is associated with a higher

resilience, $\rho = 0.29 \pm 0.10$, [0.16 ; 0.44]). Several other activity features also had correlations nearly

consistently above or below zero, but these correlations stayed relatively low on average.

Model Calibration. The stepwise linear regression models included 6 to 24 SF (both activity and

milk yield features) and had R^2_{adj} values between 0.2 and 0.76 and $RMSE_{TR}$ values between 0.128

and 0.24. The number of activity features retained in the final models was between 2 and 10, so the

activity sensors seemed to be of added value for all of the farms in predicting their RR. Including

activity features gave a higher R^2_{adj} and a lower $RMSE_{TR}$ in the calibration, while the number of

features retained was sometimes higher and sometimes lower. Again, there was very little consistency

over the different farms in which features were included in the final models. None of the SF was kept

in the model of all farms. The number of activity peaks and DIM of the first peak were retained most

often (respectively 8 and 11 out of 13 times) and with a consistently positive regression coefficient

(respectively 0.036 to 0.122 and 0.042 to 0.134). Three of the SF (6.6%) were never retained in any

of the individual farm models.

Cross validation. The CV, using the same cross-validation sets for these farms as in DS1 (i.e.,

the same animals were included in each set), showed reasonable performance, with an RMSECV of

0.22 ± 0.03 (range 0.15 to 0.26). On average $55.5 \pm 12.1\%$ (range 43.5 to 84.0%) of the cows were

predicted in the correct category (H, M or L) and $2.3 \pm 2.1\%$ (range 0.0 to 6.7%) of them were

predicted 'high' where they were actually 'low' or vice versa. This means that from the wrongly

classified animals respectively 91.8% and 94.1% were predicted in an adjacent category. Over all the

farms, including activity features improved the correct classification with $9.3 \pm 7.9\%$ ($P < 0.01$). The

classification worsened in only two farms compared to when only milk yield features were included.

The proportion classified in the opposite category decreased with on average $3.5 \pm 4.5\%$, ranging

from 5.9% to 2.3% (significant difference, $P < 0.01$).

Despite the variability between and within farms and the fact that we could not find SF that were

commonly informative to predict RR over all farms, the prediction and classification performance of

the individual farm models was in many cases significantly higher than the product of a random

classification (i.e., a third correctly classified). Furthermore, including the activity features

demonstrated a significant added value compared with using the daily milk yield features alone ($P <$

0.01). A correct classification of up to 84% of the animals suggests that at least part of the variability

in the RR is correctly captured by the SF.

One way to explain the lack of a common model structure and the observed differences in

prediction performance is the variability in culling, reproduction and health management between

farms and even within farms. For example, management practices might have differed over the

considered time span because of changing motivations and preferences of the farm staff, the economic

context, the animals' genotypes and phenotypes, the farm facilities, the feed etc. Besides the time-

varying component, the following factors can also explain part of the limited prediction performance

on some farms: (1) for this study, only features of the first lactation were included to ensure

applicability of the model for decision support; (2) the lifetime resilience ranking is based on the

limited data available in the commercial situation and was defined by experts; (3) there is a large

difference in the number of animals included per farm, possibly affecting the results.

Model Use and Implications

Predicting Lifetime Resilience Ranking Supports Breeding and Culling Decisions. Reliable

prediction of lifetime resilience within a farm would allow for a more consistent approach to the

management actions concerning advanced breeding (e.g., sexed semen, embryo transfer, ovum pick-

up, use of beef semen, or selection of animals not to breed the replacement heifers from) or culling

decisions after the first lactation (Mapletoft and Hasler, 2002; Vandeweerd et al., 2012; Boichard et

al., 2015). In this way, breeding decisions for cows in the second parity and higher could be taken

using both the genetic or genomic (available once the animal is born) and the phenotypic sensor-

derived information (once an animal completed her first lactation). The latter would provide

information on how well the animal performs in her specific farm environment, which optimizes

sustainable productivity from the *available* animals on farm. In practice, discrimination between cows

with high and low lifetime resilience would benefit a farmer even when the exact rankings remain

unknown because farmer's decision would not generally be different for e.g., the 5th or the 10th ranked

cow in the herd. Moreover, prediction of lifetime resilience also allows for the identification of

animals with a low expected RR. These cows can be targeted for more detailed monitoring in higher

lactations.

This study also advocates the need for evidence-based decision making on modern dairy farms

supporting more economically-sound and sustainable management actions. Despite the high

prediction performance on some of the farms, the lack of a common model structure and the low

performance on other farms suggest that further data-based rationalization of decisions is needed. In

order to do so, dedicated data processing in which the biology of the cows within their farm contexts

is taken into account is essential, and ideally also other key farm context indicators should be included

in the resulting tools (e.g., herd demographics, robot or parlor capacity, economic environment, etc.).

Model use beyond decision support. With the collection and re-evaluation of this sensor-based

information over many years, also general phenotypic information on complex traits for future

breeding goals is collected at herd level. From this, sires that perform well under many different

environmental conditions can be identified. In this way, the proposed tool can be used in the context

of precision phenotyping of traits that are the combined result of physiological well-being and

performance, and future genetic selection based on these new traits becomes possible.

CONCLUSIONS

In this study, we demonstrate that resilience ranking and productive lifespan of modern dairy cows

on AMS farms in Belgium and the UK could be predicted using farm-individual models based on

first lactation sensor data. With the milk yield and activity SF selected at farm level, we reached

classification performances (low, moderate, high resilience) of up to 84%, while only 2.3 ± 2.1%

(mean ± SD) of the cows were predicted in the opposite category. This shows the potential of high-

frequency milk yield and activity sensor data to rationalize evidence-based breeding and culling

decisions. However, a common model structure across all farms could not be found, which shows the

variability between farms and which highlights the need for biologically sound and context-dependent

data processing tools. Once a lifetime resilience-predicting tool is established, the farmer and the

livestock sector could not only benefit from it for management and decision support, but also at

genetic level in the context of new precision phenotyping proxies for complex traits.

ACKNOWLEDGEMENTS

Ines Adriaens received funding from the Research Foundation Flanders through grants No.

11ZG916N and V423719N and from a KU Leuven postdoctoral mandate grant No. PDM/19/132.

The data from this study were collected in the format of farm software back-up files by the authors

and the resulting database used is owned by the authors. Part of the data were collected in the context

of the VLAIO LA-trajectory 'MastiMan', grant No. HBC.2016.0774 by DVM Igor Van den Brulle

and dr. DVM Sofie Piepers from the University of Ghent, Flanders (Department of Reproduction,

Obstetrics and Herd Health, M-Team and Mastitis and Milk Quality Research Unit). This work is

part of the GenTORE project that has received funding from the European Union's Horizon 2020

research and innovation program under grant agreement No. 727213. Also Katherine Lumb (RAFT

Solutions Ltd., Ripon, United Kingdom) contributed to the data collection. We thank dr. Carmen

596   **REFERENCES**

Adriaens, I., T. Huybrechts, B. Aernouts, K. Geerinckx, S. Piepers, B. De Ketelaere, and W. Saeys.

2018. Method for the short-term prediction of milk yield at the quarter level to improve udder

health monitoring. J. Dairy Sci. 101:10327-10336. doi:10.3168/jds.2018-14696

Ahlman, T., B. Berglund, L. Rydhmer, and E. Strandberg. 2011. Culling reasons in organic and

conventional dairy herds and genotype by environment interaction for longevity. J. Dairy Sci.

94:1568–1575. doi:10.3168/jds.2010-3483.

Andersen, F., O. Østerås, O. Reksen, and Y.T. Gröhn. 2011. Mastitis and the shape of the lactation

curve in Norwegian dairy cows.. J. Dairy Res. 78:23–31. doi:10.1017/S0022029910000749.

Ben Abdelkrim, A., L. Puillet, P. Gomes, and O. Martin. 2019. Lactation curve model with explicit

representation of perturbations as a phenotyping tool for dairy livestock precision farming

661249. PCI for animal science. doi:http://dx.doi.org/10.1101/661249.

Bjerre-Harpøth, V., N.C. Friggens, V.M. Thorup, T. Larsen, B.M. Damgaard, K.L. Ingvartsen and

K.M. Moyes. 2012. Metabolic and production profiles of dairy cows in response to decreased

nutrient density to increase physiological imbalance at different stages of lactation. J. Dairy Sci.

95:2362-2380. doi: http://dx.doi.org/ 10.3168/jds.2011-4419.

Boichard, D., and M. Brochard. 2012. New phenotypes for new breeding goals in dairy

cattle. Animal. 6(4):544-550. doi:10.1017/S1751731112000018

Boichard, D., V. Ducrocq, and S. Fritz. 2015. Sustainable dairy cattle selection in the genomic era. J.

Anim. Breed. Genet. 132:135–143. doi:10.1111/jbg.12150.

Cabrera, V.E. 2018. Invited review: Helping dairy farmers to improve economic performance

utilizing   data-driving   decision   support   tools.   Animal   12:134–144.

doi:10.1017/S1751731117001665.

Codrea, M.C., S. Højsgaard and N.C. Friggens. 2011. Differential smoothing of time-series

measurements to identify disturbances in performance and quantify animal response

characteristics: An example using milk yield profiles in dairy cows. J. Anim. Sci. 89:3089-3098.

doi:10.2527/jas.2010-3753.

Elgersma, G.G., G. de Jong, R. van der Linde, H.A. Mulder. 2018. Fluctuations in milk yield are

heritable and can be used as a resilience indicator to breed healthy cows. J. Dairy Sci.

101(2):1240-1250. doi:10.3168/jds.2017-13270.

Friggens, N.C., and Y. De Haas. 2019. Resilience: reference measures based on longer-term

consequences are needed to unlock the potential of precision livestock farming technologies for

quantifying this trait. Available on www.gentore.eu.

Friggens, N.C., and V.M. Thorup. 2015. From Monitoring to Precision Phenotyping: Towards a

Systemic Use of Precision Livestock Measures in Dairy Herds. Proceedings of the New Zealand

Society of Animal Production. 15:146-148

Gröhn, Y.T., D.J. Wilson, R.N. González, J. a Hertl, H. Schulte, G. Bennett, and Y.H. Schukken.

2004. Effect of pathogen-specific clinical mastitis on milk yield in dairy cows. J. Dairy Sci.

87:3358–74. doi:10.3168/jds.S0022-0302(04)73472-4.

Højsgaard, S., and N.C. Friggens. 2010. Quantifying degree of mastitis from common trends in a

panel of indicators for mastitis in dairy cows. J. Dairy Sci. 93:582-592. doi:10.3168/jds.2009-

2445.

Hudson, C., J. Kaler, and P. Down. 2018. Using big data in cattle practice. In Pract. 40:396–408.

doi:10.1136/inp.k4328.

König, S., and K. May. 2019. Invited review: Phenotyping strategies and quantitative-genetic

background of resistance, tolerance and resilience associated traits in dairy

cattle. Animal, 13(5):897-908. doi:10.1017/S1751731118003208

Lavon, Y., G. Leitner, H. Voet, and D. Wolfenson. 2010. Naturally occurring mastitis effects on

timing of ovulation, steroid and gonadotrophic hormone concentrations, and follicular and luteal

growth in cows. J. Dairy Sci. 93:911–921. doi:10.3168/jds.2009-2112.

Mapletoft, R., and J. Hasler. 2002. Assisted reproductive technologies in cattle: a review. Sci. Tech.

Rev. Off. Int. des Epizoot. 24:393–403. doi:10.1057/palgrave.development.1100303.

Mulder, H.A., H. Rashidi. 2017. Selection on resilience improves disease resistance and tolerance to

infections. J. Anim. Sci. 95(8):3346-3358. doi:10.2527/jas.2017.1479.

Pinedo, P.J., A. De Vries, and D.W. Webb. 2010. Dynamics of culling risk with disposal codes

reported by Dairy Herd Improvement dairy herds. J. Dairy Sci. 93:2250–2261.

doi:10.3168/jds.2009-2572.

Rajala-Schultz, P.J., Y.T. Gröhn, C.E. McCulloch, and C.L. Guard. 1999. Effects of clinical mastitis

on milk yield in dairy cows.. J. Dairy Sci. 82:1213–20. doi:10.3168/jds.S0022-0302(99)75344-

0.

Royal, M.D., A.O.R. Darwash, A.P.F.F. Flint, R. Webb, J.A. Woolliams, and G.E. Lamming. 2000.

Declining fertility in dairy cattle: Changes in traditional and endocrine parameters of fertility.

Anim. Sci. 70:487–501. doi:10.1017/S1357729800051845.

Rutten, C.J., a G.J. Velthuis, W. Steeneveld, and H. Hogeveen. 2013. Invited review: sensors to

support health management on dairy farms. J. Dairy Sci. 96:1928–1952. doi:http://dx.doi.org/

10.3168/jds.2012-6107.

Santos, J.E.P., R.S. Bisinotto, and E.S. Ribeiro. 2016. Mechanisms underlying reduced fertility in

anovular dairy cows. Theriogenology 86:254–262. doi:10.1016/j.theriogenology.2016.04.038.

Sorg, D., M. Wensch-Dorendorf, K. Schöpke, G. Martin, R. Schafberg, N. Reinhold, S. Pache, and

H. Swalve. 2017. Genetic analysis of new progesterone-based fertility traits in dairy cows

measured on-farm. J. Dairy Sci. 100:8205–8219. doi:10.3168/jds.2016-11865.

Steeneveld, W., and H. Hogeveen. 2015. Characterization of Dutch dairy farms using sensor systems

for cow management. J. Dairy Sci. 98:709–717. doi:10.3168/jds.2014-8595.

Tenghe, A.M.M., A.C. Bouwman, B. Berglund, E. Strandberg, J.Y. Blom, and R.F. Veerkamp. 2015.

Estimating genetic parameters for fertility in dairy cows from in-line milk progesterone profiles..

J. Dairy Sci. 98:5763–5773. doi:10.3168/jds.2014-8732.

Vandeweerd, J.M., N. Kirschvink, P. Clegg, S. Vandenput, P. Gustin, and C. Saegerman. 2012. Is

evidence-based medicine so evident in veterinary research and practice? History, obstacles and

perspectives. Vet. J. 191:28–34. doi:10.1016/j.tvjl.2011.04.013.

van Knegsel, A.T.M., H.M. Hammon, U. Bernabucci, G. Bertoni, R.M. Bruckmaier, R.M.A.

Goselink, J.J.Gross, B. Kuhla, C.C. Metges, H.K. Parmentier, E. Trevisi, A. Tröscher and A.M.

van Vuuren. 2014. Metabolic adaptation during early lactation: key to cow health, longevity and

a sustainable dairy production chain. CAB reviews 9:1-15. doi:10.1079/PAVSNNR20149002.

Wathes, D.C. 2012. Mechanisms linking metabolic status and disease with reproductive outcome in

the dairy cow. Reprod. Domest. Anim. 47:304–312. doi:10.1111/j.1439-0531.2012.02090.x.

Wolfenson, D., G. Leitner, and Y. Lavon. 2015. The disruptive effects of mastitis on reproduction

and fertility in dairy cows. Ital. J. Anim. Sci. 14:650–654. doi:10.4081/ijas.2015.4125.

Wood, P. 1967. Algebraic model of the lactation curve in cattle. Nature 216:164–165.

 APPENDICES

Appendix A – Definition and Calculation of Milk Yield Sensor Features

In Table A1, the details of the different sensor features (**SF**) calculated from daily milk yield data

included in the prediction models are given. The Iterated Theoretical Wood model (**ITW**) is the result

of the iterative fitting and refitting procedure excluding perturbations to estimate the shape of the

theoretical lactation curve. All SF are standardized at herd level before entering them in the models,

by $= \text{SF}_{\text{standardized}} = \frac{SF - \overline{SF}}{SD(SF)}$ with SF each sensor feature, $\overline{SF}$ the average of each SF for a herd, and

SD(SF) the standard deviation for that SF for a herd.

Table A1. Lactation sensor features (**SF**) and their calculation included in the prediction models for

lifetime resilience ranking

SF No.	Name	No. of times (%) retained[5] Lactation SF / activity & lactation SF	How calculated?
SF1	305-day yield	**6 (22.2%)** **4 (30.8%)**	Total milk yield in the first 305 days of lactation
SF2	RMSE ITW[1]	**5 (18.5%)** **2 (15.4%)**	Root mean squared error of the iterated theoretical Wood model
SF3	Peak yield ITW[1]	**5 (18.5%)** **3 (23.1%)**	Maximal value of the iterated theoretical Wood model
SF4	DIM[4] peak yield ITW[1]	**4 (14.8%)** **0 (0.0%)**	Days in milk of the peak milk yield (SF3)
SF5	Rate increase ITW[1]	**5 (18.5%)** **4 (30.8%)**	Peak yield (SF3) divided by days in milk of peak milk yield (SF4)
SF6	Persistency ITW[1]	**10 (37.0%)** **5 (38.5%)**	Persistency, slope of the linear decreasing phase of the iterated theoretical Wood model
SF7	Sum of strong negative residuals of ITW[1]	**2 (7.4%)** **4 (30.8%)**	Sum of the residuals smaller than 85% of the iterated theoretical Wood model
SF8	Ratio of positive vs. negative residuals of ITW[1]	**5 (18.5%)** **4 (30.8%)**	Ratio of the number of positive vs. negative residuals calculated from the iterated theoretical Wood model
SF9	Number of major[2] perturbations	**5 (18.5%)** **2 (15.4%)**	Number of perturbations with milk production at least one day below 80% of the iterated theoretical Wood model and lasting at least 10 days
SF10	Average number of days needed for recovery from major[2] perturbations	**5 (18.5%)** **3 (23.1%)**	Number of days needed for recovery from a major perturbation (milk production at least one day below 80% of the iterated theoretical Wood model and lasting at least 10 days)
SF11	Average milk loss during recovery phase of major[2] perturbations	**7 (25.9%)** **2 (15.4%)**	Average milk losses during recovery phase of a major perturbation (milk production at least one day below 80% of the iterated theoretical Wood model and lasting at least 10 days)
SF12	Average milk loss during development phase of major[2] perturbations	**6 (22.2%)** **3 (23.1%)**	Average milk losses during development phase of a major perturbation (milk production at least one day below 80% of the iterated theoretical Wood model and lasting at least 10 days)

SF13	Average number of days needed for development of major[3] perturbations	**6 (22.2%)** **5 (38.5%)**	Number of days needed to reach the minimum of a major perturbation (milk production at least one day below 80% of the iterated theoretical Wood model and lasting at least 10 days)
SF14	Average minimum milk yield of the major[2] perturbations	**7 (25.9%)** **3 (23.1%)**	Average minimal milk yield of the major perturbations (milk production at least one day below 80% of the iterated theoretical Wood model and lasting at least 10 days)
SF15	Number of minor[3] perturbations	**6 (22.2%)** **1 (7.7%)**	Number of perturbations with milk production at least one day between 90 and 80% of the iterated theoretical Wood model and lasting at least 5 days
SF16	Average number of days needed for recovery from minor[3] perturbations	**5 (18.5%)** **1 (7.7%)**	Number of days needed for recovery from a minor perturbation (milk production at least one day between 90 and 80% of the iterated theoretical Wood model and lasting at least 5 days)
SF17	Average milk loss during recovery phase of minor[3] perturbations	**4 (14.8%)** **0 (0.0%)**	Average milk losses during recovery phase of a minor perturbation (milk production at least one day between 90 and 80% of the iterated theoretical Wood model and lasting at least 5 days)
SF18	Average milk loss during development phase of minor[3] perturbations	**6 (22.2%)** **2 (15.4%)**	Average milk losses during development phase of a minor perturbation (milk production at least one day between 90 and 80% of the iterated theoretical Wood model and lasting at least 5 days)
SF19	Average number of days needed for development of minor[3] perturbations	**3 (11.1%)** **4 (30.8%)**	Number of days needed to reach the minimum of a minor perturbation (milk production at least one day between 90 and 80% of the iterated theoretical Wood model and lasting at least 5 days)
SF20	Average minimum milk yield of the minor[3] perturbations	**5 (18.5%)** **5 (38.5%)**	Average minimal milk yield of the minor perturbations (milk production at least one day between 90 and 80% of the iterated theoretical Wood model and lasting at least 5 days)
SF21	Number of periods in which milk yield drops below 85% of the expected yield (ITW) and that last > 5 days	**6 (22.2%)** **5 (38.5%)**	Number of periods in which milk yield drops below 85% of the expected yield (iterated theoretical Wood model) and that last > 5 days
SF22	Number of periods in which milk yield drops below 85% of the expected yield and that last > 10 days	**6 (22.2%)** **3 (23.1%)**	Number of periods in which milk yield drops below 85% of the expected yield (iterated theoretical Wood model) and that last > 5 days
SF23	Percentage of days in which milk yield drops below 85% of the expected yield (ITW)	**5 (18.5%)** **2 (15.4%)**	Number of periods in which milk yield drops below 85% of the expected yield (iterated theoretical Wood model)
SF24	Average milk loss of the 3 largest perturbations	**4 (14.8%)** **1 (7.7%)**	Average milk loss of the 3 largest perturbations
SF25	Largest negative residual	**6 (22.2%)** **1 (7.7%)**	Largest negative residual
SF26	Number of sign changes of the residuals throughout the lactation	**6 (22.2%)** **2 (15.4%)**	Number of sign changes of the residuals of the iterated theoretical Wood model throughout the lactation
SF27	Average residual	**9 (33.3%)** **7 (53.8%)**	Average of the residuals
SF28	Average absolute value of the residuals	**7 (25.9%)** **5 (38.5%)**	Average of the absolute values of the residuals
SF29	Variance of the residuals of the initial Wood model	**6 (22.2%)** **5 (38.5%)**	Variance of the residuals
SF30	Ratio of the initial Wood curve fitted on all data and the ITW[1]	**3 (11.1%)** **4 (30.8%)**	Ratio of the initial Wood curve fitted on all the data and the iterated theoretical Wood curve

[1] ITW: Iterated Wood Model

[2] Major perturbations: periods of at least 10 days of successively negative residuals with at least one day of milk production lower

than 80% of the theoretical production

[3] Minor perturbations: periods of at least 5 days of successively negative residuals with at least one day of milk production between

90% and 80% of the expected production

[4] DIM: days in milk

[5] For each SF, it was evaluated how often it was retained in the individual farm models of data set 1 using the stepwise selection

procedure. The more often a certain SF was retained, the more informative it was for predicting the lifetime resilience ranking of the

cows. Data set 1 included 27 farms in total.

Appendix B. Definition and calculation of activity sensor features (SF)

First, the two-hourly activity data were aggregated in daily sums. Next, a moving median using a

window of 4 days was calculated on these daily data time series to identify short spikes associated

with estrous behavior (level 1; all spikes > 0.4* the maximal residual of the time series minus the

moving median). A moving median of 20 days was calculated to identify periods with generally lower

or higher activity (possibly associated with health events). A threshold of 20% of the minimal or

maximal activity residuals was set to identify these altered activity period. The below explained

calculated SF are based on the deviations from these median windows.

Table B1. Activity sensor features (**SF**) and their calculation included in the prediction models for

lifetime resilience ranking

SF No.	Activity feature	No. of times (%) retained[1]	How calculated?
SF31	Daily mean	**6 (46.2%)**	Daily mean activity value
SF32	Daily overall skewness	**1 (7.7%)**	Skewness of the daily activity time series
SF33	Daily overall variance	**9 (69.2%)**	Overall variance of the daily activity time series
SF34	Number of first level peaks	**11 (84.6%)**	Number of sharp peaks, larger than 0.4 times the maximal peak height after correction for 4-day median activity level
SF35	Average peak height of first level peaks	**5 (38.5%)**	Average peak height of peaks larger than 0.4 times the maximal peak height after correction for 4-day median activity level
SF36	Maximal peak height of first level peaks	**5 (38.5%)**	Maximal peak height of peaks larger than 0.4 times the maximal peak height after correction for 4-day median activity level
SF37	Days in milk of first peak	**8 (61.5%)**	Days in milk of first peak larger than 0.4 times the maximal peak height after correction for 4-day median activity level
SF38	Number of periods with higher activity compared to long-term median activity	**3 (23.1%)**	Number of periods with activity level higher than 0.2 times the maximal level compared to long-term 20-day median activity level
SF39	Average activity level of periods with higher activity compared to long-term median activity	**3 (23.1%)**	Average activity level of periods with activity level higher than 0.2 times the maximal level compared to long-term 20-day median activity level
SF40	Maximal activity level in periods with higher activity compared to long-term median activity	**2 (15.4%)**	Maximal activity level of periods with activity level higher than 0.2 times the maximal level compared to long-term 20-day median activity level
SF41	Duration of periods with higher activity compared to long-term median activity	**4 (30.8%)**	Duration of periods with activity level higher than 0.2 times the maximal level compared to long-term 20-day median activity level
SF42	Number of periods with lower activity compared to long-term median activity	**2 (15.4%)**	Number of periods with activity level lower than 1.2 times the minimal level compared to long-term 20-day median activity level
SF43	Average activity level of periods with lower activity compared to long-term median activity	**3 (23.1%)**	Average activity level of periods with activity level lower than 1.2 times the minimal level compared to long-term 20-day median activity level

SF44	Minimal activity level in periods with lower activity compared to long-term median activity	**0 (0.0%)**	Minimal activity level of periods with activity level lower than 1.2 times the minimal level compared to long-term 20-day median activity level
SF45	Duration of periods with lower activity compared to long-term median activity	**2 (15.4%)**	Duration of periods with activity level lower than 1.2 times minimal level compared to long-term 20-day median activity level

[1] For each SF, it was evaluated how often it was retained in the individual farm models of data set 2 using the stepwise selection

procedure. The more often a certain SF was retained, the more informative it was for predicting the lifetime resilience ranking of the

cows. Data set 2 included 13 farms in total.

Figure 1. Example of a lactation curve with daily milk yields, the corresponding initial Wood model

fitted on all daily milk yield data and the final Wood model fitted iteratively by excluding daily milk

yields lower than 85% of the estimated curve. The residuals of the latter curve are used to characterize

perturbations (e.g., representing health events).

Figure 2. Example of the relation between the lifetime resilience scores of all the animals on an

example farm (No. of cows = 110) against the parity number in which each animal exits the herd.

Higher resilience scores generally correspond to higher last lactation numbers.

Figure 3. Pearson linear correlation coefficients between the lifetime resilience rank at farm level

and the 30 lactation sensor features (**SF**) calculated on first parity data. Each individual thin line

represents the correlations for a particular farm (total No. = 27). The shading represents the 95%

confidence interval on the correlation coefficients over all farms. The details of the SF are described

in Appendix A. The lack of consistency in the correlations' signs and the magnitude demonstrate the

large variability in the relation between the lifetime resilience and the lactation SF.

Figure 4. Pearson linear correlation coefficients between the lifetime resilience rank at farm level

and the 15 activity sensor features (**SF**) calculated on first parity data. Each individual thin line

represents the correlations of a particular farm (total No. = 13). The shading represents the 95%

confidence interval on the correlation coefficients over all farms. The details of the activity SF are

described in Appendix B. The lack of consistency in the correlations' signs and the magnitude

demonstrate the large variability in the relation between the lifetime resilience and the activity SF.

Only SF34 (i.e., number of sharp activity peaks corresponding to estrus) have a consistently positive

correlation with the lifetime resilience rank.

Adriaens, Figure 1
747

749   **Adriaens,** Figure 2
750

751
752

753   **Adriaens,** Figure 3

754
755

756   **Adriaens,** Figure 4

757
758

759 **Adriaens,** Table 1**.** Overview of the available data sets. Data set 2 (**DS2**) is a subset of data set 1

760 (**DS1**) for which also daily activity data were available besides milk meter data.

Descriptor	Data set 1 (DS1) Average ± SD [range]	Data set 2 (DS2) Average ± SD [range]
No. of days data per farm	3001 ± 828 [1830 ; 4977]	3020 ± 815 [1837 ; 4101]
No. of cows in total	3754	2075
No. of cows per farm	139 ± 82 [24 ; 308]	160 ± 94 [57 ; 308]
No. of lactations in total	9395	5286
No. of lactations per farm	348 ± 229 [44 ; 799]	407 ± 264 [113 ; 799]
Age at first calving per farm (years)	2.2 ± 0.1 [2.0 ; 2.4]	2.2 ± 0.1 [2.0 ; 2.3]
Average 305-day milk yield per farm (kg)	9557 ± 1007 [7931 ; 11739]	9691 ± 1061 [8244 ; 11739]
Average calving interval per farm (days)	407 ± 16 [377 ; 433]	403 ± 15 [377 ; 432]
Average dry period length per farm (days)	56 ± 13 [35 ; 91]	59 ± 16 [35 ; 91]

761
762

763   **Adriaens,** Table 2**.** Overview of a selection of lactation and activity sensor features calculated on first

764   parity data for data set 1 (**DS1**) and data set 2 (**DS2**). A detailed overview for all the included features

765   is given in Appendices A and B.

	Data set 1 (DS1) Average ± SD [range]	Data set 2 (DS2) Average ± SD [range]
No. of cows included	**3754**	**2075**
Theoretical 305-day yield	8261 ± 1772 [2688 ; 13755]	8317 ± 1874 [2689 ; 13755]
RMSE[1] [kg.day^{-1}]	3.4 ± 1.4 [1 ; 28]	3.8 ± 1.5 [1.2 ; 28]
Peak yield [kg]	34 ± 6 [15 ; 61]	34 ± 6 [14 ; 61]
Days in milk of peak lactation [day]	59 ± 25 [1 ; 248]	63 ± 27 [1 ; 207]
Rate increasing part of the lactation curve [kg.day^{-1}]	0.64 ± 0.28 [0.08 ; 6.7]	0.62 ± 0.31 [0.08 ; 6.7]
Persistency of the lactation curve [kg.day^{-1}]	-0.05 ± 0.03 [-0.15 ; 0.12]	-0.04 ± 0.03 [-0.15 ; 0.12]
No. of perturbations per lactation	6.0 ± 3.1 [0 ; 17]	6.3 ± 3.2 [0 ; 17]
Losses in perturbations per lactation [kg]	148 ± 124 [0 ; 1137]	154 ± 132 [0 ; 1129]
Activity skewness		-0.05 ± 1.64 [-5.19 ; 8.77]
No. of sharp and short activity peaks		3.9 ± 2.7 [0 ; 18]
Days in milk of first activity peak [days]		56 ± 64 [1 ; 305]
No. of general activity changes		24 ± 16 [2 ; 94]

766   1 RMSE: Root mean squared error of the Wood model fitted on each lactation curve

Adriaens, Table 3. Prediction results of the individual stepwise regression models in calibration and 10-times cross-validation (**CV**)

	Calibration, milk yield features only					Calibration, milk yield & activity features								CV[8], milk yield features only				CV, milk yield & activity features			
FARM ID	R^2_{adj}	N^1	$N_{LAC}{}^2$	$N_{GOF}{}^3$	$N_{PERT}{}^4$	R^2_{adj}	N	N_{LAC}	N_{GOF}	N_{PERT}	$N_{GEN}{}^5$	$N_{L1}{}^6$	$N_{L2}{}^7$	$RMSE_{TR}{}^9$	$RMSE_{CV}{}^{10}$	$P_C{}^{11}$	$P_O{}^{12}$	$RMSE_{TR}$	$RMSE_{CV}$	P_C	P_O
FARM1	0.10	8	2	4	2									0.26	0.27	45.9	3.7				
FARM2	0.16	5	0	2	3									0.24	0.27	44.3	1.1				
FARM3	0.05	4	1	1	2									0.25	0.27	41.5	3.6				
FARM4	0.27	3	0	1	2									0.21	0.21	50.0	0.0				
FARM5	0.29	8	0	2	6									0.22	0.25	45.8	6.8				
FARM6	0.53	5	1	1	3									0.17	0.18	70.0	0.0				
FARM7	0.31	4	0	2	2									0.23	0.24	59.0	2.6				
FARM8	0.42	11	2	5	4									0.19	0.23	45.6	6.3				
FARM9	0.08	2	0	1	1									0.25	0.25	44.6	2.2				
FARM10	0.16	8	2	2	4									0.24	0.27	48.9	5.6				
FARM11	0.14	7	2	0	5									0.24	0.26	37.1	3.4				
FARM12	0.40	9	1	3	5	0.51	8	1	2	2	1	2	0	0.20	0.24	55.9	1.7	0.18	0.20	52.5	1.7
FARM13	0.21	8	2	3	3	0.32	7	1	3	0	1	2	0	0.23	0.25	45.0	11.2	0.21	0.22	52.6	1.0
FARM14	0.28	6	1	3	2									0.21	0.23	45.6	1.3				
FARM15	0.06	3	2	0	1									0.19	0.19	41.8	1.8				
FARM16	0.10	9	3	2	4	0.30	15	3	2	2	1	4	3	0.23	0.24	44.8	7.4	0.20	0.21	51.5	1.8
FARM17	0.08	6	1	2	3	0.27	14	1	3	5	2	2	1	0.26	0.27	35.8	6.1	0.23	0.26	45.3	4.0
FARM18	0.53	7	2	2	3	0.76	13	0	3	6	2	1	1	0.16	0.19	55.0	5.0	0.10	0.15	77.5	0.0
FARM19	0.18	6	3	1	2	0.34	11	2	4	2	0	2	1	0.23	0.25	55.0	8.3	0.19	0.24	52.3	1.5
FARM20	0.14	5	0	2	3									0.24	0.25	40.7	5.0				
FARM21	0.21	4	1	1	2	0.33	5	0	1	1	1	0	2	0.22	0.22	35.9	5.3	0.20	0.22	43.5	0.0
FARM22	0.30	6	1	0	5	0.67	23	3	5	9	2	2	2	0.18	0.20	45.6	1.4	0.11	0.17	66.2	0.0
FARM23	0.11	5	0	1	4	0.24	14	0	1	4	1	4	4	0.26	0.26	39.3	2.8	0.23	0.25	44.4	6.7
FARM24	0.08	5	1	1	3	0.36	13	1	3	1	2	4	2	0.26	0.26	43.4	5.9	0.21	0.22	53.2	0.5
FARM25	0.03	3	0	2	1	0.20	12	1	4	1	2	2	2	0.26	0.26	41.9	16.5	0.23	0.24	49.4	5.2
FARM26	0.61	12	1	2	9	0.62	8	1	1	4	1	1	0	0.15	0.21	62.0	2.0	0.16	0.18	84.0	4.2
FARM27	0.15	6	1	3	2	0.32	13	2	2	5	0	3	1	0.26	0.26	40.4	2.5	0.22	0.24	48.5	3.7

[1] **N**: Total number of sensor features retained in the stepwise linear regression;

[2] N_{LAC}: Number of features linked with the milk yield shape characteristics retained in the final regression models;

[3] N_{GOF}: Number of features linked with the goodness-of-fit of the iterated Wood curve fitted on the lactation data retained in the final models;

[4] N_{PERT}: Number of features linked with the identified perturbations retained in the final models;

[5] N_{GEN}: Number of features linked with the general activity time series retained in the final models;

[6] N_{L1}: Number of features linked with the activity peaks assumed to relate to estrus behavior retained in the final models;

[7] N_{L2}: Number of features linked with periods of high or low average activity assumed to relate to general deviation in behavior or activity level in the final models;

[8] **CV**: Results of the 10-fold cross-validation

[9] $RMSE_{TR}$: Average root mean squared error of the fitted scaled lifetime resilience ranking of the cows in the training set;

[10] $RMSE_{CV}$: Average root mean squared error of cross-validation for the predicted scaled lifetime resilience ranking of the cows in the validation set;

[11] P_C: Proportion of the animals on each farm predicted in the correct category (low, **L**, moderate, **M** or high, **H**);

[12] P_O: Proportion of the animals on each farm predicted in the opposite category: cows predicted as being in the lowest third while being in the highest third and vice versa