Title:

Inhibiting IRE1α-endonuclease activity in hepatic stellate cells decreases tumor cell

proliferation and metastasis in hepatocellular carcinoma

Authors: Natasa Pavlovic[1], Carlemi Calitz[1], Kessarin Thanapirom[3], Giuseppe

Mazza[3], Krista Rombouts[3], Pär Gerwins[1,2] and Femke Heindryckx[1]

Affiliations:

(1) Department of Medical Cell Biology, Uppsala University, Uppsala, Sweden

(2) Department of Radiology, Uppsala University Hospital, Uppsala, Sweden

(3) Regenerative Medicine & Fibrosis Group, Institute for Liver and Digestive Health,

University College London (UCL), London, UK

Running title: Inhibiting ER-stress decreases hepatocellular carcinoma

Keywords: Endoplasmic reticulum stress; fibrosis; hepatic stellate cells; IRE1α;

tumor-stroma interactions;

Abbreviations: αSMA - α-smooth muscle actin; DEN – diethylnitrosamine; DMEM -

Dulbecco modified eagle medium; ELISA - Enzyme-Linked immune Sorbent Assay*,*

ER - Endoplasmic reticulum; FBS - fetal bovine serum; HCC - Hepatocellular

carcinoma, H&E - Haematoxilin-eosin; TBS - tris-buffer saline; TGFβ - tumor growth

factor β; UPR - unfolded protein response;

Corresponding author: Femke Heindryckx, Department of Medical Cell Biology,

BOX 571, Husargatan 3, 75123 Uppsala, Sweden. Phone: +46768542025 Email:

femke.heindryckx@mcb.uu.se

ABSTRACT

Hepatocellular carcinoma (HCC) is a liver tumor that arises in patients with cirrhosis.

One of the key players in the progression of cirrhosis to HCC is the hepatic stellate

cell, which is activated during liver damage. Activated stellate cells play an essential

role in the pathogenesis of HCC by creating a fibrotic micro-environment that sustains

tumor growth and by producing growth factors and cytokines that enhance tumor cell

proliferation and migration. We assessed the role of endoplasmic reticulum (ER) stress

in the cross-talk between hepatic stellate cells and HCC-cells. Mice with a fibrotic HCC

were treated with the IRE1α-inhibitor 4μ8C, which led to a significant reduction in tumor

burden and collagen deposition. By co-culturing HCC-cells with hepatic stellate cells,

we found that HCC-cells induce ER-stress in hepatic stellate cells, thereby contributing

to their activation. Inhibiting IRE1α blocked stellate cell activation, which inhibited

tumor cell proliferation and migration in different *in vitro* 2D and 3D co-cultures. Our

results suggest that IRE1α is an important mediator in the communication between

stellate cells and cancer cells and components of the ER-stress pathway may be

therapeutically relevant for slowing down the progression of HCC.

INTRODUCTION

Hepatocellular carcinoma (HCC) is a primary liver tumor that typically arises in a background of chronic liver disease and cirrhosis (1). One of the key players in the progression of cirrhosis to HCC is the hepatic stellate cell, which activates during liver damage and differentiates towards a contractile myofibroblast-like cell responsible for the deposition of extracellular matrix proteins (ECM) such as collagen (2). Activated stellate cells can induce phenotypic changes in cancer cells through the production of growth factors and cytokines that stimulate tumor cell proliferation and induce a pro-metastatic phenotype (4). One of the key factors in the cross talk between tumor cells and stellate cells is tumor growth factor beta (TGFβ) (5-7). Malignant hepatocytes secrete high levels of TGFβ, which can contribute to the activation of stellate cells in the nearby stroma. These activated stellate cells are then responsible for the deposition of ECM. Several of the ECM components such as proteoglycans, collagens, laminin, and fibronectin interact with tumor cells and cells in the stroma, which can directly promote cellular transformation and metastasis (10, 11). The ECM can also act as a reservoir for growth factors and cytokines, which can be rapidly released to support the tumor´s needs. In addition, activated stellate cells contribute to a highly vascularized tumor micro-environment, by secreting pro-angiogenic molecules and by recruiting pro-angiogenic (and pro-tumoral) myeloid and lymphoid derived cell types (12). By constricting the hepatic microvasculature, they also cause hypoxia, which contributes to the angiogenic switch and can induce a more aggressive tumor phenotype (13). It is therefore not surprising that tumor cells actively secrete growth factors (such as TGFβ) to induce activation and migration of stellate cells, which creates a fibrotic environment that further supports and enhances tumor progression (2, 16, 17). Since activated stellate cells play an essential role in the onset and

progression of HCC, blocking their activation has been proposed as a potential therapy for patients with HCC (3). One strategy to block stellate cell activation, is by targeting the unfolded protein response (UPR).

The unfolded protein response serves to cope with misfolded or unfolded proteins in the ER in an attempt to restore protein folding, increase ER-biosynthetic machinery and maintain cellular homeostasis (18). It can exert a cytoprotective effect by re-establishing cellular homeostasis, while apoptotic signaling pathways will be activated in case of severe and/or prolonged ER-stress (19). The presence of misfolded proteins is sensed via 3 transmembrane proteins in the ER: IRE1α, PERK and ATF6a. Actors of the ER-stress pathways have been described to play a role in the progression of solid tumors, such as breast cancer (20), colon cancer (21) and HCC (22). Activation of the UPR has also been shown to affect different fibrotic diseases (23), including non-alcoholic fatty liver disease (24-26), hepatitis B-induced carcinogenesis (27) and biliary cirrhosis (28). We have previously shown that inhibiting the IRE1α-branch of the UPR-pathway using 4µ8C, blocks TGFβ-induced activation of fibroblasts and stellate cells *in vitro* and reduces liver fibrosis *in vivo* (29)*.* In the current study, our aim was to define the role of ER-stress in the cross-talk between hepatic stellate cells and tumor cells in liver cancer. We show that pharmacologic inhibition of the IRE1α signaling pathway decreases tumor burden in a chemically induced mouse model for HCC. Using several *in vitro* co-culturing methods, we identified that tumor cells induce ER-stress in hepatic stellate cells. Blocking ER-stress in these hepatic stellate cells prevents their activation and decreases proliferation and migration of tumor cells co-cultured with hepatic stellate cells.

MATERIAL AND METHODS

Mouse model

A chemically induced mouse model for HCC was used, as previously described (30, 31). Briefly, 5-week-old male sv129 mice received intraperitoneal injections once per week with 35mg/kg bodyweight DEN diluted in saline. From week 10, mice were injected twice per week with 10μg/g bodyweight 4μ8C (Sigma) in saline. After 25 weeks mice were euthanized and samples were taken for analysis. This method was approved by the Uppsala ethical committee for animal experimentation (C95/14). Each group contained 8 mice.

Olink multiplex proximity extension assay

Cells were detached with 100 μL ice-cold RIPA containing protease inhibitors (Sigma Aldrich), and cells were scraped off and aspired. Cell suspensions were on ice for 20–30 min, whilst mixing vigorously to enhance disruption of the cell membranes. The cell suspensions were centrifuged (20 min, 13 000 rpm, 4°C) and supernatant containing protein was collected. Supernatant was stored at -20°C until protein measurement. Protein concentration was measured using the BCA kit (ThermoFisher) and all samples were diluted to 1 mg/mL protein in RIPA. All samples were analyzed with a multiplex proximity extension assay for ninety-two biomarkers in the murine exploratory panel (Olink Bioscience, Uppsala, Sweden) [30].

Cell culture and reagents

The HCC-cell lines (HepG2, ATCC and Huh7, kind gift from Dilruba Ahmed, Karolinska Institute) and hepatic stellate cell-line LX2 (Sigma-Aldrich, Darmstadt, Germany) were cultured at 37°C with 5% CO_2 in Dulbecco modified eagle medium (DMEM)

supplemented with 10% fetal bovine serum (FBS) (ThermoFisher, Stockholm, Sweden). No FBS was used during starvation and stimulation with growth factors. Cells were detached using trypsin-EDTA (ThermoFisher), re-suspended in growth medium and plated at a density of $5x10^3$ cells/cm^2. Cells were allowed to attach and left undisturbed for 8h before being starved for 16h. Afterwards, fresh starvation medium containing indicated growth factors or substances were added. Cells were exposed for 48h to 100µM 4µ8C (Sigma-Aldrich) or 10µM SB-431541 (Tocris, Abingdon, UK) as previously described (29). For transwell co-culturing experiments, cells were grown on 12-well Corning Tissue Culture-plates with transwell inserts (Sigma-Aldrich) with 0,4µm-pore size, allowing the exchange of soluble factors, but preventing direct cell contact.

3D-tumor spheroids were generated on 96-well ultra-low attachment plates (Sigma-Aldrich) (32). After 6 days, spheroids had reached approximately 1mm^2 and 4µ8C was added. Proliferation was monitored during the subsequent 4 days. Tumor spheroids were retrieved from the plates after 10 days.

Fluorescent labeling of cells was done in imaging experiments by using CellTracker (ThermoFisher), according to manufacturer´s instructions. Cell pellets were incubated 30 minutes with 1µM of CellTracker™ Red CMTPX or 1µM of CellTracker™ Green CMFDA. Cells were washed twice in PBS prior to co-culturing.

Human liver scaffold decellularization and cell culture usage

Human healthy livers were obtained under the UCL Royal Free BioBank Ethical Review Committee (NRES Rec Reference: 11/WA/0077) approval. Informed consent

was obtained for each donor and confirmed via the NHSBT ODT organ retrieval

pathway (33). Liver 3D-scaffolds, were decellularized, sterilized and prepared for cell

culture use as preciously described (33). LX2 and HepG2-cells, as either mono-

cultures or mixed co-culture, were released on top of each scaffold as $2.5*10^5$ cells in

20μL (34).

Proliferation

Cell proliferation was monitored via a resazurin reduction assay. A 1%-resazurin

solution was added in 1/80 dilution to the cells and incubated for 24h, after which

fluorescent signal was measured with a 540/35 excitation filter and a 590/20 emission

filter on a Fluostar Omega plate reader.

Transfections

Nucleofection with 0,1μM siIRE1α (s200432, ThermoFisher), or 0,1 μM siCtrl

(4390843, ThermoFisher) was done using Amaxa Nucleofector program S-005 in

Ingenio electroporation solution (Mirus Bio LLC, Taastrup, Denmark).

Migration and chemotaxis

Non-directional migration was assessed using a scratch wound assay on fluorescently

labelled LX2-cells and HepG2-cells. Scratch size was measured by analyzing light

microscopy images in ImageJ, using the MRI Wound Healing Tool plug-in

(http://dev.mri.cnrs.fr/projects/imagej-macros/wiki/Wound_Healing_Tool). Image

analysis was done in ImageJ.

Directional migration was assessed using CellDirector-devices (GradienTech,

Uppsala, Sweden). HepG2 and LX2-cells were labelled with CellTracker-dye and left

to adhere overnight in the CellDirector-devices. Non-adherent cells were washed away with DMEM and cells were starved for 1h prior to commencing experiments. A gradient of 0 to 10% FBS was created with a flow rate of 1.5 µl/minute. Cell movement was recorded using an Axiovision 200M microscope (Zeiss, Stockholm, Sweden) for 4h and tracked using Axiovision software (Zeiss). During the assay cells were kept at 37˚C with 5% CO_2.

Quantitative RT-PCR of mRNA

RNA was isolated from tissue or cell culture using the EZNA RNA isolation Kit (VWR, Spånga, Sweden) or using TRIzol reagent and RNeasy Universal Mini Kit (Qiagen, Sollentuna, Sweden) for human liver scaffolds (33). RNA-concentration and purity were evaluated using Nanodrop. Afterwards, 500ng of mRNA was reverse transcribed using iScript cDNA synthesis kit (Bio-rad, Solna, Sweden). Amplifications were done using primers summarized in supplementary table 1. mRNA-expression was normalized to *18S, GAPDH* and/or *TBP1.* Fold change was calculated via the delta-delta-CT method*.*

The procedure to detect the spliced and unspliced isoforms of XBP1 was done by digesting RT-PCR product with the restriction enzyme *Pst*-I (ThermoFisher). This cleaves unspliced-XBP1 containing the Pst-I-cleavage site (CTGCA^G), but leaves the spliced isoform intact. The digestion reaction was stopped after 18h by 0,5M EDTA (pH 8.0) and run on a 1,5% agarose gel for 1h at 180V.

Stainings and immunocytochemistry

Tissue samples were fixed in 4% paraformaldehyde for 24h and subsequently embedded in paraffin. Cells and tumor spheroids were fixed for 10 minutes in 4%

paraformaldehyde and stored at 4°C. Detailed staining protocols are provided as supplementary methods.

Enzyme-Linked immune Sorbent Assay (ELISA)

Medium samples from cells and from the engrafted scaffolds were used to measure TGFβ using ELISA (88-8350-22, ThermoFisher), following manufacturer´s guidelines.

SDS-PAGE and western blot

Protein lysates in lysis buffer were mixed with 2x laemmli buffer and heated to 95°C for 5 minutes before being loaded onto a 10% polyacrylamide gel. After separation, the proteins were transferred to an Immobilon-Fl membrane (Millipore). The membrane was blocked using the Odyssey blocking buffer (Licor) diluted 1:4 in PBS, and then incubated with primary and secondary antibodies. After primary and secondary antibody incubation the membrane was washed 3x15 minutes in PBS-T (Phosphate buffered saline (Gibco), 0.1% Tween-20). Primary antibodies used were Bip (ab21685, abcam), p-IRE1α (PAB12435, Abnova) or vinculin (14-9777-82, ThermoFisher**)** all added in blocking buffer with 0.1% Tween-20. Secondary antibodies used were: goat-anti-mouse alexa 680 (Invitrogen) and goat-anti-rabbit IRDye 800 (Rockland) 1:20 000 diluted in blocking buffer with 0.1% Tween-20 and 0.01% SDS. All incubations were carried out at room temperature for 1h or overnight at 4°C. The membranes were scanned using an Odyssey scanner (LI-COR Biotechnology) and band intensities quantified using the Odyssey 2.1 software and normalized to the vinculin signal in each sample.

Gene-set enrichment analysis

Gene expression profiles of HCC with a fibrous stroma and without fibrous stroma was

accessed through PubMed´s Gene Expression Omnibus via accession number

GSE31370(35). A gene-set containing 79 proteins involved in the unfolded protein

response was downloaded from The Harmonizome (36) and GSEA software was used

to perform a gene-set enrichment assay (37).

Human protein atlas

Images from biopsies from HCC patients stained with antibodies against WIPI1 (38),

SHC1 (39), PPP2R5B (40) and BiP (41) were obtained through the Human Protein

Atlas (42).

Statistics

Data are presented as mean ± s.e.m. Statistical significance was determined using an

unpaired, two-tailed Student's T-test or one-way analysis of variance (ANOVA)

followed by Tukey´s multiple comparison test. Survival curves were generated with the

Kaplan-Meier method and statistical comparisons were made using the log-rank

method. P-values <0.05 were considered statistically significant. *In vitro* experiments

were done in at least 3 independent replicates, *in vivo* experiments were done on at

least 5 independent animals.

RESULTS

Pharmacological inhibition of IRE1α reduces tumor burden in a chemically

induced mouse model for HCC

Hepatocellular carcinoma was induced in mice by weekly injections with diethylnitrosamine (DEN) for 25 weeks (30). From week 10, IRE1α-endonuclease activity was pharmacologically inhibited with 4µ8C. Histological analysis of liver tissue confirmed presence of liver tumors in a fibrotic background (Figure 1A). Treatment with 4µ8C significantly reduced tumor burden (Figure1B), as measured on H&E-stained liver slides (Figure1A). Stellate cell activation and liver fibrosis was quantified by Sirius Red staining (Figure 1A and 1C) and immunohistochemical staining with αSMA-antibodies (Figure 1A and 1D) on liver sections. Mice with HCC had a significant increase in the percentage of collagen (Figure 1C) and αSMA-staining (Figure 1D), compared to healthy mice. Treatment with 4µ8C restored collagen (Figure 1C) and αSMA-levels (Figure 1D and Figure 1E) to a similar level as healthy livers. mRNA expression levels of PCNA were determined on tumor nodules and surrounding non-tumor stromal tissue (Figure 1E). As expected, proliferation of cells was increased within the tumor itself, compared to the levels seen in healthy liver tissue and stromal tissue. Treatment with 4µ8C significantly decreased the levels of PCNA mRNA expression within the tumor, suggesting a decrease in tumor cell proliferation. A proteomics array using the Olink Mouse Exploratory assay revealed that DEN-induced murine tumors had a significantly increased protein expression of 20 oncogenic proteins compared to healthy controls (Figure 1F and table 1). In the 4µ8C-treated group, only 11 oncogenic proteins were increased compared to healthy controls (Figure 1F and table 1). Treatment with 4µ8C also significantly reduced the expression of two HCC promotors, Prdx5 and DDah1 (Figure 1F and table 1).

Markers of the unfolded protein response are upregulated in HCC and mainly located in the tumor stroma.

mRNA-levels of the ER-stress-chaperone BiP were measured in tumor and surrounding non-tumor tissue of mice with DEN-induced HCC (Supplementary figure 1A). BiP-mRNA-expression was increased in the surrounding non-tumor tissue of DEN-induced mice, while there was no difference within the tumor, compared to healthy controls. Western blot confirmed the increase of BIP-protein expression in DEN-induced livers, which was reduced after treatment with 4µ8C (Supplementary figure 1 B). Co-staining of liver tissue with αSMA and p-IRE1α-antibodies (Supplementary figure 1C and D) or BiP-antibodies (Supplementary figure 1E) in untreated mice with HCC, revealed that expression of ER-stress markers was mainly localized within activated stellate cells in the liver.

A gene-set enrichment assay on microarray data from HCC-patients with fibrotic septae and without fibrotic septae showed an increase of genes involved in the UPR in the fibrotic HCC samples compared to non-fibrous HCC (supplementary figure 2A). Several actors of the IRE1α-branch of the UPR are amongst the genes that contribute to the core-enrichment of this analysis (table 2). Immunohistochemical staining of liver biopsies from HCC-patients further confirmed presence of IRE1α-mediated ER-stress markers BiP, PPP2R5B, SHC1 and WIPI1 localized in the fibrotic scar tissue and near hepatic blood vessels (Supplementary figure 2B). In addition, increased expression of these markers was significantly correlated with poor survival in patients with liver cancer (Supplementary figure 2C).

Tumor cells secrete factors that induce ER-stress in hepatic stellate cells

Hepatic stellate cell-lines (LX2) and HCC-cell lines (HepG2 and Huh7) were grown in different compartments using a transwell assay. This confirmed that tumor cells secrete factors that increase mRNA-expression of CHOP (Figure 2A), spliced XBP1

(Figure 2B and D) and BiP (Figure 2C), as well as protein expression of p-IRE1α (Figure 2E) in hepatic stellate cells co-cultured with tumor cells, indicating the presence of ER-stress. This also led to their activation, as measured by mRNA-expression of αSMA (Figure 2F) and collagen (Figure 2G) in LX2-cells grown with HepG2 or Huh7 cells in a transwell assay. The mRNA-expression of αSMA and collagen was restored to baseline levels when 4μ8C was added to the transwell co-cultures.

De-cellularised human liver 3D-scaffolds were engrafted with hepatic stellate cells (LX2) and tumor cells (HepG2). Sirius red staining and H&E staining confirmed that that LX2-cells and HepG2-cells successfully engrafted the collagen-rich matrix of the decellularized human liver scaffolds (Figure 3A). Engrafting both LX2-stellate cells and HepG2-cancer cells led to a significant increase of mRNA-expression of collagen, BiP and spliced XBP1 (Figure 3B) compared to scaffolds that were only engrafted with LX2-cells. Adding 4μ8C significantly decreased mRNA expression of collagen and BiP-mRNA-expression in the LX2 and HepG2 co-cultured scaffolds (Figure 3B).

Tumor cells are important sources of TGFβ, which is a known activator of stellate cells. Surprisingly, measuring TGFβ in mono-cultures lead to undetectable levels of TGFβ in Huh7-cells and low-levels in HepG2-cells (Supplementary figure 3A). These levels increased when LX2-cells were added to the co-cultures (Supplementary figure 3A). Engrafting both LX2-stellate cells and HepG2-cancer cells in the human liver scaffolds, slightly increased TGFβ-levels in the medium compared to scaffolds engrafted by only one cell type, but overall no significant differences were seen (Supplementary figure 3B). It is important to note that the baseline TGFβ-levels were markedly higher in the mono-cultured scaffolds, compared to the levels measured in cells grown in a standard 2D *in vitro* set-up (Supplementary figure 3A). Blocking TGFβ-receptor signaling with SB-431541 significantly reduced mRNA-expression of ER-stress markers CHOP

(Supplementary figure 3C), spliced XBP1 (Supplementary figure 3D-E) and BiP

(Supplementary figure 3F) in stellate cells co-cultured with tumor cells using transwells.

Adding a TGFβ-receptor-inhibitor to stellate cell – tumor cell co-cultures also reduced

stellate cell activation, as measured by mRNA-expression of αSMA (Supplementary

figure 3G) and collagen (Supplementary figure 3H). This indicates that TGFβ-secretion

by tumor cells could be responsible for activating stellate cells and for inducing the

UPR.

Pharmacological inhibition of IRE1α decreases tumor cell proliferation in stellate cell – tumor cell co-cultures

In transwell co-culturing assays, we found that co-culturing HepG2 or Huh7-tumor cells

with LX2-stellate cells significantly increased PCNA-mRNA-expression in HepG2 and

Huh7-tumor cell lines (Figure 4A). Adding 4µ8C significantly decreased mRNA-

expression of PCNA in Huh7-cells grown in a transwell co-culture with LX2-cells, while

not affecting PCNA-expression in tumor cell mono-cultures (Figure 4A). PCNA-levels

in HepG2-LX2 transwell co-cultures were slightly decreased, but this was not

significant. Proliferation was measured 24h after exposure to 4µ8C in tumor cells

(HepG2 and Huh7) grown as mono-cultures and in co-culture with LX2-stellate cells.

While 4µ8C induced a significant increase in proliferation of HepG2-monocultures, no

difference was seen in LX2-monocultures and a significant decrease was seen in the

HepG2-LX2 co-cultures (Figure 4B). In the Huh7 tumor cell line, 4µ8C significantly

decreased cell number compared to untreated controls and a similar reduction was

seen in the Huh7-LX2 co-cultures (Figure 4C). Immunohistochemical staining with

antibodies against Epcam and Ki67 show that the effect on proliferation is mainly

localized in the tumor cell population of these co-cultures (Figure 4D).

3D-spheroids were generated using tumor cells alone (HepG2 or Huh7) or in combination with LX2-cells. While the HepG2-spheroids experienced a lower proliferation rate when co-cultured with LX2 stellate cells (Figure 4E), there was no difference in proliferation between spheroid-monocultures and spheroid-co-cultures in the Huh7-cells (Figure 4F). Treatment with 4μ8C significantly decreased proliferation of the tumor spheroids consisting of tumor cells (Huh7 or HepG2) and stellate cells (LX2), while tumor spheroid monocultures were not affected by 4μ8C. Similarly, PCNA-mRNA-expression significantly increased in human liver scaffolds engrafted with HepG2 and LX2-cells, compared to those engrafted with only tumor cells (Figure 5A). Treatment with 4μ8C significantly decreased PCNA-mRNA-expression in the LX2+HepG2 liver scaffolds, whilst not affecting those engrafted with only tumor cells. This further confirms our hypothesis that 4μ8C affects tumor cell proliferation indirectly, namely by blocking the activation of stellate cells and thus impairing the interaction between tumor and stroma.

We measured the mRNA-expression of hepatocyte-nuclear-factor-4-alpha (Hnf4-α), which is a liver function marker that is correlated to a favorable outcome for HCC-patients (43). While co-engraftment of LX2 and HepG2-cells in the liver scaffolds only lead to a marginal increase of Hnf4-α, treatment with 4μ8C significantly increased Hnf4-α-mRNA-expression, thus suggesting an overall improvement of liver function and possibly improved prognosis (Figure 5B). Immunohistochemical staining of Epcam and ki67, showed that the HCC-cells have successfully engrafted the entire surface of the scaffolds and that 4μ8C decreases proliferation (Figure 5C).

Pharmacological inhibition of IRE1α decreases tumor cell migration in stellate cell – tumor cell co-cultures

Co-culturing HepG2 and Huh7-tumor cells with LX2-cells in the transwell assays

significantly increased mRNA-expression of the pro-metastatic marker MMP9 in

HepG2-cells (Figure 6A) and MMP1 in HepG2 and Huh7-cells (Figure 6B). Adding

4µ8C significantly decreased the mRNA-expression of MMP1 in HepG2+LX2 and

Huh7+LX2 transwell co-cultures, while a non-significant decrease of MMP9 mRNA-

expression was seen in Huh7+LX2 transwell co-cultures. To assess whether this

reduction in mRNA-expression of pro-metastatic markers has a functional effect on cell

migration, a scratch wound assay was performed on confluent monolayers of mono-

cultures (HepG2 or LX2) or tumor cell (HepG2) – stellate cell (LX2) co-cultures (Figure

6C). To visualize closing of the scratch wound by each individual cell type, cells were

fluorescently labeled using CellTracker Green (tumor cells) or CellTracker Red (LX2

stellate cells) (Figure 6D). Tumor-stellate cell co-cultures were the most efficient to

close the scratch wound (Figure 6E). This was significantly inhibited when co-cultures

were treated with 4µ8C. We also observed a direct effect of 4µ8C on LX2 and HepG2-

migration, since treatment with 4µ8C lead to a significant reduction in wound closure

after 24h, compared to untreated controls. It is important to note that traditional scratch

wound assays cannot distinguish between proliferation and migration (44). To

overcome this limitation (45), we counted the individual number of cells in the middle

of the wound area (Figure 6F and G). No significant difference was seen between

HepG2 or LX2-cells within the wound area of HepG2-LX2 co-cultures after 24 hours

(Figure9F). However, 4µ8C-treatment significantly decreased migration of HepG2-

cells and LX2-cells inside the scratch wound in co-cultures, while not affecting mono-

cultures (Figure 6G).

Metastasis is usually a result of directed migration and chemotaxis toward physical

and biochemical gradients within the tumor stroma (46). We used a microfluidic-based

device for studying cell migration towards a stable gradient of chemotactic factors, such as FBS. 4µ8C significantly decreased total migration (Supplementary figure 4A-C) and directional migration towards FBS (Supplementary figure 4B and D) of HepG2-cells co-cultured with LX2-cells. Similarly, inhibition of ER-stress with 4µ8C significantly decreased total migration (Supplementary figure 4E and G) and directional migration towards FBS (Supplementary figure 4F and H) of LX2-cells co-cultured with HepG2-cells. Overall, these data suggest that stellate cells increase proliferation and pro-metastatic potential of tumor cells and blocking the IRE1α-RNase activity decreases tumor cell proliferation and migration.

Silencing of IRE1α in stellate cells decreases tumor cell proliferation and migration in co-cultures

To investigate whether the effect of blocking IRE1α is due to a direct effect on the tumor cells or because of an indirect effect via stellate cells, we transfected the stellate-line LX2 with an IRE1α-siRNA prior to co-culturing them in a transwell assay with HepG2-cells. Transfection efficiency was determined via qPCR and showed a 50% reduction in the IRE1α-mRNA-expression (Figure 7A) compared to non-transfected (Ctrl) or mock-transfected (Scr) controls. In the transwell co-culturing assay, we found that silencing IRE1α in the LX2-cells significantly decreased PCNA-mRNA-expression in HepG2-cells (Figure 7B). Silencing IRE1α in the LX2-cells lead to a significant reduction of proliferation in LX2-HepG2 co-cultures (Figure 7C) and LX2-HepG2 spheroids (Figure 7D). Immunocytochemical staining with αSMA-antibodies (Figure 7E), confirmed a significant reduction of αSMA after si-IRE1α-transfection of LX2-stellate cells in HepG2-LX2 spheroid co-cultures (Figure 7F). A scratch wound assay on HepG2-LX2 co-cultures verified that silencing of IRE1α in LX2-cells significantly

reduced wound closure compared to non-transfected and mock-transfected stellate cells (Figure 7G - H). Overall, these data confirm that blocking the IRE1α-pathway in hepatic stellate cells decreases proliferation and pro-metastatic potential of tumor cells.

DISCUSSION

There is increasing evidence that ER-stress and activation of the UPR play an essential role during hepatic inflammation and chronic liver disease. We have previously shown that inhibition of IRE1α prevents stellate cell activation and reduces liver cirrhosis *in vivo* (29). In this report, we further define a role of ER-stress and the UPR in the interaction between tumor cells and hepatic stellate cells. We also show that IRE1α could form a valuable therapeutic target to slow down the progression of hepatocellular carcinoma.

Activated stellate cells play an important role in promoting tumorigenesis and tumors are known to secrete cytokines such as TGFβ, which induce myofibroblast activation and creates an environment that sustains tumor growth (47). Since over 80% of HCC arises in a setting of chronic inflammation associated with liver fibrosis, targeting the fibrotic tumor micro-environment is often proposed as a valuable therapeutic strategy for HCC-patients (2). We and others have shown that ER-stress plays an important role in stellate cell activation and contributes to the progression of liver fibrosis (29, 48-50). The mechanisms by which the UPR promotes stellate cell activation have been attributed to regulating the expression of c-MYB (29), increasing the expression of SMAD-proteins (48) and/or by triggering autophagy (50).

In our study, we show that ER-stress plays an important role in stellate cell – tumor cell interactions and that pharmacological inhibition of IRE1α-endoribonuclease activity slows down the progression of HCC *in vivo*. We demonstrate that tumor cells induce ER-stress in hepatic stellate cells, thereby contributing to their activation and creating an environment that is supportive for tumor growth and metastasis. Activated stellate cells are known to enhance migration and proliferation of tumor cells *in vitro (11)* and *in vivo* (51), possibly by producing extracellular matrix proteins and by producing growth factors. Extracellular matrix proteins such as collagen can act as a scaffold for tumor cell migration (52), alter the expression of MMP´s (11) and induce epithelial-mesenchymal transition (53). Activated stellate cells are also an important source of hepatocyte growth factor, which promotes proliferation, cell invasion and epithelial-mesenchymal transition via the c-MET signaling pathway (54). Interestingly, blocking ER-stress in the stellate cell population reduced tumor-induced activation towards myofibroblasts, which then decreases proliferation and migration of tumor-cells in co-cultures. This suggests that targeting the microenvironment using an ER-stress inhibitor could be a promising strategy for patients with HCC.

The UPR has been described as an essential hallmark of HCC (55), although its role within tumorigenesis remains controversial (22). While a mild to moderate level to ER-stress leads to activation of the UPR and enables cancer cells to survive and adapt to adverse environmental conditions, the occurrence of severe or sustained ER-stress leads to apoptosis. Both ER-stress inhibitors as ER-stress inducers have therefore been shown to act as potential anti-cancer therapies (56). A recent study by Wu *et al*, demonstrated that IRE1α promotes progression of HCC and that hepatocyte specific ablation of IRE1α results in a decreased tumorigenesis (57). In contrast to their study,

we found a greater upregulation of actors of the IRE1α-branch within the stroma than in the tumor itself and identified that expression of ER-stress markers was mainly localized within the stellate cell population. An important difference between both studies is the mouse model that is used. While Wu *et al* used a single injection of DEN, we performed weekly injections, causing tumors to occur in a background of fibrosis, similar to what is seen in patients (30). Our *in vitro* studies with mono-cultures confirm that 4μ8C also has a direct effect on proliferation and migration of HCC cells – similar to the findings of Wu *et al* - and the response seems to depend on the tumor cell line. Adding 4μ8C to HepG2-cells significantly increased proliferation, while a significant decrease was seen in the Huh7-cells. This difference in response could be due IRE1α´s function as a key cell fate regulator. On the one hand it can induce mechanisms that restore protein homeostasis and promote cytoprotection, while on the other hand IRE1α also activates apoptotic signaling pathways. How and when IRE1α exerts its cytoprotective or its pro-apoptotic function remains largely unknown. The duration and severity of ER-stress seems to be a major contributor to the switch towards apoptosis, possibly by inducing changes in the conformational structure of IRE1α (58). The threshold at which cells experience a severe and prolonged ER-stress that would induce apoptosis could differ between different cell lines, depending on the translational capacity of the cells (e.g. ER-size, number of chaperones and the amount of degradation machinery) and the intrinsic sources that cause ER-stress (59). A study of Li *et al*, has specifically looked at how IRE1α regulates cell growth and apoptosis in HepG2-cells (60). Similar to our findings, they discovered that inhibiting IRE1α enhances cell proliferation, while over-expression of IRE1α increases the expression of polo-like kinase, which leads to apoptosis. Interestingly, polo-like kinases have divergent roles on HCC-cell growth depending on which cell line is used, which could

explain the different response to 4µ8C in Huh7 and HepG2-cells (61). Studies on glioma cells show that IRE1α regulates invasion through MMP´s (62). In line with these results, we also detected a reduction of MMP1-mRNA expression after 4µ8C-treatment and observed a direct effect on wound closure in HepG2-cells. These results indicate that ER-stress could play a direct role in regulating tumor cell invasion, in addition to its indirect effect via stellate cells.

In conclusion, the aim of this study was to define the role of ER-stress in the cross-talk between hepatic stellate cells and tumor cells in liver cancer. We show that pharmacologic inhibition of the IRE1α-signaling pathway decreases tumor burden in a DEN-induced mouse model for HCC. Using several *in vitro* 2D and 3D co-culturing methods, we identified that tumor cells induce ER-stress in hepatic stellate cells and that this contributes to their activation. Blocking ER-stress in these hepatic stellate cells prevents their activation, which then decreases proliferation and migration of tumor cells.

ACKNOWLEDGEMENTS

This research was funded through grants obtained from the Swedish Cancer Foundation (Cancerfonden, CAN2017/518 and CAN2013/1273), The Swedish children's cancer foundation (Barncancerfonden), the Swedish society for medical research (SSMF, S17-0092), the O.E. och Edla Johanssons stiftelse. These funding sources were not involved in the study design; collection, analysis and interpretation of data; writing of the report; and in the decision to submit the article for publication. We would like to thank visiting students Kim Vanhollebeke and Justine Dobbelaere for

their technical assistance; GradienTech for providing us with their CellDirector assays

and Paul O´Callaghan for his valuable input on our project.

Competing interest: The authors have no conflict of interest to report.

REFERENCES

1.   Calderaro J, Ziol M, Paradis V, Zucman-Rossi J. Molecular and histological
correlations in liver cancer. Journal of hepatology. 2019.doi: 10.1016/j.jhep.2019.06.001
2.   Coulouarn C, Clement B. Stellate cells and the development of liver cancer:
therapeutic potential of targeting the stroma. Journal of hepatology. 2014;60(6):1306-9.doi:
10.1016/j.jhep.2014.02.003
3.   Carloni V, Luong TV, Rombouts K. Hepatic stellate cells and extracellular matrix in
hepatocellular carcinoma: more complicated than ever. Liver international : official journal of
the International Association for the Study of the Liver. 2014.doi: 10.1111/liv.12465
4.   Yu G, Jing Y, Kou X, Ye F, Gao L, Fan Q, et al. Hepatic stellate cells secreted
hepatocyte growth factor contributes to the chemoresistance of hepatocellular carcinoma.
PloS one. 2013;8(9):e73312.doi: 10.1371/journal.pone.0073312
5.   Giannelli G, Villa E, Lahn M. Transforming Growth Factor-beta as a Therapeutic
Target in Hepatocellular Carcinoma. Cancer research. 2014;74(7):1890-4.doi: 10.1158/0008-
5472.CAN-14-0243
6.   Nitta T, Kim JS, Mohuczy D, Behrns KE. Murine cirrhosis induces hepatocyte
epithelial mesenchymal transition and alterations in survival signaling pathways. Hepatology.
2008;48(3):909-19.doi: 10.1002/hep.22397
7.   Dooley S, Weng H, Mertens PR. Hypotheses on the role of transforming growth
factor-beta in the onset and progression of hepatocellular carcinoma. Digestive diseases.
2009;27(2):93-101.doi: 10.1159/000218340
8.   Bogaerts E, Heindryckx F, Vandewynckel YP, Van Grunsven LA, Van Vlierberghe
H. The roles of transforming growth factor-beta, Wnt, Notch and hypoxia on liver progenitor
cells in primary liver tumours (Review). Int J Oncol. 2014;44(4):1015-22.doi:
10.3892/ijo.2014.2286
9.   Ikushima H, Miyazono K. TGFbeta signalling: a complex web in cancer progression.
Nat Rev Cancer. 2010;10(6):415-24.doi: 10.1038/nrc2853
10.   Lin N, Chen Z, Lu Y, Li Y, Hu K, Xu R. Role of activated hepatic stellate cells in
proliferation and metastasis of hepatocellular carcinoma. Hepatology research : the official
journal of the Japan Society of Hepatology. 2014.doi: 10.1111/hepr.12356
11.   Song Y, Kim SH, Kim KM, Choi EK, Kim J, Seo HR. Activated hepatic stellate cells
play pivotal roles in hepatocellular carcinoma cell chemoresistance and migration in
multicellular tumor spheroids. Sci Rep. 2016;6:36750.doi: 10.1038/srep36750
12.   Zhang F, Hao M, Jin H, Yao Z, Lian N, Wu L, et al. Canonical hedgehog signalling
regulates hepatic stellate cell-mediated angiogenesis in liver fibrosis. Br J Pharmacol.
2017;174(5):409-23.doi: 10.1111/bph.13701
13.   Taura K, De Minicis S, Seki E, Hatano E, Iwaisako K, Osterreicher CH, et al. Hepatic
stellate cells secrete angiopoietin 1 that induces angiogenesis in liver fibrosis.
Gastroenterology. 2008;135(5):1729-38.doi: 10.1053/j.gastro.2008.07.065

14.   Xia Y, Chen R, Ye SL, Sun R, Chen J, Zhao Y. Inhibition of T-cell responses by
intratumoral hepatic stellate cells contribute to migration and invasion of hepatocellular
carcinoma. Clin Exp Metastasis. 2011;28(7):661-74.doi: 10.1007/s10585-011-9399-3
15.   Zhao W, Zhang L, Xu Y, Zhang Z, Ren G, Tang K, et al. Hepatic stellate cells
promote tumor progression by enhancement of immunosuppressive cells in an orthotopic liver
tumor mouse model. Lab Invest. 2014;94(2):182-91.doi: 10.1038/labinvest.2013.139
16.   Caja L, Dituri F, Mancarella S, Caballero-Diaz D, Moustakas A, Giannelli G, et al.
TGF-beta and the Tissue Microenvironment: Relevance in Fibrosis and Cancer. Int J Mol Sci.
2018;19(5).doi: 10.3390/ijms19051294
17.   Lu Y, Lin N, Chen Z, Xu R. Hypoxia-induced secretion of platelet-derived growth
factor-BB by hepatocellular carcinoma cells increases activated hepatic stellate cell
proliferation, migration and expression of vascular endothelial growth factor-A. Mol Med
Rep. 2015;11(1):691-7.doi: 10.3892/mmr.2014.2689
18.   Schroder M, Kaufman RJ. ER stress and the unfolded protein response. Mutat Res-
Fund Mol M. 2005;569(1-2):29-63.doi: 10.1016/j.mrfmmm.2004.06.056
19.   Sovolyova N, Healy S, Samali A, Logue SE. Stressed to death - mechanisms of ER
stress-induced cell death. Biol Chem. 2014;395(1):1-13.doi: 10.1515/hsz-2013-0174
20.   Liang H, Xiao J, Zhou Z, Wu J, Ge F, Li Z, et al. Hypoxia induces miR-153 through
the IRE1alpha-XBP1 pathway to fine tune the HIF1alpha/VEGFA axis in breast cancer
angiogenesis. Oncogene. 2018.doi: 10.1038/s41388-017-0089-8
21.   Li XX, Zhang HS, Xu YM, Zhang RJ, Chen Y, Fan L, et al. Knockdown of IRE1alpha
inhibits colonic tumorigenesis through decreasing beta-catenin and IRE1alpha targeting
suppresses colon cancer cells. Oncogene. 2017;36(48):6738-46.doi: 10.1038/onc.2017.284
22.   Vandewynckel YP, Laukens D, Geerts A, Bogaerts E, Paridaens A, Verhelst X, et al.
The paradox of the unfolded protein response in cancer. Anticancer Res. 2013;33(11):4683-
94.doi:
23.   Heindryckx F, Li JP. Role of proteoglycans in neuro-inflammation and central
nervous system fibrosis. Matrix Biol. 2018;68-69:589-601.doi: 10.1016/j.matbio.2018.01.015
24.   Bandla H, Dasgupta D, Mauer AS, Nozickova B, Kumar S, Hirsova P, et al. Deletion
of endoplasmic reticulum stress-responsive co-chaperone p58(IPK) protects mice from diet-
induced steatohepatitis. Hepatology research : the official journal of the Japan Society of
Hepatology. 2018.doi: 10.1111/hepr.13052
25.   Kwanten WJ, Vandewynckel YP, Martinet W, De Winter BY, Michielsen PP, Van
Hoof VO, et al. Hepatocellular autophagy modulates the unfolded protein response and
fasting-induced steatosis in mice. American journal of physiology Gastrointestinal and liver
physiology. 2016;311(4):G599-G609.doi: 10.1152/ajpgi.00418.2015
26.   Lebeaupin C, Vallee D, Hazari Y, Hetz C, Chevet E, Bailly-Maitre B. Endoplasmic
reticulum stress signalling and the pathogenesis of non-alcoholic fatty liver disease. Journal of
hepatology. 2018;69(4):927-47.doi: 10.1016/j.jhep.2018.06.008
27.   Li J, He J, Fu Y, Hu X, Sun LQ, Huang Y, et al. Hepatitis B virus X protein inhibits
apoptosis by modulating endoplasmic reticulum stress response. Oncotarget.
2017;8(56):96027-34.doi: 10.18632/oncotarget.21630
28.   Sasaki M, Yoshimura-Miyakoshi M, Sato Y, Nakanuma Y. A possible involvement of
endoplasmic reticulum stress in biliary epithelial autophagy and senescence in primary biliary
cirrhosis. J Gastroenterol. 2015;50(9):984-95.doi: 10.1007/s00535-014-1033-0
29.   Heindryckx F, Binet F, Ponticos M, Rombouts K, Lau J, Kreuger J, et al. Endoplasmic
reticulum stress enhances fibrosis through IRE1alpha-mediated degradation of miR-150 and
XBP-1 splicing. EMBO Mol Med. 2016;8(7):729-44.doi: 10.15252/emmm.201505925

30.   Heindryckx F, Mertens K, Charette N, Vandeghinste B, Casteleyn C, Van Steenkiste C, et al. Kinetics of angiogenic changes in a new mouse model for hepatocellular carcinoma. Mol Cancer. 2010;9:219.doi: 10.1186/1476-4598-9-219

31.   Heindryckx F, Bogaerts E, Coulon SH, Devlies H, Geerts AM, Libbrecht L, et al. Inhibition of the placental growth factor decreases burden of cholangiocarcinoma and hepatocellular carcinoma in a transgenic mouse model. Eur J Gastroenterol Hepatol. 2012;24(9):1020-32.doi: 10.1097/MEG.0b013e3283554219

32.   Calitz C, Hamman JH, Fey SJ, Viljoen AM, Gouws C, Wrzesinski K. A sub-chronic Xysmalobium undulatum hepatotoxicity investigation in HepG2/C3A spheroid cultures compared to an in vivo model. J Ethnopharmacol. 2019;239:111897.doi: 10.1016/j.jep.2019.111897

33.   Mazza G, Al-Akkad W, Telese A, Longato L, Urbani L, Robinson B, et al. Rapid production of human liver scaffolds for functional tissue engineering by high shear stress oscillation-decellularization. Sci Rep. 2017;7(1):5534.doi: 10.1038/s41598-017-05134-1

34.   Thanapirom K, Frenguelli L, Al-Akkad W, Zhang ZZ, Pinzani M, Mazza G, et al. Optimization and validation of a novel three-dimensional co-culture system in decellularized human liver scaffold for the study of liver fibrosis and cancer. Journal of hepatology. 2019;70:E24-E.doi: Doi 10.1016/S0618-8278(19)30042-8

35.   Seok JY, Na DC, Woo HG, Roncalli M, Kwon SM, Yoo JE, et al. A fibrous stromal component in hepatocellular carcinoma reveals a cholangiocarcinoma-like gene expression trait and epithelial-mesenchymal transition. Hepatology. 2012;55(6):1776-86.doi: 10.1002/hep.25570

36.   Rouillard AD, Gundersen GW, Fernandez NF, Wang Z, Monteiro CD, McDermott MG, et al. The harmonizome: a collection of processed datasets gathered to serve and mine knowledge about genes and proteins. Database (Oxford). 2016;2016.doi: 10.1093/database/baw100

37.   Subramanian A, Tamayo P, Mootha VK, Mukherjee S, Ebert BL, Gillette MA, et al. Gene set enrichment analysis: a knowledge-based approach for interpreting genome-wide expression profiles. Proc Natl Acad Sci U S A. 2005;102(43):15545-50.doi: 10.1073/pnas.0506580102

38.   https://www.proteinatlas.org/ENSG00000070540-WIPI1/pathology/liver+cancer.doi: . The Human Protein Atlas. 2019;V19.doi:

39.   https://www.proteinatlas.org/ENSG00000160691-SHC1/pathology/liver+cancer. The Human Protein Atlas.V19.doi:

40.   https://www.proteinatlas.org/ENSG00000068971-PPP2R5B/pathology/liver+cancer#imid_9419368. The Human Protein Atlas.V19.doi:

41.   https://www.proteinatlas.org/ENSG00000044574-HSPA5/pathology/liver+cancer. The Human Protein Atlas.V19.doi:

42.   Uhlen M, Fagerberg L, Hallstrom BM, Lindskog C, Oksvold P, Mardinoglu A, et al. Proteomics. Tissue-based map of the human proteome. Science. 2015;347(6220):1260419.doi: 10.1126/science.1260419

43.   Hang HL, Liu XY, Wang HT, Xu N, Bian JM, Zhang JJ, et al. Hepatocyte nuclear factor 4A improves hepatic differentiation of immortalized adult human hepatocytes and improves liver function and survival. Experimental cell research. 2017;360(2):81-93.doi: 10.1016/j.yexcr.2017.08.020

44.   Cormier N, Yeo A, Fiorentino E, Paxson J. Optimization of the Wound Scratch Assay to Detect Changes in Murine Mesenchymal Stromal Cell Migration After Damage by Soluble Cigarette Smoke Extract. J Vis Exp. 2015(106):e53414.doi: 10.3791/53414

45.   Bise R, Kanade T, Yin Z, Huh SI. Automatic cell tracking applied to analysis of cell migration in wound healing assay. Conf Proc IEEE Eng Med Biol Soc. 2011;2011:6174-9.doi: 10.1109/IEMBS.2011.6091525

46.   Oudin MJ, Weaver VM. Physical and Chemical Gradients in the Tumor Microenvironment Regulate Tumor Cell Invasion, Migration, and Metastasis. Cold Spring Harb Symp Quant Biol. 2016;81:189-205.doi: 10.1101/sqb.2016.81.030817

47.   Heindryckx F, Gerwins P. Targeting the tumor stroma in hepatocellular carcinoma. World J Hepatol. 2015;7(2):165-76.doi: 10.4254/wjh.v7.i2.165

48.   Koo JH, Lee HJ, Kim W, Kim SG. Endoplasmic Reticulum Stress in Hepatic Stellate Cells Promotes Liver Fibrosis via PERK-Mediated Degradation of HNRNPA1 and Up-regulation of SMAD2. Gastroenterology. 2016;150(1):181-93 e8.doi: 10.1053/j.gastro.2015.09.039

49.   Hernandez-Gea V, Hilscher M, Rozenfeld R, Lim MP, Nieto N, Werner S, et al. Endoplasmic reticulum stress induces fibrogenic activity in hepatic stellate cells through autophagy. Journal of hepatology. 2013;59(1):98-104.doi: 10.1016/j.jhep.2013.02.016

50.   Kim RS, Hasegawa D, Goossens N, Tsuchida T, Athwal V, Sun X, et al. The XBP1 Arm of the Unfolded Protein Response Induces Fibrogenic Activity in Hepatic Stellate Cells Through Autophagy. Sci Rep. 2016;6:39342.doi: 10.1038/srep39342

51.   Amann T, Bataille F, Spruss T, Muhlbauer M, Gabele E, Scholmerich J, et al. Activated hepatic stellate cells promote tumorigenicity of hepatocellular carcinoma. Cancer Sci. 2009;100(4):646-53.doi: 10.1111/j.1349-7006.2009.01087.x

52.   Han W, Chen S, Yuan W, Fan Q, Tian J, Wang X, et al. Oriented collagen fibers direct tumor cell intravasation. Proceedings of the National Academy of Sciences of the United States of America. 2016;113(40):11208-13.doi: 10.1073/pnas.1610347113

53.   Kumar S, Das A, Sen S. Extracellular matrix density promotes EMT by weakening cell-cell adhesions. Mol Biosyst. 2014;10(4):838-50.doi: 10.1039/c3mb70431a

54.   Liu WT, Jing YY, Yu GF, Chen H, Han ZP, Yu DD, et al. Hepatic stellate cell promoted hepatoma cell invasion via the HGF/c-Met signaling pathway regulated by p53. Cell Cycle. 2016;15(7):886-94.doi: 10.1080/15384101.2016.1152428

55.   Shuda M, Kondoh N, Imazeki N, Tanaka K, Okada T, Mori K, et al. Activation of the ATF6, XBP1 and grp78 genes in human hepatocellular carcinoma: a possible involvement of the ER stress pathway in hepatocarcinogenesis. Journal of hepatology. 2003;38(5):605-14.doi:

56.   Corazzari M, Gagliardi M, Fimia GM, Piacentini M. Endoplasmic Reticulum Stress, Unfolded Protein Response, and Cancer Cell Fate. Front Oncol. 2017;7:78.doi: 10.3389/fonc.2017.00078

57.   Wu Y, Shan B, Dai J, Xia Z, Cai J, Chen T, et al. Dual Role for Inositol-requiring Enzyme 1alpha in Promoting the Development of Hepatocellular Carcinoma during Diet-induced Obesity. Hepatology. 2018.doi: 10.1002/hep.29871

58.   Ghosh R, Wang L, Wang ES, Perera BG, Igbaria A, Morita S, et al. Allosteric inhibition of the IRE1alpha RNase preserves cell viability and function during endoplasmic reticulum stress. Cell. 2014;158(3):534-48.doi: 10.1016/j.cell.2014.07.002

59.   Cubillos-Ruiz JR, Bettigole SE, Glimcher LH. Tumorigenic and Immunosuppressive Effects of Endoplasmic Reticulum Stress in Cancer. Cell. 2017;168(4):692-706.doi: 10.1016/j.cell.2016.12.004

60.   Li X, Zhu H, Huang H, Jiang R, Zhao W, Liu Y, et al. Study on the effect of IRE1a on cell growth and apoptosis via modulation PLK1 in ER stress response. Molecular and cellular biochemistry. 2012;365(1-2):99-108.doi: 10.1007/s11010-012-1248-4

61.   Pellegrino R, Calvisi DF, Ladu S, Ehemann V, Staniscia T, Evert M, et al. Oncogenic
and tumor suppressive roles of polo-like kinases in human hepatocellular carcinoma.
Hepatology. 2010;51(3):857-68.doi: 10.1002/hep.23467
62.   Auf G, Jabouille A, Guerit S, Pineau R, Delugin M, Bouchecareilh M, et al. Inositol-
requiring enzyme 1alpha is a key regulator of angiogenesis and invasion in malignant glioma.
Proceedings of the National Academy of Sciences of the United States of America.
2010;107(35):15553-8.doi: 10.1073/pnas.0914072107
8

Figure legends

Fig. 1. Inhibiting IRE1α reduces tumor burden *in vivo*. *(A)* Representative images of liver slides stained with hematoxylin and eosin (H&E), Sirius red and αSMA-antibodies. ***(B)*** tumor burden of mice with DEN-induced HCC treated with 4μ8C or vehicle-treated controls. ***(C)*** Quantification of percentage of collagen and ***(D)*** αSMA on liver slides. ***(E)*** mRNA expression of PCNA in liver tissue from mice with HCC treated with 4μ8C ***(F)*** Heatmap showing protein expression levels in healthy liver, DEN-induced HCC and DEN-induced HCC treated with 4μ8C. P-values were calculated via the Student´s T-test, scale bars = 120μm.

Fig. 2. Tumor cells secrete factors that induce ER-stress in stellate cells, which contributes to their activation. *(A)* mRNA-expression of ER-stress markers CHOP, ***(B)*** spliced XBP1, ***(C)*** BiP in stellate cells (LX2) co-cultured with cancer cells (HepG2 or Huh7) and treated with 4μ8C or control. ***(D)*** Detection of spliced (XBP1s) and unspliced XBP1 (XBP1u) visualized by digestion of XBP1u by *Pst-I*. ***(E)*** protein expression of p-IRE1α and vinculin in stellate cells (LX2) co-cultured with cancer cells (HepG2 or Huh7) in transwell assays and treated with 4μ8C or control. ***(F)*** mRNA-expression of stellate cell activation markers αSMA and ***(G)*** collagen in LX2-cells co-cultured with HepG2 or Huh7-cells and treated with or without 4μ8C. P-values were calculated via the Student´s T-test.

Fig. 3. Inhibiting IRE1α decreases stellate cell activation in human liver 3D scaffolds engrafted with stellate cells and tumor cells. *(A)* Representative images of H&E and Sirius red stained slides of decellularized human liver scaffolds engrafted with LX2 stellate cells and HepG2-tumor cells treated with 4μ8C or control. ***(B)*** mRNA-expression of the stellate cells activation marker collagen and ER-stress markers BiP, spliced XBP-1 (XBP1-S) and CHOP in liver scaffolds engrafted with stellate cells (LX2) and cancer cells (HepG2), treated with 4μ8C or control. P-values were calculated via the Student´s T-test, scale bars = 100μm.

**Fig. 4. Inhibition of IRE1α decreases tumor cell proliferation. *(A)* PCNA mRNA-expression of HepG2 or Huh7-cells grown with LX2-cells in transwell inserts and treated with the IRE1α-inhibitor 4µ8C or control. *(B)* Relative cell number of LX2 and HepG2 or *(C)* LX2 and Huh7-cells treated with 4µ8C or control. *(D)* Representative images of tumor cells (HepG2 or Huh7) and LX2-stellate cells stained with antibodies against the HCC-marker Epcam and the proliferation marker ki67. *(E)* Cell proliferation of HepG2 or HepG2+LX2 spheroids and *(F)* Huh7 or Huh7+LX2 spheroids treated with 4µ8C or control. P-values were calculated via the Student´s T-test, scale bars = 50µm.

**Fig. 5. Inhibition of IRE1α decreases cell proliferation and improves liver function in human liver scaffolds engrafted with stellate cells and tumor cells. *(A)* PCNA and *(B)* Hnf4a expression of human liver scaffolds engrafted with HepG2-tumor cells and LX2-stellate cells, treated with 4µ8C or control. *(C)* Representative images of tumor cells (HepG2) and LX2-stellate cells stained with antibodies against the HCC-marker Epcam and the proliferation marker ki67. P-values were calculated via the Student´s T-test, scale bars = 100µm.

**Fig. 6. Inhibition of IRE1α decreases cell migration. *(A)* mRNA-expression of pro-metastatic markers MMP9 and *(B)* MMP1 in HepG2 and Huh7-cells co-cultured with LX2-cells and treated with 4µ8C or control. *(C)* Scratch wound on HepG2-cells and LX2-cells treated with 4µ8C or control. *(D)* Images of Cell Tracker stained HepG2-cells (Green) and LX2-cells (Red) invading the scratch area. *(E)* Quantification of wound size in HepG2-cells and LX2-cells treated with 4µ8C or control. *(F)* Number of HepG2-cells and LX2-cells invading the scratch wound after 24h in co-cultures and *(G)* mono-cultures. P-values were calculated via the Student´s T-test, scale bars = 120µm.

**Fig. 7. Silencing IRE1α in LX2-cells mimics 4µ8C. *(A)* IRE1α-mRNA-expression of LX2-cells transfected with IRE1α-siRNA (si-IRE1α), mock-transfected (Scr) or untransfected (Ctrl). *(B)* PCNA-mRNA-expression of HepG2-cells co-cultured with IRE1α-silenced LX2-cells or controls *(C).* Relative cell numbers in co-cultures of HepG2-cells and IRE1α-silenced LX2-cells or controls. *(D)* Proliferation of spheroids of HepG2-cells and IRE1α-silenced LX2-cells or controls *(E)* Images and *(F)* quantification

of αSMA-stained spheroids with HepG2-cells and IRE1α-silenced LX2-cells or controls. **(G)** Images and **(H)** quantification of scratch wound of HepG2-cells co-cultured with IRE1α-silenced LX2-cells or controls. P-values were calculated via the Student´s T-test, scale bars = 50μm (E) or 120μm (G).

Supplementary figure 1. Activation of the unfolded protein response is mainly located in the stroma of mice with HCC. *(A)* mRNA-expression of the ER-stress marker BiP in tumor and surrounding non-tumoral liver tissue in mice with DEN-induced HCC treated with or without the IRE1α-inhibitor 4μ8C or healthy mice. *(B)* Representative western blot image showing protein expression of BiP in liver tissue from mice with HCC treated with or without 4μ8C **(C)** Quantification of p-IRE1α staining on murine liver sections. *(D)* Co-staining of liver tissue with antibodies against αSMA and p-IRE1α. *(E)* Co-staining of liver tissue with antibodies against αSMA and BiP. P-values were calculated via the Student´s T-test, scale bars = 50μm.

Supplementary figure 2. Activation of the unfolded protein response pathway is increased in patients with fibrotic HCC. *(A)* Heat map showing gene-set enrichment analysis results from samples from fibrous HCC versos non-fibruous HCC. *(C)* Immunohistochemically stained liver biopsies from HCC-patients obtained from the human protein atlas, using antibodies against IRE1α-mediated actors of the unfolded protein response: WIPI1, SHC1, PPP2R5B and BiP. (D) Kaplan-Meier survival curves of HCC-patients with high or low expression of WIPI1, SHC1, PPP2R5B and BiP. P-values were calculated via a Log-Rank test.

Supplementary Figure 3. Secretion of TGFβ by tumor cells activates stellate cells and induces ER-stress. *(A)* concentration of TGFβ in medium from tumor cells (HepG2 or Huh7) grown in mono-culture or co-cultured with LX2-stellate cells, treated with 4μ8C or control. *(B)* concentration of TGFβ in medium from liver scaffolds engrafted with stellate cells *(C)* (LX2) and tumor cells (HepG2) treated with 4μ8C or control. mRNA-expression of the ER-stress markers CHOP, *(D)* spliced XBP1, *(E)* unspliced XBP1 and *(F)* BiP in hepatic stellate cells (LX2) grown as mono-culture or in co-cultures with the cancer cell lines HepG2 and Huh7 treated with the TGFβ receptor inhibitor SB-431541 or control**. (G)** mRNA-

expression of stellate cell activation markers αSMA and (**H**) collagen in LX2-cells grown with HepG2 or

Huh7-cells and treated with SB-431541 or control. P-values were calculated via the Student´s T-test.

Supplementary Figure 4. Inhibiting IRE1α decreases chemotaxis. *(A)* migration plots of LX2-cells

co-cultured with HepG2-cells exposed to an FBS-gradient (increasing towards the right) and treated

with control or *(B)* 4µ8C *(C)* Quantification of total migration and *(D)* directional migration of LX2-cells

(co-cultured with HepG2-cells) towards an FBS-gradient with or without 4µ8C. *(E)* Migration plots of

HepG2-cells co-cultured with LX2-stellate cells and exposed to an FBS-gradient and treated with control

or *(F)* 4µ8C. *(G)* Quantification of total migration and *(H)* directional migration of HepG2-cells (co-

cultured with LX2-cells) towards an FBS-gradient with or without 4µ8C. P-values were calculated via

the Student´s T-test.

Table 1: A proteomics array using the Olink Mouse Exploratory assay

Protein name	Biological process	CTL		Den		DEN+4u8c		Statistical significance		
		Aver age	St. Dev	Aver age	St. Dev	Aver age	St. Dev	DEN vs Ctrl	DEN vs 4u8C	Ctrl vs 4u8C
Clmp	Not prognostic in HCC	1,68	0,14	2,97	1,00	2,48	0,64	*		
Yes1	HCC promotor	7,11	0,29	7,51	0,20	7,44	0,19	*		
Foxo1	Tumor suppressor	4,15	0,06	4,12	0,73	3,87	0,49			
Pla2g4a	HCC promotor	3,42	0,38	5,70	1,36	5,04	0,80	*		*
Prdx5	HCC promotor	7,37	0,49	7,23	0,26	6,67	0,34		*	
Tgfa	Tumor growth factor	5,36	0,52	6,81	0,64	6,93	0,88	*		*
Epo	Unfavorable prognotic marker	3,20	0,34	3,71	0,35	3,37	0,33			
Axin1	HCC promotor	4,24	0,38	4,80	0,37	4,39	0,35			
Fst	HCC promotor	5,87	0,31	8,04	0,73	7,50	0,71	*		*
Nadk	Not prognostic in HCC	10,10	0,13	10,14	0,18	10,30	0,27			
Snap29	Not prognostic in HCC	7,70	0,32	7,87	0,32	7,62	0,30			
S100a4	HCC promotor	2,73	0,74	7,01	0,62	6,85	0,97	*		*
Kitlg	Metastasis	2,48	0,42	3,74	0,62	3,31	0,98	*		
Gfra1	HCC promotor	4,40	0,35	5,07	0,40	4,92	0,39	*		
Ppp1r2	Not prognostic in HCC	4,37	0,16	4,86	0,46	4,47	0,43			
Cyr61	HCC promotor	2,40	0,53	4,14	1,64	3,13	1,22	*		
Ahr	Not prognostic in HCC	6,95	0,46	7,68	0,74	7,38	0,64			
Ccl2	HCC promotor	4,59	0,58	9,69	2,04	8,93	1,56	*		*

Qdpr	Not prognostic in HCC	7,71	0,11	7,72	0,14	7,54	0,15			
Fas	HCC promotor	8,66	0,18	8,83	0,18	8,70	0,18			
Riox2	HCC promotor	7,10	0,15	7,71	0,38	7,59	0,14	*		*
Epcam	HCC promotor	1,56	0,33	3,16	1,14	3,27	0,89	*		
Ccl3	Prognostic marker	1,49	0,39	4,42	1,86	3,73	1,07	*		*
Crim1	HCC promotor	2,46	0,28	3,71	1,09	3,21	0,56	*		*
Hgf	Tumor growth factor	6,69	0,35	7,94	1,01	7,41	0,71	*		
Sez6l2	HCC promotor	-0,29	0,15	0,61	0,53	0,19	0,29	*		
Il1a	Inflammation and fibrosis	6,65	0,51	8,35	0,65	7,62	0,54	*		*
Ddah1	HCC promotor	8,04	0,22	8,18	0,05	7,84	0,18		*	
Acvrl1	Not prognostic in HCC	2,09	0,18	3,44	1,31	2,81	0,47			
Cxcl9	Inflammation and fibrosis	3,68	0,86	7,71	1,68	6,65	1,58	*		*
Map2k6	Not prognostic in HCC	7,75	0,15	7,98	0,41	7,88	0,28			
Casp3	Tumor surrpressor	9,22	0,19	9,74	0,35	9,43	0,26			
Pdgfb	Tumor growth factor	3,52	0,31	4,96	1,27	3,97	0,40	*		
Igsf3	Unfavorable prognotic marker	3,12	0,28	4,19	0,82	3,64	0,72			
Cxcl1	HCC promotor	3,77	0,40	5,74	0,78	5,06	0,51	*		*
Pak4	HCC promotor	3,47	0,42	4,39	0,68	3,93	0,54			
Lpl	Not prognostic in HCC	1,66	0,40	2,44	0,45	2,02	0,60			
Dctn2	Unfavorable prognotic marker	5,48	1,31	5,67	0,70	4,98	0,55			
Ntf3	Not prognostic in HCC	2,16	0,27	2,80	0,71	2,27	0,40			
Tnfsf12	HCC promotor	5,28	0,35	6,00	0,76	5,59	0,62			
Ccl20	Unfavorable prognotic marker	5,20	0,34	5,92	0,81	5,53	0,66			
Fli1	HCC promotor	1,91	0,22	3,73	1,38	2,98	0,83			
Tpp1	Unfavorable prognotic marker	3,67	0,38	4,24	0,64	3,73	0,50			
Parp1	Unfavorable prognotic marker	10,30	0,72	10,93	0,49	10,51	0,62			

1 **Table 2: Genes the contributed to the core-enrichment of the GSEA**

Probe	Description	Rank Gene list	Rank Metric score	Core enrichment	UPR branch
ASNS	asparagine synthetase (glutamine-hydrolyzing) [Source:HGNC Symbol;Acc:HGNC:753]	207	0.940	Yes	Perk
PPP2R5B	protein phosphatase 2 regulatory subunit B'beta [Source:HGNC Symbol;Acc:HGNC:9310]	423	0.821	Yes	Ire1a
CCL2	C-C motif chemokine ligand 2 [Source:HGNC Symbol;Acc:HGNC:10618]	847	0.689	Yes	Ire1a and Perk
EXOSC9	exosome component 9 [Source:HGNC Symbol;Acc:HGNC:9137]	1004	0.654	Yes	Ire1a and Perk
WIPI1	WD repeat domain, phosphoinositide interacting 1 [Source:HGNC Symbol;Acc:HGNC:25471]	1022	0.649	Yes	Ire1a
KDELR3	KDEL endoplasmic reticulum protein retention receptor 3 [Source:HGNC Symbol;Acc:HGNC:6306]	1106	0.635	Yes	Ire1a
SHC1	SHC adaptor protein 1 [Source:HGNC Symbol;Acc:HGNC:10840]	2691	0.432	Yes	Ire1a
TPP1	tripeptidyl peptidase 1 [Source:HGNC Symbol;Acc:HGNC:2073]	2884	0.414	Yes	Ire1a
HDGF	heparin binding growth factor [Source:HGNC Symbol;Acc:HGNC:4856]	3235	0.386	Yes	Ire1a
TLN1	talin 1 [Source:HGNC Symbol;Acc:HGNC:11845]	3264	0.384	Yes	Ire1a
EXTL3	exostosin like glycosyltransferase 3 [Source:HGNC Symbol;Acc:HGNC:3518]	3488	0.365	Yes	Ire1a
TSPYL2	TSPY like 2 [Source:HGNC Symbol;Acc:HGNC:24358]	3680	0.350	Yes	Ire1a
MBTPS1	membrane bound transcription factor peptidase, site 1 [Source:HGNC Symbol;Acc:HGNC:15456]	3996	0.327	Yes	Atf6
PDIA5	protein disulfide isomerase family A member 5 [Source:HGNC Symbol;Acc:HGNC:24811]	4530	0.294	Yes	Ire1a
DCTN1	dynactin subunit 1 [Source:HGNC Symbol;Acc:HGNC:2711]	4638	0.287	Yes	Ire1a
DNAJC3	DnaJ heat shock protein family (Hsp40) member C3 [Source:HGNC Symbol;Acc:HGNC:9439]	4761	0.281	Yes	Ire1a
SULT1A4	sulfotransferase family 1A member 4 [Source:HGNC Symbol;Acc:HGNC:30004]	4938	0.272	Yes	Ire1a
PARN	poly(A)-specific ribonuclease [Source:HGNC Symbol;Acc:HGNC:8609]	5037	0.266	Yes	Perk
ADD1	adducin 1 [Source:HGNC Symbol;Acc:HGNC:243]	5375	0.250	Yes	Ire1a
ERN1	endoplasmic reticulum to nucleus signaling 1 [Source:HGNC Symbol;Acc:HGNC:3449]	5411	0.248	Yes	Ire1a

2

3 **Supplementary Table 1. Primer sequences**

4

5 **Supplementary methods 1: histology and immunohistochemistry**

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7