1

Regulation of apical constriction via microtubule- and Rab11-dependent apical transport during tissue invagination

Thao Phuong Le[1] and SeYeon Chung[1, *]

[1] Department of Biological Sciences, Louisiana State University, Baton Rouge, LA 70803, USA

*Author for correspondence: seyeonchung@lsu.edu

Running Head: MTs and Rab11 regulate apical constriction

Abstract

The formation of an epithelial tube is a fundamental process for organogenesis. During *Drosophila* embryonic salivary gland (SG) invagination, Folded gastrulation (Fog)-dependent Rho-associated kinase (Rok) promotes contractile apical myosin formation to drive apical constriction. Microtubules (MTs) are also crucial for this process and are required for forming and maintaining apicomedial myosin. However, the underlying mechanism that coordinates actomyosin and MT networks still remains elusive. Here, we show that MT-dependent intracellular trafficking regulates apical constriction during SG invagination. Key components involved in protein trafficking, such as Rab11 and Nuclear fallout (Nuf), are apically enriched near the SG invagination pit in a MT-dependent manner. Disruption of the MT networks or knockdown of *Rab11* impairs apicomedial myosin formation and apical constriction. We show that MTs and Rab11 are required for apical enrichment of the Fog ligand as well as the apical determinant protein Crumbs (Crb) and the key adherens junction protein E-Cadherin (E-Cad). Targeted knockdown of *crb* or *E-Cad* in the SG disrupts apical myosin networks and results in apical constriction defects. Our data suggests a role of MT- and Rab11-dependent intracellular trafficking in regulating actomyosin networks and cell junctions, to coordinate cell behaviors during tubular organ formation.

Introduction

Formation of three-dimensional tubes by invagination of flat epithelial sheets is a fundamental process in forming organs such as the lungs and kidneys (Andrew and Ewald, 2010). To enter the third dimension, cells must change their shapes and positions relative to each other. A major cellular process during epithelial tube formation is apical constriction, a universal cell shape change that is linked to tissue bending, folding and invagination (Martin and Goldstein, 2014; Sawyer et al., 2010). During apical constriction, the apical side of an epithelial cell constricts, causing a columnar or cuboidal cell to become wedge-shaped (Martin and Goldstein, 2014; Sawyer et al., 2010). Manipulation of apical constriction in a group of cells impacts both local and global tissue shape directly, suggesting a critical role for apical constriction in forming proper tissue architecture (Chung et al., 2017; Guglielmi et al., 2015; Izquierdo et al., 2018).

Apical constriction is driven by actin filament (F-actin) networks and the molecular motor non-muscle myosin II (hereafter referred to as myosin). Over the past decade, important functions of different actomyosin structures in epithelial tissue morphogenesis have been discovered. Particularly, studies in *Drosophila* revealed a distinct population of pulsatile apical medial actomyosin (hereafter referred to as apicomedial myosin) that generates a pulling force that exerts on adherens junctions to drive apical constriction (Booth et al., 2014; Chung et al., 2017; Martin et al., 2009; Rauzi et al., 2010). Further studies in early *Drosophila* embryos discovered that apicomedial myosin is created in response to signaling by the Folded gastrulation (Fog) ligand and its G protein-coupled receptors (GPCRs) (Kerridge et al., 2016; Manning et al., 2013) and is regulated by apical Rho-associated kinase (Rok) (Mason et al., 2013).

Apical junctions and apical determinants also have an important role in the formation of functional actomyosin complexes during developmental processes. During *Drosophila* dorsal closure, the apical polarity regulators Par-6, aPKC and Bazooka/Par3 (Baz/Par3) (altogether known as the Par complex) regulate pulsed actomyosin contractions in amnioserosa cells (David et al., 2010). In the *Drosophila* embryonic trachea, the apical

protein Crumbs (Crb) is required for proper organization of the actomyosin complex

(Letizia et al., 2011).

Emerging evidence suggests that microtubules (MTs) play a critical role in tissue

invagination (Booth et al., 2014; Ko et al., 2019). MTs serve as tracks in intracellular

transport (Aguilar-Aragon et al., 2020; Khanal et al., 2016; Le Droguen et al., 2015),

raising the possibility that MTs regulate apical constriction through the endo- and

exocytosis of membrane receptors and adhesion molecules. Several lines of evidence

also suggest the importance of the endocytic pathway in apical constriction in both in vitro

and in vivo systems. In *Drosophila* S2 cells, RhoGEF2 travels to the cell cortex by

interaction with the MT plus-end protein EB1 to stimulate cell contraction (Rogers et al.,

2004). During *Xenopus* gastrulation, disrupting endocytosis with dominant-negative

dynamin or Rab5 perturbs apical constriction and invagination of cell sheets (Lee and

Harland, 2010). During *Drosophila* gastrulation, the apical surface of cells is reshaped via

Rab35 and RabGEF Sbf, which direct the plasma membrane to Rab11-positive recycling

endosomes through a dynamic interaction with Rab5 endosomes to reshape actomyosin

networks (Miao et al., 2019). Moreover, in the developing neural tube in *Xenopus*,

asymmetric enrichment of Rab11 at the medial apical junctions is critical for apical

constriction, suggesting that membrane trafficking has a key role in apical constriction

(Ossipova et al., 2014). However, exactly how vesicle trafficking, MTs, and actomyosin

networks are linked during tissue invagination remains to be discovered.

To determine how these three cellular attributes contribute to tissue invagination, we use

the *Drosophila* embryonic SG as a model. We and others showed that apical constriction

is regulated in a highly coordinated and spatiotemporally controlled manner during SG

tube formation (Booth et al., 2014; Chung et al., 2017; Myat and Andrew, 2000; Sanchez-

Corrales et al., 2018). The *Drosophila* embryo forms two SG tubes via invagination of two

epithelial placodes on the ventral surface (Chung et al., 2014; Myat and Andrew, 2002).

Before invagination, a small group of cells in the dorsal/posterior region of each SG

placode begin to constrict. As those cells internalize to form the invagination pit, more

cells anterior to the pit undergo clustered apical constriction in a Fog signaling-dependent

manner (Chung et al., 2017) (Figure 1A). In the absence of *fog*, SG cells fail to accumulate

Rok and myosin in the apicomedial region of the cells (Chung et al., 2017). MTs aid in the formation and maintenance of the apicomedial myosin network during SG invagination (Booth et al., 2014). The MT cytoskeleton near the invagination pit forms a network of longitudinal MT bundles, with the minus ends of MTs facing the apical domain of the cells and interacting with the apicomedial myosin (Booth et al., 2014). Disruption of MTs causes loss of apicomedial myosin and disrupted apical constriction during SG invagination (Booth et al., 2014). However, it is still unknown how intracellular trafficking affects these two processes.

In this study, we demonstrate that key proteins involved in intracellular trafficking - including Rab11, its binding partner Nuclear fallout (Nuf), and dynein heavy chain - are enriched in the apical domain of SG cells during invagination. Moreover, disruption of MTs results in mislocalization of Rab11 to the basolateral region of SG cells. Reducing *Rab11* in the SG leads to a decrease and a dispersal of Rok and myosin in the apical domain and causes uncoordinated apical constriction. Our data suggest that apical localization of the Fog ligand, the apical determinant protein Crb, and the key adherens junction protein E-Cadherin (E-Cad) is compromised when the MT networks are disrupted or *Rab11* is knocked down. We further show that reducing *crb* or *E-Cad* in the SG leads to defects that are reminiscent of *Rab11* knockdown. Altogether, our work mechanistically links MT- and Rab11-dependent intracellular trafficking to the regulation of actomyosin networks during tubular organ formation.

Results

Intracellular trafficking components are apically enriched near the invagination pit

To test a role for the intracellular trafficking machinery in spatially biased signaling activation and protein accumulation during SG invagination, we analyzed the subcellular localization of proteins that are involved in vesicle trafficking in the SG. Several endosome markers and their interacting partners were tested, including Rab5 (an early endosome marker) (Gorvel et al., 1991), Rab7 (a late endosome marker) (Meresse et al., 1995;

Wichmann et al., 1992), Rab11 (a recycling endosome marker) (Ullrich et al., 1996), Nuclear fallout (Nuf, a putative binding partner for Rab11) (Riggs et al., 2003), and Sec15 (an exocyst complex component and effector for Rab11) (Langevin et al., 2005; Zhang et al., 2004). Using labeling with E-Cad, an adherens junction marker, and CrebA, a SG nuclear marker, we segmented apical cell outlines and calculated intensity mean of fluorescence signals of each endosomal marker. We observed apical enrichment of several of them in the SG, including Rab11 and Nuf (Figure 1, B-B''''). Similar enrichment of Rab11 signals was also observed in the endogenously tagged Rab11 protein (Rab11-EYFP; (Dunst et al., 2015) (Supplemental Figure S1, A-A''''). Importantly, the enrichment of Rab11 was more pronounced in the SG over time (Figure 1, D-F''''). Before invagination, only low levels of Rab11 were detected in the apical domain of all SG cells throughout the entire SG placode (Figure 1, D-D''''). Stronger signals of Rab11 were soon detected in the SG as cells begin to undergo apical constriction in the dorsal/posterior region of the placode (Figure 1, E-E''''), which intensified as invagination proceeded (Figure 1, F-F''''). Quantification of the total intensity of Rab11 in the whole SG placode showed significantly higher levels of Rab11 in invaginating SGs compared to before invagination (Figure 1H), suggesting upregulated Rab11 levels in SG cells during invagination. Furthermore, compared to before invagination (Figure 1G, top), intensity mean of Rab11 in the apical domain of each cell showed a stronger negative correlation with the apical area of cells in the entire SG during invagination (Figure 1G, bottom), suggesting a close link between intracellular trafficking activities and apical constriction.

EYFP-tagged Rab5, an early endosome marker, was also enriched in the apical domain in the invaginating SG, suggesting active endocytosis in the apical domain during SG invagination (Supplemental Figure S1, B-B'''). Rab7, a late endosome marker, however, did not show apical enrichment but localized as large punctate structures in the cytoplasm of the entire SG placode (Supplemental Figure 1, D-D'''). Dynein heavy chain 64C, a subunit of the dynein motor complex that transports cargos along MTs toward their minus ends, also showed apical enrichment (Figure 1, C-C'''). This is consistent with previous report that the minus end of MTs faces the apical domain of SG cells near the invagination pit (Booth et al., 2014). Sec15, an exocyst complex component, also showed similar apical enrichment (Supplemental Figure S1, E-E'''). Similar to Rab11 (Figure 1G),

intensity mean of Rab5-EYFP and Sec15 in the apical domain of each cell showed a

negative correlation with the apical area of cells in the entire SG during invagination

(Supplemental Figure S1, C and F). Compared to a more uniform distribution of other

endosomal markers in the entire apical domain of SG cells, Sec15 signals appeared to

be enriched at adherens junctions (Supplemental Figure S1E'). Overall, our results

suggest active intracellular trafficking, possibly both endo- and exocytosis, in the apical

domain of SG cells near the invagination pit during invagination.

Apical enrichment of Rab11 and Nuf is MT-dependent

Strong apical enrichment of Rab11 and other trafficking components in SG cells near the

invagination pit led us to test whether these vesicle markers are associated with vertically

aligned MTs that could facilitate their polarized distribution. As in many epithelial cells,

MT minus- and plus-ends face the apical and the basal domain of the SG cells,

respectively (Booth et al., 2014; Myat and Andrew, 2002). Indeed, co-immunostaining of

tyrosinated α-tubulin, a marker of dynamic or newly polymerized MTs (Westermann and

Weber, 2003), and Rab11 showed a partial overlap of the two proteins at the apical region

of the SG cells (Figure 2, A-B'').

To test whether Rab11 apical enrichment in SG cells is dependent on MT networks, we

disrupted MTs in the SG by overexpressing spastin, a MT-severing protein (Sherwood et

al., 2004), using the SG-specific *fkh-Gal4* driver. In control SGs, tyrosinated α-tubulin and

acetylated α-tubulin, a marker of stable and longer-lived MTs (Westermann and Weber,

2003), were observed abundantly in the apical domain of cells in the whole placode

(Figure 2, C'' and C'''; Supplemental Figure S2A''). In spastin-overexpressing SGs, both

tyrosinated and acetylated α-tubulin signals were strongly reduced, revealing loss of MT

filaments (Booth et al., 2014) (Figure 2, D'' and D''' and Supplemental Figure S2B''). The

apical enrichment of Rab11 (Figure 2C') was also disrupted and Rab11 was mislocalized

basolaterally (Figure 2D'). SGs with disrupted MTs showed much lower Rab11 and Nuf

intensity in the apical domain compared to control SGs (Figure 2, E-G). To compare the

relative variability of apical enrichment of Rab11/Nuf in WT and spastin-overexpressing

SGs, we calculated coefficient of variation, the ratio of standard deviation of Rab11/Nuf

intensity to the mean Rab11/Nuf intensity. Compared to WT, spastin-overexpressing SGs

showed a lower degree of relative variability of intensity (Figure 2G). Overall, these data

suggest that MT networks are required for apical enrichment of Rab11 and Nuf during SG

invagination.

Rab11 and dynein functions are required for apical constriction in the SG

We next asked if modulation of Rab11 levels could compromise apical constriction. To

test this possibility, we disrupted the function of Rab11 in the SG by either overexpressing

a dominant-negative form of Rab11 (Rab11S25N-YFP; (Zhang et al., 2007); hereafter

referred to as Rab11-DN) or knocking down *Rab11* using RNAi lines. Reduced Rab11

levels upon *Rab11* knockdown were confirmed using a Rab11 antibody in both stage 11

and stage 13 SGs (Supplemental Figure S3, A-D'). Compared to control (Figure 3A), SGs

overexpressing Rab11-DN or knocking down *Rab11* showed more cells with larger apical

areas (Figure 3, B, C and F), suggesting a role for Rab11 in apical constriction during SG

invagination.

As dynein heavy chain was also enriched apically in the SG during invagination (Figure

1, C' and C'''), we tested if dynein function is also required for apical constriction during

SG invagination. Indeed, knockdown of *Dynein heavy chain 64C* (*Dhc64C*) in the SG

using RNAi resulted in more cells with larger apical areas during invagination, suggesting

defective apical constriction upon *Dhc64C* knockdown (Figure 3, D, D' and G). As dynein

binds to and clusters the minus ends of microtubules (Burbank et al., 2006; Elting et al.,

2014; Goshima et al., 2005; Heald et al., 1996; Khodjakov et al., 2003; Tan et al., 2018),

we asked if *Dhc64C* knockdown affects the MT networks in the SG. Knocking down

Dhc64C in the SG resulted in a slight reduction of tyrosinated α-tubulin (Figure 3J'). These

data suggest that apical constriction defects observed in *Dhc64C* knockdown is, at least

in part, due to affected MT networks.

Klarsicht (Klar), the *Drosophila* Klarsicht-Anc-Syne Homology (KASH) domain protein,

mediates apical transport in the SG (Myat and Andrew, 2002) via the MT motor

cytoplasmic dynein (Gross et al., 2000). We therefore tested if *klar* also influences apical

constriction during SG invagination. *klar* null mutant embryos showed SGs with mild

apical constriction defects compared to wild type (Figure 3, E-F' and H). Importantly,

unlike *Dhc64C* knockdown, no significant difference in the MT networks was observed in

SGs in *klar* mutants (Figure 3L'), suggesting that apical constriction defects in *klar*

mutants are due to disrupted apical transport. Overall, our data suggest essential roles of

MT-, Rab11- and dynein-dependent intracellular transport in regulating apical constriction

during SG invagination.

Reduced Rab11 function leads to reduction of apicomedial myosin formation and

failure in accumulation of apicomedial Rok in SG cells

The apicomedial myosin structure generates the pulling force to drive apical constriction

in SG cells (Booth et al., 2014; Chung et al., 2017). Disruption of MTs by spastin

overexpression inhibits formation of apicomedial myosin during SG invagination (Booth

et al., 2014). To test whether apicomedial myosin is affected in SG cells when *Rab11* is

knocked down, we measured the overall intensity of myosin in the apicomedial region of

SG cells using sqh-GFP, a functional GFP-tagged version of the myosin regulatory light

chain (Royou et al., 2004). In control SGs, sqh-GFP showed strong myosin signals with

clear web-like structures in the apicomedial region of cells near the invagination pit

(Figure 4, A-A''''). Knockdown of *Rab11*, however, caused a significant reduction of

apicomedial myosin intensity in SG cells in the same area (Figure 4, B-B'''' and C).

Moreover, unlike clear web-like structures of apicomedial myosin in control SGs (Figure

4, A''-A''''), myosin was more dispersed and fragmented in *Rab11* RNAi SGs (Figure 4,

B''-B''''), resulting in a decrease in areas of myosin accumulation in the apicomedial

region of SG cells (Figure 4E). These data suggest roles for Rab11 in both upregulating

apical myosin and forming and/or maintaining apicomedial myosin web.

We also tested whether junctional myosin, a myosin pool that is closely associated with

adherens junctions in SG cells, is affected in *Rab11* RNAi SGs. Consistent with previous

studies (Chung et al., 2017; Roper, 2012), junctional myosin showed strong signals in

wild type SG cells, often with stronger signals at vertices (Figure 4A'''). Intriguingly, the

intensity of junctional myosin slightly increased when *Rab11* was knocked down (Figure

4, B''' and D). The ratio of apicomedial to junctional myosin intensity was significantly

reduced in *Rab11* RNAi SGs (Figure 4D), suggesting imbalance of contractile forces in

the SG.

We previously showed that accumulation of apicomedial Rok is required for apicomedial

myosin formation for coordinated apical constriction in SG cells (Chung et al., 2017;

Roper, 2012). To test whether Rok accumulation is dependent on Rab11, we quantified

accumulation of apicomedial Rok-GFP signals using a ubiquitously expressed GFP-

tagged Rok transgene (Rok-GFP; (Abreu-Blanco et al., 2014) in wild type and *Rab11*

RNAi SGs (Figure 4, F-G''). Indeed, reduction of *Rab11* levels led to more dispersed Rok

signals along the apical domain of SG cells near the pit (Figure 4H). Overall, our data

suggest Rab11-dependent apical apicomedial myosin formation and accumulation of Rok

during SG invagination.

Reduced dynein function leads to failure in proper organization of apicomedial

myosin in SG cells

We next tested whether apicomedial myosin formation is affected in *Dhc64C* RNAi and

klar mutant SGs. Despite slight disruption of the MT networks in *Dhc64C* RNAi SGs

(Figure 3J'), the overall intensity of apicomedial or junctional myosin was not significantly

changed in *Dhc64C* RNAi SGs (Figure 5C), suggesting that subtle defects in the MT

networks do not affect apical upregulation of myosin. Similar to *Dhc64C* knockdown, no

significant difference in the intensity of apicomedial and junctional myosin was observed

in *klar* mutant SGs (Figure 5H). However, the ratio of apicomedial to junctional myosin

was reduced in both genotypes (Figure 5, D and I). Moreover, the area of apicomedial

myosin accumulation was significantly reduced in *Dhc64C* RNAi or *klar* mutant SGs

(Figure 5, E and J), suggesting defects in forming proper apicomedial myosin web in
these genotypes. Our data suggests that defective apical constriction in *Dhc64C* RNAi
and *klar* mutant SGs could be due to failure in proper organization of myosin structures
in SG cells.

We also tested for accumulation of apicomedial Rok-GFP signals in spastin-
overexpressing, *Dhc64C* RNAi and *klar* mutant SGs. Disruption of MTs by spastin
overexpression abolished apicomedial accumulation of Rok in cells near the invagination
pit (Figure 5, L' and P), found in control SGs (Figure 5K'). Reduction of *Dhc64C* levels
also led to more dispersed Rok signals along the apical domain of SG cells near the pit
(Figure 5, M' and P). In *klar* mutants, Rok-GFP tended to accumulate less profoundly and
less uniformly in the apicomedial region of SG cells (Figure 5O') although not statistically
significant compared to control (Figure 5Q). Overall, our data suggest that MT- and
dynein-dependent apical accumulation of Rok and apicomedial myosin formation during
SG invagination.

286 MT- and Rab11-dependent apical enrichment of Fog, Crb and E-Cad is required for
287 apical constriction during SG invagination

We next tested potential cargos of MT- and Rab11-dependent apical transport during SG
invagination. Rok accumulation, apicomedial myosin formation and subsequent clustered
apical constriction in the SG is dependent on Fog signaling (Chung et al, 2017). We
therefore tested if apical transport of the Fog ligand is MT- and Rab11-dependent.
Staining using an antibody against Fog in control SGs showed upregulated Fog signals
in the apical domain of SG cells (Figure 6, A-A'''). However, in spastin-overexpressing or
Rab11 RNAi SGs, apical Fog signals were reduced (Figure 6, B-C'''). To compare apical
enrichment of Fog between WT and spastin-overexpressing or *Rab11* RNAi SGs, we
calculated the overall intensity of apical Fog signals in the entire SG placode in
corresponding genotypes. Compared to control, Fog levels in the apical domain of the
SG were reduced when spastin was overexpressed or *Rab11* was knocked down (Figure
6G). We also calculated coefficient of variation, the ratio of standard deviation of Fog

intensity to the mean intensity of Fog in the SG. Compared to WT, spastin-overexpressing or *Rab11* RNAi SGs had a lower degree of relative variability of intensity (Figure 6H). These data suggest that apical transport of Fog in the SG is dependent of the MT networks and Rab11.

We further tested MT- and Rab11-dependent apical transport of other apical and junctional proteins during SG invagination. Transport of several key apical and junctional proteins is dependent on MTs and Rab11 (Jouette et al., 2019; Khanal et al., 2016; Le Droguen et al., 2015). One such protein is an apical transmembrane protein Crb, and Rab11 helps maintain apical Crb in the *Drosophila* ectoderm (Roeth et al., 2009). In the *Drosophila* SG and follicle cells, Crb is apically transported along MTs by the dynein motor (Myat and Andrew, 2002). Importantly, in *Drosophila* tracheae, loss of *crb* impairs apical constriction during the internalization process (Letizia et al., 2011). The key junctional protein E-Cad is also trafficked by MTs in *Drosophila* embryonic trachea (Le Droguen et al., 2015). During apical constriction, contractile forces generated by the actomyosin complex are exerted on adherens junctions, with E-Cad being a core component that integrates contractile forces to generate tension (Martin et al., 2009).

We therefore asked if MTs and Rab11 have roles in apical enrichment of Crb and E-Cad in the SG during invagination. Compared to control (Figure 6, D' and D'''), spastin-overexpressing SG cells showed discontinuous Crb signals (Figure 6, E' and E'''). Although the number of gaps was not significantly different between control and spastin-overexpressing SGs (Figure 6H), the ratio of the length of gap over junctional length was significantly increased in SGs that overexpress spastin (Figure 6H). Moreover, whereas E-Cad signals were mainly detected at adherens junctions (Figure 6D''), disruption of MTs resulted in mislocalization of E-Cad to the basolateral region (Figure 6E''). Importantly, *Rab11* knockdown resulted in similar E-Cad mislocalization and discontinuous Crb signals (Figure 6, F-F'''' and H). These data suggest that apical enrichment of Crb and E-Cad in the invaginating SG is dependent on MTs and Rab11.

To test for a role of Crb and E-Cad in regulating apical constriction during SG invagination, we knocked down either gene in the SG using RNAi. Knockdown of *crb* or *E-Cad* resulted

in reduction of junctional Crb or E-Cad levels, respectively, compared to control

(Supplemental Figure S3, E-F''). Reduction of *crb* or *E-Cad* increased the number of cells

that have larger apical areas compared to control (Figure 7, A-D'). Quantification of the

percentage of cells and cumulative percentage of cells of different apical areas showed

a significant decrease in the number of constricting cells in *crb* knockdown SGs (Figure

7D). *E-Cad* knockdown SGs also displayed a similar trend although with less statistical

significance (Figure 7D). These data suggest roles for Crb and E-Cad in regulating apical

constriction during SG invagination.

Apical localization of Fog, Crb and E-Cad is MT- and Rab11-dependent in the SG throughout development

To further investigate the role of MTs in Rab11-dependent apical trafficking during SG

development, we overexpressed spastin in the SG throughout development and analyzed

late stage SGs. Analysis of spastin-overexpressing SGs at stage 16 revealed that

disruption of MTs resulted in short SG tubes with a thin lumen compared to control SGs

of the same stage (Supplemental Figure S5, A and B). Strikingly, whereas the majority of

Rab11- and Nuf-positive vesicles localized in the apical region of control SG cells

(Supplemental Figure S5, A-A'''), Rab11 and Nuf were observed as large aggregates in

the cytoplasm in spastin-overexpressing cells, overlapping with each other (Supplemental

Figure S5, B-B'''). Our data suggest that MTs are required for apical enrichment of

Rab11/Nuf vesicles throughout SG tube formation.

We next tested whether MTs and Rab11 are required for apical transport of Fog, Crb and

E-Cad also at later stages. Compared to the control SG (Supplemental Figure S5C''),

apical Fog signals were significantly reduced in SGs that overexpress spastin or knock

down *Rab11* (Supplemental Figure S5, D'' and E''), suggesting that Fog is trafficked

apically in a MT- and Rab11-dependent manner in the SG throughout tube formation. No

significant changes were detected in apical Crb localization in SGs that overexpress

spastin or knock down *Rab11* (data not shown). Compared to strong E-Cad enrichment

in adherens junctions in control SG cells (Supplemental Figure S5C'), SGs that

overexpress spastin or knock down *Rab11* showed stronger E-Cad signals in the lateral

domain of cells (Supplemental Figure S5, D' and E').

To further test MT-dependent apical transport of Crb or E-Cad in the SG, we used a

genetic suppression strategy in the overexpression background: we co-overexpressed

Crb or E-Cad along with spastin in the SG to test whether disruption of MTs affects apical

transport of Crb or E-Cad. A similar approach was taken in a recent study, where

overexpression of Crb provided a highly sensitive genetic background for identifying

components involved in Crb trafficking (Aguilar-Aragon et al., 2020). Consistent with

previous studies (Chung and Andrew, 2014; Wodarz et al., 1993), Crb overexpression in

the SG caused a dramatic increase of the apical membrane as well as mislocalization of

Crb basolaterally along the entire membrane (Supplemental Figure S4, F-F'). Co-

overexpression of spastin along with Crb suppressed the overexpression phenotypes for

Crb (Supplemental Figure S4, G-G'). E-Cad overexpression in the SG caused

mislocalization of E-Cad as punctate structures in the basolateral region (Supplemental

Figure S4, H'-H''). Importantly, in SGs that co-overexpressed spastin along with Crb or

E-Cad, each protein mislocalized as cytoplasmic aggregates near the basolateral domain,

which were largely overlapping with mislocalized Rab11/Nuf (Supplemental Figure S4,

G'-G''' and I'-I'''). Overall, our data suggest that Fog, Crb and E-Cad are trafficked apically

in a MT- and Rab11-dependent manner in the SG throughout tube formation.

Reducing apical and junctional proteins affects apicomedial myosin formation

To test whether apical constriction defects observed in *crb* and *E-Cad* RNAi SGs are

linked to apicomedial myosin formation, we quantified myosin levels and areas of

apicomedial myosin in SGs with these knockdowns. Compared to control SGs (Figure 8,

A-A''), *crb* RNAi SGs showed a slight decrease in apicomedial myosin levels (Figure 8,

B-B'' and D). No significant difference in the junctional myosin intensity was observed in

crb RNAi SGs (Figure 8, B-B'' and D). The ratio of apicomedial to junctional myosin was

reduced (Figure 8E), suggesting that apical constriction defects in SGs depleted of *crb*

are, at least in part, due to failure of apicomedial myosin formation. Supporting this idea,

areas of apicomedial myosin web-like structures were significantly reduced in *crb* RNAi

SGs compared to WT (Figure 8F).

In SGs knocked down *E-Cad*, myosin levels at each domain did not change significantly

(Figure 8, C-C'' and E), but areas of apicomedial myosin web-like structures were

significantly decreased (Figure 8F). The ratio of apicomedial to junctional myosin was

also significantly reduced in *E-Cad* RNAi SGs compared to control (Figure 8, C-C'' and

E). Therefore, apical constriction defects caused by *E-Cad* knockdown might be due to

defective apicomedial myosin formation and the imbalance of contractile forces.

Consistent with reduced apicomedial myosin (Figure 8, B' and D), knockdown of *crb* also

resulted in Rok-GFP signals less accumulating in the apical region of cells near the

invagination pit (Figure 8, F-G'). Quantification of areas occupied by Rok-GFP puncta

showed significant reduction in accumulation of apicomedial Rok when *crb* was knocked

down (Figure 8I). Consistent with little effect of *E-Cad* knockdown on apicomedial myosin

intensity (Figure 8, C-C'' and E), Rok accumulation was not significantly affected by *E-*

Cad knockdown (Figure 8, H-H' and I). Taken together, our results suggest a role for Crb

and E-Cad in regulating apical constriction by apicomedial myosin activation during SG

invagination.

Discussion

MTs and Rab11 regulate Fog signaling activity during SG invagination

MTs have a crucial role in stabilizing apical myosin during epithelial morphogenesis both

in early *Drosophila* embryos and in the *Drosophila* SG (Booth et al., 2014; Ko et al., 2019).

In the SG, MTs interact with apicomedial myosin via Short stop, the *Drosophila*

spectraplakin, emphasizing a direct interplay between the MT and the apical myosin

networks (Booth et al., 2014). Our data reveals another key role of MTs in regulating

protein trafficking to control the apical myosin networks during tissue invagination. During

SG invagination, a network of longitudinal MT bundles is observed near the invagination

pit (Booth et al., 2014). Our data shows apical enrichment of Rab11 in the same area is

MT-dependent (Figures 1 and 2) and that this enrichment is important for forming the apicomedial myosin networks (Figure 4), suggesting a link between localized intracellular trafficking along MTs to apical myosin regulation.

The dorsal/posterior region of the SG, where Rab11 is apically enriched in a MT-dependent manner (Figures 1 and 2), correlates with localized Fog signaling activity that promotes clustered apical constriction (Chung et al., 2017). Disruption of MTs or *Rab11* knockdown reduces Fog signals in the apical domain of SG cells (Figure 6) and causes dispersed Rok accumulation and defective apicomedial myosin formation (Figure 4). It is consistent with our previous study that absence of Fog signal results in dispersed apical Rok and defects in apicomedial myosin formation (Chung et al., 2017). We therefore propose that MT- and Rab11-dependent apical trafficking regulates Fog signaling activity to control apical constriction during epithelial tube formation, through transporting the Fog ligand. As recycling of membrane receptors to the cell surface plays an important role in the regulation of overall signaling activity, it is possible that Rab11 is involved in recycling of the as yet unidentified SG receptor(s) of Fog to regulate Fog activity in the SG. Indeed, several GPCRs are recycled via Rab11 (Anborgh et al., 2000; Cerniello et al., 2017; Dale et al., 2004; Fan et al., 2003; Hamelin et al., 2005; Hunyady et al., 2002; Innamorati et al., 2001; Volpicelli et al., 2002). During epithelial invagination in early *Drosophila* embryogenesis, the concentration of the Fog ligand and receptor endocytosis by β-arrestin-2 have been shown as coupled processes to set the amplitude of apical Rho1 and myosin activation (Jha et al., 2018). It is possible that the movement of Fog receptor(s) that have internalized as a stable complex with β-arrestin is recycled back to the cell surface by Rab11. The Fog signaling pathway represents one of the best-understood signaling cascades controlling epithelial morphogenesis (Manning and Rogers, 2014). Although best studied in *Drosophila*, the pathway components have been also identified in other insects, suggesting a more widely conserved role of Fog signaling in development (Benton et al., 2019; Urbansky et al., 2016). Further work needs to be done to fully understand the regulatory mechanisms underlying trafficking of Fog and its receptor(s) during epithelial morphogenesis.

Our analysis of apicomedial myosin shows that reduced Rab11 function not only causes

a decrease of the myosin intensity but also causes myosin to be dispersed rather than

forming proper myosin web structures in the apicomedial domain of SG cells (Figure 4).

These data support the idea that Rab11 function is required for both concentration and

spatial organization of apicomedial myosin. This can be explained by the combined effect

of multiple cargos that are transported by Rab11, including Fog, Crb and E-Cad (as

discussed below). Time-lapse imaging of myosin will help determine how dynamic

behavior of apicomedial myosin is compromised when Rab11 function is disrupted.

453 Integrating apical and junctional proteins with actomyosin networks during SG

454 invagination

During branching morphogenesis in *Drosophila* trachea, MTs and dynein motors have a

critical role in proper localization of junctional proteins such as E-Cad (Le Droguen et al.,

2015). This is consistent with our observations with MT-dependent apical localization of

E-Cad in the invaginating SG (Figure 6), suggesting a conserved role of MT-dependent

intracellular trafficking in junctional remodeling and stabilization during epithelial tube

formation. Our data further suggests that the MT networks and Rab11 have key roles in

apical localization of Crb and E-Cad in the SG (Figure 6) and that proper levels of apical

and junctional proteins are important for apical constriction during SG invagination (Figure

7). Based on these data, we propose that MT- and Rab11-dependent apical trafficking of

Crb and E-Cad is critical for apical constriction during SG invagination. Alternatively, MTs

have additional role in assembling/anchoring these apical components, through

regulation of unidentified molecules. Recent studies in *Drosophila* mesoderm invagination

showed that MTs help to establish actomyosin network linked to cell junction to facilitate

efficient force transmission to promote apical constriction (Ko et al., 2019). In Ko et al.,

(2019), however, MT-interfering drugs and RNAi of CAMSAP end-binding protein were

used to prevent MT functions and the effect cannot be directly compared to our data

where spastin was used to sever existing MTs. Direct monitoring MT-dependent transport

of Crb and E-Cad during SG invagination will help clarify the mechanism.

Upon knockdown of *crb* or *E-Cad,* less prominent apicomedial myosin web structures are
observed in invaginating SGs (Figure 8), suggesting a requirement of Crb and E-Cad in
proper organization of apical actomyosin networks during SG tube formation. Crb acts as
a negative regulator of actomyosin dynamics during *Drosophila* dorsal closure (Flores-
Benitez and Knust, 2015) and during SG invagination (Roper, 2012). It is possible that
proper Crb levels are required for modulating myosin activity both in the apicomedial
domain and at junctions during SG invagination, which contribute to apical constriction
and cell rearrangement, respectively (Roper, 2012; Sanchez-Corrales et al., 2018).
Anisotropic localization of Crb and myosin was observed at the boundary of the SG
placode, where myosin accumulates at edges where Crb is lowest (Roper, 2012). Planar
polarization of Rok at this boundary is modulated through phosphorylation by Pak1
downstream of Crb (Sidor et al., 2020). Further test will help understanding whether and
how Crb might affect junctional myosin dynamics and SG invagination. As contractile
actomyosin structures exert forces on adherens junction to drive apical constriction, we
speculate that apical constriction defects upon *E-Cad* RNAi might be due to reduction of
cell adhesion and/or of improper force transmission. It will be interesting to determine if
coordination of apical and junctional proteins and apical cytoskeletal networks through
intracellular trafficking is conserved during tubular organ formation in general.

A role for dynein in apical constriction

Dhc is also apically enriched in the dorsal/posterior region of the invaginating SG (Figure
1). Our data shows that knockdown of *Dhc64C* not only affect Rok accumulation and
apicomedial myosin formation (Figure 5) but also disrupts MT organization in the SG
(Figure 3). This data is consistent with previous findings that cytoplasmic dynein is
associated with cellular structures and exerts tension on MTs. For example, dynein
tethered at the cell cortex can apply a pulling force on the MT network by walking towards
the minus end of a MT (Laan et al., 2012). Dynein also scaffolds the apical cell cortex to
MTs to generate the forces that shape the tissue into a dome-like structure (Takeda et

al., 2018). In interphase cells, force generated by dynein also regulates MT turnover and
organization (Yvon et al., 2001).

In *klar* mutants, on the other hand, MT organization is not affected in the SG (Figure 3),
suggesting that reduction of dynein-dependent trafficking by loss of *klar* does not cause
changes in the MT networks. Notably, although the intensity of apicomedial myosin does
not change upon *Dhc64C* knockdown or in the *klar* mutant background, formation of
apicomedial myosin web structures is affected (Figure 5). These data suggest a possible
scenario that dynein function is not required for concentration of myosin in the apical
domain but is only needed for spatial organization of apicomedial myosin. We cannot rule
out the possibility, however, that zygotic knockdown of *Dhc64C* by RNAi is not strong
enough to affect the intensity of apicomedial myosin. *Dhc64C* has strong maternal
expression and is essential for oogenesis and early embryo development (Li et al., 1994).
Embryos with reduced maternal and zygotic pools of *Dhc64C* showed a range of
morphological defects in the entire embryo, some of which were severely distorted (data
not shown). Precise roles for dynein and dynein-dependent trafficking in regulating
apicomedial myosin formation remain to be elucidated.

Materials and Methods

Fly stocks and husbandry

Fly lines used in our experiments were listed in a separate table. All crosses were
performed at 25°C, unless stated otherwise.

Antibody staining and confocal microscopy

Antibodies used in our experiments were listed in a separate table. Embryos were
collected on grape juice-agar plates and processed for immunofluorescence using
standard procedures. Briefly, embryos were dechorionated in 50% bleach, fixed in 1:1
heptane:formaldehyde for 40 min and devitellinized with 80% EtOH, then stained with
primary and secondary antibodies in PBSTB (1X PBS, 0.1% Triton X-100, 0.2% BSA).

For acetylated α-tubulin, tyrosinated α-tubulin, sqh-GFP, Rok-GFP, Fog and phalloidin

staining, embryos were hand-devitellinized. All images were taken with a Leica SP8

confocal microscope.

Cell segmentation and apical area quantification

Embryos immunostained with E-Cad and CrebA were imaged using a Leica SP8 confocal

microscope. As Rok accumulation, apicomedial myosin and apical constriction depend

on the depth of invagination in the SG, SGs that were invaginated within the range of 5.1-

9.9 μm depth were used for quantification for proper comparison between different

genotypes. Maximum intensity projection was generated from three apical focal planes

with highest E-Cad signals for all genotypes (0.3 μm apart for each focal plane). Cells

were segmented along E-Cad signals and cell areas were calculated using the Imaris

Program (Bitplane). Since the Imaris Program calculated the areas of both the front and

the back of the projected cell layer, we divided the measured areas by two to get the true

values of apical areas of SG cells.

Negative correlation between apical area and Rab11/Nuf intensity

Cell segmentation for five WT SGs within the range of 5.1- 9.9 μm invagination was

performed as described above. All experiments were carried out in the same condition

and same settings for confocal imaging were used. Three confocal sections in the apical

region that show strongest Rab11/Nuf signals were used to produce the maximum

intensity projection. Intensity mean was measured for Rab11/Nuf signals for each

segmented cell using the Imaris Program (Bitplane) and plotted. Correlation (Pearson)

and P values were calculated using the GraphPad Prism software.

Total intensity of Rab11, Nuf and Fog signals

For measuring the total intensity of Rab11/Nuf signals in the apical region of cells in the
whole SG placode, the integrated density of Rab11/Nuf of each SG cell was calculated
(integrated density= intensity mean x area). Intensity means and SG cell areas were
measured using the Imaris software. The total intensity was calculated as the sum of
integrated densities of all cells in the placode. Five SGs were used for quantification. P
values were calculated using Welch's t-test in the GraphPad Prism software.

For the total intensity of Fog signals, background corrections were carried out first. Using
the Fiji software, mean gray values of Fog signals of 10 epidermal cells outside of the SG
placode were measured. The background intensity was calculated as the average value
of the intensity means of those 10 cells. The intensity mean of Fog signals of all SG cells
were measured using the Imaris program, and the background intensity was subtracted.
After background correction, the intensity mean was multiplied by SG cell areas to
calculate the total Fog intensity. Five SGs were used for quantification. P values were
calculated using Welch's t-test in the GraphPad Prism software.

Coefficient of variation for Rab11, Nuf and Fog signals

Coefficient of variation was calculated as the ratio of deviation of Rab11/Nuf intensity to
the mean Rab11/Nuf intensity. The mean value of Rab11/Nuf intensity was calculated as
the average mean intensity of Rab11/Nuf signals in all SG cells. Deviation of Rab11/Nuf
intensity is the difference between the mean intensity of Rab11/Nuf in each cell and the
mean value. Five WT SGs (690 cells) and five spastin-overexpressed SGs (496 cells)
were analyzed and plotted. P values were calculated using the Mann-Whitney U test in
the GraphPad Prism software.

Coefficient of variation for Fog signals of control, spastin-overexpressing and Rab11 RNAi
SGs were calculated using the same methods as for Rab11/Nuf. Five SGs for each
genotype were analyzed (control (*fkh-Gal4/+*), 475 cells; *fkh-Gal4/UAS-Spastin*, 507
cells; *fkh-Gal4/UAS-Rab11* RNAi, 487 cells).

Quantification of intensities of myosin, Crb and E-Cad

For myosin quantification, maximum intensity projections that span the apical and the junctional region of SG cells were used (Leica LasX) and measurements were performed using the Fiji software. Five SGs were used for quantification for each genotype. A group of 20 cells in the dorsal/posterior region of the SG placode near the invagination pit was selected for quantification of myosin intensity. Regions were drawn manually along the inner or outer boundary of E-Cad signals of each cell to calculate the mean gray value of apicomedial and junctional myosin.

For background correction, mean gray values of apicomedial myosin in ten cells outside of the SG placode was measured. The average value of mean gray values of apicomedial myosin in these ten cells was used to subtract the background of the cells inside the placode from the same embryo. The mean intensity of apicomedial/junctional myosin was normalized by the median deviation. SuperPlots (Lord et al., 2020) were used to display quantification. For myosin intensity, each dot in the graph represents the average value of 20 cells in each SG. For Crb and E-Cad intensities, each dot in the graph represents the average value of 10 cells for each SG. Five SGs were used for quantification of all three proteins. P values were calculated using Welch's t-test in the GraphPad Prism software.

Quantification of area of Rok-GFP and apicomedial myosin puncta

For quantification of area of Rok-GFP puncta, a single confocal section that had the strongest medial Rok signals was selected. Cell boundaries were labeled by immunostaining with the antibody against E-Cad. To analyze Rok distribution, we performed Particle Analysis using the Fiji software. 15 cells in the dorsal/posterior region near the invagination pit were selected for quantification. Rok-GFP signals were converted into black-and-white using the *Threshold* tool in Photoshop before analysis. Using the *Analyze particles* tool in Fiji, Rok-GFP puncta with areas equal or larger than

0.02 µm^2 were measured. Five SGs were used for quantification. P values were

calculated using Welch's t-test in the GraphPad Prism software.

For measuring areas of apicomedial myosin puncta, a single confocal section with

strongest apicomedial myosin signals was selected. A group of 10 cells in the

dorsal/posterior region of SG was analyzed. Junctional myosin was excluded manually

and only apicomedial myosin signals were used for quantification. Measuring the area of

apicomedial myosin were carried out with the same method used for quantifying the area

of Rok-GFP puncta. Five SGs were used for quantification. P values were calculated

using Welch's t-test in the GraphPad Prism software.

Quantification of length and number of gaps of junctional Crb

Length of gaps and junctional length were measured using the Fiji software. Gaps that

have the length equal or more than 0.2 µm were quantified. If there were more than one

gap per junction, the length of gaps was calculated as a sum of all gaps in a given junction.

For each SG, 10 cells in the dorsal/posterior region were used for quantification. For the

number of gaps, the total number of gaps in those 10 cells were counted. Five SGs (~110

junctions) were used for quantification. P values were calculated using Welch's t-test in

the GraphPad Prism software.

Fly strains	Source	RRID	References
Rab11-EYFP	Bloomington Stock Center	62549	(Dunst et al., 2015)
Rab5-EYFP	Bloomington Stock Center	62543	(Dunst et al., 2015)
Rab7-EYFP	Bloomington Stock Center	62545	(Dunst et al., 2015)
UAS-Spastin (on II)	N. Sherwood, Duke		(Sherwood et al., 2004)
UAS-Spastin-CFP (on III)	N. Sherwood, Duke		(Du et al., 2010)
fkh-Gal4 (on II)	D. Andrew (John Hopkins University)		(Chung et al., 2017)
fkh-Gal4 (on III)	D. Andrew (John Hopkins University)		(Chung et al., 2017)
ubi-Rok-GFP	S. Parkhurst (Fred Hutchinson Cancer Research Center)		(Abreu-Blanco et al., 2014)
sqh-GFP			(Royou et al., 2004)

UAS-Rab11S25N-EYFP	Bloomington Stock Center	9792, 23261	
UAS-Dicer-2	Bloomington Stock Center	60008	
UAS-Dhc64C RNAi	Bloomington Stock Center Vienna *Drosophila* Resource Center	76941, 28749, v28053, v28054. 4 lines worked equally. v28054 was used for all of the data shown in this work.	
klar^1	Bloomington Stock Center	3256	
klar^{mCD4}	Bloomington Stock Center	25097	
klar^{mCD4} ubi-Rok-GFP	Recombinant line generated from *klar^{mCD4}* and *ubi-Rok-GFP*		
UAS-Crb RNAi	Bloomington Stock Center	38373, 38903, 40869, 34999.	

		4 lines worked equally. 2 lines (38373, 34999) were used for the data shown in this work.	
UAS-E-Cad RNAi	Bloomington Stock Center	32904	
UAS-Shg-GFP	Bloomington Stock Center	58445	
UAS-Rab11 RNAi	Bloomington Stock Center	42709, 27730. Line 42709 was used for testing myosin and Rok-GFP signals. Line 27730 was used for the rest of the experiments.	

Antibody	Source	RRID	Dilution
α-E-Cad (rat)	DSHB (deposited by T. Uemura, Kyoto University)	DCAD2 (AB_528120)	1:50
α-CrebA (rat)	Andrew lab (John Hopkins University)		1:3000
α-CrebA (rabbit)	Andrew lab (John Hopkins University)		1:5000
α-Dhc (mouse)	DSHB (deposited by J.M. Scholey, University of California)	2C11-2 (AB_2091523)	1:50
α-Rab11 (rabbit)	Andrew lab (John Hopkins University)		1:500
α-Nuf (guinea pig)	Sotillos lab (CABD)		1:500
α-acetylated α-tubulin (mouse)	Invitrogen	32-2700 (AB_2533073)	1:1000
α-tyrosinated α-tubulin (rat)	Invitrogen	MA1-80017 (AB_2210201)	1:1000
α-GFP (chicken)	Invitrogen	A10262 (AB_2534023)	1:500

α-β-galactosidase (rabbit)	Invitrogen	A11132 (AB_221539)	1:500
α-Crb (mouse)	DSHB (deposited by E. Knust, Max Planck Institute)	Cq4 (AB_528181)	1:10
α-Fog (rabbit)	N. Fuse (Fuse et al., 2013)		1:1000
α-Sec15 (guinea pig)	Bellen lab (Baylor College of Medicine)		1:2000
Alexa Fluor 488/568/647-coupled secondary antibodies	Invitrogen		1:500

Acknowledgments

We thank the members of the Chung laboratory for comments and suggestions. We thank

A. Martin, S. Parkhurst and N. Sherwood and the Bloomington stock center for fly stocks,

and D. Andrew, H. Bellen, S. Sotillos and the Developmental Studies Hybridoma Bank

for antibodies. We thank Flybase for the gene information. We are grateful to A. Bohnert,

C. Hanlon, A. Johnson and C. O'Kane for their helpful comments on the manuscript. This

work is supported by start-up fund from Louisiana State University and the grant from the

Board of Regents Research Competitiveness Subprogram GR-00005224 to S.C.

Figure legends

Figure 1. Intracellular trafficking components are apically enriched in invaginating SGs. (A) A schematic drawing of the anterior region of the *Drosophila* embryo for stage 11 (ventral and lateral views) and stage 16 (lateral view) with SGs shown in green. Top right, magnified view of a stage 11 SG. The region where SG cells undergo clustered apical constriction during invagination is shown in purple. (B-B'') En face (top) and lateral (bottom) views of a wild type stage 11 SG immunostained for E-Cad, Rab11 and Nuf. Rab11 and Nuf show apical enrichment near the invagination pit (yellow arrowheads). Red arrowheads, Rab11 and Nuf signals near the segmental groove. (B'''-B''''') Heat maps of apical area (B''') and intensity of Rab11 (B'''') and Nuf (B'''''). Cells with smaller apical areas near the invagination pit (dark blue cells in B''') show high intensity of Rab11 and Nuf (red signals in B'''' and B'''''). (C-C''') En face (top) and lateral (bottom) views of a wild type stage 11 SG immunostained for E-Cad (C) and Dhc64C (C'). Corresponding heat maps for apical areas (C'') and intensity of Dhc64C signals (C''') are shown. (D-F''') Wild type stage 11 SGs immunostained for E-Cad and Rab11 at different timepoints of invagination. (G) Total intensity of Rab11 in the whole SG placode before and during invagination. Before invagination, n= 4 SGs; during invagination, n= 5 SGs. (H) Negative correlation between Rab11 intensities and apical areas of SG cells before (top) and during invagination (bottom). R, Pearson correlation coefficient. p< 0.0001 for all samples. Before invagination, n= 4 SGs, 506 cells; during invagination, n= 4 SGs, 561 cells. Asterisks: invagination pit. White lines: SG boundary. SG boundaries are marked based on CrebA (SG nuclear marker) signals in the basal region of SG cells (not shown).

Figure 1

Figure 2. Disruption of MTs results in mislocalization of Rab11 and Nuf to the basolateral domain in SG cells. (A-A''') A wild type SG immunostained for Rab11 (green) and tyrosinated α-tubulin (Tyr-α-Tub; purple). Higher magnification of the yellow boxed area is shown in A'-A'''. Yellow arrowheads, co-localized Rab11 and tyrosinated α-tubulin. (B-B'') Z sections along the yellow line in A. (C-C''') En face (top) and lateral (bottom) views of a control (*fkh-Gal4/+*) SG show abundant levels of tyrosinated α-tubulin (C''; Tyr-α-Tub) and acetylated α-tubulin (C'''; Ace-α-Tub). (D-D''') Overexpression of spastin in the SG using *fkh-Gal4* leads to reduction of tyrosinated α-tubulin (D'') and loss of acetylated α-tubulin (D'''). Whereas Rab11 is enriched in the apical domain of WT SG cells (yellow arrowheads in C'), Rab11 mislocalizes basolaterally when spastin is overexpressed (red arrowheads in D'). (E-F'') Confocal images of WT (E and E') and spastin-overexpressing SGs (F, F') stained for Rab11 (E, F) and Nuf (E', F'). Maximum projection of three confocal sections for the apical domain is shown. Compared to strong Rab11 and Nuf signals in the apical domain in WT SGs (yellow arrowheads in E and E'), Rab11 and Nuf are reduced in spastin-overexpressing SGs (red arrowheads in F and F'). (G) Total intensity of Rab11 and Nuf in the whole SG placode in control and spastin-overexpressing SG. n= 5 SGs for each genotype. (H) Coefficient of variation of Rab11 (left) and Nuf (right) intensities in the SG (n= 5 SGs for both genotypes; WT, 690 cells; *fkh-Gal4/UAS-Spastin*, 496 cells). ****, p<0.0001, Mann-Whitney U test. Asterisks: invagination pit. White lines: SG boundary.

Figure 2

Figure 3. Rab11 and dynein functions are required for coordinated apical
constriction in the SG. (A-F) Confocal images of control (A), Rab11-DN-overexpressing
(B), *Rab11* RNAi (C), *Dhc64C* RNAi (D), wild type (E) and *klar* mutant (F) SGs
immunostained for E-Cad. (A'-F') Heat maps corresponding to images shown in A-F. (G,
H) Percentage (G) and cumulative percentage (H) of SG cells with different apical areas.
Mann-Whitney U test (for percentage of cells) and Kolmogorov-Smirnov test (for
cumulative percentage of cells). n= 5 SGs (control (*fkh-Gal4/+*), 517 cells; *fkh>Rab11-*
DN, 536 cells; *fkh>Rab11* RNAi, 474 cells; *fkh>Dhc64C* RNAi, 499 cells) and 6 SGs (WT,
690 cells; *klar[1]*, 705 cells). *, p<0.05; ****, p<0.0001. (I-J'') Control (I-I'') and *Dhc64C* RNAi
(J-J'') SGs stained for tyrosinated-α-Tub (I', J') and phalloidin (I'', J''). Compared to control
(I'), tyrosinated-α-Tub (Tyr-α-Tub) is decreased in *Dhc64C* RNAi SGs (J'). (I-J'') WT (K-
K'') and *klar* mutant (L-L'') SGs stained for acetylated-α-Tub (Ace-α-Tub; K', K') and
phalloidin (L'', L''). No significant changes in acetylated-α-Tub levels are detected in *klar*
mutant SGs. Asterisks: invagination pit. White lines: SG boundary.

Figure 3

Figure 4. Compromised Rab11 functions lead to a reduction of apicomedial myosin formation and impairs apicomedial Rok accumulation in SG cells. (A-B''') sqh-GFP signals in control (A-A''') and *Rab11* RNAi (B-B''') SGs immunostained for E-Cad and GFP. Higher magnification of the yellow boxed area in A and B are shown in A''-A''' and B''-B'''. Compared to strong signals of apicomedial myosin web structures in the control SG (yellow arrowheads in A''), reduced and dispersed myosin signals are detected upon *Rab11* knockdown (red arrowheads in B''). (A'''' and B'''') Inverted sqh-GFP signals are used for measuring areas of apicomedial myosin structures. (C-E) Quantification of the intensity of apicomedial (AM) and junctional (J) myosin (C), the ratio of apicomedial to junctional myosin (D) and areas of apicomedial myosin (E) in control and *Rab11* RNAi SG cells. n= 5 SGs for both genotypes; 10 cells in the dorsal/posterior region of each SG. *, p< 0.05; **, p< 0.01. Welch's t-test. (F-G'') Rok-GFP signals in control (F, F') and Rab11 RNAi (G, G') SGs immunostained for E-Cad and GFP. Higher magnification of the yellow boxed area in F and G are shown in F' and G'. Compared to strong accumulation of Rok-GFP signals in the apicomedial region of control SG cells (yellow arrowheads in F'), Rok-GFP signals are more dispersed when *Rab11* is knocked down (red arrowheads in G'). (F'' and G'') Inverted Rok-GFP signals are used for measuring areas of apicomedial Rok accumulation. (H) Quantification of areas of Rok-GFP puncta. *, p< 0.05; **, p<0.01. Welch's t-test. n= 5 SGs for all genotypes; 15 cells in the dorsal/posterior region for each SG. Asterisks: invagination pit.

Figure 4

| Sqh-GFP E-Cad | Sqh-GFP | Sqh-GFP E-Cad | Sqh-GFP | |

A **A'** **A''** **A'''** **A''''**

5µm   1µm

+

B **B'** **B''** **B'''** **B''''**

sqh-GFP/+; fkhGal4/+

UAS-Dicer-2/+;
UAS-Rab11 RNAi

C
myosin intensity (RFU/µm²)

60
40
20
0

AM   J   AM   J
cont.   *Rab11* RNAi

D
ratio of myosin intensity (apicomedial/junctional)

1.5
1.0
0.5
0.0

cont.   *Rab11* RNAi

E
areas of apicomedial myostin (µm²)

0.4
0.3
0.2
0.1
0.0

cont.   *Rab11* RNAi

| Rok-GFP E-Cad | | |

F **F'** **F''**

5µm   1µm

+

fkh-Gal4 ubi-Rok-GFP/+

G **G'** **G''**

UAS-Dicer-2/+;
UAS-Rab11 RNAi/+

H
area of Rok-GFP puntae (µm²)

0.15
0.10
0.05
0.00

cont.   *Rab11* RNAi

726
727

Figure 5. Compromised dynein function leads to a reduction of apicomedial myosin formation and impairs apicomedial Rok accumulation in SG cells. (A-B', F-G') sqh-GFP signals in control (A, A', F, F'), *Dhc64C* RNAi (B, B') and *klar* mutant (G, G') SGs immunostained for E-Cad and GFP. Higher magnification of the yellow boxed area in A, B, F and G are shown in A', B', F' and G'. Yellow and red arrowheads indicate apicomedial myosin. (C-E, H-J) Quantification of the intensity of apicomedial and junctional myosin (C, H), the ratio of apicomedial to junctional myosin (D, I) and areas of apicomedial myosin (E, J) in SG cells. n= 5 SGs for all genotypes; 10 cells in the dorsal/posterior region of each SG. *, p< 0.05; **, p< 0.01, Welch's t-test. (A-A'') (K-O') Rok-GFP signals in control (K, K', N, N'), spastin-overexpressing (L, L'), *Dhc64C* RNAi (M, M') and *klar* mutant (O, O') SGs immunostained for E-Cad and GFP. Higher magnification of the yellow boxed area in K-O are shown in K'-O'. (P, Q) Quantification of areas of Rok-GFP puncta. *, p< 0.05; **, P<0.01, Welch's t-test. n= 5 SGs for all genotypes; 15 cells in the dorsal/posterior region for each SG. Asterisks: invagination pit.

Figure 5

Figure 6. MTs and Rab11 are required for apical transport of Fog, Crb and E-Cad during SG invagination. (A-C') Confocal images for control (A-A'), spastin-overexpressing (B-B') and *Rab11* RNAi (C-C') SGs immunostained for E-Cad and Fog. (A''-C''') Corresponding heat maps for apical area (A''-C'') and Fog intensity (A'''-C'''). (D-F'''') Confocal images for control (A-A'), spastin-overexpressing (B-B') and *Rab11* RNAi (C-C') SGs immunostained for E-Cad and Crb. Higher magnification of boxed areas in D-F are shown in D'''-F''''. Compared to junctional enrichment of E-Cad signals in the control SG (D''), E-Cad mislocalizes to the basolateral region in spastin-overexpressing (E'') and *Rab11* RNAi (F'') SGs (red arrowheads). Compared to relatively continuous Crb signals in the control SG (D'''), Crb signals show gaps (red arrows) in spastin-overexpressing (E''') and *Rab11* RNAi (F''') SGs. (G) Total intensity of apical Fog signals in the entire SG placode in control, spastin-overexpressing and *Rab11* knocked down SG. n= 5 SGs for each genotype. (H) Coefficient of variation for Fog signals. n= 5 SGs (control (*fkh-Gal4/+*), 475 cells; *fkh-Gal4/UAS-spastin*, 507 cells; *fkh-Gal4/UAS-Rab11* RNAi, 487 cells). Kolmogorov-Smirnov test. (I) Quantification of the ratio of length of gaps to junctional length (Welch's t-test) in SGs immunostained for Crb. (J) Quantification of the number of gaps (Welch's t-test) in SGs immunostained for Crb. n= 5 SGs; 10 cells for each SG. Asterisks: invagination pit.

Figure 6

Figure 7. Crb and E-Cad have roles in regulating apical constriction during SG

invagination. (A-C) Confocal images of control (A), *crb* RNAi (B) and *E-Cad* RNAi (C)

SGs immunostained for E-Cad. (A'-C') Corresponding heat maps for apical areas in SGs

shown in A-C. (D) Percentage and cumulative percentage of cells with different apical

areas. Mann-Whitney U test (percentage of cells) and Kolmogorov-Smirnov test

(cumulative percentage of cells). n= 5 SGs (control, 517 cells; *crb* RNAi, 583 cells; *E-Cad*

RNAi, 712 cells). *, p<0.05; ****, p<0.0001. Asterisks: invagination pit. White lines: SG

boundary.

Figure 8. Knockdown of *crb* or *E-Cad* results in reduced apicomedial myosin formation and dispersed apicomedial Rok. (A-C") sqh-GFP signals in control (A-A"), *crb* RNAi (B-B") and *E-Cad* RNAi (C-C") SGs immunostained for E-Cad and GFP. (A"-C") Higher magnification of yellow boxed areas in A-C. (D-F") Rok-GFP signals in control (D-D"), *crb* RNAi (E-E") and *E-Cad* RNAi (F-F") SGs immunostained for E-Cad and GFP. (D"-F") Higher magnification of yellow boxed areas in D-F. (G-I) Quantification of the intensity of apicomedial and junctional myosin (G), the ratio of apicomedial to junctional myosin (H) and areas of apicomedial myosin structures (I) in SG cells from different genotypes shown in A-C (n=5 SGs for each genotype; 10 cells in the dorsal/ posterior region). *, p<0.05; **, p<0.01. Welch's t-test. (F-I") (J) Quantification of areas of Rok-GFP puncta in SG cells from different genotypes shown in D-F (n=5 SGs; 15 cells in the dorsal/posterior for each genotype). *, p<0.05; **, p< 0.01. Welch's t-test. Asterisks: invagination pit.

Figure 8

Figure 9. A proposed model for MT-dependent trafficking to promote apical
constriction during SG invagination. Vesicular transport is essential for apical
localization of Fog, Crb and E-Cad to regulate apical myosin networks and subsequent
apical constriction. Precise roles for the dynein motor in regulating apical constriction
remain to be determined. Red question mark represents a potential role of dynein in
trafficking cargos to the apical region of SG cells during apical constriction.

Figure 9

797
798

Figure S1. Rab11-EYFP, Rab5-EYFP and Sec15 are apically enriched in the invaginating SG. (A-A") *Rab11-EYFP* (an EYFP insertion at the N-terminus under the control of Rab11 regulatory sequences) SG labeled with antibodies against E-Cad (A), GFP (A') and Nuf (A"). (A'''-A'''') Corresponding heat maps for apical areas of SG cells (A''') and intensities of GFP (A'''') and Nuf (A''''') signals. (B, B') Confocal images of a *Rab5-EYFP* (an EYFP insertion at the N-terminus under the control of Rab5 regulatory sequences) SG stained for E-Cad (B) and GFP (B'). (B", B''') Corresponding heat maps for apical areas of SG cells (B") and intensity of Rab5-EYFP signals (B'''). (C) Negative correlation between Rab5-EYFP intensity and apical areas of SG cells (n= 4 SGs; 396 cells). (D-D') *Rab7-EYFP* (an EYFP insertion at the N-terminus under the control of Rab7 regulatory sequences) SG immunostained for E-Cad (D) and GFP (D'). (D", D''') Corresponding heat maps for apical areas of SG cells (D") and intensity of Rab7-EYFP (D'''). (E-E''') Wild type SG immunostained for E-Cad (E) and Sec15 (E'). (E", E''') Corresponding heat maps for apical areas of SG cells (E") and intensity of Sec15 signals (E'''). (F) Negative correlation between intensity of Sec15 and apical areas of SG cells (n= 5 SGs; 527 cells). Asterisks: invagination pit. White lines, SG boundary.

Supplemental Figure 1

815
816

817 **Figure S2. Overexpression of spastin in the SG results in loss of acetylated-α-**
818 **tubulin.** (A-B") Confocal images of control (A-A") and spastin-CFP-overexpressing (B-
819 B") SGs stained for phalloidin (red) and an antibody against acetylated α-tubulin (Ace-α-
820 Tub; magenta). Spastin-CFP (green) signals show SG-specific overexpression of spastin
821 by *fkh-Gal4*, which leads to a loss of MTs in the SG placode. Asterisks, invagination pit.

Supplemental Figure 2

823
824

Figure S3. Verification of RNAi knockdown in the SG. (A-D') Control (A-B') and *Rab11*

RNAi (C-D') SGs immunostained for Sage (SG nuclei) and Rab11. Stage 11 (A, A', C, C')

and stage 13 (B, B', D, D') SGs are shown. (E, E') Stage 11 control (E) and *crb* RNAi (C-

D') SGs immunostained for Crb. (E'') Quantification of junctional intensity of Crb. (F, F')

Stage 11 control (E) and *E-Cad* RNAi (C-D') SGs immunostained for E-Cad. (F'')

Quantification of junctional intensity of E-Cad. n= 5 SGs for both in E'' and F''; 10 cells in

the dorsal/posterior region of each SG. **, p< 0.01 (Welch's t-test). Asterisks, invagination

pit.

Supplemental Figure 3

Figure S4. MTs play a role in apical localization of Rab11 and apical transport of Crb and E-Cad throughout SG formation. (A-B''') Confocal images of stage 16 control (A-A''') and spastin-overexpressing (B-B''') SGs immunostained for Rab11 and Nuf. In control, Rab11 and Nuf localize in the apical region of SG cells (red arrowheads in A'' and A'''). In spastin-overexpressing SGs, Rab11 and Nuf are mislocalized to aggregates in the cytoplasm of cells (yellow arrowheads in B'-B'''). (C-E'') Confocal images of stage 13 control (C-C''), spastin-overexpressing (D-D'') and *Rab11* RNAi (E-E'') SGs immunostained for Sage (SG nuclei), E-Cad and Fog. (C'-E') Compared to strong E-Cad signals primarily at adherens junctions in the control SG (C'), strong lateral E-Cad signals are shown in spastin-overexpressing and *Rab11* RNAi SG (blue arrows in E'). Insets, higher magnification of the yellow boxed regions in C'-E'. (C''-E'') Compared to strong apical Fog signals in the control SG (C''), Fog signals are very weak in spastin-overexpressing (D'') and *Rab11* RNAi (E'') SGs. (F-G''') Confocal images of stage 16 SGs stained for Crb and Rab11. (F-F''') Overexpression of Crb causes mislocalization of Crb (green) to all membrane domains and expands membranes. Rab11 is still enriched in the apical domain (red arrowheads). (G-G''') Co-overexpression of spastin and Crb results in mislocalization of Crb to large aggregates in the cytoplasm, which overlap with Rab11 (yellow arrowheads). (H-I''') Confocal images of stage 16 SGs immunostained for E-Cad and Rab11. (H-H''') In the control SG, the majority of E-Cad signals are detected at adherens junctions (red arrowheads). Small punctate E-Cad signals are occasionally shown in the basolateral region (blue arrows). (I-I''') Co-overexpression of E-Cad and spastin causes mislocalization of E-Cad to cytoplasmic aggregates, which overlap with Nuf (yellow arrowheads). White dashed line, cell and nuclear boundaries (A', B') and apical membrane (F', H', I'). Nu, nucleus.

Supplemental Figure 4

 Fly genotypes

Figures	Genotypes
Figure 1B-F''''	*Oregon R*
Figure 2A-B'', E-E'	*Oregon R*
Figure 2C-C'''	*fkh-Gal4/+*
Figure 2D-D''', F-F'	*fkh-Gal4/UAS-Spastin-CFP*
Figure 3A-A'	*fkh-Gal4/+*
Figure 3B-B'	*fkh-Gal4/UAS-Rab11-S25N-YFP*
Figure 3C-C'	*fkh-Gal4/UAS-Rab11* RNAi (TRiP.JF02812)
Figure 3D-D'	*UAS-Dicer-2/+; fkh-Gal4/UAS-Dhc64C* RNAi (GD12258)
Figure 3E-E'	*Oregon R*
Figure 3F-F'	*klar^1*
Figure 3I-I''	*UAS-Dicer-2/+; fkh-Gal4/+*
Figure 3J-J''	*UAS-Dicer-2/+; fkh-Gal4/UAS-Dhc64C RNAi* (GD12258)
Figure 3K-K''	*Oregon R*
Figure 3L-L''	*klar^{mCD4}*
Figure 4A-A''''	*sqh-GFP/+; fkh-Gal4/+*
Figure 4B-B''''	*sqh-GFP/UAS-Dicer-2; fkh-Gal4/UAS-Rab11* RNAi (UAS-Rab11.dsRNA.WIZ)
Figure 4F-F''	*fkh-Gal4 ubi-Rok-GFP/+*

Figure 4G-G''	*UAS-Dicer-2/+; fkh-Gal4 ubi-Rok-GFP/UAS-Rab11* RNAi (UAS-Rab11.dsRNA.WIZ)
Figure 5A-A'	*sqh-GFP/+; fkhGal4/+*
Figure 5B-B'	*sqh-GFP/UAS-Dicer-2; fkh-Gal4/UAS-Dhc64C* RNAi (GD12258)
Figure 5F-F'	*sqh-GFP; sqh^{AX3}*
Figure 5G-G'	*sqh-GFP; klar^1*
Figure 5K-K'	*fkh-Gal4 ubi-Rok-GFP/+*
Figure 5L-L'	*UAS-Spastin/+; fkh-Gal4 ubi-Rok-GFP/+*
Figure 5M-M'	*UAS-Dicer-2/+; fkh-Gal4 ubi-Rok-GFP/UAS-Dhc64C* RNAi (GD12258)
Figure 5N-N'	*ubi-Rok-GFP*
Figure 5O-O'	*klar^{mCD4} ubi-Rok-GFP*
Figure 6A-A''', D-D''''	*fkh-Gal4/+*
Figure 6B-B''', E-E''''	*fkh-Gal4/UAS-Spastin*
Figure 6C-C''', F-F''''	*fkh-Gal4/UAS-Rab11* RNAi (TRiP.JF02812)
Figure 7A-A'	*fkh-Gal4/+*
Figure 7B-B''	*fkh-Gal4/UAS-Crb RNAi* (TRiP. HMS01409)
Figure 7C-C'	*fkh-Gal4/UAS-E-Cad* RNAi (TRiP.HMS00693)
Figure 8A-A''	*sqh-GFP/+; fkh-Gal4/+*

Figure 8B-B''	*sqh-GFP/+; fkh-Gal4/UAS-Crb* RNAi (TRiP.HMS01409)
Figure 8C-C''	*sqh-GFP/+; fkh-Gal4/UAS-E-Cad* RNAi (TRiP.HMS00693)
Figure 8D-D''	*fkh-Gal4 ubi-Rok-GFP/+*
Figure 8E-E''	*fkh-Gal4 ubi-Rok-GFP/UAS-Crb* RNAi (TRiP.HMS01409)
Figure 8F-F''	*fkh-Gal4 ubi-Rok-GFP/UAS-E-Cad* RNAi (TRiP.HMS00693)
Figure S1A-A'''''	*Rab11-EYFP*
Figure S1B-B'''	*Rab5-EYFP*
Figure S1D-D'''	*Rab7-EYFP*
Figure S1E-E'''	*Oregon R*
Figure S2A-A''	*fkh-Gal4/+*
Figure S2B-B''	*fkh-Gal4/UAS-Spastin-CFP*
Figure S3A-A', B-B', E, F	*fkh-Gal4/+*
Figure S3C-C', D-D'	*fkh-Gal4/UAS-Rab11* RNAi (TRiP.JF02812)
Figure S3E'	*fkh-Gal4/UAS-Crb* RNAi (TRiP.HMS01409)
Figure S3F'	*fkh-Gal4/UAS-E-Cad* RNAi (TRiP.HMS00693)
Figure S4A-A''', C-C''	*fkh-Gal4/+*
Figure S4B-B''', D-D''	*fkh-Gal4/UAS-Spastin*
Figure S4E-E''	*fkh-Gal4/UAS-Rab11* RNAi (TRiP.JF02812)

Figure S4F-F'''	*UAS-Crb/+; fkh-Gal4/+*
Figure S4G-G'''	*UAS-Crb/+; fkh-Gal4/UAS-Spastin*
Figure S4H-H'''	*fkh-Gal4/UAS-Shg-GFP*
Figure S4I-I'''	*UAS-Spastin/+; fkh-Gal4/UAS-Shg-GFP*

References

Abreu-Blanco, M.T., Verboon, J.M., and Parkhurst, S.M. (2014). Coordination of Rho family GTPase activities to orchestrate cytoskeleton responses during cell wound repair. Curr Biol *24*, 144-155.

Aguilar-Aragon, M., Fletcher, G., and Thompson, B.J. (2020). The cytoskeletal motor proteins Dynein and MyoV direct apical transport of Crumbs. Dev Biol *459*, 126-137.

Anborgh, P.H., Seachrist, J.L., Dale, L.B., and Ferguson, S.S. (2000). Receptor/beta-arrestin complex formation and the differential trafficking and resensitization of beta2-adrenergic and angiotensin II type 1A receptors. Mol Endocrinol *14*, 2040-2053.

Andrew, D.J., and Ewald, A.J. (2010). Morphogenesis of epithelial tubes: Insights into tube formation, elongation, and elaboration. Dev Biol *341*, 34-55.

Benton, M.A., Frey, N., Nunes da Fonseca, R., von Levetzow, C., Stappert, D., Hakeemi, M.S., Conrads, K.H., Pechmann, M., Panfilio, K.A., Lynch, J.A*., et al.* (2019). Fog signaling has diverse roles in epithelial morphogenesis in insects. Elife *8*.

Booth, A.J.R., Blanchard, G.B., Adams, R.J., and Roper, K. (2014). A dynamic microtubule cytoskeleton directs medial actomyosin function during tube formation. Dev Cell *29*, 562-576.

Burbank, K.S., Groen, A.C., Perlman, Z.E., Fisher, D.S., and Mitchison, T.J. (2006). A new method reveals microtubule minus ends throughout the meiotic spindle. J Cell Biol *175*, 369-375.

Cerniello, F.M., Carretero, O.A., Longo Carbajosa, N.A., Cerrato, B.D., Santos, R.A., Grecco, H.E., and Gironacci, M.M. (2017). MAS1 Receptor Trafficking Involves ERK1/2 Activation Through a beta-Arrestin2-Dependent Pathway. Hypertension *70*, 982-989.

Chung, S., and Andrew, D.J. (2014). Cadherin 99C regulates apical expansion and cell rearrangement during epithelial tube elongation. Development *141*, 1950-1960.

Chung, S., Hanlon, C.D., and Andrew, D.J. (2014). Building and specializing epithelial tubular organs: the Drosophila salivary gland as a model system for revealing how epithelial organs are specified, form and specialize. Wiley Interdiscip Rev Dev Biol *3*, 281-300.

Chung, S., Kim, S., and Andrew, D.J. (2017). Uncoupling apical constriction from tissue invagination. Elife *6*.

Dale, L.B., Seachrist, J.L., Babwah, A.V., and Ferguson, S.S. (2004). Regulation of angiotensin II type 1A receptor intracellular retention, degradation, and recycling by Rab5, Rab7, and Rab11 GTPases. J Biol Chem *279*, 13110-13118.

David, D.J., Tishkina, A., and Harris, T.J. (2010). The PAR complex regulates pulsed actomyosin contractions during amnioserosa apical constriction in Drosophila. Development *137*, 1645-1655.

Du, F., Ozdowski, E.F., Kotowski, I.K., Marchuk, D.A., and Sherwood, N.T. (2010). Functional conservation of human Spastin in a Drosophila model of autosomal dominant-hereditary spastic paraplegia. Hum Mol Genet *19*, 1883-1896.

Dunst, S., Kazimiers, T., von Zadow, F., Jambor, H., Sagner, A., Brankatschk, B., Mahmoud, A., Spannl, S., Tomancak, P., Eaton, S*., et al.* (2015). Endogenously tagged rab proteins: a resource to study membrane trafficking in Drosophila. Dev Cell *33*, 351-365.

Elting, M.W., Hueschen, C.L., Udy, D.B., and Dumont, S. (2014). Force on spindle microtubule minus ends moves chromosomes. J Cell Biol *206*, 245-256.

Fan, G.H., Lapierre, L.A., Goldenring, J.R., and Richmond, A. (2003). Differential regulation of CXCR2 trafficking by Rab GTPases. Blood *101*, 2115-2124.

Flores-Benitez, D., and Knust, E. (2015). Crumbs is an essential regulator of cytoskeletal dynamics and cell-cell adhesion during dorsal closure in Drosophila. Elife *4*.

Gorvel, J.P., Chavrier, P., Zerial, M., and Gruenberg, J. (1991). rab5 controls early endosome fusion in vitro. Cell *64*, 915-925.

Goshima, G., Nedelec, F., and Vale, R.D. (2005). Mechanisms for focusing mitotic spindle poles by minus
end-directed motor proteins. J Cell Biol *171*, 229-240.
Gross, S.P., Welte, M.A., Block, S.M., and Wieschaus, E.F. (2000). Dynein-mediated cargo transport in
vivo. A switch controls travel distance. J Cell Biol *148*, 945-956.
Guglielmi, G., Barry, J.D., Huber, W., and De Renzis, S. (2015). An Optogenetic Method to Modulate Cell
Contractility during Tissue Morphogenesis. Dev Cell *35*, 646-660.
Hamelin, E., Theriault, C., Laroche, G., and Parent, J.L. (2005). The intracellular trafficking of the G
protein-coupled receptor TPbeta depends on a direct interaction with Rab11. J Biol Chem *280*, 36195-
36205.
Heald, R., Tournebize, R., Blank, T., Sandaltzopoulos, R., Becker, P., Hyman, A., and Karsenti, E. (1996).
Self-organization of microtubules into bipolar spindles around artificial chromosomes in Xenopus egg
extracts. Nature *382*, 420-425.
Hunyady, L., Baukal, A.J., Gaborik, Z., Olivares-Reyes, J.A., Bor, M., Szaszak, M., Lodge, R., Catt, K.J., and
Balla, T. (2002). Differential PI 3-kinase dependence of early and late phases of recycling of the
internalized AT1 angiotensin receptor. J Cell Biol *157*, 1211-1222.
Innamorati, G., Le Gouill, C., Balamotis, M., and Birnbaumer, M. (2001). The long and the short cycle.
Alternative intracellular routes for trafficking of G-protein-coupled receptors. J Biol Chem *276*, 13096-
13103.
Izquierdo, E., Quinkler, T., and De Renzis, S. (2018). Guided morphogenesis through optogenetic
activation of Rho signalling during early Drosophila embryogenesis. Nat Commun *9*, 2366.
Jha, A., van Zanten, T.S., Philippe, J.M., Mayor, S., and Lecuit, T. (2018). Quantitative Control of GPCR
Organization and Signaling by Endocytosis in Epithelial Morphogenesis. Curr Biol *28*, 1570-1584 e1576.
Jouette, J., Guichet, A., and Claret, S.B. (2019). Dynein-mediated transport and membrane trafficking
control PAR3 polarised distribution. Elife *8*.
Kerridge, S., Munjal, A., Philippe, J.M., Jha, A., de las Bayonas, A.G., Saurin, A.J., and Lecuit, T. (2016).
Modular activation of Rho1 by GPCR signalling imparts polarized myosin II activation during
morphogenesis. Nat Cell Biol *18*, 261-270.
Khanal, I., Elbediwy, A., Diaz de la Loza Mdel, C., Fletcher, G.C., and Thompson, B.J. (2016). Shot and
Patronin polarise microtubules to direct membrane traffic and biogenesis of microvilli in epithelia. J Cell
Sci *129*, 2651-2659.
Khodjakov, A., Copenagle, L., Gordon, M.B., Compton, D.A., and Kapoor, T.M. (2003). Minus-end capture
of preformed kinetochore fibers contributes to spindle morphogenesis. J Cell Biol *160*, 671-683.
Ko, C.S., Tserunyan, V., and Martin, A.C. (2019). Microtubules promote intercellular contractile force
transmission during tissue folding. J Cell Biol *218*, 2726-2742.
Laan, L., Pavin, N., Husson, J., Romet-Lemonne, G., van Duijn, M., Lopez, M.P., Vale, R.D., Julicher, F.,
Reck-Peterson, S.L., and Dogterom, M. (2012). Cortical dynein controls microtubule dynamics to
generate pulling forces that position microtubule asters. Cell *148*, 502-514.
Langevin, J., Morgan, M.J., Sibarita, J.B., Aresta, S., Murthy, M., Schwarz, T., Camonis, J., and Bellaiche, Y.
(2005). Drosophila exocyst components Sec5, Sec6, and Sec15 regulate DE-Cadherin trafficking from
recycling endosomes to the plasma membrane. Dev Cell *9*, 365-376.
Le Droguen, P.M., Claret, S., Guichet, A., and Brodu, V. (2015). Microtubule-dependent apical restriction
of recycling endosomes sustains adherens junctions during morphogenesis of the Drosophila tracheal
system. Development *142*, 363-374.
Lee, J.Y., and Harland, R.M. (2010). Endocytosis is required for efficient apical constriction during
Xenopus gastrulation. Curr Biol *20*, 253-258.
Letizia, A., Sotillos, S., Campuzano, S., and Llimargas, M. (2011). Regulated Crb accumulation controls
apical constriction and invagination in Drosophila tracheal cells. J Cell Sci *124*, 240-251.

Li, M., McGrail, M., Serr, M., and Hays, T.S. (1994). Drosophila cytoplasmic dynein, a microtubule motor
that is asymmetrically localized in the oocyte. J Cell Biol *126*, 1475-1494.
Lord, S.J., Velle, K.B., Mullins, R.D., and Fritz-Laylin, L.K. (2020). SuperPlots: Communicating
reproducibility and variability in cell biology. J Cell Biol *219*.
Manning, A.J., Peters, K.A., Peifer, M., and Rogers, S.L. (2013). Regulation of epithelial morphogenesis by
the G protein-coupled receptor mist and its ligand fog. Sci Signal *6*, ra98.
Manning, A.J., and Rogers, S.L. (2014). The Fog signaling pathway: insights into signaling in
morphogenesis. Dev Biol *394*, 6-14.
Martin, A.C., and Goldstein, B. (2014). Apical constriction: themes and variations on a cellular
mechanism driving morphogenesis. Development *141*, 1987-1998.
Martin, A.C., Kaschube, M., and Wieschaus, E.F. (2009). Pulsed contractions of an actin-myosin network
drive apical constriction. Nature *457*, 495-499.
Mason, F.M., Tworoger, M., and Martin, A.C. (2013). Apical domain polarization localizes actin-myosin
activity to drive ratchet-like apical constriction. Nat Cell Biol *15*, 926-936.
Meresse, S., Gorvel, J.P., and Chavrier, P. (1995). The rab7 GTPase resides on a vesicular compartment
connected to lysosomes. J Cell Sci *108 (Pt 11)*, 3349-3358.
Miao, H., Vanderleest, T.E., Jewett, C.E., Loerke, D., and Blankenship, J.T. (2019). Cell ratcheting through
the Sbf RabGEF directs force balancing and stepped apical constriction. J Cell Biol.
Myat, M.M., and Andrew, D.J. (2000). Organ shape in the Drosophila salivary gland is controlled by
regulated, sequential internalization of the primordia. Development *127*, 679-691.
Myat, M.M., and Andrew, D.J. (2002). Epithelial tube morphology is determined by the polarized growth
and delivery of apical membrane. Cell *111*, 879-891.
Ossipova, O., Kim, K., Lake, B.B., Itoh, K., Ioannou, A., and Sokol, S.Y. (2014). Role of Rab11 in planar cell
polarity and apical constriction during vertebrate neural tube closure. Nat Commun *5*, 3734.
Rauzi, M., Lenne, P.F., and Lecuit, T. (2010). Planar polarized actomyosin contractile flows control
epithelial junction remodelling. Nature *468*, 1110-1114.
Riggs, B., Rothwell, W., Mische, S., Hickson, G.R., Matheson, J., Hays, T.S., Gould, G.W., and Sullivan, W.
(2003). Actin cytoskeleton remodeling during early Drosophila furrow formation requires recycling
endosomal components Nuclear-fallout and Rab11. J Cell Biol *163*, 143-154.
Roeth, J.F., Sawyer, J.K., Wilner, D.A., and Peifer, M. (2009). Rab11 helps maintain apical crumbs and
adherens junctions in the Drosophila embryonic ectoderm. PLoS One *4*, e7634.
Rogers, S.L., Wiedemann, U., Hacker, U., Turck, C., and Vale, R.D. (2004). Drosophila RhoGEF2 associates
with microtubule plus ends in an EB1-dependent manner. Curr Biol *14*, 1827-1833.
Roper, K. (2012). Anisotropy of Crumbs and aPKC drives myosin cable assembly during tube formation.
Dev Cell *23*, 939-953.
Royou, A., Field, C., Sisson, J.C., Sullivan, W., and Karess, R. (2004). Reassessing the role and dynamics of
nonmuscle myosin II during furrow formation in early Drosophila embryos. Mol Biol Cell *15*, 838-850.
Sanchez-Corrales, Y.E., Blanchard, G.B., and Roper, K. (2018). Radially patterned cell behaviours during
tube budding from an epithelium. Elife *7*.
Sawyer, J.M., Harrell, J.R., Shemer, G., Sullivan-Brown, J., Roh-Johnson, M., and Goldstein, B. (2010).
Apical constriction: a cell shape change that can drive morphogenesis. Dev Biol *341*, 5-19.
Sherwood, N.T., Sun, Q., Xue, M., Zhang, B., and Zinn, K. (2004). Drosophila spastin regulates synaptic
microtubule networks and is required for normal motor function. PLoS Biol *2*, e429.
Sidor, C., Stevens, T.J., Jin, L., Boulanger, J., and Roper, K. (2020). Rho-Kinase Planar Polarization at
Tissue Boundaries Depends on Phospho-regulation of Membrane Residence Time. Dev Cell *52*, 364-378
e367.
Takeda, M., Sami, M.M., and Wang, Y.C. (2018). A homeostatic apical microtubule network shortens
cells for epithelial folding via a basal polarity shift. Nat Cell Biol *20*, 36-45.

Tan, R., Foster, P.J., Needleman, D.J., and McKenney, R.J. (2018). Cooperative Accumulation of Dynein-
Dynactin at Microtubule Minus-Ends Drives Microtubule Network Reorganization. Dev Cell *44*, 233-247
e234.
Ullrich, O., Reinsch, S., Urbe, S., Zerial, M., and Parton, R.G. (1996). Rab11 regulates recycling through
the pericentriolar recycling endosome. J Cell Biol *135*, 913-924.
Urbansky, S., Gonzalez Avalos, P., Wosch, M., and Lemke, S. (2016). Folded gastrulation and T48 drive
the evolution of coordinated mesoderm internalization in flies. Elife *5*.
Volpicelli, L.A., Lah, J.J., Fang, G., Goldenring, J.R., and Levey, A.I. (2002). Rab11a and myosin Vb regulate
recycling of the M4 muscarinic acetylcholine receptor. J Neurosci *22*, 9776-9784.
Westermann, S., and Weber, K. (2003). Post-translational modifications regulate microtubule function.
Nat Rev Mol Cell Biol *4*, 938-947.
Wichmann, H., Hengst, L., and Gallwitz, D. (1992). Endocytosis in yeast: evidence for the involvement of
a small GTP-binding protein (Ypt7p). Cell *71*, 1131-1142.
Wodarz, A., Grawe, F., and Knust, E. (1993). CRUMBS is involved in the control of apical protein targeting
during Drosophila epithelial development. Mech Dev *44*, 175-187.
Yvon, A.M., Gross, D.J., and Wadsworth, P. (2001). Antagonistic forces generated by myosin II and
cytoplasmic dynein regulate microtubule turnover, movement, and organization in interphase cells. Proc
Natl Acad Sci U S A *98*, 8656-8661.
Zhang, J., Schulze, K.L., Hiesinger, P.R., Suyama, K., Wang, S., Fish, M., Acar, M., Hoskins, R.A., Bellen,
H.J., and Scott, M.P. (2007). Thirty-one flavors of Drosophila rab proteins. Genetics *176*, 1307-1322.
Zhang, X.M., Ellis, S., Sriratana, A., Mitchell, C.A., and Rowe, T. (2004). Sec15 is an effector for the Rab11
GTPase in mammalian cells. J Biol Chem *279*, 43027-43034.

1028