How do thalamocortical interactions shape causal structure among resting state brain networks during aging?

Moumita Das, Vanshika Singh, Arpan Banerjee, Dipanjan Roy

Cognitive Brain Dynamics Lab, National Brain Research Centre, NH 8,

Manesar, Gurgaon 122052, India

#Email: arpan@nbrc.ac.in, dipanjan.nbrc@gov.in

Abstract

The human brain undergoes both significant structural and functional changes across the lifespan. It is important to understand the underlying causal relationship of the emerging dynamical changes in functional connectivity with age. On average, functional connectivity within resting-state networks weakens in magnitude while connections between resting-state networks tend to increase with age. Further, few recent studies show that effective connectivity within and between large scale resting-state functional networks changes over the healthy lifespan. Motivated by these findings we move one step forward to investigate the effect of the thalamus in the context of healthy aging. Using directed connectivity and weighted net causal outflow measures on resting-state fMRI data, we examine the age-related changes in both cortical and thalamocortical causal interactions within and between resting-state networks. The three of core neurocognitive networks DMN, SN, CEN networks are identified independently by carrying out ICA as well as spatially matching of hub regions with the important RSNs previously reported in the literature. Thereafter, multivariate GCA was performed to test for

causality index between ROIs with and without the inclusion of left and right thalamus. There are two major findings, firstly, we observe that within network causal connections become progressively weaker with age, however, between network causal connections are getting stronger with age among core neurocognitive networks, primarily a reflection of within and between network resting-state functional connectivity. Secondly, significant modifications were found in causal connections and net causal outflows in the presence of thalamus. Finally, we found that the thalamus plays a crucial role as an exogenous drive in the reorganization of within network causal outflow, while Salience network plays a critical role in mediating be-tween network causal outflow with age among cortical networks. Our findings with the weighted causal outflow measures strengthen the hypothesis that balancing within and between network connectivity is perhaps critical for the preservation of cognitive functions with aging.

Keywords: Aging, Effective Connectivity, Multivariate Granger Causal, Neurocognitive Net-works, Salience Network, Thalamus, Weighted Net Causal Outflow.

1.Introduction

In the last decade, there has been an enormous interest in studying the coordinated activity in distributed brain areas when the human being is engaged in internally driven tasks, such as meandering through self-referential thoughts while seemingly at rest or more specifically not engaged in a state of goal directed action/ perception (Raichle, 2009; (Bressler & Menon, 2010) ; (Deco, Jirsa, & McIntosh, 2011)). The large scale neurocognitive networks (Bressler Menon, 2010) that chaperons the activity of resting state has been broadly classified in three distinct sub-networks based on the correlation patterns estimated from BOLD time series signals,

namely, the default mode network (DMN), saliency network (SN), and central executive net-

work (CEN) (Menon 2011) . The DMN comprises posterior cingulate cortex (PCC) and medial

prefrontal cortex (MPFC) and is implicated in self-referential mental activities (Buckner,

Andrews-Hanna, & Schacter, 2008), (Andrews-Hanna, Reidler, Sepulcre, Poulin, & Buckner,

2010) , (Raichle, 2015)). The CEN, comprises rostral and caudal bilateral middle frontal cortex

(MFC), bilateral superior Parietal Cortex (SPC) and implicated in decision making and exec-

utive functions (Corbetta & Shulman, 2002); (Fox, Corbetta, Snyder, Vincent, & Raichle,

2006)). The SN, which comprise bilateral anterior insula (AI) and caudal, rostral bilateral an-

terior cingulate cortices (ACC) is important for detection and mapping of salient inputs and

routing these inputs to control areas for mediating cognitive control ((Menon & Uddin,

2010);(Uddin, 2015)).

Interestingly, modification of interconnections within and between DMN, CEN, SN is altered

in a large number of psychiatric and neurological disorders, for instance, Alzheimer's disease,

Autism Spectrum Disorder (ASD), Attention Deficit Hyperactivity Disorder (ADHD), psycho-

sis and depression (Woodward & Cascio, 2015);(Abi-Dargham & Horga, 2016)). SN, in addi-

tion to detecting salient stimuli, plays important role in switching between DMN and CEN in

the task condition as well as in resting state ((Devarajan, Levitin, & Menon, 2008)). Thus, it

integrates and balances internal mental processes with external stimulus-driven cognitive and

affective processes. During the performance of many tasks, correlation among the nodes of SN

in tandem with CEN increases while the corresponding correlation among the DMN nodes

decreases.

An important caveat to all the resting brain network studies are that they mostly concentrate on

cortical nodes, and ignores the crucial thalamocortical interactions. The thalamus, a centrally

located relay station for transmitting information throughout the brain, participates in communication with many associative brain regions and involves global multifunctional pathways. Incorporating 10,449 meta-studies, Hwang, Bertolero, Liu, & D'Esposito, 2017 recently showed that the thalamus is engaged in multiple cognitive functions and is a critical integrative hub for functional brain networks. Cross-sectional studies of normal aging have also reported smaller thalamic volumes in older than younger adults (Cherubini, Péran, Caltagirone, Sabatini, & Spalletta, 2009, Hughes et al., 2012;). However, exactly how normal aging affects the thalamic interconnections with other brain networks and its implication in cognitive changes are not completely understood, and thus warrant further investigation. From a methodological standpoint if thalamus acts as a common source to cortical inputs during both rest and task as proposed by theories such as thalamocortical dysryhtmia ((Llinás, Ribary, Jeanmonod, Kronberg, & Mitra, 1999),(Vanneste, Song, & De Ridder, 2018)), a causality analysis that ignores thalamocortical contribution to brain dynamics is of very limited scope and possibly paint an inaccurate account of underlying complexity. Thus, the primary goal of this article is to re-evaluate the interactions among the RSNs: DMN, SN and CEN by taking into consideration the thalamocortical interactions and as a proof of concept characterize how lifespan sculpts the causal neuro-circuity.

2. Material and Methods

2.1 Participants

25 young and 24 elderly individuals participated in this study after providing the written consent. Young (13 females and 12 males) ranged in age from 18-33 years (mean age = 25.68 years). Old (18 females and 6 males) ranged in age from 39-80 years (mean age=58.1 years). All participants gave written informed consent and the study was performed under the compliance of laws and guidelines approved by the ethics committee of Charité University, Berlin.

2.2 Data Acquisition

RS fMRI as well as corresponding diffusion weighted (dw) MRI data were collected from 49 healthy participants at the Charité University Berlin, Germany (Schirner, Rothmeier, Jirsa, McIntosh, & Ritter, 2015). Each fMRI dataset amounts to 661 time points recorded at TR=2s, i.e. about 22 minutes. In the same session, EEG was also recorded, but we do not use the EEG data for our current analysis. No other controlled task was performed. Resting state BOLD activity was recorded while subjects were asked to stay awake with their eyes closed, using a 3T Siemens Trim Trio scanner and a 12 channel Siemens head coil (voxel size $3mm \times 3mm \times 3mm$). Voxel time courses are averaged inside ROIs defined by the Desikan-Killiany atlas (Desikan et al., 2006) as implemented in FreeSurfer. The empirical BOLD time series signals from Regions of Interest (ROI) used in this paper for net causal flow estimations is generated by using an automated pipeline ((Schirner, Rothmeier, Jirsa, McIntosh, & Ritter, 2015)).

2.3 Data Analysis

2.3.1 rs-fMRI preprocessing

The major pre-processing steps on T1 anatomical images were skull stripping, removal of non-brain tissue, brain mask generation, cortical reconstruction, motion correction, intensity normalization, WM and subcortical segmentation, cortical tessellation generating GM–WM and GM–pia interface surface-triangulations and probabilistic atlas-based cortical and subcortical parcellation. These parcellations, segmentations and masks were then used to guide the probabilistic tractography algorithm to estimate connection strengths (a value in the range 0 to 1) between each pair of areas in the cortical gray matter parcellation. The parcellation used in this study is Desikan–Killiany parcellation (Desikan et al., 2006) which consists of 68 cortical regions of interest (ROIs) with 34 ROIs in each hemisphere and 32 subcortical regions. For our

122 present analysis along with 68 cortical regions, two thalamic regions, left and right thalamus

123 were selected based on this parcellation. FC matrices generated from each subject's MRI data

124 are averaged element-wise to obtain an averaged FC matrix for younger and elderly cohort.

2.3.2 Selection and extraction of three large-scale resting state networks

To identify RSN activity a spatial Group ICA decomposition was performed for the fMRI data

of all subjects using FSL MELODIC ((Beckmann & Smith, 2004)) (MELODIC v4.0; FMRIB

Oxford University, UK) with the following parameters: high pass filter cut off: 100 s,

MCFLIRT motion correction, BET brain extraction, spatial smoothing, normalization to

MNI152, temporal concatenation, dimensionality restriction to 30 output components. ICs that

correspond to RSNs were automatically identified by spatial correlation with the 9 out of the

10 well-matched pairs of networks of the 29,671-subject Brain Map activation database as de-

scribed in (Smith et al., 2009) (excluding the cerebellum network). Subsequently, the three

key intrinsic networks were identified by spatially matching with pre-existing tem-

plates following widely accepted seven network resting state parcellation proposed by Buck-

ner and colleagues (Buckner, Krienen, Castellanos, Diaz, & Yeo, 2011), each of the cortical

regions were classified further down to seven parcellated resting networks. In our subsequent

analysis, we considered three resting state networks, namely DMN, SN, CEN (Node details are

provided in **Table: 1**). We chose for our study Inferior parietal lobule (IPL), PCC and

MOF as they constitute core part of DMN ((Andrews-Hanna et al., 2010);(Dixon et

al., 2017)) consistently showed activation during mental rumination and self-related

processing, mind wandering. The SN, which comprises the anterior insula and caudal,

rostral anterior cingulate cortex , is important for detection of salience events and

switching between other large scale networks ((Menon & Uddin, 2010);(Uddin,

2015)). Bilateral Rostral and Caudal middle frontal gyrus, superior parietal lobule

were selected as nodes of CEN. The core DMN regions selected in this work consist-

ently showed anticorrelation with SNs ((Fransson, 2005); (Uddin, Kelly, Biswal,

Castellanos, & Milham, 2009);(Dixon et al., 2017))

For within network analysis, we used the extracted time series for each six ROI within each

respective network. For between network analysis, we combined time series of six ROIs, using

principal component analysis, to create a single representative time series of every network.

2.2.4 Effective Connectivity: Multivariate Granger Causal Analysis

Brain signals of different nodes within and between networks are expected to follow a multi-

variate probability distribution. For instance, let X_t, Y_t, Z_t be a time series of 3 nodes in a net-

work. If there are joint dependencies between X_t, Y_t, Z_t and if we calculate unconditional

granger causality between X and Y, spurious causalities may occur due to common dependency

on Z. Thus, to eliminate the possibility of spurious causalities between two time series

Multivariate Granger Causality analysis (GCA) was performed to assess the causal influence

between nodes of SN, CEN and DMN based on the methods described in ((Barrett, Barnett, &

Seth, 2010)). In MVGC spurious causalities are eliminated by conditioning out the common

dependencies. According to MVGC approach, if we wish to test the causality from

Y to X, we have to consider the full and reduced regressions of the following form

$$X_t = \sum_{k=1}^{p} A_{xx,k} \, X_{t-k} + \sum_{k=1}^{p} A_{xy,k} \, Y_{t-k} + \sum_{k=1}^{p} A_{xz,k} \, Z_{t-k} + \varepsilon_{x,t} \qquad (1)$$

$$X_t = \sum_{k=1}^{p} A'_{xx,k} \, X_{t-k} + \sum_{k=1}^{p} A'_{xz,k} \, Z_{t-k} + \varepsilon'_{x,t} \qquad (2)$$

In full regression (1), the dependence of X on the past of Y, given its own past and the past of

Z is incorporated in the coefficients $A_{xy,k}$. If in particular there is no conditional dependence

of X on the past of Y if all the coefficients associated with Y values is zero. This leads to the

reduced regression (2), where past values of Y are omitted. Therefore, the null hypothesis of

zero causality is :

$H_0 : A_{xy.1} = A_{xy.2} = \ldots\ldots.. = A_{xy.p} = 0$

vs H_1: $at\ least\ one\ A_{xy.k} \neq 0\ for\ k = 1(1)p$ (3)

The granger causality value quantifies the degree to which the full regression model is a better

candidate compared to the reduced regression to model X_t. Appropriate measure for the model

comparison is the logarithm of the ratio of their likelihood values. The joint likelihood of the

var model is $L = \left| \Sigma^{\frac{-(m-p)}{2}} \right|$ (where m = total number of nodes, p = estimated model order).

This motivates the definition of conditional G- causality statistic as the appropriate log-likeli-

hood ratio. The conditional G -causality from Y to X is defined by

$F_{y \to x} = \ln \left| \frac{\Sigma'_{xx}}{\Sigma_{xx}} \right|$ (4)

Where $cov(\varepsilon_{x,t}) = \Sigma_{xx}$, $cov(\varepsilon'_{x,t}) = \Sigma'_{xx}$ are the error variances of full model and reduced

model respectively.

Multivariate granger causality toolbox (Barnett et al.(2014)) was employed to perform the anal-

ysis. For each model order p up to the maximum model order (in our case we set it 20), joint

likelihood $L = \left| \Sigma^{\frac{-(m-p)}{2}} \right|$ was calculated. Akaike's information criterion and Bayesian infor-

mation criterion was calculated using L. The order of the vector autoregressive (VAR) model

used for computation of the influence measure was selected using the Bayesian information

criterion (BIC). Corresponding VAR model parameters, var model coefficients, and covariance

matrices were estimated for the estimated model order. Using the reverse solution of the Yule

Walker equations, the autocovariance sequences was calculated. For granger causal estimation,

var parameters were calculated for both the full and reduced regressions. Granger causality

value was calculated using (4). Significance was tested using F-test. Since multiple causalities

were tested simultaneously, FDR correction was used to adjust for multiple hypotheses.

2.2.5 Network Analysis

We further calculated the weighted net granger causal flow to characterise the causal networks

in young and elderly groups. To calculate the weighted net granger causal flow, we selected

the significant causal outflow connections between the nodes. Weighted out-degree of a node

is defined by sum of the strength (granger causal indexes) of significant causal connections

from a node in a network to any other node. Likewise weighted in-degree of a node is sum of

the strength of causal inflow connections to a node in the network from any other node.

Weighted net causal flow was defined as weighted (Out-In) degree. For example, weighted net

granger causal outflow of node X, say Δ_x can be expressed using the following formula:

$\Delta_x = (F_{x \to y} + F_{x \to z}) - (F_{y \to x} + F_{z \to x})$ …………… (5), provided all F values are significant.

Non-significant F-values should be replaced by zero. Similarly, one can calculate Δ_y , Δ_z .

Though F values are positive, weighted net granger causal values can be positive as well as

negative also. Positive Δ for a particular node implies, higher causal influence of that node on

the other nodes of that network, furthermore negative Δ signifies, that particular node is caus-

ally influenced by other nodes of the network.

To compare the net flow of different nodes between young and old age groups we generated

100 bootstrap samples of granger causality index matrices via nonparametric sampling.

For the purpose of comparison, we constructed the distribution of weighted net causal outflow

based on the significant causal connections (p<0.05, FDR corrected for multiple comparisons).

The distribution of weighted net causal outflow was calculated for different nodes within and

between network and Wilcoxon signed rank tests were performed to test the significant differ-

ences in the net causal outflows for different age group.

3.0 Results

3.1 Reorganization of causality within intrinsic networks with age

3.1.1 Central Executive Network

In CEN, we extracted time series from six nodes (see **figure 2**), namely lRMFG, rRMFG, lCMFG, rCMFG, lSPL, rSPL. The strength of the causal interaction from the rRMFG to the rCMFG was highest among all causal interactions for both elderly and young groups (**figure 3A, 3D**). In addition, significant causal connections ($p < 0.01$, false discovery rate (FDR) corrected) were found from the lRMFG to the lCMFG, and the lSPL for young individuals (**figure 3D**). On the other hand, for elderly group, significant directed causal connectivity was found from the rRMFG to the lSPL and rSPL (in addition to the rCMFG) (**figure 3A**). In contrast to DMN, visually inspecting greater number of inter hemispheric connections were found in elderly individuals in CEN (**figure 3A**).

To further investigate the network properties, weighted net causal outflow for within CEN nodes were calculated. lRMFG and rRMFG acted as a causal outflow hub for young and elderly individuals respectively. Causal outflows were significantly different for all nodes (p<0.01). Except lRMFG and rRMFG, all nodes had negative causal outflow (causal inflow) for young individuals (**figure 4A**). In old cohorts, other than rRMFG, only lCMFG had small positive outflow. All the remaining nodes had causal inflow in old group.

3.1.2 Salience Network

Next we proceed with analysis of salience network which comprise of six nodes (see **figure 2**),

namely lInsula, rInsula, lcACC, rcACC, lrACC, rrACC. For within network analysis, multi-

variate granger causality analysis revealed a smaller number of significant directed causal in-

fluences compared to other two networks. The lInsula influenced the lCACC , and the lRACC

in young and elderly groups (**figure 3B, 3E**). Additionally, in young significant causal influ-

ences were found from the rInsula to the rCACC and from the lRACC to the lInsula . The

rInsula also causally influenced the rCACC in young but not in elderly age group (**figure 3E**).

Weighted net causal outflow analysis revealed lInsula as a causal outlow hub in the salience

network for both groups. Other than that, rInsula had positive causal outflow in both groups.

lRACC had positive outflow for young cohorts (**figure 4B**). All other nodes had negative

outflow (inflow) for both groups. Causal outflows/inflows were significantly different (p<0.01)

for all the nodes.

3.1.3 Default Mode Network

We performed multivariate granger causal analysis on the extracted time series for each of the

six DMN nodes (see **figure 2**) for both young and elderly individuals to depict the age related

alteration in dominant direction of influence ($p < 0.05$, false discovery rate (FDR) cor-

rected).While in the younger group, GCA revealed significant directed causal connectivity

from the lMOF to the lIPL, lPCC and rMOF to the rIPL, rPCC, lMOF, and lIPL to rIPL, lMOF

as shown in **figure 3F**, in elderly individuals, such causal connections from lMOF to lIPL, and

lPCC, and lIPL to rIPL were completely absent (**figure 3C**) suggesting an age related decrease

in causal drive within DMN networks. Furthermore, the causal connections between rMOF and

rIPL was reversed. Visual inspection revealed more number of inter hemispheric causal con-

nections were present in young group compared to old one.

Next, we estimated weighted net causal outflow or weighted (Out-In) degree in both young and

elderly group. For the young cohort, rMOF acted as a causal outflow hub among the nodes in

DMN, whereas rIPC acted as same for elderly group (**figure 4C**). Based on the 100 bootstrap

samples the distribution of weighted net causal outflow was calculated for each of the 6 nodes

in DMN. Causal outflows were significantly different in young compared to old (Mann-Whit-

ney test p value < 0.001).

3.2 Comparison of within network causality in three resting state networks

in presence of thalamus in young versus old individuals

To investigate the effect of thalamus in shaping within and between network causality, we

included thalamus as an additional node in our analysis. For each of the three resting state

networks, for within network analysis, we included the right and the left thalami, as the seventh

and the eighth nodes. Since the mechanism of the MVGC changes with the number of nodes,

the causal strengths were not directly comparable between two networks having different num-

ber of nodes. In other words, it is not meaningful to quantify the difference between causal

strengths of the same RSN networks before and after the inclusion of the thalami. Instead of

that, we focused our analysis to find out the changes in the pattern of causal connections and

net causal outflows.

3.2.1 Central Executive Network

In elderly group all the connections except, rRMFG to lSPL remained significant ($p < 0.05$,

false discovery rate (FDR) corrected FDR corrected). The connections from the rrMFG to the

lSPL was mediated by the lThal. Some significant causal connections were emerging from both

the thalami, from the lthal to the lRMFG, the rRMFG, the lSPL, the rSPL, from the rthal to the

lRMFG, the rSPL, and the lSPL for the elderly groups (**figure 5A**). For young groups, all the

connections without thalamus also found significant after inclusion of the thalamus (figure).

Additionally, the left thalamus causally influenced the lRMFG, the lCMFG, and the rRMFG

(**figure 5D**). Right thalamus influenced the node lRMFG.

Net granger causal outflows were significantly changed after inclusion of thalamus for both

the young and elderly groups. Effects of thalamus was greater in the younger individual's net

causal values (**figure 6A**). Left thalamus acted as a causal outflow hub for both the groups. But

the causal outflow in the left thalamus was higher in young compared to elderly group. Causal

outflows were significantly different in both the groups for all the eight nodes ($p<0.01$).

3.3.2 Salience Network

Among three resting state networks, the SN was least affected after inclusion of thalamus. No

changes were found in causal structure after inclusion of thalamus in elderly individuals (**figure**

5B). In young individuals, significant causal connections were found from the lThal to the

lInsula and from the rThal to the rInsula (**figure 5E**). No significant changes were found in the

net causal outflow pattern in both the groups. Left thalamus exhibited positive outflow, higher

in the case of old individuals ($p<0.05$). Right thalamus received marginally small negative out-

flow (inflow) for both groups (**figure 6B**).

3.3.3 Default Mode Network

After inclusion of the thalamus in default mode network, there were restructuring in the causal

connectivity pattern found in the elderly individuals. On performing multivariate granger

causal analysis on DMN after addition of thalami, some of the earlier significant causal con-

nections disappeared while some thalamo-cortical causal connections emerged as important

connections for both the age groups (**figure 5C, 5F**). Causal connection from rIPL to rMOF

continued to be the strongest connection in elderly group as found in the previous section with-

out thalamus (**figure 5C**). Other than that, GCA revealed significant causal connections from

the rMOF to the rPCC, from the rThal to the lIPl, from the lIPL to the rThal, from the rThal to

the lMOF for the elderly people (**figure 5C**). After the introduction of thalami, some of the

connections between the left hemispheric nodes disappeared in elderly. For the young group,

the effect of the inclusion of thalamus was more pronounced compared to the elderly individ-

uals (**figure 5F**). Instead of the connection from the lMOF to the lPCC, the connection from

the lThal to the lPCC emerged as the strongest one after the inclusion of the thalami (though

the lMFC to the lPCC connection also remained significant). In addition to that, other signifi-

cant connections were from the lMOF to the lIPL, from the lThal to the lMOF, from the lThal

to rThal, from the lThal to the rPCC, from the rThal to the lMFC, from the rThal to the rMFC,

from the rMFC to the rPCC, and from the rMFC to the rIPC (**figure 5F**) suggesting substantial

effect of thalamus in reorganizing within network causality in cortical networks and also re-

vealing the effect is the strongest in the younger group compared to elderly. This could be

related to the thalamic decline with aging.

Net granger causal outflows were significantly changed after inclusion of the thalamus for the

young individuals. Left thalamus emerged as a causal outflow hub for young group. Patterns

in the net causal outflows were unchanged in elderly group (**figure 6C**). rIPL continued to be

causal outflow hub for elderly group, even after inclusion of thalamus. Causal outflows were

significantly different in both groups (p<0.01).

3.2 Reorganization of between network causality with age

For between network analysis, we employed principal component analysis, to combine the time

series from each of the resting state networks. We took first principal component of the all

nodal time series of a network, as one node consisting of the all information of that particular

resting state network and performed multivariate granger causality analysis among three nodes.

In between network analysis, salience network exhibited prominent causal influence on both

default mode network and central executive network in both groups (**figure 7A and 7B**).

Causal strengths were significantly higher in old groups compared to young ($p<0.01$). Other

directed connectivity between these three nodes were also significant, with presence of higher

causal strength in old compared to young suggesting an increase in between network causality

with aging.

Next we estimated the weighted net granger causal outflow in between network nodes. Among

the three RSNs, the SN was causal outflow hub in the between network analysis. Causal out-

flows were significantly different in young and elderly groups ($p< 0.01$) (**figure 8A**).

We repeated between network analysis, with second principal component of all nodal time

series of a network taken as a node. No causal connection was found significant, confirming

the fact that, first principal component sufficiently explained all the variabilities present in the

time series of three intrinsic networks.

3.4 Comparison of between network effective connectivity in presence of
Thalamus in young versus old individuals
362
Next we investigated between network causality in presence of the thalamus for both groups.

Causal connections from the thalamus to all three network nodes, namely the DMN, the SN,

and the CEN in both age groups were found significant ($p< 0.05$, false discovery rate (FDR)

corrected). Thalamus was also causally driven by CEN and SN for both elderly and young

groups (**figure 7C and 7D**).

Weighted net causal outflows were significantly affected after inclusion of thalamus. Now thal-

amus acts as a causal outflow hub for elderly group. SN received highest causal outflow for

young group (**figure 8B**). Unlike the within network results, in between network analysis

causal outflows were greater in the elderly group. Among other three networks, the SN had

positive outflow and the DMN had negative outflow for both the groups. After inclusion of

thalamus, outflows in the CEN changed its direction in old cohorts.

Overall, thalamus had not changed the causal connectivity pattern between three resting state

networks. After inclusion of thalamus, the causality dynamics between three resting state net-

works remained unaltered. However, thalamus received causal influences from SN and CEN,

also influenced resting state networks. Considering the net causal outflows, causal outflows for

the elderly group was affected with inclusion of the thalamus. The effect of thalamus was dis-

similar in between network analysis compared to within network analysis. Effect was higher

in elderly group.

3.5 Replication Analysis

We identified a group of 50 young and 50 elderly participants from the publicly available Cam-

bridge Aging Neuroscience dataset (https://camcan-archive.mrc-cbu.cam.ac.uk//dataaccess/)

in the age range of 18–80 years who did not differ in mean age, gender distribution, or IQ from

Berlin dataset. Using this new dataset for independent verification, we conducted identical

functional and effective connectivity analyses as in the original dataset and also analysed those

measures by varying parcellation and node numbers in each of the three-core neurocognitive

resting state networks of interest.

In the replication analysis using the Harvard Oxford atlas, frontoparietal network areas were

chosen as the nodes for CEN. These included bilateral FP, LPFC, IFG, SFG and MFG.

Bilateral SPL and PPC were also included in the CEN network in consensus with existing

literature. The younger cohort exhibited greater number of causal interactions than the older

group (**figure 9A, 9D**). Significant causal connections ($p<0.05$, false discovery rate (FDR) cor-

rected) emerged from left and right thalami both in the young and old groups. However, the

strength of causal connections between thalamus and CEN nodes were greater in the younger

cohort (**figure 10A, 10D**)

ACC and bilateral RPFC, AInsula, SMG were defined as the nodes of SN in the Harvard Ox-

ford parcellation. As compared to DMN and CEN, multivariate granger causality analysis in

Salience network showed lesser number of directed causal influences. (**figure 9B, 9E**) Moreo-

ver, right thalamus causally influenced RPFCL in the younger cohort but not in the older cohort

(**figure 10B, 10E**).

For the Default mode network, the nodes selected were MPFC, bilateral aMTG, pMTG, to

MTG, AG and PCC. Multivariate Granger Causality was performed from the extracted time

series to evaluate within network reorganization in young and old cohort.

In the younger group, GCA revealed extensive causal connections between all the DMN nodes

(**figure 9F**). Fewer causal connections were observed within the DMN network in the older

cohort (**figure 9C**), suggesting an age-related decrease in causal drive. Presence of causal con-

nections of the DMN nodes with thalamus in the young (**figure 10F**) but not in the old cohort

reiterates the notion of thalamic decline with ageing.

In between network analysis, salience network emerged to have the highest causal strength,

exhibiting prominent causal influence on DMN and CEN in both young and old cohort. Net

causal outflows were significantly different for both age group.

4. Discussion

In the recent past, several studies have endeavoured to unravel the changes in the brain's structural and functional connectivity with aging and their cognitive implications. A vast majority of these studies have used temporal correlation across different brain regions as measure of functional connectivity to characterize ageing related changes in brain's large-scale functional connectivity patterns. In the present study, we employ multi variate granger causality analysis to probe within- and between- network causal relationships among three key intrinsic resting state brain networks with the hope to facilitate more biologically meaningful interpretations with healthy aging. Every cortical region receives feedforward projections from the thalamus and in turn sends outputs to one or multiple thalamic nuclei (Obeso et al.,2008);(McFarland et al.,2002). Thalamocortical projections relay nearly all incoming information to the cortex as well as mediate corticocortical communication (Sherman and Gui). Thus, a fuller insight into brain functional characterization certainly requires knowledge of the organization and properties of thalamocortical interactions. A recent study by ((Hwang, Bertolero, Liu, & D'Esposito, 2017)) showed thalamus as an integrative hub for functional networks. A handful of studies also observed disrupted thalamic resting state functional networks, in brain injury, schizophrenia((Tang et al., 2011);(Wang, Rau, Li, Chen, & Yu, 2015a)). (Goldstone, Mayhew, Hale, Wilson, & Bagshaw, 2018) investigated the association of thalamic functional connectivity with behavioural performance in elderly population. In continuation of earlier works on thalamic functional connectivity, we extend it to thalamic effective connectivity by not concentrating entirely on cortical nodes, and pinpointing how such interactions are driven with age by common exogeneous thalamocortical inputs during endogenous activities.

 We investigate the changes in the causal dynamics in RSNs with the inclusion of thalamus in our analysis. Overall, we found significant reconfiguration of between- and within-network

effective connectivity with aging. Our study also reveals the salience network's role as a me-

diator of switching between DMN and CEN with aging, a finding that is in line with the extant

literature ((Menon & Uddin, 2010); (Bonnelle et al., 2012)). Beyond this, we also find that the

role of salience network as a mediator is getting even more prominent with aging, leading to

reduced segregation and increased integration of resting state networks with aging. Finally, we

demonstrate that how thalamus plays a crucial role in restructuring cortical network causality

with aging.

According to the dedifferentiation hypothesis of cognitive aging, age-related impairments in

cognitive function arise from reduced distinctiveness of neural representations ((Li,

Lindenberger, & Sikström, 2001)). Historically, the concept of dedifferentiation was intro-

duced by Baltes and colleagues ((Baltes & et, 1980)) to account for age-related increases in the

correlation between levels of performance on different cognitive tasks. At the neural level,

numerous brain-imaging studies have shown that the aging brain adapts to the age-related

changes by exhibiting more global activation compared to younger individuals while perform-

ing a cognitive/motor task ((Cabeza, 2002); (Reuter-Lorenz, 2002);(Serrien, Ivry, & Swinnen,

2007); (Seidler et al., 2010)). In line with these finding, at the network level, several studies

have found a decrease in within-network functional connectivity and an increase in between-

network functional connectivity in RSNs ((Andrews-Hanna et al., 2007);(Betzel et al.,

2014);(Ferreira & Busatto, 2013);(Ferreira et al., 2016);(Geerligs et al., 2015);(Ng, Lo, Lim,

Chee, & Zhou, 2016)) with aging. Overall, our study reproduces these results with effective

connectivity analysis and find that the reorganization in causal connectivity patterns with age

primarily reflects functional connectivity patterns. More specifically, younger individuals

show increase in both the number and the strength of causal connections within DMN, CEN,

and SN. At the between-network level, we find, using principal component analysis, that causal

strengths are significantly higher in the elderly individuals compared to the young thus further

substantiating the dedifferentiation hypothesis. Moreover, the present study also uncovers sev-

eral novel observations, e.g., the reversal of direction of causal connections between rMOF and

rIPL with aging, age-related changes in the causal outflow in key nodes of the RSNs and re-

configuration of causal outflow hubs and so on that are inaccessible to a functional connectivity

analysis. Thus, future studies should continue to investigate brain networks across the lifespan

employing MVGC and other-directed connectivity measures.

The role of salience network in mediating switching between DMN and CEN is well estab-

lished ((Sridharan et al., 2008); (Menon & Uddin, 2010); (Goulden et al., 2014)). In agreement

with these observations, we find SN to exert strong causal influence on both DMN and CEN

in both the age groups. In between network analysis, SN is found to act as a causal outflow hub

among three RSNs we investigated. Interestingly, the causal influence of SN on both DMN

and CEN increases with aging. An increase in the between network causality in elderly group

emerges as a general trend in our analysis, and the above finding may reflect this general trend.

Thalamus has extensive connections with the entire cerebral cortex. By performing graph-the-

oretic analyses on thalamocortical functional connectivity data, ((Hwang et al., 2017)) demon-

strated that thalamus integrates multimodal information across diverse cortical functional net-

works and act as an integrative hub for functional brain networks. The association between

thalamocortical functional connectivity abnormalities and cognitive deficits in clinical condi-

tions like schizophrenia ((Wang, Rau, Li, Chen, & Yu, 2015b),) is already known. Also known

is the fact that thalamic volume significantly decreases with aging ((Walhovd et al.,

2005);(Cherubini, Péran, Caltagirone, Sabatini, & Spalletta, 2009);(Zheng et al., 2018)). How-

ever, only a handful of studies (e.g.(Goldstone et al., 2018)) have investigated the thalamic

influence in reorganizing resting state networks with aging and none has employed the causal

analysis. In our analysis with effective connectivity measures thalamus emerges as an im-

portant node to critically influence both within- and between- network connectivity patterns

among RSNs. After inclusion of thalamus in the between network analysis, thalamus acts as a

causal outflow hub and causally drives all three network nodes for both the age groups. In

contrast, in within network analysis, the influence of thalamus in reorganizing within network

causality is much more prominent in younger age group compared to the elderly. In DMN, Left

thalamus comes out as a causal outflow hub for young group while patterns in the net causal

outflows were unchanged in elderly group compared to the without thalamus analysis. After

inclusion of thalamus in CEN within network analysis, though left thalamus acts as a causal

outflow hub for both the groups, the causal outflow values are higher in young compared to

elderly group. This preferential influence of thalamus on younger individual is consistent with

the thalamic decline with aging as discussed above. Also, among three resting state networks,

the SN remains least affected after inclusion of thalamus, both in within- and between- network

analysis. No changes were found in causal architecture within SN after inclusion of thalamus

in elderly individuals. In young individuals also changes are minimal. The net causal outflow

pattern also does not change after inclusion of thalamus in both the groups. These findings are

in concurrence with the observations made by the previous studies ((Wang et al., 2015b);(Cao

et al., 2014); (Sakaki et al., 2016); (Xiao et al., 2018)) that in contrast to DMN and CEN, within

network connectivity is preserved or increased in SN with aging. Our result suggests that this

preservation of connectivity within SN may be driven by thalamus.

We have performed our replication analysis, with different cohort and different brain parcella-

tion atlas. Even though, different nodes were used for three RSN, some of the major findings

with the earlier dataset were replicated. In within network analysis, greater number of causal

connections with higher strengths in younger group compared to old individuals are consistent

with our findings with Berlin data set. Effect of thalamus is also revealed by higher number of

causal connections from thalamus to other nodes in younger group compared to old individuals.

We acknowledge several limitations of our study that should be addressed by the future studies.

The result should be replicated in a larger lifespan cohort comprise of middle young and middle

elderly groups to get additional insights and true estimate of lifespan trend in causal outflow in

resting state networks. Future research should also investigate the sensitivity of the analysis to

the choice of different brain parcellation schemes.

Finally, thalamus is a heterogenous structure composed of several nuclei; each of which sends

distinct afferent inputs to cortical regions as well as driven by cortical outputs (add reference).

Thus, probing the influence of different nuclei of thalamus on reorganization of within- and

between- network causality of different RSNs would help to better describe the complex neu-

rophysiological processes taking place in the brain with aging. In general, the analysis could

be extended to other subcortical regions to understand how cortical-subcortical connectivity

impact the cognitive ability across age.

In conclusion, the results of the present study demonstrate that effective connectivity analysis

can provide crucial insights regarding within- and between- network information flow across

lifespan over and above insights provided by functional connectivity measures. This study also

establishes thalamus as a common driver of organization and reorganization of RSNs with ag-

ing, a conclusion that should encourage future research to explore the influences exerted by

subcortical structures on cortical networks and their clinical implications.

Acknowledgement

This study was supported by NBRC Core funds, Ramalingaswami Fellowships (Department of Biotechnology, Government of India) to DR (BT/RLF/Re-entry/07/2014) and AB (BT/RLF/Re601entry/31/2011) and Innovative Young Biotechnologist Award (IYBA) to AB (BT/07/IYBA/2013).AB also acknowledges the support of the Centre of Excellence in Epilepsy and MEG. DR was also supported by SR/CSRI/21/2016 extramural grant from the Department of Science and Technology (DST) Ministry of Science and Technology, Government of India. Data collection and sharing for this project was provided by the Cambridge Centre for Ageing and Neuroscience (CamCAN). CamCAN funding was provided by the UK Biotechnology and Biological Sciences Research Council (grant number BB/H008217/1), together with support from the UK Medical Research Council and University of Cambridge, UK.

References

Abi-Dargham, A., & Horga, G. (2016). The search for imaging biomarkers in psychiatric disorders. *Nature Medicine*, *22*(11), 1248–1255. https://doi.org/10.1038/nm.4190

Andrews-Hanna, J. R., Reidler, J. S., Sepulcre, J., Poulin, R., & Buckner, R. L. (2010). Functional-Anatomic Fractionation of the Brain's Default Network. *Neuron*, *65*(4), 550–562. https://doi.org/10.1016/j.neuron.2010.02.005

Andrews-Hanna, J. R., Snyder, A. Z., Vincent, J. L., Lustig, C., Head, D., Raichle, M. E., & Buckner, R. L. (2007). Disruption of large-scale brain systems in advanced aging. *Neuron*, *56*(5), 924–935. https://doi.org/10.1016/j.neuron.2007.10.038

Baltes, P. B., & et, al. (1980). Integration versus differentiation of fluid/crytallized intelligence in old age. *Developmental Psychology*, *16*(6), 625–635. https://doi.org/10.1037/0012-1649.16.6.625

Barrett, A. B., Barnett, L., & Seth, A. K. (2010). Multivariate Granger causality and generalized variance. *Physical Review E - Statistical, Nonlinear, and Soft Matter Physics*, *81*(4). https://doi.org/10.1103/PhysRevE.81.041907

Beckmann, C. F., & Smith, S. M. (2004). Probabilistic Independent Component Analysis for

Functional Magnetic Resonance Imaging. *IEEE Transactions on Medical Imaging*, *23*(2), 137–152. https://doi.org/10.1109/TMI.2003.822821

Betzel, R. F., Byrge, L., He, Y., Goñi, J., Zuo, X. N., & Sporns, O. (2014). Changes in structural and functional connectivity among resting-state networks across the human lifespan. *NeuroImage*, *102*(P2), 345–357. https://doi.org/10.1016/j.neuroimage.2014.07.067

Bonnelle, V., Ham, T. E., Leech, R., Kinnunen, K. M., Mehta, M. A., Greenwood, R. J., & Sharp, D. J. (2012). Salience network integrity predicts default mode network function after traumatic brain injury. *Proceedings of the National Academy of Sciences*, *109*(12), 4690–4695. https://doi.org/10.1073/pnas.1113455109

Bressler, S. L., & Menon, V. (2010, June). Large-scale brain networks in cognition: emerging methods and principles. *Trends in Cognitive Sciences*, Vol. 14, pp. 277–290. https://doi.org/10.1016/j.tics.2010.04.004

Buckner, R. L., Andrews-Hanna, J. R., & Schacter, D. L. (2008). The brain's default network: Anatomy, function, and relevance to disease. *Annals of the New York Academy of Sciences*, *1124*, 1–38. https://doi.org/10.1196/annals.1440.011

Buckner, R. L., Krienen, F. M., Castellanos, A., Diaz, J. C., & Yeo, B. T. T. (2011). The organization of the human cerebellum estimated by intrinsic functional connectivity. *Journal of Neurophysiology*, *106*(5), 2322–2345. https://doi.org/10.1152/jn.00339.2011

Cabeza, R. (2002). Hemispheric asymmetry reduction in older adults: The HAROLD model. *Psychology and Aging*, *17*(1), 85–100. https://doi.org/10.1037/0882-7974.17.1.85

Cao, W., Luo, C., Zhu, B., Zhang, D., Dong, L., Gong, J., … Yao, D. (2014). Resting-state functional connectivity in anterior cingulate cortex in normal aging. *Frontiers in Aging Neuroscience*, *6*, 280. https://doi.org/10.3389/fnagi.2014.00280

Cherubini, A., Péran, P., Caltagirone, C., Sabatini, U., & Spalletta, G. (2009). Aging of

subcortical nuclei: Microstructural, mineralization and atrophy modifications measured

in vivo using MRI. *NeuroImage*, *48*(1), 29–36.

https://doi.org/10.1016/j.neuroimage.2009.06.035

Corbetta, M., & Shulman, G. L. (2002). Control of goal-directed and stimulus-driven

attention in the brain. *Nature Reviews Neuroscience*, *3*(3), 201–215.

https://doi.org/10.1038/nrn755

Deco, G., Jirsa, V. K., & McIntosh, A. R. (2011). Emerging concepts for the dynamical

organization of resting-state activity in the brain. *Nature Reviews. Neuroscience*, *12*(1),

43–56. https://doi.org/10.1038/nrn2961

Desikan, R. S., Ségonne, F., Fischl, B., Quinn, B. T., Dickerson, B. C., Blacker, D., …

Killiany, R. J. (2006). An automated labeling system for subdividing the human cerebral

cortex on MRI scans into gyral based regions of interest. *NeuroImage*, *31*(3), 968–980.

https://doi.org/10.1016/j.neuroimage.2006.01.021

Dixon, M. L., Andrews-Hanna, J. R., Spreng, R. N., Irving, Z. C., Mills, C., Girn, M., &

Christoff, K. (2017). Interactions between the default network and dorsal attention

network vary across default subsystems, time, and cognitive states. *NeuroImage*, *147*,

632–649. https://doi.org/10.1016/j.neuroimage.2016.12.073

Ferreira, L. K., & Busatto, G. F. (2013). Resting-state functional connectivity in normal brain

aging. *Neuroscience and Biobehavioral Reviews*, *37*(3), 384–400.

https://doi.org/10.1016/j.neubiorev.2013.01.017

Ferreira, L. K., Regina, A. C. B., Kovacevic, N., Martin, M. D. G. M., Santos, P. P., Carneiro,

C. D. G., … Busatto, G. F. (2016). Aging effects on whole-brain functional connectivity

in adults free of cognitive and psychiatric disorders. *Cerebral Cortex*, *26*(9), 3851–3865.

https://doi.org/10.1093/cercor/bhv190

Fox, M. D., Corbetta, M., Snyder, A. Z., Vincent, J. L., & Raichle, M. E. (2006).

Spontaneous neuronal activity distinguishes human dorsal and ventral attention systems. *Proceedings of the National Academy of Sciences of the United States of America*, *103*(26), 10046–10051. https://doi.org/10.1073/pnas.0604187103

Fransson, P. (2005). Spontaneous low-frequency BOLD signal fluctuations: An fMRI investigation of the resting-state default mode of brain function hypothesis. *Human Brain Mapping*, *26*(1), 15–29. https://doi.org/10.1002/hbm.20113

Geerligs, L., Rubinov, M., Tyler, L. K., Brayne, C., Bullmore, E. T., Calder, A. C., … Henson, R. N. (2015). State and trait components of functional connectivity: Individual differences vary with mental state. *Journal of Neuroscience*, *35*(41), 13949–13961. https://doi.org/10.1523/JNEUROSCI.1324-15.2015

Goldstone, A., Mayhew, S. D., Hale, J. R., Wilson, R. S., & Bagshaw, A. P. (2018). Thalamic functional connectivity and its association with behavioral performance in older age. *Brain and Behavior*, *8*(4). https://doi.org/10.1002/brb3.943

Goulden, N., Khusnulina, A., Davis, N. J., Bracewell, R. M., Bokde, A. L., McNulty, J. P., & Mullins, P. G. (2014). The salience network is responsible for switching between the default mode network and the central executive network: replication from DCM. *NeuroImage*, *99*, 180–190. https://doi.org/10.1016/j.neuroimage.2014.05.052

Hwang, K., Bertolero, M. A., Liu, W. B., & D'Esposito, M. (2017). The Human Thalamus Is an Integrative Hub for Functional Brain Networks. *The Journal of Neuroscience*, *37*(23), 5594–5607. https://doi.org/10.1523/jneurosci.0067-17.2017

Li, S. C., Lindenberger, U., & Sikström, S. (2001, November 1). Aging cognition: From neuromodulation to representation. *Trends in Cognitive Sciences*, Vol. 5, pp. 479–486. https://doi.org/10.1016/S1364-6613(00)01769-1

Llinás, R. R., Ribary, U., Jeanmonod, D., Kronberg, E., & Mitra, P. P. (1999). Thalamocortical dysrhythmia: A neurological and neuropsychiatric syndrome

characterized by magnetoencephalography. *Proceedings of the National Academy of*

Sciences of the United States of America, *96*(26), 15222–15227.

https://doi.org/10.1073/pnas.96.26.15222

McFarland, N. R., & Haber, S. N. (2002). Thalamic relay nuclei of the basal ganglia form

both reciprocal and nonreciprocal cortical connections, linking multiple frontal cortical

areas. *Journal of Neuroscience*, *22*(18), 8117-8132.

Menon, V., & Uddin, L. Q. (2010). Saliency, switching, attention and control: a network

model of insula function. *Brain Structure & Function*, Vol. 214, pp. 655–667.

https://doi.org/10.1007/s00429-010-0262-0

Ng, K. K., Lo, J. C., Lim, J. K. W., Chee, M. W. L., & Zhou, J. (2016). Reduced functional

segregation between the default mode network and the executive control network in

healthy older adults: A longitudinal study. *NeuroImage*, *133*, 321–330.

https://doi.org/10.1016/j.neuroimage.2016.03.029

Obeso, J. A., Rodríguez-Oroz, M. C., Benitez-Temino, B., Blesa, F. J., Guridi, J., Marin, C.,

& Rodriguez, M. (2008). Functional organization of the basal ganglia: therapeutic impli-

cations for Parkinson's disease. *Movement disorders: official journal of the Movement*

Disorder Society, *23*(S3), S548-S559.

Raichle, M. E. (2015). The Brain's Default Mode Network. *Annual Review of Neuroscience*,

38(1), 433–447. https://doi.org/10.1146/annurev-neuro-071013-014030

Reuter-Lorenz, P. A. (2002). New visions of the aging mind and brain für mehr Artikel.

Trends in Cognitive Sciences, *6*(9), 394–400. https://doi.org/10.1016/S1364-

6613(02)01957-5

Sakaki, M., Yoo, H. J., Nga, L., Lee, T. H., Thayer, J. F., & Mather, M. (2016). Heart rate

variability is associated with amygdala functional connectivity with MPFC across

younger and older adults. *NeuroImage*, *139*, 44–52.

https://doi.org/10.1016/j.neuroimage.2016.05.076

Schirner, M., Rothmeier, S., Jirsa, V. K., McIntosh, A. R., & Ritter, P. (2015). An automated

pipeline for constructing personalized virtual brains from multimodal neuroimaging

data. *NeuroImage*, *117*, 343–357. https://doi.org/10.1016/j.neuroimage.2015.03.055

Seidler, R. D., Bernard, J. A., Burutolu, T. B., Fling, B. W., Gordon, M. T., Gwin, J. T., …

Lipps, D. B. (2010). Motor control and aging: links to age-related brain structural,

functional, and biochemical effects. *Neuroscience and Biobehavioral Reviews*, *34*(5),

721–733. https://doi.org/10.1016/j.neubiorev.2009.10.005

Serrien, D. J., Ivry, R. B., & Swinnen, S. P. (2007). The missing link between action and

cognition. *Progress in Neurobiology*, *82*(2), 95–107.

https://doi.org/10.1016/j.pneurobio.2007.02.003

Smith, S. M., Fox, P. T., Miller, K. L., Glahn, D. C., Fox, P. M., Mackay, C. E., …

Beckmann, C. F. (2009). Correspondence of the brain's functional architecture during

activation and rest. *Proceedings of the National Academy of Sciences of the United*

States of America, *106*(31), 13040–13045. https://doi.org/10.1073/pnas.0905267106

Sridharan, D., Levitin, D. J., & Menon, V. (2008). A critical role for the right fronto-insular

cortex in switching between central-executive and default-mode networks. *Proceedings*

of the National Academy of Sciences of the United States of America, *105*(34), 12569–

12574. https://doi.org/10.1073/pnas.0800005105

Tang, L., Ge, Y., Sodickson, D. K., Miles, L., Zhou, Y., Reaume, J., & Grossman, R. I.

(2011). Thalamic resting-state functional networks: Disruption in patients with mild

traumatic brain injury. *Radiology*, *260*(3), 831–840.

https://doi.org/10.1148/radiol.11110014

Uddin, L. Q. (2015). Salience processing and insular cortical function and dysfunction.

Nature Reviews Neuroscience, *16*(1), 55–61. https://doi.org/10.1038/nrn3857

Uddin, L. Q., Kelly, A. M. C., Biswal, B. B., Castellanos, F. X., & Milham, M. P. (2009).

Functional Connectivity of Default Mode Network Components: Correlation,

Anticorrelation, and Causality. *Human Brain Mapping*, *30*(2), 625–637.

https://doi.org/10.1002/hbm.20531

Vanneste, S., Song, J. J., & De Ridder, D. (2018). Thalamocortical dysrhythmia detected by

machine learning. *Nature Communications*, *9*(1). https://doi.org/10.1038/s41467-018-

02820-0

Walhovd, K. B., Fjell, A. M., Reinvang, I., Lundervold, A., Dale, A. M., Eilertsen, D. E., …

Fischl, B. (2005). Effects of age on volumes of cortex, white matter and subcortical

structures. *Neurobiology of Aging*, *26*(9), 1261–1270; discussion 1275-8.

https://doi.org/10.1016/j.neurobiolaging.2005.05.020

Wang, H. L. S., Rau, C. L., Li, Y. M., Chen, Y. P., & Yu, R. (2015a). Disrupted thalamic

resting-state functional networks in schizophrenia. *Frontiers in Behavioral*

Neuroscience, *9*(FEB), 1–9. https://doi.org/10.3389/fnbeh.2015.00045

Wang, H. L. S., Rau, C. L., Li, Y. M., Chen, Y. P., & Yu, R. (2015b). Disrupted thalamic

resting-state functional networks in schizophrenia. *Frontiers in Behavioral*

Neuroscience, *9*(FEB). https://doi.org/10.3389/fnbeh.2015.00045

Woodward, N. D., & Cascio, C. J. (2015). Resting-State Functional Connectivity in

Psychiatric Disorders. *JAMA Psychiatry*, *72*(8), 743–744.

https://doi.org/10.1001/jamapsychiatry.2015.0484

Xiao, T., Zhang, S., Lee, L. E., Chao, H. H., van Dyck, C., & Li, C. S. R. (2018). Exploring

age-related changes in resting state functional connectivity of the Amygdala: From

young to middle adulthood. *Frontiers in Aging Neuroscience*, *10*(JUL).

https://doi.org/10.3389/fnagi.2018.00209

Zheng, F., Cui, D., Zhang, L., Zhang, S., Zhao,

Y., Liu, X., … Qiu, J. (2018). The volume of hippocampal subfields in relation to decline of

memory recall across the adult lifespan. *Frontiers in Aging Neuroscience*, *10*(OCT).

https://doi.org/10.3389/fnagi.2018.00320

729   **Table 1**

730 **Coordinates of selected nodes of three resting state networks according to**
731 **Desikan-Killiany parcellation atlas**

Network	Region	MNI coordinates (x,y,z)	
		Left (l)	Right(r)
Salience Network	Insula (Insula)	(-41, 13, -6)	(43,12,-6)
	Caudal anterior cingulate cortex (CACC)	(-2,21,27)	(3, 21, 27)
	Rostral anterior cingulate cortex (RACC)	(-2,39,6)	(4, 38,4)
Central Executive Network	Rostral middle frontal gyrus (RMFG)	(-34, 53, 17)	(43, 45, 21)
	Caudal middle frontal gyrus (CMFG)	(-45, 18, 46)	(43, 14, 43)
	Superior parietal lobule (SPL)	(-25, -62, 63)	(17,-65,59)
Default Mode Network	Medial orbitofrontal (MOF)	(-4, 44, -14)	(7, 45, -13)
	Inferior parietal lobule (IPL)	(-47, -70, 31)	(48, -67,29)
	Posterior cingulate cortex (PCC)	(-1, -18, 38)	(1, -16, 37)

Figure 1. Flow chart describing major steps employed in our pipeline for estimation of causal

connectivity and weighted outflow analysis using MVGC

Figure 2. ROIs selected based on seven networks resting state parcellation proposed

by Yeo et al. (2011) (cite). The ROIs (circles) related 3 prominent brain networks are

overlaid on the spatial distribution maps derived from group ICA of multiple resting

state networks of interest that is, the default mode network ROIs (DMN), the

salience network ROIs (SN), and the Control network or Central Executive Network

(CEN)

Figure 3. Effective Connectivity between the nodes of three resting state networks. Effective

connectivity between **A.** six key nodes of CEN (green), **B.** six key nodes of SN (yellow),

C. six key nodes of DMN (red) for elderly group. **D.** six key nodes of CEN (green), **E.** six

key nodes of SN (yellow), **F.** six key nodes of DMN (red) for young group.

Figure 4.   Weighted net causal flow for within network effective connectivity analysis. A.

Weighted net causal outflow in nodes of the central executive network **B.** Weighted Net causal

outflow in nodes of the salience network **C.** Weighted net causal outflow in nodes of default

mode network. Weighted net causal outflows were significantly different in few nodes in each

of the three RSN young and elderly group ($p < 0.05$ is indicated by '*', $p<0.01$ is indicated

by '**', No significant difference is indicated by 'NS')

Figure 5.   Effective Connectivity between the nodes of three resting state networks and

the thalamus. **A.** Effective connectivity between six key nodes of CEN (green), **B.** six key

nodes of SN (yellow), **C.** six key nodes of DMN (red) for elderly group in presence of thalamic

nodes (blue). **D.** Effective connectivity between six key nodes of CEN (green), **E.** six key

nodes of SN (yellow), **F.** six key nodes of DMN (red) for elderly group in presence of thalamic

nodes (blue) for young group in presence of thalamic nodes (blue).

Figure 6. Weighted net causal outflow for within network effective connectivity analysis

A. Weighted net causal outflow in nodes of the central executive network and two thalamic

regions (left, right). **B.** Weighted net causal outflow in nodes of the salience network and two

thalamic regions (left,right). **C.** Weighted Net causal outflow for default mode network and

two thalamic regions (left, right). Weighted net causal outflows were significantly different in

few nodes in each of the three RSN young and elderly group ($p < 0.05$ is indicated by '*',

$p<0.01$ is indicated by '**', No significant difference is indicated by 'NS') .

Figure 7. Effective connectivity between three resting state networks in absence and presence

of the thalamus. **A.** Effective connectivity between three nodes representing three RSN, SN

(yellow), DMN (red), CEN (green) for old population. **B.** Results for Effective connectivity

between three nodes representing three RSN for young population **C.** Effective connectivity

between three nodes representing three RSN and fourth node representing the thalamus (blue)

for old population **D.** Effective connectivity between three nodes representing three RSN and

fourth node representing the thalamus for young population

Figure 8. Weighted net causal outflow for between network effective connectivity analysis

A. Weighted net causal outflow in three RSN **B.** Weighted net causal outflow in three RSN

and the thalamus. Weighted net causal outflows were significantly different in few nodes in

each of the three RSN young and elderly group ($p<0.01$ is indicated by '**', No significant

difference is indicated by 'NS') .

Figure 9. Effective Connectivity between the nodes of three resting state networks using Harvard Oxford Atlas. Effective connectivity between **A.** key nodes of CEN (green), **B.** key nodes of SN (yellow), **C.** key nodes of DMN (red) for elderly group. **D.** key nodes of CEN (green), **E.** key nodes of SN (yellow), **F.** key nodes of DMN (red) for young group.

Figure 10. Effective Connectivity between the nodes of three resting state networks and the thalamus using Harvard Oxford Atlas. Effective connectivity between **A**. key nodes of CEN (green), **B**. key nodes of SN (yellow), **C.** key nodes of DMN (red) for elderly group in presence of thalamic nodes (blue). **D.** Effective connectivity between key nodes of CEN (green), **E.** key nodes of SN (yellow), **f.** key nodes of DMN (red) for young group in presence of thalamic nodes (blue).

Figure 1.

Control   Default Mode   Dorsal Attention   Salience

Limbic   Somatomotor   Visual

Figure 2.

845   **Figure 3.**

Figure 4.

866 **Figure 5.**

877 **Figure 6.**

A. Between RSN's connectivity for old population

B. Between RSN's connectivity for young population

C. Between RSN's and Thalamus's connectivity for old population

D. Between RSN's and Thalamus's connectivity for young population

888   **Figure 7.**

Figure 8.

Figure 9.

942
943
944
945   **Figure 10.**