Preparatory delta phase response is correlated with naturalistic speech comprehension performance

Jiawei Li[1, 3], Bo Hong[2, 3], Guido Nolte[4], Andreas K. Engel[4], Dan Zhang[1, 3*]

[1]Department of Psychology, School of Social Sciences, Tsinghua University, Beijing, China

[2]Department of Biomedical Engineering, School of Medicine, Tsinghua University, Beijing, China

[3]Tsinghua Laboratory of Brain and Intelligence, Tsinghua University, Beijing, China

[4]Department of Neurophysiology and Pathophysiology, University Medical Center Hamburg Eppendorf, Hamburg, Germany

[*]**Correspondence**:

Dan Zhang, Ph.D.

Room 334, Mingzhai Building, Tsinghua University, Beijing 100084, China

E-mail: dzhang@tsinghua.edu.cn; Tel: +86-10-62796737

Abstract

While human speech comprehension is thought to be an active process that involves top-down predictions, it remains unclear how predictive information is used to prepare for the processing of upcoming speech information. We aimed to identify the neural signatures of the preparatory processing of upcoming speech. Participants selectively attended to one of two competing naturalistic, narrative speech streams, and a temporal response function (TRF) method was applied to derive event-related-like neural responses from electroencephalographic data. The phase responses to the attended speech at the delta band (1–4 Hz) were correlated with the comprehension performance of individual participants, with a latency of -200–0 ms before onset over the fronto-central and left-lateralized parietal regions. The phase responses to the attended speech at the alpha band also correlated with comprehension performance, but with a latency of 650–980 ms post-onset over fronto-central regions. Distinct neural signatures were found for the attentional modulation, taking the form of TRF-based amplitude responses at a latency of 240–320 ms post-onset over the left-lateralized fronto-central and occipital regions. Our findings reveal how the brain gets prepared to process an upcoming speech in a continuous, naturalistic speech context.

Keywords

preparatory processing, attention, speech comprehension, electroencephalogram, temporal response function

1. Introduction

Humans can effectively comprehend complex and rapidly changing speech in challenging conditions, e.g., in a cocktail party scenario with multiple competing speech streams and high background noise. To achieve such a capacity, the human brain is equipped with an efficient neural architecture that is dedicated to bottom-up processing of perceived speech information, from the low-level acoustics, to the phoneme, syllable, and sentence levels (Pisoni and Luce 1987; DeWitt and Rauschecker 2012; Friederici 2012; Hickok 2012; Verhulst et al. 2018). In recent years, increasing evidence has also suggested that human speech comprehension is an active process that involves top-down predictions (Rao and Ballard 1999; Federmeier 2007; Arnal et al. 2011; Hickok et al. 2011; Kutas and Federmeier 2011; Fries 2015; Tian et al. 2018). In the cocktail party scenario, it is believed that a listener should continuously predict what their attended speaker is going to say next in order to efficiently understand the corresponding speech (Cherry 1953; Ding and Simon 2012; Zion Golumbic, Cogan, et al. 2013; O'Sullivan et al. 2015; Bednar and Lalor 2020).

Although the idea of prediction in human speech comprehension is gaining popularity, it remains unclear how the brain gets prepared for the processing of upcoming speech information. The available findings on prediction in speech, however, are not sufficient to determine the neural mechanisms underlying preparation. For instance, the classic studies of active speech prediction have mainly focused on neural activity in response to prediction errors. Event-related potential (ERP) components such as the N400 and P600 are frequently reported when the perceived word violates semantic and syntactic

congruency of the preceding speech context, respectively (Lau et al. 2008; Kutas and

Federmeier 2011; Van Petten and Luka 2012; Wang et al. 2018). These ERP

components normally occur >400 ms after the presentation of the perceived speech,

and so provide only indirect support for the preparatory process.

Recent studies have reported evidence of the brain's pre-activation before the onset of

the upcoming speech. Some studies have focused on preparatory attentional orientation

to specific acoustic features (e.g., spatial location, pitch, etc.) in speech or general

auditory tasks (Hill and Miller 2010; Lee et al. 2013; Holmes et al. 2016, 2018; ElShafei

et al. 2018; Nolden et al. 2019). It has been demonstrated that an external cue (e.g., a

visual symbol) prior to the onset of the auditory or speech stimuli could elicit distinct

cue-specific neural responses, depending on the cue-instructed attentional task. In the

meanwhile, researchers have reported pre-activations that are more specific for speech

processing (DeLong et al. 2005; Dikker and Pylkkänen 2013; Söderström et al. 2016,

2018): event-related neural responses to a preceding speech unit (e.g., words) were

found to be informative about possible upcoming speech units in the continuous speech

materials (e.g., sentences). While these speech-related pre-activations could reflect the

brain's preparation for processing the upcoming speech, they were represented by

event-related responses to either attention-related cues or preceding speech units.

Therefore, the possible findings are dependent upon the nature of these preceding

events, which are substantially different from the upcoming, to-be-processed speech

unit. More importantly, it could be further argued that these pre-activations might only

reflect a neural processing phase prior to the actual preparatory speech processing, and

there should be a more direct preparatory phase during the processing of the upcoming

speech. Ideally, the most direct preparatory processing should be linked to the to-be-

processed speech but occur before its onset.

While this direct evidence has not been investigated in the speech domain, studies on

general sensory processing have provided support for the existence of such a

preparatory process. In the visual domain, the amplitude and phase of pre-stimulus

oscillatory activities, especially in the alpha band, have been reported to have a

significant impact on subsequent perceptual consequences, (Van Dijk et al. 2008; Kok

et al. 2017; Harris et al. 2018; Galindo-Leon et al. 2019; Rassi et al. 2019), such as

threshold-level perception, attention orientation, visual search performance, etc.

Auditory information processing has also been shown to be affected by pre-stimulus

oscillations, mainly at lower frequency bands such as theta and delta (Ng et al. 2012;

Kayser et al. 2016; Zoefel et al. 2018). However, these studies have mainly employed

simple and abstract sensory stimuli, therefore limited in explaining possible neural

mechanisms underlying the preparatory processing of human speech. Recently, there is

an emergence of naturalistic stimuli in auditory studies (Sonkusare et al. 2019).

Researchers adopted naturalistic, continuous stimuli (e.g., poetry, long sentences) as

the stimulus (Etard and Reichenbach 2019; Donhauser and Baillet 2020; Teng et al.

2020). Naturalistic speech presents listeners with a variety and multitude of different

linguistic contents (Alexandrou et al. 2020), catering for different levels of preparation.

Thus, our study utilized naturalistic and continuous materials to study the preparatory

processing.

One crucial issue that needs to be considered is the possible dependence of the

preparatory process on top-down selective attention. As attention regulates the

processing of the input sensory information, it can be expected to affect prediction and

consequently preparation (Schröger, Kotz, et al. 2015; Schröger, Marzecová, et al.

2015). A number of recent studies have indeed reported attention-dependent neural

responses to prediction errors (Kok et al. 2012; Auksztulewicz and Friston 2015; Hisagi

et al. 2015; Marzecová et al. 2017; Smout et al. 2019), with ongoing debates on the

direction of the interplay and the involved sensory processing stages. Most of these

studies have been conducted within the visual domain, with limited exploration in the

auditory domain, let alone speech processing.

The present study aimed to identify neural signatures that directly reflect the

preparatory processing of human speech. Naturalistic, narrative speech materials were

presented auditorily to the participants with a 60-channel electroencephalogram (EEG)

recording; this procedure is believed to be of high ecological validity and thus to

provide necessary contextual information for the engagement of top-down prediction

and therefore preparation (Rao and Ballard 1999; Friston 2005; Federmeier 2007; Jehee

and Ballard 2009). A cocktail party paradigm was used to introduce a complex

perceptual environment that imposed further demands on prediction and preparation

(Cherry 1953; Ding and Simon 2012; Zion Golumbic, Cogan, et al. 2013; O'Sullivan

et al. 2015; Broderick et al. 2018). To characterize the neural responses to the

continuous, naturalistic speech streams, a temporal response function (TRF) method

was used to derive event-related-like neural responses from the EEG signals, based on

the speech power envelop of both the attended and the unattended speech streams

(Lalor et al. 2006; Crosse et al. 2016). The TRF-based responses revealed the temporal

dynamics of neural activities underlying human speech processing, and the responses

with latencies earlier than the onset of speech power envelop fluctuations are regarded

to represent the preparatory phase. Specifically, the TRF-based responses were further

decomposed into amplitude and phase responses, as amplitude and phase have been

proposed to play unique roles in networks underlying human cognition (Bonnefond and

Jensen 2012; Engel et al. 2013; Fries 2015). Following the studies on the perceptual

influence of pre-stimulus neural activities (Smith et al. 2006; Iemi et al. 2019; Rassi et

al. 2019; Avramiea et al. 2020), we were interested in whether the TRF-based responses

at the preparatory stage could be correlated to speech comprehension performance, as

measured by speech-content-related questionnaires, and how amplitude and phase

responses contributed to speech preparation. With the employment of the cocktail party

paradigm, we also addressed the issue about the attention-dependency of the to-be-

explored performance-related preparatory activities. Our study is expected to reveal the

neural mechanisms underlying how the brain gets prepared to process an upcoming

speech in a continuous, naturalistic speech context.

2.   Materials and Methods

2.1.   Ethics statement

The study was conducted in accordance with the Declaration of Helsinki and was

approved by the local Ethics Committee of Tsinghua University. Written informed

consent was obtained from all participants.

2.2. Experimental model and participant details

Twenty college students (10 females; mean age: 24.7 years; range: 20–43 years) from

Tsinghua University participated in the study as paid volunteers. The sample size (N =

20) was decided following previous TRF-based studies on human speech processing

(Di Liberto et al. 2015; Mirkovic et al. 2015; Broderick et al. 2018). All participants

were native Chinese speakers, reported having normal hearing, and normal or

corrected-to-normal vision.

2.3. Stimuli

The speech stimuli were recorded from two male speakers using the microphone of an

iPad2 mini (Apple Inc., Cupertino, CA) at a sampling rate of 44,100 Hz. The speakers

were college students from Tsinghua University, who had more than four years of

professional training in broadcasting. Both speakers were required to tell 28 1-min

narrative stories in Mandarin Chinese; the stories were either those about daily-life

topics recommended by the experimenter and told by the speaker improvising on their

own (14 stories), or those selected from the National Mandarin Proficiency Test (14

stories). The speakers were presented with the recommended topic or story materials

on the computer screen. They were allowed to prepare for as long as required before

telling the story (usually ~3 min). When they were ready, the speakers pressed the

SPACE key on the computer keyboard and the recording began with the presentation

of three consecutive pure-tone beep sounds at 1000 Hz (duration: 1000 ms; inter-beep

interval: 1500 ms). The beep sounds served as the event marker to synchronize the

speech audios in the main experiment, in which two speech streams were presented

simultaneously. The speakers were asked to start speaking as soon as the third beep had

ended (within around 3 sec). The speakers were allowed to start the recording again if

the audio did not meet the requirements of either the experimenter or the speakers

themselves (which mainly concerned speech coherence). The actual speaking time per

story ranged from 51 to 76 sec.

Two four-choice questions per story were then prepared by the experimenter and two

college students who were familiar with comprehension performance assessment.

These questions and the corresponding choices concerned story details that required

significant attentional efforts. For instance, one question following a story about one's

hometown was, "What is the most dissatisfying thing about the speaker's hometown?

(推测讲述人对于家乡最不满意的地方在于?)", and the four choices were A) There is no

heating in winter; B) There are no hot springs in summer; C) There is no fruit in autumn;

D) There are no flowers in spring (A. 冬天没暖气；B. 夏天没温泉；C. 秋天没水果；D.

春天没鲜花). Both the speech audio and corresponding questions are available for

downloads.

2.4. Experimental procedure

The main experiment consisted of 4 blocks of 7 trials per block. In each trial, two

narrative stories were presented simultaneously, one to the left ear and the other to the

right ear. The participants were instructed to attend to one spatial side. The two speech

streams within each trial were from the two different male speakers to facilitate

selective attention. Considering the possible duration difference between the two audio

streams, the trial ended after the longer speech audio had ended. Each trial began when

participants pressed the SPACE key on the computer keyboard. Participants were

instructed which side to attend to by plain text ("Please pay attention to the

[LEFT/RIGHT]") displayed on the computer screen. A white fixation cross was also

displayed throughout the trial. The speech stimuli were played immediately after the

keypress, and were preceded by the three beep sounds to allow participants to prepare.

At the end of each trial, four questions (two for each story) were presented sequentially

in a random order on the computer screen, and the participants made their choices using

the computer keyboard. After completing these questions, participants scored their

attention level of the attended stream, the experienced difficulty of performing the

attention task, and the familiarity with the attended material using three 10-point Likert

scales. No feedback was given to the participants about their performance during the

experiment. Throughout the trial, participants were required to maintain visual fixation

on the fixation cross while listening to the speech and to minimize eye blinks and all

other motor activity. The participants were recommended to take a short break (of

around 1 min) after every trial within one block, and a long break (no longer than 10

min) between blocks.

The to-be-attended side was fixed within each block (two blocks for attending to the

left side and two for attending to the right side). Within each block, the speaker identity

remained unchanged for the left and right sides. In this way, the to-be-attended spatial

side and the corresponding speaker identity were balanced within the participant, with

seven trials per side for both speakers. The assignment of the stories to the four blocks

was randomized across the participants.

The experiment was carried out in a sound-attenuated, dimly lit, and electrically

shielded room. The participants were seated in a comfortable chair in front of a 19.7-

inch Lenovo LT2013s Wide LCD monitor. The viewing distance was approximately 60

cm. The experimental procedure was programmed in MATLAB using the

Psychophysics Toolbox 3.0 extensions (Brainard and Brainard 1997). The speech

stimuli were delivered binaurally via an air-tube earphone (Etymotic ER2, Etymotic

Research, Elk Grove Village, IL, USA) to avoid possible electromagnetic interferences

from auditory devices. The volume of the audio stimuli was adjusted to be at a

comfortable level that was well above the auditory threshold. Furthermore, the speech

stimuli driving the earphone were used as an analog input to the EEG amplifier through

one of its bipolar inputs together with the EEG recordings. In this way, the audio and

the EEG recordings were precisely synchronized, with a maximal delay of 1ms (at a

sampling rate of 1000 Hz).

Figure 1. The analysis workflow.

(A) The experimental paradigm. Participants attended to one of two simultaneously presented naturalistic, narrative speech streams while 60-channel EEG was recorded. (B) EEG data analysis. Neural responses were characterized using a TRF-based modeling method. The TRF-based neural responses were decomposed into the amplitude and the phase responses using the Hilbert transform. This procedure was conducted separately for attended (Att-) and unattended (Unatt-, not shown) speech streams, and separately for EEG data filtered at delta, theta, alpha and beta bands. (C) Comprehension performance. The participants completed a comprehension task after each speech comprehension trial. The average response accuracy over all trials per participant was taken as his/her comprehension performance. (D) Correlation analysis for comprehension performance-related neural responses. We calculated the correlation between either amplitude or phase responses and comprehension performance for each channel-latency bin. We defined neural activity before 0 ms as preparatory-processing and activity after 0 ms as post-processing. The results of the delta band were illustrated here. The colored channel-latency bins showed uncorrected significant correlation with comprehension performance. The dashed box in the 'Corr-Phase' plot indicates a significant channel-latency cluster by a cluster-based permutation test.

2.5. Data acquisition and pre-processing

EEG was recorded from 60 electrodes (FP1/2, FPZ, AF3/4, F7/8, F5/6, F3/4, F1/2, FZ,

FT7/8, FC5/6, FC3/4, FC1/2, FCZ, T7/8, C5/6, C3/4, C1/2, CZ, TP7/8, CP5/6, CP3/4,

CP1/2, CPZ, P7/8, P5/6, P3/4, P1/2, PZ, PO7/8, PO5/6, PO3/4, POZ, Oz, and O1/2),

which were referenced to an electrode between Cz and CPz, with a forehead ground at

Fz. A NeuroScan amplifier (SynAmp II, NeuroScan, Compumedics, USA) was used to

record EEG at a sampling rate of 1000 Hz. Electrode impedances were kept below 10

kOhm for all electrodes.

The recorded EEG data were first notch filtered to remove the 50 Hz powerline noise

and then subjected to an artifact rejection procedure using independent component

analysis. Independent components (ICs) with large weights over the frontal or temporal

areas, together with a corresponding temporal course showing eye movement or muscle

movement activities, were removed. The remaining ICs were then back-projected onto

the scalp EEG channels, reconstructing the artifact-free EEG signals. While the

relatively long duration of the speech trials in the present study (~1 minute per story,

see Experimental procedure) has made it more difficult for the participants to avoid

inducing movement-related artifacts as compared to the classical ERP-based studies, a

temporally continuous, non-interrupted EEG segment per trial was preferred for the

employment of the TRF method. Therefore, any ICs with artifact-like EEG activities

for more than 20% of the trial time (i.e. ~12 sec) were rejected, leading to around 4–11

ICs rejected per participant. The cleaned EEG data was used for the TRF analysis

without any further artifact rejection procedures. Then the EEG signals were re-

referenced to a common average reference.

Next, the EEG data were segmented into 28 trials according to the markers representing

speech onsets. The analysis window for each trial extended from 10 to 55 s (duration:

45 s) to avoid the onset and the offset of the stories.

2.6. Temporal response function modeling

The analysis workflow is shown in Figure 1. The neural responses to the speech stimuli

were characterized using a temporal response function (TRF)-based modeling method.

The TRF response describes the impulse response to fluctuations of an input signal, and

is based on system identification theories (Lalor et al. 2006; Crosse et al. 2016). We

used the power envelope of the speech signal as the input signal required by TRF, which

has been demonstrated to be a valid index by which to extract speech-related neural

responses (Ding and Simon 2012; Mirkovic et al. 2015; O'Sullivan et al. 2015; Bednar

and Lalor 2018; Broderick et al. 2018; Huang et al. 2018). TRF-based responses before

the onset of the speech power envelope fluctuations are considered to represent

preparatory activity, whereas post-onset responses reflect post-processing of the speech

stream.

Prior to the modeling, the preprocessed EEG signals were re-referenced to the average

of all scalp channels and then downsampled to 128 Hz. Then, the EEG data were filtered

in delta (1–4 Hz), theta (4–8 Hz), alpha (8–12 Hz) and beta (12–30 Hz) bands (filter

order: 64, one-pass forward filter). The use of a causal FIR filter ensured that filtered

EEG signals were decided only by the current and previous data samples (de Cheveigné

and Nelken 2019), which is important for the present research aim of preparatory

speech processing. The filter order of 64 was chosen to keep a balance of temporal

resolution and filter performance: the filtered EEG signals were therefore calculated

based on the preceding 500 ms data (64 at 128 Hz).

The power envelopes of the speech signals were obtained using a Hilbert transform and

then downsampled to the same sampling rate of 128 Hz. When denoting $R_{i,k}^{j}(t)$ as

the downsampled EEG signals from channel i, trial k filtered at one specific frequency

band j (representing the four frequency bands) and $S_k(t)$ as the input speech power

envelope corresponding to trial k, the corresponding neural response $TRF_{i,k}^{j}(t)$ can be

formulated as follows:

$$R_{i,k}^{j}(t) = TRF_{i,k}^{j}(t) * S_k(t) \tag{1}$$

Where * represents the convolution operator. The latency in the neural response models

$TRF_{i,k}^{j}(t)$ was set to vary from -1000 ms to 1000 ms post-stimulus. To control for

overfitting, we varied the lambda from 10^{-1} to 10^3 (lambda $= 10^{-1}, 10^0, \ldots , 10^3$) in the

ridge regression (Di Liberto et al. 2015; Crosse et al. 2016; Broderick et al. 2018). The

lambda value corresponding to the backward decoder that produced the highest cross-

validated speech envelope reconstruction accuracy, averaged across trials, was selected

as the regularization parameter for all trials per participant (Broderick et al. 2019). All

TRF-based responses were further transformed into z-scores within the -1000 ms to

1000 ms time window for each channel separately per participant to account for across

participants and across session differences (Pasley et al. 2012; Kleen et al. 2016). These

z-scores were then used for the following analyses.

TRF models were calculated for attended and unattended speech processing separately

using the corresponding speech streams as the input signal. TRF models were obtained

for each EEG channel per participant, at the four frequency bands. It should be noted

that we did not consider the input lateralization for the TRF models, as the observed

behaviorally related findings were insensitive to the physical origin of the speech audios,

but rather likely to reflect the lateralization of the human speech network. Fig. S1

provides the topographical information of our main results; it is similar to Fig. 2, but

all results were calculated separately for speech stimuli from the left and right sides.

The topographies were comparable.

Amplitude and phase were calculated using the Hilbert transform of the TRF-based

neural responses at the single-trial level. Hereby, the instantaneous amplitude $A_{i,k}^{j}(t)$

and the instantaneous phase $\phi_{i,k}^{j}(t)$ can be computed as

$$A_{i,k}^{j}(t) = \sqrt{\left(TRF_{i,k}^{j}(t)\right)^2 + h\left(TRF_{i,k}^{j}(t)\right)^2} \tag{2}$$

$$\phi_{i,k}^{j}(t) = \tan^{-1}\left(\frac{h\left(TRF_{i,k}^{j}(t)\right)}{TRF_{i,k}^{j}(t)}\right) \tag{3}$$

where $h(TRF_{i,k}^{j}(t))$ represents its Hilbert transform.

These single-trial instantaneous amplitude and phase were then averaged across all

trials per participants to reflect one's overall EEG responses to the naturalistic speech

streams. Specifically, the single-trial amplitudes were averaged by taking their

arithmetic mean value, whereas the single-trial phase responses were averaged by

computing their circular mean (i.e., the mean phase angle).

In addition, the inter-trial phase locking (ITPL) was also calculated to evaluate the

phase consistency across trials within each participant, as follows:

$$ITPL(t) = \left| \sum_{k} exp\left(TRF_{i,j}^{k}(t)\right) \right| \tag{4}$$

The phase-related ITPL value varies between 0 and 1; 0 refers to a situation in which

the phase responses of different trials are uniformly distributed between 0 and 2π, and

1 means the phase responses from all trials are entirely locked to a fixed phase angle.

The TRF analysis was conducted in MATLAB using the Multivariate Temporal

Response Function (mTRF) toolbox (Crosse et al. 2016). All the other EEG processing

procedures, as well as the statistical analyses, were conducted using the FieldTrip

toolbox (Oostenveld et al. 2011).

2.7.   Quantification and statistical analysis

The extracted TRF-based amplitude and phase responses were used to correlate with

the speech comprehension performance of the attended speech at the participant level.

The Spearman's correlation was calculated between the amplitude response and the

comprehension performance at each EEG channel and each individual latency across

the participants. The correlations between the phase response and the comprehension

performance were evaluated by computing the circular linear correlation using the

CircStat toolbox (Berens 2009). Both the TRFs to the attended and unattended speech

at the four frequency bands were included for this analysis.

Statistical analysis was performed to examine the significance of correlations over all

channel-latency bins by computing the correlation *r*-values. To account for multiple

comparisons, a nonparametric cluster-based permutation analysis was applied (Maris

and Oostenveld 2007). In this procedure, neighboring channel-latency bins with an

uncorrected correlational p-value below 0.01 were combined into clusters, for which

the sum of the correlational t-statistics corresponding to the correlation r-values were

obtained. A null-distribution was created through permutations of data across

participants ($n = 1,000$ permutations), which defines the maximum cluster-level test

statistics and corrected p-values for each cluster.

To investigate the attention modulation effect, we performed paired t-tests contrasting

the TRFs to the attended speech versus the unattended speech. Both amplitude and

phase were included in the analysis. The phase difference was calculated as the phase

angle difference by the CircStat toolbox as well. A similar cluster-based permutation

was used to control for the multiple comparison problem ($p < .01$ as the threshold, $n =$

1, 000 permutations).

2.8.   Data and code availability statement

Dataset   generated   in   our   study   has   been   uploaded   to   OSF

(https://osf.io/87srv/?view_only=13f01e1f1f7b4cf98555ffacd878a53b). We have also

provided data of individual participant, including the original EEG data and behavioral

performance data.

Where existing toolboxes were used for data analysis, citations have been provided.

3.   Results

3.1. Behavioral results

The average comprehension performance was significantly better for the 28 attended stories than for the 28 unattended stories (67.0±2.5% (standard error) vs. 36.0±1.6%, $t(19) = 10.948$, $p < .001$; the four-choice chance level: 25%). The participants reported a moderate level of attention (8.146±0.343 on a 10-point Likert scale) and attention difficulties (2.039±0.530 on a 10-point Likert scale). The accuracy for the attended story was significantly correlated with both the self-reported attention level ($r = .476$, $p = .043$) and attention difficulty ($r = -.677$, $p = .001$). The self-reported story familiarity level was low for all the participants (0.860±0.220 on a 10-point Likert scale) and was not correlated with comprehension performance ($r = -.224$, $p = .342$). These results suggest that participants' selective attention was effectively manipulated, as well as good reliability of the measured comprehension performance. Most importantly, there was a large inter-individual difference in the participant-wise average comprehension performance for the attended stories; the response accuracy varied from 48.2% to 91.1%, which supports the feasibility of using these accuracy values as a behavioral indicator of comprehension-relevant neural signatures.

3.2. Speech comprehension performance related TRF-based responses

The nonparametric cluster-based permutation analysis revealed a significant correlation between the multi-channel amplitude and phase representation of the TRFs and individual speech comprehension performance of the attended speech. This corresponded to two clusters in the observed data (cluster-based permutation $p < .05$).

Both the two significant clusters are reflected by TRF-based phase responses to

attended speech, one at the delta band and the other at the alpha band (Figure 2A and

2B). No significant correlations were found for the TRFs to the unattended speech.

As shown in Figure 2C.I and II, the delta cluster is represented by TRF-based phase

responses to attended speech at -200–0 ms relative to the onset of speech power

envelope fluctuations over the fronto-central and left lateralized parietal regions

(cluster-based permutation p = .012, mean circular linear correlation r = .787). The

participants with their comprehension performance ranking in top, middle, and bottom

tertiles were associated with different delta phase angles during this pre-speech-onset

time window (average Φ_{top} = 73.411°, Φ_{middle} = 80.175°, and Φ_{bottom} = -134.767°,

respectively, see Figure 2C.III). Specifically, the top-performing participants showed a

negative peak within the time window of the cluster, whereas the bottom-performing

participants showed a positive peak during this time. However, the participants' ITPL

did not significantly correlate with their comprehension performance (r = .127, p = .

593, Figure 2C.IV).

The alpha cluster was represented by TRF-based phase responses to attended speech at

650–980 ms post speech onset, with a fronto-central distribution (cluster-based

permutation p = .031, mean r = .784, see Figure 2D.I and II). The individual difference

in the phase angle of their responses was large, with average Φ_{top} = 41.904° for the top-

performing participants and Φ_{middle} = -135.461°, Φ_{bottom} = -115.916° for the other two

groups (Figure 2D.III). Again, there was no significant correlation between the ITPL

and the comprehension performance (r = .267, p = .255, Figure 2D.IV).

Figure 2. Speech comprehension performance related TRF-based responses to naturalistic speech.

(A) Distributions of the cluster-level correlational t-statistics (log-transformation) from the 1, 000 permutated calculations of the correlation between the comprehension performance and the TRF-based amplitude response at the four frequency bands. The upper and lower panel for the attended and unattended responses, respectively. Vertical lines indicate the t-statistics of the clusters from the real data. N.C. means no cluster was formed and N.D. means the permutated distribution could not be generated (i.e., no cluster formed during the permutation calculation). (B) Distributions of the cluster-level correlational t-statistics from the 1, 000 permutated calculations of the correlation between the comprehension performance and the TRF-based phase response at the four frequency bands. The upper and lower panel for the attended and unattended responses, respectively. The two vertical lines with asterisks indicate statistically significant clusters from the real data. (C) Illustration of the significant phase-response cluster at

the delta band. I. The time course of the TRFs at one representative channel (FC1). The
three waveforms represent the average responses over the participants with
comprehension performance of the attended speech ranking in the top (red), middle
(orange), and bottom (yellow) tertiles (7, 6, and 7 participants, respectively). The
shaded region depicts the time window of interest with a significant correlation. II.
Topography of the average correlation *r*-values in the time window of interest. Black
dots indicate the channels of interest in the cluster. III. The polar plot showing the
average response at channel FC1 per participant. The vector lengths indicate the
response amplitude and the vector directions indicate the phase angle. The three
different colors show the comprehension performance rankings as in I. IV. The scatter
plot showing the participants' comprehension performance (accuracy in percentage)
versus their inter-trial phase-locking (ITPL) values. The color of the dots indicates the
comprehension performance rankings as in I and III. 'n.s.' means no significant
correlation. (D) Illustration of the significant phase-response cluster at the alpha band.
The plots in I and III are from channel FC1. The explanations of the sub-plots follow
(C).

3.3. Attention related TRF-based responses

The nonparametric cluster-based permutation analysis revealed a significant difference

between the TRFs to the attended and unattended speech. The difference was

manifested as a cluster involving a set of channels covering the left lateralized fronto-

central and occipital regions (cluster-based permutation $p < .001$) at a latency of 240–

320 ms post the onset of speech power envelope fluctuations (Figure 3C). This

attentional difference is reflected on TRF-based amplitude response at the theta band.

No significant differences were observed in amplitude responses at other frequency

bands and phase responses at all frequency bands (Figure 3A and 3B).

Figure 3. Attention related TRF-based responses to naturalistic speech.

(A) Distributions of the cluster-level t-statistics (log-transformation) from the 1, 000 permutated calculations of the paired comparisons between the TRF-based attended and unattended amplitude responses at the four frequency bands. Vertical lines indicate the max cluster-level t-statistics in real data and the asterisk shows the statistically significant cluster from the real data. N.C. means no cluster was formed. (B) Distributions of the cluster-level t-statistics from the 1, 000 permutated calculations of the paired comparisons between the TRF-based attended and unattended phase responses at the four frequency bands. (C) Illustration of the significant amplitude-response cluster at the theta band. I. The time course of the TRFs at one representative channel (FC1). The two waveforms represent the average responses to the attended (dark blue) and unattended (light blue) speech. The shaded region depicts the time window of interest with significant differences between the two waveforms. II. Topography of the average amplitude response differences in the time window of interest. Black dots indicate the channels of interest in the cluster. III. The amplitude response difference per participant.

4. Discussion

The present study aimed to identify neural signatures that directly reflect the preparatory processing of upcoming speech. We used naturalistic narrative speech materials in a selective attention paradigm and a TRF-based approach for modeling the

neural activity and observed preparatory neural activities before the onset of speech

power envelope fluctuations. We found the phase responses to the attended speech at

the delta band (1–4 Hz) were correlated with the comprehension performance of

individual participants, with a latency of -200–0 ms before onset over the fronto-central

and left lateralized parietal regions. The phase responses to the attended speech at the

alpha band also correlated with comprehension performance, but with a latency of 650–

980 ms post onset over fronto-central regions. Distinct neural signatures were found for

the attentional modulation, taking the form of TRF-based amplitude responses at a

latency of 240–320 ms post onset over the left lateralized fronto-central and occipital

regions. Our results provide direct neural evidence for how the brain prepares for the

processing of upcoming speech.

Before detailed discussions, it is necessary to state that our assumption for a preparatory

process is based on the observation that the TRF-based neural activities before the onset

of speech power envelop fluctuations during the continuous speech stream were

significantly correlated with comprehension performance. Recent TRF-based studies

using naturalistic stimuli have reported reasonable latencies that resembled their ERP

counterparts for describing selective auditory attention (~200 ms) (Mirkovic et al. 2015;

O'Sullivan et al. 2015), semantic violation processing (~400 ms) (Broderick et al. 2018),

and visual working memory (200–400 ms) (Huang et al. 2018). Although our findings

have mainly focused on the window of < 0 ms, these studies support the rationale of

using the TRF-based responses to reflect the time course of information processing in

general. In addition, the present study also replicated the timing of previously reported

attentional modulation of TRF-based responses to speech stimuli (Mirkovic et al. 2015;

O'Sullivan et al. 2015). Therefore, the pre-onset latencies observed in the present study

can be considered to represent a preparatory state that precedes speech processing.

Moreover, the TRF method has enabled us to investigate the neural dynamics in an

event-related-like manner but with naturalistic speech materials that are expected to

better resemble real-world speech comprehension tasks.

Our results highlight in particular that the delta band phase at -200–0ms before the

speech onset determines the comprehension accuracy of the listeners, which serves for

the preparation of up comping speech information. As the analysis was performed on

the neural responses to the to-be-processed speech at the pre-onset stage rather than

those related to the preceding speech-related information (i.e., attentional cue,

preceding words), our study has focused on a distinct processing phase other than

previous studies that have mainly explored either the preparatory attention orientation

and the speech-specific pre-activations by the preceding speech unit (DeLong et al.

2005; Söderström et al. 2016, 2018). Notably, the employment of the naturalistic speech

materials has ensured sufficient variations of the interval between the preceding and the

to-be-processed speech unit. Therefore, the present results are not likely to reflect the

neural responses to the preceding speech unit. Although there was one study that has

reported a sustained difference in the neural activities at around 400 ms prior to the

onset of the upcoming speech (Lee et al. 2013), their 'pre-onset' activities mainly

reflected the preparatory attention orientation by a visually-presented abstract

attentional cue. In contrast, the results of the present study are expected to reflect how

the human brain makes use of the rich contextual information in the naturalistic speech

materials to infer and prepare for the upcoming speech information.

The timing and the spatial distribution of the reported pre-onset preparatory activity is

better compared with the studies on pre-stimulus oscillatory activities. Visual pre-

stimulus studies have generally reported perceptual-relevant neural activities with

timings of 100–400 ms prior to stimulus onset (Harris et al. 2018; Wöstmann et al.

2019). In our research, we have found a similar time window for those studies. Our

results, therefore, suggest that approximately -200–0 ms prior to stimulus onset might

have a general implication for the preparation of sensory information processing. As

the present study employed naturalistic (speech) materials that are expected to provide

much richer contextual information as compared to the simple and abstract stimuli in

most of the previous studies, our results might provide more reliable support for the

observed timing on preparatory sensory processing. The result is also in line with a

recent study, which indicated that the delta band entrainment at 100 ms before the

stimulus is related to the noise-induced comprehension difference (Etard and

Reichenbach 2019). Meanwhile, the central-parietal regions found in our study

involved in the preparatory processing are also consistent with the previous auditory

pre-stimulus studies (Stefanics et al. 2010; Kayser et al. 2016). The central-parietal

responses could be related to the predictive processing of speech meaning and could

recruit a mechanism that is similar to that underlying the classical central-parietal N400

response (Federmeier 2007; Lau et al. 2008; Szewczyk and Schriefers 2018) or possibly

preparatory attentional orientation as well (Holmes et al. 2016). The left lateralized

parietal regions could also be linked to the speech-specific processing, e.g. the

Wernicke's area (Hickok and Poeppel 2007).

Our study expands on findings from the studies about the functional role of the delta

band in speech preparatory processing. The delta band could help the segmentation or

identification of intonation phrases, which is essential for the preparation and prediction

of upcoming speech (Giraud and Poeppel 2012; Ding et al. 2015; Kösem et al. 2018;

Meyer 2018; Morillon et al. 2019). In particular, the phase of the delta band before the

stimulus is related to the hit rate afterward (Ng et al. 2012), or the behavioral

consequence (i.e., the reaction time) in the auditory studies using simple, isolated

stimulus (Lakatos et al. 2008; Stefanics et al. 2010; Henry and Obleser 2012) . It is in

contrast to visual studies in which the alpha band is mostly pronounced in the pre-

stimulus stage (Busch et al. 2009; Mathewson et al. 2011; Milton and Pleydell-Pearce

2016), suggesting active auditory perception is dominated by lower-frequency (Ng et

al. 2012; VanRullen 2016). In addition, the brain activity dynamically tracks speech

streams using the delta phase (Zion Golumbic, Cogan, et al. 2013) and temporal

predictions are encoded by delta neural oscillations (Morillon and Baillet 2017;

Auksztulewicz et al. 2018). Our findings are in line with the previous studies about the

delta band phase's role in preparatory processing and unambiguously show that the

comprehension performance could be elevated when the delta is better prepared at a

particular phase angle.

The neural mechanisms of the preparatory process were investigated using correlation

analysis of the TRF-based neural activity. In line with recent TRF-based studies, we

observed attention-related neural responses (Mirkovic et al. 2015, 2016; O'Sullivan et

al. 2015), with the peak attention effect represented by theta amplitude activities at 250–

320 ms post-stimulus onset over the central and occipital regions. In contrast, the

comprehension-related post-onset neural signatures were in the alpha band at 670–980

ms. The result could be interpreted for a functional role of alpha-band for a top-down

control mechanism to achieve the preparatory process, as the phase of alpha oscillations

has an active role in attentional temporal predictions (Händel et al. 2011; Bonnefond

and Jensen 2012; Samaha et al. 2015). Meanwhile, in auditory studies, the alpha

frequency band has also been suggested to be associated with working memory

(Bonnefond and Jensen 2012; Meyer 2018), capable of storing semantics in sentences

(Haarmann and Cameron 2005) and syntax information (Bonhage et al. 2017) and

lexical decision (Strauß et al. 2015), which are also closely related to speech

comprehension.

The neural mechanisms of the preparatory process were further explored by inspecting

their relationship with attention. While our results are in line with previous research

that has reported low-frequency phase track the envelope of attended speech (Ding and

Simon 2012; Zion Golumbic, Ding, et al. 2013), we provide further evidence on how

such interactions could affect behavior (i.e., comprehension). Indeed, we only found

the correlation between the attended TRF-based neural activities and the

comprehension performance in the pre-onset stage, suggesting a possible attentional

facilitation of preparation of speech processing.

This study has some limitations that should be noted. While the use of naturalistic

speech materials could better resemble the real-world speech comprehension scenarios,

it is difficult to infer which specific types of information (e.g. timing, phoneme, etc.)

were the main contributor for the observed preparatory activities. The distributed brain

regions involved in the preparatory process may provide a guidance for designing

further experiments to have an in-depth exploration. The present study used the speech

power envelope as the reference signal from which the TRF models were derived,

which could reflect the speech information at all linguistic levels due to the highly

redundant information shared across levels (Di Liberto et al. 2015; Daube et al. 2019).

While such an operation has the advantage of providing a general overview about

preparatory processing, further investigations are necessary to differentiate possible

contributions at different linguistic levels (Di Liberto et al. 2015; Broderick et al. 2018).

Meanwhile, caution must be taken when interpreting the timing of the preparatory

activities. While the preparatory activity as early as ~200 ms before speech onset could

be the result of an optimized utilization of the rich contextual information provided by

the naturalistic speech materials, such timings may be dependent upon the materials per

se. Further studies are necessary to investigate the possible material dependence of

these timings, for instance, by employed an extended amount of speech materials. Also,

denser MEG recordings together with source localization methods are expected to more

precisely identify the brain regions for preparatory speech processing (Mazaheri et al.

2009; Nolte and Müller 2010). Besides, an inter-individual level regression analysis

method was chosen, as the average comprehension questionnaire accuracies across all

stories within each participant were believed to provide a more reliable estimation of

the speech comprehension performance than the single-trial accuracies. Thus, our results do not necessarily imply that the observed neural signatures reflect the participants' trait-like, stable speech processing style. Alternatively, it could be more plausible to consider these neural signatures to reflect a more or less efficient speech processing state. More theoretical and empirical research is needed to clarify the underlying mechanisms.

Supporting information

Figure S1. Topographies of the two responses as shown in Figure 2 (A) and calculated separately for the speech stimuli delivered to the left side (B) and the right side (C) only.

Acknowledgments

This work was supported by the National Natural Science Foundation of China (NSFC)

and the German Research Foundation (DFG) in project Crossmodal Learning (grant

number: NSFC 61621136008/DFG TRR-169/C1, B1), the National Key Research and

Development Plan (grant number: 2016YFB1001200), the National Natural Science

Foundation of China (grant number: 61977041 and U1736220), the National Social

Science Foundation of China (grant number: 17ZDA323) and Tsinghua University

Initiative Scientific Research Program (grant number: 20197010006).

The authors would like to thank Prof. Dr. Xiaoqin Wang and Dr. Yue Ding for providing

the shielded room for the experiment as well as necessary technical support.

Declaration of interests

The authors declare no competing interests.

References

Alexandrou AM, Saarinen T, Kujala J, Salmelin R. 2020. Cortical entrainment: what

we can learn from studying naturalistic speech perception. Lang Cogn Neurosci.

35:681–693.

Arnal LH, Wyart V, Giraud AL. 2011. Transitions in neural oscillations reflect

prediction errors generated in audiovisual speech. Nat Neurosci. 14:797–801.

Auksztulewicz R, Friston K. 2015. Attentional enhancement of auditory mismatch

responses: A DCM/MEG study. Cereb Cortex. 25:4273–4283.

Auksztulewicz R, Schwiedrzik CM, Thesen T, Doyle W, Devinsky O, Nobre AC,

Schroeder CE, Friston KJ, Melloni L. 2018. Not all predictions are equal: "what"

644 and "when" predictions modulate activity in auditory cortex through different

645 mechanisms. J Neurosci. 38:8680–8693.

646 Avramiea AE, Hardstone R, Lueckmann JM, Bim J, Mansvelder HD, Linkenkaer-

647 Hansen K. 2020. Pre-stimulus phase and amplitude regulation of phase-locked

648 responses is maximized in the critical state. Elife. 9:1–17.

649 Bednar A, Lalor EC. 2018. Neural tracking of auditory motion is reflected by delta

650 phase and alpha power of EEG. Neuroimage. 181:683–691.

651 Bednar A, Lalor EC. 2020. Where is the cocktail party? Decoding locations of

652 attended and unattended moving sound sources using EEG. Neuroimage.

653 205:116283.

654 Berens P. 2009. CircStat : A MATLAB Toolbox for Circular Statistics. J Stat Softw.

655 31:293–295.

656 Bonhage CE, Meyer L, Gruber T, Friederici AD, Mueller JL. 2017. Oscillatory EEG

657 dynamics underlying automatic chunking during sentence processing.

658 Neuroimage. 152:647–657.

659 Bonnefond M, Jensen O. 2012. Alpha oscillations serve to protect working memory

660 maintenance against anticipated distracters. Curr Biol. 22:1969–1974.

661 Brainard DH, Brainard DH. 1997. The Psychophysics Toolbox. In: Spatial vision. p.

662 433–436.

663 Broderick MP, Anderson AJ, Di Liberto GM, Crosse MJ, Lalor EC. 2018.

Electrophysiological Correlates of Semantic Dissimilarity Reflect the

Comprehension of Natural, Narrative Speech. Curr Biol. 28:803-809.e3.

Broderick MP, Anderson AJ, Lalor EC. 2019. Semantic Context Enhances the Early

Auditory Encoding of Natural Speech. J Neurosci. 39:7564–7575.

Busch NA, Dubois J, VanRullen R. 2009. The phase of ongoing EEG oscillations

predicts visual perception. J Neurosci. 29:7869–7876.

Cherry EC. 1953. Some experiments on the recognition of speech, with one and with

two ears. J Acoust Soc Am. 25:975–979.

Crosse MJ, Di Liberto GM, Bednar A, Lalor EC. 2016. The Multivariate Temporal

Response Function (mTRF) Toolbox: A MATLAB Toolbox for Relating Neural

Signals to Continuous Stimuli. Front Hum Neurosci. 10:604.

Daube C, Ince RAA, Gross J. 2019. Simple Acoustic Features Can Explain Phoneme-

Based Predictions of Cortical Responses to Speech. Curr Biol. 29:1924-1937.e9.

de Cheveigné A, Nelken I. 2019. Filters: When, Why, and How (Not) to Use Them.

Neuron. 102: 280–293.

DeLong KA, Urbach TP, Kutas M. 2005. Probabilistic word pre-activation during

language comprehension inferred from electrical brain activity. Nat Neurosci.

8:1117–1121.

DeWitt I, Rauschecker JP. 2012. Phoneme and word recognition in the auditory

ventral stream. Proc Natl Acad Sci U S A. 109:505–514.

Di Liberto GM, O'Sullivan JA, Lalor EC. 2015. Low-frequency cortical entrainment

to speech reflects phoneme-level processing. Curr Biol. 25:2457–2465.

Dikker S, Pylkkänen L. 2013. Predicting language: MEG evidence for lexical

preactivation. Brain Lang. 127:55–64.

Ding N, Melloni L, Zhang H, Tian X, Poeppel D. 2015. Cortical tracking of

hierarchical linguistic structures in connected speech. Nat Neurosci. 19:158–164.

Ding N, Simon JZ. 2012. Emergence of neural encoding of auditory objects while

listening to competing speakers. Proc Natl Acad Sci U S A. 109:11854–11859.

Donhauser PW, Baillet S. 2020. Two Distinct Neural Timescales for Predictive

Speech Processing. Neuron. 105:385-393.e9.

ElShafei HA, Bouet R, Bertrand O, Bidet-Caulet A. 2018. Two Sides of the Same

Coin: Distinct Sub-Bands in the α Rhythm Reflect Facilitation and Suppression

Mechanisms during Auditory Anticipatory Attention. eneuro. 5:1–14.

Engel AK, Gerloff C, Hilgetag CC, Nolte G. 2013. Intrinsic Coupling Modes:

Multiscale Interactions in Ongoing Brain Activity. Neuron. 80:867–886.

Etard O, Reichenbach T. 2019. Neural Speech Tracking in the Theta and in the Delta

Frequency Band Differentially Encode Clarity and Comprehension of Speech in

Noise. J Neurosci. 39:5750–5759.

Federmeier KD. 2007. Thinking ahead: The role and roots of prediction in language

comprehension. Psychophysiology. 44:491–505.

Friederici AD. 2012. The cortical language circuit: From auditory perception to

sentence comprehension. Trends Cogn Sci. 16:262–268.

Fries P. 2015. Rhythms for Cognition: Communication through Coherence. Neuron.

88:220–235.

Friston K. 2005. A theory of cortical responses. Philos Trans R Soc B Biol Sci.

360:815–836.

Galindo-Leon EE, Stitt I, Pieper F, Stieglitz T, Engler G, Engel AK. 2019. Context-

specific modulation of intrinsic coupling modes shapes multisensory processing.

Sci Adv. 5:1–13.

Giraud AL, Poeppel D. 2012. Cortical oscillations and speech processing: Emerging

computational principles and operations. Nat Neurosci. 15:511–517.

Haarmann HJ, Cameron KA. 2005. Active maintenance of sentence meaning in

working memory : Evidence from EEG coherences. 57:115–128.

Händel BF, Haarmeier T, Jensen O. 2011. Alpha oscillations correlate with the

successful inhibition of unattended stimuli. J Cogn Neurosci. 23:2494–2502.

Harris AM, Dux PE, Mattingley JB. 2018. Detecting Unattended Stimuli Depends on

the Phase of Prestimulus Neural Oscillations. J Neurosci. 38:3092–3101.

Henry MJ, Obleser J. 2012. Frequency modulation entrains slow neural oscillations

and optimizes human listening behavior. Proc Natl Acad Sci. 109:20095–20100.

Hickok G. 2012. Computational neuroanatomy of speech production. Nat Rev

Neurosci. 13:135–145.

Hickok G, Houde J, Rong F. 2011. Sensorimotor Integration in Speech Processing:

Computational Basis and Neural Organization. Neuron. 69:407–422.

Hickok G, Poeppel D. 2007. The cortical organization of speech processing. Nat Rev

Neurosci. 8:393–402.

Hill KT, Miller LM. 2010. Auditory attentional control and selection during cocktail

party listening. Cereb Cortex. 20:583–590.

Hisagi M, Shafer VL, Strange W, Sussman ES. 2015. Neural measures of a Japanese

consonant length discrimination by Japanese and American English listeners:

Effects of attention. Brain Res. 1626:218–231.

Holmes E, Kitterick PT, Summerfield AQ. 2016. EEG activity evoked in preparation

for multi-talker listening by adults and children. Hear Res. 336:83–100.

Holmes E, Kitterick PT, Summerfield AQ. 2018. Cueing listeners to attend to a target

talker progressively improves word report as the duration of the cue-target

interval lengthens to 2,000 ms. Attention, Perception, Psychophys. 80:1520–

1538.

Huang Q, Jia J, Han Q, Luo H. 2018. Fast-backward replay of sequentially

memorized items in humans. Elife. 7:e35164.

Iemi L, Busch NA, Laudini A, Haegens S, Samaha J, Villringer A, Nikulin V V.

2019. Multiple mechanisms link prestimulus neural oscillations to sensory

responses. Elife. 8:e43620.

Jehee JFM, Ballard DH. 2009. Predictive Feedback Can Account for Biphasic

Responses in the Lateral Geniculate Nucleus. PLoS Comput Biol. 5:e1000373.

Kayser SJ, McNair SW, Kayser C. 2016. Prestimulus influences on auditory

perception from sensory representations and decision processes. Proc Natl Acad

Sci. 113:4842–4847.

Kleen JK, Testorf ME, Roberts DW, Scott RC, Jobst BJ, Holmes GL, Lenck-Santini

PP. 2016. Oscillation phase locking and late ERP components of intracranial

hippocampal recordings correlate to patient performance in a working memory

task. Front Hum Neurosci. 10:1–14.

Kok P, Jehee JFM, de Lange FP. 2012. Less Is More: Expectation Sharpens

Representations in the Primary Visual Cortex. Neuron. 75:265–270.

Kok P, Mostert P, De Lange FP. 2017. Prior expectations induce prestimulus sensory

templates. Proc Natl Acad Sci U S A. 114:10473–10478.

Kösem A, Bosker HR, Takashima A, Meyer A, Jensen O, Hagoort P. 2018. Neural

Entrainment Determines the Words We Hear. Curr Biol. 28:2867-2875.e3.

Kutas M, Federmeier KD. 2011. Thirty Years and Counting: Finding Meaning in the

N400 Component of the Event-Related Brain Potential (ERP). Annu Rev

Psychol. 62:621–647.

Lakatos P, Karmos G, Mehta AD, Ulbert I, Schroeder CE. 2008. Entrainment of

Neuronal Oscillations as a Mechanism of Attentional Selection. Science.

320:110–113.

Lalor EC, Pearlmutter BA, Reilly RB, McDarby G, Foxe JJ. 2006. The VESPA: A

method for the rapid estimation of a visual evoked potential. Neuroimage.

32:1549–1561.

Lau EF, Phillips C, Poeppel D. 2008. A cortical network for semantics:

(de)constructing the N400. Nat Rev Neurosci. 9:920–933.

Lee AKC, Rajaram S, Xia J, Bharadwaj H, Larson E, Hämäläinen MS, Shinn-

Cunningham BG. 2013. Auditory selective attention reveals preparatory activity

in different cortical regions for selection based on source location and source

pitch. Front Neurosci. 6:1–9.

Maris E, Oostenveld R. 2007. Nonparametric statistical testing of EEG- and MEG-

data. J Neurosci Methods. 164:177–190.

Marzecová A, Widmann A, SanMiguel I, Kotz SA, Schröger E. 2017. Interrelation of

attention and prediction in visual processing: Effects of task-relevance and

stimulus probability. Biol Psychol. 125:76–90.

Mathewson KE, Lleras A, Beck DM, Fabiani M, Ro T, Gratton G. 2011. Pulsed out

of awareness: EEG alpha oscillations represent a pulsed-inhibition of ongoing

cortical processing. Front Psychol. 2:1–15.

Mazaheri A, Nieuwenhuis ILC, Van Dijk H, Jensen O. 2009. Prestimulus alpha and

mu activity predicts failure to inhibit motor responses. Hum Brain Mapp.

30:1791–1800.

Meyer L. 2018. The neural oscillations of speech processing and language

comprehension: state of the art and emerging mechanisms. Eur J Neurosci.

48:2609–2621.

Milton A, Pleydell-Pearce CW. 2016. The phase of pre-stimulus alpha oscillations

influences the visual perception of stimulus timing. Neuroimage. 133:53–61.

Mirkovic B, Bleichner MG, De Vos M, Debener S. 2016. Target speaker detection

with concealed EEG around the ear. Front Neurosci. 10:1–11.

Mirkovic B, Debener S, Jaeger M, Vos M De. 2015. Decoding the attended speech

stream with multi-channel EEG: implications for online, daily-life applications. J

Neural Eng. 12:046007.

Morillon B, Arnal LH, Schroeder CE, Keitel A. 2019. Prominence of delta oscillatory

rhythms in the motor cortex and their relevance for auditory and speech

perception. Neurosci Biobehav Rev. 107:136–142.

Morillon B, Baillet S. 2017. Motor origin of temporal predictions in auditory

attention. Proc Natl Acad Sci. 114:E8913–E8921.

Ng BSW, Schroeder T, Kayser C. 2012. A Precluding But Not Ensuring Role of

Entrained Low-Frequency Oscillations for Auditory Perception. J Neurosci.

32:12268–12276.

Nolden S, Ibrahim CN, Koch I. 2019. Cognitive control in the cocktail party:

Preparing selective attention to dichotically presented voices supports distractor

suppression. Attention, Perception, Psychophys. 81:727–737.

Nolte G, Müller KR. 2010. Localizing and estimating causal relations of interacting

brain rhythms. Front Hum Neurosci. 4:1–5.

O'Sullivan JA, Power AJ, Mesgarani N, Rajaram S, Foxe JJ, Shinn-Cunningham BG,

Slaney M, Shamma SA, Lalor EC. 2015. Attentional Selection in a Cocktail

Party Environment Can Be Decoded from Single-Trial EEG. Cereb Cortex.

25:1697–1706.

Oostenveld R, Fries P, Maris E, Schoffelen J. 2011. FieldTrip: Open Source Software

for Advanced Analysis of MEG, EEG, and Invasive Electrophysiological Data.

Comput Intell Neurosci. 2011:1–9.

Pasley BN, David S V., Mesgarani N, Flinker A, Shamma SA, Crone NE, Knight RT,

Chang EF. 2012. Reconstructing Speech from Human Auditory Cortex. PLoS

Biol. 10:e1001251.

Pisoni DB, Luce PA. 1987. Acoustic-phonetic representations in word recognition.

Cognition. 25:21–52.

Rao RPN, Ballard DH. 1999. Predictive coding in the visual cortex: A functional

interpretation of some extra-classical receptive-field effects. Nat Neurosci. 2:79–

87.

Rassi E, Wutz A, Müller-Voggel N, Weisz N. 2019. Prestimulus feedback

connectivity biases the content of visual experiences. Proc Natl Acad Sci.

116:16056–16061.

Samaha J, Bauer P, Cimaroli S, Postle BR. 2015. Top-down control of the phase of

alpha-band oscillations as a mechanism for temporal prediction. Proc Natl Acad

Sci. 112:8439–8444.

Schröger E, Kotz SA, SanMiguel I. 2015. Bridging prediction and attention in current

research on perception and action. Brain Res. 1626:1–13.

Schröger E, Marzecová A, Sanmiguel I. 2015. Attention and prediction in human

audition: A lesson from cognitive psychophysiology. Eur J Neurosci. 41:641–

664.

Smith JL, Johnstone SJ, Barry RJ. 2006. Effects of pre-stimulus processing on

subsequent events in a warned Go/NoGo paradigm: Response preparation,

execution and inhibition. Int J Psychophysiol. 61:121–133.

Smout CA, Tang MF, Garrido MI, Mattingley JB. 2019. Attention promotes the

neural encoding of prediction errors. PLoS Biol. 17:1–22.

Söderström P, Horne M, Frid J, Roll M. 2016. Pre-Activation Negativity (PrAN) in

Brain Potentials to Unfolding Words. Front Hum Neurosci. 10:1–11.

Söderström P, Horne M, Mannfolk P, van Westen D, Roll M. 2018. Rapid syntactic

pre-activation in Broca's area: Concurrent electrophysiological and

haemodynamic recordings. Brain Res. 1697:76–82.

Sonkusare S, Breakspear M, Guo C. 2019. Naturalistic Stimuli in Neuroscience:

Critically Acclaimed. Trends Cogn Sci. 1–16.

Stefanics G, Hangya B, Hernadi I, Winkler I, Lakatos P, Ulbert I. 2010. Phase

Entrainment of Human Delta Oscillations Can Mediate the Effects of

Expectation on Reaction Speed. J Neurosci. 30:13578–13585.

Strauß XA, Henry MJ, Scharinger XM, Obleser XJ. 2015. Alpha Phase Determines

Successful Lexical Decision in Noise. 35:3256–3262.

Szewczyk JM, Schriefers H. 2018. The N400 as an index of lexical preactivation and

its implications for prediction in language comprehension. Lang Cogn Neurosci.

33:665–686.

Teng X, Ma M, Yang J. 2020. Constrained Structure of Ancient Chinese Poetry

Facilitates Speech Content Grouping. Curr Biol. 30:1–7.

Tian X, Ding N, Teng X, Bai F, Poeppel D. 2018. Imagined speech influences

perceived loudness of sound. Nat Hum Behav. 2:225–234.

Van Dijk H, Schoffelen JM, Oostenveld R, Jensen O. 2008. Prestimulus oscillatory

activity in the alpha band predicts visual discrimination ability. J Neurosci.

28:1816–1823.

Van Petten C, Luka BJ. 2012. Prediction during language comprehension: Benefits,

costs, and ERP components. Int J Psychophysiol. 83:176–190.

VanRullen R. 2016. Perceptual Cycles. Trends Cogn Sci. 20:723–735.

Verhulst S, Altoè A, Vasilkov V. 2018. Computational modeling of the human

auditory periphery: Auditory-nerve responses, evoked potentials and hearing

loss. Hear Res. 360:55–75.

Wang L, Kuperberg G, Jensen O. 2018. Specific lexico-semantic predictions are

associated with unique spatial and temporal patterns of neural activity. Elife.

7:1–24.

Wöstmann M, Waschke L, Obleser J. 2019. Prestimulus neural alpha power predicts

confidence in discriminating identical auditory stimuli. Eur J Neurosci. 49:94–

105.

Zion Golumbic E, Cogan GB, Schroeder CE, Poeppel D. 2013. Visual input enhances

selective speech envelope tracking in auditory cortex at a "cocktail party". J

Neurosci. 33:1417–1426.

Zion Golumbic EM, Ding N, Bickel S, Lakatos P, Schevon CA, McKhann GM,

Goodman RR, Emerson R, Mehta AD, Simon JZ, Poeppel D, Schroeder CE.

2013. Mechanisms underlying selective neuronal tracking of attended speech at a

"cocktail party." Neuron. 77:980–991.

Zoefel B, Archer-Boyd A, Davis MH. 2018. Phase Entrainment of Brain Oscillations

Causally Modulates Neural Responses to Intelligible Speech. Curr Biol. 28:401-

408.e5.

884