Maternal Odor Reduces the Neural Threat Response in Human Infants

Sarah Jessen

Department of Neurology, University of Lübeck, Lübeck, Germany

Short title: Maternal Odor and Fear Processing in Infancy

Sarah Jessen, Department of Neurology, University of Lübeck, Ratzeburger Allee 160, 23562 Lübeck, Germany

Email: sarah.jessen@neuro.uni-luebeck.de

Phone: +49 451 3101 7449

ABSTRACT

Maternal odor is known to play an important role in mother-infant-interaction in

many altricial species such as rodents. However, we only know very little about

its role in early human development. The present study therefore investigated the

impact of maternal odor on infant brain responses to threat signals. We recorded

the electroencephalographic (EEG) signal of seven-month-old infants watching

happy and fearful facial expressions. While infants in two control groups showed

the expected EEG fear response, this response was markedly absent in the group

exposed to their mother's odor. Thus, infants respond differently to fear signals in

the presence of their mother and the mother's odor is a sufficiently strong signal

to elicit this effect. Our data suggest that olfaction, a sensory modality that has

been largely neglected as a social signal in our own species, might function as a

crucial modulator in early social learning.

<u>Keywords:</u> infancy, emotion perception, odor, breastfeeding, fear processing, EEG

 INTRODUCTION

As members of an altricial species, newborn humans completely rely on their
social environment for survival. To foster and support the care they receive,
newborns show a number of mechanisms to support social bonding, including a
strong preference for faces (Johnson, Dziurawiec, Ellis, & Morton, 1991) and their
mother's voice (DeCasper & Fifer, 1980). One modality however that is potentially
of equal importance but has received far less attention in human research is
olfaction as a form of extremely potent chemosensory signaling (Semin & Groot,
2013).

The importance of olfaction for early social development has been amply
investigated in other species, especially rodents. Rat pups form a strong
attachment to the dam which is crucially supported by a (learned) preference for
the maternal odor (Landers & Sullivan, 2012).

Initial research in human development suggests that olfaction and especially
maternal odor are important components for social development also in our
species (Lubke & Pause, 2015). Maternal odor cues for instance play a crucial role
in the successful initiation of breastfeeding (Porter & Winberg, 1999; Schaal et al.,
2009), essential for the survival of mammalian offsprings but also an important
player in the development of a secure bonding between mother and infant.
Furthermore, recent studies suggest that maternal odor impacts face perception,
in older infants (Durand, Baudouin, Lewkowicz, Goubet, & Schaal, 2013; Leleu et
al., 2019).

One setting in which this social learning is put to the test is in responses to adverse
situations such as stress and threat. Again, rodent research has provided us with
insights on the developmental trajectory of threat response and their modulation
by maternal presence and signals thereof. Very young pups who are highly
dependent on their mother show little fear responses but rather responses that
facilitate the formation of close bonding to their mother (Leon, 1992). Older pups
in contrast start to show more adult-like fear responses (Debiec & Sullivan, 2017).
Moreover, during a transition period, pups exhibit both types of responses; if their
mother is present, they show affiliative responses facilitating mother-infant-

bonding, if their mother is absent, they show fear learning typically seen in older

pups (Debiec & Sullivan, 2017; Landers & Sullivan, 2012; Moriceau & Sullivan,

2006). Hence, maternal presence appears to be a key player in determining the

processing of and response to threat in early development of rodents, and maternal

odor is assumed to be a strong signal of maternal presence.

Studying responses to threat signals in human infants is more challenging. One

well-investigated phenomenon in this context is the development of an attentional

bias towards fearful expressions around 7 months of age (Vaish, Grossmann, &

Woodward, 2008). Interestingly, recent work suggests that this phenomenon can

be strongly influenced by secondary factors, such as parental sensitivity (Taylor-

Colls & Pasco Fearon, 2015), infant temperament (Martinos, Matheson, & de Haan,

2012), and breastfeeding experience (Krol, Rajhans, Missana, & Grossmann, 2014).

Crucially, all these factors are linked to the interplay between the infant and her

social environment, providing initial evidence for a modulation by social factors.

However, all the above-mentioned components are typically stable factors relating

to interindividual differences rather than flexible changes in a given situation. It is

therefore unclear whether threat responses in human infants can be modulated in

the same way as has been suggest for rodents. Does maternal presence, as a social

short-term rather than long-term factor, impact infants' responses to threat

signals? And is maternal odor a sufficient strong cue to elicit an effect of maternal

presence also in humans?

To address these questions, we designed an experiment to investigate the impact

of maternal odor on the neural response to threat signals in human infants. In an

electroencephalographic (EEG) set-up, infants were presented with happy and

fearful facial expressions while they were exposed to either the familiar maternal

odor, to an unfamiliar mother's odor, or to no specific odor at all. To quantify

infants' fear response, we investigated the amplitude of the *Nc*, an infant event-

related potential (ERP) component observed between 400 and 800 ms after the

onset of a stimulus at frontocentral electrodes. The Nc amplitude has been linked

to the allocation of attention and is typically enhanced in response to fearful faces

in 7-month-old infants (Peltola, Leppänen, Mäki, & Hietanen, 2009). Seven months

marks a crucial transitional period in human infants, and is associated not only

with the onset of stronger fear responses but also with the onset of locomotion

(Leppänen & Nelson, 2012). Hence, this age may be a period in humans analogous

to the transitional period described for rodents above; a shift from dependence to

increasing independence.

We therefore predict that infants show an increased threat response in the absence

of their mother (or an odor signaling her presence), but we expect this response to

be reduced in their mother's presence, even if only signaled via maternal odor.

METHODS

Participants. 76 7-month-old infants were included in the final sample (age: 213 ± 8

days [mean ± standard deviation (SD)]; range: 200-225, 38 female, see Table 1 for

description of the individual groups). An additional 15 infants had been tested but

were not included in the final sample because they did not provide at least 10

artifact-free trials per condition (n=11); had potential neurological problems (n=1);

were erroneously invited too young (n=1); the mean ERP response in the time-

window and electrodes of interest was more than 4 standard deviations from the

mean (n=1, see below); or because of technical problems during the recording

(n=1).

The sample size was determined by statistical considerations and practical

conventions in the field. First, for practical considerations and the known high

attrition rates in infant EEG studies, we had planned a priori to keep collecting

data until 25 useable data sets per each of the three experimental manipulation

groups were obtained. Second, as outlined in Albers & Lakens (2018), a smallest

effect size of interest was critical here, as too small true effects sizes for odor

manipulations would not be of practical or translational relevance. In the present

study, a total sample size of n=75 in three groups, was thus powered with 80% or

more to detect medium and large effects (i.e., Cohen's d of 0.8 or larger) at a

conventional type I error level of 5 %.

Infants were recruited via the maternity ward at the local hospital

(Universitätsklinikum Schleswig-Holstein), were born full-term (38–42 weeks

gestational age), had a birth weight of at least 2500 g, and had no known

neurological deficits. The study was conducted according to the Declaration of Helsinki, approved by the ethics committee at the University of Lübeck, and parents provided written informed consent.

Table 1. Overview of participants included in the final analysis. An additional 15 infants were tested but not included in the final analysis for various reasons (see text).

	N	female	age (in days)*	still breastfed	trials (happy)*	trials (fearful)*	Inf Neg Temp*#	EPDS*#
Maternal odor	25	13	213 ± 7	14	38 ± 18	38 ± 17	3.03 ± 0.68	4.32 ± 3.74
No odor	26	9	214 ± 8	18	45 ± 21	45 ± 23	3.34 ± 0.68	5.08 ± 4.77
Stranger odor	25	16	215 ± 7	13	37± 18	37± 17	3.25 ± 0.76	4.54 ± 3.96

*mean ± standard deviation; # excluding one participant in the Stranger odor group, who did not fill in the questionnaire; Inf Neg Temp = Infant Negative Temperament, see text; EPDS = Edinburgh Postpartum Depression Screening, see text

Stimulus. As emotional face stimuli, we used colored photographs of happy and fearful facial expressions by 6 actresses from the FACES database (Ebner, Riediger, & Lindenberger, 2010 [actress-ID 54, 63, 85, 90, 115, 173]). Photographs were cropped so that only the face was visible in an oval shape, and have successfully been used in prior studies to investigate fear processing in infancy (Jessen & Grossmann, 2015, 2017).

Odor manipulation. Prior to a scheduled experimental recording, all infants' mothers were given a white cotton t-shirt and instructed to wear this t-shirt for three nights in a row. The mother was asked to store the t-shirt in a provided zip-lock bag during the day, and use her normal shampoo, soap, deodorant etc. as usual but refrain from using new products. Before the t-shirt was given to the mother, it had been washed with the same detergent for all t-shirts.

If the mother wore the t-shirt during the three nights directly preceding the experiment, the mother was asked to simply bring the t-shirt along. If the three nights did not directly precede the experiment, the mother was asked to store the t-shirt in a zip-lock bag in the freezer, as freezing has been shown to conserve odor (Lenochova, Roberts, & Havlicek, 2009).

Randomization. Infants were randomly assigned to either the *Maternal odor* group or one of the control groups (*No odor* group or *Stranger odor* group; Figure 1). As only constraint to fully random assignment, we monitored as the study proceeded

that groups did not differ in gender, age, or breastfeeding experience. Infants in the *Maternal odor* group were administered the t-shirt previously worn by their mother during the experiment. Infants in the *No odor* group were administered an unworn t-shirt. Infants in the *Stranger odor* group were administered a t-shirt previously worn by the mother of one of the other infants. The t-shirt of their own mother was stored in a freezer to be used as a stimulus for a different infant in the *Stranger odor* group. Except in one case, both, parents and the experimenter administering the t-shirt, were blind to the group assignment.

Figure 1. Overview of experimental set-up. A) Mothers were asked to wear a provided t-shirt for 3 nights in a row prior to the experiment. The infant was randomly assigned to one of three groups; a Maternal odor group (exposed to the t-shirt worn by the infant's mother), a Stranger odor group (exposed to a t-shirt worn by a different infant's mother), or a No odor group (exposed to an unworn t-shirt). We recorded the EEG signal while the infants were seated in a car seat with the t-shirt positioned over their chest area and watched happy and fearful facial expressions. B) Example of fearful and happy faces used as stimulus material, the colored circles are for illustration purpose only and correspond to the color coding used in the following figures.

Procedure and experimental design. Before the laboratory visit, families were sent the t-shirt (as described above) as well as a set of questionnaires, in particular the EPDS (Cox, Holden, & Sagovsky, 1987), the IBQ-R (Gartstein & Rothbart, 2003; Vonderlin, Ropeter, & Pauen, 2012), and a lab-internal questionnaire assessing demographic information as well as feeding and sleeping routines of the infant (One family, whose infant was assigned to the *Stranger odor* group, did not fill in the IBQ-R and the EPDS and is therefore not included in the control analyses with these two factors). After arriving in the laboratory, parents and infant were familiarized with the environment and parents were informed about the study and signed a consent form. The EEG recording was prepared while the infant was sitting on his/her parent's lap. For recording, we used an elastic cap (BrainCap, Easycap GmbH) in which 27 AgAgCl-electrodes were mounted according to the

international 10-20-system. An additional electrode was attached below the infant's right eye to record the electrooculogram. The EEG signal was recorded with a sampling rate of 250 Hz using a BrainAmp amplifier and the BrainVision Recorder software (both Brain Products).

For the EEG recording, the infant was sitting in an age appropriate car seat (Maxi Cosi Pebble) positioned on the floor. The t-shirt was positioned over the chest area of the infant, folded along the vertical axis of the t-shirt and with the armpit region of the t-shirt directed towards the infant's face. The t-shirt was fixated using the safety straps of the car seat as closely to the chin of the infant as possible and adjusted during the experiment if necessary.

In front of the infant (approximately 60 cm from the infant's feet), a 24-inch monitor with a refresh rate of 60 Hz was positioned at a height of about 40 cm (bottom edge of the screen). The parent was seated approximately 1.5 m behind the infant and instructed not to interact with the infant during the experiment.

The experiment was programmed using the Presentation software (Version 18.1). Faces were presented for 800 ms, preceded by a fixation cross presented for 300 ms, and followed by an intertrial interval jittered between 800 and 1200 ms. The faces had a height of approximately 28 cm. If necessary, short video clips containing colorful moving shapes and ringtones were played during the experiment to redirect the infant's attention to the screen. Each infant saw a maximum of 216 trials, arranged in miniblocks of 24 trials containing 12 happy and 12 fearful faces and played consecutively without interruption. Trials were presented in a pseudorandomized order, ensuring that no stimulus category (happy, fearful) was repeated more than once. The experiment continued until the infant had seen all trials or became too fussy to continue the experiment. During the experiment, the infant was video-recorded using a small camera mounted on top of the monitor to offline exclude trials in which the infant did not attend to the screen.

Data Analysis. We analyzed the data using Matlab 2013b (The MathWorks, Inc., Natick, MA), the Matlab toolbox FieldTrip (Oostenveld, Fries, Maris, & Schoffelen, 2011), and for statistical analysis the package JASP (JASP Team, version 0.10.2).

EEG Preprocessing. For purposes of artefact removal including an independent
component analysis (ICA) routine, all data were first referenced to the average of
all electrodes (average reference), filtered using a 100-Hz lowpass and a 1-Hz
highpass filter, and segmented into 1-sec-epochs. To detect epochs obviously
contaminated by artifacts, the standard deviation was computed in a sliding
window of 200 msec. If the standard deviation exceeds 100 µV at any electrode,
the entire epoch was discarded. Next, an independent component analysis (ICA)
was computed on the remaining concatenated data. Components were inspected
visually and rejected if classified as artefactual (4 ± 2 components per participants
[mean ± SD], range 0–10 components).
After removal of ICA components, the data was re-segmented into epochs ranging
from 200 ms before to 800 ms after the onset of the stimulus, re-referenced to the
linked mastoids (mean of TP9 and TP10), and a 0.2 to 20 Hz bandpass filter was
applied. A last step of automatic artifact detection was applied, rejecting all epochs
in which the standard deviation exceeded 80 µV. Data was inspected visually for
remaining artifacts, and all trials in which the infant did not attend to the screen
(as assessed via the video recording during the experiment) were rejected (see
Table 1 for number of remaining trials).

ERP analysis. To analyze the Nc response, we computed the mean response in a
time-window of 400–800 ms after stimulus onset across frontocentral electrodes
(F3, Fz, F4, C3, Cz, C4; see Supplementary Material for an analysis of occipital
electrodes, where no significant effect was found). One participant was rejected
from further analysis because the difference in the mean response to happy and
fearful faces in this time-window and electrode cluster was more than 4 standard
deviations from the mean across all other participants. Mean responses were
entered into a repeated measures ANOVA with the within-subject factor Emotion
(happy, fear) and the between-subject factor Odor (maternal, stranger, no odor).
Furthermore, we included the infant's current breastfeeding status (whether s/he
was still breastfed at the time of testing or not) as reported by the mother (Breastfed
[yes,no]) as a covariate, as lactation may impact the mother's body odor
(McClintock et al., 2005). Student's t-tests are computed as post-how tests and
effect sizes are reported as partial eta squared (η_p^2) and Cohen's *d*. In addition, we

also performed the equivalent analysis using Bayesian statistics; BF_{10} values above

1 are interpreted as anecdotal evidence, above 3 as moderate evidence, and above

10 as strong evidence for the research hypothesis (Wagenmakers et al., 2018).

To further analyze the Emotion effect without any a priori assumption about its

latency and topography, we ran a cluster-based permutation test (Maris &

Oostenveld, 2007) with 1000 permutations contrasting responses to happy and

fearful faces separately for each *Odor* group. A cluster had to comprise at least 2

adjacent electrodes, was computed across time and electrode position, and a type-

1-error probability of less than 0.05 at the cluster-level was ensured.

Negative Affect. Negative affect was computed as the mean of the IBQ-R scales

Sadness, Fear, and Distress to Limitations (Aktar et al., 2018).

RESULTS

Influence of maternal odor on the Nc response. We observed an overall enhanced Nc

amplitude in response to fearful faces, confirming the expected heightened threat

response (Peltola et al., 2009; significant main effect of Emotion [$F(1,72) = 11.60$, p

$= .001$, $\eta_p^2 = 0.14$; $BF_{10} = 2.578$]).

Most importantly, however, the threat response differed between odor groups.

That is, this emotion effect critically depended on the odor group an infant had

been assigned to (significant interaction Emotion × Odor [$F(2,72) = 5.57$, $p = .006$,

$\eta_p^2 = 0.13$; $BF_{10} = 4.564$; Figure 2]).

Figure 2. ERP response in the different odor groups. A) Shows the Nc response at frontocentral

electrodes (F3, Fz, F4, C3, Cz, C4, marked by black dots) to fearful (orange) and happy (blue) facial

expressions. While no difference in response was observed in the Maternal odor group, infants in

the No odor and the Stranger odor group showed a significantly enhanced Nc response to fearful

faces. Topographic representations averaged between 400 and 800 ms after face onset are shown at

Follow-up tests confirmed the Nc effect to fear signals was critically absent in the *Maternal odor* group [$t(24) = -0.95$, $p = .35$, $d = -.19$; $BF_{10} = 0.32$; fearful: -6.51 ± 2.99 µV, happy: -8.53 ± 3.17 µV]. In contrast, the typical enhancement of the Nc response to fearful (compared to happy) faces was present in the *Stranger odor* group [$t(24) = 2.51$, $p = .019$, $d = .50$; $BF_{10} = 2.78$; fearful: -10.57 ± 2.34 µV (mean ± SE), happy: -5.66 ± 1.78 µV] as well as in the *No odor* group [$t(25) = 3.50$, $p = .002$, $d = .68$; $BF_{10} = 21.02$; fearful: -16.94 ± 2.19 µV, happy: -11.43 ± 2.53 µV].

Corroborating analysis using a cluster-based permutation approach. While the electrode and time window selection for this analysis had not been data derived but followed standards set by previous studies (Jessen & Grossmann, 2014, 2016, 2019), we aimed to corroborate this main result by a more data-driven search for potential effects using a cluster-based permutation test (Figure 3). In both, the *No odor* group and the *Stranger odor* group, nearly identical clusters indicating a significantly response enhancement to fearful (compared to happy) faces was found (No odor: $p = .006$, $T_{sum} = 3063.8$; Stranger odor: $p = .021$, $T_{sum} = 1272.8$). Importantly, both clusters exhibit the latency and topographic distribution typical for an Nc response. Most importantly, no such cluster of significant differences was found in the *Maternal odor* group when contrasting responses to happy and fearful faces.

Hence, while both control groups (*No group* and *Stranger odor* group) showed the age-typical enhanced Nc response to fearful, threat-signalling faces, a heightened fear response was absent in the *Maternal odor* group. Our results suggest that maternal odor, as a signal of familiarity and maternal presence, reduces infant's attention allocation to threat signals.

No effect of potential confounds. Importantly, we did not find a difference between the three groups with respect to a number of potential confounds: There were no group differences in the number of included trials per infant in either Emotion condition [happy: $F(2,73) = 1.49$, $p = .23$, $BF_{10} = 0.355$; fearful: $F(2,73) = 1.25$, $p = .29$, $BF_{10} = 0.296$]; age [$F(2,73) = 0.49$, $p = .61$, $BF_{10} = 0.165$]; in maternal depression scores as assessed via the EPDS [$F(2,72) = 0.22$, $p = .80$, $BF_{10} = 0.136$]; or in infant negative temperament as assessed via the IBQ-R [$F(2,72) = 1.23$, $p = .30$, $BF_{10} = 0.294$].

Effect of Breastfeeding. A last finding supported our general line of reasoning. Namely, we did observe an interaction between Nc response to the emotional expression of the presented face and whether the infant was still breastfed or not [Emotion × Breastfeeding, $F(1,72) = 5.06$, $p = .028$, $\eta_p^2 = 0.07$; $BF_{10} = 1.632$; Figure 4]. Only the infants who were not breastfed any more at the time of testing showed an enhanced Nc response to fearful faces [$t(30) = 3.55$, $p = .001$, $d = .64$; $BF_{10} = 26.54$; fearful: -13.35 ± 2.18 μV, happy: -7.90 ± 2.00 μV], while this enhancement was absent in the infants who were still breastfed [$t(44) = 0.65$, $p = .52$, $d = 0.1$; $BF_{10} = 0.20$; fearful: -10.08 ± 2.08 μV, happy: -9.05 ± 2.10 μV].

Importantly, this was independent of (i.e., additionally true but not interacting

with) the *odor group* manipulation, as there was no meaningful Emotion ×

Breastfeeding × Odor interaction [$F(2,70) = 2.20$, $p = .12$, $\eta_p^2 = 0.06$, $BF_{10} = 1.081$].

Figure 4. Nc response depending on breastfeeding status. Nc response is depicted at frontocentral

electrodes (F3, Fz, F4, C3, Cz, C4, marked by black dots) to fearful (orange) and happy (blue) facial

expressions for infants who are still breastfed (left) and not breastfed anymore (right). Infants who

are not breastfed any more show an enhanced Nc response to fearful faces, while this effect was

absent in the group of infants who were still breastfed. Topographic representations averaged

between 400 and 800 ms after face onset are shown at the bottom.

DISCUSSION

Our results demonstrate that maternal odor is a sufficiently strong signal to reduce

an established neural marker of the fear response in 7-month-old infants. A highly

consonant effect was found for breastfeeding, suggesting that not only momentary

states but also longer-lasting effects of maternal presence impact fear processing

in infants. Our results point to a previously unknown flexibility in fear processing

in infancy and its dependency on context and experience, which is in line with

findings in other species (Landers & Sullivan, 2012).

Maternal odor as a momentary modulator of infant fear processing

We suggest that such a response pattern might be characteristic for a developing

system that on the one hand needs to establish a close bonding to a caregiver,

typically the mother, while on the other hand learning to respond to potential

threat signals in the environment. This has been indirectly suggested by studies in

older children (Gee et al., 2014) as well as rodent research (Landers & Sullivan,

2012). Our findings, however, provide first evidence for flexible processing of

threat-related information depending on maternal signals in early human

development.

A diminished response to threat signals in maternal presence might facilitate

bonding, as a positive evaluation of information and less attention to potential

negative aspects may increase positive affect towards the caregiver even in the

presence of negative signals. In addition, if maternal presence works as a "safety

signal", requiring the infant to allocate less attention to potential threats, this might

also free cognitive capacities in the infant for other processes, akin to previously

reported improved cognitive performance in rat pups in the presence of familiar

odor (Wigal, Kucharski, & Spear, 1984).

At the same time, as the infant grows more independent, detecting and responding

to threat becomes of growing importance, especially if the mother is not present.

Crucially, 7 months is an important turning point in early human development,

characterized not only by qualitative changes in emotion development, but also by

the onset of locomotion, an important step towards growing independence

(Leppänen & Nelson, 2012). During this period, flexible threat processing might

be of particular importance, akin to what has been suggested in the rodent

literature (Landers & Sullivan, 2012).

Our results further underscore the importance of odor in early social development.

Two recent studies have suggested a modulation of infant face processing in

general by the presence of maternal odor (Durand et al., 2013; Leleu et al., 2019).

Our study is the first to show that the absence or presence of maternal odor can

impact highly specific aspects of infant face processing. Maternal odor might

therefore be an important guiding factor in emotional learning in infancy.

Specifically, we found an impact on the attention-related Nc component (Webb,

Long, & Nelson, 2005) but no influence on early visual processing (see

supplementary material) or on the number of trials the infants watched. Therefore,

we found no evidence for a general impact of maternal odor on sensory processing
or compliance with the experiment, but rather odor specifically impacted the
evaluation of facial information, further underscoring its potential role in early
social learning.

Importantly, we assume that the observed effects are due to a *learned* odor
preference for familiar maternal signals rather than a specific preference for
maternal body odor. This assumption is in line with rodent work, suggesting that
effects akin to the ones reported for maternal odor can also be found other types
of highly familiar odor (Wigal et al., 1984). Accordingly, the present manipulation
did not differentiation between body odor and other odor components (such as
deodorant used or specific food consumed by the mother), thereby reflecting the
mélange of odors the infant experiences in maternal presence in everyday life.

Breastfeeding as a long-term modulator of infant fear processing

While maternal odor as a direct or phasic signal of maternal presence influenced
fear processing, so did the more tonic variable of an infant's breastfeeding
experience. In line with our interpretation of the above-mentioned odor effect, this
finding suggests a further modulatory impact of mother-infant-bonding in fear
processing: Our findings are in line with prior studies reporting an increased bias
towards expressions of happiness with increasing breastfeeding experience (Krol,
Monakhov, Lai, Ebstein, & Grossmann, 2015; Krol et al., 2014). How exactly
breastfeeding experience interacts with emotion processing is not certain, but a
possible explanation is an increased closeness between mother and infants;
breastfed infants on average spend more time interacting with their mother (Smith
& Forrester, 2017) and show a higher attachment security (Gibbs, Forste, &
Lybbert, 2018).

Maternal presence may therefore not only modulate fear processing directly, as
suggested by the influence of maternal odor, but might also exert a longer-lasting
impact.

Future Directions

An important aspect in further assessing the relevance of maternal odor in infancy

are potential changes across development. If the impact of maternal presence on

fear responses follows a similar trajectory as has been reported for rodents

(Landers & Sullivan, 2012), this predicts a specific time-window during which

infants show flexible fear responses depending on the presence and by extension

the odor of their mother. This time-window might be around seven months of age,

when infants first acquire the ability to locomote (Leppänen & Nelson, 2012), but

future studies tracing the impact of maternal odor longitudinally are clearly

warranted to solidify this suggestion.

In the present approach, we assume that the reported influence of maternal odor

is the result of early learning processes, during which the infants learns to associate

her mother with a certain odor, the maternal odor, consisting of a combination of

the mother's body odor and familiar home scents. Hence, future studies need to

consider the differential contributions of body odor in comparison to other types

of odor the infant might be highly familiar with.

Finally, the generalizability to other types of adverse signals needs to be assessed

in future work. We show that maternal odor reduced attention to fearful faces, and

we predict that maternal odor should also impact infant responses to other

negative signals, such as pain or aversive sounds. However, since recent studies

suggest a link between maternal odor and the processing of faces in infancy

(Durand et al., 2013; Leleu et al., 2019), one could also expect a more specific effect

of maternal odor on negative social information.

Conclusions

The current study demonstrates that maternal odor influences fear processing in

infancy. While infants in two control groups of different specificity (a stranger's

odor or no specific odor at all) showed an expectably enhanced attentional

response to fear signals, this response was absent in infants who could smell their

mother. Our results establish that the mother's presence, even if just signaled by

the mother's familiar odor, can result in a marked reduction of the neurobiological

fear response in infants. Furthermore, our data provide evidence for the potency
of odor as a social signal in humans and in particular in early ontogeny.

Acknowledgements

This work was supported by funding of the German Research Foundation (DFG,
grant-number JE 781/1-1 & 2). We thank Leonie Emmerich and Aylin Ulubas for
help with the data acquisition, Jonas Obleser for helpful comments on the
manuscript, and all the families for participating.

BIBLIOGRAPHY

Aktar, E., Mandell, D. J., de Vente, W., Majdandžić, M., Oort, F. J., van Renswoude, D. R.,
… Bögels, S. M. (2018). Parental negative emotions are related to behavioral and
pupillary correlates of infants' attention to facial expressions of emotion. *Infant*
Behavior and Development. https://doi.org/10.1016/j.infbeh.2018.07.004

Albers, C., & Lakens, D. (2018). When power analyses based on pilot data are biased:
Inaccurate effect size estimators and follow-up bias. *Journal of Experimental Social*
Psychology. https://doi.org/10.1016/j.jesp.2017.09.004

Cox, J. L., Holden, J. M., & Sagovsky, R. (1987). Detection of postnatal depression.
Development of the 10-item Edinburgh Postnatal Depression Scale. *British Journal of*
Psychiatry, *150*, 782–786. Retrieved from
http://www.ncbi.nlm.nih.gov/pubmed/3651732

Debiec, J., & Sullivan, R. M. (2017). The neurobiology of safety and threat learning in
infancy. *Neurobiology of Learning and Memory*.
https://doi.org/10.1016/j.nlm.2016.10.015

DeCasper, A. J., & Fifer, W. P. (1980). Of human bonding: newborns prefer their mothers'
voices. *Science*, *208*(4448), 1174–1176. Retrieved from
http://www.ncbi.nlm.nih.gov/pubmed/7375928

Durand, K., Baudouin, J. Y., Lewkowicz, D. J., Goubet, N., & Schaal, B. (2013). Eye-
catching odors: olfaction elicits sustained gazing to faces and eyes in 4-month-old
infants. *PloS One*, *8*(8), e70677. https://doi.org/10.1371/journal.pone.0070677

Ebner, N. C., Riediger, M., & Lindenberger, U. (2010). FACES--a database of facial
expressions in young, middle-aged, and older women and men: development and
validation. *Behavior Research Methods*, *42*(1), 351–362.
https://doi.org/10.3758/BRM.42.1.351

Gartstein, M. A., & Rothbart, M. K. (2003). Studying infant temperament via the Revised
Infant Behavior Questionnaire. *Infant Behavior and Development*, *26*(1), 64–86.

Gee, D. G., Gabard-Durnam, L., Telzer, E. H., Humphreys, K. L., Goff, B., Shapiro, M., …
Tottenham, N. (2014). Maternal Buffering of Human Amygdala-Prefrontal Circuitry
During Childhood but Not During Adolescence. *Psychological Science*.
https://doi.org/10.1177/0956797614550878

Gibbs, B. G., Forste, R., & Lybbert, E. (2018). Breastfeeding, Parenting, and Infant

Attachment Behaviors. *Maternal and Child Health Journal*.

https://doi.org/10.1007/s10995-018-2427-z

Jessen, S., & Grossmann, T. (2014). Unconscious discrimination of social cues from eye

whites in infants. *Proceedings of the National Academy of Sciences of the United States of*

America, *111*(45), 16208–16213. https://doi.org/10.1073/pnas.1411333111

Jessen, S., & Grossmann, T. (2015). Neural signatures of conscious and unconscious

emotional face processing in human infants. *Cortex*, *64*, 260–270.

https://doi.org/10.1016/j.cortex.2014.11.007

Jessen, S., & Grossmann, T. (2016). The developmental emergence of unconscious fear

processing from eyes during infancy. *Journal of Experimental Child Psychology*, *142*,

334–343. https://doi.org/10.1016/j.jecp.2015.09.009

Jessen, S., & Grossmann, T. (2017). Exploring the role of spatial frequency information

during neural emotion processing in human infants. *Frontiers in Human*

Neuroscience, *11*. https://doi.org/10.3389/fnhum.2017.00486

Jessen, S., & Grossmann, T. (2019). Neural evidence for the subliminal processing of facial

trustworthiness in infancy. *Neuropsychologia*, *126*, 46–53.

https://doi.org/10.1016/j.neuropsychologia.2017.04.025

Johnson, M. H., Dziurawiec, S., Ellis, H., & Morton, J. (1991). Newborns' preferential

tracking of face-like stimuli and its subsequent decline. *Cognition*, *40*(1–2), 1–19.

Retrieved from http://www.ncbi.nlm.nih.gov/pubmed/1786670

Krol, K. M., Monakhov, M., Lai, P. S., Ebstein, R. P., & Grossmann, T. (2015). Genetic

variation in CD38 and breastfeeding experience interact to impact infants' attention

to social eye cues. *Proceedings of the National Academy of Sciences of the United States of*

America, *112*(39), E5434-42. https://doi.org/10.1073/pnas.1506352112

Krol, K. M., Rajhans, P., Missana, M., & Grossmann, T. (2014). Duration of exclusive

breastfeeding is associated with differences in infants' brain responses to emotional

body expressions. *Frontiers in Behavioral Neuroscience*, *8*, 459.

https://doi.org/10.3389/fnbeh.2014.00459

Landers, M. S., & Sullivan, R. M. (2012). The development and neurobiology of infant

attachment and fear. *Developmental Neuroscience*, *34*(2–3), 101–114.

https://doi.org/000336732

Leleu, A., Rekow, D., Poncet, F., Schaal, B., Durand, K., Rossion, B., & Baudouin, J.-Y.

(2019). Maternal odor shapes rapid face categorization in the infant brain. *Dev Sci*.

Retrieved from https://doi.org/10.1111/desc.12877

Lenochova, P., Roberts, S. C., & Havlicek, J. (2009). Methods of human body odor

sampling: the effect of freezing. *Chemical Senses*, *34*(2), 127–138.

https://doi.org/10.1093/chemse/bjn067

Leon, M. (1992). The Neurobiology Of Filial Learning. *Annual Review of Psychology*.

https://doi.org/10.1146/annurev.psych.43.1.377

Leppänen, J. M., & Nelson, C. A. (2012). Early development of fear processing. *Current*

Directions in Psychological Science, *21*, 200–204.

Lubke, K. T., & Pause, B. M. (2015). Always follow your nose: the functional significance

of social chemosignals in human reproduction and survival. *Hormones and Behavior*,

68, 134–144. https://doi.org/10.1016/j.yhbeh.2014.10.001

Maris, E., & Oostenveld, R. (2007). Nonparametric statistical testing of EEG- and MEG-

data. *Journal of Neuroscience Methods*, *164*(1), 177–190.

Martinos, M., Matheson, A., & de Haan, M. (2012). Links between infant temperament

and neurophysiological measures of attention to happy and fearful faces. *Journal of*

Child Psychology and Psychiatry and Allied Disciplines, *53*(11), 1118–1127.

https://doi.org/10.1111/j.1469-7610.2012.02599.x

McClintock, M. K., Bullivant, S., Jacob, S., Spencer, N., Zelano, B., & Ober, C. (2005).

Human body scents: Conscious perceptions and biological effects. In *Chemical*

Senses. https://doi.org/10.1093/chemse/bjh151

Moriceau, S., & Sullivan, R. M. (2006). Maternal presence serves as a switch between

learning fear and attraction in infancy. *Nature Neuroscience*.

https://doi.org/10.1038/nn1733

Oostenveld, R., Fries, P., Maris, E., & Schoffelen, J.-M. (2011). FieldTrip: Open source

software for advanced analysis of MEG, EEG, and invasive electrophysiological

data. *Computational Intelligence and Neuroscience*, *2011*, 156869.

Peltola, M. J., Leppänen, J. M., Mäki, S., & Hietanen, J. K. (2009). Emergence of enhanced

attention to fearful faces between 5 and 7 months of age. *Social Cognitive and*

Affective Neuroscience, *4*, 134–142.

Porter, R. H., & Winberg, J. (1999). Unique salience of maternal breast odors for newborn

infants. *Neuroscience & Biobehavioral Reviews*, 23(3), 439–449. Retrieved from

http://www.ncbi.nlm.nih.gov/pubmed/9989430

Schaal, B., Coureaud, G., Doucet, S., Delaunay-El Allam, M., Moncomble, A. S.,

Montigny, D., … Holley, A. (2009). Mammary olfactory signalisation in females and

odor processing in neonates: ways evolved by rabbits and humans. *Behavioural Brain*

Research, *200*(2), 346–358. Retrieved from

http://www.ncbi.nlm.nih.gov/pubmed/19374020

Semin, G. R., & Groot, J. H. (2013). The chemical bases of human sociality. *Trends Cogn*

Sci, *17*(9), 427–429. https://doi.org/10.1016/j.tics.2013.05.008

Smith, J. P., & Forrester, R. (2017). Maternal Time Use and Nurturing: Analysis of the

Association between Breastfeeding Practice and Time Spent Interacting with Baby.

Breastfeeding Medicine. https://doi.org/10.1089/bfm.2016.0118

Taylor-Colls, S., & Pasco Fearon, R. M. (2015). The effects of parental behavior on infants'

neural processing of emotion expressions. *Child Development*, *86*(3), 877–888.

https://doi.org/10.1111/cdev.12348

Vaish, A., Grossmann, T., & Woodward, A. (2008). Not all emotions are created equal: the

negativity bias in social-emotional development. *Psychological Bulletin*, *134*(3), 383–

403. https://doi.org/10.1037/0033-2909.134.3.383

Vonderlin, E., Ropeter, A., & Pauen, S. (2012). Erfassung des fruehkindlichen

Temperaments mit dem Infant Behavior Questionnaire Revised. Psychometrische

Merkmale einer deutschen Version. *Zeitschrift Fuer Kinder- Und Jugendpsychiatrie*

Und Psychotherapie, *40*(5), 307–314.

Wagenmakers, E. J., Marsman, M., Jamil, T., Ly, A., Verhagen, J., Love, J., … Morey, R. D.

(2018). Bayesian inference for psychology. Part I: Theoretical advantages and

practical ramifications. *Psychonomic Bulletin and Review*.

https://doi.org/10.3758/s13423-017-1343-3

Webb, S. J., Long, J. D., & Nelson, C. A. (2005). A longitudinal investigation of visual

event-related potentials in the first year of life. *Dev Sci*, *8*(6), 605–616.

https://doi.org/10.1111/j.1467-7687.2005.00452.x

Wigal, T., Kucharski, D., & Spear, N. E. (1984). Familiar contextual odors promote

discrimination learning in preweanling but not in older rats. *Developmental*

Psychobiology. https://doi.org/10.1002/dev.420170512