A model to study orienting responses in zebrafish, and applications towards the emotion-cognition interaction

Bianca Gomes do Nascimento[1,2,3], Hingrid Suzzan Tarso Oliveira e Oliveira[1,3,4], Hadda Tercya Lima Silva[1,3,4], Diógenes Henrique de Siqueira-Silva[1,3,4], Monica Gomes Lima-Maximino[2,5], Caio Maximino[1,2*]

1 – Laboratório de Neurociências e Comportamento "Frederico Guilherme Graeff", Faculdade de Psicologia, Instituto de Estudos em Saúde e Biológicas, Universidade Federal do Sul e Sudeste do Pará, Marabá/PA, Brazil

2 – Grupo de Pesquisas em Neurofarmacologia e Psicopatologia Experimental, Universidade Federal do Sul e Sudeste do Pará, Marabá/PA, Brazil

3 – Faculdade de Biologia, Instituto de Estudos em Saúde e Biológicas, Universidade Federal do Sul e Sudeste do Pará, Marabá/PA, Brazil

4 – Grupo de Pesquisas em Reprodução de Peixes Amazônicos, Universidade Federal do Sul e Sudeste do Pará, Marabá/PA, Brazil

5 – Laboratório de Neurofarmacologia e Biofísica, Departamento de Morfologia e Ciências Fisiológicas, Universidade do Estado do Pará, Campus VIII, Marabá/PA, Brazil

* Corresponding author

Caio Maximino

Laboratório de Neurociências e Comportamento "Frederico Guilherme Graeff" – Faculdade de Psicologia, Instituto de Estudos em Saúde e Biológicas, Universidade Federal do Sul e Sudeste do Pará, Unidade III

Av. dos Ipês, S/N, s/ CEP, Bairro Cidade Jardim, Marabá/PA, Brazil

cmaximino@unifesspa.edu.br

Abstract

Orienting responses (ORs) are whole-organism reflexes that are elicited by innocuous stimuli, and which decrease in magnitude after stimulus repetition. ORs represent relatively simple responses that can be used to study attentional processes, and are modulated by the organism's state, including arousal and activation levels, as well as by emotional processes. Here we describe a simple method to study ORs in zebrafish, a model organism increasingly being used in behavioural neuroscience. After presentation of a static visual stimulus, an OR is elicited, characterized by approaching the stimulus and orienting towards it. After repeated stimulation, OR decreases, suggesting habituation. These responses are qualitatively altered by exposure to a fear-eliciting alarm substance, since exposed animals avoid the visual stimulus and orient either away from the stimulus or towards it, but at a distance. The protocol can be used to study orienting responses, as well as the impact of fear and arousal on these reflexes.

1. Introduction

Zebrafish is an emerging model organism in behavioural neuroscience, and is increasingly being favoured due to its advantages both as a model organism (i.e., ease of upkeep, handling, and reproduction; Key & Devine, 2003) and intermediate physiological and behavioural complexity (Kalueff, Echevarria, & Stewart, 2014; Kokel & Peterson, 2008; Stewart et al., 2015). As a result, zebrafish has been introduced as a model organism to study the neurochemistry and circuitry of cognition, including executive function and attention (Parker, Brock, Walton, & Brennan, 2013; Stewart & Kalueff, 2012). Attention can be described as selective or sustained; the first case refers to the ability to select a target from an array of distractors, while the second is defined as the ability to detect the presence of a target over a prolonged period of time (Parker et al., 2013). Both selective and sustained attention can be measured in different behavioural models, including zebrafish. In the 3-choice serial reaction time task, fish are taught to swim into one of three apertures after maintaining attention to the stimuli (Parker, Millington, Combe, & Brennan, 2012), and is useful to study sustained attention. In the virtual object recognition task (Braida, Ponzoni, Martucci, & Sala, 2014), selective attention is evaluated by observing the time the animal spends exploring novel shapes presented in a screen. In both cases, drugs which increase attention are able to improve performance.

Orienting responses (ORs) involve a set of behavioural and physiological adjustments that are elicited by novel and/or significant stimuli, leading to the motivational activation of a cascade of perceptual and motor processes that facilitate behavioural selection (M. M. Bradley, 2009). ORs are whole-organism reflexes that are elicited by innocuous stimuli, and which decrease in magnitude (habituate) after stimulus repetition (Pavlov, 1927; Sokolov, 1963). The virtual object recognition task (Braida et al., 2014) described above is a classical repetition-change paradigm, in which one "truly novel" stimulus is repeatedly presented, and responses to this stimulus are recorded as "orienting responses", and the decreased responsiveness after a number of presentations is recorded

as habituation; at certain intervals, a different stimulus is presented, and orienting responses to this

novel stimulus can also be recorded (dishabituation). While ORs are classically interpreted in terms

of stimulus novelty, the significance of the stimulus (including its relationship to unconditioned

appetitive or aversive cues) is highly important to orienting (Barry, 2009; Donchin, 1981). The

relationship between stimulus significance and ORs is an important factor in judging the attention-

emotion interaction.

Attention and emotion can interact through the influence of the first on emotional

processing, or through the emotional modulation of the attention processes (Vuilleumier, 2005). In

that case, once a threat is detected and the initial (automatic) fear response is elicited, further action

is needed, demanding the redirection of attentional resources to the threatening stimulus and the

engagement of flexible behavioural repertoires. The influence of emotion on spatial attention has

been investigated in humans (Beck, Stanley, Averill, Baldwin, & Deagle, 1992; B. P. Bradley,

Mogg, & Millar, 2000), and is relevant to understanding psychiatric disorders, given that patients

suffering from anxiety disorders present changes in attention and selectivity to threat-associated

stimuli (Asmundson, Sandler, Wilson, & Walker, 1992; Beck et al., 1992; B. P. Bradley et al., 2000;

Lautenbacher, Spernal, & Krieg, 2002). Developing methods to study attention and ORs that are

sensitive to emotional modulation is useful from a translational point of view, increasing the breadth

of affected domains in an integrative perspective (Kalueff, Ren-Patterson, LaPorte, & Murphy,

2008).

Conspecific alarm substance (CAS) is a complex mixture produced by epidermal club cells

and released in the water after these cells are injured by, for example, a predator (Maximino et al.,

2019). The substance is detected by conspecifics, eliciting defensive (antipredator) behaviour

(Døving & Lastein, 2009; Maximino et al., 2019; von Frisch, 1938, 1941). These behavioural

adjustments include erratic swimming and freezing, which increase the probability of detecting a

potential threat and decrease the probability of being detected by a predator (Maximino et al.,

2019). In that sense, CAS appears to act partially by modulating ORs, either increasing orienting

towards novel and/or significant stimuli to promote appraisal, or producing faster responses. In the

first case, freezing responses can promote conditions of focused attention towards environmental

novelty, increasing detection of the predator; in the second case, fast responses can in fact be

detrimental to attention, producing false alarms that are nonetheless counterbalanced by the higher

probability of escaping the predator.

The aims of the present article were to 1) propose a simple task to assess ORs in zebrafish

towards a novel neutral stimulus (a blue circle); 2) assess whether CAS increases orienting or elicits

escape responses away form this novel stimulus. We find that the stimulus induced an approach

response, accompanied by orienting towards the stimulus, that habituated after 5 trials. We also find

that CAS increased erratic swimming throughout the task, and inverted the direction of movement

during stimulus presentation: instead of approaching the stimulus, animals now flew from it. This

manuscript is a complete report of all the studies performed to test the effect of a visual stimulus on

ORs in zebrafish and of CAS on these orienting responses. We report all data exclusions (if any), all

manipulations, and all measures in the study (Simmons, Nelson, & Simonsohn, 2012).

2. Methods

2.1. Animals and housing

36 adult (~4 mo.) zebrafish from the longfin phenotype were acquired from local creators

and group-housed in tanks with a stock density of 5 animals/2 L. Temperature was kept at 28±2 °C,

hardness 75-200 mg/L $CaCO_3$, salinity 0.25-0.75 ppt, dissolved oxygen ~7,8 mg/L at 28 °C, and

nitrite levels <0.01 ppm. Animals were kept in the tank for at least two weeks to acclimate before

beginning experiments. Environmental characteristics (physico-chemical and health profile) for the

housing can be found at GitHub (https://github.com/lanec-unifesspa/lanec-welfare). All

manipulations and animal care were in accordance with Brazilian guidelines (Conselho Nacional de

Controle de Experimentação Animal - CONCEA, 2017)

2.2. Sampling plan

Effect sizes for sample size calculation were based on studies on the visual object recognition test (Braida et al., 2014). Since fixed effects (trial and group) were considered more important than the random (subject) effect, sample sizes were calculated based on two-way ANOVAs, with power = 80%, and considering $p < 0.05$ as a target. After calculation, 12 animals per group were considered sufficient.

2.3. Behavioural setup

The experimental setup consisted of two 13 x 13 x 17 (depth x length x depth cm each) glass tanks, positioned side by side, and filled with 2 L system water. The tanks were positioned at the front of the computer screen (Samsung T20c310lb, 20", LED screen, nominal brightness 20 cd/m³) such that the stimulus appeared in front of the subjects. The computer screen was located at 8.5 cm of the tank centre. All sides of the tanks, except the one facing the screen, were covered in opaque white plastic sheets, isolating animals from each other and improving contrast for tracking.

2.4. Experimental procedures

A blue circle (R: 0; G: 0; B: 128) was used, based on the ability of zebrafish to discriminate blue from green (Mueller & Neuhauss, 2012). Stimulus size was 72º of visual angle, calculated considering the diameter of the stimulus in the screen when it is at the from of the animal, and the animal is in the centre of the tank. The size was based on previous work which show that this size is insufficient to produce an escape response in zebrafish larvae (Dunn et al., 2016).

Animals were transferred to the tank and left to acclimate for 3 min before the presentation

of the stimuli. The computer screen was turned on for the entire experiment. Using an animation

based on LibreOffice Presentation (v. 6.0.7.3), the stimulus was presented in 1-min intervals

("trials") interspersed with 1-min stimulus-free periods. 10 trials were made; total session length

was 22 minutes per animal.

2.5. Behavioural tracking

A video camera (Sony Handycam® DCR-DVD 610) was positioned above the tanks to

allow recording and offline behavioural scoring. All focal fish behaviours were therefore tracked

from   a   top-down   perspective,   using   both   TheRealFishTracker

(http://www.dgp.toronto.edu/~mccrae/projects/FishTracker/)   and   fishtracker

(https://github.com/joseaccruz/fishtracker). For each behavioural video, a 2D region (arena) was

defined for tracking, comprising the inner area of the tank. A region-of-interest (ROI) of 13 x 8.5

cm   was   determined   as   "stimulus   zone".   The   following   variables   were   recorded,   using

TheRealFishTracker:

• Total time in the stimulus zone (s);

• Absolute turn angle (°), a measure of erratic swimming (Tran, Chatterjee, Facciol, & Gerlai,

2016);

• Speed (cm/s)

Following Abril-de-Abreu, Cruz, & Oliveira (2015), a region of interest (ROI) of 13 x 3 cm

(approximately 25% of the tank), corresponding to the width of the tank and the mean body length

of an adult zebrafish, was defined in the area of the arena closer to the computer screen in

fishtracker. Animal orientation is defined as a mean projection vector R_{proj} (Abril-de-Abreu et al.,

2015), determined by the transformation of each orientation angle in a vector $r_i = (cos\alpha_i, sin\alpha_i)$,

where α_i represents the angle formed by the direction of the centroid-head axis relative to the

horizontal axis in each frame. The resulting mean vector, with length $R = \|r\|$, is a measure of

directional focus, and varies between 0 and 1. The projection of this vector R in the stimulus

direction (arbitrarily defined as 180º) is defined as $R_{proj} = -Rcos\alpha$. Positive values indicate direction

towards the stimulus, while negative values indicate direction away from the stimulus, and values

close to 0 indicate absence of directionality.

2.6. CAS extraction and exposure

CAS extraction was made according to a protocol described in full detail elsewhere (do

Carmo Silva, Rocha, Lima-Maximino, & Maximino, 2018). Briefly, after sacrifice, a single fish was

subjected to 15 shallow cuts at both sides of the body, which were subsequently washed with 10 mL

distilled water. After removing debris, 7 mL of this solution was separated, and kept on ice until

used; this was referred to as "1 unit CAS". For half of the animals ($n = 12$), CAS was poured

directly into the tank water immediately after the animal was transferred to the tank. For the other

half ($n = 12$), distilled water was used.

2.7. Statistics

For data on time in stimulus zone, absolute turn angle, and speed, a mixed-effects model

was applied, with subject as random factor, trial as within-subjects factor, and group as between-

subjects factor. A temporal autocorrelation structure of order 1 was modelled within levels of the

random factor. Analyses were made using the R package 'nlme' (v. 3.1-141; https://CRAN.R-

project.org/package=nlme). For projection angle data, values were averaged across equal trials (i.e.,

stimulus "ON" vs. stimulus "OFF"), and a Watson-Williams test for homogeneity of circular data

was applied, using the 'circular' package (v. 0.4-93; https://r-forge.r-project.org/projects/circular/).

For R_{proj} data, a two-way (treatment X trial type) analysis of variance (ANOVA) was applied,

followed by Tukey's HSD post-hoc test whenever $p < 0.05$. All analyses were run on R 3.6.1 (R

Core Team, 2019).

2.8. Open science practices

Experiments were not formally preregistered. Data and analysis scripts can be found at a

GitHub repository (https://github.com/lanec-unifesspa/5-HT-CAS/tree/master/data/behavioral/

orienting). Preprint versions of the manuscripts were uploaded to bioRxiv.

3. Results

A linear model including subject as random factor was favored in relation to a model

without the random factor (likelihood ratio = 139.5079, p < 0.0001, AIC = 3986.042 vs 4121.55).

Main effects of group (χ^2 = 134.443, p < 2.2 x 10^{-16}) and trial (χ^2 = 64.747, p = 8.52 x 10^{-16}), as well

as a interaction effect (χ^2 = 27.905, p = 1.27 x 10^{-7}), were found for time in the stimulus zone

(Figure 1A). Animals in the control group spend more time near the stimulus when it was "ON",

showing evidence of an orienting response that decreased as the stimulus was repeated, suggesting

habituation. Animals exposed to CAS, on the other hand, spent *less* time in the stimulus zone when

the stimulus was "ON", suggesting that the stimulus was interpreted as threatening. A main effect of

group ($F_{[1, 524]}$ = 305.39, p < 0.0001), but not trial ($F_{[1, 524]}$ = 1.209, p = 0.272), was found for absolute

turn angle (Figure 1B); CAS increased absolute turn angle during the entire test. No effects of group

($F_{[1, 525]}$ = 0.019, p = 0.891) or trials ($F_{[1,525]}$ = 0.621, p = 0.431) were found for swimming speed

(Figure 1C).

For R_{proj} data, 2 animals from the control group, and three animals from the CAS-exposed

group, were removed because the software was not able to detect them. A significant treatment X

trial type interaction effect was found for angle ($F_{[1,36]}$ = 13.383, p = 8.07x10^{-4}; Figure 2A). A

significant interaction was also found for R_{proj} ($F_{[1, 34]}$ = 88.639, p = 5.34x10^{-11}; Figure 2B); in the

control group, the presentation of the stimulus increased R_{proj} ($p < 0.0001$), while in the CAS group

stimulus presentation decreased R_{proj} ($p = 0.000006$).

4. Discussion

In this work we demonstrate a simple method to elicit orienting responses in zebrafish

towards a visual stimulus. The orienting response was characterized by an approach to the stimulus,

as well as body orientation towards the stimulus when it was presented. These orienting responses

follow the classical pattern of habituation after repeated stimulus presentation, suggesting that they

accurately capture orienting. Moreover, we show that high increases in arousal by exposing animals

to an alarm substance abolish the orientation response and substitute it for a defence response.

Orienting responses are relatively simple and amenable to manipulations of different

components, and have translational value to understand the emotion-cognition interaction (M. M.

Bradley, 2009). Orienting responses allow a preliminary processing of the environment in terms of

stimulus significance, and selectively filter irrelevant information that arrives through sensory

channels (Barry, 2009; Sokolov, 1963). During stimulus presentation, zebrafish in the control group

responded with an orienting response, approaching the stimulus and changing body orientation to

face it. As stimulus presentation continued, this response showed signs of habituation. Thus, in the

initial trials, orienting responses allowed sensory information to be analysed by perceptual areas,

which results in information about the stimulus being available to perceptual processes. As the

novelty of the stimulus decreases, habituation processes lead to a decline in the orienting response

(Pavlov, 1927; Sokolov, 1963).

Orienting responses are in part mediated by the locus coeruleus, and noradrenaline release in

the forebrain will facilitate sensory processing, enhance cognitive flexibility and executive function,

and promote memory consolidation in limbic structures (Sara & Bouret, 2012). Stednitz et al.

(2018) used orienting responses towards conspecifics to dissect mechanisms of social attention in

zebrafish, and found that lesions in cholinergic neurons of the ventromedial telencephalon

decreased the orienting response towards conspecifics. While it is not known whether these

cholinergic neurons also participate non-social orienting responses, acetylcholine and noradrenaline

both have been implicated in the effects of arousal in attention.

Arousal levels are thought to amplify the phasic orienting response by sensitizing the

organism to both significant and non-significant stimuli (Barry, 2009). In the present work,

however, orienting was not simply amplified by CAS, but qualitatively changed it. During stimulus

presentation, animals avoided it, spending more time away from the stimulus and either orienting

away from it or towards it. Moreover, this response did not habituate throughout trials, and was

observed also in the absence of the stimulus – although it was significantly increased by stimulus

presentation. CAS also elicited an increase in absolute turn angle, suggesting erratic swimming, as

observed in zebrafish exposed to CAS and other fear-eliciting stimuli (Ahmed, Fernandes, & Gerlai,

2012; Gerlai, Fernandes, & Pereira, 2009; Maximino et al., 2019). CAS is known to elicit fear-like

behaviour in fish from the Ostariophysan superorder, decreasing the probability of predation by

promoting escape or freezing, and increasing the probability that conspecifics detect a predator by

inducing increased arousal and attention (Maximino et al., 2019; Smith, 1992). These results

suggest that CAS-induced arousal did not merely amplify the orienting response, but induced a state

of hypervigilance, in which a non-significant stimulus is interpreted as threatening.

Taken in conjunction, our results suggest a protocol to evoke orienting responses in

zebrafish as a method to study attention, as well as an activating effect of the alarm substance in

these responses. The results underline an important emotion-cognition interface in zebrafish that

can be exploited to study behaviour and that has implications to using zebrafish behavioural models

to study biological psychiatry. Future work is needed to understand the relationship between these

effects and arousal-associated molecules.

Acknowledgements

This work was supported by a Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq/Brazil) grant (Edital Universal 2016, #400726/2016-6). BGN was the recipient of a CNPq/Brazil scholarship.

References

Abril-de-Abreu, R., Cruz, J., & Oliveira, R. F. (2015). Social eavesdropping in zebrafish: Tuning of attention to social interactions. *Scientific Reports, 5,* 12678. https://doi.org/10.1038/srep12678

Ahmed, T. S., Fernandes, Y., & Gerlai, R. (2012). Effects of animated images of sympatric predators and abstract shapes on fear responses in zebrafish. *Behaviour, 149,* 1125–1153. https://doi.org/10.1163/1568539X-00003011

Asmundson, G. J. G., Sandler, L. S., Wilson, K. G., & Walker, J. R. (1992). Selective attention toward physical threat in patients with panic disorder. *Journal of Anxiety Disorders, 6*(4), 295–303. https://doi.org/10.1016/0887-6185(92)90001-N

Barry, R. J. (2009). Habituation of the orienting reflex and the development of Preliminary Process Theory. *Neurobiology of Learning and Memory, 92,* 235–242. https://doi.org/10.1016/j.nlm.2008.07.007

Beck, J. G., Stanley, M. A., Averill, P. M., Baldwin, L. E., & Deagle, E. A. (1992). Attention and memory for threat in panic disorder. *Behaviour Research and Therapy, 30*(6), 619–629. https://doi.org/10.1016/0005-7967(92)90007-4

Bradley, B. P., Mogg, K., & Millar, N. H. (2000). Covert and overt orienting of attention to emotional faces in anxiety. *Cognition and Emotion, 14,* 789–808. https://doi.org/10.1080/02699930050156636

Bradley, M. M. (2009). Natural selective attention: Orienting and emotion. *Psychophysiology, 46,* 1–11. https://doi.org/10.1111/j.1469-8986.2008.00702.x

Braida, D., Ponzoni, L., Martucci, R., & Sala, M. (2014). A new model to study visual attention in zebrafish. *Progress in Neuro-Psychopharmacology & Biological Psychiatry, 55,* 80–86. https://doi.org/10.1016/j.pnpbp.2014.03.010

Conselho Nacional de Controle de Experimentação Animal - CONCEA. *Diretriz brasileira para o cuidado e a utilização de animais para fins científicos e didáticos - DBCA. Anexo I. Peixes mantidos em instalações de instituições de ensino ou pesquisa científic*, Resolução Normativa n. 30/2016 (2017).

do Carmo Silva, R. X., Rocha, S. P., Lima-Maximino, M. G., & Maximino, C. (2018). Extraction of alarm substance in zebrafish. *Protocols.Io*. https://doi.org/10.17504/protocols.io.tr3em8n

Donchin, E. (1981). Surprise! ... Surprise? *Psychophysiology*, *18*, 493–513. https://doi.org/10.1111/j.1469-8986.1981.tb01815.x

Døving, K. B., & Lastein, S. (2009). The alarm reaction in fishes - Odorants, modulations of responses, neural pathways. *Annals of the New York Academy of Sciences*, *1170*, 413–423. https://doi.org/10.1111/j.1749-6632.2009.04111.x

Dunn, T. W. W., Gebhardt, C., Naumann, E. A. A., Riegler, C., Ahrens, M. B. B., Engert, F., & Del Bene, F. (2016). Neural circuits underlying visually evoked escapes in larval zebrafish. *Neuron*, *89*, 613–628. https://doi.org/10.1016/j.neuron.2015.12.021

Gerlai, R., Fernandes, Y., & Pereira, T. (2009). Zebrafish (Danio rerio) responds to the animated image of a predator: Towards the development of an automated aversive task. *Behavioural Brain Research*, *201*, 318–324. https://doi.org/10.1016/j.bbr.2009.03.003

Kalueff, A. V, Echevarria, D. J., & Stewart, A. M. (2014). Gaining translational momentum: More zebrafish models for neuroscience research. *Progress in Neuro-Psychopharmacology & Biological Psychiatry*, *55*, 1–6. https://doi.org/10.1016/j.pnpbp.2014.01.022

Kalueff, A. V, Ren-Patterson, R. F., LaPorte, J. L., & Murphy, D. L. (2008). Domain interplay concept in animal models of neuropsychiatric disorders: A new strategy for high-throughput neurophenotyping research. *Behavioural Brain Research*, *188*, 243–249. https://doi.org/10.1016/j.bbr.2007.11.011

Key, B., & Devine, C. A. (2003). Zebrafish as an experimental model: Strategies for developmental and molecular neurobiology studies. *Methods in Cell Science*, *25*, 1–6. https://doi.org/10.1023/B:MICS.0000006849.98007.03

Kokel, D., & Peterson, R. T. (2008). Chemobehavioural phenomics and behaviour-based psychiatric drug discovery in the zebrafish. *Briefings in Functional Genomics and Proteomics*, *7*, 483–490.

Lautenbacher, S., Spernal, J., & Krieg, J.-C. (2002). Divided and selective attention in panic disorder: A comparative study of patients with panic disorder, major depression and healthy controls. *European Archives of Psychiatry and Clinical Neuroscience*, *252*(5), 210–213. https://doi.org/10.1007/s00406-002-0382-5

Maximino, C., do Carmo Silva, R. X., dos Santos Campos, K., de Oliveira, J. S., Rocha, S. P., Pyterson, M. P., … Lima-Maximino, M. G. (2019). Sensory ecology of ostariophysan alarm substances. *Journal of Fish Biology*, *95*, 274–286. https://doi.org/10.1111/jfb.13844

Mueller, K. P., & Neuhauss, S. C. F. (2012). Automated visual choice discrimination learning in zebrafish (*Danio rerio*). *Journal of Integrative Neuroscience*, *11*, 73–85. https://doi.org/10.1142/S0219635212500057

Parker, M. O., Brock, A. J., Walton, R. T., & Brennan, C. H. (2013). The role of zebrafish (*Danio*
rerio) in dissecting the genetics and neural circuits of executive function. *Frontiers in Neural*
Circuits, *7*, Article 63. https://doi.org/10.3389/fncir.2013.00063

Parker, M. O., Millington, M. E., Combe, F. J., & Brennan, C. H. (2012). Development and
implementation of a three-choice serial reaction time task for zebrafish (*Danio rerio*).
Behavioural Brain Research, *227*, 73–80. https://doi.org/10.1016/j.bbr.2011.10.037

Pavlov, I. P. (1927). *Conditioned reflexes*. Londres: Oxford University Press.

R Core Team. (2019). *R: A language and environment for statistical computing*. Retrieved from
http://www.r-project.org/

Sara, S. J., & Bouret, S. (2012). Orienting and reorienting: The locus coeruleus mediates cognition
through arousal. *Neuron*, *76*, 130–141. https://doi.org/10.1016/j.neuron.2012.09.011

Simmons, J. P., Nelson, L. D., & Simonsohn, U. (2012). A 21-word solution. *Dialogue*, *26*, 4–7.

Smith, R. J. F. (1992). Alarm signals in fishes. *Reviews in Fish Biology and Fisheries*, *2*, 33–63.
https://doi.org/10.1007/BF00042916

Sokolov, E. N. (1963). Higher Nervous Functions: The Orienting Reflex. *Annual Review of*
Physiology, *25*, 545–580. https://doi.org/10.1146/annurev.ph.25.030163.002553

Stednitz, S. J., McDermott, E. M., Ncube, D., Tallafuss, A., Eisen, J. S., & Washbourne, P. (2018).
Forebrain Control of Behaviorally Driven Social Orienting in Zebrafish. *Current Biology*, *28*,
2445–2451. https://doi.org/10.1016/j.cub.2018.06.016

Stewart, A. M., & Kalueff, A. V. (2012). The developing utility of zebrafish models for cognitive
enhancers research. *Current Neuropharmacology*, *10*, 263–271.

Stewart, A. M., Ullmann, J. F. P., Norton, W. H. J., Parker, M. O., Brennan, C. H., Gerlai, R., &
Kalueff, A. V. (2015). Molecular psychiatry of zebrafish. *Molecular Psychiatry*, *20*, 2–17.
https://doi.org/10.1038/mp.2014.128

Tran, S., Chatterjee, D., Facciol, A., & Gerlai, R. (2016). Concentration, population, and context-
dependent effects of AM251 in zebrafish. *Psychopharmacology*, *233*, 1445–1454.
https://doi.org/10.1007/s00213-016-4240-y

von Frisch, K. (1938). Zur psychologie des Fische-Schwarmes. *Naturwissenschaften*, *26*, 601–606.

von Frisch, K. (1941). Über einen Schreckstoff der fischhaut und seine biologische Bedeutung.
Zeitschrift Für Vergleichende Physiologie, *29*, 46–145. https://doi.org/10.1007/BF00304445

Vuilleumier, P. (2005). How brains beware: Neural mechanisms of emotional attention. *Trends in*
Cognitive Sciences, *9*(12), 585–594. https://doi.org/10.1016/j.tics.2005.10.011
364

Figure captions

Figure 1 – A visual stimulus induces approach responses in zebrafish, while animals exposed to CAS avoid the stimulus and show erratic swimming. Time series scatterplots (n = 12/treatment) with (A) time spent near the stimulus, (B) absolute turn angle (erratic swimming), and (C) swimming speed, in 1-min blocks. Stimuli were presented at minutes 4, 6, 8, 10, 12, 14, 16, 18, 20, and 22. Lines and bands represent group averages and bootstrapped 95% confidence intervals around the mean.

Figure 2 – Orienting responses towards a visual stimulus in zebrafish, exposed or not to an alarm substance. (A) Polar plots of the focal fishes' individual mean resultant vector's angles (0º to 360º, stimulus positioned at 180º), combined with corresponding vector lengths R (0 to 1) for each treatment (control – black dots; CAS – red dots); when the stimulus is on, animals from the control group significantly deviate from a uniform distribution, clustering around its group mean resultant vector. (B) Scatter plots (n = 9-12/treatment) of the individual (coloured dots) resultant vector's lengths R projected onto the stimulus direction (R_{proj}), and corresponding group summaries (boxplots), for each treatment; positive values indicate directional focus towards the stimulus, zero indicates no directionality, and negative values indicate focus opposite to the stimulus.